Leaf economics explained by optimality principles

Han Wang[1*], I. Colin Prentice[1,2,3], Ian J. Wright[3], Shengchao Qiao[1], Xiangtao Xu[4], Kihachiro Kikuzawa[5], Nils Chr. Stenseth[1,6]

[1] Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing 100084, China

[2] Department of Life Sciences, Imperial College London, Silwood Park Campus, Buckhurst Road, Ascot SL5 7PY, UK

[3] Department of Biological Sciences, Macquarie University, North Ryde, NSW 2109, Australia

[4] Ecology and Evolutionary Biology, Cornell University, E139 Corson Hall, Ithaca, NY 14850, USA

[5] Laboratory of Plant Ecology, Ishikawa Prefectural University, Nonoichi, Ishikawa, 921-8836 Japan

[6] Centre for Ecological and Evolutionary Synthesis (CEES), Department of Biosciences, University of Oslo, P.O. Box 1066 Blindern NO-0316 Oslo, Norway

*Correspondence to: wang_han@mail.tsinghua.edu.cn

SUMMARY

The worldwide leaf economics spectrum relates leaf lifespan (LL) to leaf dry mass per unit area (LMA)[1]. By combining three well-supported principles[2-4], we show that an isometric relationship between these two quantities maximizes the leaf's net carbon gain. This theory predicts a spectrum of equally competent LMA-LL combinations in any given environment, and how their optimal ratio varies across environments. By analysing two large, independent leaf-trait datasets for woody species[1,5], we provide quantitative empirical support for the predicted dependencies of LL on LMA and environment in evergreen plants, and for the distinct predicted dependencies of LMA on light, temperature, growing-season length and aridity in evergreen and deciduous plants. We thereby resolve the long-standing question of why deciduous LMA tends to increase (with increasing LL) towards the equator, while evergreen LMA and LL decrease[6]. We also show how the statistical distribution of LMA within communities can be modelled as an outcome of environmental selection on the global pool of species with diverse values of LMA and LL.

Main

Plants are subject to powerful selective forces in every aspect of their lives, favouring economic efficiency of vegetative processes as a necessary condition of reproductive success in a competitive milieu. We will show here how the worldwide leaf economics spectrum (LES) – the generalization that thicker and/or denser leaves (high LMA) have longer lifespans (high LL)[1] – arises as a consequence of leaf-level maximization of net carbon gain.

LL and LMA vary several hundredfold across vascular plants, and up to tenfold among co-occurring species[1,7]. Central values of LL, LMA and their ratio also shift along climatic gradients[8-10], but the reasons are not well understood. The environmental controls on LL and LMA remain unclear, as do the different responses of deciduous and evergreen species[11]. In particular, why do the leaves of deciduous trees and shrubs tend to become thinner (as their lifespan declines) poleward, while those of evergreen species become thicker[12] and longer-lived[13]? Better understanding of the LES is required: not least for more robust modelling of the global carbon cycle and its interactions with climate, as LL and LMA directly influence carbon cycling in terrestrial ecosystems by determining the carbon requirements for leaf construction and turnover[14-16].

Recent research has shown that eco-evolutionary optimality hypotheses can predict both

known and unexpected patterns of relationship among plant traits, and between traits and

environment[17]. The increasing availability of large plant trait data sets allows quantitative

assessment of such predictions[18]. Much research has focused on leaf physiological traits, like

photosynthetic capacity, that can change within the same plants (and even the same leaves) in

response to seasonal environmental cues[19,20]. Unlike physiological traits, however, LMA and

LL generally show more muted intraspecific responses to temporal changes and geographic

gradients in environmental factors. Differences in the phenotypic plasticity and/or genotypic

adaptability of traits are revealed by comparison of within-species and between-community

trends along environmental gradients[21]. In a continental-scale analysis, intraspecific shifts in

LMA showed only about one-third of the response to environmental variation shown by

community-mean LMA[21], compared to (for example) > 90% for the ratio of leaf-internal to

ambient CO_2, a key physiological trait involved in photosynthesis[22]. Nonetheless, the LES can

be demonstrated across species worldwide[1] and the associated traits shift with environment in

consistent ways, strongly suggesting the existence of a common organizing principle[8-10,23].

An optimality theory for the leaf economic spectrum

An eco-evolutionary optimality approach is developed here to account for the environmental

controls on leaf economics for woody species in a quantitative framework. First, we consider

the relationship between LMA and LL that maximizes the leaf life-cycle average net carbon

gain, i.e. net photosynthesis minus the associated (amortized) tissue construction costs[2]. This

relationship depends on the rate at which photosynthesis declines with leaf age[3]. Second, the

ageing rate increases in proportion to the leaf's initial photosynthetic capacity, as measured by

the maximum rate of carboxylation at a standard temperature of 25˚C (V_{cmax25}), and decreases

in proportion to LMA[3]. Third, photosynthetic capacity itself is optimized to the physical

environment, following the coordination hypothesis[4,24,25]. Combining these three principles

(see Methods for derivations) we obtained the following general relationship between LMA

and optimal LL (LL_{ev}) in evergreen leaves:

LL_{ev} = $(LMA/A_0) \sqrt{(2 u C_c h_T m_c / f)}$ (1)

where A_0 is the leaf's initial photosynthetic rate; f is the growing-season length, as a fraction

of the year; u is a constant that relates the ageing rate to LMA and photosynthetic capacity; C_c

is a multiplier for the total carbon costs, including required investments in non-leaf tissues,

associated with leaf construction; h_T is the Arrhenius function relating V_{cmax} to temperature (T),

equal to unity when $T = 298.15$ K; and m_c is the ratio of the Rubisco-limited photosynthetic

rate to V_{cmax}, which is a function of the intercellular CO_2 partial pressure and the affinity of

Rubisco for CO_2 versus O_2. The coordination hypothesis leads to the prediction that A_0 is

proportional to the photosynthetic photon flux density (PPFD) absorbed by the leaf[20].

This theory predicts the existence of a spectrum of values of both quantities, from low LMA

and LL to high LMA and LL, resulting in equal rates of net carbon gain (see Methods). The

optimal LL in a given environment is proportional to LMA, and inversely proportional to

absorbed PPFD and the square root of growing-season length. We show here (Figure 1a-c,

Table S1) that these predictions are similar to the corresponding partial effects independently

inferred from a multiple regression fitted to the measurements on evergreen species in the

GlopNet data set[1], which provides the largest available worldwide compilation of paired data

on LMA and LL. The regression for evergreen species captures 42% of the variation in LL.

Evergreen LL is proportional to LMA (log-log slope ≈ 1) as predicted. The relationship of LL

to PPFD is negative as predicted, and the relationship of the LL residuals to growing-season

length is also negative ($P < 0.1$) as predicted, with predicted slopes similar in magnitude to

fitted values (although falling just outside their predicted confidence intervals).

Fig. 1 | Partial residual plots from the regression of observed leaf longevity (LL) against explanatory

variables for evergreen and deciduous species. a, LMA: leaf mass per area. **b**, I: site-mean leaf absorbed

Photosynthetic Photon Flux Density. **c and d**, f: growing-season length as a fraction of the year. Predicted

relationships in red, fitted in black with 95% confidence interval in grey; evergreen species in green,

deciduous in orange. All axes are on log scales. Data from the GlopNet trait data set[1].

Due to the different behaviour of evergreen and deciduous species in the unfavourable season,

the same optimality framework leads to a different equation that describes net carbon-gain

maximization in deciduous species (see Methods for derivation):

$LL_{de} = (LMA/A_0)\,(u\,h_T\,m_c)$ (2)

In addition to this optimality criterion, LL_{de} is tightly constrained by growing-season length.

Among the deciduous species in GlopNet, growing-season length is indeed nearly proportional

to LL (0.79 ± 0.08, 1 s.e.), and explains about half of the observed variation in LL (Figure 1d,

Table S1).

Equations (1) and (2) imply that for any given LL, both evergreen and deciduous LMA should

increase in direct proportion to absorbed PPFD. All else equal, LMA should also decrease by

2.6% K^{-1} with increasing temperature in evergreen species, and at about twice that rate in

deciduous species (see Methods). The response of LMA to growth temperature is mainly a

consequence of the thermal acclimation (down-regulation) of carboxylation capacity, which

implies slower leaf ageing at higher temperatures, and therefore reduces the carbon investment

needed to produce a unit of photosynthate. High LMA is also expected (and found) in plants

of arid environments, because maintaining turgor at low leaf water potential requires leaves to

be stiff[26], and because thin leaves have insufficient heat capacity to prevent overheating during

transient low-wind episodes[27].

The environmental dependencies of leaf mass per area

This theory predicts different environmental dependencies of LMA in evergreen and deciduous

plants. To quantify these dependencies, we must consider how environmental constraints act

on prior probability distributions of LMA and LL. The principle is illustrated in Figure S2,

where straight lines in the log-log plot of LL versus LMA – representing the optimal LES for

two sites in different environments – intersect lognormal distributions of both variables. In

deciduous plants, site-mean LL is constrained by growing-season length. In evergreen plants,

site-mean LMA is predicted at the mean of the distribution formed by the intersection of the

lines with the prior distribution. The resulting predicted sensitivities of evergreen LMA to

temperature and PPFD are only half those for deciduous LMA (see Methods). For growing-

season length, they are only a quarter of those for deciduous LMA.

These predictions are fully supported by analysis of data in the China Plant Trait Database

(CPTD)[5], which provides a larger LMA data set than GlopNet for temperate and boreal

deciduous species, many more observations for deciduous species in particular, and more than

three times as many sites. Growing-season PPFD, temperature and an index of plant-available

moisture (see Methods) all influence the LMA of evergreen species (Figure 2a-c, Table S2).

The predictive power of this relationship is limited (13%), however, presumably because environmental predictors alone can only explain a limited proportion of the variation in LMA (additional variation being driven by variation in LL that is independent of environment) because the species span a wide range of LL. Growing-season length is also a significant predictor of the residual variation in evergreen LMA (Figure 2d, Table S2). The same four environmental predictors account for more than half of the variation in LMA among deciduous species in the CPTD (Figure 2e-h, Table S2), where growing-season length is expected to largely determine LL and so to be proportional to LMA, as observed. The confidence intervals of the fitted slope coefficients for PPFD, temperature and growing-season length all include the predicted values for both evergreen and deciduous species.

Fig. 2 | Partial residual plots from the regression of observed leaf mass per area (LMA) against environmental variables for evergreen and deciduous species. a and e, I: site-mean leaf absorbed Photosynthetic Photon Flux Density. **b and f**, T_g: the mean temperature of the growing season, when daily temperature is above 0˚C. **c and g**, α: moisture index (the ratio of actual to equilibrium evapotranspiration). **d and h**, f: growing-season length as a fraction of the year. Predicted relationships in red, fitted in black with 95% confidence interval in grey; evergreen species in green, deciduous in orange. All axes except for T_g are on log scales. Data from an extended version of the China Plant Trait database[5,28].

For evergreen species, whose LL is not directly constrained by the growing-season length, it is useful to predict not only a community-mean value for LMA but also a statistical distribution representing the site-specific LES. We fitted independent lognormal distributions of LMA and LL based on GlopNet data, and imposed equation (1) as a constraint. This method successfully predicted within-community LMA distributions in evergreen temperate forests, tropical

rainforests and woodlands (Figure 3). The community-mean LMA values of temperate forests and tropical rainforests are close to the global mean value of LMA derived from the GlopNet data set. They are predicted to be similar because growth temperature and growing-season PPFD and duration are positively correlated, but to have opposite effects on LMA. The community-mean LMA of woodlands (Figs 3e,f) however shifts far from the global mean, because of the additional ecophysiological constraint imposed by aridity.

Fig. 3 | Observed and predicted within-site distributions of leaf mass per area (LMA). a and b, temperate forests. **c and d**, tropical rainforests. **e and f**, woodlands[29]. Means and standard errors of the observed (obs) and predicted (pre) distributions are shown. Predicted probability distribution functions (red curve) are scaled in height to match the observed frequency distribution. Data from the GLOPNET trait data set [1]. Sites with the largest sample size are selected; see detailed site information in Supplementary Table S3.

This theory accounts for the contrasting latitudinal patterns in LMA between deciduous and evergreen species[6], as shown in Figures 4 and S3. The expected effect of increasing temperature towards the equator on LMA is negative for both leaf types. However, in deciduous species, this influence is outweighed by the effects of increasing growing-season PPFD and length, both of which favour increased LMA.

Fig. 4 | Observed and predicted trends of leaf mass per area (LMA) from tropical to polar regions. Data and fitted regression lines with 95% confidence intervals are in green for evergreen and orange for deciduous species. Dashed lines are theoretical predictions. Data from the GlopNet trait data set[1].

Carbon-balance optimality provides a quantitative explanation for the existence of the LES, and for its systematic variations with environmental conditions. The resulting patterns are key to understanding the biogeographical dimension of leaf economics. Unlike variations in physiological traits, these patterns are primarily determined by environmental selection among species rather than plasticity and genetic adaptation within species[22,30]. However, plants do have some plasticity for LMA, as has been shown experimentally[31]. Experimental effects on LMA include positive responses to PPFD, CO_2 and aridity and a negative response to growth temperature[31,32] – all responses in the same direction as predicted here, but generally smaller in magnitude.

Plants have been subject to profound changes in climate during their evolutionary history. Large, and sometimes rapid, changes in recent geological times have resulted in repeated large-scale reassortments of species and consequent major changes in community composition[33,34]. Current anthropogenic environmental changes are adding to the forces shaping plant

communities, in ways that remain only partly understood[16]. Optimality theory may serve not

only to solve the puzzles of global phytogeography, but also to move plant functional ecology

from empirical description to robust theoretical understanding[17].

Methods

An optimality framework for leaf economics

The optimality model of leaf longevity (LL, day) proposed by Kikuzawa[2] rests on two

assumptions. First, daily photosynthesis declines linearly from an initial maximum (denoted

by A_0, g biomass m^{-2} day^{-1}) with increasing age. Second, LL maximizes the leaf's lifetime-

average net carbon gain (accumulated daily photosynthesis averaged over LL, minus the initial

structural investment amortized over LL). Building on Kikuzawa's original work, we propose

a new, unified optimality model for both deciduous and evergreen species. For deciduous

species, we extend the period for maximizing average net carbon gain from the leaf life span

to the whole leaf life cycle (LC, day), i.e. the period from the formation of a leaf until its

replacement by a new one. LC is thus equal to LL for evergreen species, but to 365 days for

deciduous species. We further simplify the derivation by disregarding leaf maintenance

respiration, as it is relatively small, and nearly proportional to photosynthetic capacity[35]. The

daily net carbon gain (g) is then:

$g = (G - C_c \text{ LMA}) / \text{LC}$ (3)

where G (g biomass m^{-2}) is the accumulated photosynthetic carbon gain over the life cycle.

The initial structural investments are represented by the product of LMA and C_c, where LMA

is leaf mass per area (g biomass m^{-2}), and C_c (> 1, gC gC^{-1}) is the total construction cost,

including non-leaf tissues, of a unit of leaf mass. G is represented differently for deciduous

and evergreen species, since deciduous leaves can carry out photosynthesis during their whole

life span, whereas evergreen leaves are inactive during the non-growing season.

We use subscripts 'ev' and 'de' to distinguish evergreen and deciduous species. Thus for

evergreen plants:

$G_{ev} = \int_0^{LC} f A(t) \, dt$ (4)

where $A(t) = A_0 (1 - t/b)$ is the assimilation rate (g biomass m^{-2} day^{-1}) as a function of age (t,

days) and b (days) is the leaf age at which carbon assimilation would decline to zero. f is the

proportion of the year that is favourable for growth (day day^{-1}). Since $LC_{ev} = LL_{ev}$, the integral

in (4) is equal to $f A_0 [LL_{ev} - (LL_{ev})^2/2b]$, and equation (3) collapses to Kikuzawa's model[2]:

$g_{ev} = f A_0 (1 - LL_{ev}/2b) - C_c LMA/LL_{ev}$, hence (5)

$\partial g_{ev}/\partial(LL_{ev}) = -f A_0/2b + C_c LMA/(LL_{ev})^2$ (6)

Setting equation (6) to zero yields an expression for optimal LL_{ev} (days) in evergreen plants:

$LL_{ev} = \sqrt{[(2b\, C_c\, LMA) / (f A_0)]}$. (7)

For deciduous plants:

$G_{de} = \int_0^{LC} A(t)\, dt = \int_0^{LL} A(t)\, dt$ (8)

The integral in equation (8) equals $A_0 [LL_{de} - (LL_{de})^2/2b]$ and thus

$g_{de} = [A_0\, LL_{de} (1 - LL_{de}/2b) - C_c\, LMA] / LC$ (9)

In deciduous species $LC = 365$, thus

$\partial g_{de}/\partial(LL_{de}) = A_0 (1 - LL_{de}/b) / 365$ (10)

Setting equation (10) to zero yields a simple equation for optimal LL (days) in deciduous plants:

$LL_{de} = b$. (11)

We implemented the model of b proposed by Xu $et\ al.$[3] in the framework described above. Xu

$et\ al.$ demonstrated that b has a positive relationship with LMA, and a negative relationship to

the maximum capacity of carboxylation at 25°C (V_{cmax25}, µmol C m^{-2} s^{-1}), with scaling

coefficients close to 1 and −1 respectively. Therefore, b can be expressed as:

$b = u\, LMA/(k_1\, k_2\, V_{cmax25})$ (12)

Here $u \approx 768 \pm 71$ (dimensionless), estimated from a meta-analysis of data on 49 species across

temperate and tropical biomes[3]. The scaling factors are $k_1 = 30$ g biomass mol C^{-1} and $k_2 =$

0.0864 s mol C day^{-1} µmol^{-1} C.

The optimality model of leaf photosynthesis

The coordination hypothesis, supported by extensive field and experimental evidence[25,35],

states that the maximum capacity of carboxylation at growth temperature ($V_{cmax.gt}$, μmol C m⁻

² s⁻¹) acclimates to the daytime environment so that the Rubisco-limited photosynthetic rate

(A_C, μmol C m⁻² s⁻¹) tends to equality with the electron transport-limited rate (A_J, μmol C m⁻²

s⁻¹). In other words, $A_C = A_J$. This acclimation is optimal because it avoids investment in excess

photosynthetic capacity, while allowing full use of the available light[4,24]. Combined with the

standard biochemical model of photosynthesis[36], the coordination hypothesis predicts $V_{cmax.gt}$:

$A_C = V_{cmax.gt}\, m_c,$ (13)

$A_J = \varphi_0\, I_{abs}\, m,$ and   (14)

$V_{cmax.gt} = \varphi_0\, I_{abs}\, m\, /\, m_c,$ where   (15)

$m = (c_i - \Gamma^*)\, /\, (c_i + 2\Gamma^*)$ and   (16)

$m_c = (c_i - \Gamma^*)\, /\, (c_i + K).$ (17)

Here, φ_0 is the intrinsic quantum efficiency of photosynthesis (μmol C μmol⁻¹ photon), which

we assume to depend on temperature as described in ref. [37]. I_{abs} is the leaf-absorbed PPFD

(μmol photon m⁻² s⁻¹) to which $V_{cmax.gt}$ acclimates, m_c and m are the CO_2 limitation terms for

A_C and A_J respectively, c_i is the leaf-internal partial pressure of CO_2 (Pa), Γ^* is the

photorespiratory compensation point (Pa), and K is the effective Michaelis-Menten coefficient

of Rubisco (Pa). Optimal stomatal regulation, according to the least-cost hypothesis[38], yields

the following relationship of c_i to environmental variables, including vapour pressure deficit

(D, Pa) and c_a:

$c_i = (\xi\, c_a + \Gamma^*\sqrt{D})\, /\, (\xi + \sqrt{D}),$ where   (18)

$\xi = \sqrt{[\beta\, (\Gamma^* + K)\, /\, (1.6\eta^*)]}$ (19)

The dimensionless parameter β in equation (19) is the ratio of unit costs for the maintenance

of carboxylation and water transport capacities, evaluated at 25°C. The term η^* is the viscosity

of water (normalized by its value at 25°C), which declines with increasing temperature, thereby

reducing the cost of water transport and thereby favouring higher c_i. The value of c_i thus

depends on temperature via η^*, Γ^* and K, as well as on D and c_a[38]. Analysis of a global leaf

stable carbon isotope dataset[20] has provided empirical support for the separate dependencies of

the $c_i{:}c_a$ ratio on temperature, vapour pressure deficit and elevation implied by equations (18)

and (19), and an estimated value of $\beta \approx 146$ (refs [20,39]).

The temperature dependencies of Γ^*, K and V_{cmax} within normal physological ranges can be

described by the Arrhenius function[40] with different activation energies (the formulation for K

is somewhat more complex as it depends on the affinities of Rubisco for CO_2 and O_2 and their

distinct activation energies, while both Γ^* and K are influenced by atmospheric pressure: see

ref. [40] for details). V_{cmax25} is related to $V_{cmax.gt}$ by:

$$V_{cmax25} \;=\; V_{cmax.gt} / h_T \tag{20}$$

where h_T is the Arrhenius function for V_{cmax}.

We further assume that the coordination hypothesis defines the initial photosynthetic capacity

of the leaf. Thus,

$$A_0 = k_1\, k_2\, V_{cmax.gt}\, m_c \tag{21}$$

Here, k_1 and k_2 together convert the unit of $V_{cmax.gt}$ from μmol C m^{-2} s^{-1} into g biomass m^{-2}

day^{-1} to match the units of A_0.

Optimal leaf longevity in evergreen species

Substituting equations (12)–(21) into (7) yields equation (1). Substituting equation (7) into (5)

also yields an expression for an evergreen leaf's average daily net carbon gain:

$$g_{ev} = f A_0 \left\{1 - \sqrt{\left[2\, C_c / (u\, h_T\, m_c\, f)\right]}\right\}, \tag{22}$$

which is independent of either LMA or LL. In other words, any combination of LMA and LL

that satisfies the optimality criterion will yield the same daily net carbon gain.

The coordination hypothesis also relates the initial rate of carbon assimilation to the electron-

transport limited photosynthetic rate:

$$A_0 = k_1\, \varphi_0\, \Sigma I_{abs}\, m \tag{23}$$

where ΣI_{abs} is the integrated instantaneous I_{abs} through a day (mol photon m^{-2} day^{-1}). It follows

that:

$LL_{ev} = LMA \, x_{T.ev} \sqrt{(2u \, C_c / f)} / (\Sigma I_{abs} \, k_1)$, where   (24)

$x_{T.ev} = \sqrt{(h_T \, m_c)} / (\varphi_0 \, m)$ (25)

The composite term $x_{T.ev}$ includes the temperature-dependent variables. We express the
parameter C_c as a function of LL, LMA and environmental predictors by rearranging equations
(24) and (25) as follows:

$C_c = f \, (LL \, \varphi_0 \, \Sigma I_{abs} \, m \, k_1 / LMA)^2 / (2u \, h_T \, m_c)$ (26)

Based on all paired observations of LL and LMA in evergreen species in GlopNet (326
samples), we estimated a median value of $C_c = 23 \pm 5.12$ gC gC^{-1} (mean $\pm$ s.e.).

Optimality leaf longevity in deciduous species

Substituting equations (12), (15) and (20) into (11) yields equation (2). Substituting equation
(2) into (9) also yields an expression for a deciduous leaf's average daily net carbon gain:

$g_{de} = f \, A_0 \, [1/2 - C_c / (u \, h_T \, m_c)]$ (27)

As in evergreen species, g_{de} is independent of either LMA or LL. The different formulae for
g_{de} and g_{ev} potentially allow modelling of competition between the two strategies.

Following similar logic to that applied to evergreens, equation (23) is then substituted into
equation (2), leading to a prediction of LL_{de} from LMA and environmental variables:

$LL_{de} = LMA \, u \, x_{T.de} / (\Sigma I_{abs} \, k_1)$, where   (28)

$x_{T.de} = (h_T \, m_c) / (\varphi_0 \, m)$ (29)

Linearized models for optimal LL

To facilitate comparisons of data with theoretical predictions, equation (24) can be linearized:

$\ln LL_{ev} = \ln LMA - \ln \Sigma I_{abs} - 0.5 \ln f + \ln x_{T.ev} + C_{ev}$, where   (30)

$C_{ev} = 0.5 \ln (2 \, u \, C_c) - \ln k_1$ (31)

By taking the partial derivative of $\ln x_{T.ev}$ with respect to temperature, we calculated that $LL_{opt.ev}$
increases with growth temperature (T_g, K) by 2.6% K^{-1} under standard environmental

conditions. Thus, $\ln x_{T.ev}$ can be replaced by the sum of $0.026\ T_g$ and $\ln x_{T.ev}$ at the reference

condition ($\ln x_{T0.ev}$). D and z are also expected to influence LL_{opt} slightly via their effects on

optimal c_i, but these effects are too small to be detected.

Similar logic yields a linearized model of LL for deciduous species from equation (28):

$\ln LL_{de} = \ln LMA - \ln \Sigma I_{abs} + \ln x_{T.de} + C_{de}$, where   (32)

$C_{de} = \ln u - \ln k_1$.   (33)

The thermal sensitivity of LL_{de} is twice of that in evergreen species. Thus, $\ln x_{T.de}$ can be

replaced by $0.052\ T_g$ plus its reference value ($\ln x_{T0.de}$).

Distributions of evergreen LMA and LL

Different combinations of LMA and LL consistent with the optimality criterion result in equal

net carbon gain, implying the existence of a LES along which species with different LMA are

equally competitive. Nonetheless, within-site distributions of LMA and LL are known to vary

systematically with climate. To account for this variation, it is necessary first to take account

of their prior distributions in the species pool. We fitted lognormal distributions to observed

LMA and LL data independently in the full GlopNet dataset (Figure S1). Equation (24) was

then imposed as a constraint, representing environmental filtering of the species pool. This

procedure generates site-specific lognormal distributions, which serve as predictions of the

probability density functions of LMA and LL subject to the optimality criterion as illustrated

in Figure S2.

Environmental dependencies of LMA for evergreen species

Equation (30) can be re-configured as an expression for the optimal LMA of evergreen species:

$\ln LMA_{ev} = \ln \Sigma I_{abs} + 0.5 \ln f - 0.026\ T_g + \ln LL_{ev} + C_{ev}'$, where   (34)

$C_{ev}' = \ln k_1 - 0.5 \ln (2u\ C_c) - \ln x_{T0.ev}$ (35)

Optimal LMA depends on LL as well as on environmental factors. For a given LL, equation

(34) implies that LMA (all else equal) should increase in proportion to absorbed PPFD and the

square root of growing-season length, but should decrease with temperature by 2.6% K^{-1}. The

prior normal distributions of $\ln$ LMA and $\ln$ LL, together with their predicted proportional

relationship, allow LMA to be predicted from environment alone (Figure S2). Here, we

consider ΣI_{abs} as an example to make the derivation, noting that the effects of the other

environmental variables (f and T_g) can be derived following the same logic. We specify

equations (34) and (35) at two sites with different light conditions (ΣI_1 and ΣI_2) but the same

growing-season length and growth temperature. The subscripts '1' and '2' in the equations

below distinguish the two sites:

$$\ln \mathrm{LMA_{ev1}} = \ln \Sigma I_1 + 0.5 \ln f - 0.026\, T_g + \ln \mathrm{LL_{ev1}} + C_{ev}' \tag{36}$$

$$\ln \mathrm{LMA_{ev2}} = \ln \Sigma I_2 + 0.5 \ln f - 0.026\, T_g + \ln \mathrm{LL_{ev2}} + C_{ev}' \tag{37}$$

Thus,

$$\ln \mathrm{LMA_{ev1}} - \ln \mathrm{LMA_{ev2}} = \ln \Sigma I_1 - \ln \Sigma I_2 + \ln \mathrm{LL_{ev1}} - \ln \mathrm{LL_{ev2}} \tag{38}$$

The normal distribution constraint induces a line perpendicular to the optimality lines for

evergreen species (Figure S2):

$$\ln \mathrm{LL_{ev1}} = k - \ln \mathrm{LMA_{ev1}} \tag{39}$$

$$\ln \mathrm{LL_{ev2}} = k - \ln \mathrm{LMA_{ev2}} \tag{40}$$

where k is the intercept of this line. Therefore,

$$\ln \mathrm{LL_{ev1}} - \ln \mathrm{LL_{ev2}} = \ln \mathrm{LMA_{ev2}} - \ln \mathrm{LMA_{ev1}} \tag{41}$$

Combining equations (38) and (41) yields:

$$\ln \mathrm{LMA_{ev2}} - \ln \mathrm{LMA_{ev1}} = 0.5\,(\ln \Sigma I_2 - \ln \Sigma I_1) \tag{42}$$

Similarly for the growing-season length and temperature:

$$\ln \mathrm{LMA_{ev2}} - \ln \mathrm{LMA_{ev1}} = 0.25\,(\ln \Sigma f_2 - \ln \Sigma f_1) \tag{43}$$

$$\ln \mathrm{LMA_{ev2}} - \ln \mathrm{LMA_{ev1}} = -0.13\,(\ln \Sigma T_{g2} - \ln \Sigma T_{g1}) \tag{44}$$

Thus, when moving from site 1 to site 2, the predicted changes in LMA are 50%, 25% and –

13% of the changes in PPFD, growing-season length and temperature respectively.

Environmental dependencies of LMA for deciduous species

As the lifespan of deciduous leaves is constrained by growing-season length, LL_{de} should also
be proportional to f ($LL_{de} = 365 f$). Re-arranging equation (32) and imposing this additional
constraint yields the following expression for deciduous LMA as a function of environment
alone:

$$\ln LMA_{de} = \ln f + \ln \Sigma I_{abs} - 0.052 \, T_g + C_{de}', \text{ where} \qquad (45)$$

$$C_{de}' = \ln k_1 + \ln 365 - \ln u - \ln x_{T0.de} \qquad (46)$$

Equation (45) implies that deciduous LMA, unlike evergreen LMA, should increase in direct
proportion to both PPFD and growing-season length, and decline with warming four times
faster than in evergreens.

The aridity constraint on LMA

Finally, we account for the expected increase in LMA with aridity as an additional, physical
constraint on LMA for both deciduous and evergreen species. This additional constraint could
in principle have been included in equations (34) and (45) as an independent predictor.
However, the current optimality framework does not predict the magnitude of this response.
We therefore estimated the aridity effect empirically and used this additional information to
modify the predicted within-site distribution of evergreen LMA, shifting the peak to higher
values in drier sites.

Leaf trait data

We tested our quantitative predictions with the woody plants represented in two non-
overlapping datasets: the GlopNet data set[1], and a version of the China Plant Trait Database
(CPTD)[5] augmented with data compiled by Geng et al.[28] from alpine shrublands. GlopNet
contains paired LMA and LL data for 326 evergreen and 179 deciduous species at 45 sites,
distributed globally. For deciduous plants we focus here on those whose growing season is
constrained by a cold winter; thus, tropical deciduous species that shed leaves in the dry season
have been excluded from the analysis. Applying this filtering criterion, the GlopNet data set
contains data on LMA alone from 1225 evergreen and 98 deciduous species at 146 sites,
distributed from 69°S to 40°N. The CPTD provides LMA data from 419 for evergreen and 398
deciduous species at 164 sites, spanning the range from alpine and boreal to tropical

environments, and from deserts and steppes through woodlands and forests. Data in the CPTD

were obtained using a defined sampling strategy to ensure adequate representation of major

species in all strata[5].

Site climate data

The Simple Process-Led Algorithms for Simulating Habitats (SPLASH) model was used to

calculate site-specific bioclimatic variables from climatologies of monthly temperature,

precipitation and sunshine hours[41]. Defining the (thermal) growing season as the period when

mean quasi-daily temperature (interpolated from monthly data) is above 0˚C, we calculated the

ratio of growing-season length to the number of days in the year and the mean values of

temperature and PPFD during growing season as estimates of site-mean f, T_g and PPFD. To

account for the fact that the field-measured trait data reflect leaves developed at a range of

irradiances at different levels in the canopy, we applied the approach used by Wang et al.[42] to

estimate I_{abs} from the site-mean PPFD, with the help of leaf area index values extracted from

SeaWiFS data[43]. A moisture index, defined as the ratio of estimated annual actual to potential

evapotranspiration (denoted α_p), was also calculated using SPLASH. For the GlopNet data set,

climatological data were extracted from Climate Research Unit data at a grid resolution of 10

minutes[44]. For the extended CPTD, climatological data were extracted from 1 km gridded data

interpolated from 1814 meteorological stations (China Meteorological Administration,

unpublished data: 740 stations have observations from 1971 to 2000, the rest from 1981 to

1990) using a three-dimensional thin-plate spline (ANUSPLIN version 4.36; Hancock and

Hutchinson, 2006).

Statistical analysis

Ordinary Least-Squares (OLS) multiple regression was used to estimate the partial

environmental dependencies of LL and LMA from data, for comparison with independent

predictions from theory. All traits and bioclimatic variables except for temperature were log-

transformed to facilitate comparisons with theoretical predictions, and to meet the assumptions

of the OLS model. For consistency with our definition of the growing season we excluded

deciduous species from sites with year-round temperatures > 0˚C. These are presumed to be

drought-deciduous. The effect of this exclusion could only be minor, however, as it merely

reduces the sample size for deciduous LMA to 292 in GlopNet and 621 in the CPTD. Climatic

controls of the relationship between LL and LMA were tested using the GlopNet paired LL

and LMA data for evergreens. The constraint of growing-season length on LL in deciduous

species was also tested using GlopNet data. The parameter C_c in equation (14) was estimated

as the median of individually estimated values. Given the larger latitudinal range covered,

GlopNet LMA data were used to investigate the opposing latitudinal trends in evergreen and

deciduous species. Climatic controls on LMA were tested using both GlopNet and CPTD data,

but CPTD-based results are given in the main text because of its larger sample size, especially

for deciduous species. Stepwise regressions, with or without evergreen LL as a predictor, for

GlopNet LMA data allowed us not only to test the explanatory power of LL alone on LMA,

but also to test its indirect environmental effects on LMA when LL was excluded. For

individual predictors that proved non-significant (probably due to the correlations between the

bioclimatic variables), we further tested their effects on the residuals from the main model.

The prior distributions of LL and LMA were derived from GlopNet. We also used GlopNet to

test the predicted within-site distributions of LMA. Six sites with adequate sample size were

selected for this test, including two sites each from evergreen-dominant vegetation in tropical

rainforests, temperate forests and woodlands (for details see Table S3). As shown in Figure S2,

the prior distribution is constrained by the site-specific optimal relationship of ln LL to ln LMA

with the slope set to unity and the intercept being the sum of the estimated value of C and the

climatic terms in equation (14). The posterior distributions of ln LMA were generated after

further imposing the aridity effect. As this effect is not theoretically quantified we used the

best-match prediction obtained from regression of ln LMA data against climatic variables in

the CPTD (Table S1).

Acknowledgements

This research was supported by the National Natural Science Foundation of China (no.

32022052, 31971495) and National Key R&D Program of China (no. 2018YFA0605400)

grants to HW. ICP acknowledges support from the ERC-funded project REALM (Re-inventing

Ecosystem And Land-surface Models, grant number 787203) and a High End Foreign Expert

award at Tsinghua University (GDW20191100161). This work is a contribution to the Imperial

College initiative on Grand Challenges in Ecosystems and the Environment (ICP). IJW and

ICP acknowledge support from the Australian Research Council (DP170103410).

Author Contributions.

HW and ICP developed the quantitative theory, building on previous work by KK and XTX.

XTX also contributed to theory development. HW carried out the analyses and constructed the

display items using datasets compiled by HW and IJW. ICP, IJW and SCQ contributed to the

interpretation of results. HW and ICP wrote the first draft and IJW, KK, XTX, SCQ and NCS

contributed to successive drafts.

References

1   Wright, I. J. *et al.* The worldwide leaf economics spectrum. *Nature* **428**, 821-827 (2004).

2   Kikuzawa, K. A cost-benefit analysis of leaf habit and leaf longevity of trees and their
geographical pattern. *The American Naturalist* **138**, 1250-1263 (1991).

3   Xu, X. *et al.* Variations of leaf longevity in tropical moist forests predicted by a trait-
driven carbon optimality model. *Ecology letters* **20**, 1097-1106 (2017).

4   Haxeltine, A. & Prentice, I. C. A general model for the light-use efficiency of primary
production. *Functional Ecology* **10**, 551-561, doi:10.2307/2390165 (1996).

5   Wang, H. *et al.* The China Plant Trait Database: toward a comprehensive regional
compilation of functional traits for land plants. *Ecology* **99**, 500-500,
doi:10.1002/ecy.2091 (2018).

6   Wright, I. J. *et al.* Modulation of leaf economic traits and trait relationships by climate.
Global Ecology and Biogeography **14**, 411-421, doi:10.1111/j.1466-
822x.2005.00172.x (2005).

7   Díaz, S. *et al.* The global spectrum of plant form and function. *Nature* **529**, 167 (2016).

8   Wright, I. J., Westoby, M. & Reich, P. B. Convergence towards higher leaf mass per
area in dry and nutrient-poor habitats has different consequences for leaf life span.
Journal of ecology **90**, 534-543 (2002).

9   Falster, D. S. *et al.* Lifetime return on investment increases with leaf lifespan among
10 Australian woodland species. *New Phytologist* **193**, 409-419 (2012).

10   van Ommen Kloeke, A., Douma, J., Ordonez, J. C., Reich, P. B. & Van Bodegom, P.
Global quantification of contrasting leaf life span strategies for deciduous and
evergreen species in response to environmental conditions. *Global Ecology and*
Biogeography **21**, 224-235 (2012).

11   Osnas, J. L. D. *et al.* Divergent drivers of leaf trait variation within species, among
species, and among functional groups. *Proceedings of the National Academy of*
Sciences **115**, 5480-5485, doi:10.1073/pnas.1803989115 % (2018).

12   Kikuzawa, K., Onoda, Y., Wright, I. J. & Reich, P. B. Mechanisms underlying global
temperature-related patterns in leaf longevity. *Global Ecology and Biogeography* **22**,
982-993 (2013).

13   Reich, P. B., Rich, R. L., Lu, X., Wang, Y.-P. & Oleksyn, J. Biogeographic variation
in evergreen conifer needle longevity and impacts on boreal forest carbon cycle
projections. **111**, 13703-13708, doi:10.1073/pnas.1216054110 %J Proceedings of the
National Academy of Sciences (2014).

14   Cui, E. *et al.* Vegetation functional properties determine uncertainty of simulated
ecosystem productivity: A traceability analysis in the East Asian monsoon region.
Global Biogeochemical Cycles **33**, 668-689 (2019).

15   Carvalhais, N. *et al.* Global covariation of carbon turnover times with climate in
terrestrial ecosystems. *Nature* **514**, 213-217, doi:10.1038/nature13731 (2014).

16 Monroe, J. G. *et al.* Ecoevolutionary Dynamics of Carbon Cycling in the Anthropocene.
Trends   in   Ecology   &   Evolution   33,   213-225,
doi:https://doi.org/10.1016/j.tree.2017.12.006 (2018).
17 Franklin, O. *et al.* Organizing principles for vegetation dynamics. *Nature plants* **6**, 444-
453 (2020).
18 Kattge, J. *et al.* TRY plant trait database – enhanced coverage and open access. *Global*
Change Biology **26**, 119-188, doi:10.1111/gcb.14904 (2020).
19 Detto, M. & Xu, X. Optimal leaf life strategies determine Vc,max dynamic during
ontogeny. *New Phytologist* **228**, 361-375, doi:https://doi.org/10.1111/nph.16712
(2020).
20 Wang, H. *et al.* Towards a universal model for carbon dioxide uptake by plants. *Nature*
plants **3**, 734-741 (2017).
21 Meng, T. T. *et al.* Responses of leaf traits to climatic gradients: adaptive variation
versus compositional shifts. *Biogeosciences* **12**, 5339-5352, doi:10.5194/bg-12-5339-
2015 (2015).
22 Dong, N. *et al.* Components of leaf-trait variation along environmental gradients. *New*
Phytologist **n/a**, doi:10.1111/nph.16558 (2020).
23 Westoby, M., Falster, D. S., Moles, A. T., Vesk, P. A. & Wright, I. J. Plant ecological
strategies: some leading dimensions of variation between species. *Annual review of*
ecology and systematics **33**, 125-159 (2002).
24 Chen, J.-L., Reynolds, J. F., Harley, P. C. & Tenhunen, J. D. Coordination theory of
leaf nitrogen distribution in a canopy. *Oecologia* **93**, 63-69 (1993).
25 Smith, N. G. *et al.* Global photosynthetic capacity is optimized to the environment.
Ecology letters **22**, 506-517, doi:10.1111/ele.13210 (2019).
26 Wright, I. J. & Westoby, M. Leaves at low versus high rainfall: coordination of
structure,   lifespan   and   physiology.   *New   Phytologist*   **155**,   403-416,
doi:https://doi.org/10.1046/j.1469-8137.2002.00479.x (2002).
27 Leigh, A. *et al.* Do thick leaves avoid thermal damage in critically low wind speeds?
New Phytologist **194**, 477-487, doi:10.1111/j.1469-8137.2012.04058.x (2012).
28 Geng, Y. *et al.* Linking above- and belowground traits to soil and climate variables: an
integrated database on China's grassland species. *Ecology* **98**, 1471-1471,
doi:10.1002/ecy.1780 (2017).
29 Ainsworth, E. A. & Long, S. P. What have we learned from 15 years of free-air CO_2
enrichment (FACE)? A meta-analytic review of the responses of photosynthesis,
canopy properties and plant production to rising CO_2. *New Phytologist* **165**, 351-372
(2005).
30 Donovan, L. A., Maherali, H., Caruso, C. M., Huber, H. & de Kroon, H. The evolution
of the worldwide leaf economics spectrum. *Trends in Ecology & Evolution* **26**, 88-95,
doi:https://doi.org/10.1016/j.tree.2010.11.011 (2011).
31 Poorter, H., Niinemets, Ü., Poorter, L., Wright, I. J. & Villar, R. Causes and
consequences of variation in leaf mass per area (LMA): a meta-analysis. *New*
Phytologist **182**, 565-588 (2009).
32 Cui, E., Weng, E., Yan, E. & Xia, J. Robust leaf trait relationships across species under
global environmental changes. *Nature communications* **11**, 2999, doi:10.1038/s41467-
020-16839-9 (2020).
33 Harrison, S. P. & Sanchez Goñi, M. F. Global patterns of vegetation response to
millennial-scale variability and rapid climate change during the last glacial period.
Quaternary Science Reviews **29**, 2957-2980 (2010).

Soh, W. K. *et al.* Palaeo leaf economics reveal a shift in ecosystem function associated with the end-Triassic mass extinction event. *Nature plants* **3**, 17104, doi:10.1038/nplants.2017.104 (2017).

Maire, V. *et al.* The coordination of leaf photosynthesis links C and N fluxes in C_3 plant species. *PloS one* **7**, e38345 (2012).

Farquhar, G. D., von Caemmerer, S. v. & Berry, J. A biochemical model of photosynthetic CO_2 assimilation in leaves of C_3 species. *Planta* **149**, 78-90 (1980).

Bernacchi, C. J., Pimentel, C. & Long, S. P. In vivo temperature response functions of parameters required to model RuBP-limited photosynthesis. *Plant, Cell & Environment* **26**, 1419-1430, doi:https://doi.org/10.1046/j.0016-8025.2003.01050.x (2003).

Prentice, I. C., Dong, N., Gleason, S. M., Maire, V. & Wright, I. J. Balancing the costs of carbon gain and water transport: testing a new theoretical framework for plant functional ecology. *Ecology letters* **17**, 82-91 (2014).

Stocker, B. D. *et al.* P-model v1.0: an optimality-based light use efficiency model for simulating ecosystem gross primary production. *Geosci. Model Dev.* **13**, 1545-1581, doi:10.5194/gmd-13-1545-2020 (2020).

Bernacchi, C., Singsaas, E., Pimentel, C., Portis Jr, A. & Long, S. Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell & Environment* **24**, 253-259 (2001).

Davis, T. W. *et al.* Simple process-led algorithms for simulating habitats (SPLASH v. 1.0): robust indices of radiation, evapotranspiration and plant-available moisture. *Geoscientific Model Development* **10**, 689 (2017).

Wang, H. *et al.* Acclimation of leaf respiration consistent with optimal photosynthetic capacity. **26**, 2573-2583, doi:10.1111/gcb.14980 (2020).

Gobron, N. *et al.* Monitoring the photosynthetic activity of vegetation from remote sensing data. *Advances in Space Research* **38**, 2196-2202, doi:https://doi.org/10.1016/j.asr.2003.07.079 (2006).

New, M., Lister, D., Hulme, M. & Makin, I. A high-resolution data set of surface climate over global land areas. *Climate research* **21**, 1-25 (2002).