Molecular and cellular insight into *Escherichia coli* SslE and its role during biofilm maturation

Paula M. Corsini[1,2†], Sunjun Wang[1†], Saima Rehman[1†], Katherine Fenn[2†], Amin Sagar[3], Slobodan Sirovica[4], Leanne Cleaver[1], Charlotte J. C. Edwards-Gayle[5], Giulia Mastroianni, Ben Dorgan[1], Lee M. Sewell[1], Steven Lynham[6], Dinu Iuga[7], W. Trent Franks[7], James Jarvis[8], Guy H. Carpenter[1], Michael. A. Curtis[1], Pau Bernadó[3], Vidya C. Darbari[2*] & James A. Garnett[1,2*]

[1]*Centre for Host-Microbiome Interactions, Faculty of Dental, Oral & Craniofacial Sciences, King's College London, London, UK*

[2]*School of Biological and Chemical Sciences, Queen Mary University of London, London, UK*

[3]*Centre de Biologie Structurale. INSERM, CNRS, Université de Montpellier, France*

[4]*Centre for Oral Bioengineering, Institute of Dentistry, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, London, UK*

[5]*Diamond Light Source Ltd., Diamond House, Harwell Science and Innovation Campus, Didcot, Oxfordshire, UK*

[6]*Proteomics Facility, Centre of Excellence for Mass Spectrometry, King's College London, London, UK*

[7]*Department of Physics, University of Warwick, Coventry, UK*

[8]*Randall Division of Cell and Molecular Biophysics and Centre for Biomolecular Spectroscopy, King's College London, London, UK*

[†]*equal contributions*

Correspondence to:

James Garnett, Centre for Host Microbiome Interactions, Dental institute, King's College London, London, UK. email: james.garnett@kcl.ac.uk

Vidya C. Darbari, Department of Biochemistry, School of Biological and Chemical Sciences, Queen Mary University of London, UK. email: v.darbari@qmul.ac.uk

Abstract

Escherichia coli **is a Gram-negative bacterium that colonizes the human intestine and virulent strains can cause severe diarrhoeal and extraintestinal diseases. The protein SslE is secreted by a range of pathogenic and some commensal** *E. coli* **strains. It can degrade mucins in the intestine, promotes biofilm maturation and it is a major determinant of infection in virulent strains, although how it carries out these functions is not well understood. Here we examine SslE from the** *E. coli* **Waksman and H10407 strains and reveal that it has a novel and dynamic structure. In response to acidification within mature biofilms we show how SslE forms a unique functional aggregate that interacts with cellulose and regulates the distribution of exopolysaccharides in macrocolony biofilms. Our data indicates that the spatial organization of SslE polymers and local pH are critical for biofilm maturation and SslE is a key factor that drives persistence of SslE-secreting bacteria during acidic stress.**

Introduction

Escherichia coli is a primary colonizer of the lower intestinal tract of humans and other warm-blooded animals. While many strains are considered beneficial to the host and help to maintain a healthy immune system, virulent strains are the cause of severe diarrhoeal diseases, including haemorrhagic colitis, and extraintestinal diseases such as neonatal meningitis, urinary tract infections, sepsis and pneumonia[1]. A wide range of pathogenic *E. coli* strains, and some commensals, use a *Vibrio*-like type II secretion system (T2SS) to translocate the protein SslE across their outer membrane and onto their extracellular surface[2, 3, 4, 5, 6, 7]. These include the Waksman (W), enterotoxigenic (ETEC), enterohemorrhagic (EHEC), enteropathogenic (EPEC), enteroaggregative (EAEC), enteroinvasive (EIEC) and neonatal meningitis *E. coli* (NMEC) strains. SslE is required for full virulence in a rabbit model of EPEC infection[4] and as a surface exposed antigen, SslE has shown great promise as a broadly protective vaccine candidate against a wide range of *E. coli* pathotypes[3, 6, 8, 9].

SslE interacts with mucosal membranes in the host intestine where it can degrade mucins[6, 10, 11, 12], a

family of heavily *O*-linked glycosylated proteins and the primary constituents of mucus[13]. This provides

nutrients during bacterial growth but also enables these *E. coli* strains to penetrate the gut mucosa to

access host cells for efficient colonisation and targeting of toxins/effectors. Furthermore, SslE is also

important for mediating maturation of EPEC biofilms[4]; microbial aggregations encased within a self-

produced extracellular matrix comprised of exopolysaccharides, adhesive proteins and nucleic acids[14].

When *E. coli* is released into the environment through faeces or wastewater effluent it can survive for

long periods within complex biofilm communities[15, 16], and these are fundamental for both the

environmental ecology of *E. coli* but also for successful colonisation of the intestinal tract[17]. However,

the specific molecular mechanisms that SslE uses to promote ecology and disease are not well

understood.

SslE is a ~165 kDa lipoprotein composed of an N-terminal periplasmic localization sequence and

lipobox motif, an unstructured ~5 kDa region, a ~110 kDa region with no significant primary sequence

homology to any other known protein, and a ~50 kDa M60-like aminopeptidase domain at its C-

terminus[18] (**Fig. 1a**). M60-like domains are metalloproteases that contain a zinc binding HExxH motif

and an additional conserved catalytic glutamate residue, which cleaves the peptide backbone of mucin-

like substrates. These and other related enzymes have been identified in both prokaryotic and eukaryotic

microbes that interact with host mucosal membranes[18] and structures of proteoglycan complexes

suggest that interactions with both the mucin peptide and *O*-linked carbohydrate sidechains are

important for specificity[19, 20]. The peptidase activity of the SslE M60 domain is effective for the

degradation of major mucins of the intestine (e.g. MUC2, MUC3, MUC5AC)[6, 10, 11, 12], however, very

little is known as to how SslE interacts with these mucins or the function of the remaining SslE

sequence.

Using electron microscopy (EM), small angle X-ray scattering (SAXS), nuclear magnetic resonance

(NMR) spectroscopy and biochemical analyses we show that SslE from the *E. coli* W strain is formed

of three defined regions (NT1, NT2 and NT3-M60), and is dynamic in solution. We also demonstrate

that SslE undergoes conformational changes under acidic conditions, and this leads to the formation of

higher order structures, which we propose is mediated by the NT2 and NT3 regions and is decorated

with NT1 domains. We directly observe acidification within mature *E. coli* W biofilms and determine

that SslE fibres can bind cellulose, a major exopolysaccharide of many *E. coli* biofilms. Furthermore,

we provide evidence that SslE regulates the localization of exopolysaccharides during maturation of

both *E. coli* W and ETEC strain H10407 biofilms and this is likely a general mechanism in all SslE-

secreting bacteria.

Results

SslE forms aggregates with amyloid-like properties. Functional amyloids are important

proteinaceous structures in biofilms[21] and analysis of the SslE primary sequence had previously

highlighted several regions across the protein that may form amyloid-like structures[22]. Furthermore,

environmental pH plays a major role initiating the formation of amyloid-like structures in

Staphylococcus aureus, *Enterococcus faecalis* and *Streptococcus mutans* biofilms[23, 24, 25], and we

speculated that pH may also stimulate similar structural changes in SslE. SslE from *E. coli* W, minus

the N-terminal lipidation motif and adjacent disordered region (rSslE; residues 67 to 1497; UniProt

ID E0IW31) (**Fig. 1a**), was therefore produced in *E. coli* K-12 and purified by nickel-affinity and size

exclusion chromatography. We then incubated rSslE across the pH range 4.0 to 7.0 and observed a ring

of protein deposited on the walls of the tube at pH ≤ 4.4, which resisted solubilization up to pH 6.2 (**Fig.**

1b). Mass spectrometry analysis of the ring identified peptides that spanned the complete sequence of

rSslE and indicated that the aggregate contained intact protein, rather than a degradation product

(**Supplementary Fig. 1**). We next used the amyloid diagnostic dye Thioflavin-T (ThT) to assess the

formation of rSslE aggregates in solution over time. We observed large increases in ThT fluorescence

emission again across the pH range 4.0 to 4.4, but no change was detected when rSslE was incubated

above pH 4.4 (**Fig. 1c**). Likewise, when rSslE was boiled and treated with sodium dodecyl sulphate

(SDS) and then analysed by immunoblotting, stable high molecular weight species were observed over

the same pH range (**Fig. 1d**).

A main feature of amyloids is the presence of a cross-β-sheet structure[26] and so we first analysed the

secondary structure of soluble rSslE using far-UV circular dichroism (CD) spectroscopy. This

suggested that at pH 7.4 rSslE is composed of approximately 30% α-helix, 12% β-sheet and 45% coil,

while under acidic conditions and prior to aggregation, there is an ~8% loss of helical secondary

structure with a ~5% increase in β-sheet structure (**Fig. 1e; Supplementary Table 1**). We next used

attenuated total reflectance (ATR) Fourier transform mid-infrared (FT-MIR) spectroscopy to probe the

secondary structure composition of soluble rSslE at pH 7.4 and rSslE fibres at pH 4.4. Examination of

the amide I absorption band showed peaks at approximately 1656 cm^{-1} and 1624 cm^{-1}, respectively

(**Supplementary Fig. 2**). The shift in the amide I absorption band to a lower frequency indicated that a

larger and more rigid structure had assembled under acid conditions. Furthermore, the presence of both

a high-intensity absorption band at ~1624 cm^{-1} and a low-intensity band at ~1695 cm^{-1} in the pH 4.4

sample was indicative of an increase in anti-parallel β-sheet structure in rSslE aggregates[27] and is

consistent with previous FT-MIR based characterisations of other anti-parallel β-sheet amyloid fibres[28]

(**Fig. 1f**).

Biofilm maturation supports polymerisation of SslE. Congo red is a dye which binds extracellular

polymeric substances (EPS), including amyloids, and we used this to assess the role of SslE during the

formation of macrocolony biofilms. We created an isogenic Δ*sslE* knockout mutant in *E. coli* W and

ETEC strain H10407 and compared these to their parental wild-type strains. When these mutants were

cultured on Congo red agar plates for 24 hrs, we observed no obvious differences, however, after 96

hrs the mutants exhibited an altered morphology with the dye retained within the centre of the colony

(**Fig. 2a**). Complementation of the Δ*sslE* mutant in *E. coli* W with a plasmid containing intact *sslE*

(Δ*sslE::sslE*) was able to restore bacterial secretion of SslE and recover macrocolony morphology

similar to that of the parental wild-type (**Fig. 2a**; **Supplementary Fig. 3**). We also generated a plasmid

expressing SslE with a truncated C-terminus (Δ*sslE::sslEΔM60*; residues 1 to 1000) to examine the role

of the M60-domain during colony formation, and this again restored SslE secretion and wild-type

colony morphology (**Fig. 2a**; **Supplementary Fig. 3**). This led us to speculate that SslE experiences

extracellular pH values <5.0 during the development of biofilms and this results in SslE being deposited

as a functional aggregate.

To test this, we first grew *E. coli* W biofilms using a microfluidic system under continuous flow and

assessed their overall morphology using confocal laser scanning microscopy (CLSM). While biofilms

grown for 48 hrs produced a relatively homogenous lawn of bacterial growth, after 96 hrs there were

clear signs of maturation, with a significant increase in biofilm mass, structural heterogeneity and the

presence of water channels (**Fig. 2b,c**). We then examined the *E. coli* W Δ*sslE* mutant under these

conditions, but in line with previous reports in EPEC strain E2348/69[4], it was unable to develop

structures beyond microcolonies (**Fig. 2c,d**). Furthermore, complementation of the mutant with either

intact *sslE* or *sslEΔM60* was again able to restore wild-type biofilm morphology. Together with our

previous observation, this indicated that the C-terminal M60-domain is not required for translocation

of SslE through its T2SS and it is dispensable for biofilm development, at least under these conditions.

We then examined pH distribution across *E. coli* W biofilms using CLSM coupled with the cell-

impermeant fluorescent ratiometric probe seminaphthorhodafluor-4F 5-(and-6) carboxylic acid (C-

SNARF-4). After growing *E. coli* W for 24 hrs under continuous flow, we observed pH values between

6.0 and 6.3 across biofilms (**Supplementary Fig. 4**), however, after 96 hrs, lower pH values were also

recorded (**Fig. 2e**). Although across the majority of biofilms examined, we detected pH values between

approximately 5.0 and 6.0, clearly defined microenvironments were also observed with pH values that

ranged from 4.2 and 4.8.

We observed that when *E. coli* W was grown overnight in liquid culture at pH 5.0, the wild-type strain

and both Δ*sslE::sslE* and Δ*sslE::sslEΔM60* complementation strains formed a clear ring on the wall of

the tube that retained congo red dye (**Fig. 3a**). This was absent in the Δ*sslE* mutant or when grown at

pH 7.0 and mass spectrometry analysis of the wild-type *E. coli* W ring again identified SslE peptides

from across its sequence (**Supplementary Fig. 1**). Examination of wild-type *E. coli* W at pH 5.0 by

immunoelectron microscopy revealed gold-labelled anti-rSslE antibodies associated with the bacterial outer membrane, but also associated with fibrous material in the extracellular milieu and emanating away from the bacterial surface; which was not observed in the Δ*sslE* mutant (**Fig. 3b,c,d**). Inspection of the *E. coli* W Δ*sslE::sslE* complemented strain under the same conditions produced less clustering of anti-rSslE antibodies, however, a wild-type phenotype was observed when cells were incubated at pH 4.0 prior to fixation and staining (**Fig. 3e,f**). This indicated that SslE can form aggregates upon secretion from *E. coli* and so we then used immunofluorescence to assess how SslE is distributed within established wild-type *E. coli* W and H10407 biofilms (**Fig. 4**). Here we observed a heterogenous distribution and clustering of SslE across biofilms formed by both strains, with SslE co-localised with bacteria, likely through association with the bacterial surface, but also clearly incorporated within the extracellular matrix.

SslE aggregates bind cellulose. We speculated that during the maturation of biofilms, SslE aggregates interact with other components of the biofilm matrix and may contribute to the structural integrity of these communities. We therefore re-examined the morphology of wild-type and Δ*sslE* mutant *E. coli* W and H10407 macrocolonies using the carbohydrate stain calcofluor white, which binds β1-3 and β1-4 polysaccharides including cellulose (**Fig. 5a**). As with Congo red staining, calcofluor white extended to the edge of the colony in wild-type strains but was retained within the centre in the Δ*sslE* mutants. We then produced rSslE aggregates at pH 4.0 and isolated them in citrate-phosphate buffer at pH 6.0, a condition in which fibres would resist solubilization (**Fig. 1b**). As controls, we used monomeric rSslE, and the C-terminal domain (CTD) of the *Porphyromonas gingivalis* gingipain protein RgpB, both of which were not expected to bind cellulose or form polymers at this pH (**Supplementary Fig. 5**). We then assayed binding of rSslE fibres to immobilised cellulose discs and observed a significant interaction, but as expected this was not detected with monomeric rSslE or RgpB-CTD (**Fig. 5b**).

Overall structural features of SslE. We next initiated structural studies of SslE. Small angle X-ray scattering (SAXS) coupled with size exclusion chromatography (SEC-SAXS) was first used to get shape information for rSslE at pH 7.4. Guinier analysis provided a radius of gyration (R_g) of 4.03 nm

and examination of the distance distribution function ($P(r)$) suggested a maximum particle dimension

(D_{max}) of 14.1 nm and R_g of 4.07 nm (**Supplementary Fig. 6; Supplementary Table 2**). Calculation

of the particle molecular mass (151 kDa) was consistent with a monomer in solution (theoretical mass

160 kDa). *Ab initio* dummy residue reconstruction generated 20 reproducible models with an average

normalized spatial discrepancy (NSD) score between reconstructions of 0.56 and a χ^2 fit between

calculated and experimental solution scattering of 1.2 (**Fig. 6a,b**). The averaged low-resolution bead

model suggested that rSslE has a slightly elongated structure, however, examination of the Kratky plot

indicated that rSslE is a dynamic particle (**Supplementary Fig. 6**) and no obvious domain features

could be assigned.

Negative-stain TEM data was next collected for rSslE at pH 7.4 and we performed single particle

analysis to generate an initial low-resolution structure, with overall dimensions of approximately

8×8×11.5 nm (**Fig. 6c; Supplementary Fig. 7**). Here, three well defined regions could be identified:

two adjacent globular domains (regions 1 and 2) with approximate diameters of 4 nm and 4.5 nm,

respectively, and a torus shaped structure (region 3) with a ~15 Å wide central channel and approximate

overall dimensions of 4.5×6.5×9 nm (**Fig. 6d**). Superposition of the SAXS bead model with the TEM

envelope supported SslE being dynamic in solution, with an apparent reduction of bead volume in all

three regions of the SAXS model. Docking analysis of a homology model of the M60 domain, based

on the IMPa M60 metalloprotease from *Pseudomonas aeruginosa* (PDB ID code 5kdv)[19], indicated that

the C-terminus of SslE is located at the base of region 3 (**Fig. 6e**). This suggested that the remaining

density in region 3 is composed of the SslE sequence directly N-terminal to the M60 domain and these

two regions pack against one another to form a stable structure around an inter-domain channel.

Furthermore, from this model the M60 HExxH active site motif faces into the channel and this could

be an important feature for its mucolytic activity. The location of the M60 domain in the TEM map was

also supported by previous reports where it was not possible to purify a folded recombinant SslE M60

fragment due to sample instability[22].

Secondary structure analysis[29] of the SslE sequence indicated two potential inter-domain boundaries

within the N-terminal region and we predicted that these represented regions 1 and 2 located at the

extreme N-terminus (NT) of SslE. We therefore renamed these as the NT1 (residues 67-211) and NT2

(residues 230-425) domains, respectively, and renamed the remaining sequence as NT3 (residues 426-

1000) (**Fig. 6f**). The NT1 and NT2 domains, along with the NT1-NT2 region (residues 67-425) and

NT3-M60 core (region 3; residues 426-1497) were then produced in *E. coli* K-12 and purified by nickel-

affinity and size exclusion chromatography. Examination of these constructs using analytical gel

filtration and nuclear magnetic resonance (NMR) spectroscopy showed that they were well folded

(**Supplementary Fig. 8**), and supported NT1, NT2 and NT3-M60 being well-defined structural

boundaries within SslE. When we examined intact rSslE by NMR, a ^{1}H^{15}N transverse relaxation

optimized spectroscopy (TROSY) HSQC spectrum yielded only ~250 strong resonances with the

remaining ~1200 peaks being either very weak or completely absent (**Fig. 6g**). Furthermore, NMR

relaxation measurements of these intense peaks provided an estimated correlation time consistent with

an ~80 kDa domain ($\tau_c \sim 35$ ns at 37°C). Comparison of rSslE with ^{1}H^{15}N-HSQC spectra from SslE

subdomains also clearly showed that the rSslE NMR spectrum is dominated by the NT1 domain but

with minor contributions from the NT2 domain (**Fig. 6g; Supplementary Fig. 9**). This further

supported this region having significant independent motion with respect to the NT3-M60 core and we

considered whether the NT1-NT2 region alone could be responsible for mediating SslE aggregation.

We therefore repeated our ring assay with SslE NT1-NT2 and the NT3-M60 constructs incubated across

the pH range 4.0 to 7.0. This resulted in significantly less protein being deposited on the walls of the

tubes and for NT3-M60 this was over a wider pH range (**Supplementary Fig. 10**). This implied that

both the NT1-NT2 and the NT3 regions are necessary for correct polymerisation of intact SslE.

Kinetics of SslE aggregation. We took advantage of SEC-SAXS to probe the global shape of rSslE at

pH 4.4 prior to fibre formation and both Guinier analysis (R_g 3.92 nm) and examination of the distance

distribution function (D_{max} 13.7 nm) produced very similar values to rSslE at pH 7.4 (**Supplementary**

Fig. 11; Supplementary Table 2). However, the presence of better-defined features in the SAXS

profile at acidic pH and the differences at large q-values in the normalized Kratky plots were indicative

of rSslE becoming more rigid under acidic conditions (**Fig. 7a**; **Supplementary Fig. 12**). *Ab*

initio dummy residue reconstruction produced 20 models with an average NSD score of 0.45 and a χ^2 fit

between calculated and experimental solution scattering of 1.0 (**Fig. 7b**; **Supplementary Fig. 13**).

Comparison of the averaged bead models indicated an increase in bead volume within the NT1 and

NT2 domains and adjacent NT3 region at pH 4.4, while little change was observed in the NT3-M60

core, which supported an increase in rigidity under acidic conditions that could trigger aggregation.

We then exploited the additive nature of SAXS along with its ability to examine structures over a large

size range to study the kinetics of fibre formation by rSslE. We acquired SAXS data at pH 4.4 during

the course of aggregation for 11 hrs and the resulting 23 profiles displayed a systematic upwards

intensity increase, indicating the presence of large aggregates in solution (**Fig. 7c**). The appearance of

these large species occurred relatively quickly as the upwards intensity increase was already observed

after 1 hr and the increase of aggregated species was concomitant to the loss of the SAXS features

linked to the monomeric species. Principal Component Analysis (PCA) indicated that the data set could

be described as a two-component system (**Supplementary Fig. 14**) and we decomposed the time-

dependent data set along with the SAXS profile of the monomer, obtained using SEC-SAXS, with

COSMiCS. COSMiCS uses a chemometric approach to decompose SAXS data sets in a model-free

manner to derive the pure SAXS curves of the co-existing species and their relative populations, which

we used here to report on the fibrillation kinetics[30]. The COSMiCS decomposition, assuming the co-

existence of two components, was able to adequately fit all the SAXS profiles with an average $<\chi^2>$ of

1.5, although some deviations from the perfect fit were observed for the SAXS curves measured in the

first 3 hrs (**Fig. 7c**). This observation suggested the presence of a small population of a third species in

the initial steps of fibrillation but this putative third species could not be captured when considering a

three-species decomposition with COSMiCS.

The extracted profile of the smallest species, and the subsequently derived R_g (3.93 nm) and D_{max} (13.3

nm) values, were almost identical to the SEC-SAXS profile for rSslE at pH 4.4 (**Fig. 7d,e**;

Supplementary Table 2), and this indicated that SslE is in a monomeric form at the beginning of the

fibrillation process. The extracted SAXS profile of the second component displayed the typical features

of a large particle and $P(r)$ analysis indicated it had an R_g of ~51.1 nm and D_{max} of ~140 nm (**Fig. 7d,e;**

Supplementary Table 2). Fractal fit analysis[31, 32] of the decomposed SAXS profiles also indicated that

rSslE aggregates into a structure that is fractal in nature (**Supplementary Fig. 15**). The radius of the

primary particle was determined to be ~3.5 nm while the mass fractal and surface fractal dimensions

were determined to be 2.9 and 3.0, respectively. Examination of the relative populations of the two

species derived from the COSMiCS decomposition over time showed that after 30 min from initiating

the measurements, the aggregated species already represented 50% of the molar fraction (**Fig. 7f**). From

this time point, this population continued to grow and reached a plateau after ~10 hrs.

Structural model of SslE fibres. Negative-stain TEM was used to visualize purified rSslE fibres

revealing two common morphologies (**Fig. 8a**). The first form appeared as short single, flexible fibres

with a core structure measuring ~4.5 nm wide by ~110 nm in length and decorated with globular

structures ~4.5 nm in diameter (**Fig. 8b,c**). The second morphology again resembled a fibrous material

but measured between ~20 to 40 nm in width by 200 to 300 nm in length (**Fig. 8d**) and appeared to be

an aggregation of the smaller fibres. Further analysis of rSslE aggregates in solution by real-time multi-

angle light scattering (RT-MALS) supported these observations (**Supplementary Fig. 16**). We

measured a smaller species with an average molecular mass of 5.4×10^3 kDa ($\pm 0.4 \times 10^3$) and a larger

more polydisperse species with an average molecular mass of 12×10^3 kDa ($\pm 0.8 \times 10^3$). In addition, we

measured an average R_g of 137.5 nm (± 1.7) and 139.8 nm (± 1.5) for these species, respectively, which

is in line with R_g measurements using COSMiCS. Examination of rSslE fibres using solid state NMR

(ssNMR) showed individual narrow signals (**Fig. 8e,f**), indicative of an ordered structure, and we could

tentatively assign approximately 40 ordered residues to specific residue types. Although we could not

assign these resonances to a specific sequence position in SslE, the pattern suggested that these residues

were localised to more than one site in SslE.

As the SAXS fibrillation data at the 10 hr time point had shown depletion of monomeric rSslE but also

had reduced fractal properties compared with later time points (**Fig. 7f**), we discarded the low q region

and used this data to carry out analysis of the fibre asymmetric unit. Geometrical shape analysis

suggested that rSslE fibres form a cylinder with radius 3.2 nm and height 16.5 nm, and cross-sectional

analysis indicated a repetitive unit with a cross-sectional R_g (R_c) of 4.78 nm. The cross-sectional $P(r)_C$

gave a cross-sectional D_{max} of 14.0 nm. *Ab initio* dummy residue reconstruction produced a final model

with a χ^2 fit between calculated and experimental solution scattering of 1.6 (**Fig. 8g; Supplementary**

Fig. 17). The width of the repeating unit varied between ~3.5 and ~5.5 nm, which is consistent with our

measurements of single fibres by TEM. As we could not identify peripheral globular domains in the

fibre bead model, this supported SslE fibres being formed from a central core structure, with folded and

dynamic domains presented on the surface (**Fig. 8h**). Comparison of the fibre reconstruction with bead

models derived from the rSslE TEM envelope also suggested that a single molecule of SslE would

satisfy the bead volume but only once the NT1 domain had been removed.

Discussion

In this study we have revealed that SslE is composed of four defined regions: two globular domains at

the N-terminus (NT1 and NT2) and a central torus shaped core, where a NT3 domain packs against an

M60-like aminopeptidase. The N-terminal region (NT1, NT2 and NT3) is composed of a unique

primary sequence that has only been identified in SslE and its homologs[33], while the C-terminal M60-

like domain is found in a wide range of proteins secreted by both prokaryotic and eukaryotic

microorganisms that interact with mucosal membranes[18, 34]. The atomic structures for three M60-like

domains in complex with *O*-glycopeptides have been described: BT4244 from *Bacteroides*

thetaiotaomicron (PDB ID code: 5kd2), IMPa from *P. aeruginosa* (PDB ID code: 5kdw) and ZmpB

from *Clostridium perfringens* (PDB ID code: 5kdn)[19]. While the active site structures in these bacterial

enzymes are highly conserved, there are variations within the remaining M60 domain, which enables

them to recognize unique glycan sequences. Structural data is not available for intact BT4244 or ZmpB,

although they are thought to contain additional carbohydrate binding domains[18] and it is likely that these

M60 augmentations have a role in the recognition of *O*-glycopeptide substrates. It also appears that the

active site in the M60 domain of SslE faces into the NT3 domain and this implies that both the NT3

and M60 domains are responsible for the recognition of mucins in SslE. With significant flexibility of

the NT1 and NT2 region, these domains may also interact with mucin substrates and may promote

processivity.

Although SslE has a clear function in the processing of mucins, it is a unique member of the M60-like

aminopeptidase family as it is also fundamental for the maturation of EPEC biofilms grown on plastics[4].

Furthermore, we have now shown that SslE is also required for the maturation of *E. coli* W and ETEC

H10407 biofilms when grown statically on agar and under shear stress on glass substrates, where it is

widely distributed throughout the extracellular matrix. We have demonstrated that this unusual function

is due to SslE forming aggregates at pH ≤ 4.4; conditions which are also observed as microenvironments

across mature wild-type *E. coli* W biofilms. It is well established that phenotypical heterogeneity

develops during the maturation of biofilms and direct measurement of extracellular pH within *E. coli*

PHL628 biofilms has also shown a heterogenous distributions of pH, with values ranging between 5.0

and 7.0[35]. Therefore, it is very likely that SslE forms functional aggregates within transiently forming

microenvironments across maturing and established biofilms, but then resists solubilisation after any

increase in the local pH.

Microbial amyloid structures are ubiquitous in biofilms and in gram negative bacteria most known

amyloid structures are synthesized by either the Curli or Fap secretion systems[36, 37, 38, 39]. These form

through spontaneous aggregation of extended peptides and provide adhesion, cohesion and contribute

to the structural integrity of the biofilm matrix. Although our biochemical and biophysical analyses

suggests that under acidic condition rSslE displays some amyloid-like features, examination of

aggregates using negative-stain TEM has revealed short fibres associated with globular structures.

SAXS reconstructions of rSslE fibres has allowed additional analysis and suggests that the repetitive

fibre unit can accommodate a single NT2-NT3-M60 region, and we therefore propose a potential model

for the fibrillation of SslE into functional aggregates via end-to-end stacking (**Fig. 8i**). In this model,

the asymmetric unit of the fibre core is composed of the NT2, NT3 and M60 domains and the NT1

domain is flexible and can move freely around the surface. Polymerisation of the fibre core is mediated

through ~40 residues within the NT2 and NT3 domains but not the M60 domain, but this would require

some rearrangement of the NT3/M60 interface.

FTIR analysis of rSslE aggregates provided an amide I maxima of 1624 cm^{-1}, which is consistent with

an increase in anti-parallel β-sheet conformation and also lies within the range of amide I maxima for

other typical amyloid structures (*e.g.* 1611-1630 cm^{-1})[40]. However, larger and more rigid amyloid

structures generally have peaks at lower frequencies, while native β-sheets show absorption bands with

maxima between 1630-1643 cm^{-1}. This suggests that the rSslE amide I band peak represents an amyloid

structure which is relatively small and is not overly rigid, although it could alternatively originate from

fibre stacking/clumping effects. Binding of ThT to rSslE at acidic pH also supports a general increase

in β-sheet content in fibres but could also indicate the formation of new cavities upon fibrillation[41]. It

is therefore still unclear whether these changes reflect the formation of a continuous amyloid-like

structure that runs the length of the fibre core or instead represent β-sheet augmentation between

adjacent SslE molecules along the fibre, upon fibrillation of SslE.

We have shown that SslE influences the localization of exopolysaccharides during the development of

macrocolony biofilms and SslE fibres can directly associate with cellulose, which is known to have a

structural function in many *E. coli* biofilms and provides cohesion and elasticity[42]. Although the precise

role of SslE during biofilm development is still vague, it could have an architectural role acting as a

hub to promote binding between other EPS or it could regulate the correct deposition of EPS within the

biofilm. As monomeric SslE does not recognise cellulose, this suggests that new binding sites are

formed after its aggregation. The presence of a surface exposed domain also implies that this domain

may be functional in binding cellulose or other yet to be identified ligands, or is functional under

different conditions, for example through interacting with mucins during intestinal colonisation.

Likewise, the fibre core may still retain its mucolytic activity and could play a role in mucin

restructuring during biofilm growth in the gut.

Although both *E. coli* W and ETEC 10407 secrete SslE, the W strain is a harmless commensal, while

ETEC is a major etiological agent in developing countries[43]. SslE is also required for colonisation in a

rabbit model of EPEC infection[4] and is actively transcribed during ETEC infection of mice[6]. Biofilm

growth and the production of enterotoxins are essential for the pathogenesis of ETEC-induced disease[43]

and secretion of SslE and the ETEC labile toxin is via the same type II secretion system[4, 44]. This

suggests that SslE increases the virulence of *E. coli* pathotypes through its ability to promote biofilm

maturation and/or through its interactions with mucosal defences. Both enterotoxigenic and

enteropathogenic *E. coli* strains cause infection in the small intestine[1] where the intraluminal pH ranges

between approximately 6.0 to 7.5[45]. However, *E. coli* possess an acid-tolerance response that supports

exponential growth at pH values as low as pH 4.2[46] and due to the synthesis of organic acids by the

residing microbiota, during colonisation of the intestine *E. coli* will experience extracellular pH values

that range between 4.0 and 6.0[47]. Although this suggests that within the intestinal tract, conditions exist

where SslE could regulate biofilm development, it remains unclear whether this role is important for

infection or during extraintestinal survival. SslE represents a unique protein and further studies are now

required to understand its complete functions during ecology and disease.

Methods

Bacterial strains and media

All bacterial strains, plasmids and primers used in this study are listed in **Supplementary**

Table 3. Lysogeny broth (LB) medium contained (per litre) 10 g tryptone, 5 g yeast extract and 10 g

NaCl. Isotopically defined M9 minimal medium (pH 7.4) contained (per litre) 6.0 g $Na_2HPO_4 \cdot 7H_2O$,

3 g KH_2PO_4, 0.5 g NaCl, 0.12 g $MgSO_4 \cdot 7H_2O$, 0.022 g $CaCl_2$, 0.04 g thiamine, 8.3 mg $FeCl_3 \cdot 6H_2O$,

0.5 mg $ZnCl_2$, 0.1 mg $CuCl_2$, 0.1 mg $CoCl_2 \cdot 6H_2O$, 0.1 mg H_3BO_3 and 13.5 mg $MnCl_2 \cdot 6H_2O$,

supplemented with 2g [U-$^{13}C_6$]glucose and/or 0.7 g $^{15}NH_4Cl$. M9 media was made up in deuterium

oxide for the production of perdeuterated protein samples and pH was adjusted using 1 M NaOH

solution. All NMR isotopes were from Sigma.

Gene deletion

A non-polar deletion of *sslE* was constructed in *E. coli* W and H10407 strains by allelic exchange

with FLP recombination target (FRT)-flanked Kan[r], using a modified protocol[48]. This fragment was

amplified from *E. coli* strain JW5925-1[49] using Platinum *Taq* DNA polymerase (Promega) and primers

PC3 and PC4 with specificity for the regions flanking *sslE*. These exchanges were facilitated by the λ

Red recombinase system carried on plasmid pKD46 and knockouts were confirmed by PCR and

sequencing.

Plasmid construction

Complementation plasmids pCPC1 and pCPC2 were generated by amplifying whole length *sslE*, or

sslE minus the M60 domain, from *E. coli* W gDNA using primer pairs PC7/PC8 or PC7/PC9,

respectively. These were digested with HindIII/NheI (NEB), ligated into HindIII/NheI-digested pBad-

cm18 vector. Synthetic genes gPC4, gPC5, gPC6 and gPC7 were synthesized by Synbio Technologies

and cloned into pET28b vector using NcoI and XhoI restriction sites to create plasmids pPC2, pPC3,

pPC4 and pPC5, respectively (**Supplementary Table 4**). Plasmid pBD1 was created by amplification

of the RgpB-CTD from *P. gingivalis* W50 gDNA using primers BD1 and BD2. This was then cloned

into pET46 Ek/LIC vector through ligation independent cloning (Novagen).

Microfluidic biofilm growth

Biofilms were grown in a BioFlux 200 device (Labtech, UK) using a protocol adapted from the

manufacturer. Single colonies of *E. coli* W and its derivatives were resuspended in 10 ml of LB (with

additional 50 µg/ml kanamycin for *Δssle* mutant and 50 µg/ml kanamycin, 25 µg/ml chloramphenicol

Δssle::pCPC1 (*sslE*) and *Δssle::pCPC2* (*sslEΔM60*)) and incubated at 37°C with shaking (200 rpm) for

16 hrs. All subsequent media contained appropriate antibiotics where required. The cultures were then

diluted (1:100) in 10 ml of LB to OD_{600nm} of 0.5. A 24 well Bioflux plate (0-20 dyn/cm^2) was pre-

warmed to 37°C on a heated stage and microfluidic channels were incubated with prewarmed LB. The

channels were then inoculated by injecting 20 µl of bacterial suspension into the output reservoir for 5

s at 2 dyne/cm^2. The microfluidic plate was incubated for 1 hr at 37°C to allow bacteria to bind to the

surface and then the flow reversed for 20 s. Prewarmed LB was then added to the input reservoir, the

flow of media was initiated at 2 dyne/cm^2 for 5 min and then decreased to 1 dyne/cm^2 for 18 hrs. The

spent media was removed from the output reservoir, fresh prewarmed media was added to the input

reservoir and the flow was lowered to 0.5 dyne/cm^2 for up to 96 hrs.

Biofilm pH measurement

The ratiometric dye seminaphthorhodafluor-4F 5-(and-6) carboxylic acid (C-SNARF-4) was used to

directly measure pH across biofilms within the Bioflux channel, using a modified method[50]. HEPES

buffer was first adjusted between pH 3.2 to 7.8 in 0.2 increments using HCl, and 95 µl of each respective

solution was then mixed with 5 µl 1mM C-SNARF-4, resulting in a final 50 µM concentration of C-

SNARF-4. This solution was flushed into the Bioflux plate channel, pre-warmed at 37°C and the

channels were imaged at x64 magnification using a DM-IRE2 confocal laser scanning microscope

(Leica Microsystems Heidelberg GmbH, Germany) and the accompanying Leica Microsystems

Confocal Software (version 2.61 Build 1537). Simultaneous images were captured using an excitation

wavelength of 488 nm and emission wavelengths of 580 nm and 640 nm, in triplicate for each pH

increment with a background image obtained after each acquisition. Acquired images were analysed

using ImageJ Software (FIJI)[51] by determining the average intensity of fluorescence and standard

deviation of each channel, minus the background control. The intensity ratio for each pH was used to

make a standard curve by plotting known pH against green/red ratio. 50µM C-SNARF-4 was then

flushed into Bioflux plates containing *E. coli* W biofilm after either 24 or 96 hrs growth and five

separate images were acquired for each Bioflux channel to calculate the pH from different areas of the

biofilm. The fluorescence intensity and standard deviation were recorded, and the ratiometric value was

used to determine pH from the standard curve. A 12×12 grid (~30 µm^2 area) was applied to each image;

each box was used to define regions of interest (ROI) and an average measure of pH was calculated

within each region. The average pH for ROIs within the biofilm fringes and centres is presented.

Biofilm immunofluorescence

Single colonies of wild-type *E. coli* W or H10407 were resuspended in 10 ml of LB and incubated at

37°C with shaking (200 rpm) for 16 hrs. Cultures were diluted (1:100) in prewarmed LB and 5 ml was

injected onto borosilicate glass coupons held within FC310 flow cells (Biosurface Technology) and

then incubated for 1 hr. Flow of LB media was initiated at 0.4 ml/min and maintained for 96 hrs.

Coupons were removed and washed three times for 10 min each in PBS-Tween. Coupons were then

blocked for 1 hr in 2% (w/v) BSA, PBS-Tween at room temperature and incubated overnight at 4°C

with polyclonal anti-rSslE antibody (rabbit; Invitrogen), diluted 1:200 in 0.1% (w/v) BSA, PBS-Tween.

After three 10 min washes in PBS-Tween, coupons were incubated in the dark for 1 hr with 10 μM

Syto9 (Invitrogen) and anti-rabbit Alexa Fluor 633 secondary antibody (goat; Invitrogen) diluted 1:500

in 0.1% (w/v) BSA, PBS-Tween. This was followed by three 10 min washes in the dark in PBS-Tween

and then fixation for 15 min at room temperature in the dark with 3% (w/v) paraformaldehyde, PBS-

Tween. After incubation for 15 min in 10 mM glycine, PBS-Tween, images were captured at 100x

magnification using an excitation wavelength of 486 nm and 632 nm, and emission wavelengths of 501

nm and 647nm, respectively, with a DM-IRE2 confocal laser scanning microscope (Leica

Microsystems Heidelberg GmbH, Germany). Negative controls consisted of biofilms being incubated

with either no primary anti-rSslE antibody or no Alexa Fluor 633 secondary antibody.

Immunoelectron microscopy

Overnight cultures were incubated for 30 min in 20 mM citrate phosphate buffer at either pH 4.0 or 5.0

and then washed in the same buffer before fixing with the 3% (w/v) paraformaldehyde for 1 hr. Cells

were then loaded onto a Glow discharged carbon coated Ni grid (Agar Scientific) for 10 mins, washed

with 50 mM glycine and then blocked with 1% (v/v) Natural Donkey Serum (Jackson Immunoresearch)

for 30 min. This was then incubated for 1 hr with primary polyclonal rSslE antibody (rabbit; Invitrogen)

diluted 1:100 in blocking buffer incubation, washed five times for 3 min each with 0.05% Natural

Donkey Serum and then incubated for 1 hr with gold-conjugated secondary antibody (donkey; Jackson

Immunoresearch) diluted 1:100 in blocking buffer. Washing was repeated with buffer only ddH$_2$O. The

sample was negatively stained with 2% (w/v) uranyl acetate for 30 s, followed by two quick washes

with double-distilled water. The grid was air dried, and images were recorded on a JEM1230 (JEOL-

Japan) at 80KV with a Morada CCD camera, iTEM software (EMSIS).

Protein purification

Recombinant SslE (residues 67-1497; numbered based on mature sequence) was expressed and

purified as described previously[12]. Likewise, SslE NT1 (residues 67-211), NT2 (residues 230-425),

NT1-NT2 (residues 67-425), NT3-M60 (residues 426-1497) and RgpB-CTD were transformed into *E.*

coli SHuffle cells (SslE; New England Biolabs) or BL21 (DE3) (RgpB-CTD; New England Biolabs),

grown at 37°C in LB media (rSslE, RgpB-CTD: 100 µg/ml ampicillin; NT1, NT2, NT1-NT2, NT3-

M60: 100 µg/ml kanamycin) and expression induced with 0.5 mM isopropyl-d-1-thiogalactopyranoside

(IPTG) at an OD$_{600nm}$ of 0.6, followed by growth overnight at 18°C. Cells were resuspended in 20 mM

Tris–HCl pH 8, 200 mM NaCl, lysed by sonication and purified using nickel affinity chromatography

followed by gel filtration with either a Superdex 75 (NT1, NT2, RgpB-CTD) or 200 (rSslE, NT1-NT2,

NT3-M60) column (GE Healthcare).

SEC-SAXS

SAXS data were collected on beamline B21[52] at Diamond Light Source (DLS, Oxford, UK), United

Kingdom at 25°C. 60 µl of rSslE (10 mg/ml) in 20 mM Tris–HCl pH 8, 200 mM NaCl was applied to

a KW403-4F column (Shodex) at 0.16 ml/min, preequilibrated in 20 mM citrate phosphate buffer, 200

mM NaCl at either pH 4.4 or 7.4, and SAXS data were measured over a momentum transfer range of

$0.003 < q < 0.44$ Å^{-1}. Peak integration and buffer subtraction were performed in CHROMIXS[53]. The

radius of gyration (R_g) and scattering at zero angle [$I(0)$] were calculated from the analysis of the Guinier

region by AUTORG[53]. The distance distribution function [$P(r)$] was subsequently obtained using

GNOM[53] yielding the maximum particle dimension (D_{max}). *Ab initio* low resolution shape restoration

was carried out by calculating 20 models in DAMMIF[53], which were subsequently averaged using

DAMAVER[53] and used as a staring model for refinement in DAMMIN[53]. An additional 20 models were

also calculated and averaged in DAMAVER[53]. CRYSOL[53] was used to compare final rSslE TEM

envelopes and models against the solution SAXS curve. Processing and refinement statistics can be

found in **Supplementary Table 2**.

TEM single particle analysis

4 µl of rSslE (625 nM) in 50 mM Tris-HCl pH 8.0, 150 mM NaCl was applied to previously glow-

discharged 300 mesh continuous carbon-coated copper grids (Agar Scientific Ltd) for 1 min and blotted

for excess liquid. 4 µl of 2% (v/v) uranyl acetate was applied for staining for 1 min. The excess liquid

was blotted and left to dry. Data was acquired using a JEOL JEM-2100 plus TEM operating at 200 kV

equipped with a OneView 16 Megapixel camera (Gatan). 50 micrographs were collected at a nominal

magnification of 50,000× with a pixel size of 2.1 Å/pixel and a range of defocus from 1 to 3 µm. Data

were processed using Relion 3.1[54]. Defocus and astigmatism parameters were estimated using

CTFFIND4[55] in Relion 3.1. An initial dataset of 41,752 particles were autopicked using 2D class

averages generated using approx. 1500 manually picked particles as reference. After a few rounds of

2D classification ignoring CTF until the first peak, 10,988 particles were taken forward for an initial

model generation using 3D initial model in Relion 3.1[54]. Following initial model generation, a few

rounds of 3D classification followed by 3D refinement was carried out. The final model was refined to

22 Å using the gold standard FSC (0.143 criterion). PHYRE2[56] was used to analyse the SslE sequence

(residues 67-1497; UniProt ID E0IW31) and generate a homology model for the C-terminal M60

domain (residues 962-1415) based on residues 421-894 of *P. aueriginosa* IMPa (PDB ID code 5kdv;

100% confidence, 21% identity)[19]. This was then docked into the rSslE TEM envelope using UCSF

CHIMERA[57].

Solution NMR spectroscopy

NMR measurements for rSslE were performed at 25°C on a 100 μM ^{2}H^{15}N-labelled sample in 50 mM

NaPO$_4$ pH 7.4, 100 mM NaCl, 10 % D$_2$O. NMR measurements for SslE NT1 (0.6 mM), SslE NT2 (1.3

mM) and SslE NT1-NT2 (0.8 mM) were performed at 25°C on ^{15}N-labelled samples in 50 mM NaPO$_4$

pH 7.0, 100 mM NaCl, 10 % D$_2$O. NMR measurements for RgpB-CTD (0.3 mM) were performed at

37°C on a ^{15}N-labelled sample in 20 mM NaPO$_4$ pH 6.0, 100 mM NaCl, 10% D$_2$O. Transverse

relaxation optimized spectroscopy (TROSY) based ^{1}H^{15}N-HSQC experiment and T_1 and T_2 relaxation

times for rSslE were recorded on a Bruker Avance III HD 950, equipped with a TXI cryoprobe. TROSY

^{1}H^{15}N-HSQC experiment for SslE NT1-NT2 was recorded on a on a Bruker Avance III HD 800,

equipped with a TCI cryoprobe. Standard ^{1}H^{15}N-HSQC spectra of SslE NT1, SslE NT2 and RgpB-CTD

were recorded on a Bruker Avance III HD 700, equipped with a TCI cryoprobe. Data were processed

in NMRPIPE[58] and analysed/visualized with ANALYSIS[59] and NMRVIEW[60].

SAXS fibrillation analysis

SAXS data were collected on beamline B21[52] at DLS (Oxford, UK) at 20°C. Immediately prior to data

collection, 10 mg/ml rSslE in 20 mM Tris–HCl pH 8, 200 mM NaCl was buffer exchanged into 20 mM

citrate-phosphate buffer, pH 4.4 using a PD10 column (GE Healthcare) and the flow through was used

as a buffer reference. Using a peristaltic pump and while constantly stirring, 5 ml of rSslE (0.7 mg/ml)

was circulated through the SAXS imaging cuvette and data were collected every 30 min over a

momentum transfer range of $0.004 < q < 0.4$ Å^{-1}, with the initial scattering data captured at 30 min after

initiating fibre growth. Data collected over the course of 11 hrs, consisting of 22 profiles, was

decomposed using COSMiCS[30], which utilizes MCR-ALS[61] to perform model free decomposition of

the entire SAXS data set. The SEC-SAXS curve collected at pH 4.4 was also included in the dataset as

the representative state of the protein at time 0 s, *i.e.* before initiating fibrillation. The time 0 s curve

was selected as one of the initial estimates and the selectivity restraint was used to ascertain that the

curve had no contribution from the other species. In addition, non-negativity restraint was imposed for

both the SAXS profiles and concentrations using FNNLS algorithm[62]. Before COSMiCS analysis,

SAXS data were scaled according to the large angle data to enhance the decomposition capacity of the

approach. Although COSMiCS was run assuming a two-component system as suggested by Principal

Component Analysis (PCA), a three-species run was also performed. The convergence criterion of <

0.01 % change in lack of fit was used with 1,000 maximum allowed iterations. The analysis of the

resulting COSMiCS curves was performed with the ATSAS suit of programs[53]. Fractal analysis was

carried out using SASVIEW (http://www.sasview.org/).

SAXS fibre modelling

The programs ATSAS[53] and Scatter were used to obtain a cross-sectional radius of gyration R_c from the

10 hr post-fibre induction scattering profile and from this the cross-sectional $P(r)_C$ was calculated.

BODIES[53] was then used to approximate the geometric shape of rSslE fibres, which suggested a

cylindrical shape with radius of 3.93 nm and a height of 16.5 nm (χ^2 1.6). These dimensions were then

used in DAMMIN[63] with data from the q range 0.03 to 0.20 to create *ab initio* shapes of the SslE fibril

repeat. 20 individual jobs were run, generating 20 independent models and were subsequently averaged

using DAMAVER[53] and then refined 5 times in DAMMIN[63]. A bead model was produced from the

rSslE negative-stain TEM envelope using the program EM2DAM[53], the NT1 bead region was removed

and then this was superposed onto the DAMMIN fibre bead model using SUPCOMB[53]. Processing and

refinement statistics can be found in **Supplementary Table 2**.

TEM fibre analysis

rSslE at 1 mg/ml was incubated in 100 mM citrate phosphate pH 3.8 overnight at room temperature

while shaking at 180 rpm. This was centrifuged at 15,000 g for 10 min, the top 80% of buffer discarded

and then 4 µl of the remaining sample was applied to a previously glow-discharged 300 mesh

continuous carbon-coated copper grids (Agar Scientific Ltd) and immediately blotted for excess liquid.

4 µl of 2% (v/v) uranyl acetate was applied for staining for 10 seconds. The excess liquid was blotted

and left to dry. Data was acquired using a JEOL JEM-1230 TEM operating at 80 kV equipped with a

Morada 2k CCD camera system and its iTEM software package (Olympus Europa, UK). Micrographs

were collected at a nominal magnification of 80,000x with a pixel size of 5.96 Å/pixel.

Solid state NMR spectroscopy

rSslE at 20 mg/ml was buffer exchanged into 10 mM citrate phosphate pH 4.0 using a PD10 column (Sigma) and then incubated overnight at room temperature while shaking at 180 rpm. This was centrifuged at 15,000 g for 10 min, the top 80% of buffer was discarded and then the remaining sample was used for solid state NMR analysis. Experiments were performed using a Bruker Neo Console operating at 850 MHz ^{1}H frequency with a 3.2 mm E-Free probe in HC mode spinning at a rate w_r= 15 kHz. A standard CP excitation ^{13}C-^{13}C DARR experiment was acquired. The direct dimension was acquired for 16.4 ms with ~55 kHz SPINAL-64 decoupling. 256 rows with 256 coadded transients and a recovery delay of 2.5 s were acquired using Time-Proportional Phase Increment (TPPI) with a 22.2 µs dwell (45 kHz sweep width, 2.84 ms total evolution) for a total of 45.5 hrs total acquisition time. The applied power was adjusted so that ^{1}H and ^{13}C hard pulses were both 4 µs ($w_{1(H,C)}$= 62.5 kHz). The initial carbon excitation was achieved with 1.5 ms of ramped Hartmann-Hahn CP, where the optimal polarization transfer was found at w_{1C}= ~70 kHz and w_{1H}= ~55 kHz with an upwards linear ramp from 70-100% on the ^{1}H channel. Homonuclear Carbon mixing was achieved with 50 ms of DARR mixing (w_{1H}= w_r= 15 kHz).

Acknowledgements

PC was supported by the Academy of Medical Sciences and Wellcome Trust (SBF002/1150) and SR was supported by the Medical Research Council (MR/R017662/1), awarded to JAG. SW and BD were supported by China Scholarship Council and Biotechnology and Biological Sciences Research Council studentships, respectively, awarded to JAG and MAC. LS was supported by the Leverhulme Trust (RPG-2017-222), awarded to JAG. KF was supported by a PhD studentship from Queen Mary University of London and VCD was supported by a start-up grant from Queen Mary University of London. This work was also supported by the Labex EpiGenMed, an « Investissements d'avenir » program (ANR-10-LABX-12-01) awarded to PB. The CBS is a member of France-BioImaging (FBI)

and the French Infrastructure for Integrated Structural Biology (FRISBI), 2 national infrastructures supported by the French National Research Agency (ANR-10-INBS-04-01 and ANR-10-INBS-05, respectively). We thank the beamline scientists at B21 of the Diamond Light Source, United Kingdom. This work was also supported by the Francis Crick Institute through provision of access to the MRC Biomedical NMR centre. The Francis Crick Institute receives its core funding from Cancer Research United Kingdom (FC001029), the United Kingdom Medical Research Council (FC001029), and the Wellcome Trust (FC001029). We also thank the Centre for Biomolecular Spectroscopy at King's College London for additional NMR access, funded by the Wellcome Trust and British Heart Foundation (ref. 202767/Z/16/Z and IG/16/2/32273 respectively). The UK 850 MHz solid-state NMR Facility used in this research was funded by EPSRC and BBSRC (contract reference EP/T015063/1), as well as the University of Warwick including via part funding through Birmingham Science City Advanced Materials Projects 1 and 2 supported by Advantage West Midlands (AWM) and the European Regional Development Fund (ERDF). Cellulose discs were gifted by Prof. Tom Ellis and Kenneth T. Walker (Imperial College London).

Author Contributions

Conceived and designed the experiments: PC, SW, SR, KF, SS, CE-G, AS, GM, BD, LMS, SL, DI, PB, VCD, JAG. Performed the experiments: PC, SW, SR, KF, AS, SS, LC, CE-G, GM, BD, LMS, SL, DI, JJ, GHC, VCD, JAG. Analyzed the data: PC, SW, SR, KF, AS, SS, CE-G, GM, BD, SL, DI, TF, PB, VCD, JAG. Contributed reagents/materials/analysis tools: SS, SL, PB, VCD, JAG. Wrote the paper: PC, SW, SR, KF, AS, SS, LC, CE-G, GM, BD, LMS, SL, DI, TF, JJ, GHC, MAC, PB, VCD, JAG.

References

1.   Kaper JB, Nataro JP, Mobley HL. Pathogenic Escherichia coli. *Nat Rev Microbiol* **2**, 123-140 (2004).

622 2.   Gu S, Shevchik VE, Shaw R, Pickersgill RW, Garnett JA. The role of intrinsic disorder and
623   dynamics in the assembly and function of the type II secretion system. *Biochim Biophys Acta*
624   *Proteins Proteom* **1865**, 1255-1266 (2017).
625

626 3.   Moriel DG*, et al.* Identification of protective and broadly conserved vaccine antigens from
627   the genome of extraintestinal pathogenic Escherichia coli. *Proc Natl Acad Sci U S A* **107**,
628   9072-9077 (2010).
629

630 4.   Baldi DL*, et al.* The type II secretion system and its ubiquitous lipoprotein substrate, SslE, are
631   required for biofilm formation and virulence of enteropathogenic Escherichia coli. *Infect*
632   *Immun* **80**, 2042-2052 (2012).
633

634 5.   Decanio MS, Landick R, Haft RJ. The non-pathogenic Escherichia coli strain W secretes SslE
635   via the virulence-associated type II secretion system beta. *BMC Microbiol* **13**, 130 (2013).
636

637 6.   Nesta B*, et al.* SslE elicits functional antibodies that impair in vitro mucinase activity and in
638   vivo colonization by both intestinal and extraintestinal Escherichia coli strains. *PLoS Pathog*
639   **10**, e1004124 (2014).
640

641 7.   Yang J, Baldi DL, Tauschek M, Strugnell RA, Robins-Browne RM. Transcriptional
642   regulation of the yghJ-pppA-yghG-gspCDEFGHIJKLM cluster, encoding the type II
643   secretion pathway in enterotoxigenic Escherichia coli. *J Bacteriol* **189**, 142-150 (2007).
644

645 8.   Naili I*, et al.* Mixed mucosal-parenteral immunizations with the broadly conserved
646   pathogenic Escherichia coli antigen SslE induce a robust mucosal and systemic immunity
647   without affecting the murine intestinal microbiota. *Vaccine* **37**, 314-324 (2019).
648

649 9.   Luo Q, Qadri F, Kansal R, Rasko DA, Sheikh A, Fleckenstein JM. Conservation and
650   immunogenicity of novel antigens in diverse isolates of enterotoxigenic Escherichia coli.
651   *PLoS Negl Trop Dis* **9**, e0003446 (2015).
652

653 10. Valeri M*, et al.* Pathogenic E. coli exploits SslE mucinase activity to translocate through the
654   mucosal barrier and get access to host cells. *PLoS One* **10**, e0117486 (2015).
655

656 11. Luo Q*, et al.* Enterotoxigenic Escherichia coli secretes a highly conserved mucin-degrading
657   metalloprotease to effectively engage intestinal epithelial cells. *Infect Immun* **82**, 509-521
658   (2014).
659

660 12. Rehman S*, et al.* Structure and functional analysis of the Legionella pneumophila chitinase
661   ChiA reveals a novel mechanism of metal-dependent mucin degradation. *PLoS Pathog* **16**,
662   e1008342 (2020).
663

664 13. Corfield AP. Mucins: a biologically relevant glycan barrier in mucosal protection. *Biochim*
665   *Biophys Acta* **1850**, 236-252 (2015).

667   14.   Garnett JA, Matthews S. Interactions in bacterial biofilm development: a structural
668   perspective. *Curr Protein Pept Sci* **13**, 739-755 (2012).
669

670   15.   Vogeleer P, Tremblay YD, Mafu AA, Jacques M, Harel J. Life on the outside: role of
671   biofilms in environmental persistence of Shiga-toxin producing Escherichia coli. *Front*
672   *Microbiol* **5**, 317 (2014).
673

674   16.   Chitlapilly Dass S*, et al.* Impact of mixed biofilm formation with environmental
675   microorganisms on E. coli O157:H7 survival against sanitization. *NPJ Sci Food* **4**, 16 (2020).
676

677   17.   Rossi E*, et al.* "It's a gut feeling" - Escherichia coli biofilm formation in the gastrointestinal
678   tract environment. *Crit Rev Microbiol* **44**, 1-30 (2018).
679

680   18.   Nakjang S, Ndeh DA, Wipat A, Bolam DN, Hirt RP. A novel extracellular metallopeptidase
681   domain shared by animal host-associated mutualistic and pathogenic microbes. *PLoS One* **7**,
682   e30287 (2012).
683

684   19.   Noach I*, et al.* Recognition of protein-linked glycans as a determinant of peptidase activity.
685   *Proc Natl Acad Sci U S A* **114**, E679-E688 (2017).
686

687   20.   Trastoy B, Naegeli A, Anso I, Sjogren J, Guerin ME. Structural basis of mammalian mucin
688   processing by the human gut O-glycopeptidase OgpA from Akkermansia muciniphila. *Nat*
689   *Commun* **11**, 4844 (2020).
690

691   21.   Taglialegna A, Lasa I, Valle J. Amyloid Structures as Biofilm Matrix Scaffolds. *J Bacteriol*
692   **198**, 2579-2588 (2016).
693

694   22.   Belousov MV*, et al.* M60-like metalloprotease domain of the Escherichia coli YghJ protein
695   forms amyloid fibrils. *PLoS One* **13**, e0191317 (2018).
696

697   23.   Taglialegna A*, et al.* The biofilm-associated surface protein Esp of Enterococcus faecalis
698   forms amyloid-like fibers. *NPJ Biofilms Microbiomes* **6**, 15 (2020).
699

700   24.   Taglialegna A*, et al.* Staphylococcal Bap Proteins Build Amyloid Scaffold Biofilm Matrices
701   in Response to Environmental Signals. *PLoS Pathog* **12**, e1005711 (2016).
702

703   25.   Chen D*, et al.* Characteristics and influencing factors of amyloid fibers in S. mutans biofilm.
704   *AMB Express* **9**, 31 (2019).
705

706   26.   Gallardo R, Ranson NA, Radford SE. Amyloid structures: much more than just a cross-beta
707   fold. *Curr Opin Struct Biol* **60**, 7-16 (2020).

709   27.   Moran SD, Zanni MT. How to Get Insight into Amyloid Structure and Formation from
710   Infrared Spectroscopy. *J Phys Chem Lett* **5**, 1984-1993 (2014).
711

712   28.   Conway KA, Harper JD, Lansbury PT, Jr. Fibrils formed in vitro from alpha-synuclein and
713   two mutant forms linked to Parkinson's disease are typical amyloid. *Biochemistry* **39**, 2552-
714   2563 (2000).
715

716   29.   Buchan DWA, Jones DT. The PSIPRED Protein Analysis Workbench: 20 years on. *Nucleic
717   Acids Res* **47**, W402-W407 (2019).
718

719   30.   Herranz-Trillo F, Groenning M, van Maarschalkerweerd A, Tauler R, Vestergaard B,
720   Bernado P. Structural Analysis of Multi-component Amyloid Systems by Chemometric
721   SAXS Data Decomposition. *Structure* **25**, 5-15 (2017).
722

723   31.   Schmidt P. Small-angle scattering studies of disordered, porous and fractal systems. *Journal
724   of Applied Crystallography* **24**, 414-435 (1991).
725

726   32.   Hurd AJ, Schaefer DW, Martin JE. Surface and mass fractals in vapor-phase aggregates. *Phys
727   Rev A Gen Phys* **35**, 2361-2364 (1987).
728

729   33.   Gish W, States DJ. Identification of protein coding regions by database similarity search. *Nat
730   Genet* **3**, 266-272 (1993).
731

732   34.   Ta BTT*, et al.* Identification, recombinant protein production, and functional analysis of a
733   M60-like metallopeptidase, secreted by the liver fluke Opisthorchis viverrini. *Parasitol Int*
734   **75**, 102050 (2020).
735

736   35.   Hidalgo G*, et al.* Functional tomographic fluorescence imaging of pH microenvironments in
737   microbial biofilms by use of silica nanoparticle sensors. *Appl Environ Microbiol* **75**, 7426-
738   7435 (2009).
739

740   36.   Van Gerven N, Van der Verren SE, Reiter DM, Remaut H. The Role of Functional Amyloids
741   in Bacterial Virulence. *J Mol Biol* **430**, 3657-3684 (2018).
742

743   37.   Shanmugam N, Baker M, Ball SR, Steain M, Pham CLL, Sunde M. Microbial functional
744   amyloids serve diverse purposes for structure, adhesion and defence. *Biophys Rev* **11**, 287-
745   302 (2019).
746

747   38.   Dueholm MS*, et al.* Functional amyloid in Pseudomonas. *Mol Microbiol* **77**, 1009-1020
748   (2010).
749

750   39.   Evans ML, Chapman MR. Curli biogenesis: order out of disorder. *Biochim Biophys Acta*
751   **1843**, 1551-1558 (2014).
752

753   40.   Zandomeneghi G, Krebs MR, McCammon MG, Fandrich M. FTIR reveals structural
754   differences between native beta-sheet proteins and amyloid fibrils. *Protein Sci* **13**, 3314-3321
755   (2004).
756

757   41.   Groenning M, Olsen L, van de Weert M, Flink JM, Frokjaer S, Jorgensen FS. Study on the
758   binding of Thioflavin T to beta-sheet-rich and non-beta-sheet cavities. *J Struct Biol* **158**, 358-
759   369 (2007).
760

761   42.   Serra DO, Richter AM, Hengge R. Cellulose as an architectural element in spatially
762   structured Escherichia coli biofilms. *J Bacteriol* **195**, 5540-5554 (2013).
763

764   43.   Qadri F, Svennerholm AM, Faruque AS, Sack RB. Enterotoxigenic Escherichia coli in
765   developing countries: epidemiology, microbiology, clinical features, treatment, and
766   prevention. *Clin Microbiol Rev* **18**, 465-483 (2005).
767

768   44.   Sjoling A, von Mentzer A, Svennerholm AM. Implications of enterotoxigenic Escherichia
769   coli genomics for vaccine development. *Expert Rev Vaccines* **14**, 551-560 (2015).
770

771   45.   Fallingborg J. Intraluminal pH of the human gastrointestinal tract. *Dan Med Bull* **46**, 183-196
772   (1999).
773

774   46.   Xu Y*, et al.* An acid-tolerance response system protecting exponentially growing Escherichia
775   coli. *Nat Commun* **11**, 1496 (2020).
776

777   47.   Lin J, Smith MP, Chapin KC, Baik HS, Bennett GN, Foster JW. Mechanisms of acid
778   resistance in enterohemorrhagic Escherichia coli. *Appl Environ Microbiol* **62**, 3094-3100
779   (1996).
780

781   48.   Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in Escherichia coli
782   K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-6645 (2000).
783

784   49.   Baba T*, et al.* Construction of Escherichia coli K-12 in-frame, single-gene knockout mutants:
785   the Keio collection. *Mol Syst Biol* **2**, 2006 0008 (2006).
786

787   50.   Schlafer S, Dige I. Ratiometric Imaging of Extracellular pH in Dental Biofilms. *J Vis Exp*,
788   (2016).
789

790   51.   Schindelin J*, et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*
791   **9**, 676-682 (2012).
792

793   52.   Cowieson NP, *et al.* Beamline B21: high-throughput small-angle X-ray scattering at Diamond
794   Light Source. *J Synchrotron Radiat* **27**, 1438-1446 (2020).
795

796   53.   Franke D, *et al.* ATSAS 2.8: a comprehensive data analysis suite for small-angle scattering
797   from macromolecular solutions. *J Appl Crystallogr* **50**, 1212-1225 (2017).
798

799   54.   Zivanov J, *et al.* New tools for automated high-resolution cryo-EM structure determination in
800   RELION-3. *Elife* **7**, (2018).
801

802   55.   Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron
803   micrographs. *J Struct Biol* **192**, 216-221 (2015).
804

805   56.   Kelley LA, Mezulis S, Yates CM, Wass MN, Sternberg MJ. The Phyre2 web portal for
806   protein modeling, prediction and analysis. *Nat Protoc* **10**, 845-858 (2015).
807

808   57.   Pettersen EF, *et al.* UCSF Chimera--a visualization system for exploratory research and
809   analysis. *J Comput Chem* **25**, 1605-1612 (2004).
810

811   58.   Delaglio F, Grzesiek S, Vuister GW, Zhu G, Pfeifer J, Bax A. NMRPipe: a multidimensional
812   spectral processing system based on UNIX pipes. *J Biomol NMR* **6**, 277-293 (1995).
813

814   59.   Vranken WF, *et al.* The CCPN data model for NMR spectroscopy: development of a software
815   pipeline. *Proteins* **59**, 687-696 (2005).
816

817   60.   Johnson BA, Blevins RA. NMR View: A computer program for the visualization and analysis
818   of NMR data. *J Biomol NMR* **4**, 603-614 (1994).
819

820   61.   Jaumot J, Gargallo R, de Juan A, Tauler R. A graphical user-friendly interface for MCR-ALS:
821   a new tool for multivariate curve resolution in MATLAB. *Chemometrics and Intelligent*
822   *Laboratory Systems* **76**, 101-110 (2005).
823

824   62.   Bro R, De Jong S. A fast non-negativity-constrained least squares algorithm. *Journal of*
825   *Chemometrics* **11**, 393-401 (1997).
826

827   63.   Volkov VV, Svergun DI. Uniqueness of ab initio shape determination in small-angle
828   scattering. *Journal of Applied Crystallography* **36**, 860-864 (2003).
829

Figures

Fig. 1 | Characterization of SslE aggregates. a, Schematic of SslE from *E. coli* W with mature

sequence numbers and structural features annotated. SS: periplasmic signal sequence; L: flexible linker;

NT: unique N-terminal region; M60: peptidase/mucinase domain. **b,** Purified rSslE forms a clear ring

of aggregated protein when incubated in citrate-phosphate buffer between pH 4.0 and 4.4 (upper panel;

white asterisks). The ring formed above the liquid line is due to shaking of the sample and it resists

solubilization up to pH 6.2 once formed (lower panel; purple asterisk), although some loss of protein

band is observed after pH 5.8 (lower panel; red asterisks). **c,** Increase in ThT fluorescence emission on

binding to rSslE fibres formed in citrate-phosphate buffer over pH 4.0 to 8.0. **d,** Immunoblot of rSslE

incubated in citrate-phosphate buffer from pH 3.8 to 4.8 and detected with anti-His$_6$ antibody. rSslE

present in the well represents aggregated protein. **e,** Far-UV CD spectra of soluble rSslE at pH 4.4 (solid

line) and 7.4 (dashed line). Secondary structure analysis is shown in the doughnut chart. **f,** ATR FT-MIR spectrum of the amide I region of rSslE fibres, displaying relatively high and low intensity absorption bands at ~ 1624 and 1695 cm^{-1} respectively, characteristic of an anti-parallel β-sheet structure.

Fig. 2 | Analysis of SslE-dependent biofilm formation by *E. coli* W and H10407 strains. a, Macrocolony biofilm phenotype of wild-type *E. coli* W and H10407 strains and their derivatives on Congo red agar medium after 24 and 96 hrs of growth. The derivatives *sslE* and *sslE*ΔM60 represent *trans*-complementation of intact *sslE* or *sslE* minus its M60 domain, respectively, into the *E. coli* W Δ*sslE* mutant. **b,** CLSM images of wild-type *E. coli* W biofilms grown for 48 and 96 hrs and stained with lipophilic probe FM 1-43 (green). **c,** Quantification of 3-dimensional biofilm intensity of wild-type *E. coli* W biofilms and its derivatives carried out in ImageJ[40]. *P <0.05; ***P <0.001; verses W 96 hrs growth by two-tailed Student's t-test. **d,** CLSM images of *E. coli* W Δ*sslE* mutant and *trans*-complementation with *sslE* (Δ*sslE::sslE*) and *sslE*ΔM60 (Δ*sslE::sslE*ΔM60) stained with FM 1-43 (green). **e,** The pH across wild-type *E. coli* W biofilms grown for 96 hrs was monitored ratiometrically using C-SNARF-4 (green). pH values were calculated over ~30 μm^2 boxes and pH values for representative regions within the biofilm fringes and centres are annotated. Dotted line outlines water channel within the biofilm, where pH values are not shown. Scale bar represents 20 μm. All data are representative of at least three independent experiments.

Fig. 3 | Analysis of SslE aggregation after bacterial secretion. a, Accumulation of SslE aggregates on the side of glass tubes containing overnight cultures of *E. coli* W strains, grown in LB media at pH 5.0 or 7.0. Arrows indicate Congo red stained ring containing SslE. **b-f,** SslE localization upon secretion from *E. coli* W strains assessed by immunoelectron microscopy. Bacteria were reacted with primary α-rSslE antibodies and secondary gold-labelled antibodies on carbon coated nickel grids and then negatively stained with uranyl acetate. **b, e,** *E. coli* W Δ*sslE* mutant washed in citrate-phosphate buffer at pH 5.0 or 4.0, respectively. **c,** Wild-type W strain washed in citrate-phosphate buffer at pH 5.0 with α-rSslE antibodies reacting with the bacterial surface and secreted fibrous material. **d,** Image highlighting clustering of antibodies to an extracellular aggregate. **f,** Δ*sslE::sslE* W strain washed in citrate-phosphate buffer at pH 4.0, again with α-rSslE antibodies reacting with the bacterial surface and secreted fibrous material. Scale bar is equivalent to 200 nm.

Fig. 4 | Localization of SslE within *E. coli* biofilms. Wild-type *E. coli* W and H10407 biofilms were formed using a continuous flow cell system over 96 hrs. Immunofluorescence were performed on fixed biofilms wherein the biofilms were incubated with α-rSslE antibodies (1:200 dilution) then incubate with goat anti-rabbit conjugated Alexa Fluor 633 (yellow). DNA was stained with Syto9 (cyan) to identify bacteria and eDNA within the biofilm matrix. Biofilms were visualized by CLSM. Scale bar is equivalent to 10 µm. All data are representative of three independent experiments.

Fig. 5 | Binding properties of SslE aggregates. a, Macrocolony biofilm phenotype of wild-type and

Δ*sslE* mutant *E. coli* W and H10407 strains on Congo red or calcofluor white agar medium after 96 hrs

of growth. Macrocolonies were visualized with white light to determine the edge of the colony, and

then either epi-red or UV light was used to determine the localization of Congo red or calcofluor white,

respectively, within the colony. **b,** ELISA analysis of binding between cellulose discs and His-tagged

rSslE monomers, His-tagged rSslE fibres and His-tagged *P. gingivalis* RgpB-CTD control, at pH 6.0

and detected with anti-His$_6$ antibody. BSA-coated wells were used as controls. $^*P$ <0.05; verses control

empty well (-) by two-tailed Student's test. All data are representative of at least three independent

experiments performed in triplicate.

Fig. 6 | Structural features of monomeric SslE. a, SAXS bead model of rSslE at pH 7.4. **b,** Fit of the SAXS bead model (teal line) and negative-stain TEM map (black line) of rSslE to the rSslE SAXS data (black open circles) with χ^2 of 1.2 and 113.7, respectively. **c,** Representative negative-stain TEM micrograph of rSslE (pH 7.4) at 50,000× nominal magnification with representative 2D classifications. Scale bar represents 20 nm. **d,** Overlay of the 22 Å resolution TEM map (grey) and SAXS bead model (teal) with the three defined regions in SslE highlighted. **e,** Docking of an SslE M60 domain homology model (orange) into the TEM map. **f,** TEM map of rSslE colored based on domain organization. The NT3-M60 interdomain channel is highlighted with an arrow. **g,** TROSY ^{1}H^{15}N-HSQC spectrum of rSslE (black) overlaid with ^{1}H^{15}N-HSQC spectra of the SslE NT1 (light green) and NT2 (olive green) domains.

Fig. 7 | Kinetics of SslE aggregation. a, SAXS normalized Kratky plots for rSslE at pH 4.4 and 7.4.

b, Overlay of rSslE SAXS bead models at pH 4.4 (red) and 7.4 (teal) with rSslE TEM map (grey). NT1

and NT2 domains are highlighted. **c,** COSMiCS decomposition of rSslE SAXS fibrillation data,

assuming the co-existence of two components. SEC-SAXS curve for rSslE at pH 4.4 is also shown for

t=0. **d,** COSMiCS decomposed scattering curves for rSslE monomer (red) and aggregates (blue). **e,**

Shape distribution [*P(r)*] functions derived from COSMiCS decomposed scattering curves for rSslE

monomer (red) and aggregates (blue). **f,** Relative populations of the two species of rSslE derived from

the COSMiCS analysis over time.

Fig. 8 | Model of SslE fibres. a, TEM analysis of negatively stained SslE fibres. Two major species are observed. **b,** Smaller filaments have an average width of ~4.5 nm and are coated with globular structures of ~4.5 nm in diameter. Scale bar, 20 nm. **c,** Schematic describing the overall dimensions of the smaller SslE fibre species. **d,** Larger species of SslE fibres appear as ~20 to 40 nm wide structures with variable lengths. Scale bar, 20 nm. **e, f,** DARR ssNMR spectrum of SslE fibres focussed on the carbonyl (**e,** left), aliphatic (**e,** right) and aromatic (**f**) regions. Likely amino acid type assignments are highlighted with red dashed lines and annotated in teal. **g,** SAXS bead model for the repeating unit of the SslE fibre core with overall dimensions given. **h,** Three units of the SslE fibre core translated along

the fibre axis (left). One unit is zoomed in and overlaid with a bead model created of the rSslE negative-stain TEM map with the NT1 domain removed (NT2-3-M60; right). **i,** Proposed model for SslE fibres. The core of the fibre contains the NT2, NT3 and M60 domains and is decorated with flexible NT1. The core fibre is formed via interactions between the NT2 and NT3 domains of adjacent SslE molecules.