Behavioural adaptations in egg laying ancestors facilitate evolutionary transitions to live birth

Amanda K. Pettersen[1*], Nathalie Feiner[1], Daniel W.A. Noble[2], Geoffrey M. While[3], Charlie K. Cornwallis[1#] & Tobias Uller[1#]

1. Department of Biology, Lund University, Lund, 22 362, Sweden

2. Division of Ecology and Evolution, Research School of Biology, The Australian National University, Canberra, 2600, Australia

3. School of Natural Sciences, University of Tasmania, Hobart, 7005, Australia

#Joint senior author

*Corresponding author: Amanda K. Pettersen. Email: amanda.pettersen@sims.org.au

Keywords: Viviparity, Oviparity, Thermal plasticity, Behavioural plasticity, Reproductive mode

Abstract

Live birth is a key innovation that has evolved from egg laying over 100 times in reptiles.

One significant feature in this transition is the thermal conditions experienced by developing

embryos. Adult lizards and snakes often have preferred body temperatures that can be lethal

to developing embryos and should prevent egg retention: how has viviparity repeatedly

evolved in the face of this pervasive mismatch? Here we resolve this paradox by conducting

phylogenetic analyses using data on thermal preference from 224 species. Thermal

mismatches between mothers and offspring are widespread but resolved by gravid females

behaviourally down-regulating their body temperature towards the thermal optimum of

embryos. Importantly, this thermoregulatory behaviour evolved in ancestral egg-laying

species before the evolutionary emergence of live birth. Maternal thermoregulatory behaviour

therefore bypasses constraints imposed by a slowly evolving thermal physiology and is likely

to have been a key requirement for repeated transitions to live birth.

Introduction

The evolution of live bearing is an important life-history transition in vertebrates (1–3). The ecological conditions that favour the transition from egg laying (oviparity) to live birth (viviparity) are relatively well understood, especially in reptiles, with particularly strong support for the adaptive value of viviparity in cool climates (3, 4). By retaining embryos throughout development, mothers can buffer offspring from suboptimal nest temperatures, thereby ensuring higher hatching success and increased offspring viability (4). Live bearing also allows females to maintain offspring at higher temperatures than those of potential nest sites, effectively increasing developmental rates to facilitate early-season hatching (5, 6). This has allowed reptile species to diversify and persist in cool climates globally (7).

Despite its adaptive advantage, it is unclear how viviparity initially evolved. A major challenge to the evolution of viviparity is that adult lizards and snakes typically have preferred body temperatures that exceed the upper lethal limit of embryos (8). For example, the average nest temperature for the Iberian emerald lizard, *Lacerta schreiberi,* is 24°C, rarely exceeding 30°C, while the preferred body temperature of females is 33°C (9). Since embryos are well adapted to the temperatures they typically experience in the nest, they generally have limited capacity to sustain development at temperatures outside a narrow thermal window (10, 11). Prolonged exposure to temperatures beyond this range can result in offspring abnormalities and even death (12). A mismatch between thermal optima of embryos and females should therefore prevent the maternal retention of embryos throughout development, and therefore evolutionary transitions to live birth (8).

Despite this apparent constraint, live birth has evolved over 100 times in squamate reptiles (13, 14). How can the repeated evolution of viviparity be reconciled with the widespread thermal mismatch between embryos and adults? There are two potential scenarios by which viviparity can evolve in the presence of maternal-embryo thermal mismatches.

First, viviparity may only evolve in species where there is no mismatch, that is, adult body

temperature (P_{bt}) and embryo thermal optima (T_{opt}) are well aligned. Second, females may

behaviourally adjust their body temperature when pregnant to close the gap between adult

and embryo thermal optima, even when substantial mismatches exist. Such plasticity may

come at a cost to female performance, but temporarily shifting body temperature while gravid

to match the thermal optima of embryos could eliminate thermal barriers to the evolution of

viviparity.

46   Elucidating how the mismatch between female and offspring thermal optima have

been resolved in viviparous lineages is challenging. Retracing the evolutionary steps that

have led to viviparity requires the reconstruction of thermal preferences of adults and

embryos at the very origins of these lineages. Here, we collate data on the thermal

preferences of female adults (163 species) and embryos (50 species) of squamate reptiles, and

use phylogenetic analyses to reconstruct the evolutionary history of live birth (Table S1).

First, we estimated the ancestral states of preferred body temperature of females and embryos

to quantify the thermal conflict between mothers and embryos at the origins of viviparity.

Second, we tested the prediction that transitions to viviparity have been restricted to

oviparous lineages with a low degree of maternal-embryo mismatch. This was assessed by

calculating if viviparity has originated more frequently from oviparous ancestors where

female *Pbt* overlaps with the thermal optima of embryos. Finally, we examined if females

adjust their body temperature when gravid to reduce the mismatch between adult and embryo

thermal optima, and if so, whether this behaviour evolved prior to the emergence of

viviparity, or afterwards in response to such mismatch.

Results and Discussion

Wide-spread mismatch in thermal optima of oviparous mothers and their embryos

Across oviparous lizards and snakes, there were many species with a significant disparity in

preferred body temperature (P_{bt}) and the temperature that maximizes hatching success (T_{opt})

(Multi-response BPMM (MR-BPMM): posterior mode (PM) of difference = 5.60, 95%

Credible Interval (CI): -0.60, 10.18, *pMCMC* = 0.06. Figure 1 & 2; Table S3). The female P_{bt}

of oviparous species is on average 4 °C higher than T_{opt} (raw data mean ± SD P_{bt} 102 species:

31.40 ± 4.31 ºC; T_{opt} 50 species: 27.25 ± 2.42 ºC). Incubation experiments have shown that

such a difference can significantly reduce hatching success, illustrating that the retention of

offspring throughout development may jeopardise embryo survival (15, 16); reviewed in

(17). P_{bt} and T_{opt} of oviparous species were both estimated to have high phylogenetic inertia

(Figure 1; MR-BPMM: P_{bt} phylo H^2: 0.99, CI: 0.91, 1.0. T_{opt} phylo H^2: 0.93, CI: 0.72, 0.98.

Table S3) and female P_{bt} and offspring T_{opt} only show weak evidence of coevolving (Figure

2; phylogenetic correlation (MR-BPMM): PM = 0.57, CI: -0.09, 0.97, *pMCMC* = 0.08. Table

S4), suggesting that this mismatch may be difficult to resolve.

Transitions to live birth occur despite mismatches between females and embryos

We found evidence that preferred body temperature was lower in viviparous, compared with

oviparous, lineages (MR-BPMM: PM = -0.20, CI: -0.43, -0.02, *pMCMC* = 0.02; Table S6).

This is consistent with the hypothesis that viviparity is an adaptation to cool climates (2, 3,

18). However, reconstructions of ancestral preferred body temperatures of adults indicated

that in the majority of lineages, lower preferred body temperatures evolved after, not before,

transitions to live bearing. This was evident from the lack of difference in P_{bt} between

ancestors of egg-laying and live-bearing lineages (MR-BPMM, PM = 0.04, CI: -3.96, 4.27,

pMCMC = 0.42; Table S7). Additionally, live-bearing lineages with egg-laying ancestors

showed higher P_{bt} than those with live-bearing ancestors (MR-BPMM, PM = 6.78, CI: -0.64,

8.74, *pMCMC* = 0.06; Table S7).

Ancestral reconstructions of *Pbt* and *Topt* of oviparous species showed that transitions

from egg laying to live bearing were no more likely to occur when adult and embryo thermal

optima were similar (% of lineages with thermal mismatches: ancestors of live bearing 84%,

ancestors of egg laying species 85%. Table S5). Only 5 of the estimated 31 evolutionary

origins of live birth had adult and embryo thermal optima that overlapped, which was similar

to the proportion of lineages where live bearing did not occur (Table S5). In the remaining 26

egg-laying ancestors of live-bearing species, the thermal optimum of embryos was

significantly lower than the preferred body temperature of adults (there were no cases of T_{opt}

being significantly higher than P_{bt}; Table S5). In summary, the degree of similarity between

adult P_{bt} and embryo T_{opt} does not predict transitions towards viviparity, and the lower P_{bt}

observed for viviparous taxa appears to have evolved after transitions to live birth, not before.

Female behavioural plasticity resolves the constraints imposed by thermal physiology

Next, we tested the hypothesis that the thermal mismatch between females and embryos

during the evolution of viviparity is resolved by females adjusting their body temperature

when gravid to meet the thermal requirements of their embryos (19). We collected data on

the preferred body temperature of gravid ($P_{bt\text{-}g}$) and non-gravid ($P_{bt\text{-}ng}$) adult females from 52

species of lizards and snakes were collected from published field and experimental studies

(live bearing n = 32, egg laying n = 20. Table S1). Using these data, an effect size of female

behavioural plasticity, Hedges' *g* was calculated (positive values = higher P_{bt} when gravid,

negative values = lower P_{bt} when gravid).

Across those 52 species, there was a negative phylogenetic correlation between P_{bt}

and the degree to which females reduced their body temperature when gravid (MR-BPMM:

PM = -0.29, CI: -0.62, -0.02, *pMCMC* = 0.02. Figure 4; Table S8). Females with high *Pbt*

have therefore evolved to behaviourally reduce their body temperature when gravid to a

greater extent than females with low *Pbt*. This relationship was not restricted to viviparous

lineages, which is expected if female behavioural plasticity is an adaptation that evolves in

response to selection for embryo retention. Instead, comparable phylogenetic correlations

between Hedges' *g* and *Pbt* were evident in both oviparous and viviparous taxa, showing that

behavioural adjustment of temperature when gravid occurs to a similar extent in both egg

laying and live bearing species (Egg laying: PM = -0.61, CI: -0.92, -0.08, *pMCMC* = 0.02.

Live bearing: PM = -0.88, CI: -0.96, -0.30, *pMCMC* = 0.005. Table S9).

  To further investigate the role of female plasticity in the evolution of live bearing, we

used phylogenetic models to reconstruct values of Hedges' *g* for the ancestors of egg-laying

and live-bearing species. Consistent with the results from our correlational analyses, we

found that, in the ancestors of both egg laying and live bearing species, thermal mismatches

between adults and embryos were resolved by females adjusting their body temperature when

gravid (Figure 4A; MR-BPMM: Egg-laying Hedges' *g* PM = -0.81, CI = -1.43, 0.05,

pMCMC = 0.03. Live bearing Hedges' *g* PM = -0.75, CI = -1.88, -0.09, *pMCMC* = 0.03.

Table S10). As a result, Hedges' *g* did not differ between the ancestors of oviparous and

viviparous species (Figure 4B; Table S9-S11), suggesting that behavioural plasticity was

likely present prior to the emergence of live birth.

  Our results illustrate that an ancient and evolutionarily conserved ability of egg laying

females to adjust their body temperature during pregnancy provides a general resolution to

the conflict over adult and embryo thermal optima that paves the way for the evolution of live

bearing. The down-regulation of female body temperature in egg-laying species while gravid

may appear surprising considering that most of these species lay their eggs within the early

stages of development (commonly around the time of limb bud formation (20)). However,

early developmental stages, involving gastrulation, neurulation and organogenesis, are

potentially even more sensitive to temperature stress than later stages, which are

predominantly associated with growth (8, 10). The temperature sensitivity of early-stage

embryos therefore suggests that resolutions to mother-offspring thermal conflicts are required

in both egg-laying and live-bearing species. If true, the key innovation of live birth may owe

its evolutionary origin to mechanisms of temperature regulation put in place long before live

birth emerged. We suggest that this behavioural flexibility of female lizards and snakes

enabled the gradual evolution of live birth by helping traverse potential fitness valleys

associated with embryo retention. In many live-bearing lineages, temperature mismatches

have subsequently been eliminated through the evolution of a lower preferred body

temperature in females. However, behavioural flexibility is frequently maintained in

viviparous lineages that have colonized cool climates, possibly as a means for mothers to

maintain embryos at temperatures significantly warmer than the external environment (2, 5,

6, 21, 22).

In conclusion, female thermoregulatory behaviour is a key adaptation that overcomes

the constraints imposed by thermal mismatches between mothers and embryos, and has

facilitated both the evolution of live birth, and expansion of reptiles into a variety of

environments.

Methods

Literature search and data collection

To investigate the relationship between maternal behavioural plasticity and embryo thermal

sensitivity, we collected existing data on reproductive mode (oviparous versus viviparous),

the female preferred body temperature (P_{bt}) ("P_{bt} dataset"), the temperature at which hatching

success was maximised (T_{opt}) ("T_{opt} dataset"), and female body temperature when gravid ($P_{bt\text{-}g}$)

$_g$) and not gravid ($P_{bt\text{-}ng}$) ("Hedges' g dataset"). Complete data are presented in Table S1. Data

were collected for each of the variables from published literature. We conducted a literature

search using ISI *Web of Science* (v.5.30) with search terms specific to each dataset.

The P_{bt} dataset provided preferred body temperatures (P_{bt}) of adult females measured at

one point in time. We used the search terms 'body temperature*', along with one of the

following: 'squamat*', 'lizard*', 'snake*' which yielded a total of 1075 papers. We only

used data from studies where P_{bt} for females was stated explicitly, unless pooled male/female

data stated no significant effect of sex. Data on females that were described to be gravid or

data collected during the reproductive season were excluded. This provided a final dataset of

female preferred body temperature for 163 species that was independent of the gravid and

non-gravid measures used to calculate Hedges' g (live bearing: n = 61 and egg laying: n =

102).

The T_{opt} dataset quantified the temperature that maximises embryonic survival (measured

as hatching success; T_{opt}) in 50 egg laying species. Hatching success data were taken from the

Reptile Developmental Database (23) and supplemented with additional studies using the

same method described for the P_{bt} dataset (search terms: 'temperature* AND incubat* AND

hatch* OR surv*', along with one of the following: 'squamat*', 'lizard*', 'snake*'), yielding

a total of 671 papers. We only included studies where three or more constant temperature

treatments were used under controlled laboratory conditions, resulting in 661 papers being

rejected due to irrelevance or overlap with the Reptile Development Database. Given that T_{opt}

had a strong phylogenetic signature (Phylogenetic Heritability (H^2) = 91.93%, 95% CI: 71.65

– 98.16%; Table S3) we fitted a Bayesian Phylogenetic Mixed Effects Model (BPMM) to

hatching success data for all species jointly. Including phylogenetic information allowed for

the T_{opt} of each species to be estimated with greater accuracy and precision given that the

range and number of temperatures across species varied (range: 10-40 ºC, mean number of

temperatures per species ± SD: 7.62 ± 4.29). Compared to non-phylogenetic models, T_{opt}

estimates produced from phylogenetic models (BPMM) showed smaller sampling error and

avoided convergence problems in estimating model parameters. Importantly, this approach

was not used to estimate T_{opt} data for species without data, only to better predict T_{opt} values

for species for which there were data. The T_{opt} BPMM model was run for 1,100,000 iterations

with a burn-in of 10,000 iterations and thinning rate of 500, leaving us with 2,000 samples

from the posterior distribution. Autocorrelation was low (lag values < 0.1) and trace plots

showed chains mixed well for all parameters. Our model included temperature as a fixed

effect (estimating both a linear and quadratic slopes) and random slopes of temperature

(linear and quadratic slopes) fitted at the phylogenetic level. From our BPMM we estimated

T_{opt}, and its corresponding sampling variance, using the posterior distribution of fixed effects

and best linear unbiased predictors (BLUPs) for the random slopes (linear and quadratic) for

each species as follows:

198
$$Topt = -\frac{\left(T_f + T_{sp}\right)}{\left(2\left(T_f^2 + T_{sp}^2\right)\right)}$$

Where T_f and T_f^2 are the posterior linear and quadratic fixed effect estimates for temperature

and T_{sp} and T_{sp}^2 are the posterior BLUPs for a given species extracted from the phylogenetic

random slopes. Calculating T_{opt} using the posterior distribution of fixed and random effects

meant that sampling error for a given species could be propagated to subsequent analyses

(see below).

For the final "Hedges' g" dataset, we collected mean, standard deviation and sample sizes

for female preferred body temperatures when gravid (P_{bt-g}) and non-gravid (P_{bt-ng})

(independent from the P_{bt} dataset) from the same study/population. Within the 'title',

'abstract' or 'keywords' we used search terms 'body temperature* AND gravid* OR

reproduct*' along with one of the following: 'squamat*', 'lizard*', 'snake*' which yielded a

total of 721 papers. We included both field and laboratory measures of female body

temperature, comprising studies that directly compared P_{bt-g} and P_{bt-ng}. Only studies that

provided both sample size and error around mean of preferred body temperature were

included. Of the 721 papers 648 papers were excluded due to irrelevance or insufficient data.

The majority of studies used artificial temperature gradients in the laboratory (n = 37) or

measured preferred basking temperature in the field (n = 36). Laboratory studies generally

measured body temperature in the same female during gestation (P_{bt-g}) and either before or

after gestation (P_{bt-ng}) as repeated measures. In contrast, field studies often measured body

temperature on a population during the reproductive season, comparing body temperatures in

gravid and non-gravid females at a single time point. Combined, this yielded a total of 73

studies published up to July 2019 from which effect sizes were calculated for 52 species (live

bearing: n = 32 and egg laying: n = 20). An effect size of female adjustment of body

temperature when gravid for each species was measured as the standardised mean differences

(Hedges' g) in preferred body temperatures between non-gravid and gravid females (P_{bt-g} -

P_{bt-ng}), adjusting for small sample sizes (24). We found no significant differences in means or

variances of Hedges' g between laboratory or field studies ("Study type").

The final full dataset (Table S1) contained a total of 224 squamate species. Two

species in our data are both oviparous and viviparous (*Saiphos equalis* and *Zootoca vivpara)*.

While for *S. equalis* all data were from an oviparous population, for *Z. vivipara* there were

data for both oviparous and viviparous populations and we treated these different

reproductive modes as being distinct in our dataset (25).

Quantification and Statistical Analysis

Data were analysed using Phylogenetic Generalised Least Squares (PGLS), Bayesian

Phylogenetic Mixed Effects Models with single (BPMM) and multiple response variables

(MR-BPMM), and Stochastic Character Mapping (SCM). All analyses were conducted in R

version 4.0.1 (26). Ancestral states were reconstructed by initially examining the mode of

evolution that best explained how traits have evolved (a model of Brownian motion best

explained all traits: Table S2), after which Bayesian Phylogenetic Mixed models (BPMM)

were used to simultaneously reconstruct ancestral states of all traits. The robustness of

ancestral reconstructions was examined using stochastic character mapping (SCM; Table

S12).

Assessing modes of evolution using Phylogenetic Generalised Least Squares (PGLS)

An assumption of many phylogenetic comparative methods is that traits evolve via a

Brownian motion process (27). This assumption was assessed by estimating the mode of

evolution that best describes the phylogenetic distribution of reproductive mode, *PBT* and

Topt using PGLS implemented with R packages 'ape' and 'nlme' (28, 29). This was done by

comparing PGLS models with different correlation structures constructed from the phylogeny

that corresponded to BM, Ornstein-Uhlenbeck (OU: corMartins in 'ape') and Accelerated

Decelerated (ACDC: corMartins in 'ape') modes of evolution. In OU models' traits undergo

a BM process but evolve towards a central tendency with a strength proportional to parameter

alpha (α), and in accelerated/decelerated models of evolution (ACDC) traits diverge under a

BM model, but rates of evolution accelerate or decelerate over time according to parameter g

(Acceleration: g < 1. BM: g = 1. Deceleration g >1). Estimating parameters, such as α and g,

directly from the data can be unstable (30, 31). Therefore, we estimated values of α (OU) and

Bloomberg's g (ACDC) by fixing α and g at 0.1 increments within the range of 0 to 10 and

used AIC values to identify best-fit models. For all traits there was significant phylogenetic

signature and BM best explained the way traits have evolved (Table S2: R code: Section 0).

Bayesian Phylogenetic Mixed Effects models (BPMM)

We implemented BPMMs in *R* with the *MCMCglmm* package (32). Unlike other generalised

linear mixed model packages, *MCMCglmm* can incorporate inverse phylogenetic correlation

matrices with missing data in response variables (i.e., trait values for both T_{opt} and P_{bt} were

not available for all species). BPMMs allow missing values to be predicted with an accuracy

related to the phylogenetic signature in traits and the strength of phylogenetic correlations

between traits. All traits had high phylogenetic signature (phylogenetic heritability >80%)

producing high correspondence between raw and predicted values (Figure S1). As a result,

our BPMMs enabled us to deal with the fact that not all traits have been measured in all

species.

Phylogenetic relationships were modelled using a published phylogeny comprising over 4000

species of squamates (33) that was pruned to the 224 species in our dataset using the *ape*

package in *R* (28). The random effect *animal*, with a covariance matrix derived from the

phylogenetic tree was included in all models (32). We calculated the phylogenetic signature

(equivalent to heritability, H^2, in the terminology of *MCMCglmm*) for each trait as the

variance explained by *animal* relative to total variance (animal and residual). In the case of

binary variables (reproductive mode) the residual variance is not identifiable and we

therefore used the intraclass coefficient of variation (ICC: phylogenetic variance /

(phylogenetic variance + residual variance +$pi^{2/3}$)*100) to examine phylogenetic signature

(34). More traditional methods were also used to validate estimates of phylogenetic signature.

Pagel's lambda (λ) from *pgls* models (35, 36) and Blomberg's K (*K*) in *phytools* using the

function *phylosig* (37, 38), which gave comparable estimates to BPMMs (see R code section

0).

Multi-response BPMMs (MR-BPMMs) can also be fitted with *MCMCglmm* that allow the

phylogenetic and within-species (residual) correlations between traits to be estimated.

Correlations between traits (e.g., A & B) were calculated as:

286
$$\frac{cov(A,B)}{\sqrt{(var(A) \cdot var(B))}}$$

Missing data imputation for missing sampling variances

The accuracy of measures of T_{opt}, P_{bt}, and Hedges' *g* varied across species due to study

design and sample sizes which can be accounted for by weighting data points by their inverse

sampling variance. However, missing values in sampling variances are not permitted in

MCMCglmm. As data on the error and sample size was missing for T_{opt}, P_{bt} and Hedges' *g* it

would not have been possible to account for sampling error in our analyses without

drastically reducing the size of our dataset. Consequently, we used multiple imputation with

predictive mean matching in the *mice* package in R to impute missing error and sample sizes

(39). To incorporate uncertainty in imputations, 20 complete datasets were generated, and all

analyses were conducted by sampling across these datasets. Each model sampled through the

20 datasets 75 times (1500 sampling events) for 2000 iterations with only the last iteration

being saved. Estimates from the last iteration of each sampling event *i* were used as the

starting parameter values for the next *i* + 1. This led to a total posterior sample of 1500

iterations, the first 500 iterations were discarded as a burn-in and the remaining 1000 (50 per

dataset) were used to estimate parameters. Pooling of posterior distributions from model

parameters from across each of the *m = 20* datasets enabled imputation uncertainty in

sampling variances to be accounted for.

Model convergence, prior settings and characterisation of posterior distributions

Non-informative uniform priors were used for fixed effects and inverse-Wishart priors for

random effects ($V = 1$, nu $= 0.002$; (32)) apart from models with binary response variables

(e.g. reproductive mode). For binary variables the residual variance was fixed to 1 and we

specified a fixed effect prior that is relatively flat on the logit scale (mu $= 0$, V $=$ residual

variance x $\pi^{2/3}$) and parameter-expanded random effect priors ($V = 1$, nu $= 0.002$,

alpha.mu$=0$, alpha.V $= 1000$). To examine model convergence we ran three independent

MCMC chains and examined autocorrelation, which was low (lag values < 0.1), trace plots,

which showed chains mixed well, and Gelman and Rubin's convergence diagnostic that

models converged (potential scale reduction factors were all below 1.1: R function

gelman.diag (40). The convergence of models of the probability of viviparity (binary

response variable) occurred more slowly and therefore were run for longer (burn-in $=$

1000000, iterations $= 6000000$, thinning $= 5000$). All other models used the run settings

specified in the data imputation section. Posterior distributions of all parameters were

characterised using modes and 95% credible intervals (CIs) throughout. Effects were

regarded as significant where CIs did not span 0. *pMCMC* (number of iterations above or

below 0 / total number of iterations) are also presented to facilitate general interpretation.

Specific analyses

We conducted six main analyses: **1)** Phylogenetic signature in P_{bt} and T_{opt} was estimated

using a MR-BPMM of P_{bt} and T_{opt} with heterogenous phylogenetic and residual covariance

matrices fitted as random effects (R code model M1.1. Table S3). **2)** The phylogenetic

correlation between P_{bt} and T_{opt} was estimated using a MR-BPMM with unstructured

phylogenetic and residual covariance matrices fitted as random effects (R code model M1.4.

Table S4). **3)** To quantify the number of ancestral egg-laying with and without mismatched

adult female and embryo thermal optima a two-step approach was used. The ancestral states

of reproductive mode were first estimated for each node in the phylogenetic tree using a

BPMM with the probability of live bearing as the response variable (R code model M3.0).

Next a MR-BPPM with P_{bt}, T_{opt} and Hedges' *g* as response variables was used (R code model

M3.1) to reconstruct ancestral states for each trait. From this model mismatches in thermal

optima (CI of $P_{bt} – T_{opt}$ not overlapping 0) and levels of female behavioural plasticity were

estimated (Table S5). Separate models were used for reproductive mode and P_{bt}, T_{opt} and

Hedges' *g* to ensure that ancestral estimates were not influenced by the strong correlation

between live bearing and P_{bt} in extant species. **4)** The phylogenetic correlation between the

probability of live bearing and P_{bt} was estimated using a MR-BPMM with unstructured

phylogenetic and residual covariance matrices fitted as random effects (R code model M2.1.

Table S6). **5)** To examine if female plasticity differed between the ancestors of oviparous and

viviparous lineages with and without mismatched mother-offspring thermal optima, we

examined estimates of Hedges' *g* in relation to predicted states of reproductive mode (R code

model M3.1. Table S7 & S10-S11). **6)** The phylogenetic correlation between P_{bt} and Hedges'

g was estimated across all species (R code model M3.1. Table S8), as well as for egg-laying

and live-bearing species separately using an MR-BPMM with separate unstructured

phylogenetic and residual covariance matrices for each reproductive mode (R code model

M4.1. Table S9).

Stochastic character mapping (SCM)

The robustness of ancestral state reconstructions of reproductive mode was examined by assessing the correspondence between the estimates from MCMCglmm with those obtained from SCM. The SCM approach calculates the conditional likelihood that each ancestral node is oviparous or viviparous based on an estimated transition rate matrix (Q) between reproductive modes and the length of the branch associated with that node. Based on these conditional likelihoods, ancestral states at each node are stochastically simulated and used in combination with observations at the tips to reconstruct a character history along each branch. Each character history is simulated using a continuous-time Markov model where changes between states and the time spent in each state is modelled as a Poisson process (see (41) for details). We found extremely good correspondence between estimates of ancestral states obtained from MCMCglmm models and SCM showing inferences of ancestral states were robustness to methodological details (Table S2).

Data and code availability

All data and code are publicly available at: https://doi.org/10.17605/OSF.IO/JT28V

References

1. J. S. Doody, J. A. Moore, Conceptual model for thermal limits on the distribution of reptiles. *Herpetological Conservation and Biology* **5**, 283–289 (2010).

2. R. Shine, "The evolution of viviparity in reptiles: An ecological analysis" in *Biology of the Reptilia*, Development B., (John Wiley & Sons, Ltd, 1985), pp. 605–694.

3. S. M. Lambert, J. J. Wiens, Evolution of viviparity: a phylogenetic test of the cold-climate hypothesis in Phrynosomatid lizards. *Evolution* **67**, 2614–2630 (2013).

4. R. Shine, A new hypothesis for the evolution of viviparity in reptiles. *The American Naturalist* **145**, 809–823 (1995).

5. D. A. Warner, R. Shine, Fitness of juvenile lizards depends on seasonal timing of hatching, not offspring body size. *Oecologia* **154**, 65–73 (2007).

6. M. Le Henanff, S. Meylan, O. Lourdais, The sooner the better: reproductive phenology drives ontogenetic trajectories in a temperate squamate (Podarcis muralis). *Biol J Linn Soc* **108**, 384–395 (2013).

7. L. Ma, L. B. Buckley, R. B. Huey, W.-G. Du, A global test of the cold-climate hypothesis for the evolution of viviparity of squamate reptiles. *Global Ecology and Biogeography* **27**, 679–689 (2018).

8. C. A. Beuchat, Temperature effects during gestation in a viviparous lizard. *Journal of Thermal Biology* **13**, 135–142 (1988).

9. C. Monasterio, L. P. Shoo, A. Salvador, P. Iraeta, J. A. Díaz, High temperature constrains reproductive success in a temperate lizard: implications for distribution range limits and the impacts of climate change. *Journal of Zoology* **291**, 136–145 (2013).

10. T. J. Sanger, J. Kyrkos, D. J. Lachance, B. Czesny, J. T. Stroud, The effects of thermal stress on the early development of the lizard Anolis sagrei. *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology* **329**, 244–251 (2018).

11. W.-G. Du, R. Shine, L. Ma, B.-J. Sun, Adaptive responses of the embryos of birds and reptiles to spatial and temporal variations in nest temperatures. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20192078 (2019).

12. R. M. Andrews, T. Mathies, D. A. Warner, Effect of Incubation Temperature on Morphology, Growth, and Survival of Juvenile Sceloporus undulatus. *Herpetological Monographs* **14**, 420–431 (2000).

13. D. G. Blackburn, Squamate Reptiles as Model Organisms for the Evolution of Viviparity. *Herpetological Monographs* **20**, 131–146 (2006).

14. D. G. Blackburn, Evolution of viviparity in squamate reptiles: Reversibility reconsidered. *J Exp Zool B Mol Dev Evol* **324**, 473–486 (2015).

15. R. Shine, Reptilian viviparity in cold climates: testing the assumptions of an evolutionary hypothesis. *Oecologia* **57**, 397–405 (1983).

16. R. M. Andrews, B. R. Rose, Evolution of Viviparity: Constraints on Egg Retention.
Physiological Zoology **67**, 1006–1024 (1994).

17. D. W. A. Noble, V. Stenhouse, L. E. Schwanz, Developmental temperatures and
phenotypic plasticity in reptiles: a systematic review and meta-analysis. *Biological*
Reviews **93**, 72–97 (2018).

18. C. M. Watson, R. Makowsky, J. C. Bagley, Reproductive mode evolution in lizards
revisited: updated analyses examining geographic, climatic and phylogenetic effects
support the cold-climate hypothesis. *Journal of Evolutionary Biology* **27**, 2767–2780
(2014).

19. C. A. Beuchat, Reproductive Influences on the Thermoregulatory Behavior of a Live-
Bearing Lizard. *Copeia* **1986**, 971–979 (1986).

20. R. M. Andrews, T. Mathies, Natural History of Reptilian Development: Constraints on
the Evolution of Viviparity. *BioScience* **50**, 227–238 (2000).

21. R. M. Andrews, Evolution of viviparity in squamate reptiles (Sceloporus spp.): a variant
of the cold-climate model. *Journal of Zoology* **250**, 243–253 (2000).

22. T. Uller, M. Olsson, Offspring size and timing of hatching determine survival and
reproductive output in a lizard. *Oecologia* **162**, 663–671 (2010).

23. D. W. A. Noble, *et al.*, A comprehensive database of thermal developmental plasticity
in reptiles. *Sci Data* **5**, 1–7 (2018).

24. M. Borenstein, L. V. Hedges, J. P. Higgins, H. R. Rothstein, "Effect Sizes Based on
Means" in *Introduction to Meta-Analysis*, (John Wiley & Sons, Ltd, 2009), pp. 21–32.

25. H. Recknagel, N. A. Kamenos, K. R. Elmer, Common lizards break Dollo's law of
irreversibility: Genome-wide phylogenomics support a single origin of viviparity and
re-evolution of oviparity. *Molecular Phylogenetics and Evolution* **127**, 579–588 (2018).

26. R Core Team, *R: A language and environment for statistical computing. R Foundation*
for Statistical Computing, Vienna, Austria. (2020).

27. L. Z. Garamszegi, Ed., *Modern Phylogenetic Comparative Methods and Their*
Application in Evolutionary Biology: Concepts and Practice (Springer-Verlag, 2014)
https:/doi.org/10.1007/978-3-662-43550-2 (December 4, 2020).

28. E. Paradis, K. Schliep, ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics* **35**, 526–528 (2019).

29. J. Pinheiro, D. Bates, S. DebRoy, D. Sarkar, R. C. Team, nlme: Linear and nonlinear
mixed effects models. *R package version* **3**, 111 (2013).

30. E. Paradis, *Analysis of Phylogenetics and Evolution with R*, 2nd Ed. (Springer-Verlag,
2012) https:/doi.org/10.1007/978-1-4614-1743-9 (December 4, 2020).

31. T. F. Hansen, Stabilizing selection and the comparative analysis of adaptation.
Evolution **51**, 1341–1351 (1997).

32. J. D. Hadfield, MCMC Methods for Multi-Response Generalized Linear Mixed Models:
The MCMCglmm R Package. *Journal of Statistical Software* **33**, 1–22 (2010).

33. R. A. Pyron, F. T. Burbrink, J. J. Wiens, A phylogeny and revised classification of
Squamata, including 4161 species of lizards and snakes. *BMC Evolutionary Biology* **13**,
93 (2013).

34. S. Nakagawa, P. C. D. Johnson, H. Schielzeth, The coefficient of determination R2 and
intra-class correlation coefficient from generalized linear mixed-effects models revisited
and expanded. *Journal of The Royal Society Interface* **14**, 20170213 (2017).

35. D. Orme, *et al*., CAPER: comparative analyses of phylogenetics and evolution in R.
Methods in Ecology and Evolution **3**, 145–151 (2013).

36. R. P. Freckleton, P. H. Harvey, M. Pagel, Phylogenetic analysis and comparative data: a
test and review of evidence. *Am. Nat.* **160**, 712–726 (2002).

37. L. J. Revell, phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution* **3**, 217–223 (2012).

38. S. P. Blomberg, T. Garland, A. R. Ives, Testing for Phylogenetic Signal in Comparative
Data: Behavioral Traits Are More Labile. *Evolution* **57**, 717–745 (2003).

39. S. van Buuren, K. Groothuis-Oudshoorn, mice: Multivariate Imputation by Chained
Equations in R. *Journal of Statistical Software* **45**, 1–67 (2011).

40. S. P. Brooks, A. Gelman, General methods for monitoring convergence of iterative
simulations. *Journal of computational and graphical statistics* **7**, 434–455 (1998).

41. J. P. Bollback, SIMMAP: Stochastic character mapping of discrete traits on
phylogenies. *BMC Bioinformatics* **7**, 88 (2006).

Figure 1. Variation in reproductive mode, preferred body temperature and plasticity in

female thermo-regulatory behaviour across 224 species of squamate reptile (lizards and

snakes). Tips and branches are coloured according to reproductive modes (red = live

bearing/viviparous, grey = egg laying/oviparous; branch colours represent predicted ancestral

values; Table S10). The size of white circles corresponds to the preferred body temperature

of female squamates (P_{bt} dataset, 163 species), the size of grey circles to the optimal

temperature for embryo survival in oviparous species (T_{opt} dataset, 50 species), and the size of

the black circles to the discrepancy between preferred body temperature in gravid versus non-

gravid females (Hedges' *g* dataset; absolute values, 52 species).

Figure 2. Mismatches in the thermal optima of mothers and embryos across 152

oviparous species of lizards and snakes. The relationship between female preferred body

temperature (P_{bt}) and the temperature that maximizes hatching success of embryos (T_{opt})

predicted by a multi-response Bayesian Phylogenetic Mixed Model across egg laying species

(regression lines ± 95% confidence intervals are plotted). Deviations from the dotted line (1:1

relationship) indicate the magnitude of the mismatch between adult and embryo thermal

optima. Corresponding measurements of T_{opt} and P_{bt} were not available for all species (n_{Pbt} =

103; n_{Topt} = 48; n_{Pbt} & n_{Topt} = 12), hence model predictions rather than raw data are shown (for

plot of raw data as well as correspondence between predicted and raw data values see Figure

S1).

Figure 3. Probability of being a live-bearing species in relation to female preferred body temperature. The line and shading represent a logistic regression curve ± 95% confidence interval.

Figure 4. Female behavioural plasticity resolves the constraints imposed by thermal physiology, such that transitions to live-bearing (viviparity) occur despite mismatches between females and embryos. (A) Adjustment of body temperature in gravid females in relation to their non-gravid preferred body temperature (P_{bt}) across 52 extant oviparous and viviparous lizards and snakes (Hedges' g; $P_{bt\text{-}g}$ - $P_{bt\text{-}ng}$). Regression lines ± 95% credible intervals are plotted. High values of Hedges' g signify an increase in body temperature when gravid while low values imply a decrease. (B) The adjustment of body temperature by females in lineages that remained egg laying (top panel) and lineages that evolved live bearing (bottom panel) according to the presence or absence of a thermal mismatch between females and their offspring (inferred from Hedges' g data shown in panel A). Abbreviations: Ovi, oviparous (egg laying); Vivi, viviparous (live bearing).