Kaposi's sarcoma-associated herpesvirus (KSHV) utilizes the NDP52/CALCOCO2 selective autophagy receptor to disassemble processing bodies

Short title: Selective targeting of processing bodies for autophagy by KSHV

inflammatory cytokine transcripts / Kaposi's sarcoma-associated herpesvirus / NDP52 / processing bodies / selective autophagy

Carolyn-Ann Robinson[1,2,3,4,5], Gillian K. Singh[1,5], Mariel Kleer[2,3,4], Elizabeth L. Castle[1,3], Bre Q. Boudreau[1], Jennifer A. Corcoran[1,2,3,4*]

[1]Department of Microbiology & Immunology, Dalhousie University, Halifax, NS, Canada
[2]Microbiology, Immunology & Infectious Diseases Department, University of Calgary, Calgary, AB, Canada.
[3]Charbonneau Cancer Research Institute, University of Calgary, Calgary, AB, Canada.
[4]Snyder Institute for Chronic Diseases, University of Calgary, AB, Canada.
[5]Equal Contribution
[*]Corresponding Author

<u>Abstract</u>

Kaposi's sarcoma-associated herpesvirus (KSHV) causes the inflammatory and angiogenic

endothelial cell neoplasm, Kaposi's sarcoma (KS). We previously demonstrated that expression

of the KSHV protein, Kaposin B (KapB), promoted inflammation via the disassembly of

cytoplasmic ribonucleoprotein granules called processing bodies (PBs). Processing bodies (PBs)

modify gene expression by silencing or degrading labile messenger RNAs (mRNAs) including

many transcripts that encode inflammatory or angiogenic proteins that are associated with KS

disease. Although our work implicated PB disassembly as one of the causes of inflammation

during KSHV infection, the precise mechanism used by KapB to elicit PB disassembly was

unclear. Here we reveal a new connection between the degradative process of autophagy and PB

disassembly. We show that both latent KSHV infection and KapB expression enhanced

autophagic flux via the phosphorylation of the autophagy regulatory protein, Beclin 1. KapB was

necessary for this effect, as infection with a recombinant virus that does not express the KapB

protein did not induce Beclin 1 phosphorylation or autophagic flux. Moreover, we reveal that PB

disassembly mediated by KSHV or KapB depended on canonical autophagy genes and the

selective autophagy receptor *NDP52/CALCOCO2* and that the PB scaffolding protein, Pat1b, co-

immunoprecipitated with NDP52. These studies reveal a new role for autophagy and the

selective autophagy receptor NDP52 in promoting PB turnover and the concomitant synthesis of

inflammatory molecules during KSHV infection.

Author Summary

Kaposi's sarcoma-associated herpesvirus (KSHV) is the causative agent of the inflammatory,

endothelial cell cancer, Kaposi's Sarcoma (KS). KSHV induces a pro-tumourigenic

inflammatory environment which aids in the establishment and maintenance of the KS lesion.

Processing bodies (PBs) are cellular structures that dampen inflammatory gene expression by

suppression or decay of their cognate RNA molecules. We previously showed that the viral

protein KapB caused PB disappearance during KSHV infection, identifying a new pathway used

by KSHV to elicit inflammation. Now, we show that KSHV and KapB hijack the cellular

degradative process of autophagy to promote PB disassembly and increase inflammatory gene

expression. This places autophagy as central to the regulation of inflammation by KSHV and

illustrates another remarkable strategy used by these viruses to create a tumourigenic

microenvironment.

Introduction

<u>Introduction</u>

Macroautophagy, hereafter referred to as autophagy, is an evolutionarily conserved degradative

process that ensures the continuous recycling of material to maintain cellular homeostasis (1–3).

Autophagy begins with the formation of a phagophore that elongates and encloses a portion of

the cytoplasm to form a double-membrane organelle, the autophagosome, which fuses with

lysosomes to form autolysosomes, where contents are degraded (4). Under basal conditions,

autophagic activity ensures the clearance of damaged macromolecular complexes and organelles,

acting as an intracellular quality control system to maintain homeostasis (2). Various changes to

the cellular microenvironment, including nutrient scarcity, hypoxia, and endoplasmic reticulum

(ER) stress can upregulate autophagy, reprioritizing resources and promoting cell survival during

adverse and stressful conditions (2).

Although previously considered a non-selective bulk degradative pathway of cytoplasmic

recycling, recent work shows differential cargo targeting by autophagy through a process known

as selective autophagy (5,6). Autophagy can selectively target specific cellular components,

including organelles (e.g mitochondria, peroxisomes, ER) and protein complexes (e.g.

ribosomes, focal adhesion complexes, protein aggregates) for degradation (5,7). In selective

autophagy, target cargo is either selected directly, by binding to lipidated microtubule associated

protein 1 light chain 3 (LC3) protein (LC3-II)/Atg8, or via a receptor that mediates cargo

recognition. These molecular bridges are termed selective autophagy receptors (SARs), and their

role is to link the selected cargo to lipidated LC3-II molecules which are anchored to the

expanding phagophore (8). Some of the best characterized SARs include p62/SQSTM1, valosin

containing protein (VCP), optineurin (OPTN), neighbor of BRCA1 gene 1 (NBR1), nuclear dot

protein 52 (NDP52/calcium binding and coiled-coil domain protein, CALCOCO2) and its

paralogue TAX1BP1/CALCOCO3 (5,9,10). Of these, OPTN, p62, NDP52, and NBR1 are

members of the sequestosome-like family and contain several conserved motifs including a LC3-

interacting region (LIR) and a ubiquitin-binding domain (UBD) to mediate cargo selection via

binding ubiquitin chains present on the cargo surface (6,11–13).

While SARs play important homeostatic roles maintaining, and at times magnifying, the turnover

of organelles and aggregates via cargo selection, emerging functions of these receptors is to

localize the initiation of autophagy and modulate cellular signalling. SARs p62 and NDP52 both

promote autophagosome formation at the cargo site through direct interactions with autophagic

initiating complexes (14–16). SARs also play important roles in fine-tuning innate immune

responses by targeting key signaling platforms for degradation (17–22). For example, NDP52 is

required for the autophagic degradation of interferon regulatory factor 3 (IRF3) and

mitochondrial antiviral signaling protein (MAVS), which supports the resolution phase of type I

interferon signaling (19,23,24). Selective autophagy can also degrade invading pathogens such

as viruses in a process called xenophagy (25). Both proviral and antiviral roles have been

attributed to SARs (26,27) and viruses can both enhance or limit SAR function accordingly,

promoting their replication (28).

Kaposi's sarcoma-associated herpesvirus (KSHV) is the infectious cause of two B-cell

malignancies and the endothelial cell cancer, Kaposi's sarcoma (KS), which features aberrant

angiogenesis, tortuous leaky blood vessels, and inflammation (29–31). Infection with KSHV has

two forms, a latent phase and a lytic phase and it is the latent phase that predominates in KS

tumours (32–34). Like many viruses, KSHV manipulates the autophagic machinery. During

latency, the viral FLICE inhibitory protein (v-FLIP) binds Atg3 to suppress the LC3 lipidation

process during autophagosome biogenesis (35). KSHV also expresses a viral cyclin D homolog

(v-cyclin), that triggers DNA damage leading to autophagy and cellular senescence (36),

facilitating the production and non-conventional secretion of pro-inflammatory cytokines/

chemokines such as IL-1β, IL-6 and CXCL8 (37). KSHV manipulation of autophagic flux

prevents xenophagy, limits interferon responses, and enhances inflammatory cytokine

production. Precisely how KSHV fine-tunes autophagy during latency to balance proviral and

antiviral activities and promote inflammation and chronic viral infection is not clear.

Processing bodies (PBs) are ubiquitous, cytoplasmic, ribonucleoprotein (RNP) granules that

regulate expression of many cytokine RNA transcripts, making them important sites of

inflammatory control. PBs contain enzymes involved in mRNA turnover, including those that

mediate mRNA decapping (Dcp2; co-factors Dcp1a, Pat1b and Edc4/Hedls), 5'-3'

exonucleolytic degradation (5'-3' exonuclease Xrn1 and RNA helicase Rck/DDX6) and some

components of the RNA-induced silencing complex (28,38–44). mRNAs are routed to PBs for

decay and/or repression by RNA-binding proteins (RBPs) through recognition of common

sequence elements (39). AU-rich elements (AREs) are destabilizing RNA regulatory elements

found in the 3' untranslated regions (UTRs) of ~8% of cellular transcripts that are responsible for

targeting RNAs to PBs (45–47). When bound by destabilizing RBPs, ARE-containing RNAs are

directed to PBs and suppressed (48–50). Our group and others showed that PB size and numbers

correlate with the constitutive turnover or translational suppression of ARE-mRNAs; when PBs

are dispersed, this suppression is reversed (51–55). Because ARE-mRNAs encode potent

regulatory molecules such as inflammatory cytokines, PBs are important post-transcriptional

sites of regulation that fine-tune the production of inflammatory cytokines whose transcripts

contain AREs, including IL-6, CXCL8, IL-1β, and TNF (46,56–62).

PBs are dynamic structures that are continuously assembled and disassembled, yet relatively

little is known about how their activity is regulated. Stimuli that activate the p38/MK2 MAP

kinase pathway as well as many virus infections elicit PB disassembly and a concomitant

reduction in ARE-mRNA suppression to promote inflammatory molecule production

(51,52,55,63–65). We have previously shown that KSHV causes PB disassembly during latent

infection, and that the viral protein Kaposin B (KapB) induces PB disassembly while enhancing

the production of an ARE-containing reporter (52,63,66). These data support the notion that PB

disassembly is likely an important contributor to inflammation associated with KS (52,67).

Although we know that KapB binds and activates the kinase MK2, and that MK2 is an important

component of the mechanism of KapB-mediated PB disassembly and ARE-mRNA stabilization,

we do not precisely understand how PB loss is elicited by KapB (52,68). The observation that

MK2 can phosphorylate the autophagy regulatory protein Beclin 1 to increase autophagic flux in

response to nutrient scarcity (69) suggested to us that KapB may drive autophagic flux through

MK2. We now show that the KSHV KapB protein enhances autophagic flux via the

phosphorylation of Beclin 1, and that KSHV infection also enhances autophagic flux and Beclin

1 phosphorylation in a KapB-dependent manner, revealing an additional layer of complexity in

viral regulation of autophagy during KSHV latency. We show that both KSHV infection and

KapB alone require autophagy proteins including the selective autophagy receptor, NDP52, to

induce PB disassembly and that KapB requires NDP52 to elevate selected ARE-containing

mRNA transcripts. PB turnover is likewise elicited by the mTOR inhibitor and potent autophagy

inducer, Torin-1, and this process is also dependent on Atg5 and NDP52. Finally, we show that

NDP52 co-immunoprecipitates with the PB scaffolding protein, Pat1b, providing a molecular

basis for the recognition of a PB component by NDP52. These data forge a new connection

between the selective autophagy receptor NDP52 and cytoplasmic PBs and show that selective

autophagy of PBs is promoted by the viral protein KapB during KSHV infection.

Results

KapB induces autophagic flux

KapB expression activates the stress-responsive kinase MAPKAPK2 (MK2) (52,68). Since MK2

phosphorylates the essential autophagy inducer Beclin 1 in response to amino acid starvation

(69–71), we questioned whether KapB expression could promote autophagic flux. Essential for

autophagosome expansion is the lipidation of LC3 to form LC3-II, permitting its incorporation

into autophagosome membranes where it is degraded along with autophagosomal contents in the

final step of autophagy (72). Since p62 is a selective autophagy receptor (SAR) that delivers

target cargo to the autophagosome, p62 is also degraded during the final step of autophagy. If

autophagic flux is elevated, blocking autophagic flux with a lysosomal inhibitor should result in

an accumulation of both p62 and LC3-II (73). To measure autophagic flux, we examined levels

of LC3-II and p62 with and without treatment with the lysosomal inhibitor, Bafilomycin A1

(BafA1). Human umbilical vein endothelial cells (HUVECs) were transduced with lentiviral

vectors that express KapB or an empty vector control and treated for increasing times with

BafA1. As expected, LC3-II and p62 accumulated in the presence of BafA1 treatment, and both

proteins showed higher accumulation after BafA1 treatment in KapB-expressing cells than

equivalently treated vector controls (Fig 1A), indicating that autophagic flux was enhanced in

KapB-expressing cells compared to vector controls. When KapB-expressing cells were treated

for 4 h with BafA1, the amount of LC3-II and p62 increased two- and four-fold, respectively

(Fig 1A). We also measured the formation of LC3 puncta as an indicator of autophagic flux by

staining KapB-expressing cells treated with BafA1 for endogenous LC3 (74,75). Untreated

KapB-expressing HUVECs displayed LC3 puncta area that was similar to that of control cells;

however, LC3 puncta area significantly increased after BafA1 treatment in KapB-expressing

cells and not in control cells, further suggesting that KapB expression enhanced autophagic flux

(Fig 1B).

KapB-mediated PB disassembly requires Atg5 and Atg14

Our group previously described that MK2 activation was an important component of KapB-

mediated PB loss (52); however, these experiments did not reveal the precise mechanism of PB

loss. Since PBs are dynamic RNP granules, we wanted to determine whether KapB-mediated PB

loss was a result of enhanced disassembly of PBs or prevention of their *de novo* assembly. To do

so, we utilized HeLa cells that express a Dox-inducible GFP-tagged version of the PB-resident

protein, Dcp1a (41). When fixed, GFP-positive puncta co-stain with endogenous PB proteins

such as the RNA helicase DDX6/Rck and the decapping co-factor Hedls/Edc4, indicating that

they represent PBs (Fig S1). KapB was transfected into these cells either before or after inducing

GFP-Dcp1a granule formation. When KapB was expressed prior to granule induction, Dcp1a

puncta formed, although appearance of GFP-positive puncta was delayed compared to vector

control cells (Fig S1A). In the presence of KapB, *de novo* PB assembly was not blocked;

however, when KapB was expressed after PB induction, GFP-positive puncta were lost (Fig

S1B), indicating that KapB expression induced PB disassembly. As further evidence that KapB

does not prevent PB formation, we treated KapB-expressing HUVECs with sodium arsenite, a

known inducer of PB assembly (76). Sodium arsenite induced endogenous PB formation in

KapB-expressing HUVECs was not significantly different from equivalently treated control cells

(Fig 1C). Taken together, these results show that KapB expression caused PB disappearance by

inducing PB disassembly but did not block PB assembly. We speculated that KapB may be

utilizing a normal cellular pathway that mediates the turnover of RNPs or their components, such

as autophagy.

To determine if autophagy was required for KapB-mediated PB disassembly, we first inhibited

autophagic flux by independently silencing two genes in the macroautophagy pathway, Atg5 and

Atg14 (Fig S2A, B). When either Atg5 or Atg14 were silenced in KapB-expressing HUVECs,

PBs were restored, whereas their silencing did not have the same effect in control cells (Fig 2A

and B). We then examined the ability of KapB to mediate PB disassembly in Atg5 -/- mouse

embryonic fibroblasts (MEFs), which (77) lack conversion of LC3-I to LC3-II (Fig S2C). KapB

failed to disassemble PBs in Atg5 -/- MEFs but did disassemble PBs in matched wild-type

controls (Fig S2D). Moreover, KapB expression resulted in a significant increase in Atg5 protein

levels in HUVECs (Fig S2A), consistent with our other data that show that KapB promotes

autophagy. Additionally, blocking autophagic degradation with BafA1 also restored PBs in

KapB-expressing cells (Fig 2C). Together, these data indicate that KapB mediates PB

disassembly by enhancing autophagy.

Autophagy, although always occurring at a basal level, can be upregulated by many different cell

stimuli including starvation or inhibition of mTOR using Torin-1, hereafter referred to as Torin

(78). It was previously reported that rapamycin, an inhibitor of mTORC1, caused a loss of PBs in

mammary epithelial cells (79). We reasoned that if we inhibited both mTOR complexes

mTORC1 and mTORC2 using Torin (80) to induce autophagy, PBs would disassemble. We

observed that the number of Hedls/Edc4 puncta were significantly reduced after Torin treatment,

indicating that chemical induction of autophagy also resulted in the disappearance of PBs (Fig

3A). To ensure that Torin treatment was inducing disassembly of PBs and not preventing

assembly, HUVECs were treated with sodium arsenite to induce PB formation; like KapB, Torin

treatment did not prevent the assembly of PBs under these conditions (Fig 3B).

Given these similarities, we wondered if KapB induced autophagy via inhibition of mTOR

complexes mTORC1 and mTORC2. To test this, we performed immunoblotting for the

phosphorylated form of the ribosomal protein S6 (Serine 235/236) (81) and the phosphorylated

forms of the Unc-51 like autophagy activating kinase (ULK1; Serine 757/555) component of the

ULK autophagy initiation complex, both of which are dephosphorylated when mTOR activity is

inhibited by Torin (82,83). Unlike Torin, KapB expression did not decrease phosphorylation of

S6 (Fig 3C) or ULK (Fig 3D). We also expressed a constitutively active form of MK2 in which

two threonine residues that are normally phosphorylated by the upstream kinase p38, have been

substituted with glutamic acid residues (T205E/T317E, called MK2EE) (84–86) and determined

the effect on phospho-S6 and phospho-ULK. We previously demonstrated that MK2EE caused

PB disassembly (52) and others showed it increased autophagic flux after starvation (69). Like

KapB, MK2EE expression did not decrease phosphorylation of S6 (Fig 3C) or ULK (Fig 3D).

These data indicate KapB does not inhibit mTOR activity to induce autophagy. Since KapB can

bind and activate MK2, we next wondered if KapB expression would mediate phosphorylation of

Beclin 1 at Serine 90, as shown by others for MK2EE (68). Like MK2EE, KapB induced

phosphorylation of Beclin 1 at Serine 90 (Fig 3E), indicating that induction of autophagy via

KapB expression is mediated by activation of Beclin 1. Taken together, these data show that

although both KapB expression and Torin treatment caused PB disassembly in an autophagy-

dependent manner, the mechanism used to upregulate autophagic flux differed in each case.

Canonical autophagy is required for KapB-mediated increases in ARE-containing cytokine transcripts

PBs are nodes of post-transcriptional mRNA regulation and microscopically visible PBs are associated with ARE-mRNA decay or suppression (43,51,52,54,55,87,88). However, in untreated HUVECs, the transcription of most ARE-mRNAs is minimal. To activate transcription of ARE-containing cytokines in a biologically relevant manner, rather than using common inflammatory stimuli such as the bacterial product, lipopolysaccharide, we treated KapB-expressing or control HUVECs with 0.22 μm-filtered conditioned media from KSHV latently infected cells. Conditioned media from latent KSHV cells would be expected to contain inflammatory cytokines, e.g. TNF, that are elevated in KS, without containing virions (89). We then examined levels of some endogenous ARE-containing RNA transcripts in KapB-expressing cells (Fig 4A). We observed significant increases of cytokine mRNA for IL-6 and IL-1β in KapB-expressing cells compared to equivalently treated controls, suggesting these increases resulted from reduced transcript decay in PBs (Fig 4A). In addition, treatment with Torin promoted the enhanced steady-state levels of IL-1β and COX-2 (Fig 4B), suggesting that inducing autophagy reduced the decay of some ARE-mRNA transcripts. To support these findings and to simplify further studies, we utilized a luciferase reporter assay that we developed, described in (66). Using this assay, we previously showed that KapB-mediated PB loss correlated with enhanced luminescence of an ARE-containing firefly luciferase (FLuc) reporter because its rapid turnover was reversed and FLuc translation was enhanced (52,66). Briefly, HeLa cells were co-transfected with a FLuc construct containing the destabilizing AU-rich element from the *CSF2* gene in its 3'UTR and a Renilla luciferase (RLuc) construct with no ARE; transcription of both reporter constructs was then stopped by the addition of doxycycline.

In control (vector transfected) cells, the FLuc-ARE mRNA decayed rapidly and FLuc

luminescence relative to the RLuc transfection control was low. Torin treatment caused a

significant increase in FLuc/RLuc relative luminescence compared to DMSO-treated control

cells, supporting our RT-qPCR findings (Fig 4C) and suggesting that enhanced autophagy

reverses the constitutive turnover/suppression of the ARE-mRNA reporter. Furthermore, KapB

expression also caused a significant increase in FLuc/RLuc relative luminescence compared to

empty vector control cells (Fig 4D) as previously shown in (52). When canonical autophagy was

perturbed by either Atg5 or Atg14 silencing (Fig S3A, B) KapB-expressing cells displayed

significantly decreased luminescence compared to KapB-expressing controls (Figs 4D, E).

Likewise, when KapB-expressing cells were treated with BafA1, relative luminescence was

significantly decreased compared to KapB-expressing untreated control cells (Fig 4F). Together,

these data showed that canonical autophagy is required for KapB-mediated increases in cytokine

mRNA levels and the enhanced expression of an ARE-mRNA reporter.

Dcp1a protein levels are decreased by KapB and Torin treatment

Both KapB and Torin mediated PB disassembly but did not prevent *de novo* PB formation. We

reasoned that KapB and Torin could have this effect because they cause the autophagic

degradation of the entire PB granule. Alternatively, since some PB proteins have important

scaffolding roles that hold granules together (EDC4/Hedls, Xrn1, DDX6/Rck, Pat1b or Dcp1a

(90)), PB disassembly could be induced by KapB or Torin if one or more key PB proteins was

missing due to accelerated degradation. To differentiate between these two scenarios, we

immunoblotted for PB proteins after Torin treatment and in the context of KapB expression (Fig

5). Immunoblotting revealed that steady-state levels of Dcp1a were reduced in Torin-treated (Fig

5A) and KapB-expressing cells (Fig 5B); however, steady-state levels of the other PB-resident

proteins we tested were not affected (Fig 5A, C). Moreover, KapB-mediated Dcp1a decreases

were reversed by BafA1 treatment (Fig 5B) and steady-state levels of the SAR, p62, included as

a measure of autophagic flux, were likewise restored (Fig 5B). Together, these data suggest that

autophagy induction with Torin or KapB expression promote the accelerated degradation of the

PB-resident protein Dcp1a, but not all PB proteins. Silencing the canonical autophagy gene,

Atg5, resulted in a reversal of KapB-mediated Dcp1a loss (Fig 5D), suggesting that autophagy is

required for the loss of Dcp1a observed in KapB-expressing cells.

KapB-mediated PB disassembly and ARE-mRNA stabilization require the selective

autophagy receptor NDP52

After establishing a link between KapB expression, autophagy, and PB disassembly, we

wondered what specific autophagy mechanisms were at work and if selective autophagy

receptors (SARs) were involved. One previous study observed the SAR, NDP52, partially co-

localized with some PB proteins (91). For this reason, we used RNA silencing to investigate

whether NDP52, p62, VCP, NBR1, or OPTN were required for KapB-mediated PB disassembly.

Despite trying several different shRNA sequences, our attempts to silence NBR1 were

unsuccessful, and shRNAs targeting VCP resulted in significant HUVEC toxicity (Fig S4A, B).

Therefore, the role of these molecules in KapB-mediated PB disassembly could not be

determined. Knockdown of NDP52, p62, and OPTN was validated in control and KapB-

expressing cells (Fig S4C-E). Silencing of p62 or OPTN did not restore PBs in KapB-expressing

cells whereas NDP52 knockdown significantly restored PBs (Fig 6A, Fig S4F). These data

pinpointed an important role for NDP52 and selective autophagy in KapB-mediated PB

disassembly. We then tested whether NDP52 silencing could prevent Torin-mediated PB

disassembly and found that PBs could not be disassembled in cells lacking NDP52 (Fig 6B,

S4G). Since steady-state levels of the decapping cofactor, Dcp1a, decreased in response to KapB

infection and Torin treatment (Fig 5), we examined the steady-state level of Dcp1a after Torin

with and without NDP52 silencing (Fig 6C). Although Dcp1a was significantly diminished after

Torin treatment, NDP52 silencing did not restore Dcp1a to control levels. We decided to

consider another PB protein, Pat1b, a decapping cofactor known to interact with several other PB

proteins including Dcp1a, DDX6, and Hedls, suggesting that Pat1b is an important scaffold for

PB components (44). Immunoblotting revealed that Torin treatment significantly diminished

steady-state levels of Pat1b, and that this decrease was restored to control levels when Torin

treatment was performed in NDP52-silenced cells (Fig 6C). These data suggest that levels of the

PB component proteins Pat1b, and to a lesser extent, Dcp1a, are regulated by NDP52-dependent

autophagic degradation.

We also examined whether these SARs were required for KapB-enhanced relative luminescence

in our ARE-mRNA reporter assay. We validated silencing of NDP52, p62, and OPTN in HeLa

cells (Fig S5A-C). We found that p62 or OPTN knockdown had no effect on KapB-mediated

ARE-containing reporter luminescence, but NDP52 silencing decreased the signal compared to

non-targeting, KapB-expressing controls (Fig 6D), further supporting the role of NDP52 in

autophagy-mediated PB disassembly. We then performed an NDP52 complementation

experiment. We first silenced endogenous NDP52 in control or KapB-expressing HUVECs using

shRNAs targeting the 3'UTR of NDP52 and verified its knockdown (Fig S6A). Next, NDP52

expression was restored by overexpressing a fluorescently tagged NDP52 construct.

Complementation of NDP52-silenced HUVECs with NDP52 restored KapB-mediated PB

disassembly compared to vector controls (Fig 6E-F; S6B, C). Notably, complementation with

NDP52 in control cells that lacked KapB failed to induce PB disassembly, suggesting that KapB-

or Torin-mediated autophagy increase is an important component of the PB disassembly

phenotype. Taken together, these data indicate that NDP52 is required for KapB- and Torin-

mediated PB disassembly and KapB-mediated ARE-mRNA stabilization.

Atg5 and NDP52 are required for KSHV-mediated PB disassembly

Autophagy regulation during KSHV latency is complex, as the viral gene products v-FLIP and v-

cyclin also modulate autophagic flux in addition to KapB (35,36). To simplify our study of

autophagy and PBs, we initially used a reductionist approach, expressing only KapB in

HUVECs. However, to confirm the relevance of our study to KSHV infection, we performed a

study in KSHV-infected HUVECs. We infected HUVECs with rKSHV.219 (92) for 96 hours to

permit latency establishment and then treated or with either BafA1 or a vehicle control for 30

minutes prior to staining infected cells for LC3 puncta as a measure of autophagic flux. We

observed a dramatic increase in LC3 puncta in KSHV infected cells after BafA1 treatment when

compared to mock infection controls (Fig 7A). These data demonstrate that autophagic flux was

upregulated during KSHV latency despite the concurrent expression of vFLIP, an inhibitor of

autophagy, during this infection phase.(35). In parallel, we immunoblotted infected cell lysates

and observed that KSHV increased phosphorylation of Ser90 of Beclin 1 (Fig 7B), consistent

with earlier observations of Ser90 phosphorylation after ectopic expression of KapB (Fig 3D).

To confirm if KapB was necessary for upregulated autophagic flux after KSHV infection, we

used a recombinant KSHV that does not express KapB (delB BAC16) and matched wild-type

control (BAC16 KSHV), described in our recent publication (93). We infected HUVECs with

WT BAC16 KSHV or delB BAC16 virus for 96 hours to permit latency establishment. WT

BAC16 KSHV latency also induced LC3 puncta formation and phosphorylation of Ser90 of

Beclin 1, consistent with our result using the KSHV.219 virus, whereas infection with delB

BAC16 virus failed to enhance autophagic flux or phosphorylation of Beclin 1 at Ser90 (Fig 7C,

D). We confirmed these data in iSLKs, a KSHV producer cell line known for tight control over

latency (92). Naïve iSLKs were infected with WT BAC16 KSHV or delB BAC16 virus for one

week to permit latency establishment, and LC3-II accumulation in the presence of BafA1

treatment was determined by immunoblotting (Fig 7E). LC3-II showed higher accumulation after

BafA1 treatment in WT KSHV-infected iSLK cells than those infected with delB BAC16 virus

(Fig 7E), confirming that autophagic flux was enhanced in a KapB-dependent manner during

KSHV latent infection of iSLK cells. Together, these data validate a necessary role for KapB in

KSHV enhancement of autophagic flux and show that during this KSHV infection phase,

autophagy proceeds via a non-canonical activation mechanism that involves the phosphorylation

of Ser90 in Beclin 1 (69).

In our previous work, we showed that KSHV infection induced PB disassembly (52) and more

recently we confirmed that KapB was not only sufficient but also necessary for this effect with

the delB BAC16 virus (93) . To determine if autophagic machinery is also required for PB

disassembly in the context of KSHV infection, we silenced *Atg5* or *NDP52* in HUVECs prior to

infecting with rKSHV.219 for 96 hours to establish latency, then stained for PBs. Consistent

with our data that showed a requirement for autophagy in KapB-mediated PB disassembly,

blocking autophagic degradation by silencing *Atg5* or *NDP52* restored PBs in KSHV-infected

cells (Fig 8A-B). We quantified PBs and compared their levels between KSHV-infected and

mock-infected cells in control cells or in silenced cells. We did not compare PB counts between

KSHV-infected and silenced cells to infected controls because silencing of autophagy regulatory

proteins influences PB homeostasis. In control cells that expressed a non-targeting shRNA,

KSHV infection decreased PBs, as expected from our previous work (Fig 8A). However, when

either *Atg5* or *NDP52* were silenced, KSHV infection had no effect on PBs relative to the

matched uninfected control, and silencing autophagy regulatory proteins restored PBs to levels

observed in the absence of infection (Fig 8A). These data confirm an essential role for autophagy

and the selective autophagy receptor NDP52 in KSHV-mediated PB disassembly.

To understand more about the molecular mechanism of NDP52-dependent PB disassembly

during KSHV infection, we first immunoblotted for steady-state levels of NDP52. KSHV

infection did not significantly alter the amount of NDP52 protein (Fig 8C). We speculated that

rather than influencing NDP52 production, enhanced autophagic flux observed after KapB

expression, KSHV infection, or Torin, would enhance the interaction of NDP52 with its target

cargo. As Torin treatment decreased Dcp1a and Pat1b protein levels, and Pat1b was restored by

NDP52 silencing (Fig 3), we tested if either Dcp1a or Pat1b were NDP52 interacting partners.

To determine this, we overexpressed FLAG-tagged NDP52 with GFP-tagged Dcp1a, GFP-

tagged Pat1b or a GFP control and immunoprecipitated using a FLAG antibody. We verified

immunoprecipitation of NDP52 and that we could detect the expression of GFP and GFP-tagged

Dcp1a in whole cell lysate; however, neither GFP alone nor GFP-Dcp1a co-immunoprecipitated

with NDP52 (Fig 8D). By contrast, we did co-immunoprecipitate GFP-Pat1b with FLAG-

NDP52. These data suggest that NDP52 and Pat1b interact which could result in delivery of

Pat1b and possibly other PB components to the nascent autophagosome for degradation. We

reasoned that if NDP52 and Pat1b interact, KSHV-mediated autophagy should decrease steady-

state levels of Pat1b. Consistent with this, Pat1b protein levels were significantly decreased after

KSHV infection (Fig 8E). KSHV infection also decreased the steady-state levels of Dcp1a, a

result consistent with earlier data that showed Dcp1a decreased in response to KapB expression

and Torin treatment. From these data, we propose the following model: KSHV infection or KapB

expression enhances autophagic flux via Beclin phosphorylation while Torin dephosphorylates

the ULK complex. In both cases, elevated autophagic flux increases the delivery of Pat1b, and

perhaps other PB components that interact with Pat1b (such as Dcp1a), to nascent

autophagosomes via recognition of Pat1b by the selective autophagy receptor, NDP52. This

interaction decreases the abundance of Pat1b, a PB scaffolding protein, resulting in PB

disassembly (Fig 8F).

<u>Discussion</u>

Here, we demonstrated that the KSHV protein KapB relies on selective autophagy to cause PB

disassembly and enhance ARE-mRNA stability. Furthermore, we identified NDP52 as the

selective autophagy receptor required for this process. To our knowledge, this is the first report

that provides a molecular basis for NDP52-mediated turnover of endogenous PBs and the first

description of the contribution of NDP52 to immune regulation by altering inflammatory

cytokine RNA turnover/suppression in PBs. We showed that PB disassembly induced by

infection with KSHV required Atg5 and NDP52, suggesting an important role for KapB-

mediated elevated autophagic flux and PB catabolism in regulating KSHV-associated

inflammation. Our major findings are: i) KapB and KSHV latent infection increased autophagic

flux by phosphorylation of the autophagy regulatory protein, Beclin 1, thereby contributing to

the complex regulation of autophagic processes during KSHV latent infection and

tumourigenesis, ii) KapB expression was necessary for KSHV-mediated autophagy increases

during latent infection, iii) KapB- and KSHV-mediated PB disassembly and enhanced translation

of an ARE-containing reporter required canonical autophagy machinery and the selective

autophagy receptor, NDP52, iv) Activation of autophagy by chemical inhibition of mTORC1

using Torin caused PB disassembly using a mechanism that was dependent on canonical

autophagy and NDP52, suggesting that KapB manipulates an existing cellular pathway that

regulates PB catabolism. These results reveal that PB turnover elicited by the KSHV KapB

protein during infection requires a host pathway that uses NDP52 to enhance PB catabolism and

promote inflammatory molecule production.

MK2 activation promotes autophagy by phosphorylating Beclin 1 (69); therefore, we

hypothesized that KapB, an MK2-activating viral protein (68), would utilize autophagy to

promote PB clearance. Our data showed that both KapB and MK2EE increased autophagic flux

via phosphorylation of Beclin 1 at Serine 90 and that KapB-mediated PB disassembly required

the canonical autophagy proteins Atg5 and Atg14, and the selective autophagy protein NDP52.

Our data are supported by previous work showing that in yeast, PB recycling requires canonical

autophagy machinery, and in mammalian cells NDP52 colocalizes with a GFP-tagged version of

Dcp1a when overexpressed (91,94). We add significantly to these studies, by performing our

work in primary human endothelial cells and by staining for endogenous PBs using

immunofluorescence for the decapping co-factor Hedls/EDC4. We revealed that when KapB was

expressed, endogenous PBs disassemble via an Atg5-, Atg14-, and NDP52-dependent

mechanism. We also complemented NDP52-silenced cells with an shRNA-resistant NDP52 and

observed the restoration of KapB-mediated PB disassembly. This occurred only in cells that

expressed KapB and not in control cells that overexpressed NDP52, an observation that suggests

that KapB-mediated autophagy increases are required for NDP52-mediated PB disassembly.

These data also suggest that NDP52 is required to select the PB protein cargo for autophagic

degradation and this is required for KSHV and KapB-mediated PB disassembly, akin to that

observed in other studies on NDP52 SAR function (95,96). Consistent with this, we showed that

the PB scaffolding protein, Pat1b, co-immunoprecipitated with NDP52, providing a molecular

basis for this interaction. In addition to identifying cargo for clearance, NDP52 has recently been

shown to orchestrate autophagosome biogenesis by recruiting essential autophagic machinery to

initiation sites (14,97,98), suggesting another possible way that NDP52 may promote KapB-

mediated PB disassembly.

Immunoelectron microscopy showed that PBs range in size from 150 to 350 nm and possess an

external area of peripheral protrusions that contain the helicase Rck/DDX6 anchored to a dense

central core of decay enzymes, including Dcp1a and Xrn1 (38). Many PB-localized enzymes are

required for PB stability including Dcp1a and Pat1b; however, only three PB proteins are

required for *de novo* PB formation, DDX6, the translation suppressor, 4E-T, and LSM14a

(44,90). Our data showed that KapB expression decreased the steady-state levels of Dcp1a, while

the steady-state levels of other PB proteins were unaffected. Inducing autophagy with the

mTORC1/2 inhibitor Torin also decreased steady-state levels of Dcp1a and Pat1b, but not other

PB proteins; however only Pat1b protein levels were restored to control levels when NDP52 was

silenced. These data support a model where the entire PB granule is not degraded by autophagy,

but PB disassembly is induced by the selective degradation of NDP52-bound Pat1b with or

without the Pat1b-interacting protein, Dcp1a. Consistent with this, we show that PBs can still

form during KapB expression if we overexpress Dcp1a-GFP or block translation using sodium

arsenite. Dcp1a is a target for other viruses that mediate PB disassembly; for example, infection

with poliovirus caused PB disassembly and the loss of Dcp1a over a time course of infection

(99); adenovirus infection has been shown to re-localize Pat1b from PBs to cytoplasmic

aggresomes during infection (100). We do not yet understand if KSHV infection or Torin

treatment alters Pat1b to promote its interaction with NDP52 and subsequent autophagic

degradation and PB disassembly; however, we speculate that post-translational modifications of

Pat1b and perhaps Dcp1a may be involved (101–107).

Infection with KSHV induces high levels of pro-inflammatory cytokines, chemokines and

angiogenic factors such as TNF, IL-1α, IL-1β, IL-6, CXCL8, IFN-γ, VEGF, COX-2 and GM-

CSF (56–62). Although several KSHV proteins promote the transcription of these products,

many harbor AREs in the 3'UTR of their cognate mRNA, meaning their transcripts are subject to

additional post-transcriptional regulation by PBs. We previously showed that KapB and the lytic

KSHV protein, viral G-protein-coupled receptor (vGPCR) enhanced the translation of an ARE-

containing reporter and elevated steady-state levels of inflammatory products, and that this

coincided with the ability of KapB and vGPCR to induce PB disassembly (51,52,66,68). We now

show that KapB expression increased steady-state levels of three ARE-containing transcripts, IL-

6, GM-CSF, and TNF, coincident with PB disassembly. However, KapB expression did not

increase CXCL8, indicating that there remains additional regulation, beyond PB disassembly, of

ARE-mRNA decay (108,109). We also show that the ability of KapB to induce PB disassembly

and enhance our ARE-mRNA reporter requires both canonical and selective autophagic

machinery. RNA transcripts for some canonical autophagy proteins in the Atg5-Atg12-Atg16

complex are also rendered labile by 3'UTR AREs, suggesting that expression of KapB or

vGPCR may cause a pro-autophagic positive feedback loop where PB disassembly culminates in

the enhanced translation of pro-autophagic products (110,111), a feature that may be aided

further by KapB-mediated upregulation of the ARE-mRNA stabilizing RBP, HuR (112). This

feed forward loop may contribute to the KapB-mediated increase in Atg5 protein levels and the

induction of autophagic flux. Taken together, our study reveals a novel way that KSHV infection

manipulates autophagic machinery to induce PB turnover and promote inflammatory molecule

production. The fact that KSHV encodes more than one protein that reverses the constitutive

repression of inflammatory mRNAs in PBs highlights its central importance.

Viral gene products both enhance and block autophagic flux during latent KSHV infection; v-

cyclin induces oncogene-induced senescence via induction of autophagy, whereas v-FLIP

counteracts v-cyclin, blocking autophagy by binding Atg3 and inhibiting LC3 lipidation (35,36).

We show herein that KapB also upregulates autophagic flux, and that KapB is necessary for the

enhanced autophagic flux we observe after KSHV infection of ECs. These data are consistent

with autophagy enhancement observed in latent B-lymphocytes (BCBL-1 cells) (113) and show

for the first time that KSHV latent infection of primary ECs culminates in an elevated autophagic

state. It is possible that KSHV products may cooperate to promote autophagy during latency,

perhaps via dysregulation of the cellular transcription factor, Nrf2, responsible for the activation

of several antioxidant response genes as well as NDP52 (114–116). Nrf2 activity is upregulated

by the viral protein vFLIP after *de novo* infection of ECs and during KSHV latency (117,118).

In viral cancers like KS, autophagic flux promotes a pro-tumourigenic and pro-inflammatory

environment (113,119–121). The data shown herein brings new understanding to this complex

relationship during KSHV infection. We show that KapB was both sufficient to induce

autophagic flux and was necessary for enhanced autophagy during KSHV latent infection. We

confirmed that autophagy machinery and NDP52 were required for PB disassembly; this

supports an important role for KapB in eliciting the pro-inflammatory environment associated

with latent KSHV infection and KS. The manipulation of NDP52 function and autophagic flux

by KSHV and KapB is yet another example of the ingenuity with which viruses usurp cellular

processes; revealing a regulatory network that links NDP52 to inflammatory molecule

production via PB catabolism.

Materials and Methods

Cell culture

All cells were grown at 37 °C with 5% CO_2 and 20% O_2. HEK293T cells (ATCC), HeLa Tet-Off

cells (Clontech), Atg5 +/+ and -/- MEFs (122), HeLa Flp-In TREx GFP-Dcp1a cells (a generous

gift from Anne-Claude Gingras), and iSLK.219 cells (a generous gift from Don Ganem) (92)

were cultured in DMEM (Thermo Fisher) supplemented with 100 U/mL penicillin, 100 µg/mL

streptomycin, 2 mM L-glutamine (Thermo Fisher) and 10% FBS (Thermo Fisher). iSLK.219

cells were additionally cultured in the presence of puromycin (10 µg/mL). HUVECs (Lonza)

were cultured in endothelial cell growth medium (EGM-2) (Lonza). HUVECs were seeded onto

gelatin (1% w/v in PBS)-coated tissue culture plates or glass coverslips. All drug treatments were

performed for the times indicated in each experiment at the concentrations listed in Table 1.

Cloning

All plasmids used in this study can be found in Table 2. All shRNAs were generated by cloning

shRNA hairpin sequences found in Table 3 into pLKO.1-TRC Puro,pLKO.1-TRC cloning vector

was a gift from David Root (Addgene plasmid # 10878; http://n2t.net/addgene:10878;

RRID:Addgene_10878) or pLKO.1-blast, pLKO.1-blast was a gift from Keith Mostov (Addgene

plasmid #26655; http://n2t.net/addgene:26655; RRID:Addgene_26655). pcDNA-FLAG-NDP52

was made by amplifying NDP52 from RFP-NDP52 (a generous gift from Dr. Andreas Till) using

the primers found in Table 4 and performing a restriction digest with EcoRI and BamHI to insert

into FLAG-HA-pCDNA, FLAG-HA-pcDNA3.1 was a gift from Adam Antebi (Addgene

plasmid # 52535; http://n2t.net/addgene:52535; RRID:Addgene_52535). pLJM1-GFP-Dcp1a

was generated from pT7-EGFP-C1-HsDcp1a, pT7-EGFP-C1-HsDCP1a was a gift from Elisa

Izaurralde (Addgene plasmid # 25030; http://n2t.net/addgene:25030; RRID:Addgene_25030)

535   through restriction digest with AgeI and EcoRI to insert into pLJM1. pLJM1-GFP-Pat1b was

536   generated from Pat1b-GFP (a generous gift from Dr. Nancy Standart) through restriction digest

537   with NheI and BamHI to insert into pLJM1.

Lentivirus Production and Transduction

All lentiviruses were generated using a second-generation system. Briefly, HEK293T cells were

transfected with pSPAX2, MD2G, and the plasmid containing a gene of interest or hairpin using

polyethylimine (PEI, Polysciences). psPAX2 was a gift from Didier Trono (Addgene plasmid

#12260; http://n2t.net/addgene:12260; RRID:Addgene_12260) and pMD2.G was a gift from

Didier Trono (Addgene plasmid #12259; http://n2t.net/addgene:12259; RRID:Addgene_12259).

Viral supernatants were harvested 48 h post-transfection, clarified using a 0.45 μ

polyethersulfone filter (VWR), and frozen at -80°C until use. For transduction, lentiviruses were

thawed at 37°C and added to target cells in complete media containing 5 μg/mL polybrene

(Sigma) for 24 h. The media was changed to selection media containing 1 μg/mL puromycin or 5

μg/mL blasticidin (Thermo) and cells were selected for 48 h before proceeding with

experiments.

Viral conditioned media treatments

iSLK.219 cells (latently infected with rKSHV.219 virus) were sub-cultured into a 10 cm cell

culture dish and grown without puromycin for 72 h. At 72 h, conditioned media was reserved

and cleared of cellular debris by centrifugation at 500 G for 5 min. Supernatant was collected

and filtered through a 0.22 μM polyethersulfone (PES) membrane filter (VWR) and stored in

aliquots at -80. Prior to the experiment, conditioned media was thawed at 37°C and combined

1:1 with fresh HUVEC EGM-2 media and used to treat vector or KapB-expressing HUVECs for

0 or 6 h prior to total RNA harvest.

Production of KSHV

Cloning and creation of the iSLK-BAC16ΔKapB cell line is described in (93). iSLK.219 or

iSLK-BAC16 cells were sub-cultured into a 10 cm cell culture dish without puromycin and

reactivated from latency with either 1 μg/mL doxycycline (Sigma-Aldrich) alone (iSLK.219) and

1 μg/mL doxycycline and 1 mM sodium butyrate (Sigma-Aldrich) (iSLK-BAC16). After 72 h,

supernatants were collected, centrifuged to remove cellular debris, aliquoted, and stored at -80°C

until use.

KSHV infection

Sub-confluent (50-70%) 12-well or 6-well plates of HUVECs or naïve iSLKs (92), respectively,

were incubated with either rKSHV.219 or KSHV BAC16 viral inoculum diluted in antibiotic and

serum free DMEM containing 5 μg/mL polybrene. Plates were centrifuged at 800 x *g* for 2 hours

at room temperature (123), the inoculum was removed and replaced with either EGM-2. For

KSHV BAC16 infections, WT and ΔKapB inoculums were normalized empirically prior to

infection to circumvent our previously observed impairment in latency establishment during

ΔKapB infection (93). To establish robust and stable latency in Figure 7, *de novo* infected iSLKs

were expanded from a 6-well to a 10cm dish, selected with 500 μg/mL hygromycin B

(ThermoFisher), split once, then selected with 1200 μg/mL hygromycin B for a combined period

of one week prior to seeding.

Immunofluorescence

Cells were seeded onto coverslips for immunofluorescence experiments and fixed for 10 mins at

37 °C in 4% (v/v) paraformaldehyde (Electron Microscopy Sciences). Samples were

permeabilized with 0.1% (v/v) Triton X-100 (Sigma-Aldrich) for 10 min at room temperature

and blocked in 1% human AB serum (Sigma-Aldrich) for 1 h at room temperature. For

autophagosome immunofluorescence cells were fixed for 10 min in ice cold methanol at -20 °C

and blocked in 1% human AB serum for 1 h at room temperature. Primary and secondary

antibodies were diluted in 1% human AB serum and used at the concentrations in Table 5.

Samples were mounted with Prolong Gold AntiFade mounting media (Thermo). Image analysis

was performed using CellProfiler (cellprofiler.org), an open source platform for image analysis

(Carpenter et al., 2006); quantification of puncta was performed as previously described, with the

exception that cells were defined by propagating out a set number of pixels from each nucleus

(124).

Co-Immunoprecipitation

HEK293T cells were transfected with plasmids expressing either FLAG-HA-pCDNA (empty

vector) or pcDNA-FLAG-NDP52 with either pLJM1-GFP (control), pLJM1-GFP-Dcp1a, or

pLJM1-GFP-Pat1b for 48 h. Cells were harvested in lysis buffer (150 mM NaCl, 10 mM Tris pH

7.4, 1 mM EDTA, 1% v/v Triton X-100, 0.5% v/v NP-40, Roche protease inhibitor tablet) and

incubated with a FLAG mouse antibody (CST) overnight at 4 °C. Protein G magnetic beads

(Biorad) were incubated overnight at 4 °C with 5 mg/mL BSA in lysis buffer without Triton X-

100 or NP-40. Immunoprecipitation was performed and samples were eluted from the magnetic

beads through boiling in 4x Laemmli buffer (Biorad) with 10% v/v beta-mercaptoethanol.

Immunoblotting

Cells were lysed in 2x Laemmli buffer and stored at -20°C until use. The DC Protein Assay

(Bio-Rad) was used to quantify protein concentration as per the manufacturer's instructions. 10-

15 µg of protein lysate was resolved by SDS-PAGE on TGX Stain-Free acrylamide gels

(BioRad). Total protein images were acquired from the PVDF membranes after transfer on the

ChemiDoc Touch Imaging system (BioRad). Membranes were blocked in 5% bovine serum

albumin (BSA) or skim milk in Tris-buffered saline-Tween20 (TBS-T). Primary and secondary

antibodies were diluted in 2.5% BSA or skim milk, dilutions can be found in Table 5.

Membranes were visualized using Clarity Western ECL substrate and the ChemiDoc Touch

Imaging system (BioRad).

Luciferase Assays

Luciferase assays were performed as previously described (66). HeLa Tet-Off cells were

transduced with recombinant lentivirus expressing different shRNAs and selected. Cells were

transfected according to Corcoran *et al.* (2011) with pTRE2 Firefly Luciferase ARE (FLuc),

pTRE2 Renilla Luciferase (RLuc), and the expression plasmid of interest using Fugene HD

(Promega). FLuc and RLuc activity were quantified using the Dual Luciferase Assay Kit

(Promega) and read on a GloMax multi-detection system (Promega).

Quantitative PCR

RNA was collected using a RNeasy Plus Mini Kit (Qiagen) according to the manufacturer's

instructions and stored at -80°C until further use. RNA concentration was determined and was

reverse transcribed using Maxima H Minus First Strand cDNA Synthesis Kit (Thermo Fisher)

using a combination of random and oligo (dT) primers according to the manufacturer's

instructions. cDNA was diluted either 1:5 or 1:10 for all qPCR experiments and GoTaq qPCR

Master Mix (Promega) was used to amplify cDNA. The ΔΔquantitation cycle (Cq) method was

used to determine the fold change in expression of target transcripts. qPCR primer sequences can

be found in Table 6.

Statistics

Data shown are the mean ± standard error of the mean (SEM). Statistical significance was

determined using a one-way ANOVA, a 2-way ANOVA or Student's t-test when appropriate

with the applicable post-test (indicated in the corresponding figure legends). All statistics were

performed using GraphPad Prism v.9.0.

Acknowledgements

The authors would like to dedicate this manuscript to Dr. Beth Levine, who was a leader in the

field of autophagy and viruses and who inspired this work. We sincerely thank the members of

the Corcoran lab for helpful discussions. We would like to thank Dr. Craig McCormick

(Dalhousie University), Dr. Eric Pringle (Dalhousie University), and Dr. Andrew Leidal (UCSF)

for valuable feedback on this manuscript and the McCormick lab for plasmids, expertise, and

advice. We would like to thank Mr. Stephen Whitefield and Ms. Mary Ann Trevors in the

Dalhousie *CORES* (Cellular and Molecular Digital Imaging and Electron Microscopy core

facilities) and Dr. Anne Vaahtokari of the Charbonneau Microscopy Facility, UCalgary for

microscopy support. We would like to thank Dr. Andreas Till (University of Bonn), Dr. Anne-

Claude Gingras (Lunenfeld-Tanenbaum Research Institute) and Dr. Don Ganem (Novartis) for

sharing valuable reagents. Model figure created with BioRender. GKS was supported by a CIHR

CGS-M scholarship and Scotia Scholars NSHRF graduate training award. ELC was supported by

a Killam predoctoral scholarship, an NSERC CGS-M scholarship, and a Nova Scotia Graduate

scholarship. MK was supported by a CSM graduate training award, a CIHR CGS-M scholarship,

and a CIHR doctoral award. Operating funds to support this work derive from a CIHR Project

Grant PJT-153210 to JAC.

Author Contributions

Carolyn-Ann Robinson: Conceptualization, Experimentation, Analysis, Paper Writing

Gillian K. Singh: Conceptualization, Experimentation, Analysis, Paper Writing

Mariel Kleer: Experimentation, Analysis, Paper Writing

Elizabeth L. Castle: Analysis

Bre Q. Boudreau: Experimentation

Jennifer A. Corcoran: Conceptualization, Experimentation, Supervision, Funding Acquisition,

Project Administration, Paper Writing

Conflict of Interest

660 The authors have no competing interests to declare.

663   **References**
664
1.   Dikic I, Elazar Z. Mechanism and medical implications of mammalian autophagy. Nat

Rev Mol Cell Biol. 2018;19:349–64.

2.   Levine B, Kroemer G. Biological Functions of Autophagy Genes: A Disease Perspective.

Cell. 2019;176: 11-42.

3.   Morishita H, Mizushima N. Diverse cellular roles of autophagy. Annu Rev Cell Dev Biol.

2019;35: 453–75.

4.   Lawrence RE, Zoncu R. The lysosome as a cellular centre for signalling, metabolism and

quality control. Nat Cell Biol. 2019;21:133-142.

5.   Beese CJ, Brynjólfsdóttir SH, Frankel LB. Selective Autophagy of the Protein

Homeostasis Machinery: Ribophagy, Proteaphagy and ER-Phagy. Front Cell Dev Biol.

2020;7: 373.

6.   Johansen T, Lamark T. Selective Autophagy: ATG8 Family Proteins, LIR Motifs and

Cargo Receptors. J Mol Biol. 2020;432: 80-103.

7.   Kirkin V, Rogov V V. A Diversity of Selective Autophagy Receptors Determines the

Specificity of the Autophagy Pathway. Mol Cell. 2019;76: 268-285.

8.   Rogov V, Dötsch V, Johansen T, Kirkin V. Interactions between Autophagy Receptors

and Ubiquitin-like Proteins Form the Molecular Basis for Selective Autophagy. Mol Cell.

2014;53:167–78.

9.   Tumbarello DA, Manna PT, Allen M, Bycroft M, Arden SD, Kendrick-Jones J, et al. The

Autophagy Receptor TAX1BP1 and the Molecular Motor Myosin VI Are Required for

Clearance of Salmonella Typhimurium by Autophagy. PLoS Pathog. 2015;11: e1005174.

10.   Sarraf SA, Shah H V., Kanfer G, Pickrell AM, Holtzclaw LA, Ward ME, et al. Loss of

TAX1BP1-Directed Autophagy Results in Protein Aggregate Accumulation in the Brain.

Mol Cell. 2020;80: 779-785.

11.   Khaminets A, Behl C, Dikic I. Ubiquitin-Dependent And Independent Signals In Selective

Autophagy. Trends Cell Biol. 2016;26: 6–16.

12.   Rogov V V, Stolz A, Ravichandran AC, Rios-Szwed DO, Suzuki H, Kniss A, et al.

Structural and functional analysis of the GABARAP interaction motif (GIM). EMBO Rep.

2017;18: 1382-1396.

13.   Marshall RS, Hua Z, Mali S, McLoughlin F, Vierstra RD. ATG8-Binding UIM Proteins

Define a New Class of Autophagy Adaptors and Receptors. Cell. 2019;177: 766-781.

14.   Ravenhill BJ, Boyle KB, von Muhlinen N, Ellison CJ, Masson GR, Otten EG, et al. The

Cargo Receptor NDP52 Initiates Selective Autophagy by Recruiting the ULK Complex to

Cytosol-Invading Bacteria. Mol Cell. 2019;74: 320-329

15.   Turco E, Witt M, Abert C, Bock-Bierbaum T, Su MY, Trapannone R, et al. How

RB1CC1/FIP200 claws its way to autophagic engulfment of SQSTM1/p62-ubiquitin

condensates. Autophagy. 2019;15: 1475-77.

16.   Vargas JNS, Wang C, Bunker E, Hao L, Maric D, Schiavo G, et al. Spatiotemporal

Control of ULK1 Activation by NDP52 and TBK1 during Selective Autophagy. Mol Cell.

2019;74: 347-362.

17.   Chen M, Meng Q, Qin Y, Liang P, Tan P, He L, et al. TRIM14 Inhibits cGAS

Degradation Mediated by Selective Autophagy Receptor p62 to Promote Innate Immune

Responses. Mol Cell. 2016;64(1):105–19.

18.   Du Y, Duan T, Feng Y, Liu Q, Lin M, Cui J, et al. LRRC25 inhibits type I IFN signaling

by targeting ISG15-associated RIG-I for autophagic degradation. EMBO J.

2018;37(3):351–66.

19.   Jin S, Tian S, Luo M, Xie W, Liu T, Duan T, et al. Tetherin Suppresses Type I Interferon

Signaling by Targeting MAVS for NDP52-Mediated Selective Autophagic Degradation in

Human Cells. Mol Cell. 2017;68:308-322.

20.   Prabakaran T, Bodda C, Krapp C, Zhang B, Christensen MH, Sun C, et al. Attenuation of

c GAS - STING signaling is mediated by a p62/ SQSTM 1-dependent autophagy pathway

activated by TBK1. EMBO J. 2018;37: e97858.

21.   Yang Q, Liu TT, Lin H, Zhang M, Wei J, Luo WW, et al. TRIM32-TAX1BP1-dependent

selective autophagic degradation of TRIF negatively regulates TLR3/4-mediated innate

immune responses. PLoS Pathog. 2017;13: e1006600.

22.   He X, Zhu Y, Zhang Y, Geng Y, Gong J, Geng J, et al. RNF 34 functions in immunity

and selective mitophagy by targeting MAVS for autophagic degradation. EMBO J.

2019;38: e100978.

23.   Seth RB, Sun L, Ea CK, Chen ZJ. Identification and characterization of MAVS, a

mitochondrial antiviral signaling protein that activates NF-κB and IRF3. Cell.

2005;122:669–82.

24.   Wu Y, Jin S, Liu Q, Zhang Y, Ma L, Zhao Z, et al. Selective autophagy controls the

stability of transcription factor IRF3 to balance type I interferon production and immune

suppression. Autophagy. 2020;17: 1376-92

25.   Orvedahl A, Levine B. Viral evasion of autophagy. Autophagy. 2008;4: 280–5.

26.   Sumpter R, Levine B. Selective autophagy and viruses. Autophagy. 2011;7: 260–5.

27.   Ahmad L, Mostowy S, Sancho-Shimizu V. Autophagy-virus interplay: From cell biology

to human disease. Front Cell Dev Biol. 2018;6: 155.

28.   Mohamud Y, Qu J, Xue YC, Liu H, Deng H, Luo H. CALCOCO2/NDP52 and

734   SQSTM1/p62 differentially regulate coxsackievirus B3 propagation. Cell Death Differ.

735   2019;26: 1062–76.

736   29.   Cesarman E, Chang Y, Moore PS, Said JW, Knowles DM. Kaposi's sarcoma-associated

737   herpesvirus-like DNA sequences in AIDS-related body-cavity-based lymphomas. N Engl

738   J Med. 1995;332: 1186–91.

739   30.   Chang Y, Cesarman E, Pessin MS, Lee F, Culpepper J, Knowles DM, et al. Identification

740   of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. Science.

741   1994;266: 1865–9.

742   31.   Soulier J, Grollet L, Oksenhendler E, Cacoub P, Cazals-Hatem D, Babinet P, et al.

743   Kaposi's sarcoma-associated herpesvirus-like DNA sequences in multicentric Castleman's

744   disease. Blood. 1995;86: 1276–80.

745   32.   Decker LL, Shankar P, Khan G, Freeman RB, Dezube BJ, Lieberman J, et al. The Kaposi

746   sarcoma-associated herpesvirus (KSHV) is present as an intact latent genome in KS tissue

747   but replicates in the peripheral blood mononuclear cells of KS patients. J Exp Med.

748   1996;184: 283–8.

749   33.   Staskus KA, Zhong W, Gebhard K, Herndier B, Wang H, Renne R, et al. Kaposi's

750   sarcoma-associated herpesvirus gene expression in endothelial (spindle) tumor cells. J

751   Virol. 1997;71: 715–9.

752   34.   Broussard G, Damania B. Regulation of KSHV Latency and Lytic Reactivation. Viruses.

753   MDPI AG; 2020;12: 1034.

754   35.   Lee J-S, Li Q, Lee J-Y, Lee S-H, Jeong JH, Lee H-R, et al. FLIP-mediated autophagy

755   regulation in cell death control. Nat Cell Biol. 2009;11: 1355–62.

36.   Leidal AM, Cyr DP, Hill RJ, Lee PWK, McCormick C. Subversion of autophagy by

Kaposi's sarcoma-associated herpesvirus impairs oncogene-induced senescence. Cell

Host Microbe. 2012;11: 167–80.

37.   Young ARJ, Narita M, Ferreira M, Kirschner K, Sadaie M, Darot JFJ, et al. Autophagy

mediates the mitotic senescence transition. Genes Dev. 2009;23: 798–803.

38.   Cougot N, Cavalier A, Thomas D, Gillet R. The Dual Organization of P-bodies Revealed

by Immunoelectron Microscopy and Electron Tomography. J Mol Biol. 2012;420: 17–28.

39.   Hubstenberger A, Courel M, Bénard M, Souquere S, Ernoult-Lange M, Chouaib R, et al.

P-Body Purification Reveals the Condensation of Repressed mRNA Regulons. Mol Cell.

2017;68:144-157.

40.   Corbet GA, Parker R. RNP granule formation: Lessons from P-bodies and stress granules.

Cold Spring Harb Symp Quant Biol. 2019;84: 203–15.

41.   Youn JY, Dunham WH, Hong SJ, Knight JDR, Bashkurov M, Chen GI, et al. High-

Density Proximity Mapping Reveals the Subcellular Organization of mRNA-Associated

Granules and Bodies. Mol Cell. 2018;69: 517-532.

42.   Riggs CL, Kedersha N, Ivanov P, Anderson P. Mammalian stress granules and P bodies at

a glance. J Cell Sci. 2020;133: jcs242487 .

43.   Luo Y, Na Z, Slavoff SA. P-Bodies: Composition, Properties, and Functions.

Biochemistry. 2018;57: 2424.

44.   Vindry C, Weil D, Standart N. Pat1 RNA-binding proteins: Multitasking shuttling

proteins. Wiley Interdiscip Rev RNA. 2019; 10: e1557

45.Barreau C, Paillard L, Osborne HB. AU-rich elements and associated factors: Are there

unifying principles? Nucleic Acids Res. Oxford University Press; 2005;33: 7138–50.

46. Bakheet T, Hitti E, Khabar KSA. ARED-Plus: An updated and expanded database of AU-

rich element-containing mRNAs and pre-mRNAs. Nucleic Acids Res. 2018;46:D218–20.

47. García-Mauriño SM, Rivero-Rodríguez F, Velázquez-Cruz A, Hernández-Vellisca M,

Díaz-Quintana A, De la Rosa MA, et al. RNA binding protein regulation and cross-talk in

the control of AU-rich mRNA Fate. Front Mol Biosci. 2017;4: 71.

48. Maitra S, Chou CF, Luber CA, Lee KY, Mann M, Chen CY. The AU-rich element mRNA

decay-promoting activity of BRF1 is regulated by mitogen-activated protein kinase-

activated protein kinase 2. RNA. 2008;14: 950–9.

49. Lai WS, Carballo E, Strum JR, Kennington EA, Phillips RS, Blackshear PJ. Evidence that

Tristetraprolin Binds to AU-Rich Elements and Promotes the Deadenylation and

Destabilization of Tumor Necrosis Factor Alpha mRNA. Mol Cell Biol. 1999;19: 4311–

23.

50. Lai WS, Blackshear PJ. Interactions of CCCH zinc finger proteins with mRNA.

Tristetraprolin-mediated AU-rich element-dependent mRNA degradation can occur in the

absence of a poly(A) tail. J Biol Chem. 2001;276: 23144–54.

51. Corcoran JA, Khaperskyy DA, Johnston BP, King CA, Cyr DP, Olsthoorn A V., et al.

Kaposi's Sarcoma-Associated Herpesvirus G-Protein-Coupled Receptor Prevents AU-

Rich-Element-Mediated mRNA Decay. J Virol. 2012;86:8859–71.

52. Corcoran JA, Johnston BP, McCormick C. Viral activation of MK2-hsp27-p115RhoGEF-

RhoA signaling axis causes cytoskeletal rearrangements, p-body disruption and ARE-

mRNA stabilization. PLoS Pathog. 2015;11:e1004597.

53. Franks TM, Lykke-Andersen J. TTP and BRF proteins nucleate processing body

formation to silence mRNAs with AU-rich elements. Genes Dev. 2007;21: 719–35.

54.   Blanco FF, Sanduja S, Deane NG, Blackshear PJ, Dixon D a. Transforming growth factor

β regulates P-body formation through induction of the mRNA decay factor tristetraprolin.

Mol Cell Biol. 2014;34: 180–95.

55.   Vindry C, Marnef A, Broomhead H, Twyffels L, Ozgur S, Stoecklin G, et al. Dual RNA

Processing Roles of Pat1b via Cytoplasmic Lsm1-7 and Nuclear Lsm2-8 Complexes. Cell

Rep. 2017;20:1187–200.

56.   Ensoli B, Nakamura S, Salahuddin SZ, Biberfeld P, Larsson L, Beaver B, et al. AIDS-

Kaposi's Sarcoma-Derived Cells Express Cytokines with Autocrine and Paracrine Growth

Effects. Science. 2010;243:223–6.

57.   Ensoli B, Stürzl M. Kaposi's sarcoma: a result of the interplay among inflammatory

cytokines, angiogenic factors and viral agents. Cytokine Growth Factor Rev. 1998;9: 63–

83.

58.   Lane BR, Liu J, Bock PJ, Schols D, Coffey MJ, Strieter RM, et al. Interleukin-8 and

growth-regulated oncogene alpha mediate angiogenesis in Kaposi's sarcoma. J Virol.

2002;76: 11570–83.

59.   Miles SA, Rezai AR, Salazar-González JF, Vander Meyden M, Stevens RH, Logan DM,

et al. AIDS Kaposi sarcoma-derived cells produce and respond to interleukin 6. Proc Natl

Acad Sci U S A. 1990;87: 4068–72.

60.   Riva G, Barozzi P, Torelli G, Luppi M. Immunological and Inflammatory Features of

Kaposi's Sarcom and Other Kaposi's Sarcoma-Associated Herpesvirus/Human

Herpesvirus 8-Associated Neoplasias. AIDS Rev. 2010;9: 25–39.

61.   Sadagopan S, Sharma-Walia N, Veettil MV, Raghu H, Sivakumar R, Bottero V, et al.

Kaposi's Sarcoma-Associated Herpesvirus Induces Sustained NF-κB Activation during

De Novo Infection of Primary Human Dermal Microvascular Endothelial Cells That Is

Essential for Viral Gene Expression. J Virol. 2007;8: 3949–68.

62. Salahuddin S, Nakamura S, Biberfeld P, Kaplan M, Markham P, Larsson L, et al.

Angiogenic properties of Kaposi's sarcoma-derived cells after long-term culture in vitro.

Science. 1988;242:430–3.

63. Corcoran JA, McCormick C. Viral activation of stress-regulated Rho-GTPase signaling

pathway disrupts sites of mRNA degradation to influence cellular gene expression. Small

GTPases. 2015;6: 178–85.

64. Standart N, Weil D. P-Bodies: Cytosolic Droplets for Coordinated mRNA Storage. Trends

Genet. 2018;34: 612–26.

65. Docena G, Rovedatti L, Kruidenier L, Fanning Á, Leakey NAB, Knowles CH, et al.

Down-regulation of p38 mitogen-activated protein kinase activation and proinflammatory

cytokine production by mitogen-activated protein kinase inhibitors in inflammatory bowel

disease. Clin Exp Immunol. 2010;162: 108–15.

66. Corcoran JA, Khaperskyy DA, McCormick C. Assays for monitoring viral manipulation

of host ARE-mRNA turnover. Methods. 2011;55: 172–81.

67. Wen KW, Damania B. Kaposi sarcoma-associated herpesvirus (KSHV): molecular

biology and oncogenesis. Cancer Lett. 2010;289: 140–50.

68. McCormick C, Ganem D. The Kaposin B Protein of KSHV Activates the p38/MK2

Pathway and Stabilizes Cytokine mRNAs. Science. 2005;307: 739–41.

69. Wei Y, An Z, Zou Z, Sumpter R, Su M, Zang X, et al. The stress-responsive kinases

MAPKAPK2/MAPKAPK3 activate starvation-induced autophagy through Beclin 1

phosphorylation. Elife. 2015;4: e05289.

70.   Shen L, Qi Z, Zhu Y, Song X, Xuan C, Ben P, et al. Phosphorylated heat shock protein 27

849   promotes lipid clearance in hepatic cells through interacting with STAT3 and activating

850   autophagy. Cell Signal. 2016;28: 1086–98.

851   71.   Gurkar AU, Chu K, Raj L, Bouley R, Lee S-H, Kim Y-B, et al. Identification of ROCK1

852   kinase as a critical regulator of Beclin1-mediated autophagy during metabolic stress. Nat

853   Commun. 2013;4: 2189.

854   72.   Klionsky DJ, Abdel-Aziz AK, Abdelfatah S, Abdellatif M, Abdoli A, Abel S, et al.

855   Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition).

856   Autophagy. 2021;17: 1-382.

857   73.   Mizushima N, Murphy LO. Autophagy Assays for Biological Discovery and Therapeutic

858   Development. Trends Biochem Sci. 2020;45: 1080-1093.

859   74.   Mizushima N, Yoshimori T, Levine B. Methods in Mammalian Autophagy Research.

860   Cell. 2010;140: 313–26.

861   75.   Mauvezin C, Neufeld TP. Bafilomycin A1 disrupts autophagic flux by inhibiting both V-

862   ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-

863   lysosome fusion. Autophagy. 2015;11:1437–8.

864   76.   Kedersha N, Anderson P. Mammalian Stress Granules and Processing Bodies. Meth

865   Enzymol. 2007;431: 61-81.

866   77.   Katayama H, Yamamoto A, Mizushima N, Yoshimori T, Miyawaki A. GFP-like proteins

867   stably accumulate in lysosomes. Cell Struct Funct. 2008;33: 1–12.

868   78.   Thoreen CC, Kang SA, Won Chang J, Liu Q, Zhang J, Gao Y, et al. An ATP-competitive

869   Mammalian Target of Rapamycin Inhibitor Reveals Rapamycin-resistant Functions of

870   mTORC1. J Biol Chem. 2009;284: 8023–32.

79.   Hardy SD, Shinde A, Wang W-H, Wendt MK, Geahlen RL. Regulation of epithelial-

mesenchymal transition and metastasis by TGF-β, P-bodies, and autophagy. Oncotarget.

2017;8: 103302–14.

80.   Liu Q, Xu C, Kirubakaran S, Zhang X, Hur W, Liu Y, et al. Characterization of Torin2, an

ATP-competitive inhibitor of mTOR, ATM, and ATR. Cancer Res. 2013;73: 2574–86.

81.   Kuang E, Tang Q, Maul GG, Zhu F. Activation of p90 Ribosomal S6 Kinase by ORF45 of

Kaposi's Sarcoma-Associated Herpesvirus and Its Role in Viral Lytic Replication. J Virol.

2008;82: 1838-50.

82.   Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through

direct phosphorylation of Ulk1. Nat Cell Biol. 2011;13: 132-41.

83.   Alers S, Löffler AS, Wesselborg S, Stork B. Role of AMPK-mTOR-Ulk1/2 in the

Regulation of Autophagy: Cross Talk, Shortcuts, and Feedbacks. Mol Cell Biol. 2012;32:

2-11.

84.   Kayyali US, Pennella CM, Trujillo C, Villa O, Gaestel M, Hassoun PM. Cytoskeletal

changes in hypoxic pulmonary endothelial cells are dependent on MAPK-activated protein

kinase MK2. J Biol Chem. 2002;277: 42596–602.

85.   Rousseau S, Houle F, Landry J, Huot J. P38 MAP kinase activation by vascular

endothelial growth factor mediates actin reorganization and cell migration in human

endothelial cells. Oncogene. 1997;15: 2169–77.

86.   Engel K, Schultz H, Martin F, Kotlyarov A, Plath K, Hahn M, et al. Constitutive

activation of mitogen-activated protein kinase-activated protein kinase 2 by mutation of

phosphorylation sites and an A-helix motif. J Biol Chem. 1995;270: 27213–21.

87.   Guo L, Louis IV-S, Bohjanen PR. Post-transcriptional regulation of cytokine expression

and signaling. Curr Trends Immunol. 2018;19: 33–40.

88.   Franks TM, Lykke-Andersen J. The control of mRNA decapping and P-body formation.

Mol Cell. 2008;32: 605–15.

89.   Sharma-Walia N, Paul AG, Bottero V, Sadagopan S, Veettil MV, Kerur N, et al. Kaposi's

Sarcoma Associated Herpes Virus (KSHV) induced COX-2: A key factor in latency,

inflammation, angiogenesis, cell survival and invasion. PLoS Pathog. 2010;6: e1000777.

90.   Ayache J, Bénard M, Ernoult-Lange M, Minshall N, Standart N, Kress M, et al. P-body

assembly requires DDX6 repression complexes rather than decay or Ataxin2/2L

complexes. Mol Biol Cell. 2015;26: 2579–95.

91.   Guo H, Chitiprolu M, Gagnon D, Meng L, Perez-Iratxeta C, Lagace D, et al. Autophagy

supports genomic stability by degrading retrotransposon RNA. Nat Commun. 2014;5:

5276.

92.   Myoung J, Ganem D. Generation of a doxycycline-inducible KSHV producer cell line of

endothelial origin: Maintenance of tight latency with efficient reactivation upon induction.

J Virol Methods. 2011;174: 12-21.

93.   Kleer M, MacNeil G, Adam N, Pringle ES, Corcoran JA. A panel of KSHV mutants in the

polycistronic kaposin locus for precise analysis of individual protein products. J Virol.

2022;96: e0156021

94.   Buchan JR, Kolaitis R, Taylor JP, Parker R. Eukaryotic Stress Granules Are Cleared by

Autophagy and Cdc48 / VCP Function. Cell. 2013;153: 1461–74.

95.   Xie X, Li F, Wang Y, Wang Y, Lin Z, Cheng X, et al. Molecular basis of ubiquitin

recognition by the autophagy receptor CALCOCO2. Autophagy. 2015;11: 1775–89.

96.   Cemma M, Kim PK, Brumell JH. The ubiquitin-binding adaptor proteins p62/SQSTM1

and NDP52 are recruited independently to bacteria-associated microdomains to target

Salmonella to the autophagy pathway. Autophagy. 2011;7: 341–5.

97.   Yamano K, Youle RJ. Two different axes CALCOCO2-RB1CC1 and OPTN-ATG9A

initiate PRKN-mediated mitophagy. Autophagy. 2020;16: 2105–7.

98.   Shi X, Chang C, Yokom AL, Jensen LE, Hurley JH. The autophagy adaptor ndp52 and the

fip200 coiled-coil allosterically activate ulk1 complex membrane recruitment. Elife.

2020;9: 1–20.

99.   Dougherty JD, White JP, Lloyd RE. Poliovirus-mediated disruption of cytoplasmic

processing bodies. J Virol. 2011;85: 64–75.

100. Friedman E, Karen K. Effects of Adenovirus Infection on the Localization of Cellular

Protein Pat1b. Georg J Sci. 2017;75: 15.

101. Bish R, Cuevas-Polo N, Cheng Z, Hambardzumyan D, Munschauer M, Landthaler M, et

al. Comprehensive Protein Interactome Analysis of a Key RNA Helicase: Detection of

Novel Stress Granule Proteins. Biomolecules. 2015;5: 1441–66.

102. Gustafson EA, Wessel GM. DEAD-box helicases: posttranslational regulation and

function. Biochem Biophys Res Commun. 2010;395: 1–6.

103. Rahman H, Qasim M, Oellerich M, Asif AR. Crosstalk between Edc4 and mammalian

target of rapamycin complex 1 (mTORC1) signaling in mRNA decapping. Int J Mol Sci.

2014;15: 23179–95.

102. Yoon J-H, Choi E-J, Parker R. Dcp2 phosphorylation by Ste20 modulates stress granule

assembly and mRNA decay in Saccharomyces cerevisiae. J Cell Biol. 2010;189: 813–27.

105. Tenekeci U, Poppe M, Beuerlein K, Busch J, Schmitz ML, Kracht Correspondence M.

K63-Ubiquitylation and TRAF6 Pathways Regulate Mammalian P-Body Formation and

mRNA Decapping. Mol Cell. 2016;62: 943–57.

106. Aizer A, Kafri P, Kalo A, Shav-Tal Y. The P Body Protein Dcp1a Is Hyper-

phosphorylated during Mitosis. Prigent C, editor. PLoS One. 2013;8: e49783.

107. Chiang PY, Shen YF, Su YL, Kao CH, Lin NY, Hsu PH, et al. Phosphorylation of mRNA

Decapping Protein Dcp1a by the ERK Signaling Pathway during Early Differentiation of

3T3-L1 Preadipocytes. PLoS One. 2013;8: e61697.

108. Palanisamy V, Park NJ, Wang J, Wong DT. AUF1 and HuR proteins stabilize interleukin-

8 mRNA in human saliva. J Dent Res. 2008;87: 772–6.

109. Palanisamy V, Jakymiw A, Van Tubergen EA, D'Silva NJ, Kirkwood KL. Control of

Cytokine mRNA Expression by RNA-binding Proteins and microRNAs. J Dent Res.

2012;91: 651–8.

110. Lystad AH, Carlsson SR, de la Ballina LR, Kauffman KJ, Nag S, Yoshimori T, et al.

Distinct functions of ATG16L1 isoforms in membrane binding and LC3B lipidation in

autophagy-related processes. Nat Cell Biol. 2019;21: 372–83.

111. Ji E, Kim C, Kang H, Ahn S, Jung M, Hong Y, et al. RNA Binding Protein HuR Promotes

Autophagosome Formation by Regulating Expression of Autophagy-Related Proteins 5,

12, and 16 in Human Hepatocellular Carcinoma Cells. Mol Cell Biol. 2019;39: e00508-

18.

112. Yoo J, Kang J, Lee HN, Aguilar B, Kafka D, Lee S, et al. Kaposin-B enhances the

PROX1 mRNA stability during lymphatic reprogramming of vascular endothelial cells by

Kaposi's sarcoma herpes virus. PLoS Pathog. 2010;6: e1001046.

113. Pringle ES, Robinson C-A, McCormick C. Kaposi's Sarcoma-Associated Herpesvirus

Lytic Replication Interferes with mTORC1 Regulation of Autophagy and Viral Protein

Synthesis. J Virol. 2019;93: e00854-19.

114.   Jo C, Gundemir S, Pritchard S, Jin YN, Rahman I, Johnson GVW. Nrf2 reduces levels of

phosphorylated tau protein by inducing autophagy adaptor protein NDP52. Nat Commun.

2014;5: 3496.

115.   Moi P, Chan K, Asunis I, Cao A, Kan YW. Isolation of NF-E2-related factor 2 (Nrf2), a

NF-E2-like basic leucine zipper transcriptional activator that binds to the tandem NF-

E2/AP1 repeat of the β-globin locus control region. Proc Natl Acad Sci U S A. 1994;91:

9926–30.

116.   Nguyen T, Nioi P, Pickett CB. The Nrf2-antioxidant response element signaling pathway

and its activation by oxidative stress. J Biol Chem. 2009;284: 13291–5.

117.   Gjyshi O, Bottero V, Veettil MV, Dutta S, Singh VV, Chikoti L, et al. Kaposi's sarcoma-

associated herpesvirus induces Nrf2 during de novo infection of endothelial cells to create

a microenvironment conducive to infection. PLoS Pathog. 2014;10: e1004460.

118.   Gjyshi O, Flaherty S, Veettil MV, Johnson KE, Chandran B, Bottero V. Kaposi's

Sarcoma-Associated Herpesvirus Induces Nrf2 Activation in Latently Infected Endothelial

Cells through SQSTM1 Phosphorylation and Interaction with Polyubiquitinated Keap1. J

Virol. 2015;89: 2268–86.

119.   Vescovo T, Pagni B, Piacentini M, Fimia GM, Antonioli M. Regulation of Autophagy in

Cells Infected With Oncogenic Human Viruses and Its Impact on Cancer Development.

Front Cell Dev Biol. 2020;8: 47.

120.   Liu J, Debnath J. The Evolving, Multifaceted Roles of Autophagy in Cancer. Adv Cancer

Res. 2016;130: 1–53.

121.   Wu WKK, Coffelt SB, Cho CH, Wang XJ, Lee CW, Chan FKL, et al. The autophagic

paradox in cancer therapy. Oncogene. 2012;31: 939–53.

122. Kuma A, Hatano M, Matsui M, Yamamoto A, Nakaya H, Yoshimori T, et al. The role of

autophagy during the early neonatal starvation period. Nature. 2004;432: 1032–6.

123. Yoo SM, Ahn AK, Seo T, Hong HB, Chung MA, Jung SD, et al. Centrifugal enhancement

of Kaposi's sarcoma-associated virus infection of human endothelial cells in vitro. J Virol

Methods. 2008;154: 160-6.

124. Castle EL, Robinson C-A, Douglas P, Rinker KD, Corcoran JA. Viral manipulation of a

mechanoresponsive signaling axis disassembles processing bodies. Mol Cell Biol.

2021;41: e0039921.

125. Horn M, Geisen C, Cermak L, Becker B, Nakamura S, Klein C, et al. DRE-1/FBXO11-

dependent degradation of BLMP-1/BLIMP-1 governs C. elegans developmental timing

and maturation. Dev Cell. 2014;28: 697-710.

126. Moffat J, Grueneberg DA, Yang X, Kim SY, Kloepfer AM, Hinkle G, et al. A Lentiviral

RNAi Library for Human and Mouse Genes Applied to an Arrayed Viral High-Content

Screen. Cell. 2006;124: 1283–98.

127. Bryant DM, Datta A, Rodríguez-Fraticelli AE, PeräCurrency Signnen J, Martín-Belmonte

F, Mostov KE. A molecular network for de novo generation of the apical surface and

lumen. Nat Cell Biol. 2010;12: 1035–45.

128. Johnston BP, Pringle ES, McCormick C. KSHV activates unfolded protein response

sensors but suppresses downstream transcriptional responses to support lytic replication.

PLoS Pathog. 2019;15: e1008185.

129. Sancak Y, Peterson TR, Shaul YD, Lindquist RA, Thoreen CC, Bar-Peled L, et al. The rag

GTPases bind raptor and mediate amino acid signaling to mTORC1. Science. 2008;320:

1496-501.

130. Tritschler F, Braun JE, Motz C, Igreja C, Haas G, Truffault V, et al. DCP1 forms

asymmetric trimers to assemble into active mRNA decapping complexes in metazoa. Proc

Natl Acad Sci U S A. 2009;106: 21591-6.

131. Marnef A, Weil D, Standart N. RNA-related nuclear functions of human Pat1b, the P-

body mRNA decay factor. Mol Biol Cell. 2012;23: 213-24.

1015

1016
1017

 Figure Legends

 Figure 1: Kaposin B increases autophagic flux. Primary human umbilical vein endothelial

 cells (HUVECs) were transduced with recombinant lentiviruses expressing either Kaposin B

 (KapB) or an empty vector control and selected with blasticidin (5 µg/mL). A: Cells were treated

 with Bafilomycin A1 (BafA1, 10 nM) or a vehicle control (DMSO) for the indicated times prior

 to harvest in 2x Laemmli buffer. Protein lysates were resolved by SDS-PAGE and immunoblot

 was performed for p62 and LC3. Samples were quantified using Image Lab (Biorad) software

 and then normalized, first to total protein and then to their respective starting time points (0 h).

 Results were plotted in GraphPad and a linear regression statistical test was performed, ±SEM;

 n=3, *=P<0.05, ***=P<0.001. B: Cells were treated with BafA1 for 30 min prior to fixation in

 methanol. Immunofluorescence was performed for LC3 (white) and DAPI (nuclei, blue). Scale

 bar=20 µm. Total LC3 area per field was quantified by identifying LC3-positive puncta using

 CellProfiler and normalizing to the number of nuclei and the vector DMSO control. Results were

 plotted in GraphPad, a 2-way ANOVA was performed with a Šidák's multiple comparison test,

 matched data from each replicate was paired for comparison, ±SEM; n=3, *=P<0.05. C:

 HUVECs were transduced with recombinant lentiviruses expressing either Kaposin B (KapB) or

 an empty vector control and selected with blasticidin (5 µg/mL). Cells were treated with sodium

 arsenite (0.25 mM) or a vehicle control for 30 min prior to fixation in 4% paraformaldehyde and

 permeabilization in 0.1% Triton X-100. Scale bar=20 µm. Hedls puncta were quantified using

 CellProfiler and presented as number of Hedls puncta per cell, all cells counted are displayed.

 Results were plotted in GraphPad and a 2-way ANOVA was performed on the main column

 effects with a Tukey's multiple comparison test, bar represents the mean; n=3, **=P<0.01,

 ****=P<0.0001.

Figure 2: Kaposin B requires canonical autophagy to disassemble PBs. A & B: HUVECs

were sequentially transduced: first with recombinant lentiviruses expressing either shRNAs

targeting Atg5 (C) or Atg14 (D) (shAtg5, shAtg14) or a non-targeting control (NS) and selected

with puromycin (1 µg/mL), and second with either KapB or an empty vector control and selected

with blasticidin (5 µg/mL). Coverslips were fixed in 4% paraformaldehyde, permeabilized in

0.1% Triton X-100, and immunostained for the PB-resident protein Hedls (white) and nuclei

(DAPI). Scale bar=20 µm. Hedls puncta were quantified using CellProfiler and presented as

number of Hedls puncta per cell, all cells counted are displayed. Results were plotted in

GraphPad and a 2-way ANOVA was performed on the main column effects with a Tukey's

multiple comparison test, bar represents the mean; n=4 (C), n=3 (D), ****=P<0.0001. C:

HUVECs were treated as in B. Coverslips were fixed in 4% paraformaldehyde, permeabilized in

0.1% Triton X-100, and immunostained for the PB-resident protein Hedls (white) and nuclei

(DAPI). Scale bar=20 µm. Hedls puncta were quantified using CellProfiler and presented as

number of Hedls puncta per cell, all cells counted are displayed. Results were plotted in

GraphPad and a 2-way ANOVA was performed on the main column effects with a Tukey's

multiple comparison test, bar represents the mean; n=3, **=P<0.01.

1057

Figure 3: Torin treatment disassembles PBs. A: HUVECs were treated with either Torin (250

nM) or a DMSO control for 2 h prior to fixation in 4% paraformaldehyde. Samples were

permeabilized in 0.1% Triton X-100. Immunofluorescence was performed for Hedls (PBs, white)

and DAPI (nuclei, blue). Scale bar=20 µm. Hedls puncta were quantified using CellProfiler and

presented as number of Hedls puncta per cell, all cells counted are displayed. Results were

plotted in GraphPad and a 2-way ANOVA was performed on the main column effects with a

Tukey's multiple comparison test, bar represents the mean; n=3, ****=P<0.0001. B: HUVECs

were treated with either Torin (250 nM) or a DMSO control for 90 min prior to the addition of

sodium arsenite (0.25 mM) for 30 min. Cells were fixed in 4% paraformaldehyde and

permeabilized in 0.1% Triton X-100. Immunofluorescence was performed for Hedls (PBs, white)

and DAPI (nuclei, blue). Scale bar=20 μm. Hedls puncta were quantified using CellProfiler and

presented as number of Hedls puncta per cell, all cells counted are displayed. Results were

plotted in GraphPad and a 2-way ANOVA was performed on the main column effects with a

Tukey's multiple comparison test, bar represents the mean; n=3, *=P<0.05. C: HUVECs were

transduced with recombinant lentiviruses expressing either KapB, constitutively active MK2

(MK2EE), or an empty vector control and selected with blasticidin (5 μg/mL). Cells were treated

with either Torin (250 nM) or a DMSO control for 2 h prior to harvest in 2x Laemmli buffer.

Protein lysates were resolved by SDS-PAGE and immunoblot was performed for phospho-S6

(Ser235/236) and total S6. D: HUVECs were transduced and treated as in C and harvested in 2X

Laemmli buffer. Protein lysates were resolved by SDS-PAGE and immunoblot was performed

for phospho-ULK (Ser757), phospho-ULK (Ser555), and total ULK. E: HUVECs were

transduced as in C and harvested in 2X Laemmli buffer. Protein lysates were resolved by SDS-

PAGE and immunoblot was performed for phospho-Beclin 1 (Ser90) and total Beclin 1. Samples

were quantified by normalizing total and phospho-Beclin 1 protein levels to the total protein in

each lane and then using Image Lab (Biorad), normalizing to the Vector control, and then

dividing phospho- over total Beclin 1. Results were plotted in GraphPad and a one-way

ANOVA was performed with a Dunnett's test, ±SEM; n=4, *=P<0.05.

1085

Figure 4: KapB-mediated ARE-mRNA reporter expression requires canonical autophagy.

A: HUVECs were transduced with recombinant lentiviruses expressing either KapB or an empty vector control and selected with blasticidin (5 µg/mL). Conditioned media harvested from r.219.KSHV latently infected iSLK cells was used to mimic the KS lesion microenvironment and induce the transcription of cytokines for 6 h prior to lysis for total RNA, normal media was used for the 0 h time point. Transcript levels were quantified by qPCR and normalized to 18S as a reference gene. Data is represented as the fold change in target transcript expression relative to the untreated vector control and was quantified using the ΔΔCq method. Results were plotted in GraphPad, a 2-way ANOVA was performed, ±SEM; n=3, *=$P<0.05$, **=$P<0.01$. B: HUVECs were treated with Torin (250 nM) or a DMSO control for 2 h prior to lysis for total RNA. Transcript levels were quantified by qPCR and normalized to HPRT as a reference gene. Data is represented as the fold change in target transcript expression relative to the untreated vector control and was quantified using the ΔΔCq method. An unpaired t-test was performed, ±SEM; n=3, *=$P<0.05$. C: HeLa Tet-Off cells were co-transfected with expression plasmids for an ARE-containing firefly luciferase plasmid (pTRE-FLuc-ARE) and a stable renilla luciferase plasmid (pTRE-RLuc). 36 h post transfection, doxycycline (Dox) was added to halt reporter gene transcription of both luciferase reporters, at the same time Torin (250 nM) or DMSO were added; 12 h after Dox addition, lysates were harvested in passive lysis buffer (Promega). Luciferase activity for both FLuc and RLuc was analyzed using the Dual-Luciferase Reporter Assay (Promega) and normalized (FLuc/RLuc) relative luciferase was calculated in relative light units (RLUs). Results were plotted using GraphPad, an unpaired t-test was performed, ±SEM; n=6, ***=$P<0.001$. D&E: HeLa Tet-Off cells were transduced with recombinant lentiviruses expressing either shRNAs targeting Atg5 or Atg14 (shAtg5, shAtg14) or a non-targeting control

(NS) and selected with puromycin (1 µg/mL). After selection, cells were co-transfected and Dox

treatment was performed as described in C, except that co-transfection also included an

expression plasmid for KapB or an empty vector control. Results were plotted using GraphPad,

an unpaired t-test was performed, ±SEM; n=3, *=P<0.05, **=P<0.01. F: Cells were co-

transfected as in E and BafA1 (10 nM) was added at the same time as Dox. Results were plotted

in GraphPad and a Student's t-test was performed, ±SEM; n=3, *=P<0.05, **=P<0.01.

1115

Figure 5: Dcp1a protein levels are decreased by KapB expression and Torin treatment. A:

HUVECs were treated with DMSO or Torin (250 nM) for 4 h prior to harvest. Samples were

lysed in 2x Laemmli buffer and resolved by SDS-PAGE before immunoblotting with Xrn1,

EDC4/Hedls, Dcp1a, and DDX6. Samples were quantified by normalizing the PB resident

protein levels to the total protein in each lane and then the DMSO control using ImageLab

(Biorad). Results were plotted in GraphPad and a one-way ANOVA was performed ±SEM; n=3,

**=P<0.01. B: HUVECs were transduced with recombinant lentiviruses expressing either KapB

or an empty vector control and selected with blasticidin (5 µg/mL). Cells were treated with

DMSO or bafilomycin A1 (BafA1, 10 nM) for 4 h prior to harvest in 2x Laemmli buffer.

Samples were resolved by SDS-PAGE and immunoblot was performed for Dcp1a or p62

(autophagy marker). Samples were quantified by normalizing Dcp1a protein levels to the total

protein in each lane using Image Lab (Biorad) and then to the Vector DMSO control.

Representative blot is from the same membrane, hashed line indicates skipped lanes. Results

were plotted in GraphPad and a 2-way ANOVA was performed, ±SEM; n=3, ***=P<0.001. C:

HUVECs were transduced with recombinant lentiviruses expressing either KapB or an empty

vector control and selected with blasticidin (5 µg/mL). Samples were harvested in 2x Laemmli

buffer, resolved by SDS-PAGE and immunoblot was performed for Xrn1, Hedls/EDC4, or

DDX6. Samples were quantified by normalizing PB protein levels to the total protein in each

lane using Image Lab (Biorad) and then to the Vector control. Results were plotted in GraphPad.

D: HUVECs were sequentially transduced: first with recombinant lentiviruses expressing either

shRNAs targeting Atg5 (shAtg5) or a non-targeting control (NS) and selected with puromycin (1

µg/mL), and second with either KapB or an empty vector control and selected with blasticidin (5

µg/mL). Samples were harvested in 2x Laemmli buffer and resolved by SDS-PAGE.

Immunoblot was performed for Dcp1a, Atg5, and KapB. Samples were quantified by

normalizing Dcp1a protein levels to the total protein in each lane using Image Lab (Biorad) and

then to the Vector NS control. Results were plotted in GraphPad and a 2-way ANOVA was

performed, ±SEM; n=3, *=P<0.05, **P=<0.01.

1143

Figure 6: KapB-mediated PB disassembly and ARE-mRNA reporter expression require the

selective autophagy receptor NDP52. A: HUVECs were sequentially transduced with

recombinant lentivirus expressing KapB or an empty vector control and selected with blasticidin

(5 µg/mL), and second with shRNAs targeting NDP52 or a non-targeting control (NS) and

selected with puromycin (1 µg/mL). Coverslips were fixed in 4% paraformaldehyde,

permeabilized in 0.1% Triton X-100 and immunostained for Hedls (PBs; white), DAPI (nuclei,

blue). Scale bar=20 µm. Hedls puncta were quantified using CellProfiler and presented as

number of Hedls puncta per cell, all cells counted are displayed. Results were plotted in

GraphPad and a 2-way ANOVA was performed on the main column effects with a Tukey's

multiple comparison test, bar represents the mean; n=3, *=P<0.05. B: HUVECs were transduced

with recombinant lentivirus expressing an shRNA targeting NDP52 and selected with puromycin

(1 µg/mL). Cells were treated with Torin (250 nM) or a DMSO control for 2 h prior to fixation in

4% paraformaldehyde and permeabilization in 0.1% Triton X-100. Samples were immunostained

for Hedls (PBs; white), DAPI (nuclei, blue). Scale bar=20 µm. Hedls puncta were quantified

using CellProfiler and presented as number of Hedls puncta per cell, all cells counted are

displayed. Results were plotted in GraphPad and a 2-way ANOVA was performed on the main

column effects with a Tukey's multiple comparison test, bar represents the mean; n=3,

****=P<0.0001. C: HUVECs were transduced with recombinant lentiviruses expressing either

an shRNA targeting NDP52 or a non-targeting control (NS) and selected with puromycin (1

µg/mL). Cells were treated with DMSO or Torin (250 nM) for 4 h prior to harvest in 2x Laemmli

buffer. Samples were resolved by SDS-PAGE and immunoblot was performed for Dcp1a, Pat1b

or DDX6. Samples were quantified by normalizing Dcp1a or Pat1b protein levels to the total

protein in each lane using Image Lab (Biorad) and then to the NS DMSO control. Results were

plotted in GraphPad and a 2-way ANOVA was performed, ±SEM; n=3 (Dcp1a) n=4 (Pat1b),

*=P<0.05. D: HeLa Tet-Off cells were transduced with recombinant lentivirus expressing

shRNAs targeting NDP52, OPTN, p62 or a NS control and selected with puromycin (1µg/mL)

then cells were co-transfected, treated with Dox and luciferase activity was recorded and

analyzed as in Figure 3. Data were plotted in GraphPad as the mean fold change in the relative

luciferase activity of each condition compared to vector NS or KapB NS; n=4. An unpaired t-test

was performed; * = P<0.05 **= P<0.01. E: HUVECs were sequentially transduced first with

recombinant lentivirus expressing shDNP52 targeting the 3'-UTR of NDP52 or a NS control and

selected with blasticidin (5 µg/mL), and second, with KapB and either mCherry control (mCh) or

RFP-NDP52. Coverslips were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton

X-100, and immunostained with Hedls (PBs, green). Scale bar=20 µm. F: Samples from E were

quantified; Hedls puncta were quantified using CellProfiler and presented as number of Hedls

puncta per cell, all cells counted are displayed. Results were plotted in GraphPad and a 2-way

ANOVA was performed on the main column effects with a Tukey's multiple comparison test,

bar represents the mean; n=3, ****=P<0.0001.

1182

Figure 7: KSHV infection induces autophagic flux and phosphorylation of Beclin 1. A:

HUVECs were infected with rKSHV.219 via spinoculation and infection was allowed to proceed

for 96 h prior to treatment with bafilomycin A1 (10 nM) for 30 min. Coverslips were fixed in

methanol and immunostained for LC3 (autophagosomes; white) and DAPI (nuclei, blue). Cells

were visually assessed for infection via viral latency associated nuclear antigen (LANA)

immunofluorescence. Scale bar=20 μm. Total LC3 area per field was quantified by identifying

LC3 puncta using CellProfiler and normalizing to the number of nuclei and the Mock DMSO

control. Results were plotted in GraphPad and a 2-way ANOVA was performed with a Šidák's

multiple comparison test, ±SEM n=5; *=P<0.05. B: Naïve iSLK cells were infected with wild-

type (WT) or delete KapB (delB) BAC16 KSHV via spinoculation, robust and stable latency was

achieved by expanding and selecting cells in hygromycin B (500 μg/mL and then 1200 μg/mL)

for one week before seeding for experiments. Cells were treated with Bafilomycin A1 (BafA1,

10 nM) or a vehicle control (DMSO) for the indicated times prior to harvest in 2x Laemmli

buffer. Protein lysates were resolved by SDS-PAGE and immunoblot was performed for p62 and

LC3. Samples were quantified using Image Lab (Biorad) software and then normalized, first to

total protein and then to their respective starting time points (0 h). Results were plotted in

GraphPad and a linear regression statistical test was performed, ±SEM; n=3, *=P<0.05. C:

HUVECs were infected as in A for 96 h and harvested in 2x Laemmli buffer. Protein lysates

were resolved by SDS-PAGE and immunoblot was performed for phosphor-Beclin 1 (S90) and

total Beclin 1. Samples were quantified using Image Lab (Biorad) software and then normalized,

first to total protein and then to the Mock control. Results were plotted in GraphPad and a

Student's t-test was performed, ±SEM n=3; *=P<0.05. D: HUVECs were infected with wild-

type (WT) or delete KapB (delB) BAC16 KSHV via spinoculation and infection was allowed to

proceed for 96 h prior to treatment with bafilomycin A1 (10 nM) for 30 min. Coverslips were

fixed in methanol and immunostained for LC3 (autophagosomes; white) and DAPI (nuclei,

blue). Cells were visually assessed for infection via viral LANA immunofluorescence. Scale

bar=20 μm. Total LC3 area per field was quantified by identifying LC3 puncta using CellProfiler

and normalizing to the number of nuclei and the Mock DMSO control. Results were plotted in

GraphPad and a 2-way ANOVA was performed with a Šidák's multiple comparison test, ±SEM

n=3; *P<0.05. E: HUVECs were infected as in C for 96 h and harvested in 2x Laemmli buffer.

Protein lysates were resolved by SDS-PAGE and immunoblot was performed for phospho-

Beclin 1 (S90) and total Beclin 1. Samples were quantified using Image Lab (Biorad) software

and then normalized, first to total protein and then to the Mock control. Results were plotted in

GraphPad and a one-way ANOVA was performed, ±SEM n=3; *=P<0.05.

1217

Figure 8: KSHV-mediated PB disassembly requires Atg5 and the selective autophagy

receptor NDP52. A: HUVECs were transduced with recombinant lentivirus expressing shRNAs

targeting Atg5, NDP52, or a non-targeting control (NS) and selected with puromycin (1 μg/mL).

Cells were infected with rKSHV.219 via spinoculation and infection was allowed to proceed for

96 h. Coverslips were fixed in 4% paraformaldehyde, permeabilized in 0.1% Triton X-100 and

immunostained for Hedls (PBs; white), LANA (KSHV latency, green), and DAPI (nuclei, blue).

Scale bar=20 μm. Hedls puncta were quantified using CellProfiler and presented as number of

Hedls puncta per cell, all cells counted are displayed. Results were plotted in GraphPad and a 2-

way ANOVA was performed on the main column effects with a Tukey's multiple comparison

test, bar represents the mean; n=3, *=P<0.05. B: HUVECs were transduced with recombinant

lentiviruses and infected with rKSHV.219 as in A. Cells were harvested in 2X Laemmli buffer.

Protein lysates were resolved by SDS-PAGE and immunoblot was performed for Atg5 and

NDP52. C: HUVECs were infected with rKSHV.219 via spinoculation and infection was

allowed to proceed for 96 h. Cells were harvested in 2x Laemmli buffer. Samples were resolved

by SDS-PAGE and immunoblot was performed for NDP52. Samples were quantified by

normalizing NDP52 protein levels to the total protein in each lane using Image Lab (Biorad) and

then to Mock. Results were plotted in GraphPad and an unpaired t-test was performed, ±SEM;

n=3, *=P<0.05. D: HEK293T cells were transfected with plasmids expressing pcDNA-FLAG-

NDP52 with either pLJM1-GFP (control), pLJM1-GFP-Dcp1a, or pLJM1-GFP-Pat1b for 48 h.

Cells were harvested in lysis buffer and incubated with a FLAG mouse antibody (CST) overnight

at 4 °C. Immunoprecipitation was performed and samples were eluted from the magnetic beads

through boiling in 4x Laemmli buffer (Biorad) with 10% v/v beta-mercaptoethanol. Samples

were resolved by SDS-PAGE. A representative blot of three independent experiments is shown.

E: HUVECs were infected with rKSHV.219 via spinoculation and infection was allowed to

proceed for 96 h. Cells were harvested in 2x Laemmli buffer. Samples were resolved by SDS-

PAGE and immunoblot was performed for Dcp1a or Pat1b. Samples were quantified by

normalizing Dcp1a or Pat1b protein levels to the total protein in each lane using Image Lab

(Biorad) and then to Mock. Results were plotted in GraphPad and an unpaired t-test was

performed, ±SEM; n=3, **=P<0.01. F:. Model of PB disassembly induced by KSHV/KapB.

KapB increases the phosphorylation of Beclin 1 at Serine 90 to enhance autophagic flux while

Torin induces autophagic flux via the inhibition of mTORC1/2 and dephosphorylation of the

ULK complex. Enhanced activation of autophagic flux by either mechanism results in PB

disassembly that is dependent on the selective autophagy receptor, NDP52. In our model, we

predict that NDP52 recruits a PB component(s) to the nascent autophagosome. Our top candidate

for this recruitment is the PB scaffolding protein, Pat1b, as NDP52 co-immunoprecipitated with

Pat1b consistent with Pat1b protein level decreases after KSHV infection and Torin treatment.

Steady state levels of Dcp1a also decreased in response to KapB expression, KSHV infection or

Torin treatment, suggesting that Dcp1a decreases may be mediated by its interaction with Pat1b.

Autophagic degradation of Pat1b and Dcp1a lead to PB disassembly.

1257

1258

1259

1260 **Tables**

1261 Table 1: Drugs

Drug	Vendor/Catalogue #	Concentration
Torin1	Sigma-Aldrich 475991	250 nM
Bafilomycin A1	Sigma-Aldrich B1793	10 nM
Chloroquine	Sigma-Aldrich C6628	25 µM
MG-132	Selleckchem S2619	5 µM
LPS	Sigma-Aldrich L3129	1 µg/mL
Sodium Arsenite	Sigma-Aldrich 1.06277	0.25 mM
Doxycycline	Sigma-Aldrich D9891	1 µg/mL
Hygromycin B	ThermoFisher 10687010	500 or 1200 µg/mL

1262
1263 Table 2: Plasmids

Plasmid	Use	Source	Mammalian Selection
RFP-NDP52	Cloning	Dr. Andreas Till (University Hospital of Bonn)	N/A
pcDNA 3.1 (+)	Empty Vector Control	Invitrogen V79020	N/A
FLAG-HA-pcDNA	Empty Vector Control Co-IP	Addgene #52535 (125)	N/A
pcDNA KapB BCBL1	ARE-mRNA stability	(52)	N/A
pcDNA FLAG-MK2EE	ARE-mRNA stability	(52)	N/A
pLKO.1-TRC	shRNA expression	Addgene #10878 (126)	Puromycin
pLKO.1-blast	shRNA expression	Addgene #26655(127)	Blasticidin
pLJM1	Empty Vector Control	(128)	Blasticidin/ Puromycin
pLJM1 KapB (pulmonary KS)	Overexpression	Cloned from pBMNIP-KapB (52) into pLJM1-BSD	Blasticidin
pLJM1 mCh	Overexpression Control	Craig McCormick (Dalhousie University)	Blasticidin
pLJM1 FLAG-MK2EE	Overexpression	Cloned into pLJM1 using BamHI and EcoRI from pBMN FLAG-MK2EE (52)	Blasticidin/ Puromycin
pLJM1 RFP-NDP52	Overexpression	Cloned into pLJM1 using NheI and BamHI from RFP-NDP52	Puromycin
pcDNA-FLAG-NDP52	Co-IP	Cloned using primers listed in Table 4 from RFP-NDP52	N/A
pLJM1-EGFP	Co-IP	Addgene #19319 (129)	Puromycin
pT7-EGFP-C1-HsDcp1a	Cloning	Addgene #25030 (130)	N/A
pLJM1-GFP-Dcp1a	Co-IP	Cloned into pLJM1 using AgeI and EcoRI from pT7-EGFP-C1-HsDcp1a	Puromycin

Plasmid	Use	Source	Mammalian Selection
Pat1b-GFP	Cloning	(131)	N/A
pLJM1-GFP-Pat1b	Co-IP	Cloned into pLJM1 using NheI and BamHI from GFP-Pat1b	Puromycin
pMD2.G	Lentivirus generation	Addgene #12259	N/A
psPAX2	Lentivirus generation	Addgene #12260	N/A
pTRE2 Firefly Luciferase-ARE	ARE-mRNA stability	(66)	N/A
pTRE2-Renilla Luciferase	ARE-mRNA stability	(66)	N/A

1264

1265   Table 3: shRNA target sequences

shRNA target	shRNA sequence 5'→3'	Region
Non-targeting (NS)	AGCACAAGCTGGAGTACAACTA	
Atg14	GTCTGGCAAATCTTCGACGAT	CDS
Atg5	CTTTGATAATGAACAGTGAGA	CDS
NDP52(1)	GAGCTGCTTCAACTGAAAGAA	CDS
NDP52(2)	GACTTGCCTATGGAAACCCAT	CDS
NDP52(3)	CCCTTTGTGAACTAAGTTCAA	3'-UTR
NDP52(4)	CCTGACTTGATACTAAGTGAT	3'-UTR
Optineurin (1)	GCACGGCATTGTCTAAATATA	CDS
Optineurin (2)	GCCATGAAGCTAAATAATCAA	CDS
p62	CCGAATCTACATTAAAGAGAA	CDS
NBR1 (1)	GCCAGGAACCAAGTTTATCAA	CDS
NBR1 (2)	CCATCCTACAATATCTGTGAA	CDS
VCP (1)	ACCGTCCCAATCGGTTAATTG	CDS
VCP (2)	AGATCCGTCGAGATCACTTTG	CDS

1266

1267   Table 4: Cloning Primers

Primer	Primer sequence
NDP52-EcoRI-Forward	5' ACCGGTCCATGGAGGAGACCATCAAA 3'
NDP52-BamHI-Reverse	5' CCGGTGGATCCTCAGAGAGA 3'

1268

1269

1270   Table 5: Antibodies

Antibody	Species	Vendor/Catalogue #	Application	Dilution
Hedls	Mouse	Santa Cruz sc-8418	Immunofluorescence Immunoblot	1:1000 1:1000
KapB	Rabbit	A generous gift from Don Ganem	Immunofluorescence	1:1000
NDP52	Rabbit	CST 60732	Immunoblot	1:1000
p62	Rabbit	CST 7695	Immunoblot	1:1000
Dcp1a	Mouse	CST 15365	Immunoblot	1:500
DDX6	Rabbit	Bethyl A300-461	Immunoblot	1:1000

Pat1b	Rabbit	CST 14288	Immunoblot	1:500
LC3B	Rabbit	CST 2775	Immunofluorescence Immunoblot	1:200 1:1000
Atg5	Rabbit	CST 2630	Immunoblot	1:1000
Atg14	Rabbit	CST 96752	Immunoblot	1:1000
OPTN	Rabbit	Abcam ab23666	Immunoblot	1:1000
Xrn1	Rabbit	Abcam ab231197	Immunoblot	1:1000
VCP	Rabbit	CST 2649	Immunoblot	1:1000
NBR1	Rabbit	CST 9891	Immunoblot	1:1000
LANA	Rabbit	A generous gift from Don Ganem	Immunofluorescence	1:1000
Phospho-S6 (Ser235/236)	Rabbit	CST 4858	Immunoblot	1:1000
Total S6	Rabbit	CST 2217	Immunoblot	1:1000
Phospho-Beclin 1 (Ser90)	Rabbit	CST 86455	Immunoblot	1:500
Total Beclin 1	Mouse	CST 4122	Immunoblot	1:1000
Phospho-ULK (Ser555)	Rabbit	CST 5869	Immunoblot	1:1000
Phospho-ULK (Ser757)	Rabbit	CST 14202	Immunoblot	1:1000
Total ULK	Rabbit	CST 8054	Immunoblot	1:1000
FLAG	Rabbit	CST 14793	Immunoblot	1:1000
FLAG	Mouse	CST 8146	Immunoprecipitation	1:100
GFP	Rabbit	CST 2956	Immunoblot	1:1000

1271

1272   Table 6: qPCR primers

Primer	Primer Sequence
HPRT-1- Forward	5' CTTTCCTTGGTCAGGCAGTATAA 3'
HPRT-1 - Reverse	5' AGTCTGGCTTATATCCAACACTTC 3'
18S - Forward	5' TTCGAACGTCTGCCCTATCAA 3'
18S - Reverse	5' GATGTGGTAGCCGTTTCTCAGG 3'
IL-6 - Forward	5' GAAGCTCTATCTCGCCTCCA 3'
IL-6 - Reverse	5' TTTTCTGCCAGTGCCTCTTT 3'
CXCL8 - Forward	5' AAATCTGGCAACCCTAGTCTG 3'
CXCL8 - Reverse	5' GTGAGGTAAGATGGTGGCTAAT 3'
IL-1β - Forward	5' CTCTCACCTCTCCTACTCACTT 3'
IL-1β - Reverse	5' TCAGAATGTGGGAGCGAATG 3'
COX-2 - Forward	5' CCCTTGGGTGTCAAAGGTAA 3'
COX-2 - Reverse	5' GCCCTCGCTTATGATCTGTC 3'

1273

Supplemental Figure Legends

Figure S1: KapB expression promotes PB disassembly but does not prevent PB assembly.

HeLa cells expressing a doxycycline-inducible GFP-Dcp1a were used to determine whether

KapB prevented granule assembly or promoted disassembly. A: HeLa cells were transfected with

either KapB or an empty vector control prior to inducing expression of GFP-Dcp1a with

doxycycline (1 µg/mL). Cells were fixed 12 h post-induction and immunostained for Hedls (PBs,

red) and KapB (blue). Scale bar=20 µm. B: GFP-Dcp1a expression was induced with

doxycycline (1 µg/mL) in HeLa cells prior to transfection with either KapB or an empty vector

control. Cells were fixed 12 h post-transfection and immunostained for Hedls (PBs, red) and

KapB (blue). Scale bar=20 µm.

Figure S2: Canonical autophagy knockdown prevents KapB-mediated PB disassembly.

A&B: HUVECs were sequentially transduced: first with recombinant lentiviruses expressing

either shRNAs targeting Atg5 or Atg14 (shAtg5, shAtg14) or a non-targeting control (NS) and

selected with puromycin (1 µg/mL), and second with either KapB or an empty vector control and

selected with blasticidin (5 µg/mL). Samples were lysed in 2x Laemmli and resolved by SDS-

PAGE. Samples were immunoblotted for Atg5 (A) or Atg14 (B). For Atg5 (A), representative

blot is from the same membrane, hashed line indicates skipped lanes. For Atg5 (A) samples were

quantified by normalizing Atg5 protein levels to the total protein in each lane using Image Lab

(Biorad) and then to Vector NS. Results were plotted in GraphPad, a 2-way ANOVA was

performed, ±SEM; n=3, ***=$P<0.001$. C: Atg5 +/+ (WT) and -/- (KO) MEFs were transduced as

in A and harvested in 2x Laemmli buffer before being resolved by SDS-PAGE. Samples were

probed for LC3, a representative blot is shown. D: Atg5 +/+ and -/- MEFs were transduced with

recombinant viruses expressing KapB or an empty vector control and selected with blasticidin (5

µg/mL). Samples were fixed in 4% paraformaldehyde and permeabilized in 0.1% Triton X-100.

Immunofluorescence was performed for Hedls (PBs, white) and DAPI (nuclei, blue). Scale

bar=20 µm. Hedls puncta were quantified using CellProfiler and presented as number of Hedls

puncta per cell, all cells counted are displayed. Results were plotted in GraphPad and a 2-way

ANOVA was performed on the main column effects with a Tukey's multiple comparison test,

bar represents the mean; n=3, **=P<0.01.

1303

Figure S3: Canonical autophagy shRNA silencing in HeLa cells. A&B: HeLas were

transduced with recombinant lentiviruses expressing either shRNAs targeting Atg5 or Atg14

(shAtg5, shAtg14) or a non-targeting control (NS) and selected with puromycin (1 µg/mL) prior

to transfection with KapB or an empty vector for luciferase assays. Samples were lysed in 2x

Laemmli buffer and resolved by SDS-PAGE before immunoblotting with Atg5 (A) or Atg14 (B).

Representative blots are shown.

1310

Figure S4: Selective autophagy receptor knockdown in HUVECs reveals that OPTN and

p62 are not required for KapB-mediated PB disassembly. A&B: HUVECs were transduced

with shRNAs targeting VCP, NBR1, or a non-targeting control (NS) and selected with

puromycin (1 µg/mL), then transduced with an empty vector control or KapB and selected with

blasticidin (5 µg/mL). Samples were lysed in 2x Laemmli buffer, resolved by SDS-PAGE, and

immunoblotted for NBR1 (A) and VCP (B). Representative blots are shown, successful VCP

knockdown was lethal, while NBR1 knockdown was not successful. C: HUVECs were

transduced with shRNAs targeting NDP52 or a non-targeting control (NS) and selected with

puromycin (1 µg/mL), then transduced with an empty vector control or KapB and selected with

blasticidin (5 µg/mL). Samples were lysed in 2x Laemmli buffer, resolved by SDS-PAGE, and

immunoblotted for NDP52. D&E: HUVECs were transduced with shRNAs targeting p62,

OPTN, or a non-targeting control (NS) and selected with puromycin (1 µg/mL), then transduced

with an empty vector control or KapB and selected with blasticidin (5 µg/mL). Samples were

lysed in 2x Laemmli buffer and resolved by SDS-PAGE. Samples were immunoblotted for p62

(D) and OPTN (E), representative blots are shown. F: HUVECs were transduced as in D and E.

Coverslips were fixed in 4% paraformaldehyde, permeabilized in 0.1% Triton X-100 and

immunostained for Hedls (PBs; white), DAPI (nuclei, blue). Scale bar=20 µm. Hedls puncta

were quantified using CellProfiler and presented as number of Hedls puncta per cell, all cells

counted are displayed. Results were plotted in GraphPad and a 2-way ANOVA was performed

on the main column effects with a Tukey's multiple comparison test, bar represents the mean;

n=3, **=P<0.01. G: HUVECs were transduced with recombinant lentiviruses targeting NDP52

or a non-targeting control (NS) and selected with puromycin (1 µg/mL). Samples were lysed in

2x Laemmli buffer, resolved by SDS-PAGE, and immunoblotted for NDP52 to confirm

knockdown for paired Torin immunofluorescence experiments.

1335

1336   **Figure S5: shRNA silencing of selective autophagy receptors in HeLa cells.** A, B&C: HeLas

1337   were transduced with shRNAs targeting NDP52, OPTN, p62, or a non-targeting control (NS) and

1338   selected with puromycin (1 µg/mL) prior to transfection with KapB or an empty vector for

1339   luciferase assays. Samples were lysed in 2x Laemmli buffer, resolved by SDS-PAGE, and

1340   immunoblotted for NDP52 (A), p62 (B), and OPTN (C). Representative blots are shown.

1341

Figure S6: NDP52 shRNA silencing and rescue in control HUVECs. A: Representative

western blot of HUVECs transduced with an shRNA targeting the 3'UTR of NDP52 or a non-

targeting control (NS) and selected with puromycin (1 µg/mL), then co-transduced with an

empty vector control and mCherry (mCh) or KapB and mCh. Samples were lysed in 2x

Laemmli, resolved by SDS-PAGE, and immunoblotted for NDP52. B: HUVECs were

sequentially transduced first with recombinant lentivirus expressing shDNP52 targeting the 3'-

UTR of NDP52 or a NS control and selected with blasticidin (5 µg/mL), and second, with vector

and either an mCherry control (mCh) or RFP-NDP52. Coverslips were fixed with 4% v/v

paraformaldehyde, permeabilized with 0.1% Triton X-100, and were immunostained with Hedls

(PBs, green). Scale bar=20 µm. C: Hedls puncta were quantified using CellProfiler and presented

as number of Hedls puncta per cell, all cells counted are displayed. Results were plotted in

GraphPad and a 2-way ANOVA was performed on the main column effects with a Tukey's

multiple comparison test, bar represents the mean; n=3.

1355

A

Vector | KapB
NS shAtg5 | NS shAtg5

Atg5

Total Protein

Atg5 protein levels
(normalized to total protein
and Vector NS)

*

● Vector
■ KapB

NS   shAtg5

B

Vector | KapB
NS shAtg14 | NS shAtg14

Atg14

Total Protein

C

WT | KO
V B | V B

LC3-I
LC3-II

Total Protein

D

Atg5 +/+   Zoom   Atg5 -/-   Zoom

Vector

KapB

Hedls puncta/cell

**   ns

Vector KapB   Vector KapB
Atg5 +/+   Atg5 -/-