**Sexual dimorphism in gene expression: coincidence and population genomics of two forms of differential expression in *Drosophila melanogaster***

**Amardeep Singh[1,2,3] and Aneil F. Agrawal[1]**

[1]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Canada, M5S 3B2

[2]Department of Integrative Biology, University of Texas at Austin, Austin, TX, 78712

[3]E-mail: amardeep.singh@utexas.edu

**Abstract**

In most species, selection favours different phenotypes in the two sexes. This type of sexual

antagonism can be resolved through the evolution of sexual dimorphism. Sex differences in gene

regulation is a proximate mechanism by which this resolution can be achieved. One form of

differential gene regulation is sex differences in the amount a gene is expressed, so called sex-

biased gene expression (SBGE). Less attention has been given to sexual dimorphism in isoform

usage (SDIU), resulting from sex-specific alternative splicing, which may be another way in

which conflict between the sexes is resolved. Here, we use RNA-seq data from two tissue types

(heads and bodies) from 18 genotypes of adult *Drosophila melanogaster* to investigate SDIU. In

our data, SBGE and SDIU are both much more prevalent in the body than the head. SDIU is less

common among sex-biased than unbiased genes in the body, though the opposite pattern occurs

in the head.  SDIU, but not SBGE, is significantly associated with reduced values of Tajima's $D$,

possibly indicating that such genes experience positive selection more frequently. SBGE, but not

SDIU, is associated with increased $\pi_N/\pi_S$, possibly indicating weaker purifying selection.

Together, these results are consistent with the idea that the SDIU and SBGE are alternative

pathways towards the resolution of conflict between the sexes with distinct evolutionary

consequences.

**Introduction**

      In sexual populations, the sexes employ different reproductive strategies to maximize fitness and as result, shared phenotypes often exhibit divergent, sex-specific trait optima. Because the sexes largely share a genome, the genetic basis of such traits is often the same between the sexes. Thus, genetic variation underpinning such shared traits will often experience divergent, sexually antagonistic selection resulting in an evolutionary conflict between the sexes termed 'intra-locus sexual conflict' (Arnqvist and Rowe 2005; Bonduriansky and Chenoweth 2009). Intra-locus sexual conflict has been well documented in a variety of plant (e.g., Hersh et al. 2015; Zemp et al. 2016) and animal (e.g., Arnqvist and Rowe 2005; Cox and Calsbeek 2009) species and may be an important mechanism in maintaining genetic variation (Kidwell et al. 1977; Rice 1984). Ultimately, intra-locus sexual conflict is thought to be resolved through the evolution of sexual dimorphism (Lande 1980; Rice 1984).

      Gene expression can be viewed as a trait in its own right or as a proximate mechanism affecting 'traditional' traits (e.g., morphological, behavioural, or physiological). Over the last ~15 years, there have been an explosion of studies investigating sexual dimorphism in gene expression, typically called 'sex-biased gene expression', SBGE (Ellegren and Parsch 2007; Parsch and Ellegren 2013; Mank 2017). SBGE refers to sex differences in the abundance of all transcripts mapping to a given gene. In addition to dimorphism in total expression of a gene, the sexes may differ in other aspects of the transcriptome. Here we focus on sexual dimorphism in isoform usage (SDIU).

      Differential isoform usage due to alternative splicing of mRNA transcripts results in the production of different isoforms from the expression of a single gene. Differential isoform usage is a widespread phenomenon among multicellular eukaryotic organisms (Ast 2004). For example, in humans, it is estimated that over half of all expressed genes exhibit alternative splicing (Mironov et al. 1999; International Human Genome Sequencing Consortium 2001). Because the function of proteins and other gene products depends on the specific exons retained in mRNA transcripts after transcription, differential isoform usage has the potential to greatly expand the functional repertoire of the genome. Indeed, in humans, the majority of genes exhibiting alternative splicing are thought to cause functionally significant alterations to gene products (Modrek et al. 2001; Modrek and Lee 2002; Sorek et al. 2004). Thus, differential

isoform usage can allow a single gene to alter or take on different functional roles across

developmental stages (e.g., Gibilisco et al. 2016) tissue types (e.g., Barberan-Soler and Zahler

2008; Telonis-Scott et al. 2009; Revil et al. 2010; Ramani et al. 2011; Tan et al. 2013) and

abiotic environments (Long et al. 2013; Jakšić and Schlötterer 2016). Across the sexes,

differential isoform usage could thus allow the sexes to express their optimal trait values through

the production of different gene products despite sharing a genome.

Widespread sexual dimorphism in isoform usage has been documented in humans

(Trabzuni et al. 2013) and other primates (Blekhman et al. 2010). A recent study documented

dimorphism in isoform usage in birds (Rogers et al. 2020). Among genes exhibiting SBGE, they

found a more even usage of alternative isoforms in the sex where the gene was more highly

expressed, a pattern that they interpreted as evidence of sex-specific selection driving SDIU.

Several microarray and RNA-seq studies across *Drosophila* species have also investigated sexual

dimorphism in isoform abundance and exon usage. These studies have generally supported a view

that sexual dimorphism in isoform usage is widespread (e.g., McIntyre et al. 2006; Telonis-Scott

et al. 2009; Chang et al. 2011), tissue-specific (Telonis-Scott et al. 2009), and evolutionarily

conserved (Gibilisco et al. 2016).

Here, we further explore SDIU using publicly available RNA-seq data from two tissue

types (head and body tissue) of 18 genotypes of adult *Drosophila melanogaster.* As SBGE and

SDIU can represent different forms of sex-differential gene regulation, we are interested in their

relationship. Resolution or mitigation of intra-locus sexual conflict may tend to require the

evolution of both SBGE and SDIU; on the other hand, SDIU and SBGE may serve as alternative

routes to mitigating sexual conflict so that the evolution of one form of sex-differential

regulation obviates selection for the other. We investigate the co-occurrence of SDIU and SBGE,

finding SDIU is less common among sex-biased than unbiased genes in the body, though the

opposite pattern occurs in the head.

Our other avenue of investigation focuses on population genetic properties of genes

exhibiting sex-differential regulation. This follows in spirit from several previous studies,

including in seed beetles (Sayadi et al. 2019) and several bird species (Wright et al. 2019), that

have examined population genomic properties such as Tajima's $D$ and $\pi_N/\pi_S$ in an attempt to

infer differences in the form of selection acting on sex-biased versus unbiased genes. We take a

similar approach but use it to compare how the two different forms of expression dimorphism—

SBGE and SDIU—are associated with the population genomic metrics used in searching for

signatures of sexual conflict resolution. Specifically, we investigate how SBGE and SDIU are

related to Tajima's $D$ and $\pi_N/\pi_S$, finding a reduction in Tajima's $D$ at genes exhibiting SDIU and

a positive relationship between $\pi_N/\pi_S$ and SBGE.

**Methods**

Obtaining RNA-seq Data and Read Mapping

We obtained publicly available raw RNA-seq data from the DNA Data Bank of Japan

Sequence Read Archive (Accession number: DRA002265). Detailed methods on animal

husbandry, RNA extraction and sequencing can be found in Osada et al. (2017). Briefly, males

from 18 randomly chosen inbred lines of the *Drosophila* Genomic Resource Panel (DGRP;

MacKay et al. 2012; Huang et al. 2014) were crossed to females from a single inbred strain from

a West African population. RNA was extracted from heads and bodies (i.e., whole body minus

heads) of 100 virgin F1 males and females from each cross. RNA from female heads and bodies

was obtained and sequenced from two replicates of each of the 18 DGRP lines while only a

single replicate was available for males. We therefore arbitrarily retained sequence data only for

female samples labelled as 'replicate 1'. Reads were inspected for quality using *FastQC*

(v0.11.9; Andrews 2010) and *MultiQC* (v0.8.3; Ewels et al. 2016). Reads were trimmed for

adapter content using *Trimmomatic* (v0.39; Bolger et al. 2014). RNA-seq reads were then

mapped to the *Drosophila melanogaster* reference genome (release 6.32; Dos Santos et al. 2015)

using *STAR* (v2.7.1a; Dobin et al. 2013) with default settings.

Differential Exon Usage Analysis

To identify a list of genes exhibiting SDIU, we performed differential exon usage

analysis using the R package *JunctionSeq* (v1.17; Hartley and Mullikin 2016). Briefly,

*JunctionSeq* splits genes into annotated exons and splice junction 'counting bins' and counts the

number of reads that overlap each counting bin within an annotated gene. To test for differential

exon usage between the sexes, a generalized linear model (GLM) with logarithmic link function

was fit at each gene feature (i.e., exon or splice junction) to assess sex differences in the number

of reads that map to a particular gene feature relative to the total number of reads that map to the

gene to which the feature belongs. Specifically, for each gene, *JunctionSeq* fits the model

$$\log(y_{ij}) = \mu + W_i + X_j + Z_{ij}$$

where $y_{ij}$ is the normalized count of reads from the $i^{th}$ sex mapping to the $j^{th}$ gene

feature/counting bin. $\mu$ is the experiment wide, mean normalized gene expression, $W_i$ reflects the

expected number of reads mapping to the focal gene in sex $i$, $X_j$ reflects the proportion of reads

mapping to counting bin $j$, and $Z_{ij}$ is an interaction term that reflects differences between the

sexes in the proportion of reads mapping to counting bin $j$. A significant effect of the interaction

term was taken as evidence of SDIU. To test the significance of the interaction term, the above

model was compared to a null model that was identical, but which omits the interaction term.

We restricted our analyses to genes with at least two annotated exons and to exons where

at least an average of 50 reads mapped in each sex. To correct for multiple testing, *P*-values were

adjusted using the Benjamini and Hochberg procedure (Benjamini and Hochberg 1995). Genes

were considered to show evidence for sexual dimorphism in isoform usage if the sexes exhibited

statistically significant differential usage of a single exon (or more) within a gene using an

adjusted *P*-value cut off of $P < 0.01$.

Ideally, genotype would be included in the model as a random effect but *JunctionSeq*

does not allow this. This is unlikely to be a major issue here. Our dataset is perfectly balanced

with each genotype being represented by only a single sample for each sex for each tissue (head

and body data were analyzed separately). More importantly, our goal is to identify a list of genes

enriched for SDIU for further analysis; our focus is not on the accuracy of *P*-values of individual

genes. As a simple check of the *JunctionSeq* pipeline, we analyzed 10 datasets in which we

switched the male/female labels for a randomly chosen half of the genotypes. In these permuted

data sets, no genes were identified as having SDIU at an adjusted *P*-value cut off of $P < 0.01$.

Gene Expression Analysis

We estimated differential gene expression between the sexes using the R package

*DESeq2* (v1.28.1; Love et al. 2014). Briefly, *DESeq2* estimates normalized read counts mapping

to genes on a $\log_2$ scale and uses a GLM framework to test for differential gene expression

between experimental conditions. Before performing differential gene expression analysis, we

performed a 'pre-filtering' procedure to remove genes that had very low numbers of reads

mapping. Pre-filtering RNA-seq data has been shown to substantially increase power in detecting

differential gene expression by reducing the burden of multiple testing but can also result in an

increase in Type I error (i.e., increased false-positive rate) (Bourgon et al. 2010). To balance

these competing issues, we removed any genes that had fewer than an average of 50 reads

mapping from all 36 samples (i.e., 18 male and 18 female genotypes). We then estimated the $\log_2$

fold change ($\log_2$FC) in male to female gene expression for each annotated gene and retained

only genes that were tested for differential exon usage. In downstream analyses, we assigned

each gene to bins based on either their average expression of the gene or based on the $\log_2$FC in

male to female gene expression. To assign genes to quantile bins based on average expression,

we determined the sex-averaged gene expression of each gene and binned genes into three equal

quantiles of increasing sex-averaged gene expression. Separately, we assigned each gene into

bins of sex-bias in gene expression. We assigned each gene to one of five bins of predefined

ranges of $\log_2$FC in gene expression ranging from highly female biased ($\log_2$FC < -2),

moderately female biased ($-2 \leq \log_2$FC $\leq -0.5$), unbiased ($0.5 \leq \log_2$FC $\leq 0.5$), moderately male

biased ($2 \geq \log_2$FC $\geq -0.5$) and strongly male biased ($\log_2$FC $\geq 2$). This procedure was

conducted separately for genes expressed in heads and bodies. Genes for which the 95%

confidence interval (CIs) in $\log_2$FC overlapped with multiple bins were excluded from analyses

using this binning procedure.

<u>Population Genomic Data</u>

We obtained whole genome sequences for 205 genotypes of the DGRP from the

*Drosophila* Genome Nexus (DGN; Lack et al. 2015, 2016). FASTA files were obtained from the

DGN website (https://www.johnpool.net/genomes.html) and were filtered to remove sites with

evidence of high relatedness and recent admixture using Perl scripts provided on the DGN

website. Two files were then created from a multi-FASTA file containing all 205 DGRP

genotypes using *SNP-sites* (v2.5.1; Page et al. 2016). The first is a typical VCF file containing

only biallelic sites. The second file was formatted identically to a VCF file but contained

genotypic information at both variant and invariant sites. Coordinates for both VCF files were

converted to release 6 of the *Drosophila* reference genome using the *UCSC liftOver* (Kent et al.
2002) tool.

<u>Estimating $\pi_N/\pi_S$ and Tajima's $D$</u>

To estimate $\pi_N/\pi_S$, per gene, we obtained a GFF file for the *D. melanogaster* genome
from Ensembl (https://useast.ensembl.org/Drosophila_melanogaster/) and scored each coding
site in the genome as 4-fold (i.e., synonymous sites) or 0-fold (i.e., non-synonymous sites)
degenerate across the *Drosophila* genome using a Python script (script available at
https://github.com/tvkent/Degeneracy). We then parsed our VCF containing variant and invariant
sites in the DGRP into two files that contained either only 4-fold sites or only 0-fold degenerate
sites. Separately for files containing 4-fold and 0-fold degenerate sites, we estimated per site
nucleotide diversity (Nei and Li 1979) using *VCFtools* (v0.1.17; Danecek et al. 2011). We then
used *bedtools* (v2.25.0 Quinlan and Hall 2010) to map 4-fold and 0-fold degenerate sites to
genes and estimated the mean $\pi_N$ within a gene as the mean nucleotide diversity at 0-fold
degenerate sites within each gene and $\pi_S$ within a gene as the mean nucleotide diversity at 4-fold
degenerate sites within each gene. Genes with fewer than 30 4-fold degenerate (i.e.,
synonymous) sites were filtered out from further analyses.

We used a similar approach to estimating gene-wide Tajima's $D$ (Tajima 1989).
Specifically, we identified biallelic SNPs at 4-fold degenerate sites in the DGRP VCF file
produced above. We then calculated Tajima's $D$ at each 4-fold degenerate site. 4-fold
degenerate, segregating sites were then mapped to genes in the *D. melanogaster* genome and
Tajima's $D$ was averaged over all 4-fold degenerate sites within each gene. Genes with fewer
than 30 4-fold degenerate sites were filtered out from further analyses.

<u>Gene Ontology (GO) Enrichment Analysis</u>

We tested for an enrichment of gene ontology (GO) terms among genes exhibiting
statistically significant SDIU. This analysis was conducted using *GOrilla* (Eden et al. 2009)
which uses a hypergeometric model to test for an enrichment of GO terms in a target set of genes
relative to a user supplied background set of genes. This analysis was conducted separately for
head and body tissue. In both analyses the target set of genes was compared to a background list
that was composed of genes that had been tested in the differential exon usage analysis

conducted with *JunctionSeq* but did not exhibit statistically significant SDIU. *P*-values for GO

terms were converted to *Q*-values and a *Q*-value threshold of 0.01 was used to assess statistical

significance.

<u>Statistical Analysis</u>

We used linear regression analysis to examine the relationship between population

genomic measures (i.e., Tajima's *D* and $\pi_N/\pi_S$) and the fixed effects of sex-averaged gene

expression, sex-bias in gene expression (measured as the absolute value the log$_2$FC of male to

female gene expression) and SDIU. To improve the distributional properties for statistical

analysis, we analyzed $\log(\pi_N/\pi_S)$ rather than $\pi_N/\pi_S$. This required the exclusion of genes where

$\pi_N = 0$, which only represented 6.01% and 5.51% of genes expressed in the body and head

respectively. For both Tajima's *D* and $\log(\pi_N/\pi_S)$, we fit a regression model using the '*lmperm*'

function of the *R* package *permuco* (v.1.1.0; Frossard and Renaud 2019) which uses a Freedman-

Lane permutation method (Freedman and Lane 1983) and type III Wald's chi-square test to

estimate *P*-values for fixed effects. *P*-values were estimated after 10,000 permutated model fits.

To ensure that the exclusion of genes with $\pi_N = 0$ did not fundamentally change the

interpretation of the data, we performed a separate logistic regression analysis including these

genes in which we treated $\pi_N/\pi_S$ as a binary trait (i.e., $x = 0$ representing $\pi_N/\pi_S = 0$ and $x = 0$

representing $\pi_N/\pi_S = 1$). The results of this analysis are described in Table S3 and add further

support to the interpretation of the main analysis.

**Results and Discussion**

<u>Identity and Functional Properties of Genes Exhibiting Sexually Dimorphic Exon Usage</u>

We find evidence of extensive sexual dimorphism in isoform usage (SDIU or 'isoform

dimorphism') among multi-exon genes expressed in both head and body tissue of adult

*Drosophila melanogaster*. Table 1 summarizes the results of our differential exon usage analysis.

Of multi-exon genes tested, 3420 and 539 genes expressed in adult body and head tissue exhibit

statistically significant SDIU respectively which accounts for 53.76% and 7.65% of all genes

tested in bodies and heads respectively (Table 1). These data suggest that SDIU usage occurs in

approximately 30.33% and 6.06% of all genes annotated in the *D. melanogaster* genome and that

are expressed in body and head tissue, respectively. Considering the union of genes expressed in

either tissue, we detected SDIU in at least one of the tissues for 29.84% of genes. These data

accord well with several previous studies that have found evidence of widespread SDIU both in

this species (e.g., McIntyre et al. 2006; Telonis-Scott et al. 2009; Chang et al. 2011) as well as

other species of *Drosophila* (Gibilisco et al. 2016). McIntyre et al. (2006) reported that 68% of

all multi-transcript genes expressed in whole body tissue exhibited significant sex-bias in the

expression of one or more alternative isoforms in a microarray study using *D. melanogaster*.

Similarly, a microarray study in this species by Telonis-Scott et al. (2009) found that ~58% of

multi-exon genes tested in whole body tissue exhibited SDIU. The lower percentage we report

here (~50% of tested genes exhibiting SDIU versus ~60-70%) may reflect differences in

technical and statistical methodologies, including the choice of false discovery rates.

**Table 1:** Summary of the differential exon usage analysis between the sexes.

|  | Genes Expressed | Genes Tested | Genes with Sexually Dimorphic Isoform Usage (% of total genes expressed) |
|---|---|---|---|
| Head | 8904 | 7006 | 540 (6.06%) |
| Body | 11,277 | 6362 | 3420 (30.33%) |
| [1]Head and Body | 11,825 | 7646 | 3528 (29.84%) |

[1]These data refer to the union of genes expressed, tested and exhibiting sexually dimorphic
isoform usage found in either head, body or both.

These results suggest heads and bodies differ substantially in the frequency of SDIU.

Although the majority of sexually dimorphic genes in head tissue was also sexually dimorphic in

body tissue, there are many genes exhibiting SDIU in body tissue where we did not detect

dimorphism in head tissue (Figure 1). An important caveat is that the power to detect differential

isoform usage is correlated with overall gene expression, indeed there is a substantial increase in

overall gene expression for genes in which we detect SDIU compared to those we do not (Figure

S1A). If this were the main driver of tissue differences we observe, then genes with similar

expression levels in both tissues should exhibit similar frequencies of SDIU. However, even for

genes with similar expression levels in head and body tissue, many more genes appear to exhibit

SDIU in body tissue than in head tissue (Figure S1B). In addition, these results comport well

with previously published data. For example, Telonis-Scott et al. (2009) reported reduced levels

of SDIU in genes when expressed in head tissue compared to carcass (i.e., head- and gonad-less

bodies) and gonadal tissue.

**Figure 1:** Venn diagrams of overlap between genes found to exhibit sexually dimorphic (SD)
exon usage when expressed in **(A)** body and **(D)** head tissue and genes with no evidence of
sexually dimorphic exon usage when expressed in **(B)** body and **(C)** head tissue. Only genes that
were tested for SDIU in both body and head are represented here.

Genes exhibiting SDIU are enriched for a variety of gene ontology (GO) terms, though

there are far more enriched GO terms for the genes identified as exhibiting SDIU in the body

(Table S1) compared to the head (Table S2). For sexually dimorphic genes expressed in head

tissue, there is an enrichment for GO terms related to cellular (including neuron specific)

components and processes as well as genes related to behaviour. Especially notable GO terms

enriched in SDIU genes in the body are GO terms associated with development and reproduction

including genes related to '*oogenesis*', '*gamete generation*' and '*germ cell generation*' as well as

genes related to cellular and organelle components and cell metabolism. The enrichment of

genes related to reproduction and development may not be surprising considering much of the

variation in isoform usage between the sexes has previously been reported as driven by gene

expression in gonadal tissue (Telonis-Scott et al. 2009; Gibilisco et al. 2016).

<u>Coincidence of two forms of sexually dimorphism in expression</u>

The majority of expression studies have examined variation in total expression for a

gene. Studies of sex-biased gene expression (SBGE) have shown repeatedly that SBGE is highly

tissue specific (e.g., Parisi et al. 2004; Yang et al. 2006; Huylmans and Parsch 2014; Dutoit et al.

2018); both our data and other studies (e.g., Telonis-Scott et al. 2009; Chang et al. 2011) indicate

that the same is true for sex differences in isoform usage. Because both of these forms of sexual

dimorphism are thought to be related to resolving intra-locus sexual conflict, we are interested in

understanding the relationship between SBGE and SDIU.

SBGE is widespread both in heads and bodies, though there is greater variation among

genes in SBGE in body tissue relative to head tissue (Figure 2A). These results are consistent

with previously published data in a number of species that find that sex-bias in gene expression

is generally more extensive in gonadal than in somatic tissue including brain tissue (e.g.,

Innocenti and Morrow 2010; Baker et al. 2011; Dutoit et al. 2018). In the head, SDIU genes are,

on average, significantly more female-biased in expression than non-SDIU genes, but there is no

difference in the body (Figure 2B).

To further explore the relationship between SDIU and SBGE, we divided genes into bins

representing different levels of sex-bias using defined ranges of $\log_2$FC in gene expression, and

only considered genes that could be assigned to bins with reasonably high certainty (see

methods). Within each bin, we then calculated the proportion of isoform dimorphic to non-

dimorphic genes in each quantile bin. For genes expressed in the body, there is a reduction in the

proportion of isoform dimorphic genes among sex-biased genes relative to unbiased genes

(Figure 2C). Conversely, for genes expressed in head tissue, female-biased and male-biased

genes are enriched for SDIU (Figure 2D). The fly body results are in line with Rogers et al.

(2020) who reported non-significant trends towards a negative relationship between SDIU and

SBGE in gonadal tissue of three bird species.

The reduction in isoform dimorphism at highly sex-biased genes expressed in the body

could be because SBGE and SDIU provide alternative routes to the resolution of sexual conflict

and thus the evolution of one reduces the selective pressure for the evolution of the other. On the

other hand, the opposite pattern emerges in genes expressed in the head with an apparent

increase in the proportion of genes exhibiting SDIU at sex-biased genes relative to unbiased

genes. One hypothesis is that the resolution of intra-locus conflict may often require both forms

of expression dimorphism whereas those genes experiencing sexually concordant selection will

evolve neither, creating a positive association between SDIU and SBGE. A somewhat more

mechanistic hypothesis for the positive association is that just as SBGE is more likely to evolve

at genes that exhibit lower pleiotropic constraint (Dean and Mank 2016), SDIU may also be able

to more easily evolve at such genes. In birds, however, Rogers et al (2020) found that genes

exhibiting SDIU exhibited greater tissue breadth in gene expression (used as an indicator of

pleiotropic constraint) than non-SDIU genes. A third, more mundane, explanation is that the

positive association of SBGE and SDIU in heads exists because of how we quantify

gene/isoform expression (i.e., counting reads mapping to genes); we may be biased towards

finding a positive association between SBGE and SIDU because we are observing the same

phenomenon in different ways. For example, if both males and females express a single isoform

of a gene at the same level but females also express an additional isoform that is not expressed

by males, we would detect both SBGE and SDIU at this gene. However, this methodological

bias must not be very strong as there are many unbiased genes that exhibit SDIU (704 in head

and 405 in body) and the association between SBGE and SDIU is negative in the body.

324
**Figure 2:** Patterns of sex-bias in gene expression (SBGE) and sexual dimorphism in isoform usage (SDIU) for genes expressed in the
body and head tissue. **(A)** Scatter plot and density distributions of $\log_2$ fold change ($\log_2$FC) in male to female gene expression of
genes exhibiting SDIU only in bodies (light blue), only in heads (red), in both heads and bodies (dark blue) or in neither heads nor
bodies (beige) when expressed in bodies (x-axis) or heads (y-axis). Density plots above and to the right of the figure correspond to the
$\log_2$FC gene expression in bodies and heads respectively. Some genes with extreme sex-bias in expression were omitted from scatter
plot, see Supplemental Figure S2 for plots with all genes. **(B)** Mean $\log_2$FC in male to female gene expression at genes exhibiting
SDIU (green) and those that do not exhibit SDIU (purple) when expressed in bodies or heads. **(C-D)** The proportion of genes
exhibiting SDIU to genes not exhibiting SDIU in bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (red),
unbiased (black) and male-biased (blue) expression when expressed in **(C)** bodies or **(D)** heads. Genes in the most strongly sex-biased
bins were excluded from plot of genes expressed in heads **(D)** because too few genes (i.e., fewer than 5) were present in these bins;
their inclusion strengthens the pattern (see Supplemental Figure S3 for plots including omitted bins). Error bars in panels B-D are 95%
confidence intervals estimated from 10,000 bootstraps resampling estimates.

Given that the X chromosome spends a disproportionate amount of its time in females,
and because of its hemizygous expression in males, there has been considerable interest in
whether it is enriched for genes that experience sexually antagonistic selection contributing to
intralocus conflict or contributing to sexual dimorphism (Rice 1987; Rowe et al. 2018; Frank and
Patten 2020) We therefore examined the genomic distribution of SDIU and SBGE genes. In the
body data, there is no evidence of an enrichment of genes exhibiting SDIU on the X
chromosome (Figure 3A). In contrast, SBGE is strongly associated with the X chromosome, with
female-biased genes being significantly overrepresented on the X chromosome relative to
unbiased genes whereas male biased genes are significantly underrepresented on the X
chromosome (Figure 3C). This observation of a "demasculinization" of the X chromosome is
consistent with previous studies of SBGE in *D. melanogaster* and several other species (Parisi et
al. 2003; Ranz et al. 2003; Khil et al. 2004; Reinke et al. 2004; Sturgill et al. 2007; Meisel et al.
2012; Albritton et al. 2014). We find no relationship between SBGE and SDIU with respect to
genomic location (Figure 3D).

These patterns are quite different for the head data. In contrast to the body data, there are
a significantly higher proportion of X-linked genes exhibiting SDIU in the head relative to non-
SDIU genes (Figure 3A). With respect to SBGE, both male- and female-biased genes are
overrepresented on the X relative to unbiased genes, with the strongest overrepresentation among
male-biased genes (Figure 3C). The over-representation of SDIU genes on the X chromosome
occurs primarily among unbiased genes rather than sex-biased genes (Figure 3E), indicating that
this pattern is not driven by a relationship between SDIU and SBGE.

**Figure 3:** Proportion of genes exhibiting sexually dimorphic isoform usage (SDIU) and sex-biased gene expression (SBGE) that are X-linked. **(A)** The proportion of total SDIU (green) and non-SDIU (purple) genes that are found on the X chromosome when expressed in body or head tissue. **(B-C)** The proportion of genes found on the X chromosome in bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue) expression when expressed in **(B)** bodies or **(C)** heads. **(D-E)** The proportion of genes exhibiting SDIU (green) and not exhibiting SDIU (purple) found on the X chromosome in bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (left most pair of bins), unbiased (middle bin) and male-biased (right most pair of bins) expression when expressed in **(D)** bodies or **(E)** heads. Genes in the most strongly sex-biased bins were excluded from plot of genes expressed in heads (i.e., panels C and E) because too few genes (i.e., fewer than 5) were present in these bins (see Supplemental Figure S4 for plots including omitted bins). Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

Population genomic patterns associated with SDIU and SBGE

To explore patterns of genetic diversity at genes with and without SDIU, we used two

population genetic metrics – Tajima's $D$ and $\pi_N/\pi_S$ – to look for clues about the form of

selection that have acted at these genes. Tajima's $D$ is a useful summary statistic of the site

frequency spectrum; selective sweeps tend to reduce Tajima's $D$ whereas balancing selection

increases it (Charlesworth and Charlesworth 2010). Recent studies exploring the relationship

between Tajima's $D$ and sexually dimorphic gene expression have been mixed: some studies

have found a decrease in Tajima's $D$ at sex-biased genes (Wright et al. 2018, 2019) while others

have reported the opposite (Dutoit et al. 2018) or a more mixed relationship (Pröschel et al.

2006; Sayadi et al. 2019). Tajima's $D$ at genes exhibiting SDIU is significantly reduced relative

to genes that do not on the basis of non-overlapping 95% confidence intervals (CIs) for genes

expressed in body and head tissue (Figure 4A). This result could indicate that balancing selection

is less common among genes with versus without SDIU; this would be consistent with the idea

that the evolution of sexually dimorphic isoform usage may represent at least a partial resolution

of sexual conflict. The Tajima's $D$ result could also indicate that selective sweeps are more

common among genes with SDIU versus those without; this would be consistent with idea that

dimorphic isoform usage removes sex-specific constraints, allowing a greater fraction of

mutations to be beneficial.

Previous studies have found that some patterns of molecular variation are correlated with

expression level. For example, highly expressed genes tend to have lower sequence variation

presumably due to stronger purifying selection (e.g., Pál et al. 2001; Subramanian and Kumar

2004; Carneiro et al. 2012; Paape et al. 2013; Williamson et al. 2014; Ho et al. 2019). To

visualize whether expression level might be a confounding factor, all genes (for a given tissue)

were first binned into equal quantiles of sex-averaged gene expression; within these bins

Tajima's $D$ tends to be lower for SDIU than non-SDIU genes (Figure 4B; Figure 4C). As others

have done (Sayadi et al. 2019; Wright et al. 2019), we also looked for a relationship between

Tajima's $D$ and sex-biased gene expression but found no obvious pattern (Figure 4E; Figure 4F).

To examine these relationships in a more statistically rigorous framework, we fit linear

models, separately for body and head tissue, modeling Tajima's $D$ as a function of sex-averaged

gene expression, sex-bias in gene expression (measured as the absolute value of the $\log_2$ fold

change of male to female gene expression) and sexual dimorphism in isoform usage (i.e., genes

that do or do not exhibit statistically significant SDIU). Consistent with the patterns described

above, for the body data, there is a significant negative relationship between Tajima's $D$ and

SDIU (Table 2). There is also a significant negative relationship with sex-averaged gene

expression but no significant relationship with sex-bias in gene expression. For genes expressed

in head tissue, the patterns are similar but marginally non-significant (Table 2), although the

direction of the effect of average gene expression and SDIU are in the same direction as for the

body data. The head dataset includes many fewer SDIU genes, so power is reduced.

**Figure 4:** Patterns of Tajima's *D* and sexual dimorphism in isoform usage (SDIU). Mean Tajima's *D* for genes exhibiting SDIU (green) or not (purple) **(A)** averaged across all genes expressed in body and head tissue. **(B, C)** Mean Tajima's *D* for genes expressed in **(B)** body or **(C)** head tissue within quantiles of sex-averaged gene expression. **(D, E)** Mean Tajima's *D* for genes expressed in **(D)** body or **(E)** head tissue averaged across genes within bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue). Genes in the most strongly sex-biased bins were excluded from plot of genes expressed in heads **(E)** because too few genes (i.e., fewer than 5) were present in these bins (see Supplemental Figure S5 for plots including omitted bins). Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

417 **Table 2:** Parameter estimates from linear model fit to test the fixed effects of overall gene expression, sex-bias in gene expression
418 (i.e., log$_2$ fold change in male to female gene expression) and sexual dimorphism in isoform usage (SDIU) on Tajima's D in body and
419 head tissue. *P*-values were obtained by a Wald's Type III Chi-square test. Bold terms indicate statistically significant effects at $\alpha$ =
420 0.05.

| Tissue | Effect | Estimate | Standard Error | *P* |
|---|---|---|---|---|
| Body Tissue | **Intercept** | **0.061** | **0.014** | **< 0.001** |
| | **Sex-averaged Gene Expression[1]** | **-2.01 x 10$^{-6}$** | **9.01 x 10$^{-7}$** | **0.025** |
| | Absolute value of SBGE[2] | -0.005 | 0.007 | 0.469 |
| | **Sexual Dimorphism in Isoform Usage (SDIU)[3]** | **-0.031** | **0.013** | **0.022** |
| Head Tissue | **Intercept** | **0.034** | **0.008** | **< 0.001** |
| | Sex-averaged Gene Expression | -6.53 x 10$^{-7}$ | 4.70 x 10$^{-7}$ | 0.154 |
| | Absolute value of SBGE[1] | 0.025 | 0.025 | 0.301 |
| | Sexual Dimorphism in Isoform Usage (SDIU) | -0.045 | 0.024 | 0.064 |

[1]Gene expression averaged across both sexes.
[2]Measured as |Log$_2$FC male:female gene expression|.
[3]Indicator of whether or not a gene exhibited statistically significant evidence of SDIU.

We also examined $\pi_N/\pi_S$, using the ratio of nucleotide diversity at 0-fold non-

synonymous sites ($\pi_N$) to 4-fold synonymous sites ($\pi_S$), which can be useful in identifying

differences in the strength of purifying selection. Overall, we do not find a significant difference

in $\pi_N/\pi_S$ between genes exhibiting SDIU relative to other genes expressed in either head or

body tissue (Figure 5A). $\pi_N/\pi_S$ tends to be lower in genes that are more highly expressed (Figure

5B; Figure 5C), but there are no obvious differences between SDIU and non-SDIU genes when

they are binned into quantiles of sex-averaged expression. This pattern of reduced $\pi_N/\pi_S$ in more

highly expressed genes has been observed in other studies and has been interpreted as resulting

from stronger purifying selection acting at highly expressed genes (e,g., Pál et al. 2001;

Subramanian and Kumar 2004; Carneiro et al. 2012; Paape et al. 2013; Williamson et al. 2014;

Ho et al. 2019).

In contrast to SDIU, SBGE is related to $\pi_N/\pi_S$, with higher $\pi_N/\pi_S$ values associated with

sex-biased genes, at least for genes expressed in the body (Figure 5D). Studies in several species

including seed beetles (Sayadi et al. 2019), several bird species (Harrison et al. 2015; Dutoit et

al. 2018) , guppies (Wright et al. 2018), and humans (Gershoni and Pietrokovski 2014) have

shown an inflation of $\pi_N/\pi_S$ at sex-biased genes. Elevated $\pi_N/\pi_S$ for sex-biased genes may

indicate weaker purifying selection. There also appears to be a negative relationship between

$\pi_N/\pi_S$ and overall gene expression (i.e., sex-averaged gene expression) whereby highly

expressed genes appear to have lower values of $\pi_N/\pi_S$. To test these relationships statistically,

we fit linear models similar to those described above separately for heads and bodies. Regression

analysis with the body dataset finds a statistically significant negative relationship between

$\pi_N/\pi_S$ and sex-averaged gene expression as well as SBGE, but no significant effect of SDIU on

$\pi_N/\pi_S$ (Table 3).

**Figure 5:** Patterns of $\pi_N/\pi_S$ and sexual dimorphism in isoform usage (SDIU). Median $\pi_N/\pi_S$ for genes exhibiting SDIU (green) or not (purple) **(A)** averaged across all genes expressed in body and head tissue. **(B, C)** Median $\pi_N/\pi_S$ for genes expressed in **(B)** body or **(C)** head tissue averaged across genes within quantiles of sex-averaged gene expression. **(D, E)** Median $\pi_N/\pi_S$ for all genes expressed in **(D)** body or **(E)** head tissue averaged across all genes within bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue). Genes in the most strongly sex-biased bins were excluded from plot of genes expressed in heads **(E)** because too few genes (i.e., fewer than 5) were present in these bins (see Supplemental Figure S6 for plots including omitted bins). Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

**Table 3:** Parameter estimates from linear model fit to test the fixed effects of overall gene expression (i.e., sex averaged gene expression), sex-bias in gene expression (i.e., $\log_2$ fold change in male to female gene expression) and sexual dimorphism in isoform usage (SDIU) on $\pi_N/\pi_S$ in body and head tissue. *P*-values were obtained by a Wald's Type III Chi-square test. Bold terms indicate statistically significant effects at $\alpha = 0.05$.

| Tissue | Effect | Estimate | Standard Error | *P* |
|---|---|---|---|---|
| Body Tissue | **Intercept** | **-2.350** | **0.004** | **< 0.001** |
| | **Sex-averaged Gene Expression[1]** | **-2.33 x 10$^{-5}$** | **3.24 x 10$^{-6}$** | **< 0.001** |
| | **Absolute value of SBGE[2]** | **0.004** | **0.014** | **0.002** |
| | Sexual Dimorphism in Isoform Usage (SDIU)[3] | -0.037 | 0.029 | 0.154 |
| Head Tissue | **Intercept** | **-2.383** | **0.022** | **< 0.001** |
| | **Sex-averaged Gene Expression[1]** | **-4.43 x 10$^{-6}$** | **91.35 x 10$^{-6}$** | **0.003** |
| | Absolute value of SBGE[2] | 0.032 | 0.066 | 0.618 |
| | Sexual Dimorphism in Isoform Usage (SDIU)[3] | 0.053 | 0.067 | 0.426 |

[1]Gene expression averaged across both sexes.

[2]Measured as |$\log_2$FC male:female gene expression|.

[3]Indicator of whether or not a gene exhibited statistically significant evidence of SDIU.

An abundance of caution is warranted in interpreting the population genomic patterns

reported here as evidence for or against particular forms of selection acting at genes exhibiting

SDIU or SBGE. Population demography (Hartl and Clark 2007; Cutter 2019), genome-wide

variation in rates of recombination (Braverman et al. 1995) and mutation (Aris-Brosou and

Excoffier 1996) are all well recognized as factors that influence patterns of genetic diversity.

Nonetheless this study provides insights into the properties of genes exhibiting two different

forms of gene regulation: sexually dimorphic isoform usage and sex-biased gene expression and

documents population genetic patterns at such genes.

SDIU and SBGE are two different ways to differentially regulate how genes are

expressed in the two sexes and are thought to be mechanisms by which sexual conflict can be

resolved (Bonduriansky and Chenoweth 2009; Stewart et al. 2010). As is well known for SBGE,

it appears that SDIU is also widespread in *Drosophila melanogaster*, particularly among genes

expressed in body tissue. Genes exhibiting SDIU are functionally diverse and involved in

processes that are likely hotspots for sexual conflict including over development and

reproduction. The negative association between SDIU and SBGE in the body suggests that to

some extent when one of these mechanisms evolves, it may negate the selective pressure to

evolve the other. However, given that we observe the opposite pattern in heads (i.e., co-

occurrence of SBGE and SDIU), it is premature to draw a strong inference.

The relationship between SDIU and SBGE in the present study suggests that these two

forms of gene expression regulation experience different forms of selection. Genes exhibiting

SDIU are negatively related to Tajima's $D$ but do not exhibit a strong relationship to $\pi_N/\pi_S$, a

pattern that is consistent with SDIU genes being more frequently subject to selective sweeps or

less frequently subject to balancing selection compared to non-SDIU genes. In contrast, SBGE is

not associated with Tajima's $D$, but is negatively associated with $\pi_N/\pi_S$, which may indicate that

that sex-biased genes experience weaker purifying selection, perhaps because selection is weak

in the sex with reduced expression (Dapper and Wade 2020). Though divergent selection

between the sexes is likely to drive the evolution of both SBGE and SDIU, the population

genomic consequences following the evolution of these alternative forms of dimorphism appears

quite different.

**Acknowledgements:** We wish to thank Tyler Kent for providing a script to identify $N$-fold

codon degeneracy as well as Julia Kreiner, and James Santangelo for helpful discussion and

bioinformatics advice. Funding was provided in the form of an Ontario Graduate Scholarship to

A.S. and an NSERC discovery grant to A.F.A.

**Literature Cited**

Albritton, S. E., A. L. Kranz, P. Rao, M. Kramer, C. Dieterich, and S. Ercan. 2014. Sex-biased gene expression and evolution of the X chromosome in nematodes. Genetics 197:865–883.

Andrews, S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.

Aris-Brosou, S., and L. Excoffier. 1996. The impact of population expansion and mutation rate heterogeneity on DNA sequence polymorphism. Mol. Biol. Evol. 13:494–504.

Arnqvist, G., and L. Rowe. 2005. Sexual Conflict. Princeton University Press, Princeton NJ.

Ast, G. 2004. How did alternative splicing evolve? Nat. Rev. Genet. 5:773–782.

Baker, D. A., T. Nolan, B. Fischer, A. Pinder, A. Crisanti, and S. Russell. 2011. A comprehensive gene expression atlas of sex- and tissue-specificity in the malaria vector, *Anopheles gambiae*. BMC Genomics 12:296.

Barberan-Soler, S., and A. M. Zahler. 2008. Alternative splicing regulation during *C. elegans* development: Splicing factors as regulated targets. PLoS Genet. 4:e1000001.

Benjamini, Y., and Y. Hochberg. 1995. Controlling the False Discovery Rate : A Practical and Powerful Approach to Multiple Testing. J. R. Stat. Soc. Ser. B 57:289–300.

Blekhman, R., J. C. Marioni, P. Zumbo, M. Stephens, and Y. Gilad. 2010. Sex-specific and lineage-specific alternative splicing in primates. Genome Res. 20:180–189.

Bolger, A. M., M. Lohse, and B. Usadel. 2014. Trimmomatic: A flexible trimmer for Illumina sequence data. Bioinformatics 30:2114–2120.

Bond[uriansky], R., and S. F. Chenoweth. 2009. Intralocus sexual conflict. Trends Ecol. Evol. 24:280–288.

Bourgon, R., R. Gentleman, and W. Huber. 2010. Independent filtering increases detection power for high-throughput experiments. Proc. Natl. Acad. Sci. U. S. A. 107:9546–9551.

Braverman, J. M., R. R. Hudson, N. L. Kaplan, C. H. Langley, and W. Stephan. 1995. The hitchhiking effect on the site frequency spectrum of DNA polymorphisms. Genetics 140:783–796.

Carneiro, M., F. W. Albert, J. Melo-Ferreira, N. Galtier, P. Gayral, J. A. Blanco-Aguiar, R. Villafuerte, M. W. Nachman, and N. Ferrand. 2012. Evidence for widespread positive and purifying selection across the european rabbit (*Oryctolagus cuniculus*) genome. Mol. Biol. Evol. 29:1837–1849.

Chang, P. L., J. P. Dunham, S. V. Nuzhdin, and M. N. Arbeitman. 2011. Somatic sex-specific transcriptome differences in *Drosophila* revealed by whole transcriptome sequencing. BMC Genomics 12:364.

Charlesworth, B., and D. Charlesworth. 2010. Elements of Evolutionary Genetics. 1st ed. Roberts and Company Publishers, Greenwood Village, CO.

Cox, R. M., and R. Calsbeek. 2009. Sexually Antagonistic Selection, Sexual Dimorphism, and the Resolution of Intralocus Sexual Conflict. Am. Nat. 173:176–187.

Cutter, A. D. 2019. A Primer of Molecular Population Genetics. Oxford University Press, Oxford.

Danecek, P., A. Auton, G. Abecasis, C. A. Albers, E. Banks, M. A. DePristo, R. E. Handsaker, G. Lunter, G. T. Marth, S. T. Sherry, G. McVean, and R. Durbin. 2011. The variant call format and VCFtools. Bioinformatics 27:2156–2158.

Dapper, A. L., and M. J. Wade. 2020. Relaxed Selection and the Rapid Evolution of Reproductive Genes. Trends Genet. 36:640–649. Elsevier Ltd.

Dean, R., and J. E. Mank. 2016. Tissue specificity and sex-specific regulatory variation permit the evolution of sex-biased gene expression. Am. Nat. 188:E74–E84.

Dobin, A., C. A. Davis, F. Schlesinger, J. Drenkow, C. Zaleski, S. Jha, P. Batut, M. Chaisson, and T. R. Gingeras. 2013. STAR: Ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.

Dos Santos, G., A. J. Schroeder, J. L. Goodman, V. B. Strelets, M. A. Crosby, J. Thurmond, D. B. Emmert, W. M. Gelbart, N. H. Brown, T. Kaufman, M. Werner-Washburne, R. Cripps, K. Broll, L. S. Gramates, K. Falls, B. B. Matthews, S. Russo, P. Zhou, M. Zytkovicz, B. Adryan, H. Attrill, M. Costa, S. Marygold, P. McQuilton, G. Millburn, L. Ponting, R. Stefancsik, S. Tweedie, and G. Grumbling. 2015. FlyBase: Introduction of the *Drosophila*

*melanogaster* Release 6 reference genome assembly and large-scale migration of genome
annotations. Nucleic Acids Res. 43:D690–D697.

Dutoit, L., C. F. Mugal, P. Bolívar, M. Wang, K. Nadachowska-Brzyska, L. Smeds, H. P. Yazdi,
L. Gustafsson, and H. Ellegren. 2018. Sex-biased gene expression, sexual antagonism and
levels of genetic diversity in the collared flycatcher (*Ficedula albicollis*) genome. Mol.
Ecol. 27:3572–3581.

Eden, E., R. Navon, I. Steinfeld, D. Lipson, and Z. Yakhini. 2009. *GOrilla*: A tool for discovery
and visualization of enriched GO terms in ranked gene lists. BMC Bioinformatics 10:48.

Ellegren, H., and J. Parsch. 2007. The evolution of sex-biased genes and sex-biased gene
expression. Nat. Rev. Genet. 8:689–698.

Ewels, P., M. Magnusson, S. Lundin, and M. Käller. 2016. MultiQC: Summarize analysis results
for multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

Frank, S. A., and M. M. Patten. 2020. Sexual antagonism leads to a mosaic of X-autosome
conflict. Evolution. 74:495–498.

Freedman, D., and D. Lane. 1983. A Nonstochastic Interpretation of Reported Significance
Levels. J. Buisness Econ. Stat. 1:292–298.

Frossard, J., and O. Renaud. 2019. *permuco*: Permutation Tests for Regression, (Repeated
Measures) ANOVA/ANCOVA and Comparison of Signals.

Gershoni, M., and S. Pietrokovski. 2014. Reduced selection and accumulation of deleterious
mutations in genes exclusively expressed in men. Nat. Commun. 5:4438.

Gibilisco, L., Q. Zhou, S. Mahajan, and D. Bachtrog. 2016. Alternative Splicing within and
between *Drosophila* Species, Sexes, Tissues, and Developmental Stages. PLoS Genet.
12:e1006464.

Harrison, P. W., A. E. Wright, F. Zimmer, R. Dean, S. H. Montgomery, M. A. Pointer, and J. E.
Mank. 2015. Sexual selection drives evolution and rapid turnover of male gene expression.
Proc. Natl. Acad. Sci. U. S. A. 112:4393–4398.

Hartl, D. L., and A. G. Clark. 2007. Principles of Population Genetics. 4th ed. Sinauer,

597  Sunderland, MA.

598 Hartley, S. W., and J. C. Mullikin. 2016. Detection and visualization of differential splicing in
599  RNA-Seq data with JunctionSeq. Nucleic Acids Res. 44:e127.

600 Hersh, E., J. A. Madjidian, S. Andersson, M. Strandh, W. S. Armbruster, and A. Lankinen. 2015.
601  Sexual antagonism in the pistil varies among populations of a hermaphroditic mixed-mating
602  plant. J. Evol. Biol. 28:1321–1334.

603 Ho, E. K. H., M. Bartkowska, S. I. Wright, and A. F. Agrawal. 2019. Population genomics of the
604  facultatively asexual duckweed *Spirodela polyrhiza*. New Phytol. 224:1361–1371.

605 Huang, W., A. Massouras, Y. Inoue, J. Peiffer, M. Ràmia, A. M. Tarone, L. Turlapati, T.
606  Zichner, D. Zhu, R. F. Lyman, M. M. Magwire, K. Blankenburg, M. A. Carbone, K. Chang,
607  L. L. Ellis, S. Fernandez, Y. Han, G. Highnam, C. E. Hjelmen, J. R. Jack, M. Javaid, J.
608  Jayaseelan, D. Kalra, S. Lee, L. Lewis, M. Munidasa, F. Ongeri, S. Patel, L. Perales, A.
609  Perez, L. L. Pu, S. M. Rollmann, R. Ruth, N. Saada, C. Warner, A. Williams, Y. Q. Wu, A.
610  Yamamoto, Y. Zhang, Y. Zhu, R. R. H. Anholt, J. O. Korbel, D. Mittelman, D. M. Muzny,
611  R. A. Gibbs, A. Barbadilla, J. S. Johnston, E. A. Stone, S. Richards, B. Deplancke, and T. F.
612  C. Mackay. 2014. Natural variation in genome architecture among 205 *Drosophila*
613  *melanogaster Genetic* Reference Panel lines. Genome Res. 24:1193–1208.

614 Huylmans, A. K., and J. Parsch. 2014. Population- and sex-biased gene expression in the
615  excretion organs of *Drosophila melanogaster*. G3 Genes, Genomes, Genet. 4:2307–2315.

616 Innocenti, P., and E. H. Morrow. 2010. The Sexually Antagonistic Genes of *Drosophila*
617  *melanogaster*. PLoS Biol. 8:e1000335.

618 International Human Genome Sequencing Consortium. 2001. Initial sequencing and analysis of
619  the human genome. Nature 409:860–921.

620 Jakšić, A. M., and C. Schlötterer. 2016. The interplay of temperature and genotype on patterns of
621  alternative splicing in *Drosophila melanogaster*. Genetics 204:315–325.

622 Kent, W. J., C. W. Sugnet, T. S. Furey, K. M. Roskin, T. H. Pringle, A. M. Zahler, and  a. D.
623  Haussler. 2002. The Human Genome Browser at UCSC. Genome Res. 12:996–1006.

Khil, P. P., N. A. Smirnova, P. J. Romanienko, and R. D. Camerini-Otero. 2004. The mouse X
chromosome is enriched for sex-biased genes not subject to selection by meiotic sex
chromosome inactivation. Nat. Genet. 36:642–646.

Kidwell, J. F., M. T. Clegg, F. M. Stewart, and T. Prout. 1977. Regions of stable equilibria for
models of differential selection in the two sexes under random mating. Genetics 85:171–
183.

Lack, J. B., C. M. Cardeno, M. W. Crepeau, W. Taylor, R. B. Corbett-Detig, K. A. Stevens, C.
H. Langley, and J. E. Pool. 2015. The *Drosophila* genome nexus: A population genomic
resource of 623 *Drosophila melanogaster* genomes, including 197 from a single ancestral
range population. Genetics 199:1229–1241.

Lack, J. B., J. D. Lange, A. D. Tang, R. B. Corbett-Detig, and J. E. Pool. 2016. A thousand fly
genomes: An expanded *Drosophila* genome nexus. Mol. Biol. Evol. 33:3308–3313.

Lande, R. 1980. Sexual Dimorphism, Sexual Selection, and Adaptation in Polygenic Characters.
Evolution. 34:292–305.

Long, Y., G. Song, J. Yan, X. He, Q. Li, and Z. Cui. 2013. Transcriptomic characterization of
cold acclimation in larval zebrafish. BMC Genomics 14:612.

Love, M. I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome Biol. 15:1–21.

MacKay, T. F. C., S. Richards, E. A. Stone, A. Barbadilla, J. F. Ayroles, D. Zhu, S. Casillas, Y.
Han, M. M. Magwire, J. M. Cridland, M. F. Richardson, R. R. H. Anholt, M. Barrón, C.
Bess, K. P. Blankenburg, M. A. Carbone, D. Castellano, L. Chaboub, L. Duncan, Z. Harris,
M. Javaid, J. C. Jayaseelan, S. N. Jhangiani, K. W. Jordan, F. Lara, F. Lawrence, S. L. Lee,
P. Librado, R. S. Linheiro, R. F. Lyman, A. J. MacKey, M. Munidasa, D. M. Muzny, L.
Nazareth, I. Newsham, L. Perales, L. L. Pu, C. Qu, M. Ràmia, J. G. Reid, S. M. Rollmann,
J. Rozas, N. Saada, L. Turlapati, K. C. Worley, Y. Q. Wu, A. Yamamoto, Y. Zhu, C. M.
Bergman, K. R. Thornton, D. Mittelman, and R. A. Gibbs. 2012. The *Drosophila*
*melanogaster* Genetic Reference Panel. Nature 482:173–178.

Mank, J. E. 2017. Population genetics of sexual conflict in the genomic era. Nat. Rev. Genet.

18:721–730.

McIntyre, L. M., L. M. Bono, A. Genissel, R. Westerman, D. Junk, M. Telonis-Scott, L.
Harshman, M. L. Wayne, A. Kopp, and S. V. Nuzhdin. 2006. Sex-specific expression of
alternative transcripts in *Drosophila*. Genome Biol. 7:R79.

Meisel, R. P., J. H. Malone, and A. G. Clark. 2012. Disentangling the relationship between sex-
biased gene expression and X-linkage. Genome Res. 22:1255–1265.

Mironov, A. A., J. W. Fickett, and M. S. Gelfand. 1999. Frequent alternative splicing of human
genes. Genome Res. 9:1288–1293.

Modrek, B., and C. Lee. 2002. A genomic view of alternative splicing. Nat. Genet. 30:13–19.

Modrek, B., A. Resch, C. Grasso, and C. Lee. 2001. Genome-wide detection of alternative
splicing in expressed sequences of human genes. Nucleic Acids Res. 29:2850–2859.

Nei, M., and W. H. Li. 1979. Mathematical model for studying genetic variation in terms of
restriction endonucleases. Proc. Natl. Acad. Sci. U. S. A. 76:5269–5273.

Osada, N., R. Miyagi, and A. Takahashi. 2017. Cis- and trans- regulatory effects on gene
expression. Genetics 206:2139–2148.

Paape, T., T. Bataillon, P. Zhou, T. J. Y. Kono, R. Briskine, N. D. Young, and P. Tiffin. 2013.
Selection, genome-wide fitness effects and evolutionary rates in the model legume
*Medicago truncatula*. Mol. Ecol. 22:3525–3538.

Page, A. J., B. Taylor, A. J. Delaney, J. Soares, T. Seemann, J. A. Keane, and S. R. Harris. 2016.
SNP-sites: rapid efficient extraction of SNPs from multi-FASTA alignments. Microb.
genomics 2:e000056.

Pál, C., B. Papp, and L. D. Hurst. 2001. Highly Expressed Genes in Yeast Evolve Slowly.
2Genetics 158:227–231.

Parisi, M., R. Nuttall, P. Edwards, J. Minor, D. Naiman, J. Lü, M. Doctolero, M. Vainer, C.
Chan, J. Malley, S. Eastman, and B. Oliver. 2004. A survey of ovary-, testis-, and soma-
biased gene expression in *Drosophila melanogaster* adults. Genome Biol. 5:R40.

Parisi, M., R. Nuttall, D. Naiman, G. Bouffard, J. Malley, J. Andrews, S. Eastman, and B. Oliver.

2003. Paucity of genes on the *Drosophila* X chromosome showing male-biased expression.
Science. 299:697–700.

Parsch, J., and H. Ellegren. 2013. The evolutionary causes and consequences of sex-biased gene
expression. Nat. Rev. Genet. 14:83–87.

Pröschel, M., Z. Zhang, and J. Parsch. 2006. Widespread adaptive evolution of *Drosophila* genes
with sex-biased expression. Genetics 174:893–900.

Quinlan, A. R., and I. M. Hall. 2010. BEDTools: A flexible suite of utilities for comparing
genomic features. Bioinformatics 26:841–842.

Ramani, A. K., J. A. Calarco, Q. Pan, S. Mavandadi, Y. Wang, A. C. Nelson, L. J. Lee, Q.
Morris, B. J. Blencowe, M. Zhen, and A. G. Fraser. 2011. Genome-wide analysis of
alternative splicing in *Caenorhabditis elegans*. Genome Res. 21:342–348.

Ranz, J. M., C. I. Castillo-Davis, C. D. Meiklejohn, and D. L. Hartl. 2003. Sex-dependent gene
expression and evolution of the *Drosophila* transcriptome. Science. 300:1742–1745.

Reinke, V., I. S. Gil, S. Ward, and K. Kazmer. 2004. Genome-wide germline-enriched and sex-
biased expression profiles in *Caenorhabditis elegans*. Development 131:311–323.

Revil, T., D. Gaffney, C. Dias, J. Majewski, and L. A. Jerome-Majewska. 2010. Alternative
splicing is frequent during early embryonic development in mouse. BMC Genomics 11:399.

Rice, W. R. 1984. Sex Chromosomes and the Evolution of Sexual Dimorphism. Evolution.
38:735–742.

Rice, W. R. 1987. The Accumulation of Sexually Antagonistic Genes as a Selective Agent
Promoting the Evolution of Reduced Recombination Between Primitive Sex Chromosomes.
Evolution. 41:911–914.

Rogers, T. F., D. H. Palmer, and A. E. Wright. 2020. Sex-Specific Selection Drives the
Evolution of Alternative Splicing in Birds. Mol. Biol. Evol. 38:519–530.

Rowe, L., S. F. Chenoweth, and A. F. Agrawal. 2018. The Genomics of Sexual Conflict. Am.
Nat. 192:274–286.

Sayadi, A., A. Martinez Barrio, E. Immonen, J. Dainat, D. Berger, C. Tellgren-Roth, B. Nystedt,

and G. Arnqvist. 2019. The genomic footprint of sexual conflict. Nat. Ecol. Evol. 3:1725–
1730.

Sorek, R., R. Shamir, and G. Ast. 2004. How prevalent is functional alternative splicing in the
human genome? Trends Genet. 20:68–71.

Stewart, A. D., A. Pischedda, and W. R. Rice. 2010. Resolving intralocus sexual conflict:
Genetic mechanisms and time frame. J. Hered. 101:94–99.

Sturgill, D., Y. Zhang, M. Parisi, and B. Oliver. 2007. Demasculinization of X chromosomes in
the *Drosophila* genus. Nature 450:238–241.

Subramanian, S., and S. Kumar. 2004. Gene expression intensity shapes evolutionary rates of the
proteins encoded by the vertebrate genome. Genetics 168:373–381.

Tajima, F. 1989. Statistical Method for Testing the Neutral Mutation Hypothesis by DNA
Polymorphism. Genetics 123:585–595.

Tan, M. H., K. F. Au, A. L. Yablonovitch, A. E. Wills, J. Chuang, J. C. Baker, W. H. Wong, and
J. B. Li. 2013. RNA sequencing reveals a diverse and dynamic repertoire of the *Xenopus*
*tropicalis* transcriptome over development. Genome Res. 23:201–216.

Telonis-Scott, M., A. Kopp, M. L. Wayne, S. V. Nuzhdin, and L. M. McIntyre. 2009. Sex-
specific splicing in *Drosophila*: Widespread occurrence, tissue specificity and evolutionary
conservation. Genetics 181:421–434.

Trabzuni, D., A. Ramasamy, S. Imran, R. Walker, C. Smith, M. E. Weale, J. Hardy, and M.
Ryten. 2013. Widespread sex differences in gene expression and splicing in the adult human
brain. Nat. Commun. 4: 2771.

Williamson, R. J., E. B. Josephs, A. E. Platts, K. M. Hazzouri, A. Haudry, M. Blanchette, and S.
I. Wright. 2014. Evidence for Widespread Positive and Negative Selection in Coding and
Conserved Noncoding Regions of Capsella grandiflora. PLoS Genet. 10:e1004622.

Wright, A. E., M. Fumagalli, C. R. Cooney, N. I. Bloch, F. G. Vieira, S. D. Buechel, N. Kolm,
and J. E. Mank. 2018. Male-biased gene expression resolves sexual conflict through the
evolution of sex-specific genetic architecture. Evol. Lett. 2:52–61.

Wright, A. E., T. F. Rogers, M. Fumagalli, C. R. Cooney, and J. E. Mank. 2019. Phenotypic
sexual dimorphism is associated with genomic signatures of resolved sexual conflict. Mol.
Ecol. 28:2860–2871.

Yang, X., E. E. Schadt, S. Wang, H. Wang, A. P. Arnold, L. Ingram-Drake, T. A. Drake, and A.
J. Lusis. 2006. Tissue-specific expression and regulation of sexually dimorphic genes in
mice. Genome Res. 16:995–1004.

Zemp, N., R. Tavares, A. Muyle, D. Charlesworth, G. A. B. Marais, and A. Widmer. 2016.
Evolution of sex-biased gene expression in a dioecious plant. Nat. Plants 2:16168.

**Supplemental Figures and Tables**

**Figure S1:** Patterns of overall gene expression and sexual dimorphism in isoform usage (SDIU) for genes expressed in the body and head tissue. **(A)** The mean of the log sex-averaged gene expression of genes exhibiting sexually dimorphic isoform usage (SDIU; green) or no SDIU (purple) for genes expressed in the body and head. Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates. **(B)** Scatter plot and density distributions of log sex-averaged gene expression of genes exhibiting SDIU only in bodies (light blue), only in heads (red), in both heads and bodies (dark blue) or in neither heads nor bodies (beige) when expressed in bodies (x-axis) or heads (y-axis). Density plots above and to the right of the figure correspond to the log sex-averaged gene expression in bodies and heads respectively.

**Figure S2:** Scatter plot and density distributions of the log$_2$ fold change (log$_2$FC) in male to female gene expression of genes exhibiting sexual dimorphism in isoform usage (SDIU) only in bodies (light blue), only in heads (red), in both heads and bodies (dark blue) or in neither heads nor bodies (beige) when expressed in bodies (x-axis) or heads (y-axis). Density plots above and to the right of the figure correspond to the log$_2$FC in gene expression in bodies and heads respectively.

**Figure S3:** The proportion of genes exhibiting SDIU to genes not exhibiting SDIU in bins of log2FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue) expression when expressed in **(C)** bodies or **(D)** heads. Faded points indicate bins with fewer than 5 genes. Error bars in panels B-D are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

**Figure S4:** Proportion of genes exhibiting sexually dimorphic isoform usage (SDIU) and sex-biased gene expression (SBGE) that are X-linked. **(A-B)** The proportion of genes found on the X chromosome in bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue) expression when expressed in **(A)** bodies or **(B)** heads. **(C-D)** The proportion of genes exhibiting SDIU (green) and not exhibiting SDIU (purple) found on the X chromosome in bins of $\log_2$FC in gene expression (see Methods) ranging from female-biased (left most pair of bins), unbiased (middle bin) and male-biased (right most pair of bins) expression when expressed in **(C)** bodies or **(D)** heads. Faded points indicate bins with fewer than 5 genes. Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

**Figure S5: (A, B)** Mean Tajima's *D* for all genes expressed in **(A)** body or **(B)** head tissue averaged across all genes within bins of log$_2$FC in gene expression (see Methods) ranging from female-biased (red), unbiased (black) and male-biased (blue). Faded points indicate bins with fewer than 5 genes. Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling estimates.

**Figure S6: (A, B)** Median $\pi_N/\pi_S$ for all genes expressed in **(A)** body or **(B)** head tissue averaged
across all genes within bins of log$_2$FC in gene expression (see Methods) ranging from female-
biased (red), unbiased (black) and male-biased (blue). Faded points indicate bins with fewer than
5 genes. Error bars are 95% confidence intervals estimated from 10,000 bootstraps resampling
estimates.

**Table S1:** Gene Ontology (GO) categories enriched for genes exhibiting sexually dimorphic exon usage when expressed in body
tissue of 18 male and 18 female genotypes from the *Drosophila* Genetic Resource Population (DGRP). Statistical significance was
assessed at an FDR *Q*-value < 0.01.

| GO Term | Description | Enrichment | FDR *Q*-value |
|---|---|---|---|
| | **Component** | | |
| GO:0005737 | Cytoplasm | 1.18 | 1.29E-11 |
| GO:0005829 | Cytosol | 1.25 | 9.20E-08 |
| GO:0044464 | Cell Part | 1.04 | 4.77E-07 |
| GO:0005886 | Plasma Membrane | 1.22 | 2.64E-05 |
| GO:0044430 | Cytoskeletal Part | 1.30 | 2.67E-05 |
| GO:0043226 | Organelle | 1.07 | 1.15E-04 |
| GO:0005819 | Spindle | 1.64 | 1.03E-04 |
| GO:0016020 | Membrane | 1.14 | 1.30E-04 |
| GO:0043228 | Non-Membrane-Bounded Organelle | 1.18 | 1.82E-04 |
| GO:0043232 | Intracellular Non-Membrane-Bounded Organelle | 1.18 | 1.64E-04 |
| GO:0044424 | Intracellular Part | 1.04 | 1.86E-04 |
| GO:0043229 | Intracellular Organelle | 1.06 | 2.42E-04 |
| GO:0005654 | Nucleoplasm | 1.36 | 3.41E-04 |
| GO:0005634 | Nucleus | 1.10 | 3.27E-04 |
| GO:0097458 | Neuron Part | 1.26 | 5.68E-04 |
| GO:0005575 | Cellular Component | 1.02 | 7.72E-04 |
| GO:0043227 | Membrane-Bounded Organelle | 1.06 | 1.35E-03 |
| GO:0005938 | Cell Cortex | 1.35 | 1.43E-03 |
| GO:0048471 | Perinuclear Region of Cytoplasm | 1.52 | 1.40E-03 |
| GO:0099568 | Cytoplasmic Region | 1.32 | 4.53E-03 |
| GO:0044456 | Synapse Part | 1.32 | 4.74E-03 |

| GO:0044444 | Cytoplasmic Part | 1.06 | 5.03E-03 |
|---|---|---|---|
| GO:0043231 | Intracellular Membrane-Bounded Organelle | 1.06 | 1.03E-02 |
| GO:0035770 | Ribonucleoprotein Granule | 1.41 | 1.27E-02 |
| GO:0070069 | Cytochrome Complex | 1.74 | 1.25E-02 |
| GO:0036464 | Cytoplasmic Ribonucleoprotein Granule | 1.41 | 1.69E-02 |
| GO:0120038 | Plasma Membrane Bounded Cell Projection Part | 1.26 | 1.81E-02 |
| GO:0044463 | Cell Projection Part | 1.26 | 1.75E-02 |
| GO:0044459 | Plasma Membrane Part | 1.17 | 1.74E-02 |

**Function**

| GO:0005515 | Protein Binding | 1.16 | 1.07E-11 |
|---|---|---|---|
| GO:0016301 | Kinase Activity | 1.31 | 5.87E-04 |
| GO:0005488 | Binding | 1.05 | 1.52E-03 |
| GO:0015318 | Inorganic Molecular Entity Transmembrane Transporter Activity | 1.30 | 3.59E-03 |
| GO:0016773 | Phosphotransferase Activity, Alcohol Group as Acceptor | 1.29 | 3.94E-03 |
| GO:0005516 | Calmodulin Binding | 1.85 | 3.42E-03 |
| GO:0019787 | Ubiquitin-Like Protein Transferase Activity | 1.34 | 4.15E-03 |
| GO:0030554 | Adenyl Nucleotide Binding | 1.18 | 7.76E-03 |
| GO:0032559 | Adenyl Ribonucleotide Binding | 1.18 | 7.69E-03 |
| GO:0016772 | Transferase Activity, Transferring Phosphorus-Containing Groups | 1.22 | 9.54E-03 |

**Process**

| GO:0048519 | Negative Regulation of Biological Process | 1.24 | 9.11E-14 |
|---|---|---|---|
| GO:0048523 | Negative Regulation of Cellular Process | 1.25 | 2.36E-12 |
| GO:0065007 | Biological Regulation | 1.11 | 4.34E-12 |
| GO:0032502 | Developmental Process | 1.18 | 4.69E-12 |
| GO:0050789 | Regulation of Biological Process | 1.11 | 1.23E-10 |
| GO:0048583 | Regulation of Response to Stimulus | 1.24 | 2.27E-09 |
| GO:0050794 | Regulation of Cellular Process | 1.11 | 4.27E-09 |

| GO:0010646 | Regulation of Cell Communication | 1.25 | 7.35E-09 |
|---|---|---|---|
| GO:0023051 | Regulation of Signaling | 1.25 | 6.53E-09 |
| GO:0009966 | Regulation of Signal Transduction | 1.26 | 1.10E-08 |
| GO:0051128 | Regulation of Cellular Component Organization | 1.24 | 1.89E-08 |
| GO:0022414 | Reproductive Process | 1.21 | 1.41E-07 |
| GO:0032879 | Regulation of Localization | 1.30 | 2.68E-07 |
| GO:0048518 | Positive Regulation of Biological Process | 1.15 | 2.94E-07 |
| GO:0048856 | Anatomical Structure Development | 1.18 | 2.79E-07 |
| GO:0048585 | Negative Regulation of Response to Stimulus | 1.33 | 3.85E-07 |
| GO:0050793 | Regulation of Developmental Process | 1.24 | 5.17E-07 |
| GO:0003006 | Developmental Process Involved in Reproduction | 1.25 | 2.03E-06 |
| GO:0022412 | Cellular Process Involved in Reproduction in Multicellular Organism | 1.23 | 1.92E-06 |
| GO:0032501 | Multicellular Organismal Process | 1.17 | 2.28E-06 |
| GO:0048869 | Cellular Developmental Process | 1.20 | 2.21E-06 |
| GO:0009987 | Cellular Process | 1.05 | 2.78E-06 |
| GO:0048522 | Positive Regulation of Cellular Process | 1.15 | 2.72E-06 |
| GO:0009653 | Anatomical Structure Morphogenesis | 1.22 | 2.75E-06 |
| GO:0009968 | Negative Regulation of Signal Transduction | 1.33 | 2.88E-06 |
| GO:0023057 | Negative Regulation of Signaling | 1.31 | 5.66E-06 |
| GO:0010648 | Negative Regulation of Cell Communication | 1.31 | 7.30E-06 |
| GO:0051246 | Regulation of Protein Metabolic Process | 1.24 | 7.43E-06 |
| GO:0010605 | Negative Regulation of Macromolecule Metabolic Process | 1.23 | 7.30E-06 |
| GO:0051239 | Regulation of Multicellular Organismal Process | 1.22 | 8.45E-06 |
| GO:0016043 | Cellular Component Organization | 1.11 | 8.98E-06 |
| GO:0060255 | Regulation of Macromolecule Metabolic Process | 1.13 | 9.89E-06 |
| GO:1902531 | Regulation of Intracellular Signal Transduction | 1.29 | 1.45E-05 |
| GO:0051493 | Regulation of Cytoskeleton Organization | 1.40 | 1.73E-05 |
| GO:0007610 | Behavior | 1.29 | 1.80E-05 |
| GO:0009892 | Negative Regulation of Metabolic Process | 1.21 | 3.00E-05 |

| GO:0051172 | Negative Regulation of Nitrogen Compound Metabolic Process | 1.24 | 3.50E-05 |
|---|---|---|---|
| GO:0032268 | Regulation of Cellular Protein Metabolic Process | 1.23 | 3.98E-05 |
| GO:0022603 | Regulation of Anatomical Structure Morphogenesis | 1.34 | 4.49E-05 |
| GO:0071840 | Cellular Component Organization or Biogenesis | 1.10 | 5.01E-05 |
| GO:2000026 | Regulation of Multicellular Organismal Development | 1.23 | 5.20E-05 |
| GO:0007010 | Cytoskeleton Organization | 1.27 | 5.32E-05 |
| GO:0065008 | Regulation of Biological Quality | 1.16 | 6.34E-05 |
| GO:0031324 | Negative Regulation of Cellular Metabolic Process | 1.23 | 6.82E-05 |
| GO:0051049 | Regulation of Transport | 1.32 | 8.06E-05 |
| GO:0051960 | Regulation of Nervous System Development | 1.28 | 1.00E-04 |
| GO:0045595 | Regulation of Cell Differentiation | 1.26 | 1.18E-04 |
| GO:0006996 | Organelle Organization | 1.16 | 1.17E-04 |
| GO:0051171 | Regulation of Nitrogen Compound Metabolic Process | 1.12 | 1.59E-04 |
| GO:0080090 | Regulation of Primary Metabolic Process | 1.12 | 1.58E-04 |
| GO:0019222 | Regulation of Metabolic Process | 1.11 | 1.64E-04 |
| GO:0006928 | Movement of Cell or Subcellular Component | 1.22 | 1.80E-04 |
| GO:0010647 | Positive Regulation of Cell Communication | 1.25 | 1.84E-04 |
| GO:0023056 | Positive Regulation of Signaling | 1.25 | 1.80E-04 |
| GO:0048609 | Multicellular Organismal Reproductive Process | 1.30 | 1.88E-04 |
| GO:0007017 | Microtubule-Based Process | 1.28 | 2.27E-04 |
| GO:0070647 | Protein Modification by Small Protein Conjugation or Removal | 1.30 | 2.46E-04 |
| GO:0033043 | Regulation of Organelle Organization | 1.23 | 3.19E-04 |
| GO:0009967 | Positive Regulation of Signal Transduction | 1.25 | 3.23E-04 |
| GO:0007276 | Gamete Generation | 1.32 | 3.48E-04 |
| GO:0008589 | Regulation of Smoothened Signaling Pathway | 1.49 | 3.82E-04 |
| GO:0051248 | Negative Regulation of Protein Metabolic Process | 1.35 | 4.38E-04 |
| GO:0048584 | Positive Regulation of Response to Stimulus | 1.22 | 4.49E-04 |
| GO:0032269 | Negative Regulation of Cellular Protein Metabolic Process | 1.35 | 5.82E-04 |
| GO:0051093 | Negative Regulation of Developmental Process | 1.34 | 7.20E-04 |

| GO:0070507 | Regulation of Microtubule Cytoskeleton Organization | 1.61 | 7.40E-04 |
|---|---|---|---|
| GO:0010604 | Positive Regulation of Macromolecule Metabolic Process | 1.17 | 7.39E-04 |
| GO:0048646 | Anatomical Structure Formation Involved in Morphogenesis | 1.27 | 7.30E-04 |
| GO:0001932 | Regulation of Protein Phosphorylation | 1.32 | 7.94E-04 |
| GO:0045596 | Negative Regulation of Cell Differentiation | 1.46 | 7.90E-04 |
| GO:0007281 | Germ Cell Development | 1.31 | 7.79E-04 |
| GO:0048149 | Behavioral Response to Ethanol | 1.64 | 8.20E-04 |
| GO:0048477 | Oogenesis | 1.36 | 8.14E-04 |
| GO:0007292 | Female Gamete Generation | 1.36 | 8.03E-04 |
| GO:0060284 | Regulation of Cell Development | 1.25 | 9.37E-04 |
| GO:0031323 | Regulation of Cellular Metabolic Process | 1.10 | 1.03E-03 |
| GO:0045880 | Positive Regulation of Smoothened Signaling Pathway | 1.61 | 1.03E-03 |
| GO:0030029 | Actin Filament-Based Process | 1.32 | 1.04E-03 |
| GO:0030036 | Actin Cytoskeleton Organization | 1.35 | 1.13E-03 |
| GO:0051130 | Positive Regulation of Cellular Component Organization | 1.25 | 1.15E-03 |
| GO:0051656 | Establishment of Organelle Localization | 1.37 | 1.18E-03 |
| GO:0080134 | Regulation of Response to Stress | 1.27 | 1.29E-03 |
| GO:0009893 | Positive Regulation of Metabolic Process | 1.15 | 1.36E-03 |
| GO:0051173 | Positive Regulation of Nitrogen Compound Metabolic Process | 1.17 | 1.35E-03 |
| GO:0032886 | Regulation of Microtubule-Based Process | 1.56 | 1.40E-03 |
| GO:0044087 | Regulation of Cellular Component Biogenesis | 1.23 | 1.44E-03 |
| GO:0016567 | Protein Ubiquitination | 1.32 | 1.81E-03 |
| GO:0016331 | Morphogenesis of Embryonic Epithelium | 1.41 | 1.82E-03 |
| GO:1902532 | Negative Regulation of Intracellular Signal Transduction | 1.37 | 1.83E-03 |
| GO:0017145 | Stem Cell Division | 1.49 | 1.87E-03 |
| GO:0048468 | Cell Development | 1.22 | 1.89E-03 |
| GO:0051241 | Negative Regulation of Multicellular Organismal Process | 1.33 | 2.06E-03 |
| GO:0051961 | Negative Regulation of Nervous System Development | 1.41 | 2.46E-03 |
| GO:0048285 | Organelle Fission | 1.51 | 2.44E-03 |

| GO:0042326 | Negative Regulation of Phosphorylation | 1.50 | 2.57E-03 |
|---|---|---|---|
| GO:0032446 | Protein Modification by Small Protein Conjugation | 1.30 | 2.91E-03 |
| GO:0034248 | Regulation of Cellular Amide Metabolic Process | 1.32 | 3.23E-03 |
| GO:0051640 | Organelle Localization | 1.29 | 3.27E-03 |
| GO:0007391 | Dorsal Closure | 1.41 | 3.46E-03 |
| GO:1903829 | Positive Regulation of Cellular Protein Localization | 1.70 | 3.47E-03 |
| GO:0001933 | Negative Regulation of Protein Phosphorylation | 1.49 | 3.66E-03 |
| GO:0050767 | Regulation of Neurogenesis | 1.27 | 3.67E-03 |
| GO:0010563 | Negative Regulation of Phosphorus Metabolic Process | 1.45 | 3.78E-03 |
| GO:0045936 | Negative Regulation of Phosphate Metabolic Process | 1.45 | 3.74E-03 |
| GO:0010629 | Negative Regulation of Gene Expression | 1.20 | 3.76E-03 |
| GO:0022604 | Regulation of Cell Morphogenesis | 1.36 | 3.84E-03 |
| GO:0022402 | Cell Cycle Process | 1.19 | 4.29E-03 |
| GO:0051179 | Localization | 1.09 | 4.31E-03 |
| GO:0042325 | Regulation of Phosphorylation | 1.26 | 4.61E-03 |
| GO:0016334 | Establishment or Maintenance of Polarity of Follicular Epithelium | 1.65 | 4.81E-03 |
| GO:0009890 | Negative Regulation of Biosynthetic Process | 1.22 | 4.78E-03 |
| GO:0031327 | Negative Regulation of Cellular Biosynthetic Process | 1.22 | 4.74E-03 |
| GO:0010608 | Posttranscriptional Regulation of Gene Expression | 1.30 | 4.90E-03 |
| GO:0031325 | Positive Regulation of Cellular Metabolic Process | 1.15 | 5.46E-03 |
| GO:0019220 | Regulation of Phosphate Metabolic Process | 1.25 | 5.42E-03 |
| GO:0051174 | Regulation of Phosphorus Metabolic Process | 1.25 | 5.37E-03 |
| GO:0008150 | Biological Process | 1.01 | 5.44E-03 |
| GO:0031399 | Regulation of Protein Modification Process | 1.23 | 5.59E-03 |
| GO:0009886 | Post-Embryonic Animal Morphogenesis | 1.22 | 5.57E-03 |
| GO:0006796 | Phosphate-Containing Compound Metabolic Process | 1.15 | 5.56E-03 |
| GO:0031400 | Negative Regulation of Protein Modification Process | 1.40 | 6.04E-03 |
| GO:0051129 | Negative Regulation of Cellular Component Organization | 1.31 | 6.12E-03 |
| GO:0097435 | Supramolecular Fiber Organization | 1.32 | 6.09E-03 |

| GO:0050768 | Negative Regulation of Neurogenesis | 1.49 | 6.56E-03 |
|---|---|---|---|
| GO:0010721 | Negative Regulation of Cell Development | 1.46 | 6.80E-03 |
| GO:0048731 | System Development | 1.27 | 7.63E-03 |
| GO:0030534 | Adult Behavior | 1.36 | 7.70E-03 |
| GO:0007051 | Spindle Organization | 1.39 | 7.70E-03 |
| GO:0010558 | Negative Regulation of Macromolecule Biosynthetic Process | 1.22 | 7.94E-03 |
| GO:2000113 | Negative Regulation of Cellular Macromolecule Biosynthetic Process | 1.22 | 7.88E-03 |
| GO:0007411 | Axon Guidance | 1.29 | 7.85E-03 |
| GO:0051301 | Cell Division | 1.31 | 8.36E-03 |
| GO:0098655 | Cation Transmembrane Transport | 1.34 | 8.60E-03 |
| GO:0060341 | Regulation of Cellular Localization | 1.32 | 9.07E-03 |
| GO:0006811 | Ion Transport | 1.22 | 9.46E-03 |
| GO:0007389 | Pattern Specification Process | 1.20 | 9.70E-03 |

805 **Table S2:** Gene Ontology (GO) categories enriched for genes exhibiting sexually dimorphic exon usage when expressed in head
806 tissue of 18 male and 18 female genotypes from the *Drosophila* Genetic Resource Population (DGRP). Statistical significance was
807 assessed at an FDR *Q*-value < 0.01.

| GO Term | Description | Enrichment | FDR *Q*-value |
|---|---|---|---|
| | **Component** | | |
| GO:0005829 | Cytosol | 1.96 | 8.51E-06 |
| GO:0044444 | Cytoplasmic Part | 1.28 | 2.27E-03 |
| GO:0097458 | Neuron Part | 1.87 | 2.06E-03 |
| GO:0005886 | Plasma Membrane | 1.69 | 2.08E-03 |
| GO:0044430 | Cytoskeletal Part | 2.15 | 3.32E-03 |
| | **Function** | | |
| GO:0005515 | Protein Binding | 1.36 | 6.14E-03 |
| | **Process** | | |
| GO:0003008 | System Process | 2.12 | 2.24E-03 |
| GO:0007010 | Cytoskeleton Organization | 2.20 | 1.25E-03 |
| GO:0007610 | Behavior | 2.02 | 8.34E-04 |
| GO:0032501 | Multicellular Organismal Process | 1.52 | 1.19E-03 |
| GO:0030036 | Actin Cytoskeleton Organization | 2.70 | 2.87E-03 |
| GO:0032879 | Regulation of Localization | 1.85 | 5.35E-03 |
| GO:0044089 | Positive Regulation of Cellular Component Biogenesis | 2.69 | 5.54E-03 |
| GO:0051049 | Regulation of Transport | 2.15 | 4.86E-03 |

810 **Table S3:** Parameter estimates from a logistic regression model fit to test the fixed effects of overall gene expression, sex-bias in gene
811 expression (i.e., log$_2$ fold change in male to female gene expression) and sexual dimorphism in isoform usage (SDIU) on $\pi_N/\pi_S$.
812 $\pi_N/\pi_S$ was modelled as a binary trait wherein genes are binned as either $\pi_N/\pi_S = 0$ or $\pi_N/\pi_S > 0$ in body and head tissue. Bold terms
813 indicate statistically significant effects at $\alpha = 0.05$.

| Tissue | Effect | Estimate | Standard Error | *P* |
|---|---|---|---|---|
| Body Tissue | **Intercept** | **2.700** | **0.112** | **< 0.001** |
| | **Sex-averaged Gene Expression[1]** | **-4.29 x 10$^{-5}$** | **5.55 x 10$^{-6}$** | **< 0.001** |
| | **Absolute value of SBGE[2]** | **0.137** | **0.062** | **0.028** |
| | Sexual Dimorphism in Isoform Usage (SDIU)[3] | 0.095 | 0.113 | 0.401 |
| Head Tissue | **Intercept** | **2.907** | **0.068** | **< 0.001** |
| | **Sex-averaged Gene Expression[1]** | **-4.68 x 10$^{-6}$** | **2.39 x 10$^{-6}$** | **0.050** |
| | Absolute value of SBGE[2] | -0.157 | 0.169 | 0.353 |
| | Sexual Dimorphism in Isoform Usage (SDIU)[3] | -0.135 | 0.195 | 0.489 |

815 [1]Gene expression averaged across both sexes.
816 [2]Measured as |Log$_2$FC male:female gene expression|.
817 [3]Indicator of whether or not a gene exhibited statistically significant evidence of SDIU.