Infant brain regional cerebral blood flow dynamics supporting emergence of vital

functional networks

Qinlin Yu[1, 2], Minhui Ouyang[1, 2], John A. Detre[2,4], Huiying Kang[1,3], Di Hu[1,3], Bo Hong[5], Fang

Fang[6], Yun Peng[3], Hao Huang[1, 2,*]

[1]Department of Radiology, Children's Hospital of Philadelphia, Philadelphia, PA, United States;

[2]Department of Radiology, [4]Department of Neurology, Perelman School of Medicine, University

of Pennsylvania, Philadelphia, PA, United States;

[3]Department of Radiology, Beijing Children's Hospital, Capital Medical University, Beijing,

China;

[5]Department of Biomedical Engineering, Tsinghua University, Beijing, China

[6]School of Psychological and Cognitive Sciences, Peking University, Beijing, China

* Corresponding author

Hao Huang, Ph.D.

34th street and Civic Center blvd,
Philadelphia, PA 19104;

Tel: 267-425-1599

Email: huangh6@chop.edu

ABSTRACT

Human infancy is characterized by most rapid cerebral blood flow (rCBF) increases across lifespan and emergence of a fundamental brain system default-mode network (DMN). However, how infant rCBF change spatiotemporally across the brain and how the rCBF dynamics support emergence of vital functional networks such as DMN remains unknown. Here, by acquiring cutting-edge multi-modal MRI including pseudo-continuous arterial-spin-labeled perfusion MRI and resting-state functional MRI of infants aged 0 to 24 months, we elucidated unprecedented 4D spatiotemporal infant rCBF framework and region-specific physiology-function coupling across infancy. We found faster rCBF increases in the DMN than other regions. We also found strongly coupled increases of rCBF and network strength specifically in the DMN, suggesting faster local blood flow increase to meet extra neuronal metabolic demands in the DMN maturation. These results offer insights into physiological mechanism of brain functional network emergence and have important implications in altered network maturation in brain disorders.

KEYWORDS: infant brain development; default mode network; perfusion; functional connectivity; metabolism; emergence

INTRODUCTION

The adult human brain receives 15-20% of cardiac output despite only representing 2% of body mass (Bouma and Muizelaar, 1990; Satterthwaite et al., 2014). Vast energy demand from the human brain starts from infancy which is characterized by fastest energy expenditure increase across lifespan (Pontzer et al., 2021). Infancy is also the most dynamic phase of brain development across entire lifespan with fastest functional and structural brain development. For example, during infancy the brain size increases dramatically in parallel with rapid elaboration of new synapses, reaching 80-90% of lifetime maximum by age of year 2 (Knickmeyer et al., 2008; Pfefferbaum et al., 1994; Ouyang et al., 2019a). Structural and functional changes of infant brain are underlaid by rapid and precisely regulated (Huang et al., 2013; Silbereis et al., 2016) spatiotemporal cellular and molecular processes including neurogenesis and neuronal migration (Rakic, 1995; Sidman and Rakic, 1973), synaptic formation (Huttenlocher and Dabholkar, 1997), dendritic arborization (Bystron et al., 2008; Ouyang et al., 2019b), axonal growth (Haynes et al., 2005; Innocenti and Price, 2005) and myelination (Miller et al., 2012; Yakovlev, 1967). These developmental processes demand rapidly increasing feed of energy to fuel the brain development. However, there have been no known whole-brain mappings of heterogeneous infant brain regional cerebral blood flow (rCBF) dynamics across landmark infant ages thus far, impeding fundamental view of energy expenditure across functional systems of early developing brain. As a result of differential neuronal growth across cortex, functional networks in the human brain develop differentially following the order from primary sensorimotor to higher-order cognitive systems (Cao et al., 2017a; Huang and Vasung, 2014; Tau and Peterson, 2010; Yu et al., 2016). The default-mode network (DMN) (Raichle et al., 2001) is widely recognized as a fundamental neurobiological system associated with cognitive processes that are directed toward the self and has important implication in typical and atypical brain development (Buckner et al., 2008). Unlike primary sensorimotor (SM) and visual (Vis) networks emerging relatively earlier around and before birth (Cao et al., 2017b; Doria et al., 2010; Smyser et al., 2010), emergence of the vital

resting-state DMN is not well established until late infancy (Gao et al., 2009). Till now, it has been

unclear how emergence of vital functional networks such as DMN is coupled with rCBF increase

during infancy.

Regional brain metabolism, including glucose utilization and oxygen consumption, is

closely coupled to regional CBF (rCBF) that delivers the glucose and oxygen needed to sustain

metabolic needs (Raichle et al., 2001; Vaishnavi et al., 2010). Infant rCBF has been

conventionally measured with positron emission tomography (PET) (Chugani and Phelps, 1986;

Chugani et al., 1987) and single-photon emission computerized tomography (SPECT) (Chiron et

al., 1992) which are not applicable to infants due to the associated exposure to radioactive tracers.

By labeling the blood in internal carotid and vertebral arteries in neck and measuring downstream

labeled arterial blood in brain, arterial-spin-labeled (ASL) (Alsop et al., 2015; Detre and Alsop,

1999) perfusion MRI provides a method for noninvasive quantifying rCBF without requiring

radioactive tracers or exogenous contrast agents. Accordingly, ASL is especially suitable for rCBF

measurements of infants (Ouyang et al., 2017; Wang et al., 2008) and children (Jain et al., 2012;

Satterthwaite et al., 2014). Phase-contrast (PC) MRI, utilizing the phase shift proportional to

velocity of the blood spins, has also been used to measure global CBF of the entire brain (Liu et

al., 2019). Through integration of pseudo-continuous ASL (pCASL) and PC MRI, rCBF measured

from pCASL can be calibrated by global CBF from PC MRI for more accurate infant brain rCBF

measurement (Aslan et al., 2010; Ouyang et al., 2017). With rCBF closely related to regional

cerebral metabolic rate of oxygen ($CMRO_2$) and glucose (CMRGlu) at the resting state in human

brain (Fox and Raichle, 1986; Gur et al., 2009; Paulson et al., 2010; Vaishnavi et al., 2010), rCBF

could be used as a surrogate measure of local cerebral metabolic level for resting infant brains.

Early developing human brain functional networks can be reproducibly measured with

resting-state fMRI (rs-fMRI). For example, a large scale of functional architecture at birth (Cao et

al., 2017b; Doria et al., 2010; Fransson et al., 2008) has been revealed with rs-fMRI. Functional

networks consist of densely linked hub regions to support efficient neuronal signaling and communication. These hub regions can be delineated with data-driven independent component analysis (ICA) of rs-fMRI data and serve as functional regions-of-interests (ROIs) for testing physiology-function relationship. Meeting metabolic demands in these hub ROIs is critical for functional network maturation. In fact, spatial correlation of rCBF to the functional connectivity (FC) in these functional ROIs was found in adult brains (Liang et al., 2013). Altered DMN plays a vital role in neurodevelopmental disorders such as autism (Doyle-Thomas et al., 2015; Lynch et al., 2013; Padmanabhan et al., 2017; Washington et al., 2014). Thus, understanding physiological underpinning of the DMN maturation offers invaluable insights into the mechanism of typical and atypical brain development.

We hypothesized heterogeneous rCBF maps at landmark infant ages and faster rCBF increase in brain regions of higher cognitive functions (namely DMN regions) during infancy than those of primary sensorimotor functions where functional networks emerge before or around birth (Cao et al., 2017a; Cao et al., 2017b; Doria et al., 2010; Fransson et al., 2007; Smyser et al., 2010; Peng et al., 2020). Furthermore, with rCBF an indicator of local metabolic level of glucose and oxygen consumption, we hypothesized strongly coupled rCBF and FC increases specifically in the DMN regions during infancy to meet extra metabolic demand of DMN maturation. In this study, we acquired multi-modal MRI including both pCASL perfusion MRI and rs-fMRI of 48 infants aged 0 to 24 months to quantify rCBF and FC, respectively. RCBF at the voxel level and in functional network ROIs were measured to test the hypothesis of spatiotemporally differential rCBF increases during infancy. Maturation of FC in the DMN was delineated. Correlation of FC increase and rCBF increase in the DMN ROIs was tested and further confirmed with data-driven permutation analysis, the latter of which was to examine if the coupling of rCBF and FC takes place only in the DMN during infancy.

RESULTS

Faster rCBF increases in the DMN hub regions during infant brain development. The labeling plane and imaging slices of pCASL perfusion MRI of a representative infant brain, reconstructed internal carotid and vertebral arteries, and four PC MR images of the target arteries are shown in Figure 1a. The rCBF maps of infant brains were calculated based on pCASL perfusion MRI and calibrated by PC MRI. As an overview, axial rCBF maps of typically developing brains at milestone ages of 1, 6, 12, 18 and 24 months are demonstrated in Figure 1b. High quality of the rCBF maps can be appreciated by a clear contrast between white matter and gray matter. A general increase of blood flow across the brain gray matter from birth to 2 years of age is readily observed. Heterogeneous rCBF distribution at a given infant age can be appreciated from these maps. For example, higher rCBF values in primary visual cortex compared to other brain regions are clear in younger infant at around 1 month. Figure 1b also demonstrates differential rCBF increases across brain regions. RCBF increases are prominent in the posterior cingulate cortex (PCC, a vital hub of DMN network), indicated by green arrows. On the other hand, rCBF in the visual cortex is already higher (indicated by blue arrows) than other brain regions in early infancy and increases slowly across infant development. The adopted pCASL protocol is highly reproducible with intraclass correlation coefficient (ICC) 0.8854 calculated from pCASL scans of a randomly selected infant subject aged 17.6 month, shown in Figure 1- figure supplement 1. As shown in Figure 1- figure supplement 2, three functional network ROIs including DMN, visual (Vis)network, and sensorimotor (SM) network were generated from infant rs-fMRI data. With rCBF measured at these functional network ROIs, Figure 1- figure supplement 3 quantitatively exhibits spatial inhomogeneity of rCBF distribution regardless of age. These quantitative measurements are consistent to the observation of heterogeneous rCBF distribution in Figure 1b. Specifically, as shown in Figure 1 – figure supplement 3, significant heterogeneity of rCBF was found across regions ($F(6, 282) = 122.6$, $p < 10^{-10}$) with ANOVA test. With further paired t-test between regions,

the highest and lowest rCBF was found in the Vis (82.1 ml/100g/min) and SM (49.1 ml/100g/min)

regions, respectively (all $ts(47) > 4.17$, $p < 0.05$), while rCBF in the DMN (67.8 ml/100g/min)

regions was in the middle (all ts (47) > 2.87, $p < 0.05$). Within the DMN, rCBF in the PCC (75.4

ml/100g/min) and LTC (72.0 ml/100g/min) regions were significantly higher than rCBF in the

MPFC (60.7 ml/100g/min) (both $ts(47) > 8.22$, $p < 0.05$) and IPL regions (59.4 ml/100g/min) (both

$ts(47) > 7.87$, $p < 0.05$).

Figure 2a shows cortical maps of linearly fitted rCBF values of infant brains from 0 to 24

months. Consistent with nonuniform profile of the rCBF maps observed in Figure 1b, the three-

dimensionally reconstructed rCBF distribution maps in Figure 2a are also not uniform at each

milestone infant age. RCBF increases from 0 to 24 months across cortical regions are apparent,

as demonstrated by relatively high rCBF-age correlation r values across the cortical surface in

Figure 2b. Heterogeneity of rCBF increases across all brain voxels can be more clearly

appreciated in Figures 2a and 2b compared to Figure 1b. With DMN functional network regions

including PCC, medial prefrontal cortex (MPFC), inferior posterior lobule (IPL) and lateral

temporal cortex (LTC) as well as Vis and SM network regions delineated in Figure 1-figure

supplement 2b as ROIs, rCBF trajectories in Figure 2c demonstrate that rCBF in these ROIs all

increase significantly with age (Vis: $r = 0.53$, $p < 10^{-4}$; SM: $r = 0.52$, $p < 10^{-4}$; DMN: $r = 0.7$, $p < 10^{-7}$;

DMN_PCC: $r = 0.66$, $p < 10^{-6}$; DMN_MPFC: $r = 0.67$, $p < 10^{-6}$; DMN_IPL: $r = 0.66$, $p < 10^{-6}$;

DMN_LTC: $r = 0.72$, $p < 10^{-8}$). Using the trajectory of primary sensorimotor (SM) (black line and

circles) in Figure 2c as a reference, rCBF increase rates across functional network ROIs are also

heterogeneous (Figure 2c). Specifically, significantly higher (all $p < 0.05$, FDR corrected) rCBF

increase rate was found in total DMN ROIs (1.59 ml/100g/min/month) and individual DMN ROIs,

including DMN_PCC (1.57 ml/100g/min/month), DMN_MPFC (1.63 ml/100g/min/month),

DMN_IPL (1.42 ml/100g/min/month) and DMN_LTC (2.22 ml/100g/min/month), compared to in

the SM ROI (0.85 ml/100g/min/month). Although the rCBF growth rate in the Vis (1.27

ml/100g/min/month) ROIs is higher than that in the SM ROIs, this difference is not significant (p

= 0.13). Collectively, Figures 1 and 2 show that the CBF increases significantly and differentially

across brain regions during infancy, with rCBF in the DMN hub regions increasing faster than

rCBF in the SM and Vis regions (Figure 2). The 4D spatiotemporal whole-brain rCBF dynamics

during infant development are presented in Video 1.

Emergence of DMN during early brain development. Figure 3a shows emergence of the DMN

in typically developing brain from 0 to 24 months as measured using rs-fMRI with a PCC seed

region indicated by the black dash line. At around birth (0 months), the DMN is still immature with

weak FC between other DMN regions (including MPFC, IPL, and LTC) and PCC (Figure 3a).

During infant brain development from 0 to 24 months, Figure 3a shows that the functional

connectivity between MPFC, IPL or ITC and PCC gradually strengthens. Figure 3b shows the FC-

age correlation r value map. It can be appreciated from Figure 3b that across the cortical surface

relatively higher r values are only located at the DMN regions (except the seed PCC).

178   As shown in Figure 3- figure supplement 1, only the FC within the DMN ($r = 0.31$, $p < 0.05$),

but not the Vis ($r = 0.048$, $p = 0.745$) or SM ($r = 0.087$, $p = 0.559$), was found to increase

significantly with age, indicating significant functional development in the DMN, but not in the Vis

or SM network.

Coupling between rCBF and FC within DMN during infant brain development. To test the

hypothesis that rCBF increases in the DMN regions underlie emergence of this vital functional

network, correlation between rCBF and FC was conducted across randomly selected voxels

within the DMN of all infants aged 0-12 months (Figure 4a) and all infants aged 12-24 months

(Figure 4b). Significant correlations ($p<0.001$) were found in both age groups. We further tested

whether functional emergence of the DMN represented by increases of FC within the DMN

(namely DMN FC) was correlated to rCBF increases specifically in the DMN regions, but not in

primary sensorimotor (Vis or SM) regions. Figure 5a shows correlations between the DMN FC

and rCBF at the DMN (red lines), Vis (green lines) or SM (blue lines) voxels. The correlations

between the DMN FC and averaged rCBF in the DMN, Vis or SM region are represented by

thickened lines in Figure 5a. A correlation map (Figure 5b) between the DMN FC and rCBF across

the entire brain voxels was generated. The procedures of generating this correlation map are

illustrated in Figure 5-figure supplement 1. The DMN, Vis and SM ROIs in Figure 5b were

delineated with dashed red, green and blue contours, respectively, and obtained from Figure 1-

figure supplement 2b. Most of significant correlations ($r > r_{crit}$) between the DMN FC and voxel-

wise rCBF were found in the voxels in the DMN regions, such as PCC, IPL and LTC, but not in

the Vis or SM regions (Figure 5b). Demonstrated in a radar plot in Figure 5c, much higher percent

of voxel with significant correlations between rCBF and the DMN FC was found in the DMN

(36.7%, $p < 0.0001$) regions, than in the SM (14.6%, $p > 0.05$) or Vis (5.5%, $p > 0.05$) regions.

Statistical significance of higher percent of voxels with significant correlations in the DMN

($p<0.0001$) was confirmed using nonparametric permutation tests with 10,000 permutations. We

also conducted the correlation between the Vis FC and rCBF across the brain as well as

permutation test. As expected, no significant correlation between the Vis FC and rCBF can be

found in any voxel in the DMN, Vis or SM ROIs, demonstrated in Figure 5-figure supplement 2a.

Similar analysis was also conducted for correlation between the SM FC and rCBF across the

brain and percent of voxels with significant correlation was close to zero, as demonstrated in

Figure 5-figure supplement 2b. Combined with the results shown in Figure 5, the results of

coupling between Vis (Figure 5-figure supplement 2a) or SM (Figure 5-figure supplement 2b) FC

and rCBF further demonstrated selected rCBF-FC coupling can be only found in the DMN ROIs,

but not in the Vis or SM network ROIs.

DISCUSSION

We revealed strongly coupled rCBF and FC increases specifically in the DMN while establishing unprecedented 4D rCBF spatiotemporal dynamics during infancy. The tight rCBF-FC relationship found with multi-modal infant MRI suggests DMN emergence is supported by faster local blood flow increase in the DMN to meet metabolic demand, offering refreshing insight into the physiological mechanism underlying early brain functional architecture emergence. The delineated 4D brain perfusion spatiotemporal framework was characterized with heterogeneous rCBF distribution across brain regions at a specific age and differential age-dependent rCBF increase rates across brain regions during infant development, and can be used as quantified standard reference for detecting rCBF alterations (e.g. the z scores) of atypically developing brains. Elucidating the ontogeny of infant brain physiology and its functional correlates could greatly advance current understanding of general principles of early brain development.

Gradient of functional network maturations in early brain development has been more extensively characterized with recent rs-fMRI studies. Differential emergence of these functional networks is distinguished by different onset time as well as different maturational rate of various brain functions in a given developmental period. For example, primary sensory and motor functional networks, such as the SM and Vis networks, appear earlier before or around birth (Cao et al., 2017a; Doria et al., 2010; Fransson et al., 2007; Smyser et al., 2010; Peng et al., 2020). Other functional networks involved in heteromodal functions appear later. The DMN (Fox et al., 2007; Greicius et al., 2003; Greicius et al., 2009; Raichle et al., 2001; Raichle, 2015; Smith et al., 2009) is a higher-order functional network. Smyser et al (Smyser et al., 2010) found that SM and Vis functional networks mature earlier and demonstrate adult-like pattern for preterm neonate brain, with the DMN much immature and incomplete around birth. Cao et al (Cao et al., 2017a) also found rapid maturation of primary sensorimotor functional systems in preterm neonates from 31 to 41 postmenstrual weeks while the DMN remained immature during that period. These previous studies suggest that significant functional maturation in primary sensorimotor networks

occur earlier in preterm and perinatal developmental period (Cao et al., 2017a; Doria et al., 2010;

Smyser et al., 2010) compared to 0-24-month infancy focused in the present study. Functional

network emergence in the DMN was found in the developmental infant cohort in Figure 3, marking

significant maturation of the DMN in infancy and distinguished network pattern from earlier

developmental period. The delineated DMN emergence in this study is also consistent to the

literature (Gao et al., 2009). Figure 3-figure supplement 1 further demonstrated significant

increase of FC only in the DMN, but not in primary sensorimotor system that already emerged in

earlier developmental period.

Glucose and oxygen are two primary molecules for energy metabolism in the brain

(Raichle et al., 2001; Vaishnavi et al., 2010). Glucose consumed by infant brain represents 30%

total amount of glucose (Raichle, 2010; Settergren et al., 1976), more than 15-20% typically seen

in adult brain (Bouma and Muizelaar, 1990; Satterthwaite et al., 2014). The cerebral metabolic

rate for glucose (CMRGlu) and oxygen (CMRO2) are direct measures of the rate of energy

consumption, which parallel the proliferation of synapses in brain during infancy (Raichle, 2010).

RCBF delivering glucose and oxygen for energy metabolism in the brain is closely related to

CMRGlu and CMRO2 and can serve as a surrogate of these two measurements (Fox and Raichle,

1986; Gur et al., 2009; Paulson et al., 2010; Vaishnavi et al., 2010). In the PET study (Chugani

and Phelps, 1986) using CMRGlu measurements, it was found that the local CMRGlu in the

sensorimotor cortex almost reaches the highest level in early infancy and then plateaus during

rest of infancy, consistent with relatively small changes of rCBF in later infancy in primary

sensorimotor ROIs found in this study (Figure 2). On the other hand, the global CBF measured

with PC MRI (Liu et al., 2019) increases dramatically during infancy with global CBF at 18 months

almost 5 times of the CBF around birth. Taken together, the literature suggests significant but

nonuniformly distributed CBF increases across the brain regions during infancy, consistent to the

measured heterogeneous rCBF increase pattern (Figures 1 and 2) in the present study.

Furthermore, the differentiated cerebral metabolic pattern reflected by measured rCBF

distribution (Figure 1) at a specific age and differential increase rates of the rCBF (Figure 2) from

0 to 24 months are strikingly consistent with spatiotemporally differentiated functional (Cao et al.,

2017a; Doria et al., 2010; Fransson et al., 2007; Gao et al., 2009; Smyser et al., 2010) and

structural (Ouyang et al., 2019a) maturational processes. In developmental brains, cellular

processes supporting differential functional emergence require extra oxygen and glucose delivery

through cerebral blood flow to meet the metabolic demand. In Hebb's principle, "neurons that fire

together wire together". Through the synaptogenesis in neuronal maturation, the neurons within

a certain functional network system tend to have more synchronized activity in a more mature

stage than in an immature stage. As shown in the diagram in Figure 6, cellular activities in

developmental brains, such as synaptogenesis critical for brain circuit formation, need extra

energy more than that in the stable and matured stage. Neurons do not have internal reserves of

energy in the form of sugar or oxygen. The demand of extra energy requires an increase in rCBF

to deliver more oxygen and glucose for formation of brain networks. In the context of infant brain

development, there is a cascade of events of CBF increase, CMRO2 and CMRGlu increase,

synaptogenesis and synaptic efficacy increase, blood oxygenation level dependent (BOLD) signal

synchronization increase, and functional connectivity increase, shown in the bottom of Figure 6.

Such spatial correlation of rCBF to the FC in the functional network ROIs was found in adults

(Liang et al., 2013). Higher rCBF has also been found in the DMN in children 6 to 20 years of age

(Liu et al., 2018). Consistent with the diagram shown in Figure 6, Figure 4 revealed significant

correlation between FC and rCBF in the DMN network, and Figure 5 identified this significant

correlation between FC and rCBF specifically in the DMN network, but not in the primary Vis or

SM networks. Collectively, these results (Figures 4 and 5) were well aligned with the hypothesis

that faster rCBF increase in the DMN underlies emergence of the DMN reflected by significant

FC increases. The revealed physiology-function relationship may shed light on physiological

underpinnings of brain functional network emergence.

It is noteworthy to highlight a few technical details below. First, present study benefits from

multimodal MRI allowing measuring functional network emergence and rCBF of the same cohort

of infants. Simultaneous rCBF and FC measurements enabled us to probe relationship of brain

physiology and function during infant development. Second, a nonparametric permutation

analysis without a prior hypothesis at the voxel level across the whole brain was conducted to

confirm the coupling of rCBF and FC is specific in the DMN regions (Figure 5), not in the primary

sensorimotor (Vis or SM) regions (Figure 5-figure supplement 2), demonstrating the robustness

of the results on physiological underpinning of functional network emergence in the DMN. Third,

the pCASL perfusion MRI was calibrated by PC MRI so that the errors caused by varying labeling

efficiency among infant subjects can be ameliorated. This calibration process therefore enhanced

accuracy of rCBF measurements for their potential use as a standard reference. Fourth, the ROIs

for rCBF measurements were obtained by data-driven ICA analysis of the same sub-cohort of

infant subjects aged 12 to 24 months instead of transferred ROIs from certain brain parcellation

atlases. Since most of parcellation atlases were built based on adult brain data and all these

atlases were established based on other subject groups, ROIs delineated from the same cohort

improve accuracy of coupling analysis. There are several limitations in the present study that can

be improved in future investigations. All data were acquired from a cross-sectional cohort. To

minimize the inter-subject variability, future study with a longitudinal cohort of infants is warranted.

With relatively small size of infant brains, spatial resolution of the pCASL and rs-fMRI can be

further improved too to improve imaging measurement accuracy. To further improve the statistical

power, larger infant sample size will be beneficial in the future studies. Finally, although the rCBF

from pCASL perfusion MRI is highly correlated with the CMRO2 and CMRGlu measured from

PET, rCBF is not direct measurement of rate of energy consumption. Physiology-function

relationship studies in infants could benefit from development of novel noninvasive MR imaging

methods as an alternative to PET to measure CMRO2 or CMRGlu without involvement of

radioactive tracer.

CONCLUSION

Novel findings in this study inform a physiological mechanism of DMN emergence during infancy

with rCBF and FC measurements from multimodal MRI in developmental infant brains. The age-

specific whole-brain rCBF maps and spatiotemporal rCBF maturational charts in all brain regions

serve as a standardized reference of infant brain physiology for precision medicine. The rCBF-

FC coupling results revealing fundamental physiology-function relationship have important

implications in altered network maturation in developmental brain disorders.

MATERIALS AND METHODS

Infant subjects. Forty-eight infants (30 males) aged 0-24 months (14.6 ± 6.32 months) were recruited at Beijing Children's Hospital. These infants were referred to MR imaging due to seizures with fever (n = 21), convulsion (n = 17), diarrhea (n = 9), or sexual precocity (n = 1). All infants had normal neurological examinations documented in medical record. The exclusion criteria include known nervous system disease, or history of neurodevelopmental or systemic illness. Every infant's parents provided signed consent and the protocol was approved by Beijing Children's Hospital Research Ethics Committee (Approval number 2016-36).

Data acquisition. All infant MR scans including pCASL, PC MRI, rs-fMRI and structural MRI were acquired with the same 3T Philips Achieva system under sedation. Earplug and headphones were used to minimize noise exposure. Pseudo-continuous arterial spin labeling (pCASL) perfusion MRI images were acquired using a multi-slice echo planar imaging with following parameters: TR = 4100 ms, TE = 15 ms, 20 slices with 5 mm slice thickness and no gap between slices, field of view (FOV) = 230 × 230 mm^2, matrix size = 84 × 84, voxel size = 2.74 × 2.74 × 5 mm^3. As shown on the left panel of Figure 1a, the labeling slab was placed at the junction of spinal cord and medulla (65 mm below central slab of imaging volume) and parallel to the anterior commissure-posterior commissure (AC-PC) line. The labeling duration was 1650 ms and the post labeling delay (PLD) was 1600 ms. 30 pairs of control and label volume were acquired for each infant. The acquisition time of the pCASL perfusion MRI images was 4.2 min. An auxiliary scan with identical readout module to pCASL but without labeling was acquired for estimating the value of equilibrium magnetization of brain tissue. Phase contrast (PC) MRI was acquired to calibrate regional cerebral blood flow (rCBF) by scaling the overall CBF of entire brain. The carotid and vertebral arteries were localized based on a scan of time-of-flight (TOF) angiography acquired with following parameters: TR = 20 ms, TE = 3.45 ms, flip angle = 30°, 30 slices, FOV= 100 × 100 mm^2, matrix size = 100 × 100, voxel size = 1 × 1 × 1 mm^3. The acquisition time of the TOF

angiography was 28 s. Based on the angiography, the slices for the PC MRI of internal carotid

arteries were placed at the level of the foramen magnum and the slices for the PC MRI of vertebral

arteries were placed between the two turns in V3 segments, as illustrated on the right panel of

Figure 1a. For left and right internal carotid and vertebral arteries, PC MRI images were acquired

with following parameters: TR = 20 ms, TE = 10.6 ms, flip angle = 15°, single slice, FOV = 120 ×

120 mm^2, matrix size = 200 × 200, voxel size = 0.6 × 0.6 × 3 mm, maximum velocity encoding=

40 cm/s, non-gated, 4 repetitions. The acquisition time of PC MRI for each artery was 24 s.

Resting state fMRI (rs-fMRI) images were acquired using echo planar imaging with following

parameters: TR = 2000 ms, TE = 24 ms, flip angle = 60°, 37 slices, FOV = 220 × 220 mm^2, matrix

size = 64 × 64, voxel size = 3.44 × 3.44 × 4 mm^3. 200 dynamics were acquired for each infant.

The acquisition time of the rs-fMRI images was 7 min. T1 weighted images of all infants were also

acquired for anatomical information and brain segmentation using MPRAGE (Magnetization

Prepared - RApid Gradient Echo) sequence with following parameters: TR = 8.28 ms, TE = 3.82

ms, flip angle = 12°, 150 slices, FOV = 200 × 200 mm^2, matrix size = 200 × 200, voxel size = 1 ×

1 × 1 mm^3. The acquisition time of the T1-weighted image was 3.7 min. Visual inspection was

carefully conducted for all MRI data by experienced pediatric radiologists (D.H. and Y.P.) with

decades of experience in clinical radiology. No significant motion artifacts were spotted with the

sedated MRI scans. Therefore, no dataset from the forty-eight infants was excluded from following

data analysis due to severe motion artifacts.

Measurement of rCBF with pCASL perfusion MRI and calibrated by PC MRI. After head

motion correction of the pCASL perfusion MRI, we estimated rCBF using the protocol similar to

that in our previous publication (Ouyang et al., 2017). Briefly, rCBF was measured using a model

described in ASL white paper (Alsop et al., 2015):

$$\text{rCBF} = \frac{6000 \cdot \lambda \cdot \Delta M \cdot e^{\frac{PLD}{T_{1a}}}}{2 \cdot \alpha \cdot M_b^0 \cdot T_{1a} \cdot \left(1 - e^{\frac{-LabelDur}{T_{1a}}}\right)} \ [ml/100g/min] \qquad\qquad [1]$$

where ΔM is the dynamic-averaged signal intensity difference between in the control and label

images; λ, the blood-brain partition coefficient, is 0.9 ml/g (Herscovitch and Raichle, 1985); PLD,

the post labeling delay time, is the cumulation of 1650 ms and the delayed time between slices;

LabelDur, the labeling duration, is 1600 ms; α, the labeling efficiency, is 0.86 predicted by the

fitting between labeling efficiency and blood velocity in the previous study (Aslan et al., 2010); T_{1a},

T_1 of arterial blood, is 1800 ms (Liu et al., 2016; Varela et al., 2011). The value of equilibrium

magnetization of brain tissue (M_b^0) was obtained from an auxiliary scan with identical readout

module of pCASL except labeling. The labeling efficiency α can vary considerably across

participants, especially in infants. Thus, we used PC MRI to estimate and calibrate rCBF

measures, as described previously (Aslan et al., 2010; Ouyang et al., 2017). To calibrate rCBF,

global CBF from PC MRI was calculated as follows:

$$f_{PC,AVG} = \int v \, dA \, / \, (\rho * brain\ volume) \qquad [2]$$

where v is the blood flow velocity in the ICAs and VAs; A is the cross-sectional area of the blood

vessel with the unit mm^2; and the brain tissue density ρ is assumed as 1.06 g/mL (Dittmer and

Dawson, 1961; Herscovitch and Raichle, 1985). Brain volume was measured from the T1-

weighted image as parenchyma volume (gray matter + white matter volume). RCBF was

calibrated by applying the scalar factor making averaged rCBF equal to global CBF from PC MRI.

To demonstrate the reproducibility of the adopted pCASL protocol, the intraclass correlation

coefficient (ICC) was calculated based on entire brain rCBF maps measured from first half and

second half of control/label series of pCASL scan of a randomly selected infant subject aged 17.6

months.

Rs-fMRI preprocessing. Same preprocessing procedures elaborated in our previous publication

(Cao et al., 2017) was used. Briefly, the normalized rs-fMRI images underwent spatially

smoothing with a Gaussian kernel (full width at half-maximum of 4mm), linear trend removal, and

temporal band-pass filtering (0.01–0.10 Hz). Several nuisance variables, including 6 rigid-body

head motion parameters and the averaged signal from white matter and cerebrospinal fluid (CSF)

tissue, were removed through multiple linear regression analysis to reduce effects of non-

neuronal signals. The CSF and white matter were segmented with the T1-weighted image using

SPM. Preprocessed rs-fMRI signals were used to estimate functional connectivity (FC), defined

as Pearson's correlation between the time courses of preprocessed rs-fMRI blood oxygenation

level dependent (BOLD) signal in two regions or two voxels (Figure 1-figure supplement 2a).

Multi-modal image registration to a customized template space from all subjects. For

integrating perfusion MRI and rs-fMRI data of all infant subjects, a customized structural template

was generated. T1-weighted image of a 12-month-old brain characterized by median brain size

at this age and straight medial longitudinal fissure was used as a single-subject template. T1-

weighted images of all subjects were registered to the single-subject template by using nonlinear

registration in Statistical Parametric Mapping (SPM 8, http://www.fil.ion.ucl.ac.uk/spm). The

averaged T1-weighted image in the template space was defined as the structural template. In

individual space, after slice timing and head motion correction, intra-subject registration of rs-fMRI

to T1-weighted image was conducted by transforming the averaged images across dynamics in

each session to the T1-weighted image of the same subject through linear registration with SPM.

This intra-subject transformation was applied to each volume of rs-fMRI. As described in the main

text, rCBF map was estimated in the individual space using Eq [1]. Intra-subject registration of

rCBF map to T1-weighted image was also conducted through linear registration with SPM. RCBF

maps and rs-fMRI images aligned to T1-weighted images in the individual space of each subject

were then normalized to the customized infant template space through the same nonlinear

registration from individual T1-weighted image to the structural template. After nonlinear inter-

subject normalization, rCBF map was smoothed spatially in the customized template space.

Identification of functional network regions-of-interests (ROIs) with rs-fMRI. Brain regions

of three functional networks, including default-mode network (DMN), visual (Vis) network, and

sensorimotor (SM) network, were used as functional ROIs for quantifying rCBF. These functional

networks were identified with rs-fMRI of 35 infants aged 12 to 24 months, as DMN can be better

delineated in later infancy than earlier infancy. Independent Component Analysis (ICA)

(Beckmann et al., 2005) in the FSL (https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/MELODIC) was used to

identify all these network regions. Individual DMN regions, including posterior cingulate cortex

(PCC), medial prefrontal cortex (MPFC), inferior parietal lobule (IPL) and lateral temporal cortex

(LTC), were extracted from the independent components (ICs) based on the spatially distributed

regions consistently identified across studies (Greicius et al., 2004; Smith et al., 2009). Vis and

SM network regions were also identified following the literature (Smith et al., 2009). The functional

network ROIs were obtained by thresholding the ICs at z value of 1.96 which corresponds to p

value of 0.05. The identified ROIs for DMN, Vis and SM networks are shown in Figure 1-figure

supplement 2b. ROIs of DMN, Vis or SM include all regions in each network, respectively.

Characterization of age-dependent changes of rCBF and FC. After multi-modal images from

all subjects were registered in the same template space, age-dependent rCBF and FC changes

were characterized using linear regression. The linear cross-sectional developmental trajectory

of voxel-wise rCBF increase was obtained by fitting rCBF measurements in each voxel with ages

across subjects. The fitted rCBF at different ages was estimated and projected onto the template

cortical surface with Amira (FEI, Hillsboro, OR) to show the spatiotemporal changes of rCBF

across cortical regions during infancy. The rCBF-age correlation coefficients r values of all voxels

were estimated and mapped to the cortical surface, resulting in the correlation coefficient map.

The functional ROIs identified above were used to quantify regional rCBF change. The rCBF in

the SM, Vis, and DMN ROIs were averaged across voxels within each ROI, and fitted with a linear

model: $\mathrm{rCBF(t)} = \alpha + \beta\, t + \varepsilon$, where α and β were intercepts and slopes for rCBF measured at

certain ROI, t was the infant age in months and ε was the error term. To test if regional rCBF

increased significantly with age, the null hypothesis was that rCBF slope in each ROI was equal

to zero. To compare regional rCBF change rates between the tested ROI and SM ROI, the null

hypothesis was that the rCBF slope of the tested ROI and rCBF slope of SM ROI were equal.

Rejection of the null hypothesis indicated a significant rCBF slope difference between two ROIs.

Nonlinear models, including exponential and biphasic models, were compared with linear

model using F-test for fitting regional rCBF in the DMN, SM and Vis ROIs. No significant difference

was found between linear and exponential (all $F(1,45) < 3.45$, $p > 0.05$, uncorrected) fitting or

between linear and biphasic (all $F(2,44) < 2.64$, $p > 0.05$, uncorrected) fitting in the regional rCBF

changes in the DMN, SM and Vis ROIs.

In the customized template space, time-dependent changes of FC to PCC and time-

dependent changes of FC within the functional network ROIs were calculated. FC of any given

voxel outside the PCC to PCC was defined as correlation between signal time course of this given

voxel and the averaged signal time course of all voxels within the PCC. Emergence of the DMN

was delineated by linear fitting of the age-related increase of FC to PCC in each voxel across

subjects. FC within each network was defined as mean of functional connectivity strength (Cao

et al., 2017b) of each voxel within this network region. The FC within the DMN, SM or Vis was

also fitted with a linear model: $\mathrm{FC(t)} = \alpha + \beta\, t + \varepsilon$, where α and β were intercepts and slopes for FC

measured within a certain network, t was the infant age in months and ε was the error term. To

test if the FC within a network increased significantly with age, the null hypothesis was that rCBF

slope in each ROI was equal to zero.

Test of heterogeneity of rCBF across functional network ROIs. To examine heterogeneity of

infant rCBF in different brain regions, rCBF measurements of all infants were averaged across

voxels in the Vis, SM and DMN (including DMN subregions DMN_PCC, DMN_MPFC, DMN_IPL

and DMN_LTC), respectively. To test significant difference of the rCBF values among different

functional network ROIs, a one-way analysis of variance (ANOVA) with repeated measures was

conducted. Paired t-tests were also conducted to test the difference of rCBF measurements

between regions. False discovery rate of each test was corrected to control the type I error.

Coupling between rCBF and FC during the infant brain development. Coupling between

rCBF and FC in the DMN was conducted with voxel-wise approach. The FC of a voxel in the DMN

was the average of correlations of rs-fMRI BOLD signal between this voxel and all other DMN

voxels. All infants were divided into two groups based on their ages, 0-12 months and 12-24

months. 4000 voxels were randomly chosen from the DMN voxels of all subjects in each age

group for regression analysis. Since the variance of both FC and rCBF cannot be ignored in this

study, Deming regression (Deming, 1943) was used to fit the trendline of coupling between FC

and rCBF. We further tested whether significant FC-rCBF coupling was specifically localized in

the DMN, but not in primary sensorimotor (Vis or SM) regions. FC within a specific network was

calculated by averaged FC of all voxels in this network. As demonstrated in Figure 5-figure

supplement 1, the correlation between FC within a network and the rCBF at each voxel resulted

in a whole-brain r map (e.g. Figure 5b). A nonparametric permutation test was then applied to

evaluate the significance of rCBF-FC correlation in the ROIs of a specific brain network (e.g. DMN,

SM or Vis). The null hypothesis is that the voxels with significant correlation ($r > 0.28$) between

rCBF and FC are distributed evenly in the brain. To test the null hypothesis, we resampled the

correlation coefficient r of all brain voxels randomly for 10,000 times to build 10,000 whole-brain

correlation coefficient distribution maps. The FC-rCBF correlation was considered significant in

certain brain network ROIs if the number of observed significant voxels in the network ROIs is

higher than the number of significant voxels corresponding to 95th percentile in the permutation

tests.

REFERENCES

Alsop, D.C., Detre, J.A., Golay, X., Günther, M., Hendrikse, J., Hernandez‑Garcia, L., Lu, H., MacIntosh, B.J., Parkes, L.M., Smits, M., et al. (2015). Recommended implementation of arterial spin‑labeled perfusion MRI for clinical applications: A consensus of the ISMRM perfusion study group and the European consortium for ASL in dementia. Magn Reson Med. 73, 102-116.

Aslan, S., Xu, F., Wang, P.L., Uh, J., Yezhuvath, U.S., Van Osch, M., and Lu, H. (2010). Estimation of labeling efficiency in pseudocontinuous arterial spin labeling. Magn Reson Med. 63, 765-771.

Beckmann CF, DeLuca M, Devlin JT & Smith SM. Investigations into resting-state connectivity using independent component analysis. *Philos Trans R Soc Lond B Biol Sci* 360, 1001-1013 (2005).

Bouma, G.J. and Muizelaar, J.P. (1990). Relationship between cardiac output and cerebral blood flow in patients with intact and with impaired autoregulation. J Neurosurg 73, 368-374..

Buckner, R.L., Andrews-Hanna, J.R., and Schacter, D.L. (2008). The brain's default network: anatomy, function, and relevance to disease. Ann. N. Y. Acad. Sci. 1124, 1–38.

Bystron, I., Blakemore, C., and Rakic, P. (2008). Development of the human cerebral cortex: Boulder Committee revisited. Nat. Rev. Neurosci. 9, 110-122.

Cao, M., He, Y., Dai, Z., Liao, X., Jeon, T., Ouyang, M., Chalak, L., Bi, Y., Rollins, N., Dong, Q., et al. (2017b). Early development of functional network segregation revealed by connectomic analysis of the preterm human brain. Cereb. Cortex 27, 1949-1963.

Cao, M., Huang, H., and He, Y. (2017a). Developmental connectomics from infancy through early childhood. Trends Neurosci. 40, 494-506.

Chiron, C., Raynaud, C., Mazière, B., Zilbovicius, M., Laflamme, L., Masure, M.C., Dulac, O., Bourguignon, M., and Syrota, A. (1992). Changes in regional cerebral blood flow during brain maturation in children and adolescents. J. Nucl. Med.: official publication, Society of Nuclear Medicine 33, 696-703.

Chugani, H.T., and Phelps, M.E. (1986). Maturational changes in cerebral function in infants determined by 18FDG positron emission tomography. Science 231, 840-843.

Chugani, H.T., Phelps, M.E., and Mazziotta, J.C. (1987). Positron emission tomography study of human brain functional development. Ann. Neurol. 22, 487-497.

Deming, W. E. (1943). Statistical adjutement of data. Wiley, NY.

Detre, J.A., and Alsop, D.C. (1999). Perfusion magnetic resonance imaging with continuous arterial spin labeling: methods and clinical applications in the central nervous system. Eur J Radiol. 30, 115-124.

Dittmer, J., and Dawson, R. (1961). The isolation of a new lipid, triphosphoinositide, and monophosphoinositide from ox brain. Biochemical Journal 81, 535.

Doria, V., Beckmann, C.F., Arichi, T., Merchant, N., Groppo, M., Turkheimer, F.E., Counsell, S.J., Murgasova, M., Aljabar, P., Nunes, R.G., et al. (2010). Emergence of resting state networks in the preterm human brain. Proc. Natl. Acad. Sci. USA 107, 20015-20020.

Doyle‑Thomas, K.A., Lee, W., Foster, N.E., Tryfon, A., Ouimet, T., Hyde, K.L., Evans, A.C., Lewis, J., Zwaigenbaum, L., Anagnostou, E., et al. (2015). Atypical functional brain connectivity during rest in autism spectrum disorders. Ann. Neurol. 77, 866-876.

Fox, M.D., Snyder, A.Z., Vincent, J.L., and Raichle, M.E. (2007). Intrinsic fluctuations within cortical systems account for intertrial variability in human behavior. Neuron 56, 171-184.

Fox, P.T., and Raichle, M.E. (1986). Focal physiological uncoupling of cerebral blood flow and oxidative metabolism during somatosensory stimulation in human subjects. Proc. Natl. Acad. Sci. USA 83, 1140-1144.

Fransson, P., Skiöld, B., Horsch, S., Nordell, A., Blennow, M., Lagercrantz, H., and Åden, U. (2007). Resting-state networks in the infant brain. Proc. Natl. Acad. Sci. USA 104, 15531-15536.

Gao, W., Zhu, H., Giovanello, K.S., Smith, J.K., Shen, D., Gilmore, J.H., and Lin, W. (2009). Evidence on the emergence of the brain's default network from 2-week-old to 2-year-old healthy pediatric subjects. Proc. Natl. Acad. Sci. USA 106, 6790-6795.

Greicius, M.D., Krasnow, B., Reiss, A.L., and Menon, V. (2003). Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. Proc. Natl. Acad. Sci. USA 100, 253-258.

Greicius MD, Srivastava G, Reiss AL & Menon V. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. *Proc. Natl. Acad. Sci. U.S.A* 101, 4637-4642 (2004).

Greicius, M.D., Supekar, K., Menon, V., and Dougherty, R.F. (2009). Resting-state functional connectivity reflects structural connectivity in the default mode network. Cereb. Cortex 19, 72-78.

Gur, R.C., Ragland, J.D., Reivich, M., Greenberg, J.H., Alavi, A., and Gur, R.E. (2009). Regional differences in the coupling between resting cerebral blood flow and metabolism may indicate action preparedness as a default state. Cereb. Cortex 19, 375-382.

Haynes, R.L., Borenstein, N.S., Desilva, T.M., Folkerth, R.D., Liu, L.G., Volpe, J.J., and Kinney, H.C. (2005). Axonal development in the cerebral white matter of the human fetus and infant. J. Comp. Neurol. 484, 156-167.

Herscovitch, P., and Raichle, M.E. (1985). What is the correct value for the brain-blood partition coefficient for water? J. Cereb. Blood Flow Metab. 5, 65-69.

Huang, H., and Vasung, L. (2014). Gaining insight of fetal brain development with diffusion MRI and histology. Int. J. Dev. Neurosci. 32, 11-22.

Huang, H., Jeon, T., Sedmak, G., Pletikos, M., Vasung, L., Xu, X., Yarowsky, P., Richards, L.J.,
Kostović, I., Šestan, N., et al. (2013). Coupling diffusion imaging with histological and gene
expression analysis to examine the dynamics of cortical areas across the fetal period of
human brain development. Cereb Cortex 23, 2620-2631.

Huttenlocher, P.R., and Dabholkar, A.S. (1997). Regional differences in synaptogenesis in human
cerebral cortex. J Comp Neurol 387, 167-178.

Innocenti, G.M., and Price, D.J. (2005). Exuberance in the development of cortical networks. Nat.
Rev. Neurosci. 6, 955-965.

Jain, V., Duda, J., Avants, B., Giannetta, M., Xie, S.X., Roberts, T., Detre, J.A., Hurt, H., Wehrli,
F.W., and Wang, D.J. (2012). Longitudinal reproducibility and accuracy of pseudo-
continuous arterial spin–labeled perfusion MR imaging in typically developing children.
Radiology 263, 527-536.

Knickmeyer, R.C., Gouttard, S., Kang, C., Evans, D., Wilber, K., Smith, J.K., Hamer, R.M., Lin,
W., Gerig, G. and Gilmore, J.H. (2008). A structural MRI study of human brain
development from birth to 2 years. J. Neurosci. 28, 12176-12182.

Liang, X., Zou, Q., He, Y., and Yang, Y. (2013). Coupling of functional connectivity and regional
cerebral blood flow reveals a physiological basis for network hubs of the human brain.
Proc. Natl. Acad. Sci. USA 110, 1929-1934.

Liu, F., Duan, Y., Peterson, B.S., Asllani, I., Zelaya, F., Lythgoe, D., and Kangarlu, A. (2018).
Resting state cerebral blood flow with arterial spin labeling MRI in developing human
brains. Eur. J. Paediatr. Neurol. 22, 642-651.

Liu, P., Chalak, L.F., Krishnamurthy, L.C., Mir, I., Peng, S.L., Huang, H., and Lu, H. (2016). T1
and T2 values of human neonatal blood at 3 Tesla: dependence on hematocrit,
oxygenation, and temperature. Magn Reson Med. 75, 1730-1735.

Liu, P., Qi, Y., Lin, Z., Guo, Q., Wang, X. and Lu, H. (2019). Assessment of cerebral blood flow in
neonates and infants: a phase-contrast MRI study. NeuroImage 185, 926-933.

Lynch, C.J., Uddin, L.Q., Supekar, K., Khouzam, A., Phillips, J., and Menon, V. (2013). Default
mode network in childhood autism: posteromedial cortex heterogeneity and relationship
with social deficits. Biol. Psychiatry 74, 212-219.

Miller, D.J., Duka, T., Stimpson, C.D., Schapiro, S.J., Baze, W.B., McArthur, M.J., Fobbs, A.J.,
Sousa, A.M., Šestan, N., Wildman, D.E., et al. (2012). Prolonged myelination in human
neocortical evolution. Proc. Natl. Acad. Sci. USA 109, 16480-16485.

Ouyang, M., Dubois, J., Yu, Q., Mukherjee, P., and Huang, H. (2019a). Delineation of early brain
development from fetuses to infants with diffusion MRI and beyond. Neuroimage 185, 836-
850.

Ouyang, M., Jeon, T., Sotiras, A., Peng, Q., Mishra, V., Halovanic, C., Chen, M., Chalak, L., Rollins, N., Roberts, T.P., et al. (2019b). Differential cortical microstructural maturation in the preterm human brain with diffusion kurtosis and tensor imaging. Proc. Natl. Acad. Sci. USA 116, 4681-4688.

Ouyang, M., Liu, P., Jeon, T., Chalak, L., Heyne, R., Rollins, N.K., Licht, D.J., Detre, J.A., Roberts, T.P., Lu, H., et al. (2017). Heterogeneous increases of regional cerebral blood flow during preterm brain development: Preliminary assessment with pseudo-continuous arterial spin labeled perfusion MRI. NeuroImage 147, 233-242.

Padmanabhan, A., Lynch, C.J., Schaer, M., and Menon, V. (2017). The default mode network in autism. Biol Psychiatry Cogn Neurosci Neuroimaging 2, 476-486.

Paulson, O.B., Hasselbalch, S.G., Rostrup, E., Knudsen, G.M., and Pelligrino, D. (2010). Cerebral blood flow response to functional activation. J. Cereb. Blood Flow Metab. 30, 2-14.

Peng, Q., Ouyang, M., Wang, J., Yu, Q., Zhao, C., Slinger, M., Li, H., Fan, Y., Hong, B., and Huang, H. (2020). Regularized-Ncut: Robust and homogeneous functional parcellation of neonate and adult brain networks. Artif Intell Med 106, 101872.

Pfefferbaum, A., Mathalon, D.H., Sullivan, E.V., Rawles, J.M., Zipursky, R.B. and Lim, K.O. (1994). A quantitative magnetic resonance imaging study of changes in brain morphology from infancy to late adulthood. J. Neurol. 51, 874-887.

Pontzer H, Yamada Y, Sagayama H, Ainslie P, Andersen LF, et a., (2021). Daily energy expenditure through the human life course. Science, 373: 808-812.

Raichle, M.E. (2010). Two views of brain function. Trends Cogn. Sci. 14, 180-190.

Raichle, M.E. (2015). The brain's default mode network. Annu. Rev. Neurosci. 38, 433-447.

Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., and Shulman, G.L. (2001). A default mode of brain function. Proc. Natl. Acad. Sci. USA 98, 676-682.

Rakic, P. (1995). Radial versus tangential migration of neuronal clones in the developing cerebral cortex. Proc. Natl. Acad. Sci. USA 92, 11323.

Satterthwaite, T.D., Shinohara, R.T., Wolf, D.H., Hopson, R.D., Elliott, M.A., Vandekar, S.N., Ruparel, K., Calkins, M.E., Roalf, D.R., Gennatas, E.D., et al. (2014). Impact of puberty on the evolution of cerebral perfusion during adolescence. Proc. Natl. Acad. Sci. USA 111, 8643-8648.

Settergren, G., Lindblad, B.S., and Persson, B. (1976). Cerebral blood flow and exchange of oxygen, glucose, ketone bodies, lactate, pyruvate and mino acides in infants. Acta Pediatrica Scand. 65, 343-353.

Sidman, R.L., and Rakic, P. (1973). Neuronal migration, with special reference to developing human brain: a review. Brain Res. 62, 1-35..

Silbereis, J.C., Pochareddy, S., Zhu, Y., Li, M., and Sestan, N. (2016). The cellular and molecular landscapes of the developing human central nervous system. Neuron 89, 248-268.

699 Smith, S.M., Fox, P.T., Miller, K.L., Glahn, D.C., Fox, P.M., Mackay, C.E., Filippini, N., Watkins,
700   K.E., Toro, R., Laird, A.R., et al. (2009). Correspondence of the brain's functional
701   architecture during activation and rest. Proc. Natl. Acad. Sci. USA 106, 13040-13045.
702

703 Smyser, C.D., Inder, T.E., Shimony, J.S., Hill, J.E., Degnan, A.J., Snyder, A.Z., and Neil, J.J.
704   (2010). Longitudinal analysis of neural network development in preterm infants. Cereb.
705   Cortex 20, 2852-2862.
706

707 Tau, G.Z., and Peterson, B.S. (2010). Normal development of brain circuits.
708   Neuropsychopharmacology 35, 147.
709

710 Vaishnavi, S.N., Vlassenko, A.G., Rundle, M.M., Snyder, A.Z., Mintun, M.A., and Raichle, M.E.
711   (2010). Regional aerobic glycolysis in the human brain. Proc. Natl. Acad. Sci. USA 107,
712   17757-17762.
713

714 Varela, M., Hajnal, J.V., Petersen, E.T., Golay, X., Merchant, N., and Larkman, D.J. (2011). A
715   method for rapid in vivo measurement of blood T1. NMR Biomed. 24, 80-88.
716

717 Wang, Z., Fernández-Seara, M., Alsop, D.C., Liu, W.C., Flax, J.F., Benasich, A.A., and Detre, J.A.
718   (2008). Assessment of functional development in normal infant brain using arterial spin
719   labeled perfusion MRI. Neuroimage 39, 973-978.
720

721 Washington, S.D., Gordon, E.M., Brar, J., Warburton, S., Sawyer, A.T., Wolfe, A., Mease‑
722   Ference, E.R., Girton, L., Hailu, A., Mbwana, J., et al. (2014). Dysmaturation of the default
723   mode network in autism. Hum Brain Mapp 35, 1284-1296.
724

725 Yakovlev, P. (1967). The myelogenetic cycles of regional maturation of the brain. Regional
726   development of the brain in early life, 3-70.
727

728 Yu, Q., Ouyang, A., Chalak, L., Jeon, T., Chia, J., Mishra, V., Sivarajan, M., Jackson, G., Rollins,
729   N., Liu, S., et al. (2016). Structural development of human fetal and preterm brain crotical
730   plate based on population-averaged templates. Cereb Cortex 26, 4381-4391.
731

ACKNOWLEDGEMENTS: This work was supported by grants from National Institute of Health:

R01MH092535, R01MH125333, R01EB031284, R21MH123930 and P50HD105354.

AUTHOR CONTRIBUTIONS: H.H. designed the research; Q.Y., M.O., H.K., D.H., Y.P. and

H.H. performed research; Q.Y. and H.H. contributed new reagents/analytic tools; Q.Y., M.O.,

J.D., B.H., F.F. and H.H. wrote the paper.

COMPETING INTERESTS: The authors declare no competing interests.

 Figures and figure supplements

740
741

Figure 1. Acquisition of high-quality infant pseudo-continuous arterial-spin-labeled (pCASL)

perfusion and phase contrast (PC) MRI and resultant axial regional cerebral blood flow (rCBF)

maps at different infant ages. **a** Labeling plane (red line) and imaging volume (blue box) of

pseudo-continuous arterial-spin-labeled perfusion MR imaging are shown on the mid-sagittal slice

of T1-weighted image of a representative infant on the left panels. Axial and sagittal view of MR

angiography with reconstructed internal carotid and vertebral arteries are shown in the middle of

the panel a. On the right of the panel a, the coronal view of the reconstructed arteries is placed in

the middle with four slices (shown as blue bars) of the PC MR scans positioned perpendicular to

the respective feeding arteries. The PC MR images are shown on the four panels surrounding the

coronal view of the angiography. These PC MR images measure the global cerebral blood flow

of internal carotid and vertebral arteries and are used to calibrate rCBF. **b** rCBF maps of

representative typically developing (TD) infant brains at 1, 6, 12, 18 and 24 months from left to

right. Axial slices of rCBF maps from inferior to superior are shown from bottom to top of the panel

b for each TD infant brain. Green arrows point to the posterior cingulate cortex (a hub of the DMN

network) characterized by relatively lower rCBF at early infancy and prominent rCBF increases

from 1 to 24 months. Blue arrows point to the visual cortex characterized by relatively higher rCBF

at early infancy and relatively mild rCBF increase from 1 to 24 months.

Figure 1-figure supplement 1. Highly reproducible pCASL protocol adopted in the present study

for measuring rCBF.

Figure 1-figure supplement 2. Identification of functional network regions-of-interests with

resting-state fMRI.

Figure 1-figure supplement 3. Heterogeneity of rCBF measurements across functional network

regions.

Figure 1-figure supplement 1. Highly reproducible pCASL protocol adopted in the present study

for measuring rCBF. Panel (a) shows reproducible rCBF maps measured from first and second

half of the control/label volumes, and panel (b) shows the intraclass correlation coefficient (ICC)

0.8854 with the 95% confident intervals of [0.88, 0.8906] calculated from entire brain rCBF maps

measured from first and second half control/label volumes. Both panels (a) and (b) are generated

based on pCASL scans of a randomly selected infant subject aged 17.6 months.

Figure 1-figure supplement 2. Identification of functional network regions-of-interests with

resting-state fMRI. (a) Functional connectivity obtained from correlation between blood

oxygenation level dependent (BOLD) fMRI time courses of two voxels; (b) Brain regions of the

visual (Vis), sensorimotor (SM) and default-mode network (DMN) identified by z statistic maps

from group independent component analysis. DMN subregions posterior cingulate cortex (PCC),

medial prefrontal cortex (MPFC), inferior posterior lobule (IPL) and lateral temporal cortex (LTC)

can be clearly appreciated.

Figure 1-figure supplement 3. Heterogeneity of rCBF measurements across functional network

regions. Significant differences ($p < 0.05$ with false discovery rate (FDR) correction) of rCBF

measurements between regions were indicated by asterisks.

788
789

790
791

Figure 2. 4D spatiotemporal regional cerebral blood flow (rCBF) dynamics and faster rCBF increases in the default-mode network (DMN) hub regions during infancy. **a** Medial (top row) and lateral (bottom row) views of fitted rCBF profiles of the infant brain at 0, 6, 12, 18 and 24 months in the custom-made infant template space demonstrate heterogeneous rCBF increase across the brain regions. **b** Medial (top) and lateral (bottom) views of rCBF-age correlation coefficient (r) map are demonstrated. **c** The scatterplots of rCBF measurements in the primary sensorimotor (SM) network (black circle and black line), visual (Vis) network (blue circle and blue line), and total and individual DMN hub regions (DMN_MPFC, DMN_PCC, DMN_IPL and DMN_LTC) (blue circle and blue line) demonstrate differential rCBF increase rates. * next to network name in each plot indicates significant (FDR-corrected $p<0.05$) differences of rCBF trajectory slopes from that of SM used as a reference and shown in a black dashed line. Abbreviations of DMN subregions: IPL: inferior posterior lobule; LTC: lateral temporal cortex; MPFC: medial prefrontal cortex; PCC: posterior cingulate cortex.

806
807
808

Figure 3. Emergence of functional connectivity (FC) within the default-mode network (DMN) during infancy. The maps of the DMN FC (posterior cingulate cortex (PCC) as a seed) at representative ages from 0 to 24 months are demonstrated in **a** and the map of correlation coefficient of FC (PCC as a seed) and age is demonstrated in **b**. In **a**, gradually emerging FC of other DMN regions (including MPFC, IPL and LTC) to the PCC from 0 to 24 months can be appreciated. The PCC is delineated by the black dashed contour. In **b**, stronger correlation between FC (PCC as a seed) and age is localized in DMN subregions IPL, ITL and MPFC. See legend of Figure 2 for abbreviations of the DMN subregions.

Figure 3-figure supplement 1. Age-dependent changes of functional connectivity (FC) within the default-mode network (DMN), visual (Vis) and sensorimotor (SM) network regions during infancy.

821 **Figure 3-figure supplement 1**. Age-dependent changes of functional connectivity (FC) within the

822 default-mode network (DMN), visual (Vis) and sensorimotor (SM) network regions during infancy.

823 Significant age-dependent increase of within-network FC was found only in the DMN ($r = 0.31$, p

824 < 0.05, indicated with an asterisk) regions, but not in the Vis ($r = 0.048$, $p = 0.745$) and SM ($r =$

825 0.087, $p = 0.559$) regions.

Figure 4. Significant correlation of regional cerebral blood flow (rCBF) and functional connectivity

(FC) at randomly selected 4000 voxels within the default-mode network (DMN) for both infants

aged 0-12 months ($p<0.001$, left scatter plot) and infants aged 12-24 months ($p<0.001$, right

scatter plot). FC is the average of FC of a certain DMN voxel to all other DMN voxels. The DMN

regions of interests obtained from a data-driven independent component analysis of resting-state

fMRI of the 12-24-month infant cohort are shown on the top panels as an anatomical reference.

See legend of Figure 2 for abbreviations of the DMN subregions.

837
838

839
840
Figure 5. Significant correlation between functional emergence of the default-mode network

(DMN) and regional cerebral blood flow (rCBF) increases specifically in the DMN regions, but not

in primary sensorimotor (visual or sensorimotor) regions. **a** Correlation of intra-default-mode-

network functional connectivity (DMN FC) and regional cerebral blood flow (rCBF) at randomly

selected voxels in the DMN (light red lines), visual (Vis, light green lines) and sensorimotor (SM,

light blue lines) network regions. Correlations of DMN FC and averaged rCBF in the DMN, Vis

and SM network regions are shown as thickened red, green and blue lines, respectively. **b**

Coupling between the DMN FC and rCBF across the brain can be appreciated by distribution of

voxel-wise correlation coefficient (r) obtained from correlation between DMN FC and rCBF at each

voxel. The short black line in the color bar indicates critical r value r_{crit} corresponding to p=0.05.

Higher r values can be appreciated in the DMN hub regions including posterior cingulate cortex

(PCC), medial prefrontal cortex (MPFC), inferior posterior lobule (IPL) and lateral temporal cortex

(LTC) with their boundaries delineated by the dashed dark red contours (from Figure 1-figure

supplement 2b). Dashed green and blue contours (also from Figure 1-figure supplement 2b)

delineate the Vis and SM network regions, respectively. **c** Radar plot shows significant correlation

between rCBF and intra-DMN FC in the DMN network (36.7%, p < 0.0001), but not in the Vis

(14.6%, p > 0.05) or SM (5.5%, p > 0.05) networks. The radius represents the percent of the

voxels with significant correlations between intra-DMN FC and rCBF in DMN, Vis and SM network

regions, respectively. The dashed line circle indicates critical percent of significant voxels with

p=0.05 from 10,000 permutation tests. *** indicates p<0.0001.

Figure 5-figure supplement 1. Procedures of generating map of correlation between the default-

mode network (DMN) functional connectivity (FC) and reginal cerebral blood flow (rCBF) across

the entire brain voxels.

Figure 5-figure supplement 2. Coupling between intra-visual network functional connectivity (Vis

FC) and regional cerebral blood flow (rCBF) and coupling between intra-sensorimotor network

(SM FC) and rCBF.

Figure 5-figure supplement 1. Procedures of generating map of correlation between the default-mode network (DMN) functional connectivity (FC) and reginal cerebral blood flow (rCBF) across the entire brain voxels. At any brain voxel indicated by a small black box in the correlation map below, correlation between the DMN FC across subjects and the rCBF at this voxel across subjects was conducted and the correlation coefficient was calculated for this voxel. The correlation map between the DMN FC and rCBF across the brain can be generated by projecting the correlation coefficient r of each voxel onto the cortical surface.

Figure 5-figure supplement 2. Coupling between intra-visual network functional connectivity

(Vis FC) and regional cerebral blood flow (rCBF) and coupling between intra-sensorimotor

network (SM FC) and rCBF. Left panels show the correlation of Vis FC (a) or SM FC (b) and

rCBF at randomly selected voxels in the default-mode (DMN) (light red lines), Vis (light green

lines) and SM (light blue lines) network regions. Correlations of Vis FC (a) or SM FC (b) and

averaged rCBF in the DMN, Vis and SM network regions are shown as thickened red, green

and blue lines, respectively. Middle panels demonstrate the distribution of voxel-wise correlation

coefficient (r) obtained from correlation between Vis FC (a) or SM FC (b) and rCBF at each

voxel. The short black line in the color bar indicates critical correlation coefficient r value r_{crit}

corresponding to p=0.05. Low r values are apparent in all network (DMN, Vis and SM) regions

with dashed red, green and blue contours (from Supplementary Fig. 2b) delineating DMN, Vis

and SM network, respectively. Right panels show radar plot with radius representing percent of

voxels with significant correlation between Vis FC or SM FC and rCBF in the DMN, Vis or SM

network regions. The radii in radar plots demonstrate that the percent of voxels with significant

correlations between intra-Vis (a) or intra-SM (b) FC and rCBF in DMN, Vis or SM network

regions is 0 or close to 0, indicating that coupling between Vis FC or SM FC and rCBF in any

network region is not significant. The dashed line circle in the radar plot indicates critical percent

of significant voxels with p=0.05 from 10,000 permutation tests.

898
899

Figure 6. A diagram illustrating hypothesized neuronal mechanism supporting coupling of regional cerebral blood flow (rCBF) and functional connectivity. Specifically, higher cerebral blood flow delivers larger amount of oxygen and glucose to neurons, leading to cascade of events of regional cerebral metabolic rate of oxygen (CMRO2) ↑ and glucose (CMRGlu) ↑, synaptogenesis ↑ and synaptic efficacy ↑, blood oxygenation level dependent (BOLD) signal synchronization ↑, and functional connectivity ↑, during infant brain development.

907   **Video 1:** Video of the 4D spatiotemporal whole-brain dynamics of regional cerebral blood flow

908   from 0 to 24 months.