Modeling features of addiction with an oral oxycodone self-administration paradigm

Caitlin Murphy[1], Yu-Hsuan Chang[1], Rajesh Pareta[1], Jun-Nan Li[1], Tom Earnest[1], Jessica Tooley[1], Yvan M. Vachez[1], Robert W. Gereau IV[1,,3,4], Bryan A. Copits[1,3], Alexxai V. Kravitz[1,2,3,4], Meaghan C. Creed[1,2,3,4]

1. Washington University Pain Center, Dept. of Anesthesiology, Washington University in St. Louis
2. Department of Psychiatry, Washington University in St. Louis
3. Department of Neuroscience, Washington University in St. Louis
4. Department of Biomedical Engineering, Washington University in St. Louis

Abstract

Prescription opioid use is an initiating factor driving the current opioid epidemic. There are several challenges with modeling prescription opioid addiction. First, prescription opioids such as oxycodone are orally self-administered and have different pharmacokinetics and dynamics than morphine or fentanyl. This oral route of administration determines the pharmacokinetic profile, which is critical for establishing reliable drug-reinforcement associations in animals. Second, intravenous opioid self-administration is typically performed with intermittent drug self-administration sessions in a separate environment from the home cage. This does not recapitulate prescription opioid use, which is characterized by continuous drug access in the patients' homes. To model features of prescription opioid use and the transition to abuse, we developed an oxycodone self-administration paradigm that is administered in the homecage. Mice voluntarily self-administer oxycodone in this paradigm without any taste modification such as sweeteners, and exhibit preference for oxycodone, escalation of intake, physical signs of dependence, reinstatement of seeking after withdrawal, and a subset of animals demonstrate drug taking that is resistant to negative consequences. This model is therefore translationally relevant and can be useful for studying the neurobiological substrates specifically relevant to prescription opioid abuse.

Introduction

Misuse of prescription opioids is a significant public health and economic burden worldwide. In 2017 in the United States, more than 10 million people reported prescription opioid misuse [1]. Despite its importance, modeling prescription opioid abuse in rodents has been challenging [2-5]. Certain aspects unique to prescription opioid use, such as route of administration and pattern of intake, may contribute to abuse liability and propensity toward relapse. As such, a representative preclinical model of prescription opioid addiction is critical for understanding the neurobiology underlying this disorder.

Pre-clinical drug self-administration studies have typically employed paradigms where drug administration is limited to discrete time periods in operant chambers, before returning animals to their homecages. In these studies, drugs are often self-administered intravenously under various reinforcement schedules, which allows experimenters to assay distinct features of addiction behavior, such as motivation for the drug [6,7], reinstatement of drug seeking [8,9] or compulsive drug taking [10-12]. However, several characteristics of prescription opioid use introduce unique challenges that may not be optimally modelled by classical self-administration paradigms.

First, in contrast to many addictive drugs (e.g., heroin, cocaine, methamphetamine or morphine), prescription opioids are most often orally self-administered: among chronic and recreational opioid users, oral self-administration is decidedly preferred over non-oral routes (e.g., insufflation, injection) [13,14]. The route of administration of drugs of abuse is highly relevant for establishing learned drug associations [15], which contribute to the subjective reinforcing properties of the drug, escalation of intake, and propensity toward reinstatement [10,11,16-20]. Indeed, the pharmacodynamic and pharmacokinetic profiles of opioids – and therefore their physiology and time-course of action – vary widely across routes of administration [21]. These differences underscore the need to model prescription opioid abuse through the oral route of administration.

Second, prescription opioids are most commonly self-administered in a familiar setting such as the home (see Caprioli et al. (2007) for a review [22]). Yet, where oral self-administration has been reported [2-4], experimental paradigms have required self-administration to take place in a novel context (e.g., operant chamber) and under food restriction. Environmental context (i.e., setting of drug taking) plays an important role in both drug taking and reinstatement, a finding that has been well-documented in humans and recapitulated by animal models [22-27]. Moreover, the nature of operant self-administration paradigms requires intermittent access to the drug during daily sessions separated by acute periods of forced abstinence. However, this intermittent pattern of intake stands in stark contrast to continuous access to prescription opiates, which is even more pronounced in the case of extended-release formulations [28]. The pattern of drug intake is critical, since intermittent access patterns exacerbates mesolimbic adaptations induced by opioid self-administration [29,30].

Previous investigations have applied the oral route of self-administration by implementing two-bottle choice procedures, where animals are free to choose between bottles containing drug- and drug-free solutions. Because the opioid alkaloid structure confers a bitter taste to opium derivatives, and typically sweeteners have been added to encourage animals to self-administer opioid via the oral route, or adulterants to the drug-free bottle to control for aversive taste. However, the addition of these extraneous reinforcers or adulterants introduce experimental confounds which limit the interpretability of two-bottle choice procedures [31]. For example, sucrose itself can support seeking after abstinence [32], disrupt

overall patterns of fluid intake [33] and stimulate dopamine and opioid receptors in the mesolimbic pathway [34-37]. Therefore, recapitulating a continuous access pattern and oral route of administration without adulterants is important to accurately model prescription opioid use disorder (OUD).

Here, we characterize a homecage-based, oral oxycodone (Oxycontin[TM] or Percocet[TM]) self-administration paradigm that models several features of prescription opioid abuse. We demonstrate that mice exhibit a preference for high concentrations of unadulterated oxycodone (1.0 mg/mL) over regular drinking water and voluntarily escalate drug intake over the course of a two-bottle choice paradigm. This model also recapitulates multiple behavioral features that have been used to model criteria for addiction as defined by the fifth version of the diagnostic and statistics manual [40] (DSM), specifically escalation of drug intake, physical signs of dependence, drug craving after withdrawal and drug use despite negative consequences. Critically, these behaviors were also accompanied by potentiation of excitatory synapses in the nucleus accumbens, which is a conserved neurobiological substrate of persistent drug reinstatement [41-43]. By understanding the neurobiological substrates specific to prescription opioid abuse, it may be possible to design effective therapies for prescription opioid misuse, and to develop novel medications to treat pain with lower abuse liability.

Materials and Methods

Subjects. A total of 92 adult mice (C57BL/6J background; age, 10–16 weeks). Both males (n=46) and females (n=46) were included in the study and balanced across experiments and treatment groups. Mice were housed in standard mouse vivarium caging and kept on a 12 h light/dark cycle (light onset, 7:00 A.M.; light offset, 7:00 P.M., temperature (22-26°C), in-cage humidity (22-50%). Mice had ad libitum access to food and were single housed during self-administration experiments. Experiments were run in successive waves of treatment- and sex-matched cohorts with up to 20 mice being run simultaneously. All studies were approved by the internal animal care and use committee at Washington University in St. Louis.

Homecage lickometer devices. All devices used in these experiments are low-cost and open-source, with detailed parts lists, code and fabrication instructions published and available on-line [44,45]. Drinking water and oxycodone solutions were provided using an in-cage two-bottle lickometer apparatus which detects interactions with two separate drinking spouts via a pair of photobeams [45]. Cages were also equipped with a wireless passive infrared (PIR) activity monitor [44] to measure mouse activity levels (Models 4430 and 4610 from MCCI Corporation, Ithaca NY). Lickometer counts on each sipper, PIR activity data, and environmental measurements (temp, humidity, light levels) were transmitted via LoRaWAN (Low frequency Radio Wide Area Network) using IoT (internet of things) infrastructure (The Things Network, Netherlands), saved in a cloud-database (InfluxDB), and visualized with an online dashboard (Grafana). The wireless lickometer and PIR sensors were also equipped with a back-up microSD cards and liquid volumes were measured manually each day.

Oxycodone self-administration. Habituation: Mice were transferred to single housing with an in-cage two-bottle choice apparatus at least three days prior to the start of the escalation protocol, during which the sipper device was the only source of drinking water (supplied in both tubes). Phase I: Mice were supplied with oxycodone hydrochloride dissolved in drinking water as their only source of liquid; 0.1 mg/mL was available for 24 hours, 0.3 mg/mL was available for 48 hours, 0.5 mg/mL was available for 48 hours, based

on an operant protocol developed by Phillips et al. [3]. The position of the bottles within the two-bottle apparatus was switched daily. Phase II: Oxycodone (1.0 mg/mL in drinking water) and drinking water were supplied ad libitum in the two-bottle choice apparatus. Both solutions were available 24 hours a day for 7 days. Withdrawal: The two-bottle lickometer was removed from the homecage and standard cage-top water bottles were replaced as the source of drinking water.

Oxycodone Seeking Test. The two-bottle lickometer devices were removed from the homecages after 7 days of free choice between drinking water and 1.0 mg/mL oxycodone (described above). After 24 hours (early) or 22-24 days (late) withdrawal, the lickometer device containing 2 empty bottles was re-introduced to the cage for a 60-minute probe trial, and interactions with the sipper tubes as detected by the photobeam were recorded. Mice were tested at both early and late withdrawal points.

Quinine Adulteration. Following the conclusion of the 7 days of two-bottle choice protocol, increasing concentrations of quinine were added to the oxycodone-containing bottle, such that mice had the choice between standard drinking water (one bottle) and 1.0 mg/mL oxycodone + quinine. Mice were given 48 hours at each quinine concentration in increasing order: 125 µM, 250 µM, 325 µM, volumes consumed from each tube were recorded daily.

Physical Withdrawal Signs. Acquisition: After either 24 hours or 22-24 days of withdrawal, mice were habituated to the testing room for approximately 2 hours. Three consecutive 5 minute videos were acquired of individual mice at 100 fps in a double-tall Von Frey chamber with an unobstructed background and overhead lighting (LUX). Videos were scored for signs of withdrawal (jumps, tremors, movement speed) and anxiety related behaviors (grooming, rearing, climbing) using markerless pose estimation and subsequent supervised machine learning predictive classifiers of behavior. Markerless pose estimation was done using DeepLabCut^{TM} (DLC, version 2.2b6 [46,47]): 12 frames of 64 videos representing all treatment conditions were extracted using k-means clustering and subsequently manually annotated for the following 9 body parts: left ear, right ear, left forepaw, right forepaw, left hindpaw, right hindpaw, snout, tail base, back. The training fraction was set to 0.95, and the resnet_50 network was trained for 1,030,000 iterations. A train error of 1.82 and test error of 11.47 were achieved with a cutoff value of p=0.6. DeepLabCut tracking data and generated videos were then imported to the Simple Behavioral Analysis (SimBA) project workflow (version 1.2 [48]). Within SimBA, single animal, 9-body part supplied configuration was used to extract behavioral features from the pose estimation data after outlier and movement correction (both parameters set to 7x outside the interaural distance. Extracted frames from four independent videos were annotated to build classifiers for the following behaviors: "climbing", "jumping", "tremor", "rearing, and "grooming". Individual models were trained using a random forest machine model with 2000 estimators, and a training fraction of 0.2 (default hyperparameters). Following validation, videos were analyzed using the random forest model, with the following p-cuttoffs and minimum behavioral bout lengths for each of the following behaviors: "climbing (p = 0.26, 35 ms)", "jumping (p = 0.4, 35 ms)", "tremor (p=0.0495, 50 ms", "rearing (p = 0.45, 70 ms)", and "grooming (p = 0.38, 70 ms)". The SimBA pipeline is built primarily on scikit-learn [49], OpenCV [50], FFmpeg45 [51], and imblearn [52]. Total time spent in each behavior was exported and the three videos were averaged to get one value for each subject to be included in the analysis. Euclidean distance of displacement of all body parts was also extracted from the pose-estimation data and averaged to achieve one 'movement' score for each subject.

Patch Clamp Electrophysiology. Coronal mouse brain slices, 220 μm in thickness were prepared in cooled artificial cerebrospinal fluid containing (in mM): NaCl 119, KCl 2.5, MgCl 1.3, CaCl2 2.5, Na2HPO4 1.0, NaHCO3 26.2 and glucose 11, bubbled with 95% O2 and 5% CO2. Slices were kept at 30- 32°C in a recording chamber superfused with 2.5 ml/min artificial cerebrospinal fluid. Visualized whole-cell voltage-clamp recording techniques were used to measure holding and synaptic responses of NAc shell MSNs. Holding potential was maintained at -70 mV, and access resistance was monitored by a depolarizing step of -10 mV each sweep, every 10 s. Currents were amplified, filtered at 5 kHz and digitized at 50 kHz. All experiments were performed in the presence of picrotoxin (100 μM) to isolate excitatory transmission, TTX (0.5 μM) was included for recordings of miniature EPSCs (mEPSCs). Internal solution contained (in mM) 130 CsCl, 4 NaCl, 5 creatine phosphate, 2 $MgCl_2$, 2 Na_2ATP, 0.6 Na_3 GTP, 1.1 EGTA, 5 HEPES and 0.1 mm spermine. Synaptic currents were electrically evoked by stimuli (50 – 100 μs) at 0.1 Hz through bipolar stainless-steel electrode placed onto the tissue. The AMPAR component was calculated as the peak amplitude at -70 mV, PPR was calculated as the ratio of the second to first AMPAR peak elicited with an ISI of 50 msec. The NMDAR component was estimated as the amplitude of the outward current at +40 mV after decay of the AMPA current, measured 50 msec after the electrical stimuli (Fig 6C).

Results

Mice exhibit preference, escalation and altered pattern of intake of oxycodone.

 We mimicked a prescribed course of opioids by including escalating concentrations of oxycodone in the drinking water in a single-bottle phase, followed by a two-bottle choice phase where mice could choose between a concentrated oxycodone solution (1.0 mg/mL) and unadulterated drinking water (Fig 1A). 1.0 mg/mL oxycodone solution is bitter, and both male and female naïve mice preferred drinking water when given both options in a two-bottle choice paradigm (Fig S1). However, after the 5 days of forced exposure to escalating doses of oxycodone in the drinking water, mice exhibited a significant preference for 1.0 mg/mL oxycodone over drinking water (Fig 1b-c). Mice increased their daily oxycodone during the single bottle choice phase, which was expected based on the increasing concentrations of oxycodone in their drinking water (Fig 1G). However, mice continued to systematically escalate their drug intake during the two-bottle choice phase, where the drug concentration remained constant (1.0 mg/mL) and drinking water was also available (Fig 1G, H).

The home-cage nature of our paradigm allows for large sample sizes with statistical power to detect sex differences (n=40 control and 52 oxycodone animals in total). While both male (n=26) and female mice (n=26) exhibited preference for oxycodone (Fig 1b-c, S1c-d), during the single bottle phase, female mice drank significantly higher volumes of oxy-containing drinking water relative to males (Fig 1E,F). Female mice continued to self-administered higher doses of oxycodone over the two-bottle phase of the protocol (Fig 1H, S1B, E, F). During the two-bottle phase, both male and female mice given the opportunity to consume oxycodone had higher levels of fluid intake than control mice. One DSM criteria for addiction is the escalation of drug consumption. Mice increased their daily oxycodone during the single bottle choice phase, which was expected based on the increasing concentrations of oxycodone in their drinking water (Fig 1G). More importantly, mice continued to systematically escalate their drug intake when the drug concentration remained constant (1.0 mg/mL) during the two-bottle choice phase, when drinking water was also freely available (Fig 1G, H). While this escalation index (oxy dose consumed during the last day

of the two-bottle choice / oxy dose consumed during the first day of two-bottle choice) did not differ between males and females

Oxycodone self-administration induces physical signs of intoxication and withdrawal.

DSM criteria for substance use disorder include tolerance to and withdrawal from to the addictive drug. We did not measure drug tolerance, although the oral self-administration protocol did induce physical signs of intoxication (hyperactivity) as well as acute and protracted physical withdrawal symptoms. Acute injections of oxycodone [53-55], or vapor exposure to opioids induce hyperactivity [56], these locomotor-activating properties of addictive drugs are thought to reflect increased dopamine release in the nucleus accumbens [57-59]. We equipped our homecages with passive infrared sensors to detect locomotor activity during the self-administration protocol (Fig 3A). We confirmed that oral intake of oxycodone induced hyperactivity in both the single-bottle phase and the two-bottle choice phases of self-administration. There were no sex differences in total activity levels; with both male and female oxycodone self-administering mice exhibiting higher locomotor activity than control mice (Fig 3A,B). This confirms that levels of oxycodone voluntarily self-administered via the oral route produce sufficient brain levels of oxycodone to induce hyperactivity [54,60,61].

We also determined whether withdrawal from oral oxycodone self-administration induced physical withdrawal signs. During withdrawal from opioids such as morphine or heroine, mice exhibit characteristic withdrawal signs including tremors, jumping, teeth chattering and diarrhea. While we were not able to reliably detect wet-dog shakes or teeth chattering in our videos, we quantified jumping and tremor, along with the total distance moved by the animal. Oxycodone self-administering mice exhibited significantly more movement, as well as increased jumping (Fig 3E) and paw tremors (Fig 3F) at both the acute (1 day) and protracted (3 weeks) withdrawal periods.

Finally, we quantified climbing (an index of escape behavior), grooming, and rearing, which have been used as a proxy for anxiety-like behavior, and have also been reported to be increased following withdrawal from opioids. Mice that had self-administered oxycodone did not exhibit significantly more rearing or grooming relative to control mice (Fig 3G,H). While there was a significant effect of drug on escape behaviors in acute oxycodone withdrawal, this effect was driven by two subjects and resulting group differences were small (Fig 3I). There were no significant differences between male and female mice on any physical symptom measured. Together, these results demonstrate that oral oxycodone self-administration induces physical signs of dependence which are specific to opioid withdrawal syndrome, and that symptoms are not related to generalized anxiety-like behavior.

The two-bottle choice paradigm induces aversion-resistant drug consumption and persistent drug seeking into withdrawal.

Drug seeking behavior can be operationally defined as performance of an action that previously led to consumption of the drug, in the absence of the drug itself [62,63]. In the case of intravenous drug self-administration, this response may be operationalized as lever-pressing or nose-poking to receive a drug infusion. In our model, oxycodone is delivered through a unique sipper device, which is distinct from the standard cage-top water bottle. Therefore, we assayed drug seeking behavior by measuring photobeam

breaks on the lickometer devices under extinction conditions, with no oxycodone or drinking water in the tubes after 24 hours and 3 weeks of forced abstinence from oxycodone (Fig 4A). Mice that had self-administered oxycodone exhibited significantly higher lickometer counts than control mice in both the acute (1 day) and protracted (3 weeks) withdrawal periods (Fig 4B). This drug seeking behavior was not significantly different between the early and the late probe trials, nor between males and females (Fig 4C-H, Fig S4A-B). This establishes that the two-bottle choice paradigm supports drug seeking behavior that persists after withdrawal.

Our final behavioral test sought to determine whether mice will consume oxycodone despite negative consequences. Negative consequences are frequently modeled by a physical punishment such as footshock [64-66], or adulteration with aversive tastants, such as quinine [67,68]. Given the analgesic properties of oxycodone that could confound responses to footshock punishment, we introduced increasing concentrations of quinine to the oxycodone-containing drinking tube, such that concentration increased by 125 µM every 48 hours (Fig 5A). Both male and female control mice who were presented with standard drinking water in both tubes avoided the quinine-containing bottle at the lowest concentration (Fig 5B). By contrast, mice persisted in their oxycodone drinking when adulterated with the lowest concentration of quinine, and a subset of mice (2/15, or 15%) persisted in oxycodone drinking at the highest concentration tested (Fig 5C, D). There were no sex differences in the AUC of the quinine preference curve, with both male and female mice showing higher consumption of oxycodone despite quinine adulteration relative to control mice (Fig 5E).

Oxycodone self-administration is characterized by enhanced excitatory transmission in the nucleus accumbens.

Addiction is a chronic, relapsing disorder, with behavioral symptoms persisting long into abstinence in humans and rodents [69,70]. The chronic nature of these symptoms is mediated by long-lasting plasticity in the mesolimbic dopamine system, which is induced by exposure to drugs of abuse. One characteristic form of this plasticity that has been causally linked to drug sensitization and persistent relapse is potentiation of excitatory input onto medium sized-spiny neurons in the nucleus accumbens (MSNs) [71-75]. This potentiation is mediated by the insertion of AMPA receptors into the post-synaptic membrane of accumbal MSNs, is dopamine-dependent, and has been observed after self-administration of opioids [41,76-79] and psychostimulants [72,75,80]. Here, we performed whole cell recordings of accumbal MSNs throughout the nucleus accumbens shell after withdrawal from oxycodone self-administration (Fig 6A,B).

The AMPA to NMDA ratio was significantly greater in mice who underwent oxycodone self-administration and withdrawal, relative to control mice, suggesting increased insertion of AMPA receptors (Fig 6D). There were no differences in the PPR between oxycodone self-administering mice and control mice, arguing that pre-synaptic excitatory transmission onto accumbal MSNs is not altered by oxycodone self-administration and withdrawal (Fig 6E). To bolster this interpretation, we recorded miniature excitatory post-synaptic currents (mEPSCs) in accumbal MSNs. The average amplitude of mEPSCs was significantly greater in oxycodone self-administering mice relative to control mice, and there was no difference in the mEPSC event frequency (Fig 6F,G). Together, these results confirm that self-administration and withdrawal from oxycodone leads to a post-synaptic potentiation of excitatory synapses onto NAc MSNs, consistent with hallmark plasticity that is induced by self-administration and withdrawal from other opioids and psychostimulants.

Discussion

Much of our current understanding of addiction has relied on animal models to understand the relationship between behavioral phenomena and their neurobiological correlates. Environmental factors, route of administration, and pattern of intake strongly influence nearly every aspect of drug misuse, yet previous models of addiction have historically been conducted under experimental parameters which differ significantly from typical conditions associated with prescription opioid misuse. For example, intermittent access to intravenous drug infusions with discrete cues in a novel environment are ideal for modeling addiction to psychostimulants or heroine. However, prescription opioids, including oxycodone, are typically administered via the oral route, with near continuous access (especially with extended-release formulations) in familiar environments. Here, we characterize a novel paradigm that more closely recapitulates factors relevant to prescription opioid abuse, which may prove valuable for studying the neurobiology of opioid use disorder in mice.

In this oral oxycodone self-administration paradigm, female mice consumed more oxycodone (mg/day) during the single bottle phase and higher total dose of oxycodone (mg/kg) compared to males (Figure 1), consistent with previous observations that female rodents self-administer higher doses of opioids than males [3,5,81-84]. The effects of sex on opioid intake, dependence, and abuse liability have been well documented [61,81,85-88], and interact with feeding conditions [89,90] and route of administration [3,38] to influence the bioavailability of oxycodone. While pharmacokinetic influences on our observed sex differences were beyond the scope of our study, others have explored sex-dependent factors that may contribute to different oral oxycodone consumption between females and males. Phillips et al. (2018) [3] demonstrate that while oral oxycodone ubiquitously resulted in elevated plasma concentrations of oxycodone and its active metabolite oxymorphone, the extent of this effect was influenced by sex, presence of sex organs, and feeding conditions, and was exacerbated following repeated administrations. Such observations are congruent with pharmacokinetic studies showing that plasma concentrations of oxycodone and its active metabolites are mediated, in part, by CYP3A4 and CYP2D6 [91,92], whose activity can be modulated by sex hormones and food intake [93,94]. Importantly, however, sex effects on oxycodone pharmacokinetics were either diminished or not observed with i.v. administration [38,95], highlighting the importance of route of administration in addiction models. To this end, it is critical that the paradigm we present achieves preference for oral consumption of oxycodone over drinking water without need for addition of sweeteners, or for adulteration of the drinking water with bitter tastants, which can influence measures of preference and drug intake.

Moreover, mice voluntarily escalate their consumption of oxycodone despite ad libitum access to drinking water (Fig 1, Fig S1), and alter their pattern of intake, both with respect to circadian cycle and inter-bout interval (Fig 2, Fig S2). This combination of drug escalation and irregularity of pattern of drug intake has been suggested to reflect 'loss of control' of drug intake, which is relevant to modeling DSM criteria for addiction [11,96,97]. Additionally, we found that this voluntary oxycodone consumption induced robust physical withdrawal signs, including hyperactivity, jumps and paw tremors which persisted to a lesser extent into protracted withdrawal (Fig 3). This is critical, because withdrawal symptoms can contribute to drug relapse through negative reinforcement (where the drug is taken to relieve the aversive state of withdrawal [98,99]. Withdrawal signs are often modelled with non-contingent drug administration and precipitation with an opioid antagonist. Our results are sufficient, as they establish that sufficient oxycodone concentrations are achieved via voluntary oral self-administration to produce physical

dependence. There was large variability of withdrawal signs, with only a subset of mice exhibiting physical symptoms into protracted withdrawal.

Drug craving is another cardinal DSM criterion for opioid use disorder (OUD) [40], which is frequently modeled with reinstatement paradigms. Typically, an instrumental response is acquired which is paired with drug infusion in self-administration paradigms. Then, following a period of extinction training, the ability of stress, drug-exposure, discriminative or contextual cues to reinstate the instrumental response in the absence of drug is measured as an index of drug craving [10,20,101]. However, this extinction training is not typically a feature of prescription drug use. Rather, abstinence is typically imposed by the cessation of opioid availability, either because of prescription termination, abrupt changes in prescribing practices or hospitalization [102]. For this reason, we opted to use a 'forced abstinence' model, where no extinction training occurs after drug self-administration, and the response that previously lead to drug consumption is measured after variable periods of withdrawal [62,103]. We found that lickometer interactions, which resulted in the delivery of oxycodone solution during the self-administration phase, were significantly greater in oxycodone-self-administering mice relative to controls at both acute (24h) and late (3 week) withdrawal phases. Interestingly, we did not find evidence for the incubation of craving in this paradigm, which refers to the tendency for craving-related behavior to increase with the length of the forced abstinence period [62]. We propose three possible explanations for this: incubation of craving is most robustly observed under long access, high concentrations of IV administration, where initial seeking responses on the first probe session are typically low [72]. This is not the case with our oral administration, where seeking behavior is quite high, even during a short assay in the acute withdrawal. Additionally, incubation of craving is frequently measured when mice are tested either in acute or late withdrawal, and behavior is compared between different subjects, with late withdrawal subjects showing increased seeking [75,104]. Here, we are testing the same subjects at two time points, and while we don't see differences between these time points, it is possible that the first session served as an extinction trial, which may mask any potential effects on incubation [105]. Finally, the time course of incubation of craving is highly dependent on the addictive substance [62], and the single three-week time point may not be adequate to capture the temporal dynamics of oxycodone seeking after withdrawal.

We modeled drug consumption despite negative consequences by adulterating oxycodone solution with increasing doses of quinine, which is commonly used to assess compulsive alcohol drinking [106,107]. While the quinine concentrations were quite low relative to ethanol studies, oxycodone is an alkaloid with a bitter taste prior to the introduction of quinine and already aversive, which we confirmed by establishing that naïve mice avoid 1.0 mg/mL oxycodone before the introduction of quinine (Fig S1A). Critically, we also observed individual differences in quinine-resistant drug consumption, with only a subset of mice persisting at the highest quinine doses (Fig 5). This is reminiscent of intravenous opioid or psychostimulant self-administration, where typically only a subpopulation (15-30%) of animals exhibit punishment-resistant drug self-administration [65,108-110]. This is critical as it also reflects human patients who are often able to use drugs recreationally, with only a subpopulation transitioning to compulsive drug use [109,111], and underscores the utility of this model for studying individual differences in aversion-resistant oxycodone consumption.

Finally, we demonstrate that oral oxycodone self-administration induces potentiation of excitatory transmission into NAc MSNs (Fig 6), which has been linked to persistence of reinstatement and physical symptoms following withdrawal from self-administration of cocaine [42,75,112] and opioids [113-117]. This plasticity is induced by drug-induced dopamine release from the ventral tegmental area (VTA) into

the nucleus accumbens [42,43,118], and can be mimicked by strong optogenetic stimulation of VTA DA neurons alone [65]. Relative to morphine, oxycodone causes even more robust release of dopamine into the NAc, presumably through mu-receptor activation of inhibitory inputs to VTA dopamine neurons [119]. Consistent with this mechanism, we also found that oxycodone self-administering mice exhibited an increase in AMPA/NMDA ratio in accumbal MSNs, which was accompanied by an increased amplitude of mEPSCs. Combined with a lack of effect on the paired pulse ratio or frequency of mEPSCs, these suggest a post-synaptic strengthening of excitatory transmission onto accumbal MSNs. Critically, we used electrical stimulation in wildtype mice, and have no cell-type specificity within the accumbens, or projection-specificity of the pathway. However, our results suggest a bimodal distribution of AMPA/NMDA ratio, raising the possibility that adaptations may be cell-type specific. Moreover, the paradigm of drug access differentially affects plasticity at discrete excitatory inputs to the NAc, with inputs from the paraventricular thalamus implicated in physical signs of withdrawal [77,78], mPFC inputs implicated in drug seeking, sensitization and reinstatement [41,72,79,120,121], and hippocampal inputs implicated in context recognition necessary for behavioral expression of drug seeking [73]. While these results demonstrate the utility of this oral paradigm for studying synaptic mechanisms underlying persistent drug-adaptive behavior into withdrawal, future work will be needed to determine how pathway- and cell-type specific plasticity contribute to discrete elements of addiction-relevant behavior in this paradigm.

Here we characterize a novel paradigm for study prescription opioid misuse. This oral, homecage two-bottle choice procedure can be used to model several features of DSM criteria that are specifically relevant to OUD, while conserving the route of administration, pattern of access and contextual factors relevant to prescription opioid intake. This model has potential utility for studying the neurobiological substrates driving drug relapse as well as aversion-resistant drug intake. Moreover, the high throughput nature of this paradigm permits high powered studies necessary for understanding sex differences, circadian biology and individual variability underlying vulnerability to OUD.

Acknowledgements

We thank M.C. Stander and E. Godynyuk for excellent technical help. We also J.J. Choong, S. Golden and M. Mathis for their commitment to open source tools and for support with supervised behavioral classification (SimBA) and markerless pose estimation (DeepLabCut[TM]), respectively. This work was supported by Brain and Behavior Research Foundation (NARSAD Young Investigator Grant 27197 to M.C.C), National Institutes of Health National Institute on Drug Abuse (R21-DA047127, R01-DA049924 to M.C.C), Whitehall Foundation Grant (2017-12-54 to M.C.C.) and Rita Allen Scholar Award in Pain (to M.C.C.). We thank MCCI Corporation (Ithaca NY) for development of wireless in-cage activity and environmental sensors and setting up the IoT cloud solution.

Author Contributions

CM, YHC, RP and YMV performed behavioral experiments. JL, JRT, BC and MCC performed patch-clamp experiments. AVK and MCC developed homecage-based lickometer devices for self-administration, TE developed custom software for analysis. CM, BC and MCC analyzed data. RG, BC, AVK and MCC supervised work, CM and MCC designed experiments and wrote the manuscript with input from all authors.

References

Prevention CfDCa.   (Centers for Disease Control and Prevention, U.S. Department of Health and Human Services, 2019).

Fulenwider HD, Nennig SE, Hafeez H, Price ME, Baruffaldi F, Pravetoni M, et al. Sex differences in oral oxycodone self-administration and stress-primed reinstatement in rats. Addict Biol. 2020;25(6):e12822.

Phillips AG, McGovern DJ, Lee S, Ro K, Huynh DT, Elvig SK, et al. Oral prescription opioid-seeking behavior in male and female mice. Addiction Biology. 2020;25(6).

Enga RM, Jackson A, Damaj MI, Beardsley PM. Oxycodone physical dependence and its oral self-administration in C57BL/6J mice. Eur J Pharmacol. 2016;789:75-80.

Zanni G, DeSalle MJ, Deutsch HM, Barr GA, Eisch AJ. Female and male rats readily consume and prefer oxycodone to water in a chronic, continuous access, two-bottle oral voluntary paradigm. Neuropharmacology. 2020;167:107978.

Hodos W. Progressive ratio as a measure of reward strength. Science (New York, NY). 1961;134(3483):943-44.

Roberts DCS, Richardson NR. Self -Administration of Psychomotor Stimulants Using Progressive Ratio Schedules of Reinforcement. Humana Press; 1992. p. 233-70.

Davis WM, Smith SG. Role of conditioned reinforcers in the initiation, maintenance and extinction of drug-seeking behavior. Pavlov J Biol Sci. 1976;11(4):222-36.

de Wit H, Stewart J. Drug reinstatement of heroin-reinforced responding in the rat. Psychopharmacology. 1983;79(1):29-31.

Venniro M, Caprioli D, Shaham Y. Animal models of drug relapse and craving: From drug priming-induced reinstatement to incubation of craving after voluntary abstinence. Prog Brain Res. 2016;224:25-52.

Venniro M, Banks ML, Heilig M, Epstein DH, Shaham Y. Improving translation of animal models of addiction and relapse by reverse translation. Nature Reviews Neuroscience. 2020;21(11):625-43.

Shalev U, Grimm JW, Shaham Y. Neurobiology of Relapse to Heroin and Cocaine Seeking: A Review. Pharmacological Reviews. 2002;54(1):1.

Gasior M, Bond M, Malamut R. Routes of abuse of prescription opioid analgesics: a review and assessment of the potential impact of abuse-deterrent formulations. Postgrad Med. 2016;128(1):85-96.

Kirsh K, Peppin J, Coleman J. Characterization of Prescription Opioid Abuse in the United States: Focus on Route of Administration. Journal of Pain & Palliative Care Pharmacotherapy. 2012;26(4):348-61.

Macenski MJ, Meisch RA. Oral Drug Reinforcement Studies with Laboratory Animals: Applications and Implications for Understanding Drug-Reinforced Behavior. Current Directions in Psychological Science. 1994;3(1):22-27.

Panlilio LV, Weiss SJ, Schindler CW. Effects of compounding drug-related stimuli: escalation of heroin self-administration. J Exp Anal Behav. 2000;73(2):211-24.

Perry CJ, Zbukvic I, Kim JH, Lawrence AJ. Role of cues and contexts on drug-seeking behaviour. Br J Pharmacol. 2014;171(20):4636-72.

Wakabayashi KT, Weiss MJ, Pickup KN, Robinson TE. Rats Markedly Escalate Their Intake and Show a Persistent Susceptibility to Reinstatement Only When Cocaine Is Injected Rapidly. The Journal of Neuroscience. 2010;30(34):11346.

Meisch RA, Carroll ME. Oral Drug Self-Administration: Drugs as Reinforcers. In: Bozarth MA, editor Methods of Assessing the Reinforcing Properties of Abused Drugs. New York, NY: Springer New York; 1987. p. 143-60.

Reiner DJ, Fredriksson I, Lofaro OM, Bossert JM, Shaham Y. Relapse to opioid seeking in rat models: behavior, pharmacology and circuits. Neuropsychopharmacology. 2019;44(3):465-77.

Leow KP, Smith MT, Watt JA, Williams BE, Cramond T. Comparative oxycodone pharmacokinetics in humans after intravenous, oral, and rectal administration. Ther Drug Monit. 1992;14(6):479-84.

Caprioli D, Celentano M, Paolone G, Badiani A. Modeling the role of environment in addiction. Prog Neuropsychopharmacol Biol Psychiatry. 2007;31(8):1639-53.

Badiani A, Belin D, Epstein D, Calu D, Shaham Y. Opiate versus psychostimulant addiction: the differences do matter. Nat Rev Neurosci. 2011;12(11):685-700.

Badiani A, Spagnolo PA. Role of Environmental Factors in Cocaine Addiction. Current Pharmaceutical Design. 2013;19(40):6996-7008.

Caprioli D, Celentano M, Dubla A, Lucantonio F, Nencini P, Badiani A. Ambience and drug choice: cocaine- and heroin-taking as a function of environmental context in humans and rats. Biol Psychiatry. 2009;65(10):893-9.

Spagnolo PA, Celentano M, Dubla A, Badiani A. in *Behavioural Pharmacology* Vol. 22   e21 (2011).

Montanari CL, De Luca MT, Meringolo M, Contu L, Badiani A. in *Behavioural Pharmacology* Vol. 22   e21 (2011).

Kalso E. Oxycodone. Journal of Pain and Symptom Management. 2005;29(5):47-56.

Cahill CM. Opioid dose regimen shapes mesolimbic adaptations. Neuropsychopharmacology. 2020;45(11):1777-78.

Lefevre EM, Pisansky MT, Toddes C, Baruffaldi F, Pravetoni M, Tian L, et al. Interruption of continuous opioid exposure exacerbates drug-evoked adaptations in the mesolimbic dopamine system. Neuropsychopharmacology. 2020;45(11):1781-92.

Winterdahl M, Noer O, Orlowski D, Schacht AC, Jakobsen S, Alstrup AKO, et al. Sucrose intake lowers μ-opioid and dopamine D2/3 receptor availability in porcine brain. Scientific Reports. 2019;9(1):16918.

Pickens CL, Airavaara M, Theberge F, Fanous S, Hope BT, Shaham Y. Neurobiology of the incubation of drug craving. Trends Neurosci. 2011;34(8):411-20.

Bainier C, Mateo M, Felder-Schmittbuhl M-P, Mendoza J. Circadian rhythms of hedonic drinking behavior in mice. Neuroscience. 2017;349:229-38.

Hajnal A, Smith GP, Norgren R. Oral sucrose stimulation increases accumbens dopamine in the rat. Am J Physiol Regul Integr Comp Physiol. 2004;286(1):R31-7.

Hoebel BG, Avena NM, Bocarsly ME, Rada P. Natural addiction: a behavioral and circuit model based on sugar addiction in rats. J Addict Med. 2009;3(1):33-41.

Rada P, Avena NM, Hoebel BG. Daily bingeing on sugar repeatedly releases dopamine in the accumbens shell. Neuroscience. 2005;134(3):737-44.

Spangler R, Wittkowski KM, Goddard NL, Avena NM, Hoebel BG, Leibowitz SF. Opiate-like effects of sugar on gene expression in reward areas of the rat brain. Brain Res Mol Brain Res. 2004;124(2):134-42.

Chan S, Edwards SR, Wyse BD, Smith MT. Sex differences in the pharmacokinetics, oxidative metabolism and oral bioavailability of oxycodone in the Sprague-Dawley rat. Clin Exp Pharmacol Physiol. 2008;35(3):295-302.

Galinkin J, Koh JL. Recognition and Management of Iatrogenically Induced Opioid Dependence and Withdrawal in Children. Pediatrics. 2014;133(1):152.

Association AP. Diagnostic and statistical manual of mental disorders. 5th ed.; 2013.

Madayag AC, Gomez D, Anderson EM, Ingebretson AE, Thomas MJ, Hearing MC. Cell-type and region-specific nucleus accumbens AMPAR plasticity associated with morphine reward, reinstatement, and spontaneous withdrawal. Brain Struct Funct. 2019;224(7):2311-24.

Mameli M, Halbout B, Creton C, Engblom D, Parkitna JR, Spanagel R, et al. Cocaine-evoked synaptic plasticity: persistence in the VTA triggers adaptations in the NAc. Nat Neurosci. 2009;12(8):1036-41.

Creed MC, Lüscher C. Drug-evoked synaptic plasticity: beyond metaplasticity. Curr Opin Neurobiol. 2013;23(4):553-8.

Matikainen-Ankney BA, Garmendia-Cedillos M, Ali M, Krynitsky J, Salem G, Miyazaki NL, et al. Rodent Activity Detector (RAD), an Open Source Device for Measuring Activity in Rodent Home Cages. eNeuro. 2019;6(4).

Godynyuk E, Bluitt MN, Tooley JR, Kravitz AV, Creed MC. An Open-Source, Automated Home-Cage Sipper Device for Monitoring Liquid Ingestive Behavior in Rodents. eNeuro. 2019;6(5).

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, et al. DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nat Neurosci. 2018;21(9):1281-89.

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW. Using DeepLabCut for 3D markerless pose estimation across species and behaviors. Nat Protoc. 2019;14(7):2152-76.

Nilsson SRO, Goodwin NL, Choong JJ, Hwang S, Wright HR, Norville ZC, et al. Simple Behavioral Analysis (SimBA) – an open source toolkit for computer classification of complex social behaviors in experimental animals. bioRxiv. 2020:2020.04.19.049452.

Pedregosa F, Varoquaux G, Gramfort A, Michel V, Thirion B, Grisel O, et al. Scikit-learn: Machine Learning in Python. J Mach Learn Res. 2011;12(null):2825–30.

Bradski G. The OpenCV Library. Dr Dobb's Journal of Software Tools. 2000;120:122-25.

Lalinský L, Wolfer P. ffmpeg version N-93294-g056a2ac168. 2006.

Lemaître G, Nogueira F, Aridas CK. Imbalanced-learn: a python toolbox to tackle the curse of imbalanced datasets in machine learning. J Mach Learn Res. 2017;18(1):559–63.

Liu YL, Liang JH, Yan LD, Su RB, Wu CF, Gong ZH. Effects of l-tetrahydropalmatine on locomotor sensitization to oxycodone in mice. Acta Pharmacol Sin. 2005;26(5):533-8.

Niikura K, Ho A, Kreek MJ, Zhang Y. Oxycodone-induced conditioned place preference and sensitization of locomotor activity in adolescent and adult mice. Pharmacol Biochem Behav. 2013;110:112-6.

Bryant CD, Guido MA, Kole LA, Cheng R. The heritability of oxycodone reward and concomitant phenotypes in a LG/J⬜×⬜SM/J mouse advanced intercross line. Addict Biol. 2014;19(4):552-61.

Gutierrez A, Nguyen JD, Creehan KM, Taffe MA. Female rats self-administer heroin by vapor inhalation. Pharmacology Biochemistry and Behavior. 2020;199:173061.

Sabeti J, Gerhardt GA, Zahniser NR. Individual differences in cocaine-induced locomotor sensitization in low and high cocaine locomotor-responding rats are associated with differential inhibition of dopamine clearance in nucleus accumbens. J Pharmacol Exp Ther. 2003;305(1):180-90.

Fennell AM, Pitts EG, Sexton LL, Ferris MJ. Phasic Dopamine Release Magnitude Tracks Individual Differences in Sensitization of Locomotor Response following a History of Nicotine Exposure. Sci Rep. 2020;10(1):173.

De Vries TJ, Cools AR, Shippenberg TS. Infusion of a D-1 receptor agonist into the nucleus accumbens enhances cocaine-induced behavioural sensitization. Neuroreport. 1998;9(8):1763-8.

Reeves KC, Kube MJ, Grecco GG, Fritz BM, Muñoz B, Yin F, et al. Mu opioid receptors on vGluT2-expressing glutamatergic neurons modulate opioid reward. Addict Biol. 2020:e12942.

Collins D, Reed B, Zhang Y, Kreek MJ. Sex differences in responsiveness to the prescription opioid oxycodone in mice. Pharmacology Biochemistry and Behavior. 2016;148:99-105.

Wolf ME. Synaptic mechanisms underlying persistent cocaine craving. Nat Rev Neurosci. 2016;17(6):351-65.

Bossert JM, Marchant NJ, Calu DJ, Shaham Y. The reinstatement model of drug relapse: recent neurobiological findings, emerging research topics, and translational research. Psychopharmacology (Berl). 2013;229(3):453-76.

Pelloux Y, Everitt BJ, Dickinson A. Compulsive drug seeking by rats under punishment: effects of drug taking history. Psychopharmacology (Berl). 2007;194(1):127-37.

Pascoli V, Terrier J, Hiver A, Lüscher C. Sufficiency of Mesolimbic Dopamine Neuron Stimulation for the Progression to Addiction. Neuron. 2015;88(5):1054-66.

Farrell MR, Ruiz CM, Castillo E, Faget L, Khanbijian C, Liu S, et al. Ventral pallidum is essential for cocaine relapse after voluntary abstinence in rats. Neuropsychopharmacology. 2019;44(13):2174-85.

Hopf FW, Lesscher HM. Rodent models for compulsive alcohol intake. Alcohol. 2014;48(3):253-64.

McCane AM, Auterson CD, DeLory MJ, Lapish CC, Czachowski CL. Differential effects of quinine-adulteration of alcohol on seeking and drinking. Alcohol. 2021.

Reichel CM, Bevins RA. Forced abstinence model of relapse to study pharmacological treatments of substance use disorder. Curr Drug Abuse Rev. 2009;2(2):184-94.

McLellan AT, Lewis DC, O'Brien CP, Kleber HD. Drug dependence, a chronic medical illness: implications for treatment, insurance, and outcomes evaluation. Jama. 2000;284(13):1689-95.

Hollander JA, Carelli RM. Abstinence from cocaine self-administration heightens neural encoding of goal-directed behaviors in the accumbens. Neuropsychopharmacology. 2005;30(8):1464-74.

Terrier J, Lüscher C, Pascoli V. Cell-Type Specific Insertion of GluA2-Lacking AMPARs with Cocaine Exposure Leading to Sensitization, Cue-Induced Seeking, and Incubation of Craving. Neuropsychopharmacology. 2016;41(7):1779-89.

Pascoli V, Terrier J, Espallergues J, Valjent E, O'Connor EC, Lüscher C. Contrasting forms of cocaine-evoked plasticity control components of relapse. Nature. 2014;509(7501):459-64.

Ma YY, Lee BR, Wang X, Guo C, Liu L, Cui R, et al. Bidirectional modulation of incubation of cocaine craving by silent synapse-based remodeling of prefrontal cortex to accumbens projections. Neuron. 2014;83(6):1453-67.

Conrad KL, Tseng KY, Uejima JL, Reimers JM, Heng LJ, Shaham Y, et al. Formation of accumbens GluR2-lacking AMPA receptors mediates incubation of cocaine craving. Nature. 2008;454(7200):118-21.

Graziane NM, Sun S, Wright WJ, Jang D, Liu Z, Huang YH, et al. Opposing mechanisms mediate morphine- and cocaine-induced generation of silent synapses. Nat Neurosci. 2016;19(7):915-25.

Russell SE, Puttick DJ, Sawyer AM, Potter DN, Mague S, Carlezon WA, Jr., et al. Nucleus Accumbens AMPA Receptors Are Necessary for Morphine-Withdrawal-Induced Negative-Affective States in Rats. J Neurosci. 2016;36(21):5748-62.

Zhu Y, Wienecke CF, Nachtrab G, Chen X. A thalamic input to the nucleus accumbens mediates opiate dependence. Nature. 2016;530(7589):219-22.

Hearing MC, Jedynak J, Ebner SR, Ingebretson A, Asp AJ, Fischer RA, et al. Reversal of morphine-induced cell-type-specific synaptic plasticity in the nucleus accumbens shell blocks reinstatement. Proc Natl Acad Sci U S A. 2016;113(3):757-62.

Murray CH, Loweth JA, Milovanovic M, Stefanik MT, Caccamise AJ, Dolubizno H, et al. AMPA receptor and metabotropic glutamate receptor 1 adaptations in the nucleus accumbens core during incubation of methamphetamine craving. Neuropsychopharmacology. 2019;44(9):1534-41.

Alexander BK, Coambs RB, Hadaway PF. The effect of housing and gender on morphine self-administration in rats. Psychopharmacology (Berl). 1978;58(2):175-9.

Hadaway PF, Alexander BK, Coambs RB, Beyerstein B. The effect of housing and gender on preference for morphine-sucrose solutions in rats. Psychopharmacology (Berl). 1979;66(1):87-91.

Klein LC, Popke EJ, Grunberg NE. Sex differences in effects of predictable and unpredictable footshock on fentanyl self-administration in rats. Exp Clin Psychopharmacol. 1997;5(2):99-106.

Moussawi K, Ortiz MM, Gantz SC, Tunstall BJ, Marchette RCN, Bonci A, et al. Fentanyl vapor self-administration model in mice to study opioid addiction. Science Advances. 2020;6(32):eabc0413.

Lynch WJ. Modeling the development of drug addiction in male and female animals. Pharmacology Biochemistry and Behavior. 2018;164:50-61.

Lofwall MR, Nuzzo PA, Walsh SL. Effects of cold pressor pain on the abuse liability of intranasal oxycodone in male and female prescription opioid abusers. Drug and Alcohol Dependence. 2012;123(1):229-38.

Lee CW-S, Ho I-K. Sex differences in opioid analgesia and addiction: interactions among opioid receptors and estrogen receptors. Molecular Pain. 2013;9(1):45.

Serdarevic M, Striley CW, Cottler LB. Sex differences in prescription opioid use. Curr Opin Psychiatry. 2017;30(4):238-46.

Benziger DP, Kaiko RF, Miotto JB, Fitzmartin RD, Reder RF, Chasin M. Differential effects of food on the bioavailability of controlled-release oxycodone tablets and immediate-release oxycodone solution. J Pharm Sci. 1996;85(4):407-10.

Webster LR, Kinzler ER, Pantaleon C, Iverson M, Aigner S. Relative Oral Bioavailability of an Abuse-Deterrent, Immediate-Release Formulation of Oxycodone, Oxycodone ARIR in a Randomized Study. Advances in Therapy. 2019;36(7):1730-40.

Samer CF, Daali Y, Wagner M, Hopfgartner G, Eap CB, Rebsamen MC, et al. The effects of CYP2D6 and CYP3A activities on the pharmacokinetics of immediate release oxycodone. Br J Pharmacol. 2010;160(4):907-18.

Grönlund J, Saari TI, Hagelberg NM, Neuvonen PJ, Olkkola KT, Laine K. Exposure to oral oxycodone is increased by concomitant inhibition of CYP2D6 and 3A4 pathways, but not by inhibition of CYP2D6 alone. Br J Clin Pharmacol. 2010;70(1):78-87.

Zanger UM, Schwab M. Cytochrome P450 enzymes in drug metabolism: regulation of gene expression, enzyme activities, and impact of genetic variation. Pharmacol Ther. 2013;138(1):103-41.

Soldin OP, Mattison DR. Sex differences in pharmacokinetics and pharmacodynamics. Clin Pharmacokinet. 2009;48(3):143-57.

Mavrikaki M, Pravetoni M, Page S, Potter D, Chartoff E. Oxycodone self-administration in male and female rats. Psychopharmacology (Berl). 2017;234(6):977-87.

Deroche-Gamonet V, Piazza PV. Psychobiology of cocaine addiction: Contribution of a multi-symptomatic animal model of loss of control. Neuropharmacology. 2014;76 Pt B:437-49.

Ahmed SH, Koob GF. Transition from moderate to excessive drug intake: change in hedonic set point. Science. 1998;282(5387):298-300.

Shurman J, Koob GF, Gutstein HB. Opioids, pain, the brain, and hyperkatifeia: a framework for the rational use of opioids for pain. Pain Med. 2010;11(7):1092-8.

Koob GF. Neurobiology of Opioid Addiction: Opponent Process, Hyperkatifeia, and Negative Reinforcement. Biol Psychiatry. 2020;87(1):44-53.

Bravo IM, Luster BR, Flanigan ME, Perez PJ, Cogan ES, Schmidt KT, et al. Divergent behavioral responses in protracted opioid withdrawal in male and female C57BL/6J mice. European Journal of Neuroscience. 2020;51(3):742-54.

Marchant NJ, Campbell EJ, Pelloux Y, Bossert JM, Shaham Y. Context-induced relapse after extinction versus punishment: similarities and differences. Psychopharmacology (Berl). 2019;236(1):439-48.

Aitken M, Kleinrock M.   (Iqvia Institute for Human Data Science, 2018).

Bossert JM, Hoots JK, Fredriksson I, Adhikary S, Zhang M, Venniro M, et al. Role of mu, but not delta or kappa, opioid receptors in context-induced reinstatement of oxycodone seeking. Eur J Neurosci. 2019;50(3):2075-85.

Scheyer AF, Loweth JA, Christian DT, Uejima J, Rabei R, Le T, et al. AMPA Receptor Plasticity in Accumbens Core Contributes to Incubation of Methamphetamine Craving. Biol Psychiatry. 2016;80(9):661-70.

Sutton MA, Schmidt EF, Choi KH, Schad CA, Whisler K, Simmons D, et al. Extinction-induced upregulation in AMPA receptors reduces cocaine-seeking behaviour. Nature. 2003;421(6918):70-5.

Darevsky D, Hopf FW. Behavioral indicators of succeeding and failing under higher-challenge compulsion-like alcohol drinking in rat. Behav Brain Res. 2020;393:112768.

Fulenwider HD, Nennig SE, Price ME, Hafeez H, Schank JR. Sex Differences in Aversion-Resistant Ethanol Intake in Mice. Alcohol Alcohol. 2019;54(4):345-52.

Kasanetz F, Deroche-Gamonet V, Berson N, Balado E, Lafourcade M, Manzoni O, et al. Transition to addiction is associated with a persistent impairment in synaptic plasticity. Science. 2010;328(5986):1709-12.

Pascoli V, Hiver A, Van Zessen R, Loureiro M, Achargui R, Harada M, et al. Stochastic synaptic plasticity underlying compulsion in a model of addiction. Nature. 2018;564(7736):366-71.

Deroche-Gamonet V, Belin D, Piazza PV. Evidence for addiction-like behavior in the rat. Science. 2004;305(5686):1014-7.

Anthony JC, Warner LA, Kessler RC. Comparative epidemiology of dependence on tobacco, alcohol, controlled substances, and inhalants: Basic findings from the National Comorbidity Survey. Experimental and Clinical Psychopharmacology. 1994;2(3):244-68.

Ortinski PI, Vassoler FM, Carlson GC, Pierce RC. Temporally dependent changes in cocaine-induced synaptic plasticity in the nucleus accumbens shell are reversed by D1-like dopamine receptor stimulation. Neuropsychopharmacology. 2012;37(7):1671-82.

Scofield MD, Heinsbroek JA, Gipson CD, Kupchik YM, Spencer S, Smith ACW, et al. The Nucleus Accumbens: Mechanisms of Addiction across Drug Classes Reflect the Importance of Glutamate Homeostasis. Pharmacological reviews. 2016;68(3):816-71.

McCutcheon JE, Wang X, Tseng KY, Wolf ME, Marinelli M. Calcium-permeable AMPA receptors are present in nucleus accumbens synapses after prolonged withdrawal from cocaine self-administration but not experimenter-administered cocaine. J Neurosci. 2011;31(15):5737-43.

Jedynak J, Hearing M, Ingebretson A, Ebner SR, Kelly M, Fischer RA, et al. Cocaine and Amphetamine Induce Overlapping but Distinct Patterns of AMPAR Plasticity in Nucleus Accumbens Medium Spiny Neurons. Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology. 2016;41(2):464-76.

Ebner SR, Larson EB, Hearing MC, Ingebretson AE, Thomas MJ. Extinction and Reinstatement of Cocaine-seeking in Self-administering Mice is Associated with Bidirectional AMPAR-mediated Plasticity in the Nucleus Accumbens Shell. Neuroscience. 2018;384:340-49.

Lüscher C, Malenka Robert C. Drug-Evoked Synaptic Plasticity in Addiction: From Molecular Changes to Circuit Remodeling. Neuron. 2011;69(4):650-63.

Wolf ME, Tseng KY. Calcium-permeable AMPA receptors in the VTA and nucleus accumbens after cocaine exposure: when, how, and why? Front Mol Neurosci. 2012;5:72.

Vander Weele CM, Porter-Stransky KA, Mabrouk OS, Lovic V, Singer BF, Kennedy RT, et al. Rapid dopamine transmission within the nucleus accumbens: dramatic difference between morphine and oxycodone delivery. Eur J Neurosci. 2014;40(7):3041-54.

Park WK, Bari AA, Jey AR, Anderson SM, Spealman RD, Rowlett JK, et al. Cocaine administered into the medial prefrontal cortex reinstates cocaine-seeking behavior by increasing AMPA receptor-mediated glutamate transmission in the nucleus accumbens. J Neurosci. 2002;22(7):2916-25.

Creed M, Pascoli VJ, Lüscher C. Addiction therapy. Refining deep brain stimulation to emulate optogenetic treatment of synaptic pathology. Science. 2015;347(6222):659-64.

Figure Captions

Figure 1. Mice voluntarily consume oxycodone in a two-bottle choice paradigm. (a) schematic of experimental paradigm (n = CTRL: 20M/20F, OXY: 26M/26F). **(b-c)** Volumes of total liquid consumed in control or oxycodone-administering mice (CTRL female 3.527 ± 0.044, male = 3.422 ± 0.049, OXY female = 4.233 ± 0.103, male = 3.862 ± 0.083). Mice with OXY available drank significantly higher volumes over the two-bottle choice phase (F_{drug} = 55.679, p < 0.001), and female mice consumed higher volumes of OXY during the self-administration protocol (F_{sex} = 9.588, p = 0.002, $F_{sex \bullet drug}$ = 1.833, p = 0.045). **(d)** Oxycodone dosage increased throughout the single bottle phase and mice continued to escalate their consumed oxycodone levels throughout the two-bottle choice phase (F_{Time} = 2.743, p = 0.002). **(e)** There was no sex difference in escalation index, calculated as the dose on the final day of two-bottle choice (day 12) divided by dose on the first day of the two-bottle choice phase (day 6; Females = 1.610 ± 0.193, Males = 1.748 ± 0.153, F = 0.313, p = 0.579). **(f)** Female mice self-administered higher total doses of oxycodone over the course of the protocol (Females = 1499.135 ± 78.745, Males = 1147.788 ± 59.855, F = 12.619, p = 0.001). **(g-h)** OXY-self-administering mice, but not control mice developed a significant preference for the oxycodone-containing bottle over the 7 days of two-bottle choice, which did not differ between males and females (F_{drug} = 227.488 p < 0.001, $F_{drug \bullet day}$ = 9.233, p = 0.003, F_{sex} = 3.308, p = 0.070). **(i)** There was no systematic preference for drinking water from the left or right bottle in CTRL mice, while mice consumed significantly more OXY solution relative to drinking water when available (CTRL female = 44.8 ± 4.9%, male = 53.9 ± 6.0% preference for right drinking bottle; OXY female = 88.2 ± 3.9%, male = 89.6 ± 3.2% preference for OXY-containing bottle, F_{drug} = 76.45 p < 0.001).

Figure 2. Homecage drinking monitoring confirms OXY preference and reveals altered patterns of OXY and water intake. (a-b) Sipper counts (photobeam breaks on each lickometer) were normalized for total Sipper Counts made over the 7 days of the two-bottle choice task and plotted in 1 hour bins in CTRL mice. There was no difference in sipper counts on the left vs. right sipper in control mice (CTRL female Left/Right = 0.531 ± 0.039/0.466 ± 0.039, male Left/Right = 0.465 ± 0.061/0.471 ± 0.048). **(c-d)** Counts were significantly higher on the oxycodone-containing bottle relative to the water containing bottle in oxycodone-administering mice (OXY female H_2O/OXY = 0.287 ± 0.075/0.641 ± 0.087, male H_2O/OXY = 0.218 ± 0.036/0.782 ± 0.036, F_{bottle} = 29.485, p < 0.001, $F_{drug \bullet bottle}$ = 38.154, p < 0.001). **(e)** Circadian index for Sipper Counts on each bottle (Dark Cycle Counts/Light Cycle Counts). Circadian index for both water and oxycodone drinking was significantly lower in OXY-self administering mice (CTRL female Left/Right = 0.740 ± 0.030/0.783 ± 0.029, male Left/Right = 0.737 ± 0.021/0.768 ± 0.029, F_{Drug} = 28.617 p < 0.001, F_{Bottle} = 0.899, p = 0.345). **(f)** Pattern of Sipper Counts was also significantly different between CTRL and OXY-self-administering mice, with OXY mice showing more variable and shorter inter-lick bout intervals (Kolmogorov-Smirnov = 1.550, p = 0.0163).

Figure 3. Oral oxycodone self-administration induces hyperactivity and persistent physical withdrawal signs. (a) Experimental schematic. Homecages were equipped with PIR sensors to monitor activity during the single- and two-bottle choice phases of OXY-self administration. Withdrawal symptoms were assayed after 24h and after 3 weeks of withdrawal. **(b-c)** Both male and female mice exhibited significantly higher homecage activity during OXY self-administration than control mice (CTRL: 2072.189 ± 196.455, OXY = 4530.370, F_{Sex} = 0.956, p = 0.333, F_{Drug}=19.549, p<0.001). **(d)** There was a significant effect of drug and time point, with OXY mice exhibiting higher levels of movement relative to control mice at acute (CTRL: 473.18 ± 20.18, OXY:

570.65 ± 43.45 cm) and protracted (CTRL: 555.56 ± 27.62, OXY: 689.67 ± 29.00 cm) withdrawal time points ($F_{TimePoint}$=10.938, p = 0.002, F_{Drug} = 13.343, p=0.001). **(e)** OXY mice exhibited greater numbers of jumps at both early (CTRL: 0.778 ± 0.298, OXY: 6.824 ± 2.254) and late withdrawal points (CTRL: 2.400 ± 0.498, OXY: 7.735 ± 1.523, F_{Drug} = 15.284, p < 0.001). **(f)** OXY mice exhibited increased paw tremors at both early (CTRL: 4.053 ± 1.217, OXY: 9.132 ± 2.423) and late time points (CTRL: 2.420 ± 0.590, OXY: 6.025 ± 1.763, F_{Drug} = 7.155, p = 0.010). **(g-h)** While there was a significant effect of time point on both rearing and grooming behavior (Rearing $F_{TimePoint}$ = 46.380, p < 0.001, Grooming $F_{TimePoint}$ = 7.757, p = 0.007), these behaviors were not different between control and OXY-self-administering mice. **(i)** There was a significant effect of drug on climbing time at the acute time point only (CTRL: 7.862 ± 1.269, OXY: 21.092 ± 6.080 sec, F_{Drug} = 5.065, p =0.028). (h-i) *p<0.05, **p<0.01, ***p<0.001.

Figure 4. Drug seeking behavior persists into protracted withdrawal. (a) Experimental timeline (CTRL n = 10M/10F, OXY n = 10M/10F). 24h (early seeking) and 24 days (late seeking) after the two-bottle choice protocol, drinking counts were measured under extinction conditions. **(b)** Time course of photobeam breaks (sipper counts) made on empty bottles over the 60 minute probe trial; mice that had self-administered OXY showed more seeking interactions relative to controls (F_{drug} = 51.062 p < 0.0001), with no difference between early and late seeking ($F_{EarlyVsLate}$ = 0.227, p = 0.653, $F_{drug•EarlyVsLate}$ = 0.927, p=0.339). **(c-e)** Both male and female mice that had self-administered OXY exhibited higher sipper counts in early withdrawal (CTRL female = 17.3 ± 3.50, male = 6.5 ± 2.50, OXY female = 62.9 ± 12.33, male = 82.6 ± 20.87, F_{drug} = 24.427, p<0.001). **(f-h)** Both male and female mice that had self-administered OXY exhibited higher sipper counts in early withdrawal (CTRL female = 15.2 ± 5.64, male = 15.9 ± 3.55, OXY female = 59.4 ± 13.58, male = 64.5 ± 8.12, F_{drug} = 29.220, p<0.001).

Figure 5. Oral OXY self-administration is resistant to quinine adulteration. (a) Experimental schematic. After seven days of two-bottle choice, the OXY-containing bottle (or one H_2O bottle in CTRL mice) was adulterated with increasing concentrations of quinine. CTRL = 7M/8F, OXY n=8M/7F. **(b-c)** Normalized volume consumed from the quinine-adulturated bottle over the final three days of two bottle choice (before quinine adulteration) and for 6 days of quinine adulteration. **(d-e)** Area under the curve of adulterated quinine drinking volumes. There was a significant effect of Drug (Fdrug = 27.908, p<0.001), of day (Fday = 66.767, p<0.001) and interaction (Fdrug•day = 7.485, p<0.001), with both male and female OXY mice exhibiting significantly higher AUC relative to CTRL mice (Fsex = 0.6896, p = 0.414, CTRL female = 0.862 ± 0.130, male = 0.989 ± 0.201, OXY female = 3.495 ± 0.662, male = 2.704 ± 0.415).

Figure 6. Excitatory transmission onto NAc MSNs is increased after withdrawal from OXY self-administration. (a) experimental schematic and representative alexofluor-488 filled MSN (green), DAPI nuclear stain shown in blue. **(b)** Location of recorded cells in control (above) and oxycodone-self administering groups (bottom); AP coordinates relative to brema are indicated (mm) **(c)** representative traces showing calculation of AMPA/NMDA and paired pulse ratio in control (top) and OXY withdrawn (bottom) groups; scale bar = 50 pA, 50 ms. **(d)** There was a significant increase in A/N following OXY withdrawal (CTRL = 1.79 ± 0.13, OXY = 3.46 ± 0.33; F_{drug} = 21.07, p<0.0001), and a significant drug by sex interaction ($F_{drug•sex}$ = 4.98, p = 0.030). **(e)** PPR was not different between groups (CTRL = 1.06 ± 0.06, OXY = 0.945 ± 0.07; F_{drug} = 1.37, p=0.36). **(f-g)** There was a significant increase in mEPSC amplitude following OXY withdrawal (Kolmogorov–Smirnov test statistic = 2.89, p < 0.05) but no difference in mEPSC frequency between groups (Kolmogorov–Smirnov test statistic = 1.12, p = 0.16). Scale bar = 20 pA, 1 sec.

Supplemental Figure 1. Individual data for oxycodone dose and preference. a) Baseline preference for 1.0 mg/ml oxycodone in naïve mice; while both males and females showed a preference for water over oxycodone, the initial preference was higher for females, which was driven by a subset of oxy-preferring mice (males: 5.21 ± 1.39, females: 24.03 ± 7.90, F_{sex} = 6.59, p = 0.014). **b)** Body weight for individual subjects; while the average weight of males was significantly higher than that of females (CTRL male = 26.09 ± 37, female = 20.96 ±0.39, OXY male = 25.57 ± 0.41, female = 22.07 ± 0.46), drug treatment did not have a significant effect on body weight (F_{sex} = 107.174, p< 0.0001, F_{Drug} = 0.474, p = 0.493, $F_{sex*drug}$ = 3.837, p = 0.054). (c-d) Daily dose escalation curves for individual subjects (n = 20 M, 20 F). **(c-d)** Preference for oxycodone for individual mice; preference is defined as [*consumed volume of oxycodone / total volume of fluid consumption*]. **(e-f)** Escalation of oxycodone dose for individual mice; dose is defined as [*mg oxycodone consumed / body weight*].

Supplemental Figure 2. Individual data for sipper counts over two-bottle choice phase in control and oxycodone self-administering mice. Individual cumulative lickometer photo-interrupter counts for each subject are shown. Counts made on the left and right bottles are shown for controls are shown separately (left, right, respectively). Counts made on the oxycodone-filled and water-filled bottles are shown separately (oxycodone, water, respectively).

Supplemental Figure 3. Oxycodone-induced activity patterns did not differ between male and female mice. PIR counts during the single bottle and two-bottle choice phases of the self-administration protocol are shown for male mice **(a)** and female mice **(b)**. **P < 0.01

Supplemental Figure 4. Individual data for sipper counts during early and late seeking tasks. Individual photobeam breaks made during the 60 minute probe test made under extinction conditions in both the early **(a)** and late **(b)** withdrawal time points. Data from males (left) and females (right) is shown.

Figure 1. Mice voluntarily consume oxycodone in a two-bottle choice paradigm.

Figure 1. Mice voluntarily consume oxycodone in a two-bottle choice paradigm. (a) schematic of experimental paradigm (n = CTRL: 20M/20F, OXY: 26M/26F). **(b-c)** Volumes of total liquid consumed in control or oxycodone-administering mice (CTRL female 3.527 ± 0.044, male = 3.422 ± 0.049, OXY female = 4.233 ± 0.103, male = 3.862 ± 0.083). Mice with OXY available drank significantly higher volumes over the two-bottle choice phase (F_{drug} = 55.679, p < 0.001), and female mice consumed higher volumes of OXY during the self-administration protocol (F_{sex} = 9.588, p = 0.002, $F_{sex \bullet drug}$ = 1.833, p = 0.045). **(d)** Oxycodone dosage increased throughout the single bottle phase and mice continued to escalate their consumed oxycodone levels throughout the two-bottle choice phase (F_{Time} = 2.743, p = 0.002). **(e)** There was no sex difference in escalation index, calculated as the dose on the final day of two-bottle choice (day 12) divided by dose on the first day of the two-bottle choice phase (day 6; Females = 1.610 ± 0.193, Males = 1.748 ± 0.153, F = 0.313, p = 0.579). **(f)** Female mice self-administered higher total doses of oxycodone over the course of the protocol (Females = 1499.135 ± 78.745, Males = 1147.788 ± 59.855, F = 12.619, p = 0.001). **(g-h)** OXY-self-administering mice, but not control mice developed a significant preference for the oxycodone-containing bottle over the 7 days of two-bottle choice, which did not differ between males and females (F_{drug} = 227.488 p < 0.001, $F_{drug \bullet day}$ = 9.233, p = 0.003, F_{sex} = 3.308, p = 0.070). **(i)** There was no systematic preference for drinking water from the left or right bottle in CTRL mice, while mice consumed significantly more OXY solution relative to drinking water when available (CTRL female = 44.8 ± 4.9%, male = 53.9 ± 6.0% preference for right drinking bottle; OXY female = 88.2 ± 3.9%, male = 89.6 ± 3.2% preference for OXY-containing bottle, F_{drug} = 76.45 p < 0.001).

Supplemental Figure 1. Individual data for oxycodone dose and preference.

Supplemental Figure 1. Individual data for oxycodone dose and preference. a) Baseline preference for 1.0 mg/ml oxycodone in naïve mice; while both males and females showed a preference for water over oxycodone, the initial preference was higher for females, which was driven by a subset of oxy-preferring mice (males: 5.21 ± 1.39, females: 24.03 ± 7.90, $F_{sex} = 6.59$, $p = 0.014$). **b)** Body weight for individual subjects; while the average weight of males was significantly higher than that of females (CTRL male = 26.09 ± 37, female = 20.96 ± 0.39, OXY male = 25.57 ± 0.41, female = 22.07 ± 0.46), drug treatment did not have a significant effect on body weight ($F_{sex} = 107.174$, $p < 0.0001$, $F_{Drug} = 0.474$, $p = 0.493$, $F_{sex*drug} = 3.837$, $p = 0.054$). (c-d) Daily dose escalation curves for individual subjects (n = 20 M, 20 F). **(c-d)** Preference for oxycodone for individual mice; preference is defined as [*consumed volume of oxycodone / total volume of fluid consumption*]. **(e-f)** Escalation of oxycodone dose for individual mice; dose is defined as [*mg oxycodone consumed / body weight*].

Figure 2. Altered patterns of oxycodone intake are revealed by homecage monitoring.

Figure 2. Homecage drinking monitoring confirms OXY preference and reveals altered patterns of OXY and water intake. (a-b) Sipper counts (photobeam breaks on each lickometer) were normalized for total Sipper Counts made over the 7 days of the two-bottle choice task and plotted in 1 hour bins in CTRL mice. There was no difference in sipper counts on the left vs. right sipper in control mice (CTRL female Left/Right = 0.531 ± 0.039/0.466 ± 0.039, male Left/Right = 0.465 ± 0.061/0.471 ± 0.048). **(c-d)** Counts were significantly higher on the oxycodone-containing bottle relative to the water containing bottle in oxycodone-administering mice (OXY female H_2O/OXY = 0.287 ± 0.075/0.641 ± 0.087, male H_2O/OXY = 0.218 ± 0.036/0.782 ± 0.036, F_{bottle} = 29.485, p < 0.001, $F_{drug \bullet bottle}$ = 38.154, p < 0.001). **(e)** Circadian index for Sipper Counts on each bottle (Dark Cycle Counts/Light Cycle Counts). Circadian index for both water and oxycodone drinking was significantly lower in OXY-self administering mice (CTRL female Left/Right = 0.740 ± 0.030/0.783 ± 0.029, male Left/Right = 0.737 ± 0.021/0.768 ± 0.029, F_{Drug} = 28.617 p < 0.001, F_{Bottle} = 0.899, p = 0.345). **(f)** Pattern of Sipper Counts was also significantly different between CTRL and OXY-self-administering mice, with OXY mice showing more variable and shorter inter-lick bout intervals (Kolmogorov-Smirnov = 1.550, p = 0.0163).

Supplemental Figure 2. Individual data for lickometer counts during two-bottle choice phase in control and oxycodone self-administering mice.

Supplemental Figure 2. Individual data for sipper counts over two-bottle choice phase in control and oxycodone self-administering mice.
Individual cumulative lickometer photo-interrupter counts for each subject are shown. Counts made on the left and right bottles are shown for controls are shown separately (left, right, respectively). Counts made on the oxycodone-filled and water-filled bottles are shown separately (oxycodone, water, respectively).

Figure 3. Oral oxycodone self-administration induces hyperactivity and persistent physical withdrawal signs.

Figure 3. Oral oxycodone self-administration induces hyperactivity and persistent physical withdrawal signs. (a) Experimental schematic. Homecages were equipped with PIR sensors to monitor activity during the single- and two-bottle choice phases of OXY-self administration. Withdrawal symptoms were assayed after 24h and after 3 weeks of withdrawal. **(b-c)** Both male and female mice exhibited significantly higher homecage activity during OXY self-administration than control mice (CTRL: 2072.189 ± 196.455, OXY = 4530.370, F_{Sex} = 0.956, p = 0.333, F_{Drug}=19.549, p<0.001). **(d)** There was a significant effect of drug and time point, with OXY mice exhibiting higher levels of movement relative to control mice at acute (CTRL: 473.18 ± 20.18, OXY: 570.65 ± 43.45 cm) and protracted (CTRL: 555.56 ± 27.62, OXY: 689.67 ± 29.00 cm) withdrawal time points ($F_{TimePoint}$=10.938, p = 0.002, F_{Drug} = 13.343, p=0.001). **(e)** OXY mice exhibited greater numbers of jumps at both early (CTRL: 0.778 ± 0.298, OXY: 6.824 ± 2.254) and late withdrawal points (CTRL: 2.400 ± 0.498, OXY: 7.735 ± 1.523, F_{Drug} = 15.284, p < 0.001). **(f)** OXY mice exhibited increased paw tremors at both early (CTRL: 4.053 ± 1.217, OXY: 9.132 ± 2.423) and late time points (CTRL: 2.420 ± 0.590, OXY: 6.025 ± 1.763, F_{Drug} = 7.155, p = 0.010). **(g-h)** While there was a significant effect of time point on both rearing and grooming behavior (Rearing $F_{TimePoint}$ = 46.380, p < 0.001, Grooming $F_{TimePoint}$ = 7.757, p = 0.007), these behaviors were not different between control and OXY-self-administering mice. **(i)** There was a significant effect of drug on climbing time at the acute time point only (CTRL: 7.862 ± 1.269, OXY: 21.092 ± 6.080 sec, F_{Drug} = 5.065, p =0.028). (h-i) *p<0.05, **p<0.01, ***p<0.001.

Supplemental Figure 3. Oxycodone induced activity patterns did not differ between male and female mice.

Supplemental Figure 3. Oxycodone-induced activity patterns did not differ between male and female mice. PIR counts during the single bottle and two-bottle choice phases of the self-administration protocol are shown for male mice **(a)** and female mice **(b)**. **P < 0.01

Figure 4. Drug seeking behavior persists into protracted withdrawal.

Figure 4. Drug seeking behavior persists into protracted withdrawal.
(a) Experimental timeline (CTRL n = 10M/10F, OXY n = 10M/10F). 24h
(early seeking) and 24 days (late seeking) after the two-bottle choice
protocol, drinking counts were measured under extinction conditions. **(b)**
Time course of photobeam breaks (sipper counts) made on empty bottles
over the 60 minute probe trial; mice that had self-administered OXY
showed more seeking interactions relative to controls (F_{drug} = 51.062 p <
0.0001), with no difference between early and late seeking ($F_{EarlyVsLate}$ =
0.227, p = 0.653, $F_{drug \bullet EarlyVsLate}$ = 0.927, p=0.339). **(c-e)** Both male and
female mice that had self-administered OXY exhibited higher sipper
counts in early withdrawal (CTRL female = 17.3 ± 3.50, male = 6.5 ± 2.50,
OXY female = 62.9 ± 12.33, male = 82.6 ± 20.87, F_{drug} = 24.427, p<0.001).
(f-h) Both male and female mice that had self-administered OXY
exhibited higher sipper counts in early withdrawal (CTRL female = 15.2 ±
5.64, male = 15.9 ± 3.55, OXY female = 59.4 ± 13.58, male = 64.5 ± 8.12,
F_{drug} = 29.220, p<0.001).

Supplemental Figure 4. Individual data for oxycodone seeking in extinction after acute and protracted withdrawal.

Supplemental Figure 4. Individual data for sipper counts during early and late seeking tasks. Individual photobeam breaks made during the 60 minute probe test made under extinction conditions in both the early **(a)** and late **(b)** withdrawal time points. Data from males (left) and females (right) is shown.

Figure 5. A subset of mice persist in oxycodone consumption despite quinine adulturation.

Figure 5. Oral OXY self-administration is resistant to quinine adulteration.
(a) Experimental schematic. After seven days of two-bottle choice, the OXY-containing bottle (or one H_2O bottle in CTRL mice) was adulterated with increasing concentrations of quinine. CTRL = 7M/8F, OXY n=8M/7F. **(b-c)** Normalized volume consumed from the quinine-adulturated bottle over the final three days of two bottle choice (before quinine adulteration) and for 6 days of quinine adulteration. **(d-e)** Area under the curve of adulterated quinine drinking volumes. There was a significant effect of Drug (F_{drug} = 27.908, $p < 0.001$), of day (F_{day} = 66.767, $p < 0.001$) and interaction ($F_{drug \bullet day}$ = 7.485, $p < 0.001$), with both male and female OXY mice exhibiting significantly higher AUC relative to CTRL mice (F_{sex} = 0.6896, p = 0.414, CTRL female = 0.862 $\pm$ 0.130, male = 0.989 $\pm$ 0.201, OXY female = 3.495 $\pm$ 0.662, male = 2.704 $\pm$ 0.415).

Figure 6. Excitatory transmission onto NAc-MSNs is increased after withdrawal from oxycodone self-administration.

Figure 6. Excitatory transmission onto NAc MSNs is increased after withdrawal from OXY self-administration. (a) experimental schematic and representative alexofluor-488 filled MSN (green), DAPI nuclear stain shown in blue. **(b)** Location of recorded cells in control (above) and oxycodone-self administering groups (bottom); AP coordinates relative to brema are indicated (mm) **(c)** representative traces showing calculation of AMPA/NMDA and paired pulse ratio in control (top) and OXY withdrawn (bottom) groups; scale bar = 50 pA, 50 ms. **(d)** There was a significant increase in A/N following OXY withdrawal (CTRL = 1.79 ± 0.13, OXY = 3.46 ± 0.33; F_{drug} = 21.07, p<0.0001), and a significant drug by sex interaction ($F_{drug \bullet sex}$ = 4.98, p = 0.030). **(e)** PPR was not different between groups (CTRL = 1.06 ± 0.06, OXY = 0.945 ± 0.07; F_{drug} = 1.37, p=0.36). **(f-g)** There was a significant increase in mEPSC amplitude following OXY withdrawal (Kolmogorov–Smirnov test statistic = 2.89, p < 0.05) but no difference in mEPSC frequency between groups (Kolmogorov–Smirnov test statistic = 1.12, p = 0.16). Scale bar = 20 pA, 1 sec.