1   POPULATION GENOMIC ANALYSIS OF *CRYPTOCOCCUS* BRAZILIAN ISOLATES REVEALS AN
2   AFRICAN TYPE SUBCLADE DISTRIBUTION
3
4

5   Corinne MAUFRAIS[1,2†], Luciana de OLIVEIRA[1†], Rafael W. BASTOS[3], Frédérique MOYRAND[1],
6   Flavia C. G. REIS[4,5], Clara VALERO[3], Bianca GIMENEZ[4], Luisa J. JOSEFOWICZ[4], Gustavo H.
7   GOLDMAN[3], Marcio L. RODRIGUES[4,6] and Guilhem JANBON[1]*
8
[1] Institut Pasteur, Unité Biologie des ARN des Pathogènes Fongiques, Département de
Mycologie, F-75015, Paris, France
[2] Institut Pasteur, HUB Bioinformatique et Biostatistique, C3BI, USR 3756 IP CNRS, F-75015, Paris,
France
[3] Departamento de Ciências Farmacêuticas, Faculdade de Ciências Farmacêuticas de Ribeirão
Preto, Universidade de São Paulo, Ribeirão Preto,Brazil
[4] Instituto Carlos Chagas, Fundação Oswaldo Cruz (FIOCRUZ), Curitiba, Brazil
[5] Centro de Desenvolvimento Tecnologico em Saude (CDTS-Fiocruz), Brazil
[6] Instituto de Microbiologia Paulo de Góes (IMPG), Universidade Federal do Rio de Janeiro, Rio
de Janeiro, Brazil
19
20
* Corresponding author
† Both authors should be considered as first authors

Abstract

The genomes of a large number of *Cryptococcus neoformans* isolates have been

sequenced and analyzed in recent years. These genomes have been used to understand the

global population structure of this opportunistic pathogen. However, only a small number of

South American isolates have been considered in these studies, and the population structure

of *C. neoformans* in this part of the world remains elusive. Here, we analyzed the genomic

sequences of 53 Brazilian *Cryptococcu*s isolates and deciphered the *C. neoformans* population

structure in this country. Our data reveal an African-like structure that suggested repeated

intercontinental transports from Africa to South America. We also identified a mutator

phenotype in one VNBII Brazilian isolate, exemplifying how fast-evolving isolates can shape the

Cryptococcus population structure. Finally, phenotypic analyses revealed wide diversity but not

lineage specificity in the expression of classical virulence traits within the set of isolates.

The current classification identifies seven human-pathogenic *Cryptococcus* species,

which are responsible for 180,000 deaths every year in the world (Hagen *et al.* 2015;

Rajasingham *et al.* 2017). *C. neoformans* and *C. deneoformans* represent the *neoformans*

group of species, whereas the *gattii* group of species comprises *C. gattii, C. deuterogattii, C.*

tetragattii, C. decagattii, and *C. bacillisporus* (Hagen *et al.* 2015). *C. neoformans* is the most

common species in clinical settings, representing more than 95% of the clinical isolates (Kwon-

Chung *et al.* 2014; Janbon *et al.* 2019). This species affects mostly immunocompromised

patients in whom it provokes meningoencephalitis. Other species, such as *C. gattii*, can be

primary pathogens and cause pulmonary infections (Kwon-Chung *et al.* 2014). *C. neoformans* is

thought to be acquired very early in life and can stay in a dormant state for years, probably in

alveolar macrophages (Garcia-Hermoso *et al.* 1999; Alanio 2020). As soon as a defect in the

cellular immunity appears, yeast cells can multiply, reach the central nervous system, and

induce meningoencephalitis, which is invariably deadly if untreated.

C. neoformans strains are ubiquitous in the environment and have been isolated from all

continents except Antarctica. Similarly, although isolates of the *gattii* group of species have

been long considered to be geographically restricted to tropical and subtropical regions and

typically associated with *Eucalyptus* trees, they have emerged as a primary pathogen in

western Canada and the United States of America (Kidd *et al.* 2004). Both species can live in a

broad range of ecological habitats and have been commonly isolated from soil, guano,

decaying wood, fruits, and insects (Lin and Heitman 2006). Pigeons (*Columba livia*) are vectors

for the worldwide dissemination of *Cryptococcus*. Indeed, dry pigeon guano seems to be an

ideal environment for *C. neoformans* development and is commonly infected with this fungus.

Nevertheless, pigeons are usually not harmed by *C. neoformans* because their body

temperature is too high (42.5°C) for its growth and disease development (Johnston *et al.*

2016).

All *neoformans* and *gattii* species are haploid species with two possible mating type alleles

(*MAT***a** and *MAT*α) and a bipolar mating system. Efficient sexual reproduction has also been

described in the laboratory. Nevertheless, *C. neoformans MAT***a** isolates are rare in clinical and

environmental isolates with the exception of some places in Africa (Kwon-Chung and Bennett

1978; Litvintseva *et al.* 2003). In addition, it has been reported that 8% of the population of *C.*

neoformans isolates are diploid (Lin *et al.* 2009). Diverse hybrids have also been isolated from

environmental and clinical sources. The most common are AD hybrids, which are produced by

hybridization of *C. neoformans* with *C. deneoformans* isolates. In addition, rare cases of

gattii/*neoformans* hybrids have been reported (Bovers *et al.* 2008; Aminnejad *et al.* 2012).

Nevertheless, population structure analysis suggested that the different *neoformans* and *gattii*

populations are largely clonal with a limited degree of recombination (Litvintseva *et al.* 2005;

Bui *et al.* 2008; Litvintseva and Mitchell 2012).

Historically, these analyses were based on serotyping, which was first replaced by PCR-based

characterization (RAPD, AFLP, mating type analysis) of the isolates and then by sequencing of a

defined set of genes by multilocus sequence typing (MLST) (Ruma *et al.* 1996; Boekhout *et al.*

2001; Matsumoto *et al.* 2007; Meyer *et al.* 2009a). Originally, five *neoformans* molecular

subtypes were thus defined (VNI, VNII, VNIV, VNIII, and VNB) (Meyer *et al.* 2009b). The new

species definition includes VNI, VNII, and VNB within *C. neoformans*, VNIV within *C.*

deneoformans, and VNIII as *C. neoformans*/*C. deneoformans* hybrids. VNI is the most common

C. neoformans molecular subtype and has a global distribution. The VNB subtype was discovered

in Africa and has a genetically variable population with 10% *MAT***a** isolates, which are very rare

in all other studied populations (Litvintseva *et al.* 2003). The VNB clade includes strains isolated

from the endemic mopane tree (*Colophospermum mopane*), whereas strains isolated from

pigeon excreta in the same area had the VNI molecular type, which is shared by most isolates

globally. This observation supported the "out of Africa" hypothesis in which few of these isolates

would have adapted their biology to multiply in pigeons to achieve global dispersion (Litvintseva

et al. 2011; Litvintseva and Mitchell 2012).

More recently, the improvement of sequencing technologies and bioinformatics tools allowed

analysis of whole genome sequences, revealing new details in the structure and global

geographical distribution of the *Cryptococcus* populations together with the identification of

additional clades and subclades (Farrer *et al.* 2019). Most of these studies focused on *C.*

neoformans, with >1000 *C. neoformans* genome sequences analyzed in four studies

(Desjardins *et al.* 2017; Rhodes *et al.* 2017b; Vanhove *et al.* 2017; Ashton *et al.* 2019; Ergin *et*

al. 2019). Through these efforts, the VNB clade has been separated into the VNBI and VNBII

subclades, which have different resistance to oxidative stress and a different ability to produce

melanin (Desjardins *et al.* 2017; Vanhove *et al.* 2017). The VNI clade has been separated into a

number of subclades with three phylogenetic groups representing the majority of the studied

isolates (Rhodes *et al.* 2017b; Ashton *et al.* 2019). Surprisingly, only 11 haploid strains isolated

in South America were included in these studies. This small number conflicts with the

relevance of South America as a source of genetic diversity in the *gattii* isolates (Engelthaler *et*

al. 2014). Up until now, the only analysis of *C. neoformans* population structure in South

America has been an MLST analysis that concluded that subtype ST93 was the most prevalent

among strains isolated from HIV patients in Southeastern Brazil with a low level of

recombination (Ferreira-Paim *et al.* 2017). Similarly, few phenotypic analyses that include

South American isolates have been published, and a recent analysis of virulence factor

expression within Brazilian *Cryptococcus* clinical isolates does not include any sequencing or

molecular typing analysis (De Sousa *et al.* 2020). In the present study, we analyzed the

phenotypes and genome sequences of 53 Brazilian isolates, revealing very high phenotypic

diversity within *Cryptococcus* isolates and a subclade distribution that suggests several waves

of *Cryptococcus* introduction from Africa to South America.

Material and Methods

Isolates used in this study

Strains 2B, 2C, 2D, 4C, 7, 33A1, 54A1, 56A1, 68B1, 69A1, 81A1, 96A1, 106A1, 144A1, 151A1,

176A1, 3Pb3, 5Pb2, 19Pb4, 23Pb2, 32Bp1, and RR 2605 were isolated from Northern Brazil (Dal

Pupo *et al.* 2019) and kindly donated by Dr. Halan Dal Pupo. Strains Cn201, Cn359, Cn894,

Cn14HC, and Cn201 were obtained from the culture collection of the Central Hospital of the

University of São Paulo. All other isolates were obtained from the Collection of Pathogenic

Fungi available at Fiocruz. No genomic information was available for any of these isolates

before the present study.

Read alignment, variant detection, and ploidy analysis

Illumina reads were aligned to the *Cryptococcus* reference genomes using Minimap2 aligner v.

2.9 (Li 2018) with the "-ax sr" parameter. BAM files were sorted and indexed using SAMtools

(Li *et al.* 2009) version 1.9. Picard version 2.8.1 (http://broadinstitute.github.io/picard) tools

were used to identify duplicate reads and assign correct read groups to BAM files. SAMtools

version 1.9 and Picard version 2.8.1 were then used to filter, sort, and convert SAM files and

assign read groups and mark duplicate reads. Single-nucleotide polymorphisms (SNPs) and

insertions/deletions (indels) were called using Genome Analysis Toolkit version 3.6 with

ploidy=1 according to the GATK Best Practices. HaploScore parameters used to filter SNPs and

indels included VariantFiltration, QD <2.0, LowQD, ReadPosRankSum<-8.0, LowRankSum, FS

>60.0, HightFS, MQRankSum<-12.5, MQRankSum, MQ <40.0, LowMQ, and HaplotypeScore

>13.0. Phyml version 20120412 (Guindon *et al.* 2010) with the GTR, NNIs substitution model

was used to infer phylogenetic relationships between the isolates using the separate dataset

of confident SNPs. To examine variations in ploidy across the genome, the sequencing depth at

all positions was computed using SAMtools (Li *et al.* 2009), and then the average depth was

computed for 1-kb windows across the genome.

MAT locus determination

The mating type of Brazilian isolates was determined as previously published (Rhodes *et al.*

2017b). Briefly, Illumina reads were aligned to the mating-type locus sequences [AF542529.2

and AF542528.2] using BWA-MEM. Computed average depths at the *SXI* and *STE20* loci using

SAMtools mpileup were used to determine the mating type.

Population inference by fastSTRUCTURE

We used fastStucture (Raj *et al.* 2014) to identify admixture in the dataset from K=2 to K=4.

After K=4, clusters with the highest number of isolates were split into sub-clusters, which likely

reflects an issue due to the number of isolates within clusters rather than true biological

significance.

Drug susceptibility testing

The minimum inhibitory concentration (MIC) of fluconazole (FLC) (Sigma-Aldrich) was

determined as described by M27-A3 (Institute 2017).

Titan cell production

Titan cell (TC) production was evaluated as described by Dambuza and colleagues with some

modifications (Dambuza *et al.* 2018). The isolates were cultivated in yeast peptone dextrose

(YPD) agar (1% yeast extract, 2% bacto-peptone, 2% glucose, 2% bacto-agar) for 48h at 30°C. A

single colony was transferred to 5 mL YNB without amino acids (Sigma Y1250) prepared

according to the manufacturer's instructions plus 2% glucose. After incubation overnight at 30°C

and 200 rpm, $1X10^3$ cells/mL were inoculated into 5 mL 1xPBS (phosphate buffer solution) + 10%

heat-inactivated (HI) fetal calf serum (FCS) and incubated for 72 h at 37°C, 5% CO_2. We visualized

the cells microscopically without any staining, and 50 cells were measured. TCs were considered

body cells (without the capsule) with a diameter >10 μm. The experiment was repeated twice.

Capsule staining

Fungal cells were cultivated in YPD agar for 24 h at room temperature. Single colonies were

transferred by sterile loop to the wells of a 96-well plate, each containing 200 μL of Roswell

Park Memorial Institute (RPMI) medium at a final cell density of 10^7 yeast/mL. The plates were

incubated for an additional 24 h at 37°C in 5% CO_2 and centrifuged to separate the cells from

the supernatants. The supernatants were stored for GXM analysis, and the cells were washed

twice in PBS for further fixation in 4% paraformaldehyde for 1 h at room temperature. Yeast

cells were then blocked (1% bovine serum albumin [BSA] in PBS for 1 h at 37°C) and incubated

with the 18B7 anti-GXM monoclonal antibody (Casadevall *et al.* 1992) (mAb 18B7; 10 µg/mL

for 1 h at 37°C), a kind gift from Dr. Arturo Casadevall (Johns Hopkins University). The cells

were washed and incubated with an Alexa Fluor 488-conjugated secondary antibody (10

µg/mL for 1 h at 37°C; Invitrogen, USA). The cells were finally stained with calcofluor white (5

µg/mL for 1 h at 37°C; Invitrogen, USA). Alternatively, the cells were counterstained with India

ink for capsular visualization. The cells were visualized on a Leica TCS SP5 confocal microscope

or analyzed with a FACS Canto II flow cytometer. Data were processed with the FACSDiva

software, version 6.1.3.

Data availability

All sequence data from this study have been submitted to GenBank under the BioProject

identification no. PRJNA702892.

Results

Description of the origin of the Brazilian isolates

We selected 53 *Cryptococcus* Brazilian isolates, of which 12 were classified as *C. gattii* and 41

as *C. neoformans.* More than 41% (n=22) were environmental isolates and 36% (n=19) were

clinical isolates. The origin of the isolate was unknown for the remaining 23% (n=12). Twenty-

one strains were isolated from pigeon excreta in Northern Brazil (Dal Pupo *et al.* 2019), 28

isolates were obtained from the culture collections available at Fiocruz, and 4 were acquired

from the Central Hospital of the University of São Paulo (Table S1).

Genome characterization

Whole genome sequencing was performed for each of the 53 Brazilian isolates considered in

this study. Between 2.7 and 10.9 million 100-nt paired-end reads were obtained and aligned to

eight *Cryptococcus* reference genomes using Minimap2 (Li 2018) (Table S1). Reference

genomes were *C. neoformans* strain H99 (Janbon *et al.* 2014), *C. deneoformans* strain JEC21

(Loftus *et al.* 2005; Gonzalez-Hilarion *et al.* 2016), *C. deuterogattii* strain R265 (Yadav *et al.*

2018; Gröhs Ferrareze *et al.* 2021), *C. gattii WM276*, *C. bacillisporus* strains CA1280 and

CA1873, *C. decagattiii* strain AFLP10, and *C. tetragattii* strain IND107 (Basenko *et al.* 2018). Of

the 53 isolates, 40 were identified as *C. neoformans* with >80% of the reads aligning to the H99

genome (<40% of reads could be aligned to any other reference genome). Among the 13

remaining isolates, 8 were *C. deuterogattii* and 5 were *C. gattii* (Table S1). Of note, we found

that isolate Cg366, which was previously identified as *C. gattii*, is actually a *C. neoformans*

isolate. Isolates Cn201 and Cn894, which were previously considered to be *C. neoformans*,

were identified as *C. deuterogattii* isolates.

In order to determine mating type, we compared the number of reads aligned to the *STE20a*,

STE20alpha, *SXI1*, and *SXI2* genes of the corresponding species. These analyses revealed that

all isolates were *MATα*, confirming the overwhelming dominance of this mating type in clinical

and environmental isolates observed in most areas in the world (Desjardins *et al.* 2017).

The analysis of the read alignment profiles provided insight into possible large duplications or

aneuploidies. All *C. neoformans* isolates displayed an even alignment profile. This observation

together with the absence of heterozygous SNPs in our analysis suggested that these isolates

are either haploid or euploid homozygotes. However, we observed a duplication of the end of

the chromosome 1 in three isolates (Cn160, Cn161, and Cn186)) (left side in Figure 1). The

isolate Cn161 apparently had additional duplications within chromosome 8. Similar patterns

were observed in the eight *C. deuterogattii* isolates, with two isolates (Cn201 and Cn894)

displaying a duplication of a region located in chromosomes 8 and 13, respectively. In contrast,

all five *C. gattii* isolates displayed an even alignment profile, suggesting that all are haploid

(data not shown).

Figure 1. Large segmental duplications within the genomes of Brazilian isolates were revealed
after aligning sequencing reads to the *Cryptococcus* reference genomes.
226

Phylogenetic analyses

To obtain insight into the population structure of the Brazilian *C. neoformans* isolates, we

constructed a phylogenetic tree including 352 isolates considered in previous studies

(Desjardins *et al.* 2017; Rhodes *et al.* 2017b; Ashton *et al.* 2019) that encompass all VNI and

VNB subclades (Table S2). We also included 10 VNII isolates as the outgroup to root the tree

(Desjardins *et al.* 2017; Rhodes *et al.* 2017b; Ashton *et al.* 2019). Overall, 945,565 variable

positions were used to construct the tree presented in Figure 2A.

VNB isolates

Whereas the vast majority (92.5%, n=37) of the *C. neoformans* Brazilian isolates belonged to

the VNI clade, we also identified one VNB isolate of the VNBII subclade (Cn225). Two other

Brazilian isolates (Cn216 and Cn14HC) appear to be associated with the VNBI and VNBII clades,

respectively, but with a close-to-the-root long branch (Figure 2A). Accordingly, the number of

SNPs compared to the H99 reference genome was slightly higher in these two isolates than in

the other VNB isolates (Figure 2B). As previously reported, analysis of the population using

fastSTRUCTURE (k=4) (Raj *et al.* 2014) readily separated VNI, VNII, VNBI, and VNBII isolates

(Desjardins *et al.* 2017) into four independent populations (Figure 2C). As expected, this

analysis showed that the two Brazilian outlier isolates (Cn216 and Cn14HC) are associated with

the VNBI and VNBII clades, respectively.

246
247

Figure 2. (A) Phylogenetic tree generated by iTOL v5 using *C. neoformans* genome sequences.
The Brazilian isolates are indicated in red. The two Brazilian outlier isolates are shown. (B)
Number of SNPs within the genomes of the different isolates analyzed in this study. (C) Results
of the structure analysis (k=4) suggesting VNBI and VNBII origin for the two Brazilian outlier
isolates (CnH14C and Cn216, respectively).

Because VNI/VNB and VNII/VNB hybrids have been previously reported within African isolates

(Rhodes *et al.* 2017b), we sought to compare their phylogenetic position with the two outlier

Brazilian isolates. We first redrew a phylogenetic tree including six of these isolates (the

VNII/VNB isolate MW-RSA852 and four VNI/VNB isolates [Bt125, Bt131, Bt162, and Bt163])

(Figure S1). As previously published, the African hybrid isolates are also positioned at the root

of the VNB and VNII branches, respectively. We then performed principal component analysis

(PCA) including this set of hybrid isolates. As previously published, this analysis readily

separated isolates positioned at an intermediary position between the defined clades (Figure

3) (Desjardins *et al.* 2017). Like most African VNI/VNB hybrids, the Bt163 isolate was

positioned between VNI and VNBII, suggesting that it is a hybrid of isolates of these two clades.

Of note, the VNI/VNB isolate Bt125 has an intermediary position between the VNBI, VNBII, and

VNI lineages (Figure 3).

Figure 3. Principal component analysis revealing the intermediary position of the Brazilian
outlier isolates. VNI/VNB and VNII/VNB hybrid isolates (Rhodes *et al.* 2017b) have been added
to this analysis. The positions of the hybrid (black) and of the two Brazilian outlier isolates (in
red) are indicated.
271
The two outlier Brazilian isolates are positioned between the VNBI and VNBII isolates,

suggesting that they are not VNB/VNI hybrids. In agreement with the location of these isolates

in the phylogenetic tree, this analysis showed that the Cn216 isolate is closer to VNBII isolates

whereas the Cn14HC isolate is closer to VNBI isolates. According to these results, Cn216 and

CnH14C could be either VNBI/VNBII hybrids or evolved VNBI and VNBII isolates, respectively. In

order to test these two hypotheses, we first determined the position and number of SNPs

specific to each clade (Figure 4A,B). This analysis revealed that the number of clade-specific

SNPs was equivalent for each clade with an even distribution along the H99 reference genome.

Of note, the VNBI clade appear to have fewer clade-specific SNPs, which might be related to

the previously reported loss of diversity in this lineage (Desjardins *et al.* 2017).

Figure 4. (A) Number of specific SNPs in each clade and in the two Brazilian outlier isolates
(Cn216 and Cn14HC). The Brazilian VNBII isolate Cn225 and the hybrid isolates Bt125 and
Bt163 (Rhodes *et al.* 2017b) were used as controls. (B) Density and position of the clade-
specific SNPs in the H99 reference genome. The results obtained with chromosome 14 are
presented.

We reasoned that we could determine the origins of the SNPs in the two outlier Brazilian

isolates by comparing them with clade-specific SNPs (Figure 5). We compared the SNPs of

Cn216 and Cn14HC SNPs (n=213,915 and n=207,438, respectively) with the VNI-, VNII-, VNBI-,

and VNBI-specific SNPs and plotted their positions in the H99 reference genome. As a control,

we performed the same analysis with the VNBII Brazilian isolate Cn215 (n=207,639 SNPs).

Figure 5. Origin of the SNPs in the Brazilian outlier isolates (Cn216 and Cn14HC). Positions of
the SNPs shared between the clade-specific SNPs were identified and plotted on each
chromosome. Chromosome 14 is presented as an example. Results obtained with the Brazilian
VNBII Cn225 isolate and the African VNB/VNI hybrid isolates Bt125 and Bt163 (Rhodes *et al.*
2017b) are shown as controls.

This analysis revealed that the Brazilian outlier isolates Cn214 and Cn14HC mostly share SNPs

with the VNBII- and VNBI-specific SNPs, respectively, confirming that they belong to these

respective lineages. We observed an even distribution of these SNPs along the chromosomes

with no sign of a recent recombination event between the VNBI and VNBII lineages. As

expected, a similar result was obtained with the Brazilian VNBII isolate Cn225. In contrast, the

African VNB/VNI hybrid Bt163 displays a SNP distribution common to the clade-specific SNPs,

supporting a recent recombination event between a VNBII isolate and a VNI isolate and thus

confirming the PCA results and previously published results (Rhodes *et al.* 2017b). For the

Bt125 isolate, the SNP pattern was more complex than previously published, suggesting a

triple VNBI/VNBII/VNI origin for this isolate, in agreement with the intermediary position of

this isolate observed by PCA (Figure 3). Overall, these analyses suggest that the Cn214 and

CnH14C isolates belong to the VNBII and VNBI clades, respectively, with no sign of

recombination with another clade. Yet, the number of the SNPs unique to these two isolates

(n=37,236 in Cn216 and n=26,925 in CnH14C) is much higher than in the VNBII Brazilian isolate

Cn225 (n=4788) and in the two VNI/VNB African hybrids (n=7485 in Bt125 and n=3511 in

Bt163) (Figure 4A). These isolate-specific SNPs are evenly distributed along the H99 reference

genome (Figure 4B). This SNP pattern may be due to an ancient separation of these isolates

from most other isolates of their respective clades or a recent but rapid evolution of their

genome sequences due to an increased mutation rate.

In order to test these hypotheses, we compared the mutation rates in both Cn216 and Cn14HC

by observing the appearance of spontaneous 5-fluorouracil (FU)-resistant colonies in

fluctuation assays (Boyce *et al.* 2017; Billmyre *et al.* 2020). We included the reference strain

KN99α and the Brazilian VNBII isolate Cn225 as controls. The mutation rate of Cn14HC was

higher than those of Cn225 and KN99α (Figure 6). In contrast, the mutation rate in Cn216 was

not significantly different from that of Cn225 (VNBII). Similar results were obtained when 5-

FOA selection was used (Figure S2).

Figure 6. CnH14C display an elevated mutation rate. (A) Spontaneous 5-FU–resistant colonies
in a standard wild type (KN99α) isolate and three Brazilian isolates (Cn216, Cn225, and
CnH14C). (B) Quantitative assessment of mutation rates in the four strains using fluctuation
analysis of the spontaneous resistance to 5-FU. The Lea-Coulson method of the median was
used to estimate the number of mutations from the observed values of mutants from 10
independent, parallel cultures (Foster 2006; Gillet-Markowska *et al.* 2015). Error bars indicate
95% confidence intervals.

Isolates with mutations in the mismatch repair pathway associated with a mutator phenotype

have been observed both in *C. deuterogattii* (Billmyre *et al.* 2017) and *C. neoformans* clinical

isolates (Boyce *et al.* 2017; Rhodes *et al.* 2017a). Although experimental in vitro analyses

suggested that only *MSH2*, *MLH1*, and *PMS1* play a major role in regulating the mutation rate

in *C. neoformans* (Boyce *et al.* 2017), recurrent isolate evolution analysis suggested that *MSH5*

and *RAD5* could also influence isolate evolution in patients (Rhodes *et al.* 2017a). Given that

we did not identify Cn14HC nonsense mutations, we looked at the Cn14HC-specific SNPs

responsible for non-conservative mutations in these mismatch repair genes. Interestingly, two

of these genes contain mutations specific to Cn14HC: *PMS1* (CNAG_07955) encodes a protein

with a S483P mutation and *RAD5* (CNAG_04733) encodes a protein with a P332S mutation.

These SNPs were not present in any of the other 391 isolates considered in this study. In

contrast, we did not identify specific SNPs responsible for a change in any of these proteins in

Cn216 or Cn225.

Overall, our results suggest that CnH14C possesses a mutator phenotype associated with

specific mutations in two mismatch repair genes. Of course, more experiments are needed to

formally establish a causal relationship between this genotype and this phenotype.

Nevertheless, the mutator phenotype of CnH14C has resulted in its rapid evolution, and could

explain its position in the phylogenic tree. In contrast, Cn216 might be the result of a long

evolution independent of the African VNBII isolate due to an ancient geographical separation.

VNI isolates

As stated above, nearly all *C. neoformans* isolates in this collection belong to the VNI clade,

mostly within the VNIa subclade. Only two isolates were identified as VNIb and none were

identified as VNIc. Interestingly, the second major lineage within the VNIa Brazilian isolates

was VNIa_Y, encompassing 27.5% (n=10) of isolates. All of these strains were isolated from

pigeon droppings in Northern Brazil. These are the first VNIa_Y strains isolated outside of

Africa; previous strains, all of which originated in the environment, were isolated in Botswana,

Malawi, and Uganda (Ashton *et al.* 2019). The population structure of the Brazilian VNIa

isolates resembles that of the African isolates. Approximately 45% of the VNIa isolates are of

the VNIa_93 lineage, similar to previous observations in Ugandan isolates (Ashton *et al.* 2019).

Similar to previous reports for African countries, only 2 of the Brazilian isolates were of the

VNIa_5 subclass and no VNIa_4 isolates were identified, confirming that these two subclasses

are specifically enriched within Southeast Asia isolates (Ashton *et al.* 2019).

Phylogenetic trees constructed using only the VNIa subclades revealed a clear geographical

clustering of the isolates (Figure 7), suggesting that the current subclade organization of *C.*

neoformans isolates predates the dissemination into South America. However, the existence of

unclustered VNI-93 Brazilian isolates suggests multiple intercontinental isolate transfers.

Figure 7. Phylogenetic trees of the VNIa subclades as visualized by ITOL revealed geographic
clustering. The Brazilian, African, and South Asian isolates are labelled in red, blue, and yellow,
respectively. The Indian isolates are labelled in purple.
377

Phenotypic analyses

We next characterized the Brazilian isolates at a phenotypic level. We focused our analysis on

virulence-associated phenotypes and drug resistance. All isolates grew well at 30°C, and only *C.*

gattii isolate Cg187 and *C. neoformans* VNIa_Y isolate 3Pb3 exhibited a growth defect at 37°C

(Figure S3). Similarly, most of these *Cryptococcus* isolates demonstrated a low MIC for

fluconazole (Table S3), with a MIC_{50} of 2 µg/mL and MIC_{90} of 8 µg/mL. The clinical isolate

Cn191 (VNIa_93) and the environmental isolate 151A1 (VNIa_Y) showed the highest MIC

values (8.0 µg/mL).

Capsule formation and exportation of GXM, a major capsular antigen, are essential for

Cryptococcus virulence. We analyzed multiple aspects of capsule formation in Brazilian

isolates, including capsular morphology, serological reactivity, and extracellular concentration

of GXM. Capsule morphology was examined by fluorescence microscopy after incubation of

the cells with an antibody to GXM for capsule visualization and calcofluor white for cell wall

staining. Alternatively, the capsule was counterstained with India ink for visualization of the

capsule by light microscopy. This analysis revealed formidable heterogeneity at morphological

and serological levels (Figure 8).

Figure 8 (A) Microscopic analysis of the cell surface of the Brazilian isolates. R265 and H99 are
standard cryptococcal isolates that were used as reference in this assay. For each isolate,
fluorescence panels are shown on the left and India ink counterstaining is shown on the right.
Red fluorescence indicates capsule staining with the mAb 18B7 antibody and blue fluorescence
represents cell wall staining. Images are illustrative of two independent analyses that

produced similar results. Corresponding data for the Northern Brazil isolates have been
previously published (Dal Pupo *et al.* 2019). (B) Flow cytometry analysis of the cryptococcal
isolates after capsule staining with mAb 18B7. At least 5000 cells were analyzed for each
isolate. Results are presented as average values of fluorescence intensity and are illustrative of
two independent analyses producing similar results. R265 and H99 are standard cryptococcal
isolates that were used as reference in this assay. Data for the Northern Brazil isolates were
obtained from (Dal Pupo *et al.* 2019).
407
408
On the basis of the visual perception that antibody recognition of the capsule was variable, we

measured the indices of fluorescence intensity after incubation of each isolate with the GXM-

binding antibody by flow cytometry (Figure 8B). Using the standard isolates H99 and R265 as

reference strains, this analysis confirmed both highly reactive isolates and poorly reactive

isolates, supporting the notion that surface GXM is highly diverse in *Cryptococcus*. Although *C.*

gattii and *C. deuterogattii* isolates were predominantly less capsulated than *C. neoformans*

(student t-test *P* value <0.05), we did not observe any statistical difference between the *C.*

neoformans subclades.

We also analyzed the concentration of polysaccharide in culture supernatants of each isolate

(Figure S4). Detection of extracellular and surface-associated GXM showed an apparent

correlation, as suggested by the statistical analysis (Figure S4), supporting the notion that

cryptococcal isolates are highly variable in their ability to form GXM-containing capsules and to

export polysaccharide.

We also examined melanin production on L-Dopa medium. We observed a variable phenotype

among the Brazilian isolates, with *C. gattii* and *C. deuterogattii* isolates displaying lower

melanin production (Figure 9). There was no obvious correlation between melanin production

among subclades, although strains isolated from bird droppings appear to produce more

melanin than other environmental or clinical isolates.

Figure 9. Melanin production of Brazilian isolates as estimated on L-Dopa medium after 72 h of incubation. Strains isolated from bird droppings are indicated by a yellow box.

Lastly, we examined production of TCs. *C. neoformans* and *C. gattii* cells can undergo an unusual transition in the host lung from the typical size of 5 to 7 µm to between 10 µm and 100 µm, a phenomenon known as titanization. TCs are known to be highly polyploid (Zaragoza and Nielsen 2013). TCs can also be induced in vitro using three different approaches (Dambuza *et al.* 2018; Hommel *et al.* 2018; Trevijano-Contador *et al.* 2018). Here, we followed the protocol described by Dambuza and colleagues (2018) to study whether Brazilian *Cryptococcus* isolates produce TCs. In our study, *C. neoformans* H99 formed nearly 17% TCs, aligning with previously published findings (Dambuza *et al.* 2018). Approximately 49% of Brazilian *Cryptococcus* isolates produced TCs (defined here as cells with a size >10 µm) (Figure 10, Table S4). Around 24% of Brazilian *Cryptococcus* isolates generated more TCs than *C. neoformans* H99 (Figure 10, Table S4). Interestingly, and in contrast to a previous study using the same protocol, all Brazilian *C. gattii* and *C. deuterogattii* isolates tested formed TCs (Figure 10, Table S4). The highest percentage of TCs (97.33%) was produced by *C. deuterogattii* isolate Cn894 (Figure 10, Table S4).

In order to address whether cells that formed more TCs than H99 are naturally larger, we compared the size of these cells before and after the titanization induction. The mean size of the cells of Brazilian *Cryptococcus* isolates varied between 4.18 to 5.70 µm before and 6.85

to 14.19 µm after cultivation in TC-inducing conditions, proving that the titanization induction

was responsible for the increase in cell size (Table S5). We did not observe any link between

subclade and the ability of the corresponding isolate to undergo this phenotypic change.

Figure 10. Cell diameter after titanization induction in Brazilian isolates. Cryptococcus isolates
were subjected to TC induction in PBS-HI-FCS. Cells with a diameter >10 µm were considered
to be TCs. ND: not determined.
454
455   Overall, this analysis revealed spectacular phenotypic variability within the Brazilian isolates,

456   although no association was detected between one clade and any of these phenotypic traits.

Discussion

Population genomics studies have resulted in a model in which clonal reproduction has shaped the global population structure of *C. neoformans* isolates, which are mostly of the VNI clade and of the *MATα* sexual type (Cuomo *et al.* 2018). These genome sequence analyses have also identified several VNI subclades with specific geographical distribution (Ashton *et al.* 2019). Currently, little is known about *Cryptococcus* species diversity in South America because only a few isolates have been sequenced (Rhodes *et al.* 2017b). In the present study, we analyzed the largest South American genome collection studied to date. As expected, most of the Brazilian isolates belonged to the VNI clade. Our analysis revealed that the VNI Brazilian *C. neoformans* population shares many features with the African population. For instance, VNI_Y isolates that were previously reported only in Africa were also prominent within this set of South American isolates. As in Africa, Brazilian VNIa_5 isolates were rare; in contrast, VNIa_5 isolates represent one of the largest subclades within Asian isolates (Ashton *et al.* 2019). Actually, in most regards, the *C. neoformans* Brazilian population structure is very different from the Asian one described by Ashton and colleagues in 2019 (Ashton *et al.* 2019). A detailed analysis of the different VNIa subclade population structures further supported the hypothesis of repeated and privileged exchanges from African to South American isolates. Thus, South American isolates do not form an independent clade or subclade but rather are scattered within nearly all global subclades, indicating repeated transport. Although we observed a perfect clustering of Brazilian isolates within the VNIa_Y and VNIa_32 subclades—possibly from an ancient importation followed by independent evolution of the ancestral isolate—some VNIa-93 isolates clustered with the other global isolates, suggesting independent and possibly more recent importations. Of course, because, all these unclustered VNI-93 Brazilian strains are clinical isolates, we cannot completely exclude the hypothesis that these patients had travelled to Africa, and had been infected there, before developing a cryptococcosis in Brazil as previously described in some rare cases (Garcia-Hermoso *et al.* 1999).

A similar pattern was observed with the VNB isolates. Two of the Brazilian isolates were

positioned close to the roots of the VNB branches, as previously observed for two sequenced

Brazilian isolates (Rhodes *et al.* 2017b). However, the third isolate clustered within the other

VNB African isolates, suggesting a more recent intercontinental exchange. In that case, it is

important to note that this last strain has been isolated in the environment.

In 2017, Casadevall and colleagues suggested that the separation of *C. neoformans* complex

species and *C. gattii* complex species occurred in the middle of the Cretaceous period

(Casadevall *et al.* 2017). This geological period was marked by the breakup of the

supercontinent west Gondawa which encompassed Africa and South America. This modern

continent formation ended much before the global dispersion of major *Cryptococcus* spreader

birds (Prum *et al.* 2015) which occurred 40×10^6 years after the end of the Cretaceous period.

Although our data cannot reliably date the separation between the *C. neoformans* subclades,

the similar structure of the African and South American population suggests an African origin

for every *C. neoformans* isolates. In this model, the transport of *C. neoformans* from Africa to

South America started only with birds migrations 25×10^6 years ago and continued until today.

This study also exemplified the impact of rapidly evolving isolates on the *C. neoformans*

population structure. Mutator phenotypes have already been observed in natural *C.*

neoformans and *C. gattii* isolates and have been shown to impact the dynamics of isolate

evolution, phenotypic diversity, and drug resistance (Billmyre *et al.* 2017; Billmyre *et al.* 2020).

A *msh2*-associated mutator phenotype was also observed in a subset of *C. deuterogattii* Pacific

Northwest outbreak isolates, named VGIIa-like (Kidd *et al.* 2004; Billmyre *et al.* 2014; Billmyre

et al. 2017). Here, the VNBI isolate CnH14C displayed a mutator phenotype, albeit less

pronounced than the previously described *C. neoformans msh2Δ*, *mlh1Δ*, and *pms1Δ* mutant

strains (Boyce *et al.* 2017). Accordingly, we did not identify nonsense mutations in any of these

genes in the CnH14C isolate. However, *PMS1*- and *RAD5*-specific alleles in CnH14C might

suggest a causal relationship. In fact, *RAD5* has been recently identified as regulating mutation

rates in a quantitative trait locus analysis in *S. cerevisiae* (Gou *et al.* 2019), and a specific *RAD5*

mutation has also been identified in a rapidly evolved *C. neoformans* recurrent clinical isolate

(Rhodes *et al.* 2017a).

However, in contrast to what has been observed in some African isolates, we did not identify

any hybrid isolates within this collection of Brazilian isolates. As previously reported

(Matsumoto *et al.* 2007; Andrade-Silva *et al.* 2018), none of these isolates were *MAT***a**.

Nevertheless, our model of multiple intercontinental transport of African isolates to South

America and the proportion of *MAT***a** isolate within the VNB isolates in Africa (Litvintseva *et al.*

2007; Litvintseva *et al.* 2011; Litvintseva and Mitchell 2012) predict that additional sampling

will probably identify such isolates.

MLST and genome sequence analysis identified South America, and in particular Brazil, as the

major source of the global genetic diversity of the *C. gattii* complex, especially within the VGII

and VGIII lineages (Hagen *et al.* 2013; Billmyre *et al.* 2014; Souto *et al.* 2016; Barcellos *et al.*

2018; Vilas-Bôas *et al.* 2020). This is associated with great phenotypic diversity within the

Brazilian isolates (Silva *et al.* 2012; Barcellos *et al.* 2018). Similarly, genotypic and phenotypic

analyses of Brazilian *C. neoformans* isolates, including the analysis in the present study, have

revealed great diversity within South American isolates (Ferreira-Paim *et al.* 2017; Andrade-

Silva *et al.* 2018; Dal Pupo *et al.* 2019; De Sousa *et al.* 2020). Nevertheless, our data suggest

that Brazil is probably not an important source of global diversity of *C. neoformans* isolates.

Rather, our findings support the model in which this diversity is mostly imported from Africa

through repeated intercontinental transport of *C. neoformans* isolates to South America.

Our studies continue the investigation of *Cryptococcus* species global diversity and emphasize

the importance of understanding the evolution of fungal pathogens. Further studies with

additional isolates from Brazil and other countries in South America will help us to clarify many

aspects of population genetics of this important fungal pathogen.

Acknowledgments

M.L.R. is currently on leave from the position of associate professor at the Microbiology

Institute of the Federal University of Rio de Janeiro, Brazil. M.L.R., G.J., and G.H.G. were

supported by a Tripartite USP-Fiocruz-Pasteur grant. FCGR was financed in part by a

scholarship from the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES,

Brazil, Finance Code 001). This work in the Janbon lab was supported in part by an Infect-ERA

grant (project Cryptoview).

References
546

547   Alanio, A., 2020 Dormancy in *Cryptococcus neoformans*: 60 years of accumulating
548   evidence. The Journal of Clinical Investigation 130**:** 3353-3360.
549   Aminnejad, M., M. Diaz, M. Arabatzis, E. Castaneda, M. Lazera *et al.*, 2012
550   Identification of novel hybrids between *Cryptococcus neoformans* var. *grubii*
551   VNI and *Cryptococcus gattii* VGII. Mycopathologia 173.
552   Andrade-Silva, L. E., K. Ferreira-Paim, T. B. Ferreira, A. Vilas-Boas, D. J. Mora *et al.*,
553   2018 Genotypic analysis of clinical and environmental *Cryptococcus*
554   *neoformans* isolates from Brazil reveals the presence of VNB isolates and a
555   correlation with biological factors. PLOS ONE 13: e0193237.
556   Ashton, P. M., L. T. Thanh, P. H. Trieu, D. Van Anh, N. M. Trinh *et al.*, 2019 Three
557   phylogenetic groups have driven the recent population expansion of
558   *Cryptococcus neoformans*. Nature Com. 10: 2035.
559   Barcellos, V. A., L. M. S. Martins, A. C. L. Fontes, J. C. V. Reuwsaat, E. D. Squizani *et al.*,
560   2018 Genotypic and phenotypic diversity of *Cryptococcus gattii* VGII Clinical
561   isolates and its Impact on Virulence. Frontiers in microbiology 9: 132-132.
562   Basenko, E. Y., J. A. Pulman, A. Shanmugasundram, O. S. Harb, K. Crouch *et al.*, 2018
563   FungiDB: An integrated bioinformatic resource for Fungi and Oomycetes. J.
564   Fungi 4: 39.
565   Billmyre, R. B., S. Applen Clancey, L. X. Li, T. L. Doering and J. Heitman, 2020 5-
566   fluorocytosine resistance is associated with hypermutation and alterations in
567   capsule biosynthesis in *Cryptococcus*. Nature Com. 11: 127.
568   Billmyre, R. B., S. A. Clancey and J. Heitman, 2017 Natural mismatch repair mutations
569   mediate phenotypic diversity and drug resistance in *Cryptococcus deuterogattii*.
570   eLife 6**:** e28802.
571   Billmyre, R. B., D. Croll, W. Li, P. Mieczkowski, D. A. Carter *et al.*, 2014 Highly
572   recombinant VGII *Cryptococcus gattii* population develops clonal outbreak
573   clusters through both sexual macroevolution and asexual microevolution. mBio
574   5**:** e01494-01414.
575   Boekhout, T., B. Theelen, M. Diaz, J. W. Fell, W. C. J. Hop *et al.*, 2001 Hybrid genotypes
576   in the pathogenic yeast *Cryptococcus neoformans*. Microbiology 147: 891-907.
577   Bovers, M., F. Hagen, E. E. Kurumae, H. L. Hoogveld, F. Dromer *et al.*, 2008 AIDS patient
578   death caused by novel *Cryptococcus neoformans* x *C. gattii* hybrid. Emerg.
579   Infect. Dis. 14: 1105-1108.
580   Boyce, K. J., Y. Wang, S. Verma, V. P. S. Shakya, C. Xue *et al.*, 2017 Mismatch repair of
581   DNA replication errors contributes to microevolution in the pathogenic fungus
582   *Cryptococcus neoformans*. mBio 8: e00595-00517.
583   Bui, T., X. Lin, R. Malik, J. Heitman and D. Carter, 2008 Isolates from *Cryptococcus*
584   *neoformans* from infected animals reveal genetics exchange in unisexual, α
585   mating type population. Eukaryot. Cell 7**:** 1771-1780.
586   Casadevall, A., J. B. Freij, C. Hann-Soden and J. Taylor, 2017 Continental drift and
587   speciation of the *Cryptococcus neoformans* and *Cryptococcus gattii* species
588   complexes. mSphere 2**:** e00103-00117.

Casadevall, A., J. Mukherjee, S. J. N. Devi, R. Schneerson, J. B. Robbins *et al.*, 1992
Antibodies elicited by a *Cryptococcus neoformans*-tetanus toxoid conjugate
vaccine have the same specificity as those elicited in infection. The Journal of
Infectious Diseases 165**:** 1086-1093.
Cuomo, C. A., J. Rhodes and C. A. Desjardins, 2018 Advances in *Cryptococcus* genomics:
insights into the evolution of pathogenesis. Mem. Inst. Oswaldo Cruz 113.
Dal Pupo, H. D., B. A. G. Sena, F. C. G. Reis, L. Machado, S. T. Fortes *et al.*, 2019
Polysaccharide diversity in VNI isolates of *Cryptococcus neoformans* from
Roraima, Northern Brazil. Fungal Biology 123**:** 699-708.
Dambuza, I. M., T. Drake, A. Chapuis, X. Zhou, J. Correia *et al.*, 2018 The *Cryptococcus*
neoformans Titan cell is an inducible and regulated morphotype underlying
pathogenesis. PLoS Pathog. 14**:** e1006978.
de Sousa, H. R., G. P. de Oliveira, S. de Oliveira Frazão, K. C. de Melo Gorgonha, C. P.
Rosa *et al.*, 2020 Faster *Cryptococcus* melanization increases virulence in
experimental and human cryptococcosis. bioRxiv**:** 2020.2007.2029.222794.
Desjardins, C. A., C. Giamberardino, S. M. Sykes, C.-H. Yu, J. L. Tenor *et al.*, 2017
Population genomics and the evolution of virulence in the fungal pathogen
Cryptococcus neoformans. Genome Res. 27**:** 1207-1219.
Engelthaler, D. M., N. D. Hicks, J. D. Gillece, C. C. Roe, J. M. Schupp *et al.*, 2014
Cryptococcus gattii in North American Pacific Northwest: Whole-Population
Genome Analysis Provides Insights into Species Evolution and Dispersal. mBio
5**:** e01464-01414.
Ergin, Ç., M. Şengül, L. Aksoy, A. Döğen, S. Sun *et al.*, 2019 *Cryptococcus neoformans*
recovered from olive trees (*Olea europaea*) in Turkey reveal allopatry with
African and South American lineages. Frontiers in Cellular and Infection
Microbiology 9.
Farrer, R. A., M. Chang, M. J. Davis, L. van Dorp, D.-H. Yang *et al.*, 2019 A new lineage
of *Cryptococcus gattii* (VGV) discovered in the central Zambezian Miombo
Woodlands. mBio 10**:** e02306-02319.
Ferreira-Paim, K., L. Andrade-Silva, F. M. Fonseca, T. B. Ferreira, D. J. Mora *et al.*, 2017
MLST-based population genetic analysis in a global context reveals clonality
amongst *Cryptococcus neoformans* var. *grubii* VNI isolates from HIV patients in
Southeastern Brazil. PLOS Neglected Tropical Diseases 11**:** e0005223.
Foster, P. L., 2006 Methods for determining spontaneous mutation rates. Methods in
enzymology 409**:** 195-213.
Garcia-Hermoso, D., G. Janbon and F. Dromer, 1999 Epidemiological evidence for
dormant *Cryptococcus neoformans* infection. Journal of Clinical Microbiology
37**:** 3204-3209.
Gillet-Markowska, A., G. Louvel and G. Fischer, 2015 *bz-rates*: A Web Tool to Estimate
Mutation Rates from Fluctuation Analysis. G3: Genes|Genomes|Genetics 5**:**
2323.
Gonzalez-Hilarion, S., D. Paulet, K.-T. Lee, C.-C. Hon, P. Lechat *et al.*, 2016 Intron
retention-dependent gene regulation in *Cryptococcus neoformans*. Sci. Rep. 6**:**
32252.
Gou, L., J. S. Bloom and L. Kruglyak, 2019 The genetic basis of mutation rate variation
in yeast. Genetics 211**:** 731-740.

Gröhs Ferrareze, P. A., C. Maufrais, R. Silva Araujo Streit, S. J. Priest, C. A. Cuomo *et al.*,
2021 Application of an optimized annotation pipeline to the *Cryptococcus*
deuterogattii genome reveals dynamic primary metabolic gene clusters and
genomic impact of RNAi loss. G3 (Bethesda) 11.
Guindon, S., J.-F. Dufayard, V. Lefort, M. Anisimova, W. Hordijk *et al.*, 2010 New
algorithms and methods to estimate maximum-likelihood phylogenies:
assessing the performance of PhyML 3.0. Systematic Biology 59**:** 307-321.
Hagen, F., P. C. Ceresini, I. Polacheck, H. Ma, F. v. Nieuwerburgh *et al.*, 2013 Ancient
dispersal of the human fungal pathogen *Cryptococcus gattii* from the Amazon
rainforest. PLOS ONE 8**:** e71148.
Hagen, F., K. Khayhan, B. Theelen, A. Kolecka, I. Polacheck *et al.*, 2015 Recognition of
seven species in the *Cryptococcus gattii/Cryptococcus neoformans* species
complex. Fungal Genet. Biol. 78**:** 16-48.
Hommel, B., L. Mukaremera, R. J. B. Cordero, C. Coelho, C. A. Desjardins *et al.*, 2018
Titan cells formation in *Cryptococcus neoformans* is finely tuned by
environmental conditions and modulated by positive and negative genetic
regulators. PLoS Pathog. 14**:** e1006982.
Institute, C. L. S., 2017 Reference method for broth dilution antifungal susceptibility
testing of yeasts; approved standard, document document M27-A3 in *National*
Committee for Clinical Laboratory Standards.
Janbon, G., K. L. Ormerod, D. Paulet, E. J. Byrnes III, G. Chatterjee *et al.*, 2014 Analysis
of the genome and transcriptome of *Cryptococcus neoformans* var. *grubii*
reveals complex RNA expression and microevolution leading to virulence
attenuation. PLoS Genet. 10**:** e1004261.
Janbon, G., J. Quintin, F. Lanternier and C. d'Enfert, 2019 Studying fungal pathogens of
humans and fungal infections: fungal diversity and diversity of approaches.
Genes & Immunity 20**:** 403-414.
Johnston, S. A., K. Voelz and R. C. May, 2016 *Cryptococcus neoformans*
thermotolerance to avian body temperature is sufficient for extracellular
growth but not intracellular survival in macrophages. Sci. Rep. 6**:** 20977.
Kidd, S. E., F. Hagen, R. L. Tscharke, M. Huynh, K. H. Bartlett *et al.*, 2004 A rare
genotype of *Cryptococcus gattii* caused the cryptococcosis outbreak on
Vancouver Island (British Columbia, Canada). Proc. Natl. Acad. Sci. U.S.A. 101**:**
17258-17263.
Kwon-Chung, K. J., and J. E. Bennett, 1978 Distribution of a and α mating types of
Cryptococcus neoformans among natural and clinical isolates. Am. J. Epidemiol.
108**:** 337-340.
Kwon-Chung, K. J., J. A. Fraser, T. L. Doering, Z. Wang, G. Janbon *et al.*, 2014
Cryptococcus neoformans and *Cryptococcus gattii*, the etiologic agents of
cryptococcosis. Cold Spring Harb Perspect Med 4**:** a019760.
Li, H., 2018 Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics
34**:** 3094-3100.
Li, H., B. Handsaker, A. Wysoker, T. Fennell, J. Ruan *et al.*, 2009 The Sequence
Alignment/Map format and SAMtools. Bioinformatics (Oxford, England) 25**:**
2078-2079.
Lin, X., and J. Heitman, 2006 The biology of the *Cryptococcus neoformans* species
complex. Ann. Rev. Microbiol. 60**:** 69-105.

Lin, X., S. Patel, A. P. Litvintseva, A. Floyd, T. G. Mitchell *et al.*, 2009 Diploids in the
Cryptococcus neoformans serotype A population homozygous for the alpha
mating type originate via unisexual mating. PLoS Pathog. 5**:** e1000283.
Litvintseva, A. P., I. Carbone, J. Rossouw, R. Thakur, N. P. Govender *et al.*, 2011
Evidence that the human pathogenic fungus *Cryptococcus neoformans* var.
grubii may have evolved in Africa. PloS One 6**:** e19688.
Litvintseva, A. P., L. Kestenbaum, R. Vilgalys and T. G. Mitchell, 2005 Comparative
analysis of environmental and clinical populations of *Cryptococcus neoformans*.
Journal of Clinical Microbiology 43**:** 556-564.
Litvintseva, A. P., X. Lin, I. Templeton, J. Heitman and T. G. Mitchell, 2007 Many
globally isolated AD hybrid strains of *Cryptococcus neoformans* originated in
Africa. PLoS Pathog. 3**:** e114.
Litvintseva, A. P., R. E. Marra, K. Nielsen, J. Heitman, R. Vilgalys *et al.*, 2003 Evidence of
sexual recombination among *Cryptococcus neoformans* serotype A isolates in
sub-Saharan Africa. Eukaryot. Cell 2**:** 1162-1168.
Litvintseva, A. P., and T. G. Mitchell, 2012 Population genetic analyses reveal the
African origin and strain variation of *Cryptococcus neoformans* var. *grubii*. PLoS
Pathog. 2012**:** e1002495.
Loftus, B., E. Fung, P. Roncaglia, D. Rowley, P. Amedeo *et al.*, 2005 The genome and
transcriptome of *Cryptococcus neoformans*, a basidiomycetous fungal pathogen
of humans. Science 307**:** 1321-1324.
Matsumoto, M. T., A. M. Fusco-Almeida, L. C. Baeza, M. d. S. C. Melhem and M. J. S.
Medes-Giannini, 2007 Genotyping, serotyping and determination of mating-
type of *Cryptococcus neoformans* clinical isolates from São Paulo State, Brazil.
Revista do Instituto de Medicina Tropical de São Paulo 49**:** 41-47.
Meyer, W., D. M. Aanensen, T. Boekhout, M. Cogliati, M. R. Diaz *et al.*, 2009a
Consensus multi-locus sequence typing scheme for *Cryptococcus neoformans*
and *Cryptococcus gattii*. Med. Mycol. 47**:** 561-570.
Meyer, W., D. M. Aanensen, T. Boekhout, M. Cogliati, M. R. Diaz *et al.*, 2009b
Consensus multi-locus sequence typing scheme for *Cryptococcus neoformans*
and *Cryptococcus gattii*. Med. Mycol. 57**:** 561-570.
Prum, R. O., J. S. Berv, A. Dornburg, D. J. Field, J. P. Townsend *et al.*, 2015 A
comprehensive phylogeny of birds (Aves) using targeted next-generation DNA
sequencing. Nature 526**:** 569-573.
Raj, A., M. Stephens and J. K. Pritchard, 2014 fastSTRUCTURE: Variational Inference of
Population Structure in Large SNP Data Sets. Genetics 197**:** 573.
Rajasingham, R., R. M. Smith, B. J. Park, J. N. Jarvis, N. P. Govender *et al.*, 2017 Global
burden of disease of HIV-associated cryptococcal meningitis: an updated
analysis. The Lancet Infectious Diseases 17**:** 873-881.
Rhodes, J., M. A. Beale, M. Vanhove, J. N. Jarvis, S. Kannambath *et al.*, 2017a A
population genomics approach to assessing the genetic basis of within-host
microevolution underlying recurrent cryptococcal meningitis infection. G3:
Genes|Genomes|Genetics 7: 1165.
Rhodes, J., C. A. Desjardins, S. M. Sykes, M. A. Beale, M. Vanhove *et al.*, 2017b Tracing
genetic exchange and biogeography of *Cryptococcus neoformans* var. *grubii* at
the global population level. Genetics**:** 10.1534/genetics.1117.203836.

Ruma, P., S. C. A. Chen, T. C. Sorrel and A. G. Brownlee, 1996 Characterization of
Cryptococcus neoformans by random DNA amplification. Letters in Applied
Microbiology 23**:** 312-316.

Silva, D. C., M. A. Martins, M. W. Szeszs, L. X. Bonfietti, D. Matos *et al.*, 2012
Susceptibility to antifungal agents and genotypes of Brazilian clinical and
environmental *Cryptococcus gattii* strains. Diagnostic Microbiology and
Infectious Disease 72**:** 332-339.

Souto, A. C. P., L. X. Bonfietti, K. Ferreira-Paim, L. Trilles, M. Martins *et al.*, 2016
Population genetic analysis reveals a high genetic diversity in the Brazilian
Cryptococcus gattii VGII population and shifts the global origin from the
amazon rainforest to the semi-arid desert in the Northeast of Brazil. PLoS
neglected tropical diseases 10**:** e0004885-e0004885.

Trevijano-Contador, N., H. C. de Oliveira, R. García-Rodas, S. A. Rossi, I. Llorente *et al.*,
2018 *Cryptococcus neoformans* can form titan-like cells in vitro in response to
multiple signals. PLoS Pathog. 14**:** e1007007.

Vanhove, M., M. A. Beale, J. Rhodes, D. Chanda, S. Lakhi *et al.*, 2017 Genomic
epidemiology of *Cryptococcus* yeasts identifies adaptation to environmental
niches underpinning infection across an African HIV/AIDS cohort. Mol. Ecol. 26**:**
1991-2005.

Vilas-Bôas, A. M., L. E. Andrade-Silva, K. Ferreira-Paim, D. J. Mora, T. B. Ferreira *et al.*,
2020 High genetic variability of clinical and environmental *Cryptococcus gattii*
isolates from Brazil. Med. Mycol. 58**:** 1126-1137.

Yadav, V., S. Sun, R. B. Billmyre, B. C. Thimmappa, T. Shea *et al.*, 2018 RNAi is a critical
determinant of centromere evolution in closely related fungi. Proc. Natl. Acad.
Sci. U.S.A. 115**:** 3108.

Zaragoza, O., and K. Nielsen, 2013 Titan cells in *Cryptococcus neoformans*: Cells with a
giant impact. Current Opinion in Microbiology 16**:** 409-413.