A new test suggests that balancing selection maintains hundreds of non-synonymous

polymorphisms in the human genome

Vivak Soni[1,3], Michiel Vos[2,3] and Adam Eyre-Walker[1,4]

1. School of Life Sciences

University of Sussex

Brighton

BN1 9QG

United Kingdom

2. European Centre for Environment and Human Health, University of Exeter Medical

School, Environment and Sustainability Institute

Penryn

TR10 9FE

United Kingdom

3. These authors contributed equally to this study

4. Correspondence: a.c.eyre-walker@sussex.ac.uk

Abstract

The role that balancing selection plays in the maintenance of genetic diversity remains

unresolved. Here we introduce a new test, based on the McDonald-Kreitman test, in which

the number of polymorphisms that are shared between populations is contrasted to those

that are private at selected and neutral sites. We show that this simple test is robust to a

variety of demographic changes, and that it can also give a direct estimate of the number of

shared polymorphisms that are directly maintained by balancing selection. We apply our

method to population genomic data from humans and conclude that more than a thousand

non-synonymous polymorphisms are subject to balancing selection.

Introduction

How genetic variation is maintained, either in the form of DNA sequence diversity or quantitative genetic variation, remains one of the central problems of population genetics. Balancing selection encapsulates several selective mechanisms that increase variability within a population. These include heterozygote advantage (also referred to as overdominance), frequency dependent selection, and selection that varies through space and time (Nielsen, 2005). However, although there are some clear examples of each type of selection (Allison, 1956; Nosil, et al., 2018), the overall role that balancing selection plays in maintaining genetic variation, either directly, or indirectly through linkage, remains unknown.

A number of methods have been developed to detect the signature of balancing selection (Hughes and Nei, 1988; Asthana et al. 2004; Bubb et al. 2006; Andres et al. 2009; Leffler et al. 2013; Degiorgio et al. 2014; Gao et al. 2015; Hunter-Zinck & Clark, 2015; Fijarczyk & Babik, 2015; Sheehan & Song, 2016; Siewert & Voight, 2017; Bitarello et al. 2018). Application of these methods have identified a number of loci subject to balancing selection, largely in the human genome, in which most of this research has taken place. However, these methods are generally quite complex to apply, often leveraging multiple population genetic signatures of balancing selection and many require simulations to determine the null distribution. Furthermore, they do not readily yield an estimate of the number of polymorphisms that are directly subject to balancing selection, as opposed to being in linkage disequilibrium. Here we introduce a method that is simple to apply and which generates a direct estimate of the number of polymorphisms subject to balancing selection.

One signature of balancing selection that has been utilised in several studies is the sharing of polymorphisms between species (Asthana et al. 2004; Leffler et al. 2013; Gao et al. 2015). If the species are sufficiently divergent that they are unlikely to share neutral polymorphisms, then shared genetic variation can be attributed to balancing selection. These studies have concluded that there are relatively few balanced polymorphisms that are shared between humans and chimpanzees (Asthana et al. 2004; Leffler et al. 2013). However, this test is likely to be weak because humans and chimpanzees diverged millions

of years in the past and it is unlikely that any shared selection pressures will be maintained

over that time period.

The major problem with approaches that consider the sharing of polymorphisms between

species or populations is differentiating selectively maintained polymorphisms from neutral

variation inherited from the common ancestor. This problem can be solved by comparing

the number of shared polymorphisms at sites which are selected, to those that are neutral.

We expect the number of shared polymorphisms at selected sites to be lower than at

neutral sites because many mutations at selected sites are likely to be deleterious, and

hence unlikely to be shared. However, we can estimate the proportion that are effectively

neutral by considering the ratio of polymorphisms, which are private to one of the two

populations or species, at selected versus neutral sites. Although the method can be applied

to any group of neutral and selected sites that are interspersed with one another we will

characterise it in terms of non-synonymous and synonymous sites. Let the numbers of

polymorphisms that are shared between two populations or species be S_N and S_S at non-

synonymous and synonymous sites respectively, and the numbers that are private to one of

the populations be R_N and R_S respectively. Let us assume that synonymous mutations are

neutral and non-synonymous mutations are either neutral or strongly deleterious. Then it is

evident that $\frac{S_N}{S_S} = \frac{R_N}{R_S} = f$, where f is the proportion of the non-synonymous mutations that

are neutral. However, if there is balancing selection acting on some non-synonymous SNPs

and this selection persists for some time such that the balanced polymorphisms are shared

between populations then $\frac{S_N}{S_S} > \frac{R_N}{R_S}$. A simple test of balancing selection is therefore

whether Z > 1 where

$$Z = \frac{S_N/S_S}{R_N/R_S},$$ (1)

a simple corollary of the McDonald-Kreitman test for adaptive divergence between species

(McDonald and Kreitman, 1991). It can be shown, under some simplifying assumptions in

which synonymous mutations are neutral and non-synonymous mutations are strongly

deleterious, neutral or subject to balancing selection, that an estimate of the proportion of

non-synonymous mutations subject directly to balancing selection is $\alpha_b = 1 - \frac{S_S R_N}{S_N R_S}$ (see

results section). In this analysis, we perform population genetic simulations to investigate

whether the method can detect the signature of balancing selection and assess whether the

method is robust to demographic change. Second, we apply the method to human

population genetic data. We show that the method is robust and we estimate that

substantial numbers of non-synonymous polymorphisms are maintained by balancing

selection in humans.

Results

Simulations

We propose a new test for balancing selection in which the ratio of selected to neutral

polymorphisms is compared between those that are shared between populations or species

and those that are private to populations or species. To explore the properties of our

method to detect balancing selection we ran a series of simulations in which an ancestral

population splits to yield two descendent populations. We initially simulated loci under a

simple stationary population size model where the ancestral population is duplicated to

form two equally sized populations (equal to each other and the ancestral population). This

is an unrealistic scenario, but it has the advantage that it involves no demographic change in

the transition from ancestral to descendent populations. We assume that synonymous

mutations are neutral and we explore the consequences of different selective models for

non-synonymous mutations. If all non-synonymous mutations are neutral, then as expected

$Z = 1$ (figure 1a), and if we make some of the non-synonymous mutations deleterious,

drawing their selection coefficients from a gamma distribution, as estimated from human

polymorphism data (Eyre-Walker et al, 2006) we find that $Z < 1$ (figure 1a). Again, this is

expected because slightly deleterious mutations (SDMs) are likely to contribute more to the

level of private than shared polymorphism. If we simulate a locus in which most non-

synonymous mutations are deleterious, drawn from a gamma distribution, but each locus

contains a single balanced polymorphism that is shared between populations then $Z >$

1 (figure 1a). It is important to note that the density of balanced polymorphisms is high in

these simulations because we have simulated a short exon, of just 288bp, the average

length in humans, and each one contains a balanced polymorphism. If we were to reduce

the density of balanced polymorphisms then Z can be less than one even if there is

balancing selection operating.

Slightly deleterious mutations tend to depress the value of Z because they are more likely to

segregate within a population, than to be shared between populations that diverged

sometime in the past. There are two potential strategies for coping with this tendency. We

can test for the presence of balancing selection as a function of the frequencies of the

polymorphisms in the population, because SDMs will tend to be enriched amongst the rarer

polymorphisms in the population. A similar approach has been used successfully to

ameliorate the effects of SDMs in the classic MK approach for estimating the rate of

adaptive evolution between species (Fay et al. 2001; Charlesworth and Eyre-Walker, 2008;

Messer and Petrov, 2013). Or we can explicitly model the generation of shared and private

polymorphisms under a realistic demographic and selection model to control for the effects

of SDMs. We focus our attention here on the first of these strategies, although we touch on

the latter strategy in the discussion. We apply the frequency filter to both the private and

shared polymorphisms; this is necessary because if we applied the filter only to the private

polymorphisms, we could be comparing high frequency private polymorphisms, with a low

ratio of R_N to R_S, because SDMs have been excluded, to low frequency shared

polymorphisms, which may contain many SDMs and hence have a high value of S_N/S_S; this

can yield artefactual evidence of of balancing selection. This could be exacerbated if some of

the SDMs are recessive. To investigate the effects of polymorphism frequency on our

estimate of Z we divided polymorphisms into 5 bins of 0.1 (we did not orient SNPs); for

shared polymorphisms we estimated their frequency as the unweighted mean frequency

from the two populations.

If we simulate a population in which non-synonymous mutations are deleterious, whose

effects are drawn from a gamma distribution, we find that $Z < 1$ but this is less marked for

the high frequency categories, as we expect. For the lowest frequency category Z decreases

as a function of the time to most recent common ancestor, whereas for the higher

frequency categories it is either unaffected or increases slightly (figure 1b). If we include a

balanced polymorphism, introduced prior to the population split and subject to strong

selection, into the model, which still also includes deleterious mutations, we find that $Z > 1$

for all frequency bins except the lowest one (figure 1c). In each case Z increases as a

function of the time since the population split; this is to be expected because Z is related to

the proportion of shared polymorphisms that are subject directly to balancing selection (see

below), and as time progresses, so neutral and SDMs go to fixation or loss in one or both of

the descendant populations. Note, once again that the level of balancing selection in these

simulations is high because every locus contains a balanced polymorphism.

Figure 1: Stationary population size simulations, in which the ancestral population is

duplicated to form two daughter populations of the same size to each other and the

ancestor. Each simulation was repeated 2 million times. The time to the most recent

common ancestor (tMRCA) is measured in N generations, where N is the population size. A Z

value of greater than 1 indicates a greater proportion of shared non-synonymous

polymorphisms than private non-synonymous polymorphism, which is a signal of balancing

selection. For b-c) private polymorphisms have been binned by minor allele frequency, in

bins of size 0.1. a) - simulations of neutral genetic variation only; a) – orange & b) neutral

and deleterious variation; a) – green & c) neutral, deleterious and balanced polymorphisms.

The simulation above does not take into account the demographic effects that a division in a

population involves. We therefore performed more realistic simulations which involve

vicariance and dispersal scenarios with and without migration between the sampled

populations (Figures S1-S10). We also simulated with and without expansion after

separation. We performed all simulations under two distributions of fitness effects (DFE),

which were estimated from human and *Drosophila melanogaster* populations. In the

vicariance scenario the ancestral population splits into two daughter populations of equal or

unequal sizes. In the dispersal scenario a single daughter population is generated by

duplicating part of the ancestral population, which remains the same size as it was before;

we vary the daughter population size. In both cases, we explore the consequences of

expansion after separation of the populations.

None of the simulated demographic scenarios is capable of generating Z values greater than

1 under either DFE - the method does not seem to generate false positives (Figures S1-S10).

However, it is worth noting that a more severe difference in the size of the descendant

populations results in depressed Z values in the smaller of the two populations, suggesting

demography can affect the value of Z. In all cases the value of Z is smallest for the lowest

frequency category, those polymorphisms with frequencies <0.1, and this frequency

category often shows a dramatic difference to the other categories. We therefore suggest

combining the polymorphisms above 0.1 when data is limited. As expected, we find that Z<1

in all simulations when we sum all polymorphisms with frequencies > 0.1 (Figures S11 and

S12).

Estimating the level of balancing selection

One of the great advantages of our method is that it gives an estimate of the number of

polymorphisms that are directly affected by balancing selection, under a simple model of

evolution. Let us assume that synonymous mutations are neutral and that non-synonymous

mutations are strongly deleterious, neutral or subject to balancing selection; we further

assume that all balanced polymorphisms arose before the two populations split. Then the

expected numbers of non-synonymous, R_N, and synonymous, R_S, private polymorphisms are

$R_S = \theta \rho W$

$R_N = \theta \rho W f$ (2)

where $\theta = 4N_e u$, N_e is the effective population size and u is the mutation rate per site per

generation. ρ is the proportion of polymorphisms that are private to the population, W is

Watterson's coefficient and f is the proportion of non-synonymous mutations that are

neutral, $(1-f)$ being deleterious or subject to balancing selection.

In deriving expressions for S_N and S_S we have to take into account that a balanced

polymorphism can maintain neutral variation in linkage disequilibrium that may also be

shared between populations. If we have b balanced non-synonymous polymorphisms and

each of those maintains x neutral mutations in linkage disequilibrium, then the expected

values of S_N and S_S are

$S_S = \theta(1 - \rho)W + bx$
$S_N = \theta(1 - \rho)Wf + b + bxf$ $\hspace{3cm}$ (3)

It is then straightforward to show that the proportion of shared non-synonymous

polymorphisms that are directly maintained by balancing selection is

$\alpha_b = 1 - Z = 1 - \dfrac{S_S R_N}{S_N R_S} = \dfrac{b}{S_N}$ $\hspace{3cm}$ (4)

This is clearly an unrealistic model in several respects. First, it can be expected that there

are slightly deleterious mutations in many populations and this will lead to an

underestimation of α_b; and second, it is likely that new balanced polymorphisms will be

arising all the time and these will contribute to private polymorphism, increasing R_N/R_S and

leading to a conservative estimate of α_b.

Data analysis - humans
We have shown that the method has the potential to detect balancing selection under

realistic evolutionary models. We therefore applied our method to human data from the

1000 genomes project (The 1000 Genomes Project Consortium, 2015) focussing on four

populations – Africans, Europeans, East Asians and South Asians. We find that Z >1 when

private polymorphisms from the African population are used for all population comparisons

if the frequency of private and shared polymorphisms >0.1; we also find that Z>1 in the

South Asian and East Asian population comparison when using South Asian private

polymorphisms and polymorphisms with frequencies > 0.1 (Figure 2). For several

comparisons we have no polymorphisms at the relevant frequencies, and many of the

confidence intervals on our estimates of Z are large. As a consequence, we combined the

data for all polymorphisms with frequencies > 0.1 (Figure 2, right-most point in each panel).

The patterns above are replicated; Z is significantly greater than one when we use private

polymorphisms from Africa, and in the comparison between the East and South Asian

populations, but Z<1 otherwise. In some cases, the Z value for the combined data can

appear inconsistent with the Z values from the individual frequency categories – for

example, in the European-South Asian comparison the combined Z value is greater than the

Z value for 0.1-0.2 despite the fact that this is the only Z value above 0.1. This is because

there are polymorphisms with frequencies >0.2, but there are not enough to yield a valid

estimate of Z.

Although, the values of Z are not consistently > 1 across populations, the results suggest

that there is balancing selection operating. Our simulations show that Z is consistently <1

when there is no balancing selection, and that the value of Z differs between the two

population comparisons if the populations have undergone different demographies. We

therefore infer that there is balancing selection maintaining polymorphisms between

populations.

267
268

Figure 2: The value of Z plotted against the frequency of shared and private polymorphisms,
calculated for 1000 genome data from Africans (AFR), East Asians (EAS), Europeans (EUR)
and South Asians (SAS). In each panel we show the value of Z for a comparison of two
populations using the private polymorphisms from each, the population used being
indicated in the plot legend. Data binned by minor allele frequency bins of size 0.1 on the x-
axis. Final bin is 0.1-0.5 (i.e. all data minus the lowest frequency bin). Confidence intervals
were generated by bootstrapping the data by gene 100 times. Only datapoints in which
there were at least 20 polymorphisms for all polymorphism categories were plotted,
because the confidence intervals were very large otherwise.

If we estimate α_b to those comparisons in which Z is significantly greater than 1, we
estimate that approximately 12% of the non-synonymous shared polymorphisms between
the African and other human populations are subject to balancing selection, as well as
between the Asian populations (Table 1). These estimates are likely to be underestimates
because there will still be SDMs segregating in our data, even though we have removed the
lowest frequency variants (see simulation results). The proportions suggest that ~500
polymorphisms, which are shared between the African and other populations, are
maintained by balancing selection. We estimate rather more are maintained between the
two Asian populations, although the confidence intervals on this estimate are large. If we
combine data across the non-African populations, we estimate that ~1400 polymorphisms
are shared between African and non-African populations in total because of balancing
selection (Table 1). The fact that the estimate from the African-non-African comparison is
larger than the estimate from the African versus each individual population, suggests that
many of the balanced polymorphisms shared between populations are unique to each
population pair – i.e. there are balanced polymorphisms shared between African and
European populations, that are not shared between Africans and East Asians.

Target population	Comparative population	α_b	α_{b_low}	α_{b_high}	b	b_{low}	b_{high}
African	nonAfrican	0.12	0.09	0.14	1420	1077	1737
African	European	0.04	0.01	0.06	577	176	926
African	East Asian	0.05	0.03	0.06	657	401	882
African	South Asian	0.03	0.01	0.05	507	182	796
South Asian	East Asian	0.09	0.00	0.19	1273	18	2594

Table 1: Estimates of the proportion of shared non-synonymous polymorphisms under

balancing selection, α_b, and the number of polymorphisms, b, being directly maintained by

balancing selection for population comparisons in which Z>1. 95% confidence intervals were

generated by bootstrapping the data by gene 100 times.

A concern in any analysis of human population genetic data is the influence of biased gene

conversion (BGC). This process tends to increase the number and allele frequencies of

AT>GC mutations, and reduce the number and allele frequencies of GC>AT mutations. If this

process differentially affects synonymous and non-synonymous sites and shared and private

polymorphisms, then it could potentially lead to Z>1. To investigate whether BGC has an

effect we performed two analyses. In the first, we divided our genes according to whether

they were in high and low recombining regions, dividing the data at the median

recombination rate (RR). Our two groups differ substantially in their mean rate of

recombination (mean RR in low group = 1.2×10^{-7} centimorgans per site and high group =

1.8×10^{-6} centimorgans per site). We find that Z is actually higher in the low RR regions,

although not significantly so (Table 2), which suggests that BGC is not responsible for the

comparisons in which Z > 1.

mean recombination rate	sN	sS	rN	rS	Z	Z_{low}	Z_{high}
1.20E-09	7023	7447	553	653	1.114	1.066	1.160
1.80E-08	7767	8514	604	702	1.060	1.021	1.103

Table 2: Estimates of *Z* for data split by median recombination rate. Confidence intervals

were generated by bootstrapping genes 100 times.

In the second test of the influence of BGC on the value of Z we limited our analysis to

mutations that are not affected by BGC – i.e. G<>C and A<>T mutations. This reduces our

dataset by about 80%. As a consequence, we summed the data for all polymorphisms with

frequencies >0.1. We find that our estimate is largely unchanged from that when all

polymorphisms are included, however the confidence intervals are increased substantially

so that Z is only significantly greater than one for the African-non-African, and the South

versus East Asian comparisons (Table 2). Our two tests suggest that our results are not

affected by BGC.

Target population	Comparative population	all polymorphism data			filtered for BGC		
		Z	Z_{low}	Z_{high}	Z	Z_{low}	Z_{high}
African	non-African	1.13	1.10	1.17	1.12	1.03	1.21
African	East Asian	1.05	1.03	1.07	1.00	0.95	1.05
African	European	1.04	1.01	1.07	1.01	0.93	1.10
African	South Asian	1.04	1.01	1.06	1.04	0.98	1.09
South Asian	East Asian	1.10	1.00	1.23	1.36	1.04	1.73

Table 3: Testing the effects of BGC for population comparisons which show Z>1 using all

polymorphisms with frequencies > 0.1. To control for BGC the analysis was restricted to

A<>T and G<>C SNPs. 95% confidence intervals were generated by bootstrapping the data

by gene 100 times.

Groups of genes

We can potentially apply our test of balancing selection to individual genes or groups of

genes, where we have enough data. Balancing selection has been implicated in the

evolution of immune related genes (e.g. Bitarello et al. 2018, Weedel and Conway, 2010,

Hughes and Nei, 1988, Hedrick and Thompson, 2002), particularly major histocompatibility

complex (MHC), or human leukocyte antigen genes (HLA) (Aguilar et al. 2004 and Paterson,

1998). To investigate whether we could detect this signature in our data, we split our

dataset into HLA and non-HLA genes **(The MHC sequencing consortium, 1999)**. Due to a lack

of private polymorphisms, we combined all frequency categories >0.1. We found Z was

significantly greater in HLA than non-HLA genes for all population comparisons except

Europeans and South Asians using European private polymorphisms (Figure S13). However,

the value of Z is greater than one in population comparisons in which African private

polymorphisms are used, consistent with balancing selection maintaining variation in the

HLA region. We estimate that a very substantial proportion of non-synonymous genetic

variation is being maintained by balancing selection, although the confidence intervals on

our estimates are large; roughly 50% of the shared non-synonymous SNPs are being

maintained by balancing selection between African and non-African populations in the HLA

region and this equates to approximately 200 polymorphisms (Table 4).

However, the signature of balancing selection that we have detected across all genes is not

simply due to the HLA genes. We find that Z>1 in non-HLA genes in most population

comparisons in which Z>1 for all genes, except in the comparison of East and South Asian

populations (Table 5); in most cases these estimates of Z are significantly greater than 1. We

estimate that >1000 of non-synonymous polymorphisms are subject to balancing selection

amongst the polymorphisms shared by African and non-African populations, with several

100 shared between each of the populations and the African population.

target	comparative	α	α_{b_low}	α_{b_low}	b	b_{low}	b_{high}
AFR	nonAFR	0.702	-0.011	0.779	299	Na	332
AFR	EAS	0.287	-0.005	0.502	134	Na	234
AFR	EUR	0.731	0.317	0.820	338	147	379
AFR	SAS	0.529	0.307	0.694	247	143	324
EAS	SAS	-0.307	Na	0.292	Na	Na	133

Table 4: Estimates of the proportion of shared non-synonymous polymorphisms under

balancing selection, α_b, and the number of polymorphisms being directly maintained by

balancing selection, b, for population comparisons in the HLA region for population

comparisons in which Z>1 when using all genes. Estimates for polymorphisms with

frequency > 0.1. Confidence intervals were generated by bootstrapping the data by gene

100 times.

target	comparative	α_b	α_{b_low}	α_{b_low}	b	b_{low}	b_{high}
AFR	nonAFR	0.101	0.077	0.127	1193	907	1502
AFR	EAS	0.033	0.016	0.049	453	227	678
AFR	EUR	0.024	-0.001	0.049	355	Na	716
AFR	SAS	0.021	0.001	0.041	293	19	587
EAS	SAS	-1.254	-1.981	-0.833	Na	Na	Na

Table 5. Estimates of the proportion of shared non-synonymous polymorphisms under balancing selection, α_b, in non-HLA genes, and the number of polymorphisms being directly maintained by balancing selection, b, for population comparisons in which Z>1 when using all genes. Confidence intervals were generated by bootstrapping the data by gene 100 times.

If we run our analysis grouping genes by their GO category and restricting the analysis to those groups that have at least 100 polymorphisms with frequencies >0.1 we find 683 categories in which Z is significantly greater than 1 in at least population comparison (Supplementary file S1) and we list those significant in 5 or more population comparisons in Table 6. One of these GO categories, "nucleic acid binding" is shared across 7 of the 14 population comparisons, "endoplasmic reticulum membrane" across 6 population comparisons; amongst those categories shared among 5 are "viral process" and "immune system process", but there are many others which are surprising including "protein import into the nucleus" (Supplementary file S1). Eighty-five categories are shared between 4 or more population comparisons, and 155 amongst three or more population comparisons. These include 7 categories related to immunity (including immune system process which is significant in 5 population comparisons), and 40 categories that are linked to antigen presentation though not classified as immune-related categories. There are also 5 viral-related categories (including viral process which is significant in 5 population comparisons).

GO category	Number of population comparisons
nucleic acid binding	7
endoplasmic reticulum membrane	6
response to stimulus	5
viral process	5
zinc ion binding	5
DNA binding	5
nucleus	5
immune system process	5
chromatin binding	5
keratinization	5
intermediate filament	5
positive regulation of transcription by RNA polymerase II	5
chromosome	5

Table 6. GO categories in which Z is significantly greater than one in at least 5 population

comparisons.

Individual genes

We applied our statistic to individual human genes, combining all frequency bins (0-0.5) due

to a lack of polymorphism. We tested for significance using a one-tailed Fisher's exact test.

Of the 14,261 genes we analysed 514 had Z>1 in at least one population comparison.

Eighteen of these were nominally significant at p<0.1 (supplementary file S2), but no gene

was individually significant when we corrected for multiple testing using a Bonferroni

correction. Eighteen genes have Z>1 in at least 9 population comparisons; note that since

populations share polymorphisms, we cannot combine the evidence for balancing selection

across these populations (Table 7). Four of these genes MUC4, RP1L1, PKD1L2 and ZAN have

Z>1 in all population comparisons.

Gene symbol	Number of population comparisons in which Z>1
MUC4	14
RP1L1	14
PKD1L2	14
ZAN	14
C1orf167	13
SPTBN5	12
MKI67	12
DNAH14	11
WDFY4	10
FAM230G	10
CMYA5	9
CRIPAK	9
SYNE2	9
FSIP2	9
GREB1	9
ALMS1	9
MUC19	9
CENPF	9

Table 7. Genes with Z>1 in multiple population comparisons.

If we use the 514 genes and do a GO enrichment analysis, we find multiple GO categories

enriched for these genes including immune response categories with 3-fold enrichment. The

most highly enriched categories are involved in energy production and conversion (including

dynein binding) and intracellular transport (including microtubule motor activity)

(Supplementary file S2).

Discussion

We propose a new method for detecting and quantifying the amount of balancing selection that is operating on polymorphisms, in which the numbers of non-synonymous and synonymous polymorphisms that are shared between populations and species are compared to those that are private. The method is analogous to the McDonald-Kreitman test used to test and quantify the amount of adaptive evolution between species (McDonald & Kreitman, 1991). We show that our test is robust to the presence of slightly deleterious mutations under simple demographic models of population division, expansion and migration. When we apply our method to data from human populations, we find evidence that hundreds of non-synonymous polymorphisms are being directly maintained by balancing selection in human populations.

Our method for detecting balancing selection is simple to apply and appears to be robust to changes in demography. The classic MK test of adaptive evolution between species can generate artefactual evidence of adaptive evolution if there are SDMs and there has been population size expansion (McDonald & Kreitman, 1991; Eyre-Walker, 2002); this is because SDMs that might have been fixed when the effective population size was small, no longer segregate once the population size is large. A similar bias does not appear to affect our test, although we have only investigated two DFEs and a limited number of demographic scenarios. Our test is likely to be more robust than the classic MK test because the shared polymorphisms are affected by the demographic changes that affect the private polymorphisms; i.e. if the population expands this will increase the effectiveness of natural selection on both the private and the shared polymorphisms. However, although our method seems to be relatively robust to changes in demography, in the sense that it does not generate artefactual evidence of balancing selection, it is evident that demography does affect the chance of balancing selection being identified, because the values of Z depend on the demography and which population the private polymorphisms are taken from (Figure 2).

The method can in principle be applied to any pair of populations or species. However, the test is likely to be weak when the populations/species are closely related for two reasons.

First, there will be relatively few private polymorphisms, and second, the proportion of

shared polymorphisms that are subject to balancing selection is likely to be low, because so

many neutral polymorphisms are shared between populations because of recent common

ancestry. As the populations/species diverge so the number of private polymorphisms will

increase, and the proportion of shared polymorphisms that are balanced will increase. Of

course, as the time of divergence increases so the selective conditions that maintained the

polymorphism are likely to change and the polymorphism might become neutral, or subject

to directional selection. Our method is also likely, like all methods, to be better at detecting

balanced polymorphisms that are common, because most populations are dominated by

large numbers of rare neutral variants. Finally, our method requires that the neutral and

selected sites are interdigitated; the method is therefore easy to apply to protein coding

sequences, but may be more difficult to apply to other types of variation, such as that which

affects gene expression.

A great advantage of our method is that it gives an estimate of the proportion and number

of shared polymorphisms that are directly subject to balancing selection, under a set of

simplifying assumptions. However, the method is likely to yield underestimates of the

proportion of balanced polymorphisms, under a more realistic models of evolution. We

have assumed, in deriving α_b, that all non-synonymous mutations are either strongly

deleterious, neutral or subject to balancing selection. However, a substantial fraction of

non-synonymous mutations appear to be slightly deleterious in humans (Cargill et al, 1999;

Fay et al, 2001; Eyre-Walker et al, 2002; Hughes, et al, 2003; Asthana, et al., 2007) and other

species (Fay et al, 2001; Eyre-Walker et al, 2002; Hughes, 2005; Charlesworth & Eyre-

Walker, 2006) – i.e. they are deleterious, but sufficiently weakly selected that they

contribute to polymorphism. Under stationary population size assumptions – i.e. in which

the ancestral population is duplicated to form the daughter populations - this will lead to an

underestimate of Z because SDMs tend to contribute more to private than shared

polymorphism, and hence inflate R_N/R_S relative to S_N/S_S (Figure 1). Under more realistic

demographic models, in which at least one of the derived populations is reduced, this is

expected to depress Z in the population that is being reduced because more SDMs will tend

to segregate in smaller populations hence inflating R_N/R_S (compare Figure 1 to Figures S2

and S3). Simulations suggest there is however a slight increase in Z using private

polymorphisms from the larger of the populations but these Z values never exceed 1

(Figures S2 and S3). The second reason that we are likely underestimating the number of

balanced polymorphisms using our simple method is that we assume that there are no

balanced polymorphisms that are private to each population; these would inflate R_N/R_S.

Private balanced polymorphisms might arise from an ancestral polymorphism that is lost

from one of the daughter populations, or one that arises de novo. A more realistic model

balancing selection is one in which balanced polymorphisms are continually generated with

the selective forces persisting for some time before they dissipate (Sellis et al, 2011) and the

balanced polymorphism is lost. The process of population division itself is likely to lead to

the loss of many balanced polymorphisms as the environment shifts in the two daughter

populations.

A potential solution to the tendency for our method to underestimate Z is to simulate data

under a realistic demographic model assuming that there is no balancing selection, and

interpret the observed values of Z that are greater than simulated values, as evidence of

balancing selection. A similar approach has been used to estimate the rate of adaptive

evolution between species (e.g. Eyre-Walker & Keightley, 2009; Boyko, et al., 2008;

Schneider et al, 2011; Galtier, 2016; Tataru et al, 2017). However, there are challenges in

this approach; in particular, we need an accurate demographic model. We have performed

simulations under the commonly used human demographic model inferred by Gravel et al.

(2011) estimating the DFE from the current African population; we chose the African

population because it has been subject to relatively modest demographic change. Our

observed Z values do not match the simulated values (Figure S14); in particular we find that

the observed values of Z are substantially greater than the simulated amongst the low

frequency polymorphisms. However, the model of Gravel et al. does not fit the SFS of the

individual populations of 1000 genome data; for example, in the African population there

are far too many singleton SNPs even amongst the putative neutral synonymous mutations

(Figure S15). The lack of fit is perhaps not surprising; Gravel et al. inferred their model using

80 chromosomes per population, whereas the 1000 genome data contains >1000

chromosomes per population. Furthermore, the inference of a demographic model should

take into account the influence of biased gene conversion and background selection, which

appear to be pervasive factors in the human genome (Pouyet et al, 2018), so these

simulations will be complex.

It might be argued that the evidence of balancing selection is weak because we typically find

Z>1 using the private polymorphism from only one of the two populations. However, we

have been unable to find a demographic model in which there is no balancing selection and

Z>1 – note that we never observe Z>1 under the Gravel et al. model of human demography

even when we change the parameters of the DFE and demographic model; furthermore, we

find that simulations which involve different demographies in the two populations generate

different Z values for the two populations, so there is an expectation in many species that

we will observe Z values that differ between populations.

Values of Z in excess of one could potentially be due to biased gene conversion; we expect

BGC to increase the allele frequency of AT>GC mutations and to decrease the frequency of

GC>AT mutations; this will tend to make AT>GC mutations more likely to be shared between

populations than GC>AT mutations. Since the GC-content at the third codon position is

typically higher than the GC content at the first two positions, this will mean that there are

more non-synonymous AT>GC mutations than synonymous, and hence more shared non-

synonymous than synonymous polymorphisms. However, our results do not appear to be

affected by BGC; results are similar between genes in high and low recombination rate

regions of the human genome (Table 2), and our point estimates of Z are largely unaffected

by restricting the analysis to SNPs which are unaffected by BGC, although the confidence

intervals increase substantially (Table 3).

We estimate that there ~500 balanced polymorphisms shared between the African and each

of the other human populations, ~1250 shared between the Asian populations and ~1400

shared between the African and non-African populations; these are likely to be

underestimates due to the presence of slightly deleterious mutations. The fact that we

estimate that ~1400 shared non-synonymous polymorphisms are being maintained

between African and non-African populations, but only ~500 between each of the individual

populations and the African population suggests, that many of the polymorphisms shared

between the African and each individual population are unique – i.e. many polymorphisms

shared between Africans and Europeans, may not be shared between Africans and Asians.

These numbers are substantial, but are consistent with those of Bitarello et al. (2018) who

estimated that ~8% of human genes show some evidence of long-term balancing selection,

and many of these signatures are shared between populations. However, their method

could not determine whether the balanced polymorphisms were coding or noncoding

mutations.

As expected, we find evidence of balancing selection affecting the HLA or MHC genes (table

4) (Hughes & Nei; 1988; Hedrick, 1998). However, we find evidence of balancing selection

even when these genes are removed from the analysis (Table 5). The analysis of GO

categories shows that numerous categories show evidence of balancing selection across

multiple population comparisons (supplementary file S1). Some of these are expected, but

many are not, such as "nucleic acid binding", which is significant in 7 of the 14 population

comparisons.

No individual gene is significant when we control for multiple testing, however, several

genes have Z>1 in multiple population comparisons including 10 which are shared across at

least 10 of the 14 population comparisons. Three of these overlap with previous genome-

wide scans of selection, namely the protein-coding gene DNAH14, implicated in brain

compression and encoding axenomal dynein (Voight et al, 2006); MUC4, implicated in biliary

tract cancer (Tennessen & Akey, 2011); and ZAN, which encodes a protein involved in sperm

adhesion, previously implicated in balancing selection and positive selection in human

populations (Gasper & Swanson, 2006). Two of these ten genes are associated with

tumours. MKI67 expression is associated with a higher tumour grade and early disease

recurrence (Yang, et al., 2017), and WDFY4 plays a critical role in the regulation of certain

viral and tumour antigens in dendritic cells (Theisen, et al., 2019). PKD1L2 is associated with

polycystic kidney disease and RP1L1 variants are associated with several retinal diseases

including occult macular dystrophy (Davidson, et al., 2013). SPTBN5 encodes for the

cytoskeletal protein spectrin, that plays a role in maintaining cytoskeletal structure (Huh,

Glantz, Je, Morrow, & Kim, 2001) and C1orf167 expresses open reading frame protein that is

highly expressed in the testis (Fagerberg, et al., 2014). Finally, FAM230G is highly expressed

in testes (Delihas, 2018).

Twenty-five of the 514 with Z>1 genes overlap with those genes identified by Bitarello et al.

(2018), but this is similar to the level of overlap expected at random; i.e. they observed that

7.9% of protein coding genes overlapped regions identified by their method as being subject

to balancing selection, and we identified 514 candidates; so we expect 0.079 x 514 = 41 by

chance alone. The lack of a significant overlap is possibly not surprising; we have applied our

method to non-synonymous variation, whereas the method of Bitarello et al. (2018)

considers all variation. Furthermore, the method of Bitarello et al. (2018) is most powerful

at detecting balancing selection over long time periods; in the case of humans, over periods

of millions of years. In contrast, we have applied our method to populations that diverged

10,000's of years ago.

It is possible that the signature of balancing selection is caused by a form of associated

overdominance, in which neutral alleles at a locus are linked to different deleterious

recessive alleles at other loci. For example, let us imagine that we have two closely linked

loci at which we have deleterious alleles; let the A2 allele be the recessive allele at the A

locus and the B2 allele at the B locus. Now consider a third neutral locus with alleles C1 and

C2. If C1 is in linkage disequilibrium (LD) with the A2 allele, and C2 is in LD with the B2 allele,

then C1C2 heterozygous individuals will have higher fitness than C1C1 and C2C2

homozygotes. This form of selection can lead to the maintenance of genetic variation (Zhao

and Charlesworth 2016) in low recombination rate regions. However, Z is not substantially

greater in regions of low recombination so AOD seems an unlikely explanation (Table 2).

Conclusion

610 We present a new approach to test for the presence of balancing selection and to the

611 number of polymorphisms that are directly affected by it. Our method appears to be robust

612 to demographic change. Application of the method to human population genetic data

613 suggests that 100s of non-synonymous polymorphisms shared between populations are

614 being maintained by balancing selection.

Methods and Materials

Human data

Human variation data was obtained from 1000 genomes Grch37 vcf files (The 1000 Genomes Project Consortium, 2015)**. Variants were annotated using Annovar's hg19 database (Wang & Li, 2010). The annotated data was then parsed to remove multi-nucleotide polymorphisms and indels. Because 1000 genomes data provides allele frequencies for the non-reference allele rather than the minor allele, the minor allele frequency for each superpopulation and also for the global minor allele frequency was calculated. We used 1000 genomes from the African, South Asian, East Asian and European populations. The American population was removed due to the fact that it is an admixed population. GO category information was obtained from Ensembl's BioMart data mining tool (Yates, et al., 2020). We used pyrho demography-aware recombination rate maps (Spence & Song, 2019) for analyses that control for recombination rate.

Simulations

All simulations were run using the SLiM software platform (Haller & Messer, 2017). Parameter values were taken from human estimates. Almost all simulations were of a 288bp locus, this being the average size of a human exon (Yates et al. 2020). Unless otherwise stated, the scaled recombination rate and scaled mutation rate were set at $r = 1.1$ x 10^{-8} (Dumont & Payseur, 2008); $\mu = 2.5$ x 10^{-8} (Nachman & Crowell, 2000)in the ancestral population. The distribution of fitness effects was assumed to be a gamma distribution and the shape and mean strength of selection estimates for humans were taken from Eyre-Walker et al. (2006) ($\beta = 0.23$; mean $N_e s = 425$). For *Drosophila* estimates were taken from Keightley and Eyre-Walker (2007) ($\beta = 0.35$; mean $N_e s = 1800$); again these were values in the ancestral population. Unless dominance was fixed, it was calculated using the model of Huber et al. (2018), which was estimated from Arabidopsis. The Huber model varies the dominance coefficient depending on the selection coefficient of the mutation, where the dominance coefficient increases with the strength of selection. It's formula is $h = f(s) = \frac{1}{\frac{1}{\theta_{intercept}} - \theta_{rate} s}$ where $\theta_{intercept}$ defines the values of h at $s = 0$ and θ_{rate} determines how quickly h approaches zero with decreasing negative selection coefficient. We set $\theta_{intercept}$

to 0.5 so that all mutations with a selection coefficient of $s = 0$ have a dominance

coefficient, $h = 0.5$, and θ_{rate} = 41225.56. This assumes an inverse relationship between h

and s, which gives the highest log likelihood score of the relationships compared by Huber

et al. (2018). For balancing selection simulations, we assume a model of heterozygote

advantage and the strength of selection was sampled from a distribution such that the

equilibrium frequency was uniformly distributed between 0 and 1; however, it should be

noted that some balanced polymorphisms with low equilibrium frequencies were lost in one

of the descendent populations. These simulations were discarded. The balanced

polymorphism is introduced at the centre of the 288bp region. Two million successful

simulation runs were conducted for each model.

For the generic simulations (i.e. not those involving the human demographic model) the

ancestral population size was set at 200. This was allowed to equilibrate for 15N

generations before a balanced polymorphism was introduced 5N generations before the

population was split into two. The descendant populations were then sampled every 0.05N

generations up to 20N generations after the split. We ran five different generic simulations:

(i) simulations in which the ancestral population was duplicated, (ii) vicariance simulations in

which the ancestral population was divided between the daughter populations in splits of

0.5N-0.5N; 0.75N-0.25N; 0.9N-0.1N, (iii) variance simulations in which the descendant

populations expanded, iv) dispersal simulations, in which some variable fraction (0.5N;

0.25N; 0.1N) of the ancestral population is duplicated to form the dispersal population, and

the ancestral population continues as the other daughter population, and v) dispersal with

population increase of the dispersal population. The dispersal population starts as 0.1N and

expands exponentially 2 to 10x its original size after 21N generations. Scenarios ii-v were

repeated with migration rates of 0.01N and 0.001N of the ancestral population size

between the descendant populations.

We also ran some simulations under the human demographic model of Gravel et al. (2011);

for details of the demographic structure of the simulation (Gravel et al. 2011). The

distribution of fitness effects for deleterious mutations was assumed to be a gamma

distribution using the parameters estimated from the African superpopulation using the

GammaZero model within the Grapes software (Galtier, 2016); the parameters are similar to

those estimated by Eyre-Walker et al (2006), and used in the generic simulations (Gamma

shape = 0.17 and Mean $N_e s$ = 1144). We chose to infer the DFE for the African

superpopulation because this is currently the largest dataset available for a population that

has been inferred to be relatively stable. Dominance was calculated using the Huber model

discussed above. Sampling of all populations (African, East Asian and European) was

conducted at the end of the simulation (i.e. the equivalent of the present day).

Acknowledgements: the work was supported by grant #????? from the Natural

Environment Research Council to MV.

Works Cited

Aguilar, A., Roemer, G., Debenham, S., Binns, M., Garcelon, D., & Wayne, R. (2004). High MHC diversity maintained by balancing selection in an otherwise genetically monomorphic mammal. *PNAS, 101*(10), 3490-4.

Allison, A. (1956). The sickle-cell and Haemoblobin C genes in some African populations. *Ann. Human Genetics, 21*, 67-89.

Andres, A., Hubisz, M., Indap, A., Torgerson, D., Degenhardt, J., Boyko, A., . . . Clark, A. N. (2009). Targets of Balancing Selection in the Human Genome. *Mol Biol Evol., 26*(12), 2755-64.

Ashburner, M., Ball, A., Blake, J., Botsein, D., Butler, H., Cherry, J., . . . Sherlock, G. (2000). Gene ontology: tool for the unification of biology. *Nature Genetics, 25*, 25-9.

Asthana, S., Noble, W., Kryukov, G., Grant, C., Sunyaev, S., & Stamatoyannopoulos, J. (2007). Widely distributed noncoding purifying selection in the human genome. *PNAS, 104*(30), 12410-5.

Asthana, S., Schmidt, S., & Sunyaev, S. (2004). A limited role for balancing selection. *Trends in Genetics, 21*(1), 30-2.

Auton, A., Fledel-Alon, A., Pfeifer, S., Venn, O., Segurel, L., Street, T., . . . Donnelly, P. (2012). A fine scale chimpanzee genetic map from population sequencing. *Science, 336*(6078), 193-8.

Bamshad, M., Mummidi, S., Gonzalez, E., Ahuja, S., Dunn, D., Watkins, W., . . . Ahuja, S. (2002). A strong signature of balancing selection in the 5' cis- regulatory region of CCR5. *Proc Natl Acad Sci, 99*(16), 10539-44.

Bepari, K., Malakar, A., Paul, P., Halder, B., & Chakraborty, S. (2015). Allele frequency for Cystic fibrosis in Indians vis-a/-vis global populations. *Bioinformation, 11*(7), 348-52.

Bitarello, B., de Filippo, C., T. J., Schmidt, J., Kleinert, P., Meyer, D., & Andres, A. (2018). Signatures of Long-Term Balancing Selection in Human Genomes. *GBE, 10*(3), 939–55.

Black, F., & Hedrick, P. (1997). Strong balancing selection at HLA loci: Evidence from segregation in South Amerindian families. *PNAS, 94*(23), 12452-6.

Boyko, A., Williamson, S., Indap, A., Degenhardt, J., Hernandez, R., Lohmueller, K., . . . Bustamante, C. (2008). Assessing the Evolutionary Impact of Amino Acid Mutations in the Human Genome. *PLoS Genetics, 4*(5), e1000083.

Breuer, K., Foroushani, A., Laird, M., Chen, C., Sribnaia, A., Lo, R., . . . Lynn, D. (2013). InnateDB: systems biology of innate immunity and beyond - recent updates and continuing curation. *Nucl. Acids Res., 41*.

Bronson, P., Mack, S., Erlich, H., & Slatkin, M. (2013). A sequence-based approach demonstrates that balancing selection in classical human leukocyte antigen (HLA) loci is asymmetric. *Hum Mol Genet., 22*(2), 252-61.

Bubb, K., Bovee, D., Buckley, D., Haugen, E., Kibukawa, M., Paddock, M., . . . Olson, M. (2006). Scan of Human Genome Reveals No New Loci Under Ancient Balancing Selection. *Genetics, 173*(4), 2165-77.

CA, J., P, T., & M, W. (2001). The major histocompatibility complex and its functions. In *Immunobiology: The Immune System in Health and Disease*. New York: Garland Science.

Cargill, M., Altshular, D., Ireland, J., Sklar, P., Ardlie, K., Patil, N., . . . Lander, E. (1999). Characterisation of single-nucleotide polymorphisms in coding regions of human genes. *Nat Genet., 22*(3), 231-8.

Charlesworth, J., & Eyre-Walker, A. (2006). The Rate of Adaptive Evolution in Enteric
Bacteria. *Mol Biol Evol., 23*(7), 1348-56.
Charlesworth, J., & Eyre-Walker, A. (2008). The McDonald–Kreitman Test and Slightly
Deleterious Mutations. *Mol Biol Evol., 25*(6), 1007-15.
Cheng, X., & DeGiorgio, M. (2018). Detection of shared balancing selection in the absence
of trans-species polymorphism. *Mol Biol Evol., 36*(1), 177-99.
Comeron, J., Ratnappan, R., & Bailin, S. (2012). The many landscapes of recombination in
Drosophila melanogaster. *PLoS Genetics, 8*(10), e1002905.
Davidson, A., Sergouniotis, P., Mackay, D., Wright, G., Waseem, N., Michaelides, M., . . .
Webster, A. (2013). RP1L1 variants are associated with a spectrum of inherited
retinal diseases including retinitis pigmentosa and occult macular dystrophy. *Hum
Mutat., 34*(3), 506-14.
Day, N., & Byar, D. (1979). Testing hypotheses in case-control studies-equivalence of
Mantel–Haenszel statistics and logit score tests. *Biometrics, 35*(3), 623–30.
DeGiorgio, M., Lohmueller, K., & Nielsen, R. (2014). A Model-Based Approach for
Identifying Signatures of Ancient Balancing Selection in Genetic Data. *PLoS
Genetics, 10*(8).
Delihas, N. (2018). A family of long intergenic non-coding RNA genes in human
chromosomal region 22q11.2 carry a DNA translocation breakpoint/AT-rich
sequence. *PLoS One, 13*(4), e0195702.
Dumont, B., & Payseur, B. (2008). Evolution of the genomic rate of recombination in
mammals. *Evolution, 62*(2), 281.
Enard, D., Cai, L., Gwennap, C., & Petrov, D. (2016). Viruses are a dominant driver of
protein adaptation in mammals. *Elife, 5*, e12469.
Eyre-Walker, A., & Keightley, P. (2009). Estimating the rate of adaptive molecular evolution
in the presence of slightly deleterious mutations and population size change. *Mol Biol
Evol., 26*(9), 2097-108.
Eyre-Walker, A., Keightley, P., Smith, N., & Gaffney, D. (2002). Quantifying the Slightly
Deleterious Mutation Model of Molecular Evolution. *Mol. Bol. Evol., 19*(12), 2142-9.
Eyre-Walker, A., Woolfit, M., & Phelps, T. (2006). The Distribution of Fitness Effects of
New Deleterious Amino Acid Mutations in Humans. *Genetics, 173*(2), 891-900.
Fagerberg, L., Hallstrom, B., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J., . . .
Nilsson, P. (2014). Analysis of the human tissue-specific expression by genome-wide
integration of transcriptomics and antibody-based proteomics. *Mol Cell Proteomics.,
13*(2), 397-406.
Fay, J., Wycoff, G., & CI, W. (2001). Positive and negative selection on the human genome.
Genetics, 158(3), 1227-34.
Ferrer-Admetlla, A., Liang, M., Korneliussen, T., & Nielsen, R. (2014). On detecting
incomplete soft or hard selective sweeps using haplotype structure. *Mol Biol Evol.,
31*(5), 1275-91.
Fijarczyk, A., & Babik, W. (2015). Detecting balancing selection in genomes: limits and
prospects. *Molecular Ecology, 24*, 3529–45.
Fu, Y., & Li, W. (1993). Statistical tests of neutrality of mutations. *Genetics, 133*, 693-709.
Gabriel, S., Brigman, K., Koller, B., Boucher, R., & Stutts, M. (1994). Cystic fibrosis
heterozygote resistance to cholera toxin in the cystic fibrosis mouse model. *Science,
266*(5182), 107-9.
Galtier, N. (2016). Adaptive Protein Evolution in Animals and the Effective Population Size
Hypothesis. *PLoS Genetics, 12*(1), e1005774.
Gao, Z., Przeworski, M., & Sella, G. (2015). Footprints of ancient-balanced polymorphisms
in genetic variation data from closely related species. *Evolution, 69*(2), 431-46.

Gasper, J., & Swanson, W. (2006). Molecular Population Genetics of the Gene Encoding the
Human Fertilization Protein Zonadhesin Reveals Rapid Adaptive Evolution. *Am J*
Hum Genet., 79(5), 820-30.
Gossett, C., Nasciemento, J., Auge, M., & Bierne, N. (2014). Evidence for adaptation from
standing genetic variation on an antimicrobial peptide gene in the mussel Mytilus
edulis. *Molecular Ecology, 23*(12).
Gravel, S., Henn, B., Guttenkunst, R., Indap, A., Marth, G., Clark, A., . . . Bustamante, C.
(2011). Demographic history and rare allele sharing among human populations.
PNAS, 108(29), 11983-8.
Gutenkunst, R., Hernandez, R., Williamson, S., & Bustamante, C. (2009). Inferring the Joint
Demographic History of Multiple Populations from Multidimensional SNP Frequency
Data. *PLoS Genetics, 5*(10).
Haller, B. &. (2017). SLiM 2: Flexible, interactive forward genetic simulations. *34*(1).
Haller, B., & Messer, P. (2017). SLiM 2: Flexible, interactive forward genetic simulations.
Mol Biol Evol., 34(1), 230-40.
Harpak, A., Bhaskar, A., & Pritchard, J. (2016). Mutation rate variation is a primary
determinant of the distribution of allele frequencies in humans. *PLoS Genetics,*
12(12).
Hedrick, P. (1998). Balancing selection and MHC. *Genetics, 104*(3), 207-14.
Hedrick, P. (2012). What is the evidence for heterozygote advantage selection? *Trends in*
Ecology and Evolution, 27(12), 698-704.
Hedrick, P., & Thompson, G. (2002). Pathogen resistance and genetic variation at MHC loci.
Evolution, 56, 1902-8.
Hodgkinson, A., Ladoukakis, E., & Eyre-Walker, A. (2009). Cryptic Variation in the Human
Mutation Rate. *PLoS Biology, 7*(2).
Huber, C., Durvasula, A., Hancock, A., & Lohmueller, K. (2018). Gene expression drives the
evolution of dominance. *Nature Communications, 9*(2750).
Hudson, R., & Kaplan, N. (1988). The coalescent process in models with selection and
recombination. *Genetics, 120*(23), 831–40.
Hudson, R., Kreitman, M., & Aguade, M. (1997). A test of neutral molecular evolution based
on nucleotide data. *Genetics, 116*(1), 153-9.
Hughes, A. (2005). Evidence for Abundant Slightly Deleterious Polymorphismsin Bacterial
Populations. *Genetics, 169*, 533-8.
Hughes, A., & Nei, M. (1988). Pattern of nucleotide substitution at major histocompatibility
complex class I loci reveals overdominant selection. *Nature, 335*(6186), 167-70.
Hughes, A., & Nei, M. (1989). Nucleotide substitution at major histocompatibility complex
class II loci: Evidence for overdominant selection. *Proceedings of the Natural*
Academy of Sciences, 958-62.
Hughes, A., Packer, B., Welch, R., Bergen, A., Chanock, S., & Yeager, M. (2003).
Widespread purifying selection at polymorphic sites in human protein-coding loci.
PNAS, 26, 15754-7.
Huh, G.-Y., Glantz, S., Je, S., Morrow, J., & Kim, J. (2001). Calpain proteolysis of αII-
spectrin in the normal adult human brain. *Neuroscience Letters, 316*(1), 41-4.
Hunter-Zinck, H., & Clark, A. (2015). Aberrant Time to Most Recent Common Ancestor as a
Signature of Natural Selection. *Mol Biol and Evol., 32*(10), 2784-97.
Johnson, P., & Hellmann, I. (2011). Mutation rate distribution inferred from coincident SNPs
and coincident substitutions. *GBE, 3*, 842-50.
Josefson, D. (1998). CF Gene May Protect against Typhoid Fever. *BMC, 316*(7143), 1481.

Keightley, P., & Eyre-Walker, A. (2007). Joint inference of the distribution of fitness effects
of deleterious mutations and population demography based on nucleotide
polymorphism frequencies. *Genetics, 177*, 2251-61.
Key, F., Teixeira, J., de Filippo, C., & Andre, A. (2014). Advantageous diversity maintained
by balancing selection in humans. *Current Opinion in Genetics & Development, 29*,
45-51.
Kimura, M. (1983). *The Neutral Theory of Molecular Evolution.* Cambridge/New York:
Cambridge Univ. Press.
Klein, J. (1987). Origin of Major Histocompatibility Complex Polymorphism: The Trans-
Species Hypothesis. *Human Immunology*(19), 155-62.
Leffler, E., Gao, Z., Pfeifer, S., Seguel, L., Auton, A., Venn, O., . . . Przeworski, M. (2013).
Multiple instances of ancient balancing selection shared between humans and
chimpanzees. *Science, 339*(6127), 1578-82.
Li, H. (2011). A statistical framework for SNP calling, mutation discovery, association
mapping and population genetical parameter estimation from sequencing data.
Bioinformatics, 27(21), 2987–93.
Li, W., Wu, H., Li, F., Tian, S., Kherraf, Z., Zhang, J., . . . Zhang, F. (2019). Biallelic
mutations in CFAP65 cause male infertility with multiple morphological
abnormalities of the sperm flagella in humans and mice. *BMJ, 10*(1136).
Lippman, Z., & Zamir, D. (2007). Heterosis: revisiting the magic. *Trends Genetics, 23*, 60-6.
Loewe, L., & Charlesworth, B. (2006). Inferring the distribution of mutational effects on
fitness in Drosophila. *Biol Lett., 2*(3), 426-30.
Mann, H., & Whitney, D. (1947). On a Test of Whether one of Two Random Variables is
Stochastically Larger than the Other. *Annals of Mathematical Statistics, 18*(1), 50-60.
Martinez, G., Kherraf, Z., Zoari, R., Mustapha, S., Saut, A., Pernet-Gallay, K., . . . Ray, P.
(2018). Whole-exome sequencing identifies mutations in FSIP2 as a recurrent cause
of multiple morphological abnormalities of the sperm flagella. *Human Reproduction,*
33(10), 1973-84.
McDonald, J., & Kreitman, M. (1991). Adaptive protein evolution at the Adh locus in
Drosophila. *Nature, 351*(6328), 652-4.
Messer, P., & Petrov, D. (2013). Frequent adaptation and the McDonald-Kreitman test. *Proc*
Natl Acad Sci, 110(21), 8615-20.
Meyer, D., & Thomson, G. (2001). How selection shapes variation of the human major
histocompatibility complex: a review. *Ann Hum Genet, 65*, 1–26.
Meyer, M., Kircher, M., Gansauge, M., Li, H., Racimo, F., Mallick, S., . . . Shendure, J.
(2012). A High-Coverage Genome Sequence from an Archaic Denisovan Individual.
Science, 338(6104), 222-6.
Nachman, M., & Crowell, S. (2000). Estimation of the mutation rate per nucelotide in
humans. *Genetics, 156*(1), 297-304.
Nielsen, R. (2001). Statistical tests of selective neutrality in the age of genomics. *Heredity,*
86, 641-7.
Nielsen, R. (2005). Molecular Signatures of Natural Selection. *Annual Review Genetics, 39*,
197-218.
Nielsen, R., Hellmann, I., Hubisz, M., Bustamante, C., & Clark, A. (2007). Recent and
ongoing selection in the human genome. *Nat Rev Genet., 8*(11), 857-68.
Nosil, P., Villoutreix, R., de Carvalho, C., Farkas, T., Soria-Carrasco, V., Feder, J., . . .
Gompert, Z. (2018). Natural selection and the predictability of evolution in Timema
stick insects. *Science, 359*(6377), 765-70.

Oliver, M., Lambin, X., Cornulier, T., & Piertney, S. (2008). Spatio-temporal variation in the
strength and mode of selection acting on major histocompatibility complex diversity
in water vole (Arvicola terrestris) metapopulations. *Molecular Ecology, 18*(1).
Pasteur, G. (1982). A classificatory review of mimicry systems. *Annual Review of Ecology*
and Systematics, 13, 169-99.
Pasvol, G., Weatherall, D., & Wilson, R. (1978). Cellular mechanism for the protective effect
of haemoglobin S against P. falciparum malaria. *Nature, 274*, 701-3.
Paterson, S. (1998). Evidence for balancing selection at the major histocompatibility complex
in a free-living ruminant. *The Journal of Heredity, 89*(4), 289-94.
Pouyet, F., Aeschbacher, S., Thiery, A., & Excoffier, L. (2018). Background selection and
biased gene conversion affect more than 95% of the human genome and bias
demographic inferences. *Elife*, e36317.
Prufer, K., de Filippo, C., Grote, S., Mafessoni, F., Korlevic, P., Hajdinjak, M., . . . Nelson,
B. (2017). A high-coverage Neandertal genome from Vindija Cave in Croatia.
Science, 358(6363), 655-8.
Prufer, K., Racimo, F., Patterson, N., Jay, F., Sankararaman, S., Sawyer, S., . . . Theunert, C.
(2013). The complete genome sequence of a Neanderthal from the Altai Mountains.
Nature, 505, 43-9.
Quinlan, A., & Hall, I. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics, 26*(6), 841-2.
Reich, D., & ES, L. (2001). On the allelic spectrum of human disease. *Trends Genet, 17*(9),
502–10.
Sabeti, P., Reich, D., Higgins, J., Levine, H., Richter, D., Schaffner, S., . . . Lander, E.
(2002). Detecting recent positive selection in the human genome from haplotype
structure. *Nature, 419*, 832-7.
Sakharkar, M., Chow, V., & Kangueane, P. (2004). Distributions of exons and introns in the
human genome. *In Silico Biology, 4*(4), 387-93.
Schneider, A., Charlesworth, B., Eyre-Walker, A., & Keightley, P. (2011). A Method for
Inferring the Rate of Occurrence and Fitness Effects of Advantageous Mutations.
Genetics, 189(4), 1427-37.
Sellis, D., Callahan, B., Petrov, D., & Messer, P. (2011). Heterozygote advantage as a natural
consequence of adaptation in diploids. *PNAS, 108*(51), 20666-71.
Sheehan, S., & Song, Y. (2016). Deep Learning for Population Genetic Inference. *PLoS*
Computational Biology.
Siewart, K., & Voight, B. (2017). Detecting Long-Term Balancing Selection Using Allele
Frequency Correlation. *Mol Biol Evol., 34*(11), 2996–3005.
Singh, N., Arndt, P., & Petrov, D. (2005). Genomic Heterogeneity of Background
Substitutional Patterns in Drosophila melanogaster. *Genetics, 169*(2), 709-22.
Smith, N., & Eyre-Walker, A. (2002). Adaptive protein evolution in Drosophila. *Nature, 415*,
1022-4.
Spence, J., & Song, Y. (2019). Inference and analysis of population-specific fine-scale
recombination maps across 26 diverse human populations. *Sci Adv., 5*(10), e9206.
Tajima, F. (1989). Statistical method for testing the neutral mutation hypothesis by DNA
polymorphism. *Genetics, 123*(3), 585–95.
Tataru, P., Mollion, M., Glemin, S., & Bataillon, T. (2017). Inference of distribution of
fitness effects and proportion of adaptive substitutions from polymorphism data.
Genetics, 207, 1103-19.
Tennessen, J., & Akey, J. (2011). Parallel adaptive divergence among geographically diverse
human populations. *PLoS genetics, 7*(6), e1002127.

The 1000 Genomes Project Consortium. (2015). A global reference for human genetic
variation. *Nature, 526*(7571), 68-74.
The Gene Ontology Consortium. (2019). The Gene Ontology Resource: 20 years and still
GOing strong. *Nucleic Acids Research, 47*(D1), D330-8.
The MHC sequencing consortium. (1999). Complete sequence and gene map of a human
major histocompatibility complex. *Nature, 401*, 921-3.
Theisen, D., Davidson, J., Briseno, C., Gargaro, M., Lauron, E., Wang, Q., . . . Murphy, T.
(2019). Wdfy4-deficiency reveals a critical role for cross-presentation in anti-viral
and anti-tumor responses. *J Immunol, 177*.
Voight, B., Kudaravalli, S., Wen, X., & Pritchard, J. (2006). A map of recent positive
selection in the human genome. *PLoS Biology, 4*(3), e72.
Voight, B., Kudaravalli, S., Wen, X., & Pritchard, J. (2006). A Map of Recent Positive
Selection in the Human Genome. *PLoS Biology, 4*(3).
Wang, H., Song, W., Hu, T., Zhang, N., Miao, S., Zong, S., & Wang, L. (n.d.). Fank1
interacts with Jab1 and regulates cell apoptosis via the AP-1 pathway. *Cell Mol Life
Sci, 68*, 2129-39.
Wang, K., & Li, M. &. (2010). ANNOVAR: Functional annotation of genetic variants from
next-generation sequencing data. *38*(164).
Wang, K., Li, M., & Hakonarson, H. (2010). ANNOVAR: Functional annotation of genetic
variants from next-generation sequencing data. *Nucleic Acids Research, 38*(16), e164.
Weedel, G., & Conway, D. (2010). Detecting signatures of balancing selection to identify
targets of anti-parasite immunity. *Trends Parasitol., 26*(7), 363-9.
Welch, J. (2006). Estimating the genomewide rate of adaptive protein evolution in
Drosophila. *Genetics, 173*, 821-37.
Woolf, L., Cranston, W., & Goodwin, B. (1967). Genetics of phenylketonuria.
Heterozygosity for phenylketonuria. *Nature, 213*, 882-3.
Yang, C., yu, T., Han, C., Qin, W., Liao, X., Yu, L., . . . Peng, T. (2017). Genome-Wide
Association Study of MKI67 Expression and its Clinical Implications in HBV-
Related Hepatocellular Carcinoma in Southern China. *Cell Physiol Biochem, 42*,
1342-57.
Yates, A., Achuthan, P., Akanni, W., Allen, J., Allen, J., Alvarez-Jarreta, J., . . . Grego, T.
(2020). Ensembl 2020. *Nucleic Acids Research, 48*(D1), D682-8.
Zhao, L., & Charlesworth, B. (2016). Resolving the Conflict Between Associative
Overdominance and Background Selection. *Genetics, 203*, 1315–34.
967
968

Supplementary Material

Target population	Comparative population	all polymorphism data			filtered for BGC		
		Z	Z_{low}	Z_{high}	Z	Z_{low}	Z_{high}
African	non-African	1.13	1.10	1.17	1.12	1.03	1.21
African	East Asian	1.05	1.03	1.07	1.00	0.95	1.05
African	European	1.04	1.01	1.07	1.01	0.93	1.10
African	South Asian	1.04	1.01	1.06	1.04	0.98	1.09
South Asian	East Asian	1.10	1.00	1.23	1.36	1.04	1.73

Supplementary Table 1: Testing the effects of BGC for population comparisons which show Z>1. Confidence intervals were generated by bootstrapping the data by gene 100 times.

target	comparative	Z	Z_{low}	Z_{high}	α	$α_{b\_low}$	$α_{b\_low}$	b	b_{low}	b_{high}
AFR	nonAFR	3.354	0.989	4.533	0.702	-0.011	0.779	299	Na	332
AFR	EAS	1.403	0.995	2.007	0.287	-0.005	0.502	134	Na	234
AFR	EUR	3.714	1.465	5.568	0.731	0.317	0.820	338	147	379
AFR	SAS	2.123	1.443	3.272	0.529	0.307	0.694	247	143	324

974 **Supplementary Table 2:** Estimates of Z for HLA genes only. Confidence intervals were

975 generated by bootstrapping the data by gene 100 times.

target	comparative	Z	Z_{low}	Z_{high}	α	$α_{b\_low}$	$α_{b\_low}$	b	b_{low}	b_{high}
AFR	nonAFR	1.112	1.083	1.146	0.101	0.077	0.127	1193	907	1502
AFR	EAS	1.034	1.017	1.052	0.033	0.016	0.049	453	227	678
AFR	EUR	1.025	0.999	1.052	0.024	-0.001	0.049	355	Na	716
AFR	SAS	1.021	1.001	1.043	0.021	0.001	0.041	293	19	587

977 **Supplementary Table 3:** Estimates of Z for non-HLA genes only. Confidence intervals were

978 generated by bootstrapping the data by gene 100 times.

982
Supplementary Figure S2: Vicariance simulations in which the ancestral population splits to
form two daughter populations of the size specified in the panel. Each column is a separate
set of simulations, with the top row plotting Z against tMRCA (measured in N generations,
where N is the population size) for the larger daughter population, and the bottom row the
smaller. Deleterious mutations are drawn from a gamma DFE with parameters inferred from
human population data.
989

Supplementary Figure S2: Dispersal simulations in which a single daughter population
disperses from the ancestral population. Each column is a separate set of simulations, with
the top row plotting Z against tMRCA (measured in N generations, where N is the population
size) for the ancestral population, and the bottom row the daughter population. Deleterious
mutations are drawn from a gamma DFE with parameters inferred from human population
data.

Supplementary Figure S3: Vicariance expansion simulations in which both daughter

populations expand. The ancestral population (of size N=200) splits to form two daughter

populations of size N=100. Both daughter populations go on to expand in size. In the left

column the daughter populations double in size. In the right panel they reach 10x their initial

size. Deleterious mutations are drawn from a gamma DFE with parameters inferred from

human population data.

1005

1006

Supplementary Figure S4: Vicariance expansion simulations in which only one daughter

population expands. The ancestral population (of size N=200) splits to form two daughter

populations of size N=100. One daughter population (upper panels) goes on to expand in

size. In the left column the daughter populations double in size. In the right panel they reach

10x their initial size. Deleterious mutations are drawn from a gamma DFE with parameters

inferred from human population data.

1013

Supplementary Figure S5: Dispersal expansion simulations in which a single daughter

population disperses from the ancestral population and then expands. The ancestral

population (of size N=200) splits to form a daughter population of size N=100, which

expands to the final population size shown in the panel. Each column is a separate set of

simulations, with the top row plotting Z against tMRCA (measured in N generations, where

N is the population size) for the ancestral population, and the bottom row the daughter

population. Deleterious mutations are drawn from a gamma DFE with parameters inferred

from human population data.

1022

1023

Supplementary Figure S6: Vicariance simulations in which the ancestral population splits to

form two daughter populations of the size specified in the panel. Each column is a separate

set of simulations, with the top row plotting Z against tMRCA (measured in N generations,

where N is the population size) for the larger daughter population, and the bottom row the

smaller. Deleterious mutations are drawn from a gamma DFE with parameters inferred from

Drosophila melanogaster population data.

1030

1031

Supplementary Figure S7: Dispersal simulations in which a single daughter population

disperses from the ancestral population. Each column is a separate set of simulations, with

the top row plotting Z against tMRCA (measured in N generations, where N is the population

size) for the ancestral population, and the bottom row the daughter population. Deleterious

mutations are drawn from a gamma DFE with parameters inferred from *Drosophila*

melanogaster population data.

1038

1039

Supplementary Figure S8: Vicariance expansion simulations in which both daughter

populations expand. The ancestral population (of size N=200) splits to form two daughter

populations of size N=100. Both daughter populations go on to expand in size. In the left

column the daughter populations double in size. In the right panel they reach 10x their initial

size. Deleterious mutations are drawn from a gamma DFE with parameters inferred from

Drosophila melanogaster population data.

1046

1047

Supplementary Figure S9: Vicariance expansion simulations in which only one daughter

population expands. The ancestral population (of size N=200) splits to form two daughter

populations of size N=100. One daughter population (upper panels) goes on to expand in

size. In the left column the daughter populations double in size. In the right panel they reach

10x their initial size. Deleterious mutations are drawn from a gamma DFE with parameters

inferred from *Drosophila melanogaster* population data.

1054

1055

Supplementary Figure S10: Dispersal expansion simulations in which a single daughter
population disperses from the ancestral population and then expands. The ancestral
population (of size N=200) splits to form a daughter population of size N=100, which
expands to the final population size shown in the panel. Each column is a separate set of
simulations, with the top row plotting Z against tMRCA (measured in N generations, where
N is the population size) for the ancestral population, and the bottom row the daughter
population. Deleterious mutations are drawn from a gamma DFE with parameters inferred
from *Drosophila melanogaster* population data.

1064

1065

Supplementary Figure S11: Simulations with for combined 0.1-0.5 minor allele

frequencies. Each panel is a separate simulated scenario, with population sizes listed in the

panel legend. The first number is for the filled in data lines, denoting the ancestral population

in dispersal scenarios, and for the larger population in the vicariance scenarios. The second

number is for the dotted data lines, denoting the daughter population in dispersal scenarios,

and the smaller population in the vicariance scenarios. For more details on each scenario

please see supplementary figures S1-10. Deleterious mutations are drawn from a gamma DFE

with parameters inferred from human population data.

1074

1075

Supplementary Figure S12: Simulations with for combined 0.1-0.5 minor allele
frequencies. Each panel is a separate simulated scenario, with population sizes listed in the
panel legend. The first number is for the filled in data lines, denoting the ancestral population
in dispersal scenarios, and for the larger population in the vicariance scenarios. The second
number is for the dotted data lines, denoting the daughter population in dispersal scenarios,
and the smaller population in the vicariance scenarios. For more details on each scenario
please see supplementary figures S1-10. Deleterious mutations are drawn from a gamma DFE
with parameters inferred from *Drosophila melanogaster* population data.

1084

1085

Supplementary Figure S13: Simulations using the Gravel model of human demography

(Gravel et al, 2011). Shown are the observed (filled circles) and simulated (crosses) values of

Z. Each column represents a different population comparison. From left to right: Africans

(AFR) and East Asians (EAS), Africans and Europeans (EUR), Europeans and East Asians.

The population name in the upper left indicates which set of private polymorphisms are used

to calculate Z in each population comparison. The x-axis represents private polymorphism

minor allele frequency bins. Confidence intervals generated by bootstrapping.

1093

1094

1095 **Supplementary Figure S14:** Comparison of simulated (under the Gravel et al. (2011) model

1096 of human demography) and observed SFS from the African population.

1097

1098