# How much does it take to be old? Modelling the time since the last harvesting to infer the distribution of mature forests in France

Lucie Thompson[1], Eugénie Cateau[2], Nicolas Debaive[2], Frédéric Bray[1], André Torre[1], Patrick Vallet[1], Yoan Paillet[1,*]

1. Univ. Grenoble Alpes, INRAE, Lessem, 2 rue de la Papeterie, BP76, 38402 Saint Martin d'Hères, France
2. Réserves Naturelles de France, La Bourdonnerie, 2 allée Pierre Lacroute, CS 67524, 21075 Dijon, France
   *corresponding author: yoan.paillet@inrae.fr

Yoan Paillet, ORCID: 0000-0001-7232-7844

# Abstract

**Aim**

The global knowledge of the distribution and proportion of forests with high conservation values such as mature forests in metropolitan France is scarce, focuses only on a few specific well-known sites, and has never been tackled countrywide and evaluated with a robust multivariate statistical approach. Here, we modelled French forest reserves' time since the last harvesting operation, then inferred the current statistical distribution of mature forests (i.e. forests over 50 years without harvesting) in metropolitan France.

**Location**

Metropolitan France

**Methods**

We used inventories of forest stand structure issued from forest reserves and managed forests to calibrate a generalised linear mixed model explaining the time since the last harvesting with selected stand structure attributes of maturity, combined with environmental variables. We chose the time since the last harvesting as a proxy for forest maturity, assuming older unmanaged plots to host older trees. We then projected this model on an independent nation-wide dataset issued from the National Forest Inventory. Thus, we obtained an updated estimation of the proportion and a rough distribution of mature forest stands (i.e. abandoned for over 50 years) in metropolitan France.

**Results**

We found that the structural variables that best characterized the time since the last harvesting were high basal area of very large trees, high volumes of both standing and downed deadwood, high diversity of tree-related microhabitats and more marginally diversity of decay stages. Variables translating legacy of past forest management also characterised more mature plots, which displayed lower volumes of stumps and high density of coppices. Our projection gave an estimated 3% of French forests over 50 years without harvesting and a promising proportion of French forests that could reach interesting levels of maturity in the near future.

**Main conclusions**

Our model showed that the time since the last harvesting is a good proxy for forest maturity in French temperate forest. It highlighted the stand structure attributes that are key in characterising those forests with high conservation values. We found that forests with higher levels of maturity seem to be located in more inaccessible places notably mountainous areas. Our projection gives the first robust statistical estimate of the proportion of mature forests in metropolitan France and may serve to report on the status of mature forests.

**Keywords**

Forests, maturity, old growth, time since the last harvesting, distribution, proportion, stand structure

# 1  |  INTRODUCTION

Mature forests have a key role in the mitigation of climate change (Achard & Hansen, 2012; Frey et al., 2016), the protection of water resources, the prevention of soil erosion  (Brockerhoff et al., 2017; Forest Europe, 2015). They also host a myriad of features with high conservation value promoting biodiversity (Bauhus et al., 2009; Burrascano et al., 2013; Larrieu et al., 2018; Lassauce et al., 2012; Paillet et al., 2015, 2017).

Forest maturity is dependant of tree maturity, which is linked to the tree's biological development and is defined as a combination of tree age and its estimated longevity. Maturity is generally quantified based on the stand's age structure, e.g. a stand is considered mature if its dominant tree species are on average older than half of the species' estimated longevity (Rossi & Vallauri, 2013; Wirth et al., 2009). Although quite straightforward, this definition is problematic in two aspects. First, tree age is best determined by core sampling which is a tedious task. Second, longevity is not a well-known species feature, as it depends on various factors and it might also be altered by the long history of forest management displayed in European forests (Cateau et al., 2015). To work around those issues, international reporting generally uses the time since the last harvesting as a proxy for forest maturity or naturalness (see indicator 4.3 in the State of Europe's forests, Forest Europe 2015). By "harvesting" we mean any human intervention extracting wood biomass from the forest.

Despite the many services they provide, forests undisturbed by man or primary forests remain scarce at the global scale, especially in western Europe and most are, as yet, unprotected (Forest Europe, 2015; Sabatini et al., 2018, 2020). Studies of other high conservation value forests such as mature forests in France are scarce and have targeted, to date, only a few emblematic reserves (Christensen et al., 2005; Mountford, 2002; Pontailler et al., 1997). There is a recognised need for a global map of old-growth forest patches (Chiarucci & Piovesan, 2020), and the "EU Biodiversity Strategy for 2030" explicitly mentions that "it will be crucial to define, map, monitor and strictly protect all the EU's remaining primary and old-growth forests" notably by designating at least 10% of Europe's land to strict protection (European Commission, 2020). The knowledge of the proportion and distribution of mature forests is the first step in facilitating their protection and conservation. Mature forests usually display a long history of harvesting of varied intensity and management regimes. The use of the time since the last harvesting is a way of assessing the degree of naturalness of the forests in question. In France, it is acknowledged that 50 years of abandonment are a minimum for a forest to display sufficient maturity features (MAAPRAT-IFN, 2011). The sole estimation for the proportion of mature forest stands in France dates back to 1993 and was evaluated to about 3% (MAAPRAT-IFN, 2011). This estimation, which has not been updated since, relied more on expertise rather than sound data or rigorous analytical approach. There is therefore a knowledge gap concerning the proportion, distribution and overall characterisation of mature forests that needs to be filled.

The occurrence of mature structural attributes increases in older unmanaged stands, which display more trees with larger diameters at breast height (Burrascano et al., 2013; Heiri et al., 2009; Paillet et al., 2015), an abundance of tree-related microhabitats (Larrieu et al., 2018; Paillet et al., 2017; Winter & Möller, 2008) along with high volumes of deadwood  (Harmon, 2009; Siitonen et al., 2000). Those specific structural attributes harboured by mature forests also

have high conservation value for biodiversity (Burrascano et al., 2013; Bauhus et al., 2009; Siitonen et al., 2000). More generally, whatever their degree of maturity, these forests also bear the legacy of past – sometimes intensive – forest use in their stand structure, notably silvicultural treatments such as coppice-with-standards (Lassauce et al., 2012).

In this study, we used inventories of forest stand structure issued from forest reserves and managed forests to calibrate a generalised linear mixed model explaining the time since the last harvesting with selected attributes of maturity, herein called structural variables, combined with environmental variables. We hypothesized that time since the last harvesting would be positively influenced by a high volume of large logs and snags and high basal area of very large trees, a high diversity of tree-related microhabitats and decay stages. We expected time since last harvesting to be negatively correlated with the volume of stumps and stem density of medium trees. We also tested other structural variables that would bear witness to past management (e.g. proportion of coppice). We then projected this model on an independent nation-wide dataset issued from the National Forest Inventory. Thus, we obtained an updated estimation of the proportion and a rough distribution of mature forest stands (i.e. abandoned for over 50 years) in metropolitan France.

# 2 | METHODS

## 2.1 | Training data

We trained the full model on a dataset issued from a monitoring program that has been implementing forest stand structure description in French forest reserves since 2005, to monitor the evolution of forest attributes. A stratified sample design encompasses the variability within a forest reserve, whose plots can be subject to different management regimes. Indeed, plots comprise strict forests reserves, where harvesting is prohibited, special forest reserves, where management targets specific habitat or species, as well as nature reserves in which harvesting is usually allowed (Table 1). One forest reserve can host up to several hundred plots depending on the size of the represented area and its layout. We kept the plots where the time since the last harvesting was recorded by forest managers, by looking into current or old management plans. Plots from this network show a larger abandonment gradient than French forests in general (see Appendix S1 in Supporting Information).

4 728 plots in 71 reserves were used for modelling (Figure 1). We only kept the plots located by spatial coordinates and classified into one of the six most abundant tree species groups. We did not keep plots over 110 years since the last harvesting as we judged the data too unreliable. We left out reserves with less than 20 remaining plots to ensure sufficient data for the averaging of the random effect (see below). We checked that forests subjected to different management regimes were equivalent in terms of site fertility, therefore not introducing bias in the dataset.

| Management regime | Number of plots | Mean basal area of living trees (m².ha⁻¹) | Mean volume of standing dead trees (m³.ha⁻¹) | Mean volume of downed deadwood (m³.ha⁻¹) | Mean time since the last harvesting (years) |
|---|---|---|---|---|---|
| **Managed forest reserves** | 228 | 34.9 | 10.6 | 17.4 | 30.8 |
| **Special forest reserves** | 2628 | 27.6 | 14.5 | 28.8 | 31.5 |
| **Strict forest reserves** | 1872 | 28.4 | 10.7 | 20.4 | 33.6 |

*Table 1: Description of the training dataset (4 728 plots and 71 forest reserves) split by the management regime of plots. Special forest reserves include future forest reserves (not yet designated), national and regional forest reserves and Natura 2000 sites*

Forest type
- Pure deciduous oak
- Mostly coniferous trees
- Mostly broadleaved trees
- Mixed broadleaved trees
- Pure beech trees
- Pure spruce and/or fir

*Figure 1 : Distribution of the 4728 plots and 71 reserves from the training data kept for modelling. Plots colour and shapes correspond to forest type.*

The protocol for the stand structure surveys is detailed in Appendix S2. Living trees with a diameter at breast height (DBH) under 30 cm were all measured within a fixed 10 m radius plot. Larger trees were measured if they were comprised in a 3% angle count sampling (Paillet et al., 2019). Coppice and standard stems were differentiated. Each shoot from the same stump was measured individually. Standing deadwood under 30 cm and standing and downed deadwood over 30 cm in diameter were measured respectively on a 10 m and 20 m radius plot, discriminating between stumps (under 130 cm tall) snags and standing dead trees. Downed deadwood with a diameter under 30 cm was surveyed on three linear 20 m long transects. Decay stage as well as diameter were recorded off all sampled deadwood. Finally, tree-related microhabitat surveys followed the protocol detailed in Paillet et al. (2019) by visually inspecting trees and recording presence of microhabitats. Different microhabitat classifications have been used over

the years. Therefore, we created a harmonized classification based on a simplified version of Larrieu et al. (2018)'s

microhabitat types to narrow it down to one homogeneous classification.

## 2.2 | Structural attributes and environmental variables

Among the attributes surveyed, we kept the basal area, volume and stem density of living trees, volume of

standing deadwood (stumps, snags and standing dead trees) and volume of downed deadwood (logs). Diversity of

decay stages of deadwood and diversity of tree-related microhabitats on living trees and standing deadwood were

calculated with the Shannon diversity index (see Table 2). We split observations into diameter classes: small trees

[DBH: 17.5, 27.5cm], medium trees [27.5, 47.5], large trees [47.5, 67.5] and very large trees over 67.5 cm.

*Table 2 : Structural features chosen as candidate variables for modelling the time since the last harvesting of forest plots, and the*
*literature supporting them. "Hypothesis" relates to the expected effect of the structural feature on the time since the last harvesting:*
*« + » (« - ») means we expect a positive (negative) effect of the feature on the time since the last harvesting*
*« ++ » means it is a very common feature found in most articles dealing with forest maturity.*
*This table is based on a literature review, but other attributes were tested considering the specificities of our dataset.*
151

| Structural features | Variable | Hypothesis | Source |
|---|---|---|---|
| Living trees | Density and basal area of large and very large trees | ++ | Paillet et al., (2015) ; Burrascano et al., (2013) |
| | Density of living trees | + | Portier et al., (2020) |
| | Total basal area of living trees | + | Heiri et al., (2009) |
| | Density of medium sized living trees | - | Paillet et al., (2015) |
| Decay stages | Diversity of decay stages | + | Wirth et al., (2009) ; Winter, Möller, (2008) ; Siitonen et al. (2000) |
| Tree-related microhabitats | Diversity of tree-related microhabitats | ++ | Paillet et al., (2017) ; Larrieu et al., (2018) ; Winter, Möller, (2008) |
| Downed deadwood | Volume of downed deadwood | ++ | Portier et al., (2020) ; Paillet et al., (2015) ; Heiri et al., (2009) ; Bauhus et al., (2009) ; Vandekerkhove et al., (2009) ; Siitonen et al., (2000) |
| Standing deadwood | Volume of stumps | - | Paillet et al., (2015) ; Siitonen et al., (2000) |
| | Volume of standing dead trees and snags | ++ | Portier et al., (2020) ; Heiri et al., (2009) ; Siitonen et al., (2000) |

Diversity of decay stages and tree-related microhabitats were calculated at the plot scale with the Shannon index, one

type corresponding to either decay stage associated with tree species and type of deadwood (stump, snag, standing

dead tree, downed deadwood) or tree-related microhabitat type associated with species and type of wood (dead or

alive).

We extracted environmental variables using plot locations (see Table 3 for the sources):

- edaphic : soil pH, C/N ratio, maximum water holding capacity;

- climatic: mean annual temperature and precipitations;

- topographic: elevation, slope, aspect.

Three other environmental composite variables were also extracted or calculated:

- a site index extracted from Toïgo et al. (2015)'s model predicting tree growth as a function of tree species and
environmental variables, to account for site productivity;

- the forested region or "*sylvoécorégions*" accounted for the regional context of the plots. It is a classification
of forested areas in metropolitan France within which factors discriminant for forestry and habitat distribution
are homogeneous (L'IF n°14, 2011) ;

- the plot's forest type which was created based on an adaptation of an existing French classification, the *BD*
*forêt* version 2 (IGN, 2016a). This classification establishes forest types for plots based on the relative tree
cover of each species present on said plot. We approximated tree cover with tree basal area, as tree cover
was not accounted for in the reserve inventories. As an example, a plot was considered "pure" when over 75%
of the total basal area of the plot was composed of a single species (or a grouping of species with similar
characteristics, see Appendix S3 for more details).

*Table 3 : Environmental covariates tested for modelling and where they were sourced*

| Category | Covariate | Units | Source |
|---|---|---|---|
| **Topographic** | Elevation | meters | BD ALTI v.2.0 : digital terrain model 25 m |
| | Slope | degrees | Calculated[1] with BD ALTI v.2.0 |
| | cosine(Aspect) | - | Calculated with BD ALTI v.2.0 |
| **Climatic** | Mean annual temperature (BIO2) | degrees Celsius | WorldClim v.2 (Fick & Hijmans, 2017) |
| | Total annual precipitations (*BIO12*) | millimetres | WorldClim v.2 |
| **Edaphic** | Bioindicated soil pH[2] | - | DIGITALIS dataset - Laboratoire SILVA (Université de Lorraine-AgroParisTech-INRA) |
| | C/N ratio[3] | - | DIGITALIS dataset |
| | Maximum water holding capacity | Millimetres/cm of soil | DIGITALIS dataset |
| **Forest type** | | - | Adaptation of the *BD forêt* v.2 (IGN, 2016a) |
| **Forested region** | SylvoEcoRegion | - | National Forest Inventory |
| **Site index** | | - | Toïgo et al. (2015) |

176   [1] « terrain » function from R package *raster* (Hijmans, 2020)

## 2.3   |   Statistical methods

We used ascendant variable selection based on AIC (Akaike Information Criterion) scores to select the best

model. Considering the large size of our dataset, we chose a conservative five-point AIC threshold instead of the

standard two-points threshold. First, we selected stand structure variables, to which we added environmental

variables and interactions. We tested first order interactions as well as a few second order interactions that made

biological sense. We favoured more specific variables over more generalized ones (e.g. standing and downed

deadwood volumes separately vs. total deadwood volume). We checked for correlation and multicollinearity of model

covariates, with variance inflation factors (VIF) under five (Zuur et al., 2010).

We validated the final model following Zuur and Ieno, (2016) by plotting residuals against fitted values,

variables included and variables excluded from the model. We checked for overfitting with 50 iterations of 10-fold

cross validation (i.e. random sets of 10% of the plots were consecutively removed from the training dataset). We

looked at estimate variations as well as prediction accuracy by measuring the coefficient of determination "r squared"

from the linear regression of predicted versus observed time since the last harvesting (Harrel, 2015). Model robustness

was also considered by leaving out reserves one by one and running the model to check for influential or problematic

reserves.

## 2.4 | Prediction on systematic nation-wide dataset

We predicted the time since the last harvesting on a systematic nation-wide National Forest Inventory (NFI)

dataset. Each year, about 6500 NFI forest plots are surveyed, recording measurements of stand structure, soil related

and floristic variables. We inferred time since the last harvesting on an NFI subset from the 2012 to 2018 survey

campaigns, assuming that most of the plots included have not been subjected to major disturbance during this six-

year period, so that our estimations can be valid currently. We selected 27 075 plots out of the 38 432 NFI plots

available, distributed over the whole French territory using the same selection criteria as for training plots (e.g. forest

types, see map Fig. S4.1). Since the sampling design of NFI plots is not homogeneous for all regions and forest types,

we weighted the time since the last harvesting estimations by the relative proportion of each forest type per forested

region (« *sylvoécorégion* ») to account for this heterogeneity.

For the prediction, the final model was simplified as the NFI data was missing some forest attributes included

in the initial model fitted on the learning dataset (e.g. tree-related microhabitats, volume of stumps). We ran a new

variable selection process with only the variables available in the NFI dataset, which led to the simplified model.

Resulting model estimates were then used for prediction, and 95% confidence intervals were calculated using the

variance covariance matrix from the fitted model.

# 3 | RESULTS

## 2.1 | Selection and description of the training dataset

In the training dataset, pure spruce and fir plots accounted for 11% of the total plots, pure beech plots accounted for 14%, mixed broadleaved 32%, mostly broadleaved [50 – 75% of broadleaved] 8%, mostly conifers [50 – 75% of coniferous] 9% and pure deciduous oaks 7% (Fig. S4.2). As a comparison, overall French production forests is comprised of 65% of broadleaved forests, 21% of coniferous forests and 11% of mixed forests (MAAF & IGN, 2016). Our modelling dataset has therefore slightly less coniferous (-8%) and slightly more broadleaved (+3%) and mixed forest types (+9%) than the average French forest.

Our training dataset plots displayed a mean volume of deadwood of 25 $m^3$/ha, which is above the national average for similar forest types - about 18 $m^3$/ha. The volume of standing deadwood was of 13 $m^3$/ha for our plots, against 8 $m^3$/ha for the equivalent National Forest Inventory plots. Finally, the mean volume of living trees on our plots was of 303 $m^3$/ha, against a national average of 208 $m^3$/ha.

Our dataset showed an overall decreasing trend for the number of plots as the time since the last harvesting got longer, and especially few plots with abandonment times over 75 years since the last harvesting (Appendix S1). The median time since the last harvesting was of 26 years.

## 3.2 | Selected model

The volume of stumps at an early decay stage was negatively correlated to the time since the last harvesting (standardised $\beta$ = -0.051, SE = 0.007) (Table 3). The second most important variable, coppice density, had a slightly milder magnitude and was positively correlated to the time since the last harvesting ($\beta$ = 0.043, SE = 0.009). Other positively correlated variables, but at a lower magnitude than coppices, were the basal area of very large trees ($\beta$ = 0.026, SE = 0.008), the volume of standing deadwood, i.e. snags and standing dead trees ($\beta$ = 0.021, SE = 0.007), the volume of downed deadwood ($\beta$ = 0.018, SE = 0.007) and the diversity of tree-related microhabitats ($\beta$ = 0.023, SE = 0.008). Furthermore, diversity of decay stages had a negative influence on the time since the last harvesting, but was only marginally significant ($\beta$ = -0.012, SE= 0.007).

Concerning the environmental covariates, soil pH had a strong positive influence on the time since the last harvesting ($\beta$ = 0.195, SE = 0.048). The same went for mean annual precipitations ($\beta$ = 0.211, SE = 0.040). Precipitations and pH were the two effects with the strongest magnitude.

247 *Table 3: Standardised estimates, standard errors, and p-values for the variables from the generalised linear mixed model with a*
248 *gamma distribution, log link and "site" random effect explaining the time since the last harvesting of French forest plots. '***' < 0.001,*
249 *'**' < 0.01, '*' < 0.05, '(*)' <0.1. Volumes and basal areas are all per hectare values.*

| | | *Standardised estimates (β)* | *Standard error (SE)* | *p-value (p)* |
|---|---|---|---|---|
| | Intercept | 3.160 | 0.093 | < 2e-16 *** |
| *Structural covariates* | Volume of stumps at early decay stages | -0.051 | 0.007 | 2.53e-13 *** |
| | Coppice density | 0.043 | 0.009 | 3.29e-06 *** |
| | Volume of standing deadwood | 0.021 | 0.007 | 0.001 ** |
| | Volume of downed deadwood | 0.018 | 0.007 | 0.011 * |
| | Diversity of tree-related microhabitats | 0.029 | 0.008 | 3.51e-04 *** |
| | Basal area of very large living trees | 0.026 | 0.008 | 0.001 ** |
| | Diversity of decay stages | -0.012 | 0.007 | 0.094 (*) |
| *Environmental covariates* | Mean annual precipitations | 0.211 | 0.040 | 1.25e-07 *** |
| | Soil pH | 0.195 | 0.048 | 5.34e-05 *** |
| *Interactions* | Volume of stumps at an early decay stage : Precipitations | 0.025 | 0.007 | 6.55e-4 *** |
| | Downed deadwood : Precipitations | -0.016 | 0.007 | 0.025 * |
| | Basal area of very large trees : Precipitations | -0.024 | 0.007 | 4.39e-4 *** |
| | Diversity of decay stages : Precipitations | -0.019 | 0.007 | 0.005 ** |
| | Volume of stumps at an early decay stage : pH | 0.013 | 0.005 | 0.015 * |
| | Volume of downed deadwood : pH | -0.012 | 0.006 | 0.044 * |
| | Precipitations : pH | -0.286 | 0.035 | < 2e-16 *** |

251 The estimates also showed that when precipitations or pH increased, the negative effect of the volume of
252 stumps on the time since the last harvesting decreased (closer to zero) ($\beta$ = 0.025, p = 0.007; $\beta$ = 0.013, SE = 0.005,

respectively) and the positive effect of the volume of downed deadwood decreased (β = -0.016, SE = 0.007 ; β = -0.012, SE = 0.006, respectively) (Figure 2). Positive effect of the basal area of very large trees also decreased with higher precipitations (β = -0.024, SE = 0.007). Furthermore, the negative influence of the diversity of decay stages on the time since the last harvesting decreased with more abundant precipitations (β = -0.019, SE = 0.007). Finally, when the soil pH increased, precipitations had a decreasing positive effect on the time since the last harvesting (β = -0.286, SE = 0.035).

*Figure 2 : Illustration of the interactions between volume of downed deadwood (β = -0.012, p = 0.044) and volume of stumps at an early decay stage (β = 0.013, p = 0.015) with soil pH. Lines represent a generalised linear model with a log link and 95% confidence interval for pH values over and under 6.*

3.3 | Model validation

The distribution of model residuals as a function of the predicted time since the last harvesting showed relative homogeneity, except for a few plots with overestimated time since the last harvesting (Fig. S5. 1). When checked, the residual distribution as a function of the variables both included and excluded from the model showed no noticeable patterns, except for some badly estimated time since the last harvesting on the lower tail of the x axis for the variables "volume of stumps at early decay stages", "volume of downed deadwood", "volume of standing deadwood", and "coppice density".

Reserves were removed one at a time from the training data to check for any particularly influential sites regarding the estimated values. Cross validation results showed that predictions were relatively constant, with an r² linking the predicted values to the observed values varying between 0.756 and 0.900 over 500 simulations (Fig. S5.2). The estimated values were quite stable except for the covariates soil pH and precipitation, as well as for the interaction between those two variables for which estimates slightly varied during simulations, but to a negligible extent (Fig. S5. 3).

3.4 | Model projection on the National Forest Inventory dataset

The simplified model used for projection (see Table S5.1) was comprised of four structural features: the
volume of standing and large downed deadwood, coppice density and basal area of very large trees, as well as the two
environmental covariates soil pH and precipitations. As for interactions, downed deadwood and basal area of very
large trees interacted negatively with precipitations, and coppice density interacted negatively with soil pH. Finally,
there was still a negative interaction between soil pH and precipitations.

We predicted an average time since the last harvesting of 26 years for the NFI plots issued from the 2012 to
2018 campaigns (including only the six forest types kept for modelling). According to this projection, about 3.1% of
French metropolitan forest has reached or surpassed 50 years of abandonment. 43% of forests were predicted to have
a time since the last harvesting of between 26 and 50 years. When varying the maturity threshold from 30 to 300 years
of abandonment, we observe the decreasing trend presented in Figure 3a. The confidence interval is very large, around
30 years without harvesting and gets narrower with higher thresholds. There are very few forests predicted above 75
years without harvesting, which is in accordance with the distribution of the time since the last harvesting in the
training dataset (French forest reserves). Plots with longer predicted time since the last harvesting are mainly
distributed in mountainous regions of eastern France (Vosges, southern Alps and Jura), as well as in the south of France
(Pyrenees and Cevennes) and Corsica (Figure 3d).

Figure 3 : (a) represents how the maturity threshold changes the predicted proportion of mature forests. The ribbon represents the 95% confidence interval. The 50-year threshold (vertical bar) is the one we chose for this study. The confidence interval narrows very fast and at 50 years without harvesting it ranges from 1 to 9%. (b) NFI plots with predicted time since the last harvesting under 26 years, (c) NFI plots with predicted time since the last harvesting between 26 and 50 years, (d) NFI plots with predicted time since the last harvesting over 50 years.

# 4 | DISCUSSION

We evaluated French forest's state of maturity using time since the last harvesting as a proxy for maturity. We showed that most old growth variables highlighted in the literature did indeed explain time since the last harvesting quite well, and other variables translating legacy of past forest management also played a role in the modelling approach. Our projection gives the first robust statistical estimate of the proportion of mature forests in metropolitan France and may serve to report on the status of mature forests here. Our multivariate modelling approach is quite original, as previous studies (e.g. (Larrieu et al., 2019; Paillet et al., 2015; Vandekerkhove, 2005)) used the opposite reasoning, i.e. modelling structural attributes as a function of the time since the last harvesting.

## 4.1 | Influence of parameters on the models

As expected, the volume of stumps at early decay stages was negatively correlated with the time since the last harvesting (Paillet et al., 2015; Siitonen et al., 2000). When plots are unharvested for a long period of time, stumps that originated from human disturbance decompose, hence the decline in abundance of fresh stumps in more mature forests. Stumps at an early decay stage are characteristic of recent harvesting, and their absence is a token of longer abandonment times.

Higher basal area of very large living trees is also a known characteristic of mature forests (Bauhus et al., 2009; Burrascano et al., 2013; Paillet et al., 2015). Additionally, larger trees often bear higher diversity of tree-related microhabitats and therefore promote biodiversity at the stand level (Larrieu et al., 2018; Paillet et al., 2017): tree-related microhabitats are the result of tree alteration by biotic and abiotic processes (Larrieu et al., 2018), their presence is thus mostly related to the timespan during which trees are subjected to those processes and thus to time since the last harvesting.

We also expected the diversity of decay stages on standing and downed deadwood to be positively correlated to the time since the last harvesting. Although it was, the observed effect was only marginally significant. Wood decomposition dynamics depends on other factors than time only, such as environmental conditions prone to microorganisms (Herrmann & Bauhus, 2013), the chemical composition of wood (i.e. species), its diameter, micro-climatic conditions as well as edaphic parameters (Heiri et al., 2009; Larrieu et al., 2019; Přívětivý et al., 2018; Zell et al., 2009). Those characteristics were not selected in the model interactions, it therefore seems that there is no univocal correlation between the time since the last harvesting and decay stages, hence the marginal effect observed here. Furthermore, Wirth et al. (2009) suggested that deadwood volume and decay stages could potentially be misleading and could not be taken as sole indicators of maturity. Indeed, deadwood dynamics follow a U-shaped pattern as a result of stand succession (Aakala, 2010), but some punctual disturbances such as storms or droughts can result in locally high amounts of deadwood (Aakala, 2010; Harmon, 2009). In our case, some plots have been subjected to major disturbances, e.g. the 1999 storms that caused damage to a large portion of the French forests, which could still bear the legacy of those events. Our results confirm that deadwood is a key component of mature French forest

reserves, but that the relationship between decay stages and the time since the last harvesting might be more complex than anticipated and would require non-linear relationships or more information on individual plot to better characterise it.

Finally, coppice density was strongly and positively correlated to the time since the last harvesting. Coppicing is a legacy of past management for fuel (heating) and is a practice that has decreased over the last century (IGN, 2016b). Many coppice-with-standards plots have now been abandoned (Unrau et al., 2018) resulting in what Lassauce et al. (2012) call "coppice-with-standard stands with an overmature component". In their study, they also found a significant increase in the basal area of large living trees and downed deadwood for more mature plots with a history of coppicing. This trend seems widespread among stands with a history of coppice-with-standard management. Indeed, (Becker et al., 2017) also noticed that after 40 years of abandonment of coppicing, legacies of this practice still remained in species composition. These mature coppices could constitute another form of mature forests (Lassauce et al., 2012), and it would be interesting to consider coppice density as another indicator for forest maturity, especially considering coppice forests currently represent up to 15% of Europe's forest resources (Unrau et al., 2018).

Environmental context (soil pH and annual precipitations) interacted significantly with stand characteristics in the model, notably interactions between deadwood volumes and pH/precipitations. Plots with higher pH and higher precipitations, where high basal areas and higher stem densities have the potential to be reached sooner, are sites with higher productivity. Nonetheless, we observed a negative interaction between soil pH and precipitations, meaning that overall, productive sites show shorter times since the last harvesting. Rainier plots and/or plots with higher pH conditions show lessened effects of the volume of stumps, the diversity of decay stages, and the volume of downed deadwood and basal area of very large trees on the time since the last harvesting. These interactions could be linked to sites with higher amounts of precipitations being more attractive for harvesting. Surprisingly, elevation was not selected in favour of mean annual precipitations (which increased model AIC of 22 points compared to elevation). Precipitation was nonetheless strongly correlated to elevation (correlation coefficient = 0.9). Precipitations seem to have a broader role and a higher explanatory power than elevation, since they condition both biotic (tree growth, decay rates, soil properties) and abiotic (atmospheric humidity) phenomena.

4.2 | The projected proportion and rough distribution of mature forests

Our model predicted that 3.1% of the national forested area has a time since the last harvesting above 50 years (95% confidence interval ranges from 1 to 9 %). This prediction is valid only for the six forest types kept for modelling (pure beech, pure deciduous oak, mixed broadleaved, pure spruce and/or fir, mostly broadleaved and mostly coniferous types). The prediction is in accordance with the early expertise that had been proposed in 1993. Forested areas abandoned more than 50 years ago seem to occur mostly in more remote and mountainous areas in southern and eastern France (Figure 3c). Indeed, it has been shown that accessibility and favourable topographic, climatic, and soil conditions characterise intensively managed areas, whereas de-intensification and abandonment trends occur in more marginal areas (Levers et al., 2014, 2018; Sabatini et al., 2018). Those highlighted areas therefore

show high conservation potential. More interestingly, our prediction also showed evidence of a large proportion of French forest with abandonment times comprised between 26 and 50 years, with "hotspots" such as Corsica, Brittany and middle-eastern France. Those forests could - in a relatively close future - display interesting conservation attributes which would deserve more attention, especially if some larger continuous and connected areas, particularly interesting for biodiversity, could be restored or managed sensibly (Bauhus et al., 2009; Portier et al., 2020; Sabatini et al., 2020). Indeed, French metropolitan forested surface area has grown since the last century and now occupies 31% of the metropolitan France or about 16.9 million hectares (IGN, 2018). Whereas strict forest reserves compose a mere 0.15% of the French forests (Cateau et al., 2017). Private forests represent 75% of the French forested area (MAAF & IGN, 2016) and considering that the harvesting in these - often small and fragmented - properties is usually low or inexistent, they could potentially account for this large portion of "in between" forest in our prediction. Those areas are not accounted for in most studies and conservation decisions. If we consider that each country is equally responsible for contributing to the "EU Biodiversity Strategy for 2030" goal for 10% of strictly protected land (European Commission, 2020), then France, with 3% of estimated mature forest – not all protected and probably quite fragmented - still has a long way to go. But the recognition that areas with high potential for achieving this goal exist and are identified, is a first step forward.

4.3 | Limits and perspectives to our modelling approach

The distribution of mature forests predicted by our model should not be taken as a precise map but as an overview of the state of French forests to this day. Indeed, the plot density is quite scarce with 27 075 plots for the whole French territory. The time since the last harvesting used for training the model could introduce some error, since the older the last known harvesting the more uncertain the estimation of the time since the last harvesting. Also, our model did not consider the intensity of harvesting (e.g. volume of wood harvested, type of management), although this knowledge would probably further refine our predictions. Additionally, removing covariates for the predictive model probably hampers the performance of the prediction, which would be more precise with information about volume of stumps and tree-related microhabitats. Nonetheless, large areas with potential for hosting forests with high conservation value were highlighted, and further validation, e.g. by crossing our findings with maps and data on previously studied mature sites, along with field validation, would enable us to refine our knowledge of the distribution of more mature forests, and evaluate the precision of our predictions.

The overall results of our study are encouraging, in the sense that a large portion of the French forest has potential to attain interesting states of maturity in a close future and therefore to contribute to the race against climate change, by acting as carbon and biodiversity pools (Portier et al., 2020; Sabatini et al., 2020; Carey et al., 2001). We expect our results to be used in ensuring that the existing 3% of mature forest are properly managed and protected, and that our work will constitute a stepping-stone to further refining our knowledge of the state of maturity of French forests.

## 5 | ACKNOWLEDGEMENTS

This study is issued of the Master's training period of LT and was funded by the French Foundation for Biodiversity Research ("Appel Masters 2019", www.fondationbiodiversite.fr) and the Direction for Water and Biodiversity of the French ministry in charge of the environment (convention "FORETMAT"). We are in debt to the reserve and forest managers who fed the databases and made this study possible. Without their daily commitment and implication in the forest surveys, we would not have reached such complete and global results. We are also grateful to F. Benest (IGN) and B. Courbaud (INRAE) for constructive discussions on the different parts of this work.

## 6 | DATA ACCESSIBILITY

The two datasets we worked with for this study were the Nature Forest Reserve data (PSDRF), ONF-RNF and the French national forest inventory data (raw data, annual surveys from 2005 to 2018, https://inventaire-forestier.ign.fr/spip.php?rubrique159).

## 7 | REFERENCES

Aakala, T. (2010). Tree mortality and deadwood dynamics in late-successional boreal forests. *Dissertationes Forestales*, *2010*(100). https://doi.org/10.14214/df.100

Achard, F., & Hansen, M. C. (2012). *Global Forest Monitoring from Earth Observation*. CRC Press.

Bauhus, J., Puettmann, K., & Messier, C. (2009). Silviculture for old-growth attributes. *Forest Ecology and Management*, *258*(4), 525–537. https://doi.org/10.1016/j.foreco.2009.01.053

Becker, T., Spanka, J., Schröder, L., & Leuschner, C. (2017). Forty years of vegetation change in former coppice-with-standards woodlands as a result of management change and N deposition. *Applied Vegetation Science*, *20*(2), 304–313. https://doi.org/10.1111/avsc.12282

Brockerhoff, E. G., Barbaro, L., Castagneyrol, B., Forrester, D. I., Gardiner, B., González-Olabarria, J. R., Lyver, P. O., Meurisse, N., Oxbrough, A., Taki, H., Thompson, I. D., van der Plas, F., & Jactel, H. (2017). Forest biodiversity, ecosystem functioning and the provision of ecosystem services. *Biodiversity and Conservation*, *26*(13), 3005–3035. https://doi.org/10.1007/s10531-017-1453-2

Burrascano, S., Keeton, W. S., Sabatini, F. M., & Blasi, C. (2013). Commonality and variability in the structural

attributes of moist temperate old-growth forests: A global review. *Forest Ecology and Management*, *291*,

458–479. https://doi.org/10.1016/j.foreco.2012.11.020

Carey, E. V., Sala, A., Keane, R., & Callaway, R. M. (2001). Are old forests underestimated as global carbon sinks?

*Global Change Biology*, *7*(4), 339–344. https://doi.org/10.1046/j.1365-2486.2001.00418.x

Cateau, E., Duchamp, L., Garrigue, J., Gleizes, L., Tournier, H., & Debaive, N. (2017). *Le patrimoine forestier des*

*réserves naturelles—Focus sur les forêts à caractère naturel* (p. 104). Réserves Naturelles de France.

http://www.reserves-naturelles.org/sites/default/files/librairie/cahier7hd.pdf

Cateau, E., Larrieu, L., Vallauri, D., Savoie, J.-M., Touroult, J., & Brustel, H. (2015). Ancienneté et maturité: Deux

qualités complémentaires d'un écosystème forestier. *Comptes Rendus Biologies*, *338*(1), 58–73.

https://doi.org/10.1016/j.crvi.2014.10.004

Chiarucci, A., & Piovesan, G. (2020). Need for a global map of forest naturalness for a sustainable future.

*Conservation Biology*, *34*(2), 368–372. https://doi.org/10.1111/cobi.13408

Christensen, M., Hahn, K., Mountford, E. P., Ódor, P., Standovár, T., Rozenbergar, D., Diaci, J., Wijdeven, S., Meyer,

P., Winter, S., & Vrska, T. (2005). Dead wood in European beech (Fagus sylvatica) forest reserves. *Forest*

*Ecology and Management*, *210*(1), 267–282. https://doi.org/10.1016/j.foreco.2005.02.032

European Commission. (2020). *EU Biodiversity Strategy for 2030* (p. 23). European Commission. https://eur-

lex.europa.eu/resource.html?uri=cellar:a3c806a6-9ab3-11ea-9d2d-

01aa75ed71a1.0001.02/DOC_1&format=PDF

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate surfaces for global land areas.

*International Journal of Climatology*, *37*(12), 4302–4315. https://doi.org/10.1002/joc.5086

Forest Europe. (2015). *State of Europe's forests*. https://www.foresteurope.org/docs/fullsoef2015.pdf

Frey, S. J. K., Hadley, A. S., Johnson, S. L., Schulze, M., Jones, J. A., & Betts, M. G. (2016). Spatial models reveal the

microclimatic buffering capacity of old-growth forests. *Science Advances*, *2*(4), e1501392.

https://doi.org/10.1126/sciadv.1501392

Harmon, M. E. (2009). *Woody Detritus Mass and its Contribution to Carbon Dynamics of Old-Growth Forests: The*

*Temporal Context*. 32.

Harrel, F. E. (2015). *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal*

*Regression, and Survival Analysis*. Springer. https://link.springer.com/book/10.1007/978-3-319-19425-7

Heiri, C., Wolf, A., Rohrer, L., & Bugmann, H. (2009). Forty years of natural dynamics in Swiss beech forests:

Structure, composition, and the influence of former management. *Ecological Applications*, *19*(7), 1920–1934.

https://doi.org/10.1890/08-0516.1

Herrmann, S., & Bauhus, J. (2013). Effects of moisture, temperature and decomposition stage on respirational

carbon loss from coarse woody debris (CWD) of important European tree species. *Scandinavian Journal of*

*Forest Research*, *28*(4), 346–357. https://doi.org/10.1080/02827581.2012.747622

Hijmans, R. J. (2020). *Package "raster"* (3.3-13) [Computer software]. https://cran.r-

project.org/web/packages/raster/raster.pdf

IGN. (2016a). *BD forêt® Version 2.0—Descriptif de contenu* (p. 51). IGN. https://inventaire-

forestier.ign.fr/IMG/pdf/dc_bdforet_2-0.pdf

IGN. (2016b). *Indicateurs de gestion durable des forêts françaises métropolitaines—Edition 2015*. *38*, 12.

https://inventaire-forestier.ign.fr/IMG/pdf/IF38.pdf

IGN. (2018). *La forêt Française—Etat des lieux et évolutions récentes—Panorama des résultats de l'inventaire*

*forestier* (Edition 2018; p. 56). https://inventaire-forestier.ign.fr/IMG/pdf/180906_publiff_bd.pdf

Larrieu, L., Cabanettes, A., Gouix, N., Burnel, L., Bouget, C., & Deconchat, M. (2019). Post-harvesting dynamics of the

deadwood profile: The case of lowland beech-oak coppice-with-standards set-aside stands in France.

*European Journal of Forest Research*, *138*(2), 239–251. https://doi.org/10.1007/s10342-019-01164-8

Larrieu, L., Paillet, Y., Winter, S., Bütler, R., Kraus, D., Krumm, F., Lachat, T., Michel, A. K., Regnery, B., &

Vandekerkhove, K. (2018). Tree related microhabitats in temperate and Mediterranean European forests: A

hierarchical typology for inventory standardization. *Ecological Indicators*, *84*, 194–207.

https://doi.org/10.1016/j.ecolind.2017.08.051

Lassauce, A., Anselle, P., Lieutier, F., & Bouget, C. (2012). Coppice-with-standards with an overmature coppice

component enhance saproxylic beetle biodiversity: A case study in French deciduous forests. *Forest Ecology*

*and Management*, *266*, 273–285. https://doi.org/10.1016/j.foreco.2011.11.016

Levers, C., Müller, D., Erb, K., Haberl, H., Jepsen, M. R., Metzger, M. J., Meyfroidt, P., Plieninger, T., Plutzar, C., Stürck,

J., Verburg, P. H., Verkerk, P. J., & Kuemmerle, T. (2018). Archetypical patterns and trajectories of land

systems in Europe. *Regional Environmental Change*, *18*(3), 715–732. https://doi.org/10.1007/s10113-015-

0907-x

Levers, C., Verkerk, P. J., Müller, D., Verburg, P. H., Butsic, V., Leitão, P. J., Lindner, M., & Kuemmerle, T. (2014).

Drivers of forest harvesting intensity patterns in Europe. *Forest Ecology and Management*, *315*, 160–172.

https://doi.org/10.1016/j.foreco.2013.12.030

L'IF n°14. (2011). *Une nouvelle partition écologique et forestière du territoire métropolitain: Les sylvoécorégions*

*(SER)*. 8. https://inventaire-forestier.ign.fr/IMG/pdf/IF_SER_web.pdf

MAAF, & IGN. (2016). *Indicateurs de gestion durable des forêts françaises métropolitaines—Edition 2015—Résultats*

(p. 243). Maaf - IGN. https://inventaire-forestier.ign.fr/IMG/pdf/IGD_2015_FR.pdf

MAAPRAT-IFN. (2011). *Indicateurs de gestion durable des forêts françaises métropolitaines—Edition 2010* (p. 202).

Inventaire Forestier National. https://inventaire-forestier.ign.fr/IMG/pdf/IGD2010-FR.pdf

Mountford, E. P. (2002). Fallen dead wood levels in the nearnatural beech forest at La Tillaie reserve, Fontainebleau,

France. *Forestry*, *75*(2), 203–208. https://doi.org/10.1093/forestry/75.2.203

Paillet, Y., Archaux, F., Boulanger, V., Debaive, N., Fuhr, M., Gilg, O., Gosselin, F., & Guilbert, E. (2017). Snags and

large trees drive higher tree microhabitat densities in strict forest reserves. *Forest Ecology and Management*,

*389*, 176–186. https://doi.org/10.1016/j.foreco.2016.12.014

Paillet, Y., Debaive, N., Archaux, F., Cateau, E., Gilg, O., & Guilbert, E. (2019). Nothing else matters? Tree diameter

and living status have more effects than biogeoclimatic context on microhabitat number and occurrence: An

analysis in French forest reserves. *PLOS ONE*, *14*(5), e0216500.

https://doi.org/10.1371/journal.pone.0216500

Paillet, Y., Pernot, C., Boulanger, V., Debaive, N., Fuhr, M., Gilg, O., & Gosselin, F. (2015). Quantifying the recovery of

old-growth attributes in forest reserves: A first reference for France. *Forest Ecology and Management*, *346*,

51–64. https://doi.org/10.1016/j.foreco.2015.02.037

Pontailler, J.-Y., Faille, A., & Lemée, G. (1997). Storms drive successional dynamics in natural forests: A case study in

Fontainebleau forest (France). *Forest Ecology and Management*, *98*(1), 1–15.

https://doi.org/10.1016/S0378-1127(97)00073-X

519 Portier, J., Wunder, J., Stadelmann, G., Zell, J., Abegg, M., Thürig, E., & Rohner, B. (2020). 'Latent reserves': A hidden

520  treasure in National Forest Inventories. *Journal of Ecology*, 1365-2745.13487. https://doi.org/10.1111/1365-

521  2745.13487

522 Přívětivý, T., Adam, D., & Vrška, T. (2018). Decay dynamics of Abies alba and Picea abies deadwood in relation to

523  environmental conditions. *Forest Ecology and Management*, *427*, 250–259.

524  https://doi.org/10.1016/j.foreco.2018.06.008

525 Rossi, M., & Vallauri, D. (2013). *Evaluer la naturalité. Guide pratique version 1.2* (p. 154). WWF.

526  http://rgdoi.net/10.13140/2.1.4973.8565

527 Sabatini, F. M., Burrascano, S., Keeton, W. S., Levers, C., Lindner, M., Pötzschner, F., Verkerk, P. J., Bauhus, J.,

528  Buchwald, E., Chaskovsky, O., Debaive, N., Horváth, F., Garbarino, M., Grigoriadis, N., Lombardi, F., Marques

529  Duarte, I., Meyer, P., Midteng, R., Mikac, S., … Kuemmerle, T. (2018). Where are Europe's last primary

530  forests? *Diversity and Distributions*, *24*(10), 1426–1439. https://doi.org/10.1111/ddi.12778

531 Sabatini, F. M., Keeton, W. S., Lindner, M., Svoboda, M., Verkerk, P. J., Bauhus, J., Bruelheide, H., Burrascano, S.,

532  Debaive, N., Duarte, I., Garbarino, M., Grigoriadis, N., Lombardi, F., Mikoláš, M., Meyer, P., Motta, R.,

533  Mozgeris, G., Nunes, L., Ódor, P., … Kuemmerle, T. (2020). Protection gaps and restoration opportunities for

534  primary forests in Europe. *Diversity and Distributions*, ddi.13158. https://doi.org/10.1111/ddi.13158

535 Siitonen, J., Martikainen, P., Punttila, P., & Rauh, J. (2000). Coarse woody debris and stand characteristics in mature

536  managed and old-growth boreal mesic forests in southern Finland. *Forest Ecology and Management*, *128*(3),

537  211–225. https://doi.org/10.1016/S0378-1127(99)00148-6

538 Spies, T. A., & Franklin, J. F. (1991). *The Structure of Natural Young, Mature, and Old-Growth Douglas-Fir Forests in*

539  *Oregon and Washington*. 21.

540 Toïgo, M., Vallet, P., Perot, T., Bontemps, J.-D., Piedallu, C., & Courbaud, B. (2015). Overyielding in mixed forests

541  decreases with site productivity. *Journal of Ecology*, *103*(2), 502–512. https://doi.org/10.1111/1365-

542  2745.12353

543 Unrau, A., Becker, G., Spinelli, R., Lazdina, D., Magagnotti, N., Nicolescu, V.-N., Buckley, P., Bartlett, D., & Kofman, P.

544  D. (Eds.). (2018). *Coppice Forests in Europe* (1. Auflage). Albert-Ludwigs-Universität Freiburg.

545 Vandekerkhove, K. (2005). Spontaneous re-establishment of natural structure and related biodiversity in a previously

546  managed beech forest in Belgium after 20 years of non intervention. *For. Snow Landsc. Res.*, 12.

Vandekerkhove, K., De Keersmaeker, L., Menke, N., Meyer, P., & Verschelde, P. (2009). When nature takes over from

man: Dead wood accumulation in previously managed oak and beech woodlands in North-western and

Central Europe. *Forest Ecology and Management*, *258*(4), 425–435.

https://doi.org/10.1016/j.foreco.2009.01.055

Winter, S., & Möller, G. C. (2008). Microhabitats in lowland beech forests as monitoring tool for nature conservation.

*Forest Ecology and Management*, *255*(3–4), 1251–1261. https://doi.org/10.1016/j.foreco.2007.10.029

Wirth, C., Messier, C., Bergeron, Y., Frank, D., & Fankhänel, A. (2009). Old-Growth Forest Definitions: A Pragmatic

View. In C. Wirth, G. Gleixner, & M. Heimann (Eds.), *Old-Growth Forests* (Vol. 207, pp. 11–33). Springer Berlin

Heidelberg. https://doi.org/10.1007/978-3-540-92706-8_2

Zell, J., Kändler, G., & Hanewinkel, M. (2009). Predicting constant decay rates of coarse woody debris—A meta-

analysis approach with a mixed model. *Ecological Modelling*, *220*(7), 904–912.

https://doi.org/10.1016/j.ecolmodel.2009.01.020

Zuur, A. F., & Ieno, E. N. (2016). A protocol for conducting and presenting results of regression-type analyses.

*Methods in Ecology and Evolution*, *7*(6), 636–645. https://doi.org/10.1111/2041-210X.12577

Zuur, A. F., Ieno, E. N., & Elphick, C. S. (2010). A protocol for data exploration to avoid common statistical problems:

Data exploration. *Methods in Ecology and Evolution*, *1*(1), 3–14. https://doi.org/10.1111/j.2041-

210X.2009.00001.x

# 8 | SUPPLEMENTARY MATERIALS

*Appendix S1: Distribution of time since the last harvesting of the PSDRF plots coloured by forest type.*

Appendix S2: Protocol for the French nature forest reserves stand structure description

On each circular plot, forest stand structure was characterized using the following protocol (Paillet et al., 2015, 2019):

- All living trees larger than 7.5 cm were measured considering, by checking if they issued from the same base or not, the difference between coppice and standard trees.

- For living trees with a diameter at breast height (DBH) above 30 cm, a fixed angle plot method was used to select the individuals within a relascopic angle of 3%. Practically, this meant that sampling distance was proportional to the apparent DBH of a tree. For example, a tree with a DBH of 60 cm was included in the sample if it was within 20 m of the center of the plot. This technique allowed us to better account for larger trees at a small scale.

- All other variables were measured on fixed-area plots. Within a fixed 10-m (314 m2) radius plot, the diameter of all living trees and standing deadwood (stumps, snags (height > 130 cm), standing dead trees) was measured from 7.5 to 30 cm DBH.

- Within a 20-m radius (1256 m2), all standing deadwood and downed deadwood (logs) with a diameter > 30 cm were recorded.

- Downed deadwood (logs) with a diameter under 30 cm was recorded along three separate 20 m long radial transects (every 127°).

- All standing trees (alive and dead) were visually inspected for the recording of tree-related microhabitats. Observers attended a training session and were given a field guide with pictures to help them better determine microhabitat types and detailed criteria to include in the inventories.

On every piece of deadwood decay stage of the bark and of the wood were recorded, respectively on a scale of one to four and one to five. Whenever possible, all trees, both alive and dead, were identified to species level.

*Appendix S3: Table presenting the classification used to derive plot forest types from plot species composition. This classification is inspired from the "BD forêt version 2" classification, which is based on the species' relative tree cover. We chose to use tree basal area instead, as tree cover was not unknown in our dataset. Each plot forest type is defined as the proportion of a specific species' basal area on the said plot. Therefore, plots are called "pure" if over 75% of the total plot basal area is occupied by one (or two in a few cases) species. We then called mixed any plots where a mixture of two or more species occupied over 75% of the total basal area. Finally, the last two broader forest types correspond to plots where a true mixture of tree species occur and the only definition we can give is if either broadleaved or coniferous trees are preponderant. In the end, only the six highlighted types out of the 22 included enough plots to be kept for modelling.*

| Code *BD forêt version 2* | Class | Label | Definition | Species |
|---|---|---|---|---|
| FF1G01-01 | Broadleaved | Pure deciduous oak | Basal area of deciduous oaks ≥ 75% | *Quercus cerris, Quercus rosacea, Quercus robur, Quercus pubescens, Quercus rubra, Quercus petraea, Quercus tauza* |
| FF1G06-06 | Broadleaved | Pure evergreen oak | Basal area of evergreen oaks ≥ 75% | *Quercus suber, Quercus ilex* |
| FF1-09-09 | Broadleaved | Pure beech | Basal area of beech trees ≥ 75% | *Fagus sylvatica* |
| FF1-10-10 | Broadleaved | Pure chestnut | Basal area of chestnut trees ≥ 75% | *Castanea sativa* |
| FF1-14-14 | Broadleaved | Pure black locust | Basal area of black locust trees ≥ 75% | *Robinia pseudoacacia* |
| FF1-49-49 | Broadleaved | Other pure broadleaved trees | Basal area of a single other broadleaved species ≥ 75% | Ash tree (*Fraxinus sp*), Maple trees (*Acer sp*), Birch (*Betula sp*), Charm (*Carpinus betulus*), Alder (*Alnus sp*), Hazel tree (*Corylus sp*) |
| FF1-00-00 | Broadleaved | Mixed broadleaved trees | Combined basal area of at least two broadleaved species that are not either deciduous oak or evergreen oak ≥ 75% (without any single species occupying a basal area ≥ 75%) | *Fagus sylvatica, Castanea sativa, Robinia pseudoacacia,* Ash trees (*Fraxinus sp*), Maple trees (*Acer sp*), Birch (*Betula sp*), Charm (*Carpinus betulus*), Alder (*Alnus sp*), Hazel tree (*Corylus sp*) |
| FF2-51-51 | Coniferous | Pure maritime pine | Basal area of maritime pine ≥ 75% | *Pinus pinaster* |
| FF2-52-52 | Coniferous | Pure scots pine | Basal area of scots pine ≥ 75% | *Pinus sylvestris* |
| FF2G53-53 | Coniferous | Pure laricio pine and/or black pine | Basal area of laricio pine and/or black pine ≥ 75% (can be either a mixture of both species or each species individually) | *Pinus nigra* |

| Code *BD forêt version 2* | Class | Label | Definition | Species |
|---|---|---|---|---|
| FF2-57-57 | Coniferous | Pure Aleppo pine | Basal area of Aleppo pine ≥ 75% | *Pinus halepensis* |
| FF2G58-58 | Coniferous | Pure mountain pine and/or swiss pine | Basal area of laricio pine and/or black pine ≥ 75% (can be either a mixture of both species or each species individually) | *Pinus mugo, Pinus cembra* |
| FF2-81-81 | Coniferous | Other pure pine | Basal area of another pine species ≥ 75% | *Pinus mugo, Pinus pinea* |
| FF2-80-80 | Coniferous | Mixed pine | Combined basal area of at least two pine species ≥ 75% (without being a mixture of black pine and laricio pine or of mountain pine and swiss pine and without any single species occupying a basal area ≥ 75%) | *Pinus pinaster, Pinus sylvestris, Pinus nigra, Pinus halepensis, Pinus mugo, Pinus cembra, Pinus mugo, Pinus pinea* |
| FF2G61-61 | Coniferous | Pure spruce and/or fir | Basal area of spruce and/or fir ≥ 75% (can be either a mixture of both species or each species individually) | *Abies cephalonica, Abies nordmanniana, Abies grandis, Abies alba, Picea abies, Picea sitchensis* |
| FF2-63-63 | Coniferous | Pure larch trees | Basal area of larch trees ≥ 75% | *Larix decidua* |
| FF2-64-64 | Coniferous | Pure douglas fir | Basal area of Douglas firs ≥ 75% | *Pseudotsuga menziesii* |
| FF2-91-91 | Coniferous | Other pure coniferous trees | Basal area of a single other coniferous species ≥ 75% | Cedar trees *(Cedrus atlantica, Cedrus libani, Thuja sp)* Cypress *(Cupressus sp)* Yew *(Taxus baccata)* |
| FF2-90-90 | Coniferous | Other mixed coniferous trees | Basal area of a mixture of at least two coniferous species which are not either a mixture of pine trees or spruce and fir ≥ 75% (without any single species occupying a basal area ≥ 75%) | Spruce, fir, cedar, cypress, yew (see above) |
| FF2-00-00 | Coniferous | Mixed pine and other coniferous trees | Basal area of a mixture of at least a pine tree and another coniferous species ≥ 75% (without any | *All coniferous species above* |

| Code *BD forêt version 2* | Class | Label | Definition | Species |
|---|---|---|---|---|
| | | | single species occupying a basal area ≥ 75%) | |
| FF31 | Broadleaved /Coniferous | Mostly broadleaved trees (mixture of broadleaved and coniferous trees) | 50% ≤ Basal area of broadleaved trees ≤ 75% | All broadleaved and coniferous species above |
| FF32 | Broadleaved /Coniferous | Mostly coniferous trees (mixture of broadleaved and coniferous trees) | 50% ≤ Basal area of coniferous trees ≤ 75% | All broadleaved and coniferous species above |

| Code *BD forêt version 2* | Class | Label | Definition | Species |
|---|---|---|---|---|

  *Appendix S4 : maps and graphs of the two datasets*

610  *Fig. S4.1: Map of the 27 075 National Forest Inventory plots from the 2012 to 2018 campaigns in metropolitan France which served*
611  *as the spatial prediction dataset. Plots are coloured by forest type.*

## Map of National Forest Inventory plots
## from campaigns 2012 to 2018

### Forest type
- Pure deciduous oak
- Mostly coniferous trees
- Mostly broadleaved trees
- Mixed broadleaved trees
- Pure beech trees
- Pure spruce and/or fir

613 *Fig. S4.2: Six most abundant forest types among the 4728 ploys kept from the PSDRF data*
614     *" Pure » types: species group represents over 75% of total basal area,*
615     *« Mostly » types: > 50% of total basal area,*
616     *« mixed » types: > 75% of total basal area*
617

618
619 *Appendix S 5 : model validation*

620 *Fig. S5. 1 : Distribution of the Pearson residuals from the generalised linear mixed model explaining the time since the last harvesting*
621 *using structural features and environmental variables. The trend is satisfyingly neutral, although a few plots have overestimated time*
622 *since the last harvesting.*

623
624
625

626 *Fig. S5.2: Distribution of the values of the correlation coefficient r squared over the 500 simulations of cross validation. The r squared*
627 *varied between 0.756 and 0.900.*

629     *Fig. S5. 3 : Estimate variation (mean and standard deviation) after 500 cross validation simulations.*

Table S5.1: Standardised estimates, standard errors and p-values rounded to three decimal places for the variables from the simplified generalised linear mixed model with a gamma distribution, log link and "site" random effect explaining the time since the last harvesting of French forest plots. '***' < 0.001, '**' < 0.01, '*' < 0.05, '(*)' <0.1. Volumes and basal areas are all per hectare values.

| | | Standardized estimates ($\beta$) | Standard error | p-value (p) |
|---|---|---|---|---|
| | Intercept | 3.172 | 0.095 | < 2e-16 *** |
| Structural features | Coppice density | 0.052 | 0.006 | 8.71e-8 *** |
| | Volume of standing deadwood | 0.024 | 0.006 | 1.36e-4 *** |
| | Volume of large downed deadwood | 0.020 | 0.006 | 1.48e-3 *** |
| | Basal area of very large trees | 0.029 | 0.008 | 2.89e-04 *** |
| Environmental covariates | Mean annual precipitations | 0.244 | 0.040 | 1.17e-09 *** |
| | Soil pH | 0.191 | 0.049 | 9.08e-05 *** |
| Interactions | Large downed deadwood : Precipitations | -0.010 | 0.006 | 7.98e-2(*) |
| | Basal area of very large trees : Precipitations | -0.025 | 0.007 | 2.65e-04 *** |

| | | | |
|---|---|---|---|
| Coppice density : pH | -0.018 | 0.011 | 9.59e-2 (*) |
| Precipitations : pH | -0.324 | 0.034 | < 2e-16 *** |