Genetic models of latent tuberculosis in mice reveal differential control by innate and adaptive immunity

Hongwei Su, Kan Lin, Divya Tiwari, Claire Healy, Carolina Trujillo, Yao Liu, Dirk Schnappinger, and Sabine Ehrt

Department of Microbiology and Immunology, Weill Cornell Medicine, New York, NY

Correspondence to Sabine Ehrt: sae2004@med.cornell.edu; Dirk Schnappinger: dis2003@med.cornell.edu

D. Tiwari's current affiliation is: Centre for Global Health, Barts and The London School of Medicine and Dentistry, Queen Mary University of London, Whitechapel, London, UK

C. Healy's current affiliation is: School of Medicine, Trinity College Dublin, Dublin, Ireland

Abstract

Studying latent *Mycobacterium tuberculosis* (Mtb) infection has been limited by the lack of a suitable mouse model. We discovered that transient depletion of protein biotin ligase (BPL) and thioredoxin reductase (TrxB2) results in latent infections during which Mtb cannot be detected but that relapse in a subset of mice. The immune requirements for Mtb control during latency, and the frequency of relapse, were strikingly different depending on how latency was established. TrxB2 depletion resulted in a latent infection that required adaptive immunity for control and reactivated with high frequency, whereas latent infection after BPL depletion was dependent on innate immunity and rarely reactivated. We identified immune signatures of T cells indicative of relapse and demonstrated that BCG vaccination failed to protect mice from TB relapse. These reproducible genetic latency models allow investigating the host immunological determinants that control the latent state and offer opportunities to evaluate therapeutic strategies in settings that mimic aspects of latency and TB relapse in humans.

Introduction

Infection with *Mycobacterium tuberculosis* (Mtb) most frequently results in latent tuberculosis infection (LTBI), a clinically asymptomatic state in which the only sign of infection is a positive test for immune reactivity to TB antigens, either a tuberculin skin test (TST) or interferon gamma release assay (IGRA). Individuals with LTBI are at risk for developing active TB, and many active TB cases that occur in someone with LTBI, especially in areas of low active TB prevalence, arise from reactivation of the Mtb strain that originally infected the individual. Although most active TB cases may represent relapses of a primary Mtb infection that occurred recently, within 1-5 years (Behr et al., 2018, 2019), even these cases proceed through a paucibacillary, clinically asymptomatic state and therefore fit within the definition of LTBI.

TB chemotherapy often fails to sterilize Mtb infections, resulting in individuals at risk of relapse TB, even in the absence of bacterial drug resistance (Malherbe et al., 2016). In fact, 5% of TB patients that have been treated with standard TB chemotherapy develop recurrent TB (Romanowski et al., 2019; Colangeli et al., 2018; Merle et al., 2014) and recurrent TB may account for up to 10-30 % of all TB cases (Chaisson and Churchyard, 2010). In patients with recurrent TB the Mtb strain that caused the primary infection regrows to lead to a second episode of active TB (relapse), or infection with a new strain causes the second episode (reinfection)(McIvor et al., 2017). The proportion of relapse and reinfection among recurrent TB cases varies with the overall TB incidence rate and HIV status (Cardona, 2016). Notwithstanding, post-treatment LTBI, in which drug susceptible Mtb persists despite chemotherapy and can cause disease relapse, is a major contributor to the spread of TB.

The standard TB mouse model exhibits some features of human TB and has informed about TB immunopathogenesis, the role of host genetics, the efficacy of antimicrobial therapy, and host-pathogen interactions (Flynn, 2006). However, it fails to mimic human LTBI, including persistent infections following TB chemotherapy. Infection of C57BL/6 or BALB/C mice by aerosol with ~100 colony forming units (CFU) of Mtb follows a reproducible disease course consisting of a three to four week-long period during which the bacteria replicate until Mtb antigen-specific T cell responses prevent replication and maintain the infection at a stable bacterial load for many months, until eventual death of the animals (Wolf et al., 2008; Urdahl et al., 2011). During this chronic, persistent infection the typical mycobacterial load is ~10^6 CFU in the lung, consisting of bacilli that likely exist in different subpopulations, which are either killed by the immune system, replicating slowly, or maintain viability in a nonreplicating state (Gill et al., 2009; Wang et al., 2019). The overall bacterial burden remains nearly constant, indicating a balance between the host immune response restricting the infection and Mtb resisting eradication. Although during LTBI in humans, the bacilli are likely in a similar standoff with the immune system, a

healthy human immune system restricts bacterial replication much more vigorously and is therefore capable of driving Mtb into a state of latency in which Mtb is believed to be contained at low numbers within granulomas (Barry et al., 2009). In contrast to the high-titer persistence that occurs during the chronic phase of the conventional mouse model, latent infections in humans are defined by paucibacillary stages, which are characterized by low, often undetectable numbers of Mtb and the corresponding lack of clinical symptoms. Recently, infection of mice with an ultra-low dose of Mtb was shown to cause outcomes that more closely model human TB (Plumlee et al., 2021). Infection with 1-3 CFU resulted in heterogeneous bacterial burdens in lungs, lymph nodes and spleens and mice established pulmonary lesions that shared features with human granulomas. Some ultra-low dose infected mice developed a single granuloma with little or no dissemination to the contralateral lung. Despite the variability in bacterial burden, all infected mice exhibited between 46 and 82,000 CFU in their lungs on day 83 post infection, and by this time point Mtb had disseminated to the spleen in all infected mice and reached a titer of approximately 10,000 CFU. Thus, while ultra-low-dose infection of mice models aspects of human TB, it does not mimic LTBI and relapse TB.

We sought to develop a mouse model that is characterized by low numbers of persisting bacteria and allows studying post-treatment latent infection. That Mtb can establish a latent infection in mice was first demonstrated in the "Cornell Model" developed by McCune and colleagues at Cornell University (McCune and TOMPSETT, 1956; McCune et al., 1956, 1966a; b). In this model, Mtb infected mice are treated with isoniazid (INH) and pyrazinamide (PZA) until no bacteria can be detected by microscopy and culture on agar plates, even when homogenates of whole lungs, spleens, and livers and every organ including bones from several mice were cultured. After cessation of drug treatment, the infection relapsed spontaneously in about 30% of animals, and relapse could be induced in the remaining mice by immunosuppression with corticosteroids. The Cornell model has been used to study various aspects of Mtb persistence and latency (Flynn et al., 1998; Pinxteren et al., 2000; Dhillon and Mitchison, 1994). However, this model is notorious for its varying relapse rates and it did not result in reproducible relapse when we tried to implement it. To overcome its limitations, particularly the lack of reproducibility but also the need for prolonged antibiotic therapy which can induce resistance conferring mutations or drug tolerance (Scanga et al., 1999), we developed models in which latent infection is established via transient genetic inactivation of essential Mtb genes. After Mtb can no longer be detected, target gene expression is restored and the timing and frequency of spontaneous or corticosteroid-induced relapse report the number of mice in which Mtb persisted in the form of a paucibacillary infection. In these mouse models, Mtb persists for months below the limit of detection and reproducibly causes TB relapse in a fraction of infected animals, mimicking aspects of post-treatment latency in humans. Unexpectedly

we discovered that the immunological requirements for control of latent Mtb in these models differ depending on the protein that was transiently depleted to induce latent infection.

Results and Discussion

Chemotherapy of Mtb infected mice did not result in consistent latent infection

In an effort to implement the Cornell model we infected C57BL/6 mice with virulent Mtb, strain H37Rv. Starting four weeks post infection, we treated the infected mice with isoniazid (INH) and pyrazinamide (PZA) for 14 weeks (Fig. S1 A). CFU in lungs, spleens, livers, lymph nodes, bone marrow and kidneys in groups of 5 mice declined with drug treatment and on day 126, when no CFU were detected in 10 mice, chemotherapy was terminated. On day 182 post infection, 8 weeks after the end of chemotherapy, a group of 10 mice was analyzed, and no Mtb was detectable in any of the mouse lungs and other organs. A group of 10 mice received dexamethasone for 4 weeks. Eight weeks later these 10 immunosuppressed mice and 10 mice that did not receive dexamethasone were analyzed for CFU, but all mice had been sterilized by the drug treatment. In a repeat experiment we increased the number of mice in each treatment group. But again, in none of the organs from 15 mice that did not receive dexamethasone was any Mtb CFU detectable, and only one out of 14 mice that had received dexamethasone relapsed with CFU in lung and spleen (Fig. S1 B). In past studies the frequency of relapse also varied substantially and sometimes relapse could not be observed at all, even with immunosuppression (Scanga et al., 1999; Lenaerts et al., 2004; Jeon et al., 2012). Therefore, we sought a different approach to reproducibly establish latent Mtb infection in mice.

Genetic depletion of essential proteins reproducibly results LTBI in mice

We have previously developed genetic switches to control gene expression in Mtb during mouse infection and demonstrated that the depletion of essential proteins kills Mtb (Ehrt et al., 2005; Kim et al., 2013; Puckett et al., 2017; Lin et al., 2016; Tiwari et al., 2018). A dual-control (DUC) switch that combines transcriptional repression with inducible proteolysis can generate more than 99.9% repression, quickly inactivates gene activities in replicating and nonreplicating mycobacteria, and provides increased robustness against phenotypic reversion (Kim et al., 2013). We hypothesized that killing of Mtb following transient depletion of essential proteins does not fully sterilize the infection but may lead to paucibacillary infection states during which low numbers of viable bacteria persist and could reactivate to cause TB relapse. We further expected that the depletion of different targets will have different consequences for establishing a latent state and result in reproducibly distinct relapse rates.

We selected DUC mutants for biotin protein ligase (BPL) and thioredoxin reductase (TrxB2) to test these hypotheses.

BPL covalently attaches biotin to acyl-CoA carboxylases and inactivation of BPL inhibits all lipid biosynthesis (Tiwari et al., 2018). Inactivating BPL is thus expected to cause pleiotropic effects, including profound changes in Mtb's envelope structure, which might make the pathogen exceptionally vulnerable to host immunity and only rarely allow Mtb to establish a persistent, paucibacillary infection. To test this, we infected C57BL/6 mice with a BPL-DUC mutant and after the mutant had established a chronic infection, eradicated it via doxycycline (doxy) mediated BPL depletion (Fig. 1 A). Following apparent sterilization, doxycycline was withdrawn to restore BPL expression and reestablish a wild type paucibacillary infection. At multiple time points post doxy treatment groups of 10 mice were sacrificed and analyzed for Mtb CFU. We could not detect Mtb CFU in lungs or spleens of the majority of mice. In three experiments the frequency of relapse varied from 0% to 10%. In two of these experiments, we treated additional groups of mice with dexamethasone (dexa) for 8 or 4 weeks. Eleven mice out of 35 mice that were treated with dexa relapsed; thus, the frequency of mice with regrowth in lung and/or spleen increased ~3-fold from an average of 10% to 31%. Table 1 summarizes relapse data from multiple experiments with BPL-DUC and reveals that targeting Mtb's BPL results in a low-frequency-relapse latency mouse model, in which 70% of relapse depends on immunosuppression.

Thioredoxin reductase (TrxB2) is essential for thiol redox homeostasis and has been considered as a drug target (Lin et al., 2016; Harbut et al., 2015). Depletion of TrxB2 has pleiotropic effects and kills Mtb *in vitro* and during acute and chronic high titer mouse infections. However, the importance of redox homeostasis for establishing paucibacillary infection was unclear. We analyzed the consequences of targeting TrxB2 using the TrxB2-DUC mutant in six independent experiments (Figure 1B shows data from four experiments). The experimental design was similar to the one used to analyze BPL-DUC. But in contrast to BPL-DUC, which caused approximately 10% relapse without immunosuppression, TrxB2-DUC regrew on average in lungs and/or spleens of 78% of immunocompetent mice after a period of at least 8 weeks during which no CFU could be recovered from lungs or spleens in the majority of mice. In two experiments we included dexa treatment which caused relapse in >90% of the mice (Fig. 1B, table 1). Targeting Mtb's TrxB2 thus results in a high-frequency-relapse (>90% relapse) latency mouse model in which relapse occurs frequently even without immunosuppression.

For both the TrxB2-DUC and the BPL-DUC model, we confirmed that in the bacteria that were recovered following relapse, the targeted genes were still regulated in an anhydrotetracycline (atc)/doxy dependent manner and thus did not represent escape mutants. Both mutants were killed with similar kinetics in mice during doxy treatment and we confirmed that the minimal inhibitory concentration (MIC)

of atc and doxy required to suppress growth in vitro was not significantly different for the two mutants (Fig. S2). Most likely, BPL depletion, which interferes with cell envelope biosynthesis synergizes more effectively with host immunity in eliminating Mtb, than depletion of TrxB2 which perturbs growth-essential pathways, including sulfur metabolism, DNA replication and antioxidant defense systems (Lin et al., 2016).

To investigate if viable Mtb was detectable in organs other than lung and spleen, we analyzed multiple organs and found Mtb in the lymph nodes, liver, kidney, bone marrow and thymus during acute infection (Fig. 1 C and D). These bacteria were eliminated with doxy treatment similarly to those in lungs and spleens and we did not detect Mtb in any of the surveyed organs during latency. We previously demonstrated that depletion of BPL and TrxB2 is accompanied by progressive healing of lesions in the lungs of infected mice (Tiwari et al., 2018; Lin et al., 2016) Previous work also demonstrated that doxy penetrates every mouse tissue and organ to the extent needed to silence Mtb genes controlled by the DUC system (Gengenbacher et al., 2020). Thus, it is unlikely that the bacilli survived in a doxy protected site.

We found that Mtb reactivated most frequently in lungs, followed by spleens and lymph nodes and thymus (Fig. 1 C and D) suggesting that the primary site of paucibacillary persistence in aerosol infected mice is the lung. We cannot distinguish whether TB relapse is caused by very few surviving organisms, that may be below the limit of detection by culture on agar plates or is due to a larger population of bacilli that are in a metabolically altered state that does not support colony formation on agar plates. We tested if Mtb was detectable with a liquid resuscitation assay (Fig. S3 A and B), that resuscitates Mtb from sputa from TB patients, chronically infected mice and *in vitro* stressed cultures (Chengalroyen et al., 2016; Hu et al., 2015; Saito et al., 2017; Imperial et al., 2018). However, this method also failed to identify viable bacilli during latency; and during active infection it yielded similar numbers as obtained by CFU determinations of infected organ homogenates on agar plates. In summary, the data presented here demonstrate that conditional depletion of an essential Mtb protein can induce latency of Mtb in mice, which is defined by a period of time lasting from weeks to months during which Mtb CFU are not detectable by culturing organ homogenates and that is followed by spontaneous or immune suppression induced regrowth of Mtb in a significant proportion of mice.

Control of TrxB2-DUC requires adaptive immunity while BPL-DUC is likely controlled by innate responses

These new models of TB latency provided an opportunity to interrogate the immune factors that control the Mtb latent state. T cells are crucial for the control of chronic Mtb infections (Jasenosky et al., 2015) and we therefore analyzed the pulmonary T cell populations in BPL-DUC and TrxB2-DUC infected mice

at different time points of relapse experiments. As expected, the proportion of CD4 and to some degree CD8 T cells in the lung increased during acute infection and decreased with the decline of bacterial titers (Fig. 2, A-C). This included antigen 85 specific CD4 T cells (Fig. 2 D). There were no significant differences in the proportions of cytokine producing T cells between mice infected with either mutant (Fig. S3 C). We interrogated whether lung infiltrating T cells resembled effector memory T cells (T_{EM}, $CD44^{hi}CD62L^{lo}$) and resident memory T cells (T_{RM}, $CD44^{hi}CD62L^{lo}CD69^{+}$) cells. This revealed that latent infection with TrxB2-DUC was accompanied by an increased fraction of CD4 lung-resident memory T cells and CD4 effector memory T cells compared to mice infected with BPL-DUC at five and seven months post infection (Fig. 2). Memory CD8 T cell populations were not significantly different between mice infected with the two mutants, except that at seven months post infection, the proportion of CD8 lung-resident memory T cells was higher in TrxB2-DUC infected mice compared to BPL-DUC infected mice. The increased percentage of CD4 memory T cells might reflect a larger latent TrxB2-DUC population that eventually causes relapse, than the persisting BPL-DUC population, which the mice are able to control better.

To assess whether adaptive immunity is required to control latent Mtb, we infected SCID mice, which lack functional B and T cells, with both mutants. Because SCID mice are highly susceptible to Mtb, we started doxy treatment at two weeks post infection, kept the mice on doxy for eight weeks, which cleared the infection, and then monitored relapse (Fig. 3). BPL-DUC reactivated only in 1 of 14 mice and in this mouse could be isolated from multiple organs (Fig. 3 A and B). In contrast, TrxB2-DUC did not establish latent infection in SCID mice as it regrew almost immediately after removal of doxy and disseminated to all organs we cultured (Fig. 3 C and D). This identifies the adaptive immune response as essential to establish latent Mtb infection following depletion of TrxB2. Control of BPL-DUC occurred even in the absence of adaptive immunity. The reactivation frequency of BPL-DUC in SCID mice was even lower than in dexamethasone treated mice (Fig. 1 A) suggesting that innate immune responses might be crucial to establish latency after inactivation of BPL. However, the differences in experimental setup, including the earlier start (2 weeks instead of 4 weeks post infection) and shorter period of doxy treatment (66 days instead of 112 days) of the SCID mice complicates the direct comparison of these experiments. Future work with identical experimental parameters is required to substantiate these findings and their interpretation.

BCG vaccination does not prevent TB relapse

Vaccination with *Mycobacterium bovis* Bacille Calmette–Guérin (BCG) protects infants from extrapulmonary manifestations of Mtb infection and against progression from infection to active disease (Roy et al., 2014; Trunz et al., 2006). In contrast, its effectiveness in preventing pulmonary TB disease

in adults is highly variable (Mangtani et al., 2014). We sought to determine if BCG vaccination reduces TB relapse in our genetic latency model. To assess this, we vaccinated mice that were latently infected with TrxB2-DUC with BCG Pasteur. We vaccinated via the intravenous (iv) route because in non-human primates high-dose iv BCG vaccination was recently demonstrated to prevent or substantially limit Mtb infection (Darrah et al., 2020). We analyzed immune responses in BCG vaccinated mice and phosphate-buffered saline (PBS) treated controls 25 days post vaccination and determined reactivation of Mtb in lungs and spleens of both groups on day 280 post infection, 20 weeks post vaccination (Fig. 4). BCG vaccination did not affect reactivation of latent Mtb in the lungs of mice; almost all mice in the vaccinated and the control group had CFU in their lungs (Fig. 4 A). In spleens however, Mtb regrew in only 5 of 21 vaccinated mice while 17 of 21 PBS-treated control mice yielded CFU from their spleens (Fig. 4 B). Thus, while systemic BCG vaccination did not prevent TB relapse it reduced Mtb regrowth in spleens. Vaccination might reduce dissemination of reactivating Mtb, if reactivation of Mtb primarily starts in the lung as our data suggest (Fig. 1 C and D). BCG vaccination induced more cytokine expressing CD4 T cells in the spleen than in the lung, in particular the number of IL-2 producing CD4 T cells only increased in the spleen of vaccinated mice (Fig. 4 C and D). This might be due to the higher BCG titers in spleens compared to lungs of the vaccinated mice (Fig. 4 E). BCG titers declined in the lungs approximately 10-fold from day 1 to day 24 and then remained stable until day 40 post immunization. In the spleen BCG CFU did not significantly decline during that period and remained approximately 1.5 log10 higher that in the lung. The proportion of effector memory CD4 and CD8 T cells increased in both lungs and spleens in response to BCG vaccination (Fig. 4 F and G). In summary, these data demonstrate that, like in humans, BCG-induced immune responses are not sufficient to prevent reactivation of latent Mtb infection. The TrxcB2-DUC latency model could be utilized to test vaccine candidates or strategies for their efficacy in preventing TB relapse.

To address the need for mouse models of LTBI and relapse TB, we developed reproducible models of paucibacillary (low levels of bacilli) infection in mice. We demonstrated that Mtb persists for months below the limit of detection and causes relapse TB in a proportion of infected animals. Although we do not claim that these mouse models represent the equivalent of human latency, we do believe that they can inform our understanding, via comparison, of analogous human TB disease states such as reactivation from latency and relapse after TB treatment. These are both states of paucibacillary infection in which bacterial and host immune factors interact to determine whether Mtb reappears or remains clinically silent. The mouse latency models described here allow to study these aspects of TB biology that are not accessible in the conventional TB mouse model. They provide opportunity to define the host cells and molecules that prevent transition from LTBI to acute TB. They could help prioritize

drug candidates that eradicate latent populations and thus could shorten chemotherapy and they could facilitate the evaluation of TB vaccine candidates.

Materials and Methods

Bacterial strains and culturing conditions

Mycobacterium tuberculosis H37Rv was used as virulent wild type strain. BPL-DUC and TrxB2-DUC have been previously described (Tiwari et al., 2018; Lin et al., 2016). BCG Pasteur was from the American Type Culture Collection strain TMC 1011 (ATCC #35734). Strains were cultured in liquid Middlebrook 7H9 medium supplemented with 0.2% glycerol, 0.05% tween80 and ADN (0.5% bovine serum albumin, 0.2% dextrose, 0.085% NaCl) and on Middlebrook 7H10 agar plates supplemented with 0.2% glycerol and Middlebrook OADC enrichment (Becton Dickinson). Antibiotics were added for selection of genetically modified strains at the following concentrations: hygromycin (50 µg/ml), kanamycin (25 µg/ml), zeocin (25 µg/ml).

Mouse infections

Female 8- to 10-week-old C57BL/6 (# 000664, Jackson Laboratory) or SCID mice (# 001913, Jackson Laboratory) were infected with the indicated Mtb strains using an inhalation exposure system (Glas-Col) with a mid-log phase Mtb culture to deliver approximately 200 bacilli per mouse. To vaccinate, mice were injected in the lateral tail vein with 100 µl BCG suspension in PBS at a concentration of 10^7/ml. Mice received doxycycline containing mouse chow (2,000 ppm; Research Diets) for the indicated periods. For the Cornell model, mice received INH (125 µg/ml) and PZA (15 g/L) in the drinking water for 14 weeks. To enumerate CFU, organs were homogenized in PBS and cultured on 7H10 agar. Charcoal (0.4 %, w/v) was added to the plates that were used to culture homogenates from doxy and antibiotic treated mice. Agar plates were incubated for 3-4 weeks at 37°C. We identified suppressor mutants by culturing half of each organ homogenate on 7H10 plates containing atc (500 ng/ml). When suppressor mutants grew up, which was very rare, the infected mice were excluded from the experiment. Mice were housed in a BSL3 vivarium. All mouse experiments were approved by and performed in accordance with requirements of the Weill Cornell Medicine Institutional Animal Care and Use Committee.

Limited dilution assay and MPN calculations

Half of each lung homogenate (3 right lung lobes resuspended in 2 mL PBS supplemented with 2% (v/v) BD MGIT PANTA antibiotic mixture, BD Cat#245114) was used to set up the limited dilution series as described previously (Saito et al., 2017). Mtb H37Rv culture filtrate was prepared and MPN calculations were performed as described (Saito et al., 2017).

Flow cytometry

Mouse lungs were isolated and placed in RPMI1640 containing Liberase Blendzyme 3 (70 µg/ml; Roche) and DNase I (50 µg/ml; Sigma-Aldrich). Lungs were then cut into small pieces and incubated at 37°C for 1 hour. The cells were filtered using cell strainers, collected by centrifugation, resuspended in ACK hemolysis buffer (ThermoFisher) and incubated for 10 minutes at room temperature. Cell were then washed with PBS and resuspended in splenocyte medium (RPMI-1640, supplemented with 10% FBS, 2 mM GlutaMax 10 mM HEPES, and 50 µM M2-mercapto-ethanol). For intracellular staining samples, cells were stimulated with PPD (20 µg/ml) in the presence of anti-CD28 antibody (37.51, BioLegend) for 1.5 hours and 10 µg/ml Brefeldin A (Sigma) and monesin were added and incubated at 37 degree for another 3 hrs. Samples were kept on ice in a refrigerator overnight. Cells were stained with Zombie-UV (BioLegend) to discriminate live and dead cells. Purified anti-CD16/32 antibody (93, BioLegend) was used to block Fc receptor before staining. PerCp-Cy5.5 anti-CD62L (MEL-14, Thermofisher), Alexa 700 anti-CD45 (30-F11, BioLegend), BV605 anti-CD4 (RM4-5, BioLegend), BV711 anti-CD8 (53-6.7, BioLegend), APC Cy7 anti-CD69 (H1.2F3, BioLegend), BUV395 anti-CD44 (IM7, BD Biosciences) were used to stain cells for 30 minutes at room temperature. Then the cells were fixed in fixation buffer (BioLegend) for 30 minutes and taken out of BSL3. The cells were incubated for 20 minutes in permeabilization buffer (eBioscience) before intracellular cytokine staining. FITC anti-TNF (MP6-XT22, BioLegend), BUV737 anti-IFNγ(XMG1.2, BD Biosciences), APC anti-IL-2 (JES6-5H4, BioLegend), BV785 anti-CD3 (17A2, BioLegend) were used to stain cells for 30 minutes. Cells were washed and resuspend in cell staining buffer (BioLegend). Flow cytometry data were acquired on a cytometer (LSR Fortessa TM; BD Biosciences) and were analyzed with FlowJoTM V10. Gating strategies are depicted in Fig. S3.

Statistical Analysis

Flow cytometry data were analyzed by one-way ANOVA with Turkey's multiple comparison test or by unpaired, two tailed t-test and differences in cell frequencies between TrxB2-DUC and BPL-DUC infected animals are reported using Prism 9.0.1 (GraphPad Software).

Acknowledgements

We thank J. Andres, N. Betancourt, I. DaSilva, J. Kim, J. McConnell, R. Aguilera Olvera, N. Song, P. Pino Tamayo for technical assistance. We thank K. Saito, T. Warrier, K. Burns, C. Nathan, M. Glickman, K. Rhee and V. Dartois for advice and helpful discussions; M. Glickman, C. Nathan and K. Rhee for critically reviewing this manuscript and constructive suggestions. This work was supported by the Tri-Institutional TB Research Unit (NIH Grant U19 AI111143).

Author contributions

H. Su, K. Lin, D. Tiwari, C. Healy, C. Trujillo, Y. Liu, D. Schnappinger, and S. Ehrt conceived the study and analyzed the data; H. Su, K. Lin, D. Tiwari, C. Healy, C. Trujillo, Y. Liu performed the experiments; S. Ehrt wrote the manuscript with input from the other authors.

The authors declare no competing financial interests.

Table 1. Data obtained from 4 experiments with BPL-DUC and 6 experiments with TrxB2-DUC, which included a dexamethasone treatment group in 2 experiments with each mutant. Reported are the average group sizes, average number of mice with regrowth in lung or spleen or both organs, the average number of mice in which Mtb regrew (relapsed mice) and the average % relapse and standard deviation (SD).

		Spontaneous regrowth						Dexamethasone induced regrowth					
		Group size	Lung only	Spleen only	Lung & spleen	Relapsed	% Relapse	Group size	Lung only	Spleen only	Lung & spleen	Relapsed	% Relapse
BPL-DUC	Mean	26.0	0.5	1.0	1.0	2.5	**10.1**	17.5	4.0	0	1.5	5.5	**31.4**
	SD	3.9	1.0	1.1	0.8	2.1	8.2	2.1	0	0	0.7	0.7	0.23
TrxB2-DUC	Mean	20.8	4.7	0	12	16.7	**78.1**	18	5.5	2.5	9.5	17.5	**96.9**
	SD	3.2	4.2	0	4.4	6.7	22.9	2.8	5.5	0.7	6.4	3.5	4.4

References

Barry, C.E., H.I. Boshoff, V. Dartois, T. Dick, S. Ehrt, J. Flynn, D. Schnappinger, R.J. Wilkinson, and D. Young. 2009. The spectrum of latent tuberculosis: rethinking the biology and intervention strategies. *Nat. Rev. Microbiol*. 7:845–855. doi:10.1038/nrmicro2236.

Behr, M.A., P.H. Edelstein, and L. Ramakrishnan. 2018. Revisiting the timetable of tuberculosis. *BMJ (Clinical research ed.)*. 362:k2738. doi:10.1136/bmj.k2738.

Behr, M.A., P.H. Edelstein, and L. Ramakrishnan. 2019. Is *Mycobacterium tuberculosis* infection life long? *BMJ*. 367:l5770. doi:10.1136/bmj.l5770.

Cardona, P.-J. 2016. Reactivation or reinfection in adult tuberculosis: Is that the question? *Int. J. Mycobacteriol*. 5:400–407. doi:10.1016/j.ijmyco.2016.09.017.

Chaisson, R.E., and G.J. Churchyard. 2010. Recurrent Tuberculosis: Relapse, Reinfection, and HIV. *J. Infect. Dis*. 201:653–655. doi:10.1086/650531.

Chengalroyen, M.D., G.M. Beukes, B.G. Gordhan, E.M. Streicher, G. Churchyard, R. Hafner, R. Warren, K. Otwombe, N. Martinson, and B.D. Kana. 2016. Detection and Quantification of Differentially Culturable Tubercle Bacteria in Sputum from Patients with Tuberculosis. *Am. J. Respir. Crit. Care Med*. 194:1532–1540. doi:10.1164/rccm.201604-0769oc.

Colangeli, R., H. Jedrey, S. Kim, R. Connell, S. Ma, U.D.C. Venkata, S. Chakravorty, A. Gupta, E.E. Sizemore, L. Diem, D.R. Sherman, A. Okwera, R. Dietze, W.H. Boom, J.L. Johnson, W.R.M. Kenzie, and D. Alland. 2018. Bacterial Factors That Predict Relapse after Tuberculosis Therapy. *N. Engl. J. Med*. 379:823–833. doi:10.1056/nejmoa1715849.

Darrah, P.A., J.J. Zeppa, P. Maiello, J.A. Hackney, M.H. Wadsworth, T.K. Hughes, S. Pokkali, P.A. Swanson, N.L. Grant, M.A. Rodgers, M. Kamath, C.M. Causgrove, D.J. Laddy, A. Bonavia, D. Casimiro, P.L. Lin, E. Klein, A.G. White, C.A. Scanga, A.K. Shalek, M. Roederer, J.L. Flynn, and R.A. Seder. 2020. Prevention of tuberculosis in macaques after intravenous BCG immunization. *Nature*. 577:95–102. doi:10.1038/s41586-019-1817-8.

Dhillon, J., and D.A. Mitchison. 1994. Effect of vaccines in a murine model of dormant tuberculosis. *Tubercle Lung Dis*. 75:61–64. doi:10.1016/0962-8479(94)90105-8.

Ehrt, S., X.V. Guo, C.M. Hickey, M. Ryou, M. Monteleone, L.W. Riley, and D. Schnappinger. 2005. Controlling gene expression in mycobacteria with anhydrotetracycline and Tet repressor. *Nucleic Acids Res*. 33:e21. doi:10.1093/nar/gni013.

Flynn, J.L. 2006. Lessons from experimental *Mycobacterium tuberculosis* infections. *Microbes Infect*. 8:1179–1188. doi:10.1016/j.micinf.2005.10.033.

Flynn, J.L., C.A. Scanga, K.E. Tanaka, and J. Chan. 1998. Effects of aminoguanidine on latent murine tuberculosis. *J. Immunol*. 160:1796–803.

Gengenbacher, M., M.D. Zimmerman, J.P. Sarathy, F. Kaya, H. Wang, M. Mina, C. Carter, M.A. Hossen, H. SU, C. Trujillo, S. Ehrt, D. Schnappinger, and V. Dartois. 2020. Tissue Distribution of Doxycycline in Animal Models of Tuberculosis. *Antimicrob. Agents Chemother*. 64:5547. doi:10.1128/aac.02479-19.

Gill, W.P., N.S. Harik, M.R. Whiddon, R.P. Liao, J.E. Mittler, and D.R. Sherman. 2009. A replication clock for *Mycobacterium tuberculosis*. *Nat. Med*. 15:211–214. doi:10.1038/nm.1915.

Harbut, M.B., C. Vilchèze, X. Luo, M.E. Hensler, H. Guo, B. Yang, A.K. Chatterjee, V. Nizet, P.G. Schultz, and F. Wang. 2015. Auranofin exerts broad-spectrum bactericidal activities by targeting thiol-redox homeostasis. *Proc. Natl. Acad. Sci*. *USA*. 112:4453–4458. doi:10.1073/pnas.1504022112.

Hu, Y., A. Liu, F. Ortega-Muro, L. Alameda-Martin, D. Mitchison, and A. Coates. 2015. High-dose rifampicin kills persisters, shortens treatment duration, and reduces relapse rate in vitro and in vivo. *Front. Microbiol*. 6:641. doi:10.3389/fmicb.2015.00641.

Imperial, M.Z., P. Nahid, P.P.J. Phillips, G.R. Davies, K. Fielding, D. Hanna, D. Hermann, R.S. Wallis, J.L. Johnson, C. Lienhardt, and R.M. Savic. 2018. A patient-level pooled analysis of treatment-shortening regimens for drug-susceptible pulmonary tuberculosis. *Nat. Med*. 24:1708–1715. doi:10.1038/s41591-018-0224-2.

Jasenosky, L.D., T.J. Scriba, W.A. Hanekom, and A.E. Goldfeld. 2015. T cells and adaptive immunity to *Mycobacterium tuberculosis* in humans. *Immunol. Rev*. 264:74–87. doi:10.1111/imr.12274.

Jeon, B.Y., J. Kwak, M.Y. Hahn, S.Y. Eum, J. Yang, S.C. Kim, and S.N. Cho. 2012. In vivo characteristics of Korean Beijing *Mycobacterium tuberculosis* strain K1 in an aerosol challenge model and in the Cornell latent tuberculosis model. *J. Med. Microbiol*. 61:1373–1379. doi:10.1099/jmm.0.047027-0.

Kim, J.-H., K.M. O'Brien, R. Sharma, H.I.M. Boshoff, G. Rehren, S. Chakraborty, J.B. Wallach, M. Monteleone, D.J. Wilson, C.C. Aldrich, C.E. Barry, K.Y. Rhee, S. Ehrt, and D. Schnappinger. 2013. A genetic strategy to identify targets for the development of drugs that prevent bacterial persistence. *Proc. Natl. Acad. Sci. USA*. 110:19095–19100. doi:10.1073/pnas.1315860110.

Lenaerts, A.J., P.L. Chapman, and I.M. Orme. 2004. Statistical limitations to the Cornell model of latent tuberculosis infection for the study of relapse rates. *Tuberculosis*. 84:361–364. doi:10.1016/j.tube.2004.03.002.

Lin, K., K.M. O'Brien, C. Trujillo, R. Wang, J.B. Wallach, D. Schnappinger, and S. Ehrt. 2016. *Mycobacterium tuberculosis* Thioredoxin Reductase Is Essential for Thiol Redox Homeostasis but Plays a Minor Role in Antioxidant Defense. *PLoS Pathog*. 12:e1005675-20. doi:10.1371/journal.ppat.1005675.

Malherbe, S.T., S. Shenai, K. Ronacher, A.G. Loxton, G. Dolganov, M. Kriel, T. Van, R.Y. Chen, J. Warwick, T. Song, M. Lee, G. Schoolnik, G. Tromp, D. Alland, J. Winter, G. Walzl, C.T. Consortium, L. Lucas, G. van der Spuy, K. Stanley, L. Thiart, B. Smith, N.D. Plessis, C.G.G. Beltran, E. Maasdorp, A. Ellmann, H. Choi, J. Joh, L.E. Dodd, B. Allwood, C. Koegelenberg, M. Vorster, and S. Griffith-Richards. 2016. Persisting positron emission tomography lesion activity and *Mycobacterium tuberculosis* mRNA after tuberculosis cure. *Nat. Med*. 22:1094–1100. doi:10.1038/nm.4177.

Mangtani, P., I. Abubakar, C. Ariti, R. Beynon, L. Pimpin, P.E.M. Fine, L.C. Rodrigues, P.G. Smith, M. Lipman, P.F. Whiting, and J.A. Sterne. 2014. Protection by BCG Vaccine Against Tuberculosis: A Systematic Review of Randomized Controlled Trials. *Clin. Infect. Dis*. 58:470–480. doi:10.1093/cid/cit790.

McCune, R.M., F.M. Feldmann, H.P. Lambert, and W. McDermott. 1966a. Microbial persistence. I. The capacity of tubercle bacilli to survive sterilization in mouse tissues. *J. Exp. Med*. 123:445–468.

McCune, R.M., F.M. Feldmann, and W. McDermott. 1966b. Microbial persistence. II. Characteristics of the sterile state of tubercle bacilli. *J. Exp. Med*. 123:469–486.

McCune, R.M., W. McDermott, and R. TOMPSETT. 1956. The fate of *Mycobacterium tuberculosis* in mouse tissues as determined by the microbial enumeration technique. II. The conversion of tuberculous infection to the latent state by the administration of pyrazinamide and a companion drug. *J. Exp. Med*. 104:763–802.

McCune, R.M., and R. TOMPSETT. 1956. Fate of *Mycobacterium tuberculosis* in mouse tissues as determined by the microbial enumeration technique. I. The persistence of drug-susceptible tubercle bacilli in the tissues despite prolonged antimicrobial therapy. *J. Exp. Med*. 104:737–762.

McIvor, A., H. Koornhof, and B.D. Kana. 2017. Relapse, re-infection and mixed infections in tuberculosis disease. *Pathog. Dis*. 75:ftx020-. doi:10.1093/femspd/ftx020.

Merle, C.S., K. Fielding, O.B. Sow, M. Gninafon, M.B. Lo, T. Mthiyane, J. Odhiambo, E. Amukoye, B. Bah, F. Kassa, A. N'Diaye, R. Rustomjee, B.C. de Jong, J. Horton, C. Perronne, C. Sismanidis, O. Lapujade, P.L. Olliaro, and C. Lienhardt. 2014. A Four-Month Gatifloxacin-Containing Regimen for Treating Tuberculosis. *N. Engl. J. Med*. 371:1588–1598. doi:10.1056/nejmoa1315817.

Pinxteren, L.A. van, J.P. Cassidy, B.H. Smedegaard, E.M. Agger, and P. Andersen. 2000. Control of latent *Mycobacterium tuberculosis* infection is dependent on CD8 T cells. *Eur. J. Immunol*. 30:3689–3698. doi:10.1002/1521-4141(200012)30:12<3689::aid-immu3689>3.0.co;2-4.

Plumlee, C.R., F.J. Duffy, B.H. Gern, J.L. Delahaye, S.B. Cohen, C.R. Stoltzfus, T.R. Rustad, S.G. Hansen, M.K. Axthelm, L.J. Picker, J.D. Aitchison, D.R. Sherman, V.V. Ganusov, M.Y. Gerner, D.E. Zak, and K.B. Urdahl. 2021. Ultra-low Dose Aerosol Infection of Mice with *Mycobacterium tuberculosis* More Closely Models Human Tuberculosis. *Cell Host Microbe*. 29:68-82.e5. doi:10.1016/j.chom.2020.10.003.

Puckett, S., C. Trujillo, Z. Wang, H. Eoh, T.R. Ioerger, I. Krieger, J. Sacchettini, D. Schnappinger, K.Y. Rhee, and S. Ehrt. 2017. Glyoxylate detoxification is an essential function of malate synthase required for carbon assimilation in *Mycobacterium tuberculosis*. *Proc. Natl. Acad. Sci. USA*. 114:E2225–E2232. doi:10.1073/pnas.1617655114.

Romanowski, K., R.F. Balshaw, A. Benedetti, J.R. Campbell, D. Menzies, F.A. Khan, and J.C. Johnston. 2019. Predicting tuberculosis relapse in patients treated with the standard 6-month regimen: an individual patient data meta-analysis. *Thorax*. 74:291–297. doi:10.1136/thoraxjnl-2017-211120.

Roy, A., M. Eisenhut, R.J. Harris, L.C. Rodrigues, S. Sridhar, S. Habermann, L. Snell, P. Mangtani, I. Adetifa, A. Lalvani, and I. Abubakar. 2014. Effect of BCG vaccination against *Mycobacterium tuberculosis* infection in children: systematic review and meta-analysis. *Bmj. Br. Medical J*. 349:g4643. doi:10.1136/bmj.g4643.

Saito, K., T. Warrier, S. Somersan-Karakaya, L. Kaminski, J. Mi, X. Jiang, S. Park, K. shigyo, B. Gold, J. Roberts, E. Weber, W.R. Jacobs, and C.F. Nathan. 2017. Rifamycin action on RNA polymerase in antibiotic-tolerant *Mycobacterium tuberculosis* results in differentially detectable populations. *Proc. Natl. Acad. Sci. USA*. 114:E4832–E4840. doi:10.1073/pnas.1705385114.

Scanga, C.A., V.P. Mohan, H. Joseph, K. Yu, J. Chan, and J.L. Flynn. 1999. Reactivation of Latent Tuberculosis: Variations on the Cornell Murine Model. *Infect. Immun*. 67:4531–4538. doi:10.1128/iai.67.9.4531-4538.1999.

Tiwari, D., S.W. Park, M.M. Essawy, S. Dawadi, A. Mason, M. Nandakumar, M. Zimmerman, M. Mina, H.P. Ho, C.A. Engelhart, T. Ioerger, J.C. Sacchettini, K. Rhee, S. Ehrt, C.C. Aldrich, V. Dartois, and D. Schnappinger. 2018. Targeting protein biotinylation enhances tuberculosis chemotherapy. *Sci. Trans. Med*. 10:eaal1803. doi:10.1126/scitranslmed.aal1803.

Trunz, B.B., P. Fine, and C. Dye. 2006. Effect of BCG vaccination on childhood tuberculous meningitis and miliary tuberculosis worldwide: a meta-analysis and assessment of cost-effectiveness. *Lancet*. 367:1173–1180. doi:10.1016/s0140-6736(06)68507-3.

Urdahl, K.B., S. Shafiani, and J.D. Ernst. 2011. Initiation and regulation of T-cell responses in tuberculosis. *Mucosal Immunol*. 4:288–293. doi:10.1038/mi.2011.10.

Wang, R., K. Kreutzfeldt, H. Botella, J. Vaubourgeix, D. Schnappinger, and S. Ehrt. 2019. Persistent *Mycobacterium tuberculosis* infection in mice requires PerM for successful cell division. *eLife*. 8:100. doi:10.7554/elife.49570.

Wolf, A.J., L. Desvignes, B. Linas, N. Banaiee, T. Tamura, K. Takatsu, and J.D. Ernst. 2008. Initiation of the adaptive immune response to *Mycobacterium tuberculosis* depends on antigen production in the local lymph node, not the lungs. *J. Exp. Med*. 205:105–115. doi:10.1084/jem.20071367.

Fig. 1. Genetic depletion of essential proteins results in reproducible paucibacillary infection. Colony forming units (CFU) in lungs and spleens of mice infected with (**A**) BPL-DUC and (**B**) TrxB2-DUC from 3 and 4 independent experiments. Mice were given doxy-containing chow (yellow area) until the infection had cleared (no detectable CFU). After that, mice were kept on regular chow. In two experiments, groups of 16-26 mice received dexamethasone for either the last 8 weeks or for 4 weeks starting at 2 months before the end of the experiment (pink area). Stippled lines indicate the limit of detection. (**C**) Analysis of TrxB2-DUC in multiple organs. CFU data in organs from 5 mice (months 1, 2, 3.25, 4,5) and 10 mice (month 7) and 19 mice (month 10). (**D**) Analysis of BPL-DUC in multiple organs. CFU data in organs from 4 mice (months 1, 1.2, 2, 4) and 10 mice (months 5.8, 8.8) and 38 mice (month 12), half of which received dexamethasone for 1 month starting at month 9 post infection. Data points from animals that received dexamethasone are marked by a dot within the symbol. The yellow area indicates time of doxy treatment. Stippled lines indicate the limit of detection.

Fig. 2. Paucibacillary infection correlates with an increased frequency of memory T cells. (**A**) CFU from lungs of mice infected with TrxB2-DUC and BPL-DUC. Data correspond to experiment 3 shown in Figure 1. The two TrxB2-infected mice that had relapsed at month 7 and 8 post infection were excluded from the analysis of T cell responses shown in panels B-H. (**B, C**) Frequency of CD4⁺ and CD8⁺ T cells in at the indicated time points. (**D**) Proportion of Ag85 specific CD4⁺ T cells. (**E, F**) CD4⁺ lung-resident memory T cells and effector memory T cells in mice infected with TrxB2-DUC and BPL-DUC. (**G, H**) CD8⁺ lung-resident memory T cells and effector memory T cells. Mice received doxycycline from month 1 to 4.5 as indicated in A. Data are from groups of 3 - 7 mice and differences in cell frequencies between TrxB2-DUC and BPL-DUC infected animals were analyzed by one-way ANOVA with Turkey's multiple comparison test * $P < 0.05$, ** $P < 0.01$, *** $P < 0.005$.

Fig. 3. TrxB2-DUC does not establish paucibacillary infection in SCID mice. (**A**) CFU in lungs of SCID mice infected with BPL-DUC The proportion of mice with CFU is indicated. (**B**) CFU in indicated organs of SCID mice infected with BPL-DUC. One mouse relapsed with CFU in lungs and other organs on day 126. CFU in spleen and liver could not be quantified due to contamination. (**C**) CFU in lungs of SCID mice infected with TrxB2-DUC The proportion of mice with CFU is indicated (**D**) CFU in indicated organs of SCID mice infected with BPL-DUC. Mice received doxy containing food (yellow area) for 8 weeks. Stippled lines indicate the limit of detection.

Fig. 4. BCG vaccination fails to prevent TB relapse. (A) CFU from lungs and (**B**) spleens of mice infected with TrxB2-DUC and vaccinated with BCG or injected with PBS on day 140 post infection (arrow). Groups of 21 mice were analyzed on day 280 post infection, the proportion of mice with CFU is indicated in brackets. Twenty-five days post vaccination immune responses in lungs (**C,F**) and in spleens (**D,G**) were analyzed. Cytokine expressing CD4 T cells were measured by intracellular cytokine staining in CD4$^+$, CD44$^+$ T lymphocytes following stimulation of lung leukocytes (**C**) and splenocytes (**D**) with purified protein derivative (PPD). **(E)** Quantification of BCG CFU in lungs and spleens. On day 1, 25 and 40 post vaccination CFU in lungs and spleens were determined by culturing organ homogenates on 7H10 agar plates. (**F,G**) Effector memory CD4 and CD8 T cells were quantified by flow cytometry in lungs (**F**) and spleens (**G**) in groups of 3-4 PBS treated and 5 BCG vaccinated mice. The differences between PBS treated and BCG vaccinated mice were analyzed by unpaired, two tailed t-test. * *P*< 0.05; ** *P*< 0.01; *** *P*<0.001.