Intermolecular interactions drive protein adaptive and co-adaptive evolution at both species and

population levels

Junhui Peng, Li Zhao

Laboratory of Evolutionary Genetics and Genomics, The Rockefeller University, New York, NY 10065,

USA

*Correspondence to: lzhao@rockefeller.edu

Abstract

Proteins are the building blocks for almost all the functions in cells. Understanding the molecular

evolution of proteins and the forces that shape protein evolution is an essential step in understanding the

basis of function and evolution. Previous studies have shown that adaptation occurs frequently at the

protein surface, such as in genes involved in host-pathogen interactions. However, it remains unclear

whether adaptive sites are distributed randomly or at regions that are associated with particular structural

or functional characteristics across the genome, since many of the proteins lack structural or functional

annotations. Here, we seek to tackle this question by combining large-scale bioinformatic prediction,

structural analysis, phylogenetic inference, and population genomic analysis of *Drosophila* protein-coding

genes. Although adaptation is more relevant to function-related rather than structure-related properties,

we observed that physical interactions may play a role in the co-adaptation of fast-adaptive proteins.

Importantly, protein-protein and protein-DNA interaction sites are hotspots for protein adaptive evolution,

regardless of the levels of intrinsic structural disorder or relative solvent accessibility. We found that

strongly differentiated amino acids across geographic regions in protein coding genes are mostly adaptive,

which may contribute to the long-term adaptive evolution. This strongly indicates that a number of

adaptive sites are repeatedly mutated and selected in evolution, in the past, present, and maybe future. Our

results suggest important roles of intermolecular interactions and co-adaptation in the adaptive evolution

of proteins both at the species and population levels.

Introduction

Natural selection plays an important role in molecular evolution of protein sequences. Recent advances in

genome sequencing and reliable inference methods at both phylogenetic and population levels have

enabled fast and robust estimation of evolutionary rates and adaptation driven by natural selection. In

addition, the increased availabilities of structural and functional data of proteins have made it possible to

study how structural and functional constraints affect protein sequence evolution and adaptation. It is now

well established that different proteins and different sites within a protein have varying rates of evolution

and adaptation due to both structural and functional constraints (Echave et al., 2016; Kosiol et al., 2008;

Lindblad-Toh et al., 2011; Zhang and Yang, 2015). For example, genes that are highly expressed or

perform essential functions are under strong purifying selection and tend to evolve slowly (Drummond et

al., 2005; Moutinho et al., 2019; Pál et al., 2001; Zhang and He, 2005; Zhang and Yang, 2015); genes

involved in host-pathogen interactions, e.g., immune responses and antivirus responses, show

exceptionally high rates of adaptive changes (Enard et al., 2016; Nielsen et al., 2005; Obbard et al., 2009;

Palmer et al., 2018; Sackton et al., 2007; Sironi et al., 2015; Uricchio et al., 2019); and residues that are

intrinsically disordered or at the protein surface are fast evolving and has been proved to be hotspots of

adaptive evolution (Afanasyeva et al., 2018; Goldman et al., 1998; Lin et al., 2007; Moutinho et al., 2019;

Ramsey et al., 2011). More recently, Slodkowicz & Goldman (Slodkowicz and Goldman, 2020)

employed genomic-scale integrated structural and phylogenetic evolutionary analysis in mammals and

showed that positively selected residues are clustered near ligand binding sites, especially in proteins that

are associated with immune responses and xenobiotic metabolism. However, vast majority of the work

focused on differences at the species level, it is unclear how much of the polymorphic changes within a

species may contribute to long-term evolution.

Although evidence have shown that adaptation is more likely to occur at intrinsically disordered

regions and clustered at the surface of proteins, the functional properties of adaptation in the genomic and

population scale remains unclear. Moreover, due to lack of structural and functional information of many

proteins in the genome, the underlying mechanism derived from current studies might be incomplete.

Here, we systematically investigated the evolution and adaptation of protein-coding genes in *Drosophila*

melanogaster by comparing it to its closely related species and their own populations, in order to

distinguish the main factors that impact the evolution and adaption at the protein-coding level. We applied

large-scale bioinformatic and structural analysis to obtain structural and functional properties of proteins.

We then classified residues into different structural and functional sites. By comparing rates of sequence

evolution and adaptation between different proteins and different sites, we were able to locate hotspots of

adaptation at the genome scale. Although adaptation is more sensitive to functional properties rather than

structural properties, we found that putative binding regions including allosteric sites at protein surface

show higher rates of adaptation than other sites. For proteins that are under fast-adaptive evolution, we

showed that they tend to interact with each other more frequently than random expectations and are often

associated with reproduction, immunity, and environmental information processing in *D. melanogaster*.

In addition, we showed that interacting proteins in *D. melanogaster* might undergo co-adaptive evolution.

Furthermore, we hypothesize that molecular interactions or physical interactions might be an important

mechanism that contribute to the adaptive and co-adaptive evolution in *D. melanogaster* genome. At last,

we showed that many non-synonymous SNPs contributing to short-term adaptation are overlapped with

SNPs contributing to long-term adaptive evolution, suggesting that a subset of SNPs on the genomes are

constantly utilized for adaptive purpose.

Results

Putative molecular interaction sites are hotspots for protein adaptive evolution

To uncover the main factors that impact the evolutionary rates of genes, we analyzed 13,528 protein-

coding genes in *D. melanogaster* using genome data from *melanogaster* subgroup species and *D.*

melanogaster population genomics data from 205 inbred lines from Drosophila Genetic Reference Panel,

Freeze 2.0, DGRP2 (Huang et al., 2014). We applied a maximum likelihood method (Yang, 2007) to

compute dN/dS ratio (ω) using the protein-coding sequences of five closely related melanogaster

subgroup species (*D. melanogaster*, *D. simulans*, *D. sechellia*, *D. yakuba* and *D. erecta*). We estimated

the proportions of adaptive changes (α) in each gene by applying an extension of MK test named

asymptotic MK (Messer and Petrov, 2013; Uricchio et al., 2019) using *D. simulans* as outgroup. We then

calculated the rate of adaptive changes (ω_a) of each gene by multiplying ω to α ($\omega_a = \alpha\omega$) (Moutinho et

al., 2019) using *D. yakuba* as the outgroup species (See methods). The rate of nonadaptive changes can be

further calculated by $\omega_{na}=\omega-\omega_a$. Finally, we successfully assigned ω to 12,118 protein coding genes and

ω_a and ω_{na} to 7,192 genes. For each of *D. melanogaster* genes subjecting the same pipeline of analysis,

we further obtained 17 different structural or functional properties (see Methods and supplementary file

S1). We calculated Pearson's correlations of ω, ω_a and ω_{na} with all these properties (Table S1). We

showed that many of these genome-wide correlations were expected according to previous studies

(Supplement Information, section *Impact of gene properties on evolution of protein-coding genes in D.*

melanogaster, Table S1). Interestingly, among these properties, we found that some previously not

reported properties, fractions of molecular-interaction sites (PPI-site ratio, ratio of residues involved in

protein-protein interactions, and DNA-site ratio, ratio of residues involved in protein-DNA interactions)

strongly positively correlated with ω, ω_a and ω_{na} (Supplement Information, section *Molecular*

interactions contribute to the variations of protein sequence evolution and adaptation, Table S1, Figure

S1). The results indicate that molecular interactions might act as an important factor that drive protein

adaptive evolution in *Drosophila* genome.

We then investigate whether residues involved in molecular interactions are targets for adaptive

evolution. To tackle this question, we predicted protein-protein interaction sites (PPI-sites) and DNA

binding sites (DNA-sites) for each of *D. melanogaster* protein sequence (see Methods). In addition, we

characterized allosteric residues as surface and interior critical residues with STRESS model (Clarke et

al., 2016) for all the structural models. We also extracted putative binding sites from STRESS Monte

Carlo (MC) simulations. We calculated ω, ω_a and ω_{na} for residues in each of the putative molecular

interaction category. Strikingly, we observed that residues involved in protein-protein interactions, DNA

binding and ligand binding exhibited higher rates of adaptive evolution compared to their corresponding

null sites (Fig. 1A-C). In addition, allosteric residues at protein surface showed higher adaptation rates

than allosteric residues at protein interior or residues that are not involved in ligand binding (Fig. 1C).

Since we observed significant positive intercorrelations between PPI and DNA binding with ISD

(intrinsic structural disorder) and RSA (relative solvent accessibility) (Table S2), we next asked whether

the increase of ω_a in protein-protein interactions sites or DNA binding sites was caused by the increase of

disorder or site exposure. We calculated and compared ω, ω_a and ω_{na} for putative PPI and DNA binding

sites with different levels of ISD or RSA. Remarkably, we found that ω_a of these binding sites remains

similar among different levels of ISD or RSA (Fig. S5AC). The results suggest that PPI or DNA binding

events in proteins can result in elevated adaptation rates regardless their structural disorder or site

exposure. While for residues that are not associated with putative PPI or DNA binding, we also observed

increase in ω_a when increasing ISD or RSA (Fig. S5BD), which could be the result of some other yet

unknown underlying mechanisms. In addition, there is possibility that binding sites in disordered regions

are not well-predicted. However, given that ISD does not show strong impact to binding sites (Fig.

S5AC), we think the inaccuracy of binding sites may not play a significant role.

In order to gain better understanding of adaptation in molecular interaction sites, we further

visualized positive selections that are associated with molecular interactions. We first investigated

whether adaptive evolution is associated with particular protein structures or protein families. To do this,

we looked into fast-adaptive proteins with the largest ~15% rates of adaptation ($\omega_a > 0.15$) that are linked

to high quality structural models. Interestingly, among these proteins, we found 45 enriched as trypsin-

like cysteine/serine peptidase domain and 17 7TM chemoreceptors, suggesting widespread adaptive

evolution acting on these protein families or protein domains in *D. melanogaster* (Table S3). Many of the

7TM chemoreceptors are olfactory and gustatory genes and show adaptive evolution in various species

such as *Drosophila* and mosquito (Hill et al., 2002; Lawniczak and Begun, 2007; McBride, 2007; Wu et

al., 2009). In addition to these two protein families, previous studies identified recurrent positive

selections acting on some other fast-adaptive proteins in *Drosophila* and mammals, and the possible

adaptive evolution mechanisms have been linked to exogenous ligand binding, for example, serine

protease inhibitors (serpin), Toll-like receptor 4 (TLR-4), and cytochrome P450 (Jiggins and Kim, 2007;

Slodkowicz and Goldman, 2020).

In order to visualize the link between adaptive evolution and molecular interactions in the two

protein families with frequent adaptive evolution, we showed significant positive selections and

molecular interactions in two representatives: CG10232 and Or67a, each for trypsin-like cysteine/serine

peptidase domain and 7TM chemoreceptors, respectively. We observed that in both cases, positively

selected sites highly overlapped with predicted or inferred binding pockets (Fig. 1D-E). Specifically, in CG10232, we found clusters of positive selected sites around NAG binding sites that are inferred from a crystal structure of serine protease (PDB code: 2XXL) (Fig. 1D), while in Or67a, positively selected sites expand around the putative odorant binding channel formed by helices S1-S6 in extracellular regions (Butterwick et al., 2018) (Fig. 1E).

Except for these examples that are associated with exogenous ligand or exogenous peptide binding, we also identified two previously not described examples where adaptive evolution might be linked to endogenous protein binding: Spaztle (spz, Fig. 1F) and Cul6 (Fig. 1G). Spaztle can bind to Toll-like receptors (TLR) and trigger humoral innate immune response. We built the missing loop in Spaztle in the crystal structure of Toll/Spaztle complex (PDB code 4BV4) according to the dimeric crystal structure of Spaztle (PDB code 3E07). In this complex structural model, we observed several positively selected sites in Toll-4/Spaztle interfaces (Fig. 1F). Cul6, another example, is a protein in cullins family in *D. melanogaster*. The cullins protein family are known as scaffold proteins that assemble multi-subunit Cullin-RING E3 ubiquitin ligase by forming SCF complex with F box and RING-box (Rbx) proteins (Zheng et al., 2002). We constructed the putative Cul6 contained SCF complex by superimposition to the crystal structure of the Cul1-Rbx1-Skp1-F box^{Skp2} SCF ubiquitin ligase complex (Zheng et al., 2002). In the structural model, we observed positive selected sites in Cul6 clustered around the binding sites of RING-box protein, Rbx1, and F-box protein, Skp1 (Fig. 1G).

Frequent adaptive evolution and co-adaptative evolution in genes involved in reproduction, immune system, and environmental information processing

To find out whether specific biological functions were associated with fast-adaptive genes, we applied DAVID Go analysis with genes that have largest ~15% rates of adaptation ($\omega_a > 0.15$). The significant Go terms are frequently linked to serine-type endopeptidase activity, reproduction, protein lysis, chemosensory and other related biological functions (Table S4). As these fast-adaptive genes tend to be enriched in similar biological functions, we asked whether these genes are evolved co-adaptively, i.e., whether these proteins are interacting with each other frequently. To test this possibility, we obtained PPI of *D. melanogaster* from STRING database (Szklarczyk et al., 2019) and analyzed protein-protein interactions among fast-adaptive proteins. We found that fast-adaptive proteins tend to interact with each other more frequently than expected (PPI enrichment p-value < 1.0e-16). In the PPI network of fast-adaptive proteins, we observed 7 strongly connected sub-clusters with at least 5 members (Fig. 2A, Table S5). Proteins in these sub-clusters are enriched in biological processes such as reproduction, immune response, defense response to bacterium and virus, RNA interference, chitin metabolic, etc., which are in line with the Go analysis of fast-adaptive genes (Table S6-S11).

We next asked whether co-adaptation plays a role in the adaptive evolution of interacting proteins

to a broader extend, including both fast- and slow-adaptive proteins. To address this question, we

analyzed and compared adaptation rates of all D. *melanogaster* PPIs available in STRING database with

high confidence and we found that protein partners of fast-adaptive proteins (ω_a>0.15) have significantly

larger maximum/average ω_a compared to slow-adaptive proteins (Figure 3). We further analyzed and

visualized adaptive evolutionary rates of proteins in PPI networks of 9 different biological pathways

extracted from KEGG pathways, including immune system, xenobiotics biodegradation, response to

environment, aging and development, genetic information processing, sensory system, transport and

catabolism, cell growth and death and metabolism. We observed that, in these PPI networks, proteins with

relatively large ω_a tend to interact with each other (Figure 4AB). We also noticed that, for pathways that

are previously known as adaptation-hotspots, e.g., immune system, fast-adaptive proteins can act as

central nodes and are co-adaptively evolved with other fast-adaptive proteins (Figure 4AC). While in

pathways such as transport and catabolism, fast-adaptive proteins are mainly at PPI periphery. In line with

these findings, we found that ω_a are larger in pathways that harbor fast-adaptive proteins as central nodes

than other pathways (Figure S6).

Physical interactions contribute to co-adaptation of fast-adaptive genes. Having established that

molecular interactions contribute to adaptive evolution of protein sequence, we then investigated whether

these physical molecular interactions could drive protein-protein co-adaptation. To do this, we looked into

interacting fast-adaptive protein pairs that are associated known or inferred complex structural models.

For inferred complex structural models, we superimposed the structural models of the pair of proteins

onto their high resolution homologous complex structures. Here we observed and illustrated co-adaptation

at PPI interface in two examples: Toll-4/Spatzle and Spn28Db/CG18563 (Fig. 2BC).

Toll-4/Spatzle. Toll-4 is a member of toll-like receptors. Previous studies have shown strong evidence of

adaptive evolution of Toll-4 in *Drosophila* and mammals (Levin and Malik, 2017; Slodkowicz and

Goldman, 2020). Toll-4 can bind to Spatzle and trigger further innate immune responses with high

confidence (inferred from STRING database). In the previous section, we showed that several positively

selected sites in Spatzle overlap with Toll-Spatzle interfaces (Fig. 1F). Here, we further showed that, in

Toll-4, considerable number of significant positively selected sites were located at interface for Spatzle

(Fig. 2B), which is in line with a previous study of Toll-4 in *D. willistoni* (Levin and Malik, 2017).

Spn28Db/CG18563. Spn28Db is one of the serine protease inhibitors in *D. melanogaster* that are

expressed in male accessory glands, while CG18563 belongs to the protein family of trypsin-like

cysteine/serine peptidase domain. The interactions between the two proteins were predicted with high

confidence from STRING database, and the molecular interactions can be inferred from existing crystal

structure of serpin and bacteria protease complex (PDB code 1EZX). We observed many positive

selected sites at the molecular interface between the two proteins (Fig. 2C), suggesting that physical

interactions might play a role in the co-adaptation of the two proteins.

Most clinally differentiated non-synonymous SNPs in protein-coding genes are adaptive

To find out the relations between short-term adaptation to local environments and long-term adaptive

evolution, we extracted residues with significant F_{ST} SNPs from clinal variations (Svetec et al., 2016). We

then computed evolutionary rates (ω), adaptation rates (ω_a) and non-adaptation rates (ω_{na}) of these

residues as in previous section. We observed that these residues have much higher ratio of adaptation

rates over non-adaptation rates than genome-wide random expectations (Fig. 5A), suggesting that these

residues have higher proportions of adaptive changes, and that they can be hotspots for adaptive

evolution. To find out whether these SNPs are related with even longer-term adaptive evolution, we

inferred positive selection sites of each protein-coding gene from phylogenic data (see Methods). We

found that the non-synonymous F_{ST} SNPs are significantly enriched in long-term positive selections

(Table S12- S13). To further characterize structural and functional properties of short-term genetic

variations, we mapped significant nonsynonymous F_{ST} residues to different structural and functional

characteristics, such as ISD, RSA, PPI-sites, DNA-sites and ligand-binding sites. We found that these

non-synonymous SNPs were enriched in disordered regions and protein surfaces and were significantly

more likely to be involved in protein-protein interactions and ligand-binding than expectation (Table S14-

S18). To better visualize the characteristics of these SNPs, we used Toll-4 as an example. We mapped

significant non-synonymous F_{ST} SNPs in Toll-4 on to its structural model. We showed that F_{ST} SNPs are

either positively selected or being very close to positively selected sites (Fig. 5BC). For example, highly

differentiated sites, N279 (FDR 3e-7) and H431 (FDR 3e-6) were predicted to be positively selected both

at probability at p=0.9. While another highly differentiated site, D424 was close to three positively

selected sites S401 (p=0.8), H431 (p=0.95) and V448 (p=0.8). We also noticed some differentiated sites

that may be located within ligand binding sites, including F297 (FDR 3e-3), S311 (FDR 3e-3), H431

(FDR 3e-6) and H462 (FDR 1e-2).

Discussion

In this study, we systematically studied the impact of structure- and function-related gene properties on

protein sequence evolution and adaptation in *D. melanogaster* genome. We found that molecular

interactions in proteins contribute to the variation of protein sequence adaptive evolution. A novel

discovery of this work is that molecular interaction sites including protein-protein interaction sites and

protein-DNA interaction sites are hotspots for adaptative evolution. We revealed that fast-adaptive

proteins tend to interact with each other frequently and protein partners of these fast-adaptive proteins

tend to have higher adaptation rates, suggesting that co-adaptive evolution might be common in *D*.

melanogaster. By looking at interacting fast-adaptive proteins, we further demonstrated that physical

interactions may contribute to the mechanisms of co-adaptative evolution of fast-adaptive proteins.

Although our results are in agreement with previous studies on the factors driving protein

sequence evolution (Zhang and Yang, 2015), we showed some complex correlations between ω, ω_a and

ω_{na} and protein length and male specificity (Supplement information, section *Complex correlations of*

protein length and male expression level with protein evolutionary rates, Fig. S2-S4, supplement file S2).

These complex correlations suggest caveat exists when we looked at protein length and gene expression

levels. For example, gene expression level was proved to be a major determinant (Zhang and Yang, 2015)

through mechanisms such as the pressure for translational robustness, i.e., robustness to translational

missense errors (Drummond et al., 2005). Previous studies have revealed that male biased or female

biased genes can be fast evolving (Yang et al., 2016). While on the other hand, many male biased genes

can be highly expressed in testis, which results in a complex correlation between protein sequence

evolutionary rate and male expression level or even mean expression level of *D*. *melanogaster*. The

unique evolutionary property of these male biased or specific genes could be caused by the unique

transcriptional scanning mechanism in testis (Xia et al., 2020). We propose that tissue specificity might be

a better quantity when considering the impact of gene expression profile on protein sequence evolution in

D. *melanogaster*. In addition to male expression level, a similar complex correlation was observed for

protein length. It has been the notion that short proteins tend to evolve faster than long proteins, which

may be biologically relevant or byproduct of other factors such as selection on buried and exposed sites

(Moutinho et al., 2019). Here, we demonstrated that, in *D*. *melanogaster*, although protein length is

strongly negatively correlated with protein sequence evolutionary rate, genes that have the slowest

evolutionary rates tend to be relatively short. This could be caused by the fact that under essential

functional constraint, genes can undergo strong purifying selections, while essential genes such as

secreted proteins are constrained to be smaller, and that essential genes could be shorter than other genes

(Chen et al., 2020).

Protein surface and intrinsic disorder regions are frequent targets for adaptive evolution and

contribute to the variations of protein sequence adaptive evolution (Afanasyeva et al., 2018; Moutinho et

al., 2019), however, the detailed mechanisms underlying these observations remains unclear. One

possible explanation would be that these regions are frequently linked to intermolecular interactions

(Afanasyeva et al., 2018; Moutinho et al., 2019). For example, Moutinho et al hypothesized that

molecular interactions involved in host-pathogen coevolution were the major driver of protein adaptation

(Moutinho et al., 2019). Here, we further identified that proportions of possible molecular interaction sites

inside proteins contribute to the variations of protein sequence adaptive evolution and that these

molecular interaction sites or regulatory sites at protein surface can be hotspots of protein adaptation.

Indeed, some specific molecular interactions have been linked to adaptive evolution in several case

studies (Bachtrog, 2008; Hughes and Nei, 1988; Levin and Malik, 2017; Schott et al., 2014) and large-

scale studies based on proteins with high quality structural models (Slodkowicz and Goldman, 2020). In

the latter study, the authors showed that positive selections in mammals tend to cluster closer to binding

sites of exogenous ligands than expected by chance (Slodkowicz and Goldman, 2020), suggesting an

important role of function important regions in adaptive evolution. Here, we extend the conclusion *to D*.

melanogaster genome, including proteins with or without high resolution structural models. We also

showed that except for exogenous ligands, endogenous ligands might also contribution to adaptive

evolution, while the latter might explain why interacting proteins tend to evolve co-adaptively.

Notably, previous studies have revealed that multi-interface proteins tend to be evolving more

slowly than single-interface proteins (Kim et al., 2006), which seems to be contradictory to our results

that proteins with more interaction sites evolve faster and have faster adaptation rates. Here, we argue

that, in our study, we used sequence profile to predict molecular interaction sites in proteins at a genomic

scale, rather than only looking into proteins with high resolution structures. In this way, we may capture

many weak or transient interactions, which are thought to be evolving faster than obligate and conserved

interactions (Mintseris and Weng, 2005). Meanwhile, we did not exclude intrinsic disordered regions

(IDR) or intrinsic disordered proteins (IDP) in our study, which are widespread in *D. melanogaster*

genome. It has been suggested that IDR/IDP tend to evolve fast due to lack of structural restraints

(Echave et al., 2016). In the functional aspect, IDR/IDP are thought to be promiscuous binders through

many multiple binding mechanisms, including forming static, semi-static, and fuzzy or dynamic

complexes (Uversky, 2019), suggesting that the evolution of IDR/IDP cannot be explained merely by the

lack of structural restraints. Actually, IDP and IDR in human genome were found to be undergoing

extensive adaptive evolution (Afanasyeva et al., 2018). At last, it has been recognized that, except for

allosteric regulations, encounter complexes (Gabdoulline and Wade, 1999) might also play an important

role in mediating intermolecular interactions, such as protein-protein association (Tang et al., 2006) and

protein-ligand binding (Re et al., 2019). Since encounter residues that are responsible for encounter

complexes do not reside in conserved binding interfaces, these residues could be under relaxed purifying

selections or even positive selections, which could be another yet-to-identify mechanism that contribute to

protein sequence adaptive evolution.

We showed that fast-adaptive proteins are enriched in molecular functions such as reproduction,

immunity and environmental information processing (Begun and Lindfors, 2005; Begun and Whitley,

2000; Lazzaro et al., 2004). We further demonstrated that fast-adaptive proteins tend to interact with each

other more frequently than random expectations, suggesting co-adaptation might be common among fast-

adaptive proteins. Mechanisms that contribute to the co-adaptation could be: (1) interacting fast-adaptive

proteins are often enriched in similar molecular functions and under similar selective pressure; (2)

interacting fast-adaptive undergo co-evolution through physical interactions. In this study we showed two

examples that adaptive evolution could occur at protein-protein interface, which suggest that physical

interactions could contribute to the co-adaptation of fast-adaptive proteins in $D. melanogaster$. Moreover,

we showed that co-adaptation might exist to a broader extend rather than only among fast-adaptive

proteins. Specifically, proteins that interact with fast-adaptive proteins tend to have higher adaptation

rates. Since molecular interactions contribute to adaptive evolution, it is reasonable to hypothesize that

co-adaptation at this broader extend could be regulated by these interactions. Actually, it has been

suggested that interacting proteins tend to have similar evolutionary rates and the possible mechanism

would be the co-evolution of physical interactions (Pazos and Valencia, 2008).

In this study, we found that loci with significant genetic variance among populations harbor

higher proportions of long-term adaptive changes and these loci follow similar patterns as adaptive

changes, i.e. they are enriched in disordered regions, protein surfaces, and functionally important regions.

These results suggest that population differentiation of protein-coding genes can be an important basis for

long-term adaptive evolution. In other word, many SNPs are repeatedly selected for adaptive process in

evolution. Importantly, our results indicate that most of the clinal amino-acid changes are adaptive,

suggesting that non-selective forces play a non-essential role in the SNPs that show strong geographic

differences. Our results also support a large effect of spatially varying selection on protein sequence and

structures (Storz and Kelly, 2008).

It should be noted that studies at the genomic scale that aim to uncover the function- or structure-

related constraints imposed on protein sequence evolution and adaptation share similar limitations that for

most of the proteins or residues, structural or functional information would be incomplete or even

missing. To overcome this, in this study, we used highly accurate neural-network based tools to predict

molecular interactions, secondary structures, intrinsic structural disorder, relative solvent accessibility for

each of the protein. In this way we were able to identify key factors that impact protein sequence

evolution and adaptation in a less accurate but rather systematic fashion. We hope that with the

availability of more and more curated structural, functional information and complex structural models of

proteins in the near future, we will be able to uncover the precise role of molecular interactions in protein

sequence adaptive evolution.

Material and Methods

d_N/d_S ratio (ω). We used a maximum likelihood method to infer d_N/d_S ratio (ω) of $D. melanogaster$

protein-coding genes using the genome sequences of five species in $melanogaster$ subgroup ($D.$

melanogaster, *D. simulans*, *D. sechellia*, *D. yakuba*, and *D. erecta*). The protein-coding sequences were

extracted from the alignments of 26 insects, which were obtained from UCSC Genome Browser

(http://hgdownload.soe.ucsc.edu/downloads.html). The sequences were further processed by GeneWise

(Birney et al., 2004) to remove possible insertions and deletions using the longest isoforms of the

corresponding *D. melanogaster* protein sequences as references (FlyBase version r6.15) (Thurmond et al.,

2019). The processed sequences were then realigned by PRANK -codon function (Löytynoja, 2014). We

used codeml in PAML (Yang, 2007) to compute gene-specific ω using M0 model. We removed

sequences that have more than 15% of their nucleotides not aligned (gaps) to *D. melanogaster* genes in

more than 2 species. To further avoid numeric errors and ensure reasonable estimations, we only retained

relatively divergent sequences that are: (1) divergent with dS larger than 0.3, (2) less divergent with dS

larger than 0.1 and dN smaller than 0.001 (dS>>dN). At last, there were 12118 genes in total passed all

the criteria and were assigned gene specific ω, containing 6,538,872 amino acids. We also calculated site-

specific ω by using likelihood ratio tests (LRT) comparing M7 model against M8 model (Yang et al.,

2005).

Rate of adaptive and nonadaptive changes. We recalled all SNPs of 205 inbred lines from

the Drosophila Genetic Reference Panel (DGRP), Freeze 2.0 (Huang et al., 2014)

(http://dgrp2.gnets.ncsu.edu). We then generated 410 alternative genomes using all monoallelic and bi-

allelic SNP data sets. We extracted the coding sequences of *D. melanogaster* genes from the generated

alternative genomes, removed all possible insertions and deletions using GeneWise (Birney et al., 2004)

as described above. We then align all the coding sequences to their corresponding aligned CDS sequences

using PRANK -codon function (Löytynoja, 2014). We removed polymorphisms segregating at

frequencies smaller than 5% to reduce possible slightly deleterious mutations (Charlesworth and Eyre-

Walker, 2008). In order to avoid possible effects of low divergence between *D. simulans* and D

melanogaster (Keightley and Eyre-Walker, 2012), we used *D. yakuba* as outgroup to estimate

nonsynonymous polymorphisms (Pn), synonymous polymorphisms (Ps), nonsynonymous substitutions

(Dn) and synonymous substitutions (Ds) by MK.pl (Begun et al., 2007; Langley et al., 2012). Similar as

Begun et al. (Begun et al., 2007), we only analyzed genes with at least six variants for each of

substitutions, polymorphisms, nonsynonymous changes and synonymous changes. We used an extension

of MK test, asymptotic MK (Messer and Petrov, 2013; Uricchio et al., 2019), to estimate the proportions

of adaptive changes (α). The rate of adaptive changes (ω_a) was then calculated as $\omega_a = \omega\alpha$ and the rate of

non-adaptive changes as $\omega_{na} = \omega - \omega_a$. Details of the asymptotic MK test were as following:

(1) Classical McDonald–Kreitman test. According to Smith and Eyre-Walker (Smith and Eyre-Walker,

2002), the proportions of adaptive changes for protein-coding genes can be calculated as following:

373
$$\alpha = 1 - \frac{DsPn}{DnPs}$$

According to this equation, we could estimate the proportion of adaptive changes and carried out classical

MK test by applying Fisher's exact test.

(2) Asymptotic estimation of α. A known problem of the classical estimation of α above is the

accumulation of slightly deleterious mutations at low frequencies. We therefore used an extension of MK

test, asymptotic MK test approach (Messer and Petrov, 2013) to estimate the proportions of adaptive

changes. As in original aMK, we defined α(x) as a function of derived allele frequency (x):

380
$$\alpha(x) = 1 - \frac{DsPn(x)}{DnPs(x)}$$

where Pn(x) and Ps(x) are number of non-synonymous and synonymous polymorphisms at frequency x,

respectively. However, the original approach may suffer from numeric errors when there were very few

polymorphic sites, which is quite common in many of *D. melanogaster* genes. To make the estimations

more robust while preserving the same asymptote, we further define Pn (x) and Ps(x) as total number of

Pn and Ps above frequency x as described in Uricchio et al (Uricchio et al., 2019). We fitted α(x) to an

exponential curve of α(x) ≈ exp(-bx)+c using lmfit (Newville and Stensitzki, 2018) and determined the

asymptotic value of α at the limit of x, 1.0. We then estimate the rate of adaptive changes (ω_a) as

388
$$\omega_a = \frac{N_a/L_N}{dS} = \frac{dN_a}{dS} = \frac{dN_a}{dN} \cdot \frac{dN}{dS} = \alpha\omega$$

where N_a is the number of adaptive changes and $dN_a = N_a/L_N$ is the number of adaptive changes per

nonsynonymous site. Finally, we calculated the rate of nonadaptive changes (ω_{na}) as $\omega_{na} = \omega - \omega_a$. The final

dataset contains 7192 protein-coding genes, with smallest ω_a being 0.00 and largest being 1.29.

Structure-/function- related properties of D. melanogaster proteins. We obtained function-related

properties mentioned in main text as following. We derived *D. melanogaster* gene ages (Kondo et al.,

2017; Zhang et al., 2010) for genes that are specific to *Drosophila*, and from GenTree (Shao et al., 2019)

for genes that are beyond *Drosophila* clade. We then assigned a pseudo-age to each of the genes.

Specifically, there are 11 age groups from "cellular organisms", assigning to a pseudo age value of 0, to

"melanogaster", assigning a pseudo age value of 10. We downloaded *D. melanogaster* protein-protein

interaction (PPI) from STRING database (Szklarczyk et al., 2019). A cut-off of combined score larger

than 0.7 was used to retain high confident PPI for further analysis. We then used BSpred (Mukherjee and

Zhang, 2011) to predict protein-protein interaction (PPI) sites and DRNApred (Yan and Kurgan, 2017) to

predict DNA binding sites. For each protein, we calculated ratios of protein interaction residues (PPI-site

ratio) and ratios of DNA binding residues (DNA-site ratio) by dividing total predicted protein interaction

sites and DNA binding sites over protein length, respectively. For structure-related properties, we used

DeepCNF (Wang et al., 2016) to predict these properties for each gene, including three-state secondary

structures (helix, sheet, and coil), structural disorder, relative solvent accessibility (RSA). Further, we

calculated the ratios of helix, sheet, helix+sheet, and coil residues of each gene from predicted secondary

structures. For each gene, we computed intrinsic structural disorder (ISD) and relative solvent

accessibility (RSA), as protein-length normalized summations of the probabilities of each residue being

disorder and exposed, respectively.

Gene expression patterns. We downloaded gene expression profile from FlyAtlas2 (Leader et al., 2018).

We converted FPKM to TPM by normalizing FPKM against the summation of all FPKMs as following:

$$TPM_i = \frac{FPKM_i}{\sum FPKM_j} \times 10^6$$

After TPM conversion, we only retained genes with expression level larger than 0.1 TPM for further

analysis. We treated male and female whole-body TPM as male and female expression levels. We

calculated mean expression level by averaging male and female TPM. We used following Z-score to

describe male specificities of *D. melanogaster* genes:

$$zscore = \frac{TPM(male\ expression) - TPM(female\ expression)}{\sqrt{sd^2(male\ expression) + sd^2(female\ expression)}}$$

We calculated tissue specificities of genes using tau values (Yanai et al., 2005) based on the expression

profiles of 27 different tissues.

High quality 3D structures of D. melanogaster proteins. We downloaded high-quality structures or

structural models of *D. melanogaster* proteins from protein data bank (PDB) (Burley et al., 2019),

SWISS-MODEL Repository (Bienert et al., 2017), and MODBASE (Pieper et al., 2011), with descending

priorities. For example, if there were 3D structures of a same protein or protein region in multiple

databases, we first considered high-resolution structures from PDB; if no structures were found in PDB,

we then considered SWISS-MODEL Repository; and at last from MODBASE. In addition, we used

blastp (Camacho et al., 2009) to search homologs of each *D. melanogaster* protein against all PDB

sequences with E-value threshold of 0.001. We further carried out comparative structural modeling using

RosettaCM (Song et al., 2013) to model high-quality structural models of proteins or protein regions that

were not available in PDB, SWISS-MODEL Repository and MODBASE. For each RosettaCM

simulation, we used no more than 5 most significant hits from blastp search. For proteins that are in

complex forms, we only extracted monomers for further analysis. At last, we obtained 14543 high quality

structural models, corresponding to 11284 genes. These structural models contain 2,691,913 unique

amino acids, 41.2% of all the residues in genes that were assigned ω.

Evolutionary rates of different structural/functional sites. We classified amino acids into different

classes of structural/functional properties. Specifically, we classified three classes for both ISD and RSA

according the probability of residues being disordered or exposed: ordered or buried (0.00 to 0.33),

medium (0.33 to 0.67), disordered or exposed (0.67 to 1.00). For both PPI and DNA binding, we

classified two classes: PPI-site or DNA-site (binding sites), None-PPI or None-DNA (corresponding null

sites for PPI or DNA binding). For residues that have 3D structures, we used STRESS (Clarke et al.,

2016) to predict putative ligand binding sites and allosteric sites from all the high-quality structures or

structural models. The allosteric sites were further classified as surface critical or interior critical

according to their locations. We then classified these residues into four groups: LIG (ligand binding sites),

Surf. Crit. (surface critical sites), Interior Crit. (interior critical sites) and Others (other sites). For each of

the site classes, we randomly sampled 100 sequences, each containing 10,000 amino acids. We computed

ω, ω_a, and ω_{na} for the randomly sampled sequences similar as the steps described in the above sections.

Acknowledgements

We thank members of the Zhao Lab for helpful discussions.

Author contribution

J.P. and L.Z. conceived the study. J.P. performed the analysis with the input from L.Z.. J.P. and L.Z.

wrote the manuscript.

Funding

The work was supported by NIH MIRA R35GM133780, the Robertson Foundation, a Monique Weill-

Caulier Career Scientist Award, an Alfred P. Sloan Research Fellowship (FG-2018-10627), a Rita Allen

Foundation Scholar Program, and a Vallee Scholar Program (VS-2020-35) to L. Z.. J.P. is supported by a

C. H. Li Memorial Scholar Fund Award at The Rockefeller University.

Declaration of interests

The authors declare no competing interests.

Reference

Afanasyeva, A., Bockwoldt, M., Cooney, C.R., Heiland, I., and Gossmann, T.I. (2018). Human long

intrinsically disordered protein regions are frequent targets of positive selection. Genome Res. *28*, 975–

982.

Bachtrog, D. (2008). Positive selection at the binding sites of the male-specific lethal complex involved in

dosage compensation in Drosophila. Genetics *180*, 1123–1129.

Begun, D.J., and Lindfors, H.A. (2005). Rapid evolution of genomic Acp complement in the

melanogaster subgroup of Drosophila. Mol. Biol. Evol. *22*, 2010–2021.

Begun, D.J., and Whitley, P. (2000). Adaptive evolution of relish, a Drosophila NF-kappaB/IkappaB

protein. Genetics *154*, 1231–1238.

Begun, D.J., Holloway, A.K., Stevens, K., Hillier, L.D.W., Poh, Y.P., Hahn, M.W., Nista, P.M., Jones,

C.D., Kern, A.D., Dewey, C.N., et al. (2007). Population genomics: Whole-genome analysis of

polymorphism and divergence in Drosophila simulans. PLoS Biol. *5*, 2534–2559.

Bienert, S., Waterhouse, A., De Beer, T.A.P., Tauriello, G., Studer, G., Bordoli, L., and Schwede, T.

(2017). The SWISS-MODEL Repository-new features and functionality. Nucleic Acids Res. *45*, D313–

D319.

Birney, E., Clamp, M., and Durbin, R. (2004). GeneWise and Genomewise. Genome Res. *14*, 988–995.

Burley, S.K., Berman, H.M., Bhikadiya, C., Bi, C., Chen, L., Di Costanzo, L., Christie, C., Dalenberg, K.,

Duarte, J.M., Dutta, S., et al. (2019). RCSB Protein Data Bank: Biological macromolecular structures

enabling research and education in fundamental biology, biomedicine, biotechnology and energy. Nucleic

Acids Res. *47*, D464–D474.

Butterwick, J.A., del Mármol, J., Kim, K.H., Kahlson, M.A., Rogow, J.A., Walz, T., and Ruta, V. (2018).

Cryo-EM structure of the insect olfactory receptor Orco. Nature *560*, 447–452.

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., and Madden, T.L.

(2009). BLAST+: Architecture and applications. BMC Bioinformatics *10*, 421.

Charlesworth, J., and Eyre-Walker, A. (2008). The McDonald-Kreitman test and slightly deleterious

mutations. Mol. Biol. Evol. *25*, 1007–1015.

Chen, H., Zhang, Z., Jiang, S., Li, R., Li, W., Zhao, C., Hong, H., Huang, X., Li, H., and Bo, X. (2020).

New insights on human essential genes based on integrated analysis and the construction of the HEGIAP

web-based platform. Brief. Bioinform. *21*, 1397–1410.

Clarke, D., Sethi, A., Li, S., Kumar, S., Chang, R.W.F., Chen, J., and Gerstein, M. (2016). Identifying

Allosteric Hotspots with Dynamics: Application to Inter- and Intra-species Conservation. Structure *24*,

826–837.

Drummond, D.A., Bloom, J.D., Adami, C., Wilke, C.O., and Arnold, F.H. (2005). Why highly expressed

proteins evolve slowly. Proc. Natl. Acad. Sci. U. S. A. *102*, 14338–14343.

Echave, J., Spielman, S.J., and Wilke, C.O. (2016). Causes of evolutionary rate variation among protein

sites. Nat. Rev. Genet. *17*, 109–121.

Enard, D., Cai, L., Gwennap, C., and Petrov, D.A. (2016). Viruses are a dominant driver of protein

adaptation in mammals. Elife *5*, e12469.

Gabdoulline, R.R., and Wade, R.C. (1999). On the protein-protein diffusional encounter complex. J. Mol.
Recognit. *12*, 226–234.

Goldman, N., Thorne, J.L., and Jones, D.T. (1998). Assessing the impact of secondary structure and
solvent accessibility on protein evolution. Genetics *149*, 445–458.

Hill, C.A., Fox, A.N., Pitts, R.J., Kent, L.B., Tan, P.L., Chrystal, M.A., Cravchik, A., Collins, F.H.,
Robertson, H.M., and Zwiebel, L.J. (2002). G Protein-Coupled Receptors in Anopheles gambiae. Science
298, 176–178.

Huang, W., Massouras, A., Inoue, Y., Peiffer, J., Ràmia, M., Tarone, A.M., Turlapati, L., Zichner, T.,
Zhu, D., Lyman, R.F., et al. (2014). Natural variation in genome architecture among 205 Drosophila
melanogaster Genetic Reference Panel lines. Genome Res. *24*, 1193–1208.

Hughes, A.L., and Nei, M. (1988). Pattern of nucleotide substitution at major histocompatibility complex
class I loci reveals overdominant selection. Nature *335*, 167–170.

Jiggins, F.M., and Kim, K.W. (2007). A screen for immunity genes evolving under positive selection in
Drosophila. J. Evol. Biol. *20*, 965–970.

Keightley, P.D., and Eyre-Walker, A. (2012). Estimating the rate of adaptive molecular evolution when
the evolutionary divergence between species is small. J. Mol. Evol. *74*, 61–68.

Kim, P.M., Lu, L.J., Xia, Y., and Gerstein, M.B. (2006). Relating three-dimensional structures to protein
networks provides evolutionary insights. Science *314*, 1938–1941.

Kondo, S., Vedanayagam, J., Mohammed, J., Eizadshenass, S., Kan, L., Pang, N., Aradhya, R., Siepel,
A., Steinhauer, J., and Lai, E.C. (2017). New genes often acquire male specific functions but rarely
become essential in Drosophila. Genes Dev. *31*, 1841–1846.

Kosiol, C., Vinař, T., Da Fonseca, R.R., Hubisz, M.J., Bustamante, C.D., Nielsen, R., and Siepel, A.
(2008). Patterns of positive selection in six mammalian genomes. PLoS Genet. *4*, e1000144.

Langley, C.H., Stevens, K., Cardeno, C., Lee, Y.C.G., Schrider, D.R., Pool, J.E., Langley, S.A., Suarez,
C., Corbett-Detig, R.B., Kolaczkowski, B., et al. (2012). Genomic variation in natural populations of
Drosophila melanogaster. Genetics *192*, 533–598.

Lawniczak, M.K.N., and Begun, D.J. (2007). Molecular population genetics of female-expressed mating-
induced serine proteases in Drosophila melanogaster. Mol. Biol. Evol. *24*, 1944–1951.

Lazzaro, B.P., Sceurman, B.K., and Clark, A.G. (2004). Genetic basis of natural variation in D.
melanogaster antibacterial immunity. Science *303*, 1873–1876.

Leader, D.P., Krause, S.A., Pandit, A., Davies, S.A., and Dow, J.A.T. (2018). FlyAtlas 2: A new version
of the Drosophila melanogaster expression atlas with RNA-Seq, miRNA-Seq and sex-specific data.
Nucleic Acids Res. *46*, D809–D815.

Levin, T.C., and Malik, H.S. (2017). Rapidly evolving Toll-3/4 genes encode male-specific Toll-like

receptors in drosophila. Mol. Biol. Evol. *34*, 2307–2323.

Lin, Y.S., Hsu, W.L., Hwang, J.K., and Li, W.H. (2007). Proportion of solvent-exposed amino acids in a

protein and rate of protein evolution. Mol. Biol. Evol. *24*, 1005–1011.

Lindblad-Toh, K., Garber, M., Zuk, O., Lin, M.F., Parker, B.J., Washietl, S., Kheradpour, P., Ernst, J.,

Jordan, G., Mauceli, E., et al. (2011). A high-resolution map of human evolutionary constraint using 29

mammals. Nature *478*, 476–482.

Löytynoja, A. (2014). Phylogeny-aware alignment with PRANK. Methods Mol. Biol. *1079*, 155–170.

McBride, C.S. (2007). Rapid evolution of smell and taste receptor genes during host specialization in

Drosophila sechellia. Proc. Natl. Acad. Sci. U. S. A. *104*, 4996–5001.

Messer, P.W., and Petrov, D.A. (2013). Frequent adaptation and the McDonald-Kreitman test. Proc. Natl.

Acad. Sci. U. S. A. *110*, 8615–8620.

Mintseris, J., and Weng, Z. (2005). Structure, function, and evolution of transient and obligate protein-

protein interactions. Proc. Natl. Acad. Sci. U. S. A. *102*, 10930–10935.

Moutinho, A.F., Trancoso, F.F., Dutheil, J.Y., and Zhang, J. (2019). The Impact of Protein Architecture

on Adaptive Evolution. Mol. Biol. Evol. *36*, 2013–2028.

Mukherjee, S., and Zhang, Y. (2011). Protein-protein complex structure predictions by multimeric

threading and template recombination. Structure *19*, 955–966.

Newville, M., and Stensitzki, T. (2018). Non-Linear Least-Squares Minimization and Curve-Fitting for

Python. Zenodo.

Nielsen, R., Bustamante, C., Clark, A.G., Glanowski, S., Sackton, T.B., Hubisz, M.J., Fledel-Alon, A.,

Tanenbaum, D.M., Civello, D., White, T.J., et al. (2005). A scan for positively selected genes in the

genomes of humans and chimpanzees. PLoS Biol. *3*, 0976–0985.

Obbard, D.J., Welch, J.J., Kim, K.W., and Jiggins, F.M. (2009). Quantifying adaptive evolution in the

Drosophila immune system. PLoS Genet. *5*, e1000698.

Pál, C., Papp, B., and Hurst, L.D. (2001). Highly expressed genes in yeast evolve slowly. Genetics *158*,

927–931.

Palmer, W.H., Hadfield, J.D., and Obbard, D.J. (2018). RNA-interference pathways display high rates of

adaptive protein evolution in multiple invertebrates. Genetics *208*, 1585–1599.

Pazos, F., and Valencia, A. (2008). Protein co-evolution, co-adaptation and interactions. EMBO J. *27*,

2648–2655.

Pieper, U., Webb, B.M., Barkan, D.T., Schneidman-Duhovny, D., Schlessinger, A., Braberg, H., Yang,

Z., Meng, E.C., Pettersen, E.F., Huang, C.C., et al. (2011). ModBase,a database of annotated comparative

protein structure models,and associated resources. Nucleic Acids Res. *39*, D465–D474.

Ramsey, D.C., Scherrer, M.P., Zhou, T., and Wilke, C.O. (2011). The relationship between relative

solvent accessibility and evolutionary rate in protein evolution. Genetics *188*, 479–488.

Re, S., Oshima, H., Kasahara, K., Kamiya, M., and Sugita, Y. (2019). Encounter complexes and hidden

poses of kinaseinhibitor binding on the free-energy landscape. Proc. Natl. Acad. Sci. U. S. A. *116*,

18404–18409.

Sackton, T.B., Lazzaro, B.P., Schlenke, T.A., Evans, J.D., Hultmark, D., and Clark, A.G. (2007).

Dynamic evolution of the innate immune system in Drosophila. Nat. Genet. *39*, 1461–1468.

Schott, R.K., Refvik, S.P., Hauser, F.E., López-Fernández, H., and Chang, B.S.W. (2014). Divergent

positive selection in rhodopsin from lake and riverine cichlid fishes. Mol. Biol. Evol. *31*, 1149–1165.

Shao, Y., Chen, C., Shen, H., He, B.Z., Yu, D., Jiang, S., Zhao, S., Gao, Z., Zhu, Z., Chen, X., et al.

(2019). GenTree, an integrated resource for analyzing the evolution and function of primate-specific

coding genes. Genome Res. *29*, 682–696.

Sironi, M., Cagliani, R., Forni, D., and Clerici, M. (2015). Evolutionary insights into host-pathogen

interactions from mammalian sequence data. Nat. Rev. Genet. *16*, 224–236.

Slodkowicz, G., and Goldman, N. (2020). Integrated structural and evolutionary analysis reveals common

mechanisms underlying adaptive evolution in mammals. Proc. Natl. Acad. Sci. U. S. A. *117*, 5977–5986.

Smith, N.G.C., and Eyre-Walker, A. (2002). Adaptive protein evolution in Drosophila. Nature *415*, 1022–

1024.

Song, Y., Dimaio, F., Wang, R.Y.R., Kim, D., Miles, C., Brunette, T., Thompson, J., and Baker, D.

(2013). High-resolution comparative modeling with RosettaCM. Structure *21*, 1735–1742.

Storz, J.F., and Kelly, J.K. (2008). Effects of Spatially Varying Selection on Nucleotide Diversity and

Linkage Disequilibrium: Insights From Deer Mouse Globin Genes. Genetics *180*, 367–379.

Svetec, N., Cridland, J.M., Zhao, L., and Begun, D.J. (2016). The Adaptive Significance of Natural

Genetic Variation in the DNA Damage Response of Drosophila melanogaster. PLoS Genet. *12*,

e1005869.

Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva,

N.T., Morris, J.H., Bork, P., et al. (2019). STRING v11: Protein-protein association networks with

increased coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic

Acids Res. *47*, D607–D613.

Tang, C., Iwahara, J., and Clore, G.M. (2006). Visualization of transient encounter complexes in protein-

protein association. Nature *444*, 383–386.

Thurmond, J., Goodman, J.L., Strelets, V.B., Attrill, H., Gramates, L.S., Marygold, S.J., Matthews, B.B.,

Millburn, G., Antonazzo, G., Trovisco, V., et al. (2019). FlyBase 2.0: The next generation. Nucleic Acids

Res. *47*, D759–D765.

Uricchio, L.H., Petrov, D.A., and Enard, D. (2019). Exploiting selection at linked sites to infer the rate

and strength of adaptation. Nat. Ecol. Evol. *3*, 977–984.

Uversky, V.N. (2019). Intrinsically disordered proteins and their "Mysterious" (meta)physics. Front.

Phys. *7*, 10.

Wang, S., Li, W., Liu, S., and Xu, J. (2016). RaptorX-Property: a web server for protein structure

property prediction. Nucleic Acids Res. *44*, W430–W435.

Wu, D.D., Wang, G.D., Irwin, D.M., and Zhang, Y.P. (2009). A profound role for the expansion of

trypsin-like serine protease family in the evolution of hematophagy in mosquito. Mol. Biol. Evol. *26*,

2333–2341.

Xia, B., Yan, Y., Baron, M., Wagner, F., Barkley, D., Chiodin, M., Kim, S.Y., Keefe, D.L., Alukal, J.P.,

Boeke, J.D., et al. (2020). Widespread Transcriptional Scanning in the Testis Modulates Gene Evolution

Rates. Cell *180*, 248-262.e21.

Yan, J., and Kurgan, L. (2017). DRNApred, fast sequence-based method that accurately predicts and

discriminates DNA-and RNA-binding residues. Nucleic Acids Res. *45*.

Yanai, I., Benjamin, H., Shmoish, M., Chalifa-Caspi, V., Shklar, M., Ophir, R., Bar-Even, A., Horn-

Saban, S., Safran, M., Domany, E., et al. (2005). Genome-wide midrange transcription profiles reveal

expression level relationships in human tissue specification. Bioinformatics *21*, 650–659.

Yang, Z. (2007). PAML 4: Phylogenetic analysis by maximum likelihood. Mol. Biol. Evol. *24*, 1586–

1591.

Yang, L., Zhang, Z., and He, S. (2016). Both Male-Biased and Female-Biased Genes Evolve Faster in

Fish Genomes. Genome Biol. Evol. *8*, 3433–3445.

Yang, Z., Wong, W.S.W., and Nielsen, R. (2005). Bayes empirical Bayes inference of amino acid sites

under positive selection. Mol. Biol. Evol. *22*, 1107–1118.

Zhang, J., and He, X. (2005). Significant impact of protein dispensability on the instantaneous rate of

protein evolution. Mol. Biol. Evol. *22*, 1147–1155.

Zhang, J., and Yang, J.R. (2015). Determinants of the rate of protein sequence evolution. Nat. Rev. Genet.

16, 409–420.

Zhang, Y.E., Vibranovski, M.D., Krinsky, B.H., and Long, M. (2010). Age-dependent chromosomal

distribution of male-biased genes in Drosophila. Genome Res. *20*, 1526–1533.

Zheng, N., Schulman, B.A., Song, L., Miller, J.J., Jeffrey, P.D., Wang, P., Chu, C., Koepp, D.M., Elledge,

S.J., Pagano, M., et al. (2002). Structure of the Cul1-Rbx1-Skp1-F boxSkp2 SCF ubiquitin ligase

complex. Nature *416*, 703–709.

Figure 1. Adaptive evolution in molecular interaction sites. Protein-protein interaction sites (A), DNA
binding sites (B) and putative ligand binding sites (C) show higher adaptation rates than none binding
sites. Examples of positive selection around molecular interaction sites in high quality structural models
of CG10232 (D), Or67a (E), spz (F), and Cul6 (G). Except for spz (PDB code 3e07), the other proteins
are obtained from SWISS model repository. Putative ligand binding pockets of CG10232 (D) and Or67a
(E) are shown in blue spheres. Ligands including interacting proteins are shown in cyan or green: NAG of
CG10232 in cyan (D), Toll receptor of spz in cyan (F), RING-box protein in cyan and F-box protein in
green for Cul6 (G). The putative odorant binding channel of Or67a is highlighted in cyan circle (E). The
ligand poses in (D, F and G) are obtained by superimposition from structure 2XXL, 4BV4 and 1LDK,
respectively.

651
Figure 2. Co-adaptation of fast-adaptive proteins. (A) Sub-clusters of PPI networks of fast-adaptive
proteins. Only proteins with at least one partner were shown. Examples of molecular interactions that
might regulate co-adaptation in fast-adaptive proteins: (B) Toll-4 (gray) and spz (orange, with green
representing the other spz monomer), (C) Spn28Db (gray, serine protease inhibitor 28Db) and CG18563
(cyan, with Go term "serine-type endopeptidase activity"). A putative N-terminus (transparent beads) of
Toll-4 were built by superimposition from 4LXR, since the N-terminus were missing in the structural
model. Complex structural model of Spn28Db and CG18563 was inferred from 1EZX.
659

660
661 Figure 3. Co-adaptation of PPIs in *D. melanogaster*. For fast-adaptive proteins, adaptation rates of their
662 partners (orange box plot) are significantly larger compared to slow adaptive proteins (blue box plot).
663 Max ω_a of protein partners are shown in (A and C) and averaged ω_a, of protein partners are shown in (B
664 and D). PPI from STRING with median confidence (combined score larger than 0.4) are shown in (A and
665 B), and PPI with high confidence (combined score larger than 0.7) are shown in (C and D).
666

Figure 4. Rates of protein sequence adaptive evolution in the PPI network of different functional pathways. The PPI networks showed the adaptive evolution in immune system (A) and transport and catabolism (B). (C) In pathways that are hotspots of adaptive evolution, fast-adaptive proteins can act as central nodes, while in conserved pathways, fast-adaptive proteins are often at the periphery of the PPI network.

696
Figure 5. Adaptive evolution in significant nonsynonymous F_{ST} SNPs. (A) The significant SNPs at
different FDR cutoffs all show much higher proportions of adaptation than genome-wide expectation. (B)
Positive selections in Toll-4 and Spaztle, related to Fig. 2B. (C) Significant nonsynonymous F_{ST} SNPs in
Toll-4. Ligands are shown in cyan by superimposing crystal structure of Toll-Spatzle (PDB code 4BV4)
on to Toll-4 structural model. N279, H431 are both highly differentiated (FDR 3e-7 and 3e-6) and
positively selected (both probability at p=0.9). Other highly differentiated sites, F297, S311, H424, H431
and H462 are located near ligand binding sites or positively selected sites.
704
705