Title Page

Thinopyrum intermedium TiAP1 interacts with a chitin deacetylase from

Blumeria graminis f. sp. *tritici* and increases the resistance to *Bgt* in wheat

Yanlin Yang[1,‡], Pan Fan[1,‡], Jingxia Liu[1,‡], Wenjun Xie[2], Na Liu[1], Zubiao Niu[1],
Quanquan Li[1], Jing Song[1], Qiuju Tian[1], Yinguang Bao[1], Honggang Wang[1], Deshun
Feng[1,*]

1. State Key Laboratory of Crop Biology, Shandong Key Laboratory of Crop Biology, College of
Agronomy, Shandong Agricultural University, Tai'an 271018 Shandong, China;
2. Plant Defence Genetics Lab, Department of Plant and Environmental Sciences, University of
Copenhagen, Thorvaldsensvej 40, Frederiksberg C, Denmark

Author for correspondence:

Deshun Feng
Tel: +86 538 8246821
Email: dsfeng@sdau.edu.cn

Thinopyrum intermedium **TiAP1 interacts with a chitin deacetylase from *Blumeria graminis* f. sp. *tritici* and increases the resistance to *Bgt* in wheat**

Yanlin Yang[1,‡], Pan Fan[1,‡], Jingxia Liu[1,‡], Wenjun Xie[2], Na Liu[1], Zubiao Niu[1], Quanquan Li[1], Jing Song[1], Qiuju Tian[1], Yinguang Bao[1], Honggang Wang[1], Deshun Feng[1,*]

1. State Key Laboratory of Crop Biology, Shandong Key Laboratory of Crop Biology, College of Agronomy, Shandong Agricultural University, Tai'an 271018 Shandong, China;
2. Plant Defence Genetics Lab, Department of Plant and Environmental Sciences, University of Copenhagen, Thorvaldsensvej 40, Frederiksberg C, Denmark
* Correspondence: dsfeng@sdau.edu.cn (D.F.); Tel.: +86-0538-8246821(D.F.)
‡,These authors contributed equally to this work.

Summary

The biotrophic fungal pathogen *Blumeria graminis* f. sp. *tritici* (*Bgt*) is a crucial factor causing reduction of global wheat production. Wild wheat relatives, e.g. *Thinopyrum intermedium,* is one of the wild-used parents in wheat disease-resistant breeding. From *T. intermedium* line, we identified the aspartic acid protein gene, *TiAP1*, which involved in resistance against *Bgt*. TiAP1 is a secreted protein that accumulates in large amounts at the infection sites of powdery mildew and extends to the intercellular space. Yeast two-hybrid showed that it interacted with the chitin deacetylase (BgtCDA1) of *Bgt*. The yeast expression, purification, and *invitro* test confirmed the chitin deacetylase activity of BgtCDA1. The bombardment and VIGS mediated host-induced gene silencing showed *BgtCDA1* promotes the invasion of *Bgt*. Transcriptome analysis showed the cell wall xylan metabolism, lignin biosynthesis-related, and defence genes involved in the signal transduction were upregulated in the transgenic *TiAP1* wheat induced by *Bgt*. The TiAP1 in wheat may inactivate the deacetylation function of BgtCDA1, cause chitin oligomers expose to wheat chitin receptor, then trigger the wheat immune response to inhibit the growth and penetration of *Bgt*, and thereby enhance the tolerance of wheat to pathogens.

Key words:

Blumeria graminis f. sp. *tritici*, chitin deacetylase, disease resistance, *Thinopyrum*

intermedium, *TiAP1*, wheat

Introduction

Wheat (*Triticum aestivum* L.) is one of the main crops in the world, therefore the

production of wheat is closely related to food security. In recent years, pathogenic

infections have seriously affected the yield of wheat. Wheat powdery mildew caused

by the obligate biotrophic fungus *Blumeria graminis* f. sp. *tritici* (*Bgt*) can result in

severe reductions in grain yield (Singh *et al*., 2016). When powdery mildew invades

the plant, it releases proteinaceous exudates that contain the mechanical and

molecular features required for the full virulence of the pathogen (Zhang *et al*., 2005).

Fungal exudates function in the attachment of conidia and in the adherence of the

penetrating appressoria (App) to invade plants (Deising *et al*., 2000). At the same time,

the plant responds immediately to the pathogen invasion, and a number of secreted

proteins, especially protease, from plant cells accumulate in the apoplastic space to

carry an apoplastic battle between hosts and pathogen (Geziel, 2015; Wang et al.,

2020).

In wheat, more than 66 powdery mildew resistance loci (*Pm1-Pm66*) located on

different chromosomes of the common wheat have been named (Li *et al*., 2020a).

Most of them originated from wild relative species of wheat, such as *Pm21* from *H.*

villosa has been introduced into the common wheat using the translocation line

T6VS·6AL (Chen *et al*., 2013). Li *et al*. (2020b) cloned *Pm41*, which was a powdery

mildew resistance gene deriving from the wild emmer wheat. *Thinopyrum*

intermedium had been hybridized extensively with wheat and proven a useful source

of resistance to various diseases of hexaploid wheat, and several powdery mildew

resistance genes, such as *Pm40*, *Pm43*, and *PmL962* (Luo et al., 2009; He et al., 2009;

Shen et al., 2015), have been identified and genetic mapped. The mining and

utilization of these new resistance genes are vital means to improve current wheat

varieties and increase their resistance to powdery mildew.

Aspartic proteases consist of four main proteolytic enzymes that are widely present

in animals, plants, yeast, microorganisms, and viruses (Rawlings and Barrett, 1995).

Xia *et al*. (2004) reported that the peptide signal system of CDR1 was involved in the

activation of the resistance mechanism, while Prasad *et al*. (2009, 2010) found that

OsCDR1/OsAP5 could enhance the disease resistance of rice and *Arabidopsis*

thaliana. The aspartic protease gene, *AP13*, from grape could promote the salicylic

acid-dependent signal transduction pathway (Guo *et al*., 2016). Moreover, Alam *et al*.

(2014) found that the rice *OsAP77* gene accumulated in the extracellular space and

sieve tubes to defend the pathogens. Hence, these studies have shown aspartic

proteases participate in the biotic stresses of plants to enhance the resistance.

To resist pathogens, plants have developed a complex immune system including the

plasma membrane receptors that recognise pathogen-related molecular patterns, such

as chitin from the fungal cell walls that can trigger defence responses. Kamakura *et al*.

(2002) found a new germination tube-specific gene, *CBP1*, in the rice blast fungus

Magnaporthe grisea. This gene encodes a chitin-binding protein (CBP) having two

similar chitin-binding domains at both sides of the central domain. This structure is

similar to the fungus chitin deacetylase, which may play a vital role in the

hydrophobic surface sensing of *M. grisea* during App differentiation. Kuroki *et al*.

(2017) confirmed *CBP1* encoded a chitin deacetylase and participated in

differentiation stage of App in *M. oryzae*. These results proved the chitin deacetylase

activity of CBP1 is necessary for the formation of App. In addition, Yang *et al*. (2019)

identified the chitinase gene *MoChia1* from *M*. *oryzae*. This gene can trigger a plant's

defence response to *M. oryzae* in rice under an inducible promoter. MoChia1 was also

a functional chitinase that is required for the growth and development of *M. oryzae*.

The MoChia1 binding to free chitin could inhibit the plant immune response, while

another protein, OsTPR1, competitively binding with MoChia 1, would re-establish

the immune response (Yang *et al*., 2019). Han *et al*. (2019) also found the mechanism

of MoChi1 that targeted the host lectin to inhibit rice immunity and promote the

colonisation. Furthermore, they also found rice protein OsMBL1 could interact with

MoChi1 to enhance the resistance against *M. oryzae*. Overexpressing of *OsMBL1* can

lead to the activation of defence response genes of rice and the burst of chitin-induced

reactive oxygen species (ROS). Moreover, Gao *et al*. (2019) identified a secreted polysaccharide deacetylase (PDA1) from soil-borne *Verticillium dahliae*. PDA1 can promote the deacetylation of chitin oligomers, whose *N*-acetyl group contributes to the host lysine motif (LysM)-containing receptor that perceives ligand-triggered immunity and facilitates virulence. Thus, the silencing of PDA1 allows the N-acetyl group of the chitin-triggered host immunity to occur.

In a previous study, we obtained an aspartic protease gene, *TiAP1*, from trititrigia SN6306 through a comparative transcriptome analysis. Further sequence analysis showed *TiAP1* originated from *T. intermedium* (Tian *et al*., 2017). Through gene expression and virus-induced gene silencing (VIGS) analyses, we found the *TiAP1* gene might be involved in the resistance to wheat powdery mildew. In addition, subcellular localisation analysis showed that TiAP1 is of a secretory protein. During the powdery mildew invasion, TiAP1 would assemble in large numbers at the infect site. The yeast two-hybrid (Y2H) verification, luciferase (LUC) complementation imaging (LCI), and bimolecular florescent complimentary (BiFC) analysis show that TiAP1 interacts with chitin deacetylase (BgtCDA1) of *Bgt*, while BgtCDA1 could promote the invasion of *Bgt*. Therefore, we propose that TiAP1 induces the plant to resist the invasion of pathogens by interacting with BgtCDA1, which makes the chitin of the fungus inherently immune to the plant.

Results

The expression of *TiAP1* mediates the *Bgt*- resistance in trititrigia SN6306 to *Bgt* invasion

Since Li *et al*. (2016) analyzed the transcriptome of SN6306 and YN15, and obtained a gene that was upregulated by *Bgt* induction, which encoded an aspartic protease, while Tian *et al*. (2017) cloned the *TiAP1* gene and found that it came from *T. intermedium*. Here, we found that the *TiAP1* gene was expressed in SN6306 not in wheat YN15, and was upregulated by the *Bgt*-induced in the early stages of the Bgt invasion (Fig. 1A).

To investigate the effect of endogenous *TiAP1* on SN6306 resistance to *Bgt*, we reduced the endogenous *TiAP1* gene levels by suppressing *TiAP1* expression using

BSMV-VIGS. The *TiAP1* expression levels were significantly suppressed in the

leaves of BMSV:TiAP1as SN6306 relative to the BMSV:00 negative control (Fig. 1B).

When the BMSV:TiAP1as SN6306 was inoculated with *Bgt* at the two-leaf stage, the

infection in BMSV:TiAP1as SN6306 was much more severe than in the BMSV:00

control at 10 days post-inoculation (dpi) (Figs. 1C, 1D). This proved that the *TiAP1*

gene was resistant to *Bgt* invasion and could play a role in the resistance.

161 The TiAP1 protein enhances resistance to *Bgt in vitro*

Western blot analysis showed that TiAP1 was induced in the precipitation through

prokaryotic expression (Figs. 1E, 1F, S1). After renaturation, the protein concentration

was determined to be 0.437 mg/mL by the Bradford method (Fig. 1G; Bradford,

1976). Moreover, regarding the TiAP1 expression effect on the *Bgt* invasion, the

number of *Bgt* conidia on YN15 leaf after smearing TiAP1 for 5 days was

significantly lower than that of YN15 with CK (pET28a). In addition, there was a

small number of hyphae in the leaves smeared with TiAP1, where the hyphal length

and the diseased area were less than those of the leaves smeared with CK (Figs. 1H,

1I). Lesion areas of the leaves treated with TiAP1 were also significantly less than

that of the control (Fig. 1J), suggesting that TiAP1 can resist the *Bgt* invasion *in vitro*.

172 *TiAP1* gene transfer can elevate the resistance to *Bgt* of the recipients wheat

To investigate the function of *TiAP1*, the pMUbi-*TiAP1* vector and the screening

marker *Bar* gene vector, were co-transformed into the susceptible hexaploid wheat

cultivar, Bobwhite. The obtained transgenic Bobwhite with the *TiAP1* gene were

subjected to PCR identification of the *Bar* and *TiAP1* genes (Figs. S2A, S2B) to

obtain the positive transgenic plants. The qRT-PCR analysis showed that the

expression level of the *TiAP1* gene was higher in OE1 and OE2 of SN6306 when

infected by *Bgt* 2 dpi, respectively. After multiple comparisons, we found that the

expression level of *TiAP1* when infected by *Bgt* in OE2-2 was significantly different

from that of OE2-0 ($P < 0.05$) (Fig. 2A), indicating that the expression of *TiAP1* could

be involved in the interaction with *Bgt*. In addition, we found that the overexpression

of *TiAP1* gene in Bobwhite (OE1 or OE2) showed moderate resistance, while the leaf

surface was still entirely green and less conidia were present at 5 dpi than the control

Bobwhite (Fig. 2B). Moreover, there were a large number of conidia in Bobwhite that

had germinated and formed hyphae 2 dpi using *Bgt*. A large number of conidia in the

OE1-2, OE2-2 and SN6306 were retained at App of 41.2 %, 43.4 % and 83.7 %, and

16.2 %, 14.5% and 0.34 % retained at hyphae stage, respectively, while in the control

Bobwhite, 28.9 % of the conidia were present in the form of App and 25.6 % of the

conidia were present in the form of hyphae (Figs. 2C, 2D). The HI in Bob-2 was

68.6%, while in OE1-2, OE2-2 and SN6306 it was 48.8 %, 53.5 % and 6.7 %,

respectively (Fig.2F). In addition, figures 2D and 2E suggested that the transgenic

TiAP1 gene could affect the formation of *Bgt* Hau, thereby resisting the invasion of

Bgt.

Meanwhile, we crossbred between the homozygous transgenic *TiAP1* Bobwhite and

YN15 to obtain the hybrid F_1. Gene amplification and phenotype investigation

confirmed the inheritance of the *TiAP1* gene and its function in resisting *Bgt* in the

hybrid F_1. A chi-square test showed that the proportion of resistant-susceptible

separation in the F_2 generation was 3:1. The subsequent F_3 generation at the seedling

stage (Figs. S2C, S3; Table S1) also double confirmed *TiAP1* endue the resistance to

Bgt for wheat.

In the adult stage, through natural infection by powdery mildew in the glasshouse,

the overexpression of *TiAP1* gene in Bobwhite showed high resistance to *Bgt* (Figs.

2F2, S4; Table S2), whereas Bobwhite (Fig. 2F1) and YN15 (Fig. 2F3) showed high

susceptibility to *Bgt* on the leaves with many lesions of more than 1 mm in diameter

and more hyphae. At the same time, the hybrid F_1 between the overexpression of

TiAP1 gene in Bobwhite and YN15, showed high resistance to *Bgt*, where the

diameter of the lesions was less than 1 mm, the amount of the leisons was little, the

whole leaf surface was still green (Fig. 2F4). .

TiAP1 is a secretory protein that accumulates at the infection site

To determine the localisation of TiAP1 in plant cells, we utilised the TiAP1-GFP,

Δsp-TiAP1-GFP, and free GFP transient transformation in *Nicotiana benthamiana*.

Through plasmolysis methods, we found that the fluorescent signals of TiAP1-GFP

were outside the cell membrane, while Δsp-TiAP1-GFP (without signal peptide) was in the cytoplasm (Figs. 3A, 3B, 3C, S5). To further confirmation whether TiAP1 is a secretory protein, the mCherry-TaSYP51, which is a member of the syntaxin superfamily on plasma membrane, was co-bombarded with TiAP1-mYFP into the barley leaf epidermal cells. We found a large numbers TiAP1 and TaSYP51 proteins accumulated at the infection site of the primary and App germination tube of *Bgh*, and TiAP1 extended to the barley intercellular space (Fig. 3D; Video S1). Integrating with our observations of the *Bgt* on the leaves of *TiAP1* gene transferred wheat, we speculated that TiAP1 restrains the powdery mildew from penetration, assisting the plants in resisting the invasion of *Bgt*.

The BgCDA1 interact with TiAP1

To clarify how TiAP1 can regulate wheat resistance to *Bgt*, CDS of TiAP1 without signal peptide was fused with the GAL4 DNA-binding domain sequence to generate the bait vector. We performed a Y2H screen assay using *Bgt* cDNAs as a prey library. Interestingly, one of the obtained interactors with highly similar (95.06%) to the chitin deacetylase (EPQ66796.1) of *Bgt*. Furthermore, the analysis of the cDNA revealed that the related protein contains a polysaccharide deacetylation domain. Therefore, we named this gene chitin deacetylase (*BgtCDA1*) of *Bgt* (Figs. S6A, S6B). The BgtCDA1 has an N-terminal 19-aminoacid signal peptide (http://www.cbs.dtu.dk/services/SignalP) and five conserved motifs that are required for hydrolysing the acetyl groups of the substrates (Figs. S6C, S6D). Moreover, the point-to-point verified the BgtCDA1 without signal peptide fused to the GAL4 activation domain interact with TiAP1 (Fig. 4A).

Subsequently, we furtherly confirmed the interaction through LCI, and BiFC. As expected, the LUC activity in the *N. benthamiana* leaves was high when co-injected using the ΔspTiAP1-nLuc and ΔspBgtCDA1-cLuc, while the other three combinations were used as negative controls (Fig. 4B). The BiFC analysis also confirmed that the co-expression of nGFP-ΔspBgtCDA1 and cGFP-ΔspTiAP1 resulted in a clear GFP signal in the cytoplasm. In contrast, no visible signal was detected in any of the

243 corresponding negative controls (Fig. 4D). Therefore, these observations indicate that

244 there is an *in vivo* interaction between TiAP1 and BgtCDA1.

HIGS analysis showed that the silencing of *BgtCDA1* inhibits the penetration and Hau formation of *Bgt*

To confirm the role of *BgtCDA1* in the growth and development of *Bgt*, we used a

transient transformation system based on the particle bombardment of RNAi

constructs (Douchkov *et al.*, 2005). Through statistical analysis, we found that after

the normalization treatment, the HI of the positive control, Mlo-RNAi, was 47.44%,

while that of *BgtCDA1* RNAi was 55.36% (Fig. 5A, 5B). And through BSMV-VIGS,

we also found that the lesions area in the leaves of BSMV: BgtCDA1as wheat

infected with powdery mildew for 5 days and the hyphal density score of powdery

mildew infection for 3 days were significantly lower than those of BSMV:00 wheat

(Figs. 5C, 5D, 5E, 5F). This indicated that *BgtCDA1* gene silencing suppressed the

growth of powdery mildew conidia and the *BgtCDA1* gene promoted the invasion of

Bgt.

The chitin deacetylase activity of BgtCDA1

We examined the chitin deacetylase activity of BgtCDA1. After Pichi yeast

expression, the purified BgtCDA1-HIS (2 uM) was incubated with 1 mM

Chitooctaose with six GlcNAc moieties (A6) at 37 □ in 50 mM Tris-HCl buffer

（pH8.0）, followed by detection of the enzymatic reaction products using

MALDI-TOF-MS. As shown in Fig. 5G, the deacetylated intermediate products A5D1

(where A is N-acetyl-D-glucosamine and D is D-glucosamine) were detected after 30

min incubation. It showed that BgtCDA1 has the deacetylase activity.

TiAP1 and BgtCDA1 co-regulate the expression of wheat pathogen-responsive genes

To evaluate the impact of TiAP1 and BgtCDA1 on the pathogen-responsive gene

expression on a genome-wide scale, we performed RNA-Seq experiments using

Bobwhite and OE2 seedlings inoculated with *E09* for two days (Fig. S7A). Genes

with more than a four-fold change in expression ($P < 0.01$) were considered to be

differentially expressed. Total 274 differentially expressed genes were detected in the

OE2-2-vs-Bob-2 after 2 dpi, of which 116 genes were upregulated when inoculated

with *E09* in the OE2-2 (Figs. S7B, S7C; Table S3). Gene ontology (GO) analysis

indicated that these differentially expressed genes were enriched in the chitin

catabolic process, cell wall macromolecule catabolic process, defence response to

bacterium and fungus, chitinase activity, and chitin-binding (Fig. S7D; Table S4).

Defence-related genes were also upregulated in OE2-2 relative to Bob-2 (Fig. 6A;

Table   S5),   such   as   *traesCS1A02G410500*,   *traesCS2B02G187500*,

traesCS2A02G161500, *traesCS3B02G379200* and *traesCS1B02G440300* that encode

proteins   containing   WRKY   domains;   *traesCS5A02G049600*   encodes

phytochrome-interacting bHLH; *traesCS7A02G201100* encodes TIFY family gene;

traesCS5D02G188600   encodes   protein   phosphatase   2C   (PP2C);

traesCS6D02G217800 and *traesCS3A02G404400* encode ERF transcription factors;

traesCS6B02G018700 encodes LigB domain-containing protein, and may involved in

betaine metabolism; *traesCS7D02G201600* and *traesCS6D02G237900* are ubiquitin

E3 ligase RING-type related genes; and *traesCS6A02G266100* that encode the

xyloglucan glycosyltransferase/hydrolase and may be involved in cell wall xylan

metabolism. By qRT-PCR, we further analysed twelve gene which was differential

expression induced of *Bgt* for two days. Among these twelve genes, the expression

levels   of   *TraesCS1B02G440300*,   *TraesCS2A02G161500*,   *TraesCS2A02G199300,*

TraesCS3B02G379200, TraesCS5D02G188600 and *TraesCS5A02G049600* which

related to plant hormone signal transduction were all upregulated in OE2-2 relative to

Bob-2, beside *TraesCS5D02G188600* which was no difference in OE2-2 relative to

Bob-2   (Fig.   6B).   The   *TraesCS3B02G259500*,   *TraesCS2B02G224300*,

TraesCS6A02G266100 and *TraesCS2D02G099900* involved in cell wall anabolism, in

which   *TraesCS3B02G259500*   and   *TraesCS6A02G266100*   were   consistent   with

transcriptome were upregulated in OE2-2 relative to Bob-2, *TraesCS2B02G224300*

and *TraesCS2D02G099900* were no difference in OE2-2 relative to Bob-2, but the

expression in OE2-2 was higher than Bob-2, and *TraesCS6B02G018700* and *TraesCS2B02G529400* genes expression pattern were similar with *TraesCS2B02G224300* and *TraesCS2D02G099900* (Fig. 6B). These quantitative results were slightly different to those at the transcriptome, possibly because of the differences between individuals. In summary, the upregulated expression of these genes provided favourable evidence to understand the disease resistance mechanism of the *TiAP1* gene.

Discussion

Aspartic proteases play a vital regulatory role in plant growth and may be involved in the senescence process (Chen *et al*., 2015), plant reproduction (Niu *et al*., 2013), programmed cell death (Chen and Foolad, 1997), resistance to pathogens (Prasad *et al*., 2010), response to stress (Guo *et al*., 2013) and endoproteases in plants (Rawlings *et al*., 2016). If the endoproteases encoding genes were deleted or inhibited, the plant's sensitivity to pathogens increases (Jashni *et al*., 2015; Díaz *et al*., 2018). In the early stage, our laboratory cloned an aspartic protease (Tian et al., 2017). Through multiple sequence comparisons, we found that the TiAP1 gene is 98% similar to scaffold 208 assembled from the *T. intermedium* genome (https://jgi.doe.gov/), and Yang and Feng (2020) analysed the aspartic protease gene family and their response under powdery mildew stress in wheat. Among those family genes, the AP gene which most similar to *TiAP1*, TRIAE_CS42_5DS_TGACv1_456657_AA1475640.1, were located on the chromosome 5DS with 70% identity. The qRT-PCR analysis showed that the *TiAP1* gene was rapidly expressed under the induction of *Bgt* in SN6306 with no expression in YN15 (Fig. 1A). *TiAP1* expression levels were significantly suppressed with more hyphae piles appearing in the leaves of BMSV:TiAP1as SN6306 (Figs. 1B, 1C, 1D) as well as lesion areas on the leaves that were smeared with the TiAP1 protein level when compared to the control (Figs. 1H–1J). These results indicated that *TiAP1* might resist the invasion of powdery mildew.

Furthermore, the identification of the *Bgt* resistance phenotype at the seedling stage

showed that the overexpression of *TiAP1* gene in wheat was moderately resistant to

powdery mildew (Fig. 2B), while the life cycle of *Bgt* in the two-day-infected leaves

of the overexpression of *TiAP1* gene in Bobwhite was primarily retained in the App

stage. In contrast, the conidia on the Bobwhite leaves were mainly in Hau stage. Hau

was wrapped in the extrahaustorial membrane (EHM), which was usually used for

signal exchange and nutrient absorption (Panstruga, 2013), where the development of

the Hau reflected the growth and metabolism of the entire pathogen to a certain extent.

It was a vital trait for evaluating host cell resistance (Bracket, 1968).

The HI of *Bgt* in Bobwhite was significantly higher than that of the transgenic

TiAP1 gene lines, indicating that overexpression of *TiAP1* gene in Bobwhite were

resistant to the *Bgt* invasion (Fig. 2E). This resistance may be because the transgenic

wheat inhibited the penetration of the powdery mildew, thereby affecting their

infection rate. Subsequently, we found that the transgenic *TiAP1* Bobwhite and the F_1

generation showed high resistance to powdery mildew in the adult-stage (Figs. 2F2,

2F4, S4). Many studies refer to this trait, which showed less significant resistance to

the disease at the seedling stage than that of the adult stage (Hautea *et al*., 1987;

Griffey and Das, 1994), or of a chronic disease resistance type (Shaner, 1973).

Moreover, Shaner (1973) found condia of Konx, a typical slow-white powder variety,

could not successfully invade. Therefore, we speculated that the transgenic wheat line

with the *TiAP1* gene might be used as a chronic powdery mildew-resistant variety to

prevent the invasion of *Bgt*. Syntaxins were members of the SNARE superfamily of

proteins that mediated membrane fusion events (Colline *et al*., 2003). Since TiAP1

was a secreted protein (Figs. 3A, 3B, 3C) which expressed in large quantities at

intercellular space of the infection site (Fig. 3D, Video S1), we speculate that syntaxin

TaSYP51 may help to transport the TiAP1 protein to the outside of the cell, where the

detailed mechanisms need to be further elucidated.

Through the Y2H, LCI, and BiFC *in vivo* verification, we determined TiAP1

interacts with BgtCDA1 (Fig. 4). Subsequently, we found that when the *BgtCDA1*

gene was silenced, the HI, lesions area and hyphal density score were significantly

reduced (Fig. 5). Similarly, Zhao *et al*. (2010) found that chitin deacetylase was

secreted when it interacted with other substances. The chitin deacetylase of *C.*
lindemuthianum was secreted during the process of fungal hyphae invading plants to
modify the chitin that can be recognised by the plant's resistance system (Tsigos and
Bouriotis, 1995). Geoghegan and Gurr (2017) also found in *M. oryzae* that CDA1 and
CDA4 were necessary for chitin deacetylation in the cell wall of the mycelium.
During infection, the ability of the fungal cells to convert chitin to chitosan was
critical for masking the cell wall chitin from being recognised by the host chitinases,
thereby promoting the fungal attack (Upadhya *et al*., 2018). Chitin oligomers can also
act as ligands that were recognised by LysM receptors on the cell surface and
triggered pathogen-associated molecular pattern (PAMP)-triggered immunity (PTI)
(Hurlburt *et al*., 2018; Gao *et al*., 2019). We found the silencing of *BgtCDA1* can
inhibit the invasion of *Bgt* and speculated that the interaction between BgtCDA1 and
TiAP1 made BgtCDA1 be degraded by TiAP1 which inhibit the deacetylation
function of BgtCDA1. Moreover, the growth and penetration of *Bgt* was inhibited and
the chitinase of the wheat was secreted to release the chitin oligomers, which were
recognised by the wheat chitin receptor, such as LysM. This led to PTI and caused the
transgenic wheat with the *TiAP1* gene to be resistant to powdery mildew (Fig. 7).
When the pathogen invades the host, an extremely complex and precise arms race of
'attack and defence' occurs in the apoplastic space. Xia *et al*. (2020) found that
soybean could secreted an apoplastic aspartic protease, GmAP5, to bind and degrade a
pathogen-secreted apoplastic endoglucanase PsXEG1 to destroy its virulence in
Phytophthora sojae invasion, and proposed a 'multi-layered immune model' of the
plants against pathogens in the extracellular region. Therefore, this study revealed the
complexity of the interaction between pathogens and hosts as well as provided a
theoretical basis to further study the role of TiAP1 in resistance mechanisms.
When chitin receptors recognise the chitin, the plant can active defence genes and
release some metabolites that initiate the signal transduction in plant cells to trigger an
immune response. Many defence-related genes were upregulated in OE2-2 relative to
Bob-2 (Fig. 6A; Table S5). WRKY TFs are key nodes in abscisic acid
(ABA)-responsive signaling networks (Rushton et al., 2012) and Zhang et al. (2015)

found the OsWRKY24, -53, and -70 are involved in gibberellic acid. Therefore, we speculate that these genes may be involved in the ABA or gibberellic acid (GA) signaling pathways. Phytochrome-interacting bHLH. These proteins are known to play important roles in red light-mediated (phyA and/or phyB-dependent) signal transduction pathways at immediate positions downstream of the photoreceptor in *A. thaliana* (Nakamura et al., 2007). In *A. thaliana*, the TIFY family may be involved in jasmonic acid (JA) pathway (Chung and Howe, 2009). PP2Cs are vitally involved in ABA signaling (Park et al., 2009). The ERF transcription factor family are involved in ethylene-activated signaling pathway (Xu et al., 2008). The metabolites include ABA, ethylene, JA, GA, and ubiquitination that caused the plants to respond to the invasion of pathogens (Fig. 7; Table S3). Moreover, RNA-Seq and qRT-PCR analysis revealed that there was a cell wall xylan metabolism gene, *traesCS6A02G266100*, whose expression in Bobwhite was sharply reduced after being induced by powdery mildew, while the gene in OE2-2 was significantly upregulated relative to that of Bob-2 (Fig. 6).

Since xylan was a type of hemicellulose that was closely related to functional proteins in the cell wall, it had a great influence on the ductility, pore elasticity, flow, and rheological properties of plant cells during growth (Zhou *et al*., 2017). Therefore, we infer that the penetration during the powdery mildew infection of the transgenic wheat with the *TiAP1* gene was blocked. We also found that the *TraesCS2B02G224300* gene encoded a phenylalanine ammonia-lyase (PAL), which was upregulated and significant difference in OE2-2 relative to Bob-2 in transcriptome, (Fig. 6A; Table S5) and in qRT-PCR the expression of *TraesCS2B02G224300* in OE2-2 was twice higher than Bob-2 (Fig. 6B) and the PAL was pivotal for the enzymes related to the biosynthesis of lignin, salicylic acid, and other phenylalanine-derived metabolites (You *et al*., 2020). As a main component of the plant cell wall, lignin contributes to the formation of the physical and chemical barriers in the plant immunity. Zhou *et al*. (2018) also found that *OsPAL1*-mediated resistance to rice blast might be due to the accumulation of lignin, while the broad-spectrum blast-resistant rice varieties *Pigm-* and *PIBP1*-mediated resistance

might be related to lignin content (Zhai *et al*., 2019). Similarly, the overexpression of

OsPAL8 in rice can lead to increased lignin content and enhanced brown plant hopper

resistance (He *et al*., 2020). Thus, these results demonstrate that the transgenic wheat

with the *TiAP1* gene could enhance resistance to *Bgt*.

In summary, we speculate the fungal chitin deacetylase BgtCDA1 can block the

PAMP processes that the fungal derived chitin oligoes are recognized by wheat chitin

receptor protein and then trigger an immune response, while TiAP1 competitively

interacting with BgtCDA1 is the causal reason to re-establish resistance to fungal

pathogen *Bgt*. Moreover, the overexpression of *TiAP1* upregulates the cell

wall-related genes in transgenic *TiAP1* wheat thereby blocks the penetration of

powdery mildew and enhances the tolerance to pathogens. Furtheremore, we conclude

many other plants may share similar mechanism and this mechanism can be used for

future genetic improvement of crops. However, the mechanism of BgtCDA1 degraded

by TiAP1, and the resulting detailed disease resistance pathway remains to be studied.

Materials and methods

Plant material and pathogen infection

The trititrigia SN6306 was highly resistant to the powdery mildew and obtained by

hybridisation between the highly susceptible to powdery mildew wheat (*Triticum*

aestivum L.) cultivar Yannong 15 (YN15) and *T. intermedium* (Li *et al*., 2016). Wheat

Bobwhite were highly susceptible to the powdery mildew. The transgenic

overexpression *TiAP1* in Bobwhite line 1 and 2 were named as OE1 and OE2,

respectively. The untreated with *Bgt* wheat lines Bobwhite, OE1 and OE2 were named

as Bob-0, OE1-0 and OE2-0, respectively. The Bobwhite, OE1 and OE2 wheat

inoculated with *E09* for two days were named as Bob-2, OE1-2 and OE2-2,

respectively. The seeds of these materials above and *N. benthamiana* were sown in a

mixed soil (loess, matrix, and vermiculite of 1:1:1, *v/v/v*) at 25°C, 70% humidity, and

a long-term photoperiod of a 16 h/8 h light/dark cycle in a glasshouse with a light

intensity of 600 mmol $m^{-2}s^{-1}$. The virulent isolate *E09* of the wheat powdery mildew

fungus (*Bgt*) was propagated on the wheat variety YN15 via a weekly inoculum

transfer. The barley (*Hordeum vulgare* L.) line Ingrid *ror2* (Collins *et al.*, 2003) was

highly susceptible to *Blumeria graminis* f. sp. *hordei* (*Bgh*) and used for the

localisation and expression profiling of *TiAP1*. The *Bgh* isolate B103 was propagated

on barley P-10 via a weekly inoculum transfer, while the *Escherichia coli* (*E. coli*)

strain BL21 (DE3) was used for the prokaryotic expression of TiAP1.

VIGS, barley stripe mosaic virus (BSMV), and host-induced gene silencing

(HIGS)

VIGS analysis was conducted according to Yuan *et al.* (2011) to insert the conserved

sequence of the target *TiAP1* and *BgtCDA1* gene (316 bp and 300 bp, respectively)

into the BSMV vector to create BSMV:TiAP1as and BSMV:BgtCDA1as and

transform the *Agrobacterium*. The detailed method was described in Methods S1. Ten

days after inoculation with BSMV, 6 leaves were harvested to test the expression level

of *TiAP1* and *BgtCDA1* gene, while the other 6 leaves were inoculated with *Bgt* race

E09 and were observed and used to evaluate the resistance to the powdery mildew

using a biological microscope (Nikon Ni-U, Japan) after the Coomassie Brilliant Blue

staining and the statistics of hyphal density score refer to the method of Norriss et al.

(2007) and three biological replicates.

BgtCDA1 gene silencing in *Bgt* was performed using the HIGS method as

described by Nowara *et al*. (2010). The RNA interference (RNAi) construct BgtCDA1

RNAi and a β-glucuronidase (GUS) reporter gene construct were co-transformed into

the leaf epidermal cells of the wheat YN15 that grew for seven days using particle

bombardment. The fungal haustorial formation was examined for the transformed

(GUS expressing) blue cells using a biological microscope (Nicon Ni-U, Japan). The

empty vectors pIPKTA30N, and the Mlo-RNAi (pIPKTA36) constructs, were used as

negative and positive controls, respectively and four biological replicates were

conducted per analysis with three technical replicates. The detailed method was

described in Methods S1 according to Ahmed (2015).

Constructing the TiAP1 prokaryotic expression plasmid and the protein

expression

To improve the expression of *TiAP1*, the gene sequence was optimised according to *E.*

coli bias, and constructed into the pET28a vector (containing 6xHis tags).

Recombinant plasmids were transformed into BL21 (DE3) chemically competent

cells where a single-colony transformant was cultured and inoculated at 15 °C

overnight induced expression with 0.25 mM isopropyl β- d-1-thiogalactopyranoside

(IPTG). The cells were harvested and subjected to a sodium dodecyl

sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) to detect the protein

expression. Then, the recombinated TiAP1 was purified according to the method

described in Methods S1.

Determination of the antifungal activity of TiAP1

To confirm the effect of TiAP1 protein on *Bgt*, we used the renatured TiAP1 and

pET28a tag protein to add 0.025% Tween-20 to smear the leaves of YN15 and

inoculate with *E09*. After five days, the infectivity of *Bgt* was observed and the leaves

were stained with the Coomassie Brilliant Blue stain, as described previously (Göllner

et al., 2008) and three biological replicates were conducted per analysis with three

technical replicates. Lesion areas on the leaf surface were calculated according to

Goodwin and Hsiang (2010) to analyse the pixels of the *Bgt* infection area relative to

the normal leaf area.

Constructing the particle bombardment genetic transformation vector to transform wheat and identify the disease resistance phenotype

The coding sequence (CDS) of *TiAP1* (KJ513672) was cloned into the *Ubi-gene*-Tnos

vector with a ubiquitin (Ubi) promoter, yielding a recombinant pMUbi-*TiAP1* vector.

The resulting plasmid pMUbi-*TiAP1* and the vector harboring the *Bar* gene were

mixed and co-bombarded with a particle bombardment of 1100 psi, 1 μg per gun, and

60 μg gold powder into the callus that was induced from the mature embryos of the

Bobwhite wheat, which was highly susceptible to *Bgt*. During the induction and

regeneration, the transformed wheat was screened using the herbicide Basta, while the

presence of the transgenes was determined by amplifying the target gene using the

pUbi-ASPF/R primers (Table S6).

Next, the disease responses to powdery mildew of the transgenic plants were tested.

When the Bobwhite and the transgenic Bobwhite with the *TiAP1* gene grew to the

three-leaf stage, the first leaves of them were spread out on an acrylic plate with the

conidium of the *Bgt* race *E09* sprayed evenly. The leaves were harvested at 48 h

post-inoculation with *E09* and stained with Coomassie Brilliant Blue as described

previously (Göllner *et al*., 2008) and three biological replicates were conducted per

stains with three technical replicates. Furthermore, to evaluate the resistance to *Bgt* of

the Bobwhite and the transgenic Bobwhite with the *TiAP1* gene in the adult stage,

they were natural infected by powdery mildew, where the infection type (IT) of each

plant was scored as described by Lu *et al*. (2020).

DNA and RNA extraction and qRT-PCR

Total DNA was extracted from the plant leaves using the cetyltrimethylammonium

bromide (CTAB) method (Allen *et al*., 2006). While total RNA (500 ng) was

extracted using the EasyPure® Plant RNA kit (TransGen Biotech Co., Ltd, Beijing,

China), reverse-transcribed (RT) using a RevertAid First Strand cDNA synthesis kit

(TransGen Biotech Co. Ltd, Beijing, China), and quantified on an ABI Quantitative

PCR Q6 Detection System (Thermo Fisher Scientific, USA) with the SYBR Premix

Ex Taq kit (TransGene Biotech Co., Ltd, Beijing, China). The wheat actin gene was

used as the reference gene, in which three independent biological replicates were

conducted per analysis with at least three technical replicates. The primers have been

listed in Table S6.

Subcellular localisation

The CDS of the *TiAP1* fragments was fused green fluorescent protein (GFP) in the

vector pCAMBIA1300. Then the constructed TiAP1-GFP or Δsp-TiAP1-GFP (with or

without signal peptide of TiAP1 respectively) were transferred into the *Agrobacterium*

tumefaciens GV3101 and transiently expressed in *N. benthamiana* leaves. Two days

after infiltration, they were observed using a confocal laser scanning microscope

(Leica SP5-X, Germany), Ten *N. benthamiana* leaves were analyzed in each of the three experiments and the detailed method was described in Methods S1.

Transient expression of the TiAP1 protein induced by *Bgh* using barley leaves

The expression vector of the TiAP1 protein was constructed using the gateway system (Thermo Fisher Scientific, USA). The two recombinant plasmids, TiAP1-mYFP and mCherry-TaSYP51, were co-bombarded onto the leaves of the 7-day-old barley by particle bombardment at a helium pressure of 1,100 psi and placed on the 1% Phytagel^{TM} petri dishes containing 40mg ml^{-1} of benzimidazole (Sigma-Aldrich, USA) to culture for 24 h, and then inoculated with *Bgh* spores. After additional 24 h of inoculation, the expression of TiAP1 and TaSYP51 proteins were observed and analysed using a confocal laser-scanning microscope (Leica SP5‑X, Germany) according to the methodology of Smigielski *et al.* (2019). And three biological replicates were conducted per co-bombarded with six leaves.

Protein Interaction Assay We sampled the leaves of YN15 and SN6306, which were infected with *Bgt* from 0 to 5 d, and then construct a primary cDNA library using the BP reaction of the CloneMiner II kit (Invitrogen, Thermo Fisher Scientific, USA) by Qingdao Oebiotech Co. Ltd. (China). The quality of the cDNA library was identified. Next, we used the nuclear system Y2H mating standard operating procedure method as described in the manufacturer's instructions (Clontech Laboratories, Inc., USA) to screen the protein that interacts with TiAP1 and further verified the obtained interaction protein, and three biological replicates were conducted per Y2H verify. The structure, amino acid sequence domain, and the signal peptide of the obtained protein BgCDA1 were analysed using the the Basic Local Alignment Search Tool (PBLAST; Altschul *et al*., 1997), Pfam (Finn *et al*., 2014), and SignalP 5.0 (Almagro Armenteros *et al*., 2019) software, respectively.

For LCI assay, by amplifying the sequence of TiAP1 and BgtCDA1 and applying homologous recombination with the nLuc and cLuc, as previously described (Chen *et al*., 2008), the fusion protein and one of the corresponding co-injected empty vectors,

nLuc/cLuc, was co-expressed in *N. benthamiana* as a negative control. At 48 h

post-infiltration, the LUC activity of the leaves was observed using an *in vivo* imaging

system (Berthold LB985 NightSHADE, Germany). Ten *N. benthamiana* leaves were

analyzed in each of the three experiments.

For BiFC assay, the sequences of TiAP1 and BgtCDA1 were amplified and fused

with the N and C terminals of the GFP, respectively, using the gateway system

(Thermo Fisher Scientific, USA). As mentioned previously (Bracha-Drori *et al*., 2004),

the fusion protein and one of the corresponding empty GFP-N/GFP-C vectors were

co-expressed in *N. benthamiana* as a negative control. The GFP signal was observed

48 h after infiltration using a confocal laser-scanning microscope (Leica SP5-X,

Germany). Ten *N. benthamiana* leaves were analyzed in each of the three

experiments.

The yeast expression, purification and chitin deacetylase activity assay of

BgtCDA1

For protein expression and purification in yeast, BgtCDA1 coding sequences with a

6XHIS tag were amplified and ligated into pPIC9K-HIS vector (Invitrogen), and then

transformated into *Pichia pastoris* strain GS115 (Invitrogen). The positive

transformants were induced protein expression using methanol. The target proteins

were purified using Ni-column affinity chromatography. Purified protein was used to

detected the enzyme activity of BgtCDA1 by measuring the amount of released

acetate using ion chromatography and qualitative analysis of deacetylated products by

MALDI-TOF by Hoogen Biotech, Shanghai, China according to Gao et al., (2019).

MALDI-TOF MS analysis was performed on a Ultraflex Extreme MALDI-TOF-TOF

mass spectrometer (Bruker Daltonics, USA). The standard reaction mixture (200 µl)

containing 2 µM BgtCDA1, 50 mM Tris-HCl (pH8.0) and 1 mM chitin oligomers

with 6 GlcNAc moieties (A6) as the substrate were incubated at 37 ℃ for 5, 30 min

followed by heating at 100 ℃ for 10 min. The mixture without BgtCDA1 was as the

control. The yeast expression, purification and enzyme activity assay of BgtCDA1

were detailed performed described in Methods S1.

RNA-Seq and data analysisRNA-Seq analysis was performed using the two-leaf

stage Bobwhite (Bob-0) and the transgenic Bobwhite with the *TiAP1* gene (OE2-0) as

well as a Bobwhite inoculated with *E09* two days later (Bob-2) and the corresponding

transgenic Bobwhite with the *TiAP1* gene (OE2-2). Three biological replicates were

used for the RNA extraction, where total RNA was extracted and treated with DNase I,

in which the quality test was performed, and then used to construct the library. The

quality of the library was assessed using an Agilent 2100 Bioanalyzer (Agilent

Technologies, Inc., USA), while the libraries were sequenced on an Illumina HiSeq

2500 platform (Illumina, Inc., USA). The RNA-Seq reads were aligned to the wheat

genome of the International Wheat Genome Sequencing Consortium with the RefSeq

v1.1 annotation (https://wheat-urgi.versailles.inra.fr/Seq-Repository/Annotations).

Differentially expressed genes across the samples were identified using the DESeq2

package (Love *et al*., 2014) using the standard parameters with an adjusted

false-discovery rate of *P*-value < 0.01 and fold change > 4.

Statistical analysis

The statistical significance of the results was calculated using a one-way analysis of

variance followed by least significant difference and Duncan's new multiple range test

at a significant difference ($P < 0.05$) using IBM SPSS software version 19.0 (Chicago,

Illinois, USA). All experiments reported include three to six biological replicates and

a minimum of three technical replicates with similar results. All primers have been

listed in Table S6.

Accession numbers

613 The cDNA and coded protein sequence of *TiAP1* sequence in this research has been

614 deposited in NCBI under the accession number KJ513672 and AJC64141.1; *Blumeria*

615 *graminis* f. sp. *tritici* BgtCDA1 was coded_by =" join (KE374986.1:122515..122759,

616 KE374986.1:122809..123142,   KE374986.1:123188..123296,

617 KE374986.1:123342..123381,   KE374986.1:123432..123502,

KE374986.1:123547..123801, KE374986.1:123919..123923, KE374986.1:123978..124142, KE374986.1:124319..124375), and had one and two additional alternative splicings compared with EPQ66796 between KE374986.1:123143 and KE374986.1:123187, KE374986.1:123802 and KE374986.1:123977, respectively; Other protein sequence data from this article can be found in the NCBI under the following accession numbers: *Saccharomyces cerevisiae CDA1 (NP_013410) and Saccharomyces cerevisiae CDA2 (NP_013411).* TaSYP51 was coded by sequence in chr5A of wheat between 67321475 and 67319313.

Acknowledgements

This work was supported by the grants from the National Natural Science Foundation of China (31771777); the Key R & D program of Shandong Province (Public Welfare Special) (2018GSF121008); Shandong 'Double Tops' Program; the Major Basic Research Project of Shandong Natural Science Foundation (2017C03); and the Overseas Visiting Programme for Graduate Mentors of Shandong Province. We are very grateful to Prof. Hans Thordal-Christensen's group, University of Copenhagen, Denmark, for providing experimental assistance, to Dr. Patrick Schweizer, IPK, Gatersleben, Germany, for supplying the RNAi vector and the Mlo-RNAi construct, to Prof. Pinghua Li, Shandong Agricultural University, for providing the LCI system, and to Prof. Dawei Li, China Agriculture University, for providing the BSMV-RNAi system. Confocal image data were collected at the Center for Advanced Bioimaging (CAB), University of Copenhagen, Denmark.

Author contributions

D.F., H.W., and Y.Y. conceived and designed the experiments. Y.Y., P.F., J.L., W.X., N.L., Z.N., Q.L., J.S., Q.T., and Y.B. performed the experiments and analysed the data; Y.Y. and D.F. wrote the paper; D.F. and H.W. conceived, directed and coordinated the project.

Competing interests

The authors have no competing interests to declare.

References

Ahmed AA, Pedersen C, Schultz-Larsen T, Kwaaitaal M, Jørgensen HJL,
Thordal-Christensen H (2015) The barley powdery mildew candidate secreted
effector protein CSEP0105 inhibits the chaperone activity of a small heat shock
protein. *Plant Physiology* **168**: 321-333.
Alam MM, Nakamura H, Ichikawa H, Miyao A, Hirochika H, Kobayashi K, Yamaoka
N, Nishiguchi M (2014) Response of anasparctic protease gene *OsAP77* to fungal,
bacterial and viral infections in rice. *Rice* **7**: 1-14.
Allen GC, Flores-Vergara MA, Krasynanski S, Kumar S, Thompson WF (2006) A
modified protocol for rapid DNA isolation from plant tissues using
cetyltrimethylammonium bromide. *Nature Protocols* **1**: 2320-2325.
Almagro Armenteros JJ, Tsirigos KD, Sønderby CK, Petersen TN, Winther O, Brunak
S, von Heijne G, Nielsen H (2019) SignalP 5.0 improves signal peptide
predictions using deep neural networks. *Nature Biotechnol* **37**: 420-423.
Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ
(1997) Gapped BLAST and PSI-BLAST: a new generation of protein database
search programs. *Nucleic Acids Research* **25**: 3389-3402.
Bracha-Drori K, Shichrur K, Katz A, Oliva M, Angelovici R, Yalovsky S, Ohad N
(2004) Detection of protein–protein interactions in plants using bimolecular
fluorescence complementation. *Plant Journal* **40**: 419-427.
Bracket CE (1968) The ultrastructure of the haustorial apparatus of *Erysiphe graminis*
and its relationship to the epidermal cell of barley. *Phytopathology* **58**: 12-30.
Bradford MM (1976) A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. *Analytical*
Biochemistry **72**: 248-254.
Chen F, Foolad MR (1997) Molecular organization of a gene in barley which encodes
a protein similar to aspartic protease and its specific expression in nucellar cells
during degeneration. *Plant Molecular Biology* **35**: 821-831.
Chen H, Zou Y, Shang Y, Lin H, Wang Y, Cai R, Tang X, Zhou JM (2008) Firefly
luciferase complementation imaging assay for protein-protein interactions in
plants. *Plant Physiology* **146**: 368-376.
Chen HJ, Huang YH, Huang GJ, Huang SS, Chow TJ, Lin YH (2015) Sweet potato
SPAP1 is a typical aspartic protease and participates in ethephon-mediated leaf
senescence. *Journal of Plant Physiology* **180**: 1-17.
Chen P, You C, Hu Y, Chen S, Zhou B, Cao A, Wang X (2013) Radiation-induced
translocations with reduced *Haynaldia villosa* chromatin at the *Pm21* locus for
powdery mildew resistance in wheat. *Molecular Breeding* **31**: 477-484.
Chung HS, Howe GA (2009) A critical role for the TIFY motif in repression of
jasmonate signaling by a stabilized splice variant of the JASMONATE
ZIM-domain protein JAZ10 in Arabidopsis. *The Plant Cell* **21**: 131-145.
Collins NC, Thordal-Christensen H, Lipka V, Bau S, Kombrink E, Qiu JL,

Hückelhoven R, Stein M, Freialdenhoven A, Somerville SC *et al.* (2003) SNARE-protein-mediated disease resistance at the plant cell wall. *Nature* **425**: 973-977.

Deising HB, Werner S, Wernitz M (2000) The role of fungal appressoria in plant infection. *Microbes and Infection* **2**: 1631-1641.

Díaz ME, Rocha GF, Kise F, Rosso AM, Guevara MG, Parisi MG (2018) Antimicrobial activity of an aspartic protease from *Salpichroa origanifolia* fruits. *Letters in Applied Microbiology* **67**: 168-174.

Douchkov D, Nowara D, Zierold U, Schweizer P (2005) A high-throughput gene-silencing system for the functional assessment of defense-related genes in barley epidermal cells. *Molecular Plant-Microbe Interactions* **18**: 755-761.

Gao F, Zhang BS, Zhao JH, Huang JF, Jia PS, Wang S, Zhang J, Zhou JM, Guo HS (2019) Deacetylation of chitin oligomers increases virulence in soil-borne fungal pathogens. *Nature Plants* **5**: 1167-1176.

Geoghegan IA, Gurr SJ (2017) Investigating chitin deacetylation and chitosan hydrolysis during vegetative growth in *Magnaporthe oryzae. Cell Microbiology* **19**: e12743.

Geziel BA (2015) *On the mechanism of action of selected effector candidates from the barley powdery mildew fungus.* PhD thesis, University of Copenhagen, Copenhagen, Denmark.

Goodwin PH, Hsiang T (2010) Quantification of fungal infection of leaves with digital images and scion image software. In: Sharon A, ed. *Molecular and Cell Biology Methods for Fungi, Methods in Molecular Biology.* Springer Science+Business Media, LLC **638:**125-134.

Göllner K, Schweizer P, Bai Y, Panstruga R (2008) Natural genetic resources of *Arabidopsis thaliana* reveal a high prevalence and unexpected phenotypic plasticity of RPW8-mediated powdery mildew resistance. *New Phytologist* **177**: 725-742.

Griffey C, Das M (1994) Inheritance of adult-plant resistance to powdery mildew in Knox62 and Massey winter wheats. *Crop Science* **34**: 641-646.

Guo R, Xu X, Carole B, Li X, Gao M, Zheng Y, Wang X (2013) Genome-wide identification, evolutionary and expression analysis of the aspartic protease gene superfamily in grape. *BMC Genomics* **14**: 554.

Guo R, Tu M, Wang X, Zhao J, Wan R, Li Z, Wang Y, Wang X (2016) Ectopic expression of a grape aspartic protease gene, *AP13*, in *Arabidopsis thaliana* improves resistance to powdery mildew but increases susceptibility to *Botrytis cinerea. Plant Science* **248**: 17-27.

Han Y, Song L, Peng C, Liu X, Liu L, Zhang Y, Wang W, Zhou J, Wang S, Ebbole D *et al.* (2019) A *Magnaporthe* chitinase interacts with a rice jacalin-related lectin to promote host colonization. *Plant Physiology* **179**: 1416-1430.

Hautea RA, Coffman WR, Sorrells ME, Bergstrom GC (1987) Inheritance of partial resistance to powdery mildew in spring wheat. *Theoretical and Applied Genetics* **73**: 609-615.

He R, Chang Z, Yang Z, Yuan Z, Zhan H, Zhang X, Liu J (2009) Inheritance and

mapping of powdery mildew resistance gene *Pm43* introgressed from *Thinopyrum intermedium* into wheat. *Theoretical and Applied Genetics* **118**: 1173-1180.

He J, Liu Y, Yuan D, Duan M, Liu Y, Shen Z, Yang C, Qiu Z, Liu D, Wen P *et al*. (2020) An R2R3 MYB transcription factor confers brown planthopper resistance by regulating the phenylalanine ammonia-lyase pathway in rice. *Proceedings of the National Academy of Sciences of the United States of America* **117**: 271-277.

Hurlburt NK, Chen LH, Stergiopoulos I, Fisher AJ (2018) Structure of the *Cladosporium fulvum* Avr4 effector in complex with (GlcNAc)$_6$ reveals the ligand-binding mechanism and uncouples its intrinsic function from recognition by the Cf-4 resistance protein. *PLoS Pathogens* **14**: e1007263.

Jashni MK, Mehrabi R, Collemare J, Mesarich CH, de Wit PJGM (2015) The battle in the apoplast: further insights into the roles of proteases and their inhibitors in plant-pathogen interactions. *Frontiers in Plant Science* **6:** 584.

Kamakura T, Yamaguchi S, Saitoh KI, Teraoka T, Yamaguchi I (2002) A novel gene, CBP1, encoding a putative extracellular chitin-binding protein, may play an important role in the hydrophobic surface sensing of *Magnaporthe grisea* during appressorium differentiation. *Molecular Plant-Microbe Intertations* **15**: 437-444.

Kuroki M, Okauchi K, Yoshida S, Ohno Y, Murata S, Nakajima Y, Nozaka A, Tanaka N, Nakajima M, Taguchi H *et al*. (2017) Chitin-deacetylase activity induces appressorium differentiation in the rice blast fungus *Magnaporthe oryzae*. *Scientific Reports* **7**: 9697.

Li H, Dong Z, Ma C, Xia Q, Tian X, Sehgal S, Koo DH, Friebe B, Ma P, Liu W (2020a) A spontaneous wheat-Aegilops longissima translocation carrying Pm66 confers resistance to powdery mildew. *Theoretical and Applied Genetics* **133**: 1149-1159.

Li M, Dong L, Li B, Wang Z, Xie J, Qiu D, Li Y, Shi W, Yang L, Wu Q, *et al*. (2020b) A CNL protein in wild emmer wheat confers powdery mildew resistance. *New Phytologist* **228**: 1027-1037.

Li Q, Niu Z, Bao Y, Tian Q, Wang H, Kong L, Feng D (2016) Transcriptome analysis of genes related to resistance against powdery mildew in wheat-Thinopyrum alien addition disomic line germplasm SN6306. *Gene* **590**: 5-17.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biology* **15**: 550.

Lu P, Guo L, Wang Z, Li B, Li J, Li Y, Qiu D, Shi W, Yang L, Wang N *et al*. (2020) A rare gain of function mutation in a wheat tandem kinase confers resistance to powdery mildew. *Nature Communications* **11**: 680.

Luo P, Luo H, Chang Z, Zhan H, Zhang M, Ren Z (2009) Characterization and chromosomal location of *Pm40* in common wheat: a new gene for resistance to powdery mildew derived from *Elytrigia intermedium*. *Theoretical and Applied Genetics* **118**:1059-1064.

Nakamura Y, Kato T, Yamashino T, Murakami M, Mizuno T (2007) Characterization of a set of phytochrome-interacting factor-like bHLH proteins in *Oryza sativa*. *Bioscience, Biotechnology, and Biochemistry* **71**:1183-1191.

Niu N, Liang W, Yang X, Jin W, Wilson ZA, Hu J, Zhang D (2013) EAT1 promotes
tapetal cell death by regulating aspartic proteases during male reproductive
development in rice. *Nature Communications* **4**: 1445.
Norriss MG, Frost-Smith TJ, Sutherland JIM (2007) Increasing endophyte alkaloid
expression in tall fescue by selecting for increased endophyte hyphal density.
Proceedings of the New Zealand Grassland Association **69**: 197-200.
Nowara D, Gay A, Lacomme C, Shaw J, Ridout C, Douchkov D, Hensel G, Kumlehn
J, Schweizer P (2010) HIGS: host-induced gene silencing in the obligate
biotrophic fungal pathogen Blumeria graminis. *Plant Cell* **22**: 3130-41.
Panstruga R (2013) Establishing compatibility between plants and obligate biotrophic
pathogens. *Current Opinion in Plant Biology* **6**: 320-326.
Park SY, Fung P, Nishimura N, Jensen DR, Fujii H, Zhao Y, Lumba S, Santiago J,
Rodrigues A, Chow TF, Alfred SE, Bonetta D, Finkelstein R, Provart NJ,
Desveaux D, Rodriguez PL, McCourt P, Zhu JK, Schroeder JI, Volkman BF,
Cutler SR (2009) Abscisic acid inhibits type 2C protein phosphatases via the
PYR/PYL family of START proteins. *Science* **324**:1068-1071.
Prasad BD, Creissen G, Lamb C, Chattool BB (2009) Overexpression of rice (*Oryza*
sativa L.) OsCDR1 leads to constitutive activation of defense responses in rice
and Arabidopsis. *Molecular Plant-Microbe Intertations* **22**: 1635-1644.
Prasad BD, Creissen G, Lamb C, Chattoo BB (2010) Heterologous expression and
characterization of recombinant OsCDR1, a rice aspartic proteinase involved in
disease resistance. *Protein Expression and Purification* **72**: 169-174.
Rawlings ND, Barrett AJ (1995) Families of aspartic peptidases, and those of
unknown catalytic mechanism. *Methods in Enzymology* **248**: 105-120.
Rawlings ND, Barrett AJ, Finn R (2016) Twenty years of the MEROPS database of
proteolytic enzymes, their substrates and inhibitors. *Nucleic Acids Research* **44**:
343-350.
Rushton DL, Tripathi P, Rabara RC, Lin J, Ringler P, Boken AK, Langum TJ, Smidt L,
Boomsma DD, Emme NJ, Chen X, Finer JJ, Shen QJ, Rushton PJ (2012) WRKY
transcription factors: key components in abscisic acid signalling. *Plant*
Biotechnology Journal **10**: 2-11.
Shaner G (1973) Evaluation of slow-mildewing resistance of Knox wheat in the field.
Phytopathology **63**: 867-872.
Shen XK, Ma LX, Zhong SF, Liu N, Zhang M, Chen WQ, Zhou YL, Li HJ, Chang ZJ,
Li X, *et al*. (2015) Identification and genetic mapping of the putative
Thinopyrum intermedium‐derived dominant powdery mildew resistance gene
PmL962 on wheat chromosome arm 2BS. *Theoretical and Applied Genetics*
128:517-528.
Singh RP, Singh PK, Rutkoski J, Hodson DP, He X, Jørgensen LN, Hovmøller MS,
Huerta-Espino J (2016) Disease impact on wheat yield potential and prospects of
genetic control. *Annual Review of Phytopathology* **54**: 303-322.
Smigielski L, Aguilar GB, Kwaaitaal M, Zhang WJ, Thordal-Christensen H (2019)
The isoelectric point of proteins influences their translocation to the
extrahaustorial matrix of the barley powdery mildew fungus. *Cellular*
Microbiology **21**: e13091.
Tian QJ, Song J, Yang YL, Xu T, Wang HG, Feng DS (2017) Cloning and

Bioinformatics of *TiAP1* Gene from *Thinopyrum intermedium*. *Shandong Agricultural Science* **49**: 1-9.

Tsigos I, Bouriotis V (1995) Purification and characterization of chitin deacetylase from *Colletotrichum lindemuthianum*. *Journal of Biological Chemistry* **270**: 26286-26291.

Upadhya R, Baker LG, Lam WC, Specht CA, Donlin MJ, Lodge JK (2018) *Cryptococcus neoformans* Cda1 and its chitin deacetylase activity are required for fungal pathogenesis. *mBio* **9**: e02087-18.

Wang Y, Wang Y, Wang Y (2020) Apoplastic proteases: powerful weapons against pathogen onfection in plants. *Plant Communication* 1, 100085.

Xia YJ, Suzuki H, Borevitz1 J, Blount J, Guo ZJ, Patel K, Dixon RA, Lamb C (2004) An extracellular aspartic protease functions in Arabidopsis disease resistance signaling. *EMBO Journal* **23**: 980-988.

Xia YQ, Ma ZC, Qiu M, Guo BD, Zhang Q, Jiang HB, Zhang BY, Lin YC, Xuan MR, Sun L, *et al*. (2020) N-glycosylation shields *Phytophthora sojae* apoplastic effector PsXEG1 from a specific host aspartic protease. *Proceedings of the National Academy of Sciences of the United States of America* **117**: 27685-27693.

Xu ZS, Chen M, Li LC, Ma YZ (2008) Functions of the ERF transcription factor family in plants. *Botany* **86**: 969-977.

Yang C, Yu Y, Huang J, Meng F, Pang J, Zhao Q, Islam MA, Xu N, Tian Y, Liu J (2019) Binding of the *Magnaporthe oryzae* chitinase MoChia1 by a rice tetratricopeptide repeat protein allows free chitin to trigger immune responses. *Plant Cell* **31**: 172-188.

Yang YL, Feng DS (2020) Genome-Wide Identification of the Aspartic Protease Gene Family and Their Response under Powdery Mildew Stress in Wheat. Mol Biol Rep **47**: 8949-8961.

You XM, Fang H, Wang RY, Wang GL, Ning YS (2020) Phenylalanine ammonia lyases mediate broad-spectrum resistance to pathogens and insect pests in plants. *Science Bulletin* **65**: 1425-1427.

Yuan C, Li C, Yan L, Jackson AO, Liu Z, Han C, Yu J, Li D (2011) A high throughput barley stripe mosaic virus vector for virus induced gene silencing in monocots and dicots. *PLoS One* **6**: e26468.

Zhang L, Gu L, Ringler P, Smith S, Rushton PJ, Shen QJ (2015) Three WRKY transcription factors additively repress abscisic acid and gibberellin signaling in aleurone cells. *Plant Science*, **236**: 214-222.

Zhang Z, Henderson C, Perfect E, Carver TLW, Thomas BJ, Skamnioti P, Gurr SJ (2005) Of genes and genomes, needles and haystacks: *Blumeria graminis* and functionality. *Molecular Plant Pathology* **6**: 561-575.

Zhai K, Deng Y, Liang D, Tang J, Yan B, Yin X, Lin H, Chen F, Yang D, Xie Z *et al*. (2019) RRM transcription factors interact with NLRs and regulate broad-spectrum blast resistance in rice. *Molecular Cell* **74**: 996-1009.

Zhao Y, Park RD, Muzzarelli RA (2010) Chitin Deacetylases: Properties and Applications. *Marine Drugs* **8**: 24-46.

Zhou S, Miao QX, Huang LL, Chen LH (2017) Research progress in

hemicellulose-based functional materials. *Chemistry and Industry of Forest*
*Product*s **37**: 10-18.
Zhou X, Liao H, Chern M, Yin J, Chen Y, Wang J, Zhu X, Chen Z, Yuan C, Zhao W *et*
al. (2018) Loss of function of a rice TPR-domain RNA-binding protein confers
broad-spectrum disease resistance. *Proceedings of the National Academy of*
Sciences of the United States of America **115**: 3174-3179.

873
874

FIGURE LEGENDS

Fig. 1 qRT-PCR and VIGS analysis of the *TiAP1* effect on the *Bgt* resistance, the

TiAP1 prokaryotic expression and purification, and the reduction of the powdery

mildew severity (lesions area) on the wheat seedlings by a recombination protein

treatment. (A) Expression levels of *TiAP1* in YN15 and SN6306 that were

induced by *Bgt*. Total RNA from the leaves of YN15 and SN6306 that were

induced by *Bgt* at 0, 12, 24, 48, 72, and 96 h were extracted as the template to be

detected by qRT-PCR. Data represent mean ± S.D. of n = 3 technical replicates.

(B) Relative quantification of expression levels of *TiAP1*. Total RNA from the

leaves of barley stripe mosaic virus (BSMV):00 and BSMV: TiAP1as inoculated

with *E09* for ten days were extracted as the template for the detection by

qRT-PCR. (C) Disease phenotype of the BSMV:00 and BSMV: TiAP1as

inoculated with *E09*. Photographs were taken at 10 dpi. (D) Microscopic

observation of the disease symptoms of BSMV:00 and BSMV: TiAP1as 10 dpi.

Bar = 100 µm. (E) The TiAP1 protein expression detection by western blotting

before and after induction. M: Pre-stained protein ladder; Lane1: Supernatant of

non-induced; Lane 2: Subside of non-induced; Lane 3: Supernatant of induced;

Lane 4: Subside of induced. (F) SDS-PAGE analysis of the inclusion body

purification. M: Low-molecular weight protein ladder; Lane 1: Inclusions after

dissolving the centrifugal supernatant; Lane 2: The effluent of supernatant was

incubated with nickel iminodiacetic acid (Ni-IDA); Lane 3-6: The effluent of the

imidazole elution of 50 mM; Lane 7-11: The effluent of the imidazole elution of

100 mM; Lane 12-13: The effluent of the imidazole elution of 300 mM; (G)

TiAP1 quality assurance after renaturation. M: Pre-stained protein ladder; Lane 1:

TiAP1 protein (1.5 µg). (H) Disease phenotype of YN15 smeared with TiAP1 and the tag protein from pET28a (CK) when inoculated with *E09* for five days. (I) Microscopic (Nikon Ni-U, Japan) analysis of the disease symptoms of YN15 smeared with TiAP1 and CK when inoculated with *E09* for five days. Bar = 100 µm. (J) The proportion of lesions in the leaf area of YN15 that was pre-treated with the recombinant TiAP1 five days before *Bgt E09* inoculation relative to the control. Control: Pre-treated with the purified recombinant label protein from the pET28a construct. Different letters above the bars indicate statistically significant differences (P <0.05) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method.

Fig. 2 Identifying the resistance of transgenic *TiAP1* wheat. (A) Expression levels of *TiAP1* in Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for two days. (B) Disease phenotype of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for five days. (C) Microscopic (Nikon Ni-U, Japan) analysis of the disease symptoms of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for two days. Bar = 100 µm. (D) Count of *Bgt* conidia in different stages of the *Bgt* when inoculated with *E09* for two days in Bobwhite and the transgenic Bobwhite with *TiAP1*. (E) The HI statistics of the *Bgt* when inoculated with *E09* for two days in Bobwhite and the transgenic Bobwhite with *TiAP1*. Different letters above the bars indicate statistically significant differences (P <0.05) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method. (F) Identifying the resistance of different wheat at the adult stage natural infected by powdery mildew in the glasshouse. (1) Bobwhite; (2) The transgenic Bobwhite with *TiAP1*; (3) YN15; (4) F_1 obtained from cross breeding between YN15 and homozygous transgenic *TiAP1* Bobwhite.

Fig. 3 Subcellular localisation of the fusion protein through transient expression of agroinfiltrated GFP, TiAP-GFP, and Δsp-TiAP-GFP in the epidermal cells of *N. benthamiana* after plasmolysis, and co-transient expression of TiAP1-mYFP and mCherry-TaSYP51 in the epidermal cells of barley through bombardment. (A)

Confocal laser scanning microscopy (Leica SP5-X, Germany) observation of the

leaf epidermal cells revealed that Δsp-TiAP-GFP appears to be localised in the

cytoplasm, while TiAP1-GFP is predominantly localised in the extracellular

region, where white arrows indicate plasmolysis and the yellow dotted line

shows the location of the cell membrane of the two adjacent cells. A GFP signal

was used as the control. Bar = 20 μm. (B) The enlargement of the blue box in

TiAP1-GFP. (C) The enlargement of the blue box in Δsp-TiAP1-GFP. （D）*B.*

graminis f. sp. *hordei* induced the accumulation of TiAP1-mYFP and

mCherry-TaSYP51 at the infection site. This figure is part of Video S1. Bar = 20

μm.

Fig. 4 Interaction *in vitro* and *in vivo* of TiAP1 with BgtCDA1. (A) Yeast two-hybrid

analysis of the interaction between TiAP1 and BgtCDA1. Yeasts expressing the

interacting combinations of bait and prey were spotted on the

SD-HAWL/X-α-Gal/ABA. Bait-53 and AD-T (prey) were the positive controls,

while Bait-Lam and AD-T as well as Bait-BD and AD-BgtCDA1 were used as

the negative controls. (B) Luciferase complementation imaging assay showing

that TiAP1 interacted with and BgtCDA1. The ΔspTiAP1-nLuc and

cLuc-ΔspBgtCDA1 were transiently co-expressed in *N. benthamiana.*

Co-infiltration of cLuc/nLuc-tagged ΔspTiAP1 protein with the

nLuc/cLuc-tagged ΔspBgtCDA1 or the empty vectors were used as negative

controls, where the images were observed using chemiluminescence imaging two

days later. (C) Bimolecular florescent complimentary assay showing TiAP1

interaction with BgtCDA1. The cGFP-ΔspTiAP1 and nGFP-ΔspBgtCDA1 were

transiently co-expressed in *N. benthamiana.* Co-infiltrations of

nGFP/cGFP-ΔspTiAP1, cGFP/nGFP- ΔspBgtCDA1, or the empty vectors group

nGFP/cGFP were used as negative controls. The positive controls were

nYFP-14-3-3/14-3-3-cCFP. The green florescent protein (GFP) signal was

visualised using confocal microscopy (Leica SP5-X, Germany). Bar = 20 μm.

Fig. 5 The silencing of *BgtCDA1* reduced *Bgt* infection and the BgtCDA1 mediated

deacetylation of chitohexaose (A6). (A) Wheat YN15 leaves, bombarded with RNAi and GUS reporter constructs, were infected with *Bgt* and calculated for fungal haustorial index. The haustorial index was calculated as the ratio of haustoria-containing transformed cells (GUS expressing cells) divided by the total number of transformed cells. Data shown are mean values, $n = 4$, and a minimum of 150 transformed cells were counted in each repeat. Relative haustorial index was computed relative to the empty vector pIPKTA30N control of each experiment, which was set to 100%. The Mlo-RNAi was as another control. Different letters above the bars indicate statistically significant differences ($P <0.05$) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method. Arrow indicates haustorium. Bar = 20 µm. (B) Wheat YN15 leaves, bombarded with RNAi and GUS reporter constructs, were infected with *Bgt*. Arrow indicates haustoria. Bar = 20 µm. (C) Disease phenotype of the BSMV:00 and BSMV: BgtCDA1as inoculated with *E09*. Photographs were taken at 5 dpi. (D) Microscopic observation of the disease symptoms of BSMV:00 and BSMV: BgtCDA1as 3 dpi using *E09*. Bar = 100 µm. (E) The proportion of lesions in the leaf area of BSMV:00 and BSMV: BgtCDA1as inoculated with *E09* five days. (F) The hyphal density score of leaf material in BSMV:00 and BSMV: BgtCDA1as 3 dpi using *E09* in microscopic observation field plots. Different letters above the bars indicate statistically significant differences ($P <0.05$) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method. (G) Purified BgtCDA1 (2 uM) was incubated with 1 mM A6 for 5, 30 min as indicated. The partially deacetylated product A5D1 was detected. A, N-acetyl-D-glucosamine; D, D-glucosamine.

Fig. 6 The transcriptome and qRT-PCR analysis of the transgenic wheat with *TiAP1*. (A) Heat map of selected pathogen-responsive genes in Bobwhite without and with induction of *Bgt* for two days (Bob-0 and Bob-2, respectively), and overexpression line OE2 as the same treatment (OE2-0 and OE2-2, respectively).

(B) qRT-PCR analysis of the differential expression gene *TraesCS1B02G440300*,

TraesCS2A02G161500, *TraesCS2A02G199300*, *TraesCS3B02G379200*,

TraesCS5D02G188600, *TraesCS5A02G049600*, *TraesCS3B02G259500*,

TraesCS2B02G224300, *TraesCS6A02G266100*, *TraesCS2D02G099900*,

TraesCS6B02G018700 and *TraesCS2B02G529400* of selected induced of *Bgt* for

two days.

Fig. 7 A proposed model of the TiAP1 interaction with chitin deacetylase to increase

the resistance to *Bgt* in wheat. (A) The *Bgt* cell walls were targeted by

wheat-secreted chitinases to release the chitin fragments that can further activate

the host immune system. (B) *Bgt* would secrete BgtCDA1 to convert the cell

wall chitin into chitosan, while the host invasion may protect the hyphae of the

pathogenic fungi from being hydrolysed by the extracellular plant chitinases,

thereby protecting the integrity of the fungal cell wall and inhibiting the

resistance response mechanism of the plant. (C) Interaction between TiAP1 and

BgtCDA1 in the transgenic *TiAP1* gene wheat, causing BgtCDA1 to inactivate

its deacetylation function, therefore the chitin in fungal cell walls would be

targeted by the wheat-secreted chitinases that liberate chitin the fragments that

can further activate the host immune system. When the transgenic *TiAP1* wheat

was infected by powdery mildew, the hormone signals related gene expression *in*

vivo would interact to activate plant disease resistance and upregulate the

expression of phenylalanine ammonia lyases in the lignin biosynthesis and

XTH30 in the xylan metabolism of the transgenic *TiAP1* wheat. Hence, it

enhances the ability of the plant cell walls to resist the penetration by pathogens.

1010

1011   **Supporting Informations**

1012   **Short Legends**

1013   Fig. S1 The TiAP1 prokaryotic expression and purification. (A) The TiAP1 protein

1014   expression was detected by SDS-PAGE. M: Low-molecular weight protein

ladder, Lane1: Uninduced TiAP1; Lane 2: 15 °C induced overnight. (B) TiAP1 quality assurance after renaturation. M: Low-molecular weight protein ladder; Lane 1: Bovine serum albumin (2.0 µg); Lane 2: TiAP1 protein (1.5 µg).

Fig. S2 Identification of the transgenic Bobwhite with the *TiAP1* gene and the subsequent hybrid generation. (A) Identification of the *Bar* gene in the transgenic *TiAP1* wheat. M: DL2000 DNA Marker [Takara Biomedical Technology (Beijing) Co., Ltd.]; Lane 1-10: transgenic lines. (B) Identification of the *TiAP1* gene in transgenic wheat. M: Trans2K$^{®}$ Plus II DNA Marker [TransGen Biotechnology (Beijing) Co., Ltd.]; Lane 1-2: Bobwhite and ddH$_2$O; Lane 3-12: transgenic lines. (C) Identification of the *TiAP1* gene in the subsequent hybrid generation of the wheat F$_3$. M: DL2000 DNA Marker [Takara Biomedical Technology (Beijing) Co., Ltd.]; Lane 1-10: subsequent hybrid generation of the F$_3$ lines; Lane 11: OE1; Lane 12: YN15 wheat.

Fig. S3 Validation of wheat powdery mildew resistance in the representative hybrid subsequent generation of F$_3$, which showed resistance to *E09* at 7 dpi in the seedling stage. Wheat YN15 and OE1 were used as the susceptible and resistant controls, respectively. Lane 1: YN15; Lane 2-11: The subsequent hybrid generation of the F$_3$ lines 1-10; Lane 12: OE1.

Fig. S4 Validation of the wheat powdery mildew resistance in 7 transgenic wheats Bobwhite with the *TiAP1* gene in the adult stage, which showed high resistance to powdery mildew. Bobwhite was used as the susceptible control. (A) Disease phenotype of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for ten days. (B) The proportion of lesions in the leaf area of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for ten days.

Fig. S5 Subcellular localisation of the fusion protein in the *N. benthamiana* leaf epidermal cells. (A) GFP, TiAP1-GFP and Δsp-TiAP1-GFP under normal conditions for GFP signal detection in *N. benthamiana* leaves, where GFP acted as a positive control signalling the nucleus and cytoplasm, while Δsp-TiAP1-GFP was localised in the cytoplasm. (B) Western blot detection of the GFP, TiAP1-GFP

and Δsp-TiAP1-GFP proteins expression.

Fig. S6 The BgtCDA1 protein bioinformatics analysis. (A) The domain of the amino

acid sequence of BgtCDA1 by Pfam; (B) The structure analysis of BgtCDA1 by

the Basic Local Alignment Search Tool (BLAST); (C) The SignalP analysis of the

amino acid sequence of BgtCDA1. (D) Alignment of the five conserved motifs of

the BgtCDA1 domain of *Saccharomyces cerevisiae* CDA1 (NP_013410) and

Saccharomyces cerevisiae CDA2 (NP_013411) as well as the five conserved

motifs (Motif 1-5) of the carbohydrate esterase family 4 (CE-4) that are indicated

by red boxes.

Fig. S7 Transcriptome analysis of the *TiAP1* and *BgtCDA1* co-regulated genes. (A)

Sample-to-sample cluster analysis results. (B) The pie chart shows the different

expression and up-regulated genes in OE2-2 relative to Bob-2. (C) Heat map

showing the differentially expressed genes (DEGs) inoculation with *E09* for two

days. (D) Gene ontology (GO) analysis of the top 30 DEGs in OE2-2 and Bob-2.

Methods S1 Additional description of methods.

Table S1 Evaluation of the resistance to *E09* in YN15, wheat lines overexpressing

TiAP1, and the hybrid subsequent generation of F_3 in the seedling stage. Lines

1-10: ten hybrid F_3 generation lines. YN15 and OE1 are as the susceptible and

resistant controls respectively

Table S2 Evaluation of the resistance to powdery mildew in Bobwhite and wheat lines

overexpressing *TiAP1* at the adult stage. OE 1-10 represent ten wheat lines

overexpressing *TiAP1*. Bobwhite is as the control

Table S3 Differential gene expression in the transgenic lines overexpression OE2 and

Bobwhite when inoculated with *E09* for two days. The default conditions of the

screening included $P < 0.01$ and fold change > 4

Table S4 Gene ontology (GO) analysis of the differentially expressed genes in OE2-2

and Bob-2

Table S5 Expression of selected pathogen-responsive genes in Bob-0, Bob-2, OE2-0,

and OE2-2 in the RNA-Seq experiments

Table S6 The primers used in this study

Video S1 Confocal microscopy (Leica SP5-X, Germany) z-scan image series of

TiAP-mYFP/mCherry-TaSYP51 on barley leaf epidermic cell under the infection

of *Bgh*. Bar = 20 μm.

1078

1079

1080

1081

1082

1083

1084

1085

1086

1087

1088

Figures and Tables

1089

1090

1091

1092

1093

Fig. 1 qRT-PCR and VIGS analysis of the *TiAP1* effect on the *Bgt* resistance, the

TiAP1 prokaryotic expression and purification, and the reduction of the powdery

mildew severity (lesions area) on the wheat seedlings by a recombination protein

treatment. (A) Expression levels of *TiAP1* in YN15 and SN6306 that were

induced by *Bgt*. Total RNA from the leaves of YN15 and SN6306 that were

induced by *Bgt* at 0, 12, 24, 48, 72, and 96 h were extracted as the template to be

detected by qRT-PCR. Data represent mean ± S.D. of *n* = 3 technical replicates.

(B) Relative quantification of expression levels of *TiAP1*. Total RNA from the

leaves of barley stripe mosaic virus (BSMV):00 and BSMV: TiAP1as inoculated

with *E09* for ten days were extracted as the template for the detection by

qRT-PCR. (C) Disease phenotype of the BSMV:00 and BSMV: TiAP1as

inoculated with *E09*. Photographs were taken at 10 dpi. (D) Microscopic

observation of the disease symptoms of BSMV:00 and BSMV: TiAP1as 10 dpi.

Bar = 100 μm. (E) The TiAP1 protein expression detection by western blotting

before and after induction. M: Pre-stained protein ladder; Lane1: Supernatant of non-induced; Lane 2: Subside of non-induced; Lane 3: Supernatant of induced; Lane 4: Subside of induced. (F) SDS-PAGE analysis of the inclusion body purification. M: Low-molecular weight protein ladder; Lane 1: Inclusions after dissolving the centrifugal supernatant; Lane 2: The effluent of supernatant was incubated with nickel iminodiacetic acid (Ni-IDA); Lane 3-6: The effluent of the imidazole elution of 50 mM; Lane 7-11: The effluent of the imidazole elution of 100 mM; Lane 12-13: The effluent of the imidazole elution of 300 mM; (G) TiAP1 quality assurance after renaturation. M: Pre-stained protein ladder; Lane 1: TiAP1 protein (1.5 µg). (H) Disease phenotype of YN15 smeared with TiAP1 and the tag protein from pET28a (CK) when inoculated with *E09* for five days. (I) Microscopic (Nikon Ni-U, Japan) analysis of the disease symptoms of YN15 smeared with TiAP1 and CK when inoculated with *E09* for five days. Bar = 100 µm. (J) The proportion of lesions in the leaf area of YN15 that was pre-treated with the recombinant TiAP1 five days before *Bgt E09* inoculation relative to the control. Control: Pre-treated with the purified recombinant label protein from the pET28a construct. Different letters above the bars indicate statistically significant differences ($P < 0.05$) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method.

Fig. 2 Identifying the resistance of transgenic *TiAP1* wheat. (A) Expression levels of *TiAP1* in Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for two days. (B) Disease phenotype of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for five days. (C) Microscopic (Nikon Ni-U, Japan) analysis of the disease symptoms of Bobwhite and the transgenic Bobwhite with *TiAP1* when inoculated with *E09* for two days. Bar = 100 µm. (D) Count of *Bgt* conidia in different stages of the *Bgt* when inoculated with *E09* for two days in Bobwhite and the transgenic Bobwhite with *TiAP1*. (E) The HI statistics of the *Bgt* when inoculated with *E09* for two days in Bobwhite and the transgenic Bobwhite with *TiAP1*. Different letters above the bars indicate statistically significant differences (P <0.05) as obtained by one-way analysis of variance with least significance difference and Duncan's multiple range test method. (F) Identifying the resistance of different wheat at the adult stage natural infected by powdery mildew in the glasshouse. (1) Bobwhite; (2) The transgenic Bobwhite with *TiAP1*; (3) YN15; (4) F_1 obtained from cross breeding between YN15 and homozygous transgenic *TiAP1* Bobwhite.

1150

1151

Fig. 3 Subcellular localisation of the fusion protein through transient expression of

agroinfiltrated GFP, TiAP-GFP, and Δsp-TiAP-GFP in the epidermal cells of *N*.

benthamiana after plasmolysis, and co-transient expression of TiAP1-mYFP and

mCherry-TaSYP51 in the epidermal cells of barley through bombardment. (A)

Confocal laser scanning microscopy (Leica SP5-X, Germany) observation of the

leaf epidermal cells revealed that Δsp-TiAP-GFP appears to be localised in the

<u>cytoplasm,</u> while TiAP1-GFP is predominantly localised in the extracellular

region, where white arrows indicate plasmolysis and the yellow dotted line

shows the location of the cell membrane of the two adjacent cells. A GFP signal

was used as the control. Bar = 20 µm. (B) The enlargement of the blue box in

TiAP1-GFP. (C) The enlargement of the blue box in Δsp-TiAP1-GFP. （D）*B.*

graminis f. sp. *hordei* induced the accumulation of TiAP1-mYFP and

mCherry-TaSYP51 at the infection site. This figure is part of Video S1. Bar = 20

µm.

1178

Fig. 4 Interaction *in vitro* and *in vivo* of TiAP1 with BgtCDA1. (A) Yeast two-hybrid

analysis of the interaction between TiAP1 and BgtCDA1. Yeasts expressing the

interacting   combinations   of   bait   and   prey   were   spotted   on   the

SD-HAWL/X-α-Gal/ABA. Bait-53 and AD-T (prey) were the positive controls, while

Bait-Lam and AD-T as well as Bait-BD and AD-BgtCDA1 were used as the negative

controls. (B) Luciferase complementation imaging assay showing that TiAP1

interacted with and BgtCDA1. The ΔspTiAP1-nLuc and cLuc-ΔspBgtCDA1 were

transiently co-expressed in *N. benthamiana*. Co-infiltration of cLuc/nLuc-tagged

ΔspTiAP1 protein with the nLuc/cLuc-tagged ΔspBgtCDA1 or the empty vectors

were   used   as   negative   controls,   where   the   images   were   observed   using

chemiluminescence   imaging   two   days   later.   (C)   Bimolecular   florescent

complimentary   assay   showing   TiAP1   interaction   with   BgtCDA1.   The

cGFP-ΔspTiAP1 and nGFP-ΔspBgtCDA1 were transiently co-expressed in *N.*

benthamiana. Co-infiltrations of nGFP/cGFP-ΔspTiAP1, cGFP/nGFP- ΔspBgtCDA1,

or the empty vectors group nGFP/cGFP were used as negative controls. The positive

controls were nYFP-14-3-3/14-3-3-cCFP. The green florescent protein (GFP) signal

was visualised using confocal microscopy (Leica SP5-X, Germany). Bar = 20 µm.

Fig. 5 The silencing of *BgtCDA1* reduced *Bgt* infection and the BgtCDA1 mediated

deacetylation of chitohexaose (A6). (A) Wheat YN15 leaves, bombarded with

RNAi and GUS reporter constructs, were infected with *Bgt* and calculated for

fungal haustorial index. The haustorial index was calculated as the ratio of

haustoria-containing transformed cells (GUS expressing cells) divided by the

total number of transformed cells. Data shown are mean values, $n = 4$, and a

minimum of 150 transformed cells were counted in each repeat. Relative

haustorial index was computed relative to the empty vector pIPKTA30N control

of each experiment, which was set to 100%. The Mlo-RNAi was as another

control. Different letters above the bars indicate statistically significant

differences (P <0.05) as obtained by one-way analysis of variance with least
significance difference and Duncan's multiple range test method. Arrow
indicates haustorium. Bar = 20 μm. (B) Wheat YN15 leaves, bombarded with
RNAi and GUS reporter constructs, were infected with *Bgt.* Arrow indicates
haustoria. Bar = 20 μm. (C) Disease phenotype of the BSMV:00 and BSMV:
BgtCDA1as inoculated with *E09.* Photographs were taken at 5 dpi. (D)
Microscopic observation of the disease symptoms of BSMV:00 and BSMV:
BgtCDA1as 3 dpi using *E09.* Bar = 100 μm. (E) The proportion of lesions in
the leaf area of BSMV:00 and BSMV: BgtCDA1as inoculated with *E09* five
days. (F) The hyphal density score of leaf material in BSMV:00 and BSMV:
BgtCDA1as 3 dpi using *E09* in microscopic observation field plots. Different
letters above the bars indicate statistically significant differences (P <0.05) as
obtained by one-way analysis of variance with least significance difference and
Duncan's multiple range test method. (G) Purified BgtCDA1 (2 uM) was
incubated with 1 mM A6 for 5, 30 min as indicated. The partially deacetylated
product A5D1 was detected. A, N-acetyl-D-glucosamine; D, D-glucosamine.

1230
1231
1232

Fig. 6 The transcriptome and qRT-PCR analysis of the transgenic wheat with *TiAP1*.

(A)   Heat map of selected pathogen-responsive genes in Bobwhite without and

with induction of *Bgt* for two days (Bob-0 and Bob-2, respectively), and

overexpression line OE2 as the same treatment (OE2-0 and OE2-2, respectively).

(B) qRT-PCR analysis of the differential expression gene *TraesCS1B02G440300*,

TraesCS2A02G161500,   *TraesCS2A02G199300*,   *TraesCS3B02G379200*,

TraesCS5D02G188600,   *TraesCS5A02G049600*,   *TraesCS3B02G259500*,

TraesCS2B02G224300,   *TraesCS6A02G266100*,   *TraesCS2D02G099900*,

TraesCS6B02G018700 and *TraesCS2B02G529400* of selected induced of *Bgt* for

two days.

1243
1244
1245

1246
1247

Fig. 7 A proposed model of the TiAP1 interaction with chitin deacetylase to increase the resistance to *Bgt* in wheat. (A) The *Bgt* cell walls were targeted by wheat-secreted chitinases to release the chitin fragments that can further activate the host immune system. (B) *Bgt* would secrete BgtCDA1 to convert the cell wall chitin into chitosan, while the host invasion may protect the hyphae of the

pathogenic fungi from being hydrolysed by the extracellular plant chitinases,

thereby protecting the integrity of the fungal cell wall and inhibiting the

resistance response mechanism of the plant. (C) Interaction between TiAP1 and

BgtCDA1 in the transgenic *TiAP1* gene wheat, causing BgtCDA1 to inactivate

its deacetylation function, therefore the chitin in fungal cell walls would be

targeted by the wheat-secreted chitinases that liberate chitin the fragments that

can further activate the host immune system. When the transgenic *TiAP1* wheat

was infected by powdery mildew, the hormone signals related gene expression *in*

vivo would interact to activate plant disease resistance and upregulate the

expression of phenylalanine ammonia lyases in the lignin biosynthesis and

XTH30 in the xylan metabolism of the transgenic *TiAP1* wheat. Hence, it

enhances the ability of the plant cell walls to resist the penetration by pathogens.

1265