DNA damage-induced phosphorylation of histone H2A at serine 15 is linked to

DNA end resection

Salar Ahmad, Valérie Côté and Jacques Côté#

St-Patrick Research Group in Basic Oncology, Centre Hospitalier Universitaire de Québec-

Université Laval Research Center, Laval University Cancer Research Center, Quebec City, QC,

G1R 3S3, Canada

Running Title: Phosphorylation of H2AS15 at DNA breaks

Corresponding authors: jacques.cote@crhdq.ulaval.ca

Word Count

Introduction, Results and Discussion: 3,138

Materials and Methods: 1,118

ABSTRACT

The repair of DNA double-strand breaks (DSBs) occurs in chromatin and several histone post-

translational modifications have been implicated in the process. Modifications of histone H2A

N-terminal tail has also been linked to DNA damage response, through acetylation or

ubiquitination of lysine residues that regulate repair pathway choice. Here, we characterize a

new DNA damage-induced phosphorylation on chromatin, at serine 15 of H2A in yeast. We

show that this SQ motif functions independently of the classical S129 C-terminal site (γH2A) and

mutant mimicking constitutive phosphorylation increases cell sensitivity to DNA damage.

H2AS129ph is induced by Tel1[ATM] and Mec1[ATR], and loss of Lcd1[ATRIP] or Mec1 signaling

decreases γH2A spreading distal to the DSB. In contrast, H2AS15ph is completely dependent on

Lcd1[ATRIP], indicating that this modification only happens when end resection is engaged. This is

supported by an increase of RPA and a decrease in DNA signal near the DSB in the *H2AS-15E*

phosphomimic mutant, indicating higher resection. This serine is replaced by a lysine in

mammals (H2AK15), which undergoes an acetyl-monoubiquityl switch to regulate binding of

53BP1 and resection. This regulation seems functionally conserved with budding yeast H2AS15

and 53BP1-homolog Rad9, using different post-translational modifications between organisms

but achieving the same function.

INTRODUCTION

The genetic information of the cell is challenged by various endogenous and exogenous cues

which leads to DNA damage. If unchecked, this can lead to mutations, translocations, loss of

genetic information and, in higher eukaryotes, be the underlying cause for diseases such as

cancer (1, 2). One of the most harmful form of DNA damage is DNA double-strand break (DSB).

To endure and mitigate such damage to the DNA, cells orchestrate cell cycle arrest and DNA

repair pathways, collectively termed the DNA damage response (DDR). Two main pathways to

repair DSBs are non-homologous end joining (NHEJ) and homologous recombination (HR). NHEJ

involves direct ligation of the two broken DNA ends which may require some processing,

making it a more error-prone pathway (3). In HR, single-stranded DNA (ssDNA) is generated

through a process called resection and involves the use of homologous sequence or sister

chromatid as a template to copy the genetic information, ensuring a lower error rate (4).

The phosphatidylinositol-3-OH kinase-related kinases (PIKKs) family plays an important

role in DDR. It includes *S. cerevisiae* Tel1 and Mec1 and mammalian ataxia telangiectasia-

mutated (ATM), AT-related (ATR) and DNA-dependent protein kinase (DNA-PK) (5, 6). One of

the first DNA damage-dependent chromatin modification to be identified was γ-H2A(X), which

is the phosphorylation of histone H2A at S129 in yeast and H2A.X at S139 in mammals (7-9).

This phosphorylation is carried out by both Tel1/ATM and Mec1/ATR. In yeast, H2AS129ph

spreads more than 50Kb on both sides flanking the DSB, whereas in mammals it covers up to a

megabase (10, 11). This histone mark is implicated in the DNA damage response and is required

for proper recruitment of several repair/signaling factors as well as chromatin

remodelers/modifiers (12-17). H2AX$^{-/-}$ mice are radiation sensitive, growth retarded, immune

deficient and show recruitment defect for a variety of repair factors such as Nbs1, 53BP1 and

BRCA1 (18).

Interestingly, budding yeast has another Tel1/Mec1-dependent SQ phosphorylation site

at H2B C-terminus, H2BT129ph, which largely correlates with H2AS129ph but is modulated by it

(10). Other DNA damage-induced phosphorylation marks have been mapped on chromatin but

are not deposited by PIKKs. These include H4S1ph, H2AS122ph and H2AT126ph in yeast, which

have been linked to processes like chromatin restoration after repair, meiosis, HR and repair of

fragile sites with CAG/CTG repeats (19-23). Apart from phosphorylation, other histone

modifications like acetylation, methylation, ubiquitylation and SUMOylation have been

implicated in DDR (16). In mammals H2AK15 is ubiquitinated by RNF168 and this mark is

responsible for recruitment and repair foci formation of 53BP1 in combination with H4K20me2,

inhibiting resection and favoring NHEJ (24). At DSBs, the TIP60 complex can acetylate H2AK15

and binds H4K20me, thus antagonizing 53BP1 and favoring repair by HR (25). In budding yeast,

Rad9 is the functional homolog of 53BP1, contains a Tudor domain which binds H3K79me

(equivalent to mammalian H4K20) and a BRCT domain which recognizes γ-H2A (17, 26, 27).

Both domains are required for the recruitment and retention of Rad9 at DSBs (17, 28). In

contrast to 53BP1, Rad9 does not have a ubiquitin-binding domain neither the yeast H2A has

lysine corresponding to human H2AK15 (**Fig. 1A**). Instead, yeast H2A has a SQ site at this

position which has been found phosphorylated in a Mec1-dependent manner upon DNA

damage, in a large phosphoproteomic study (29).

This led us to investigate whether H2AS15 had a role in the repair of DNA breaks. Our

results indicate that H2AS15ph is induced by Mec1[ATR] over a large domain of chromatin around

the DSB. We also present data arguing that H2AS15ph regulates Rad9 binding and DNA end

resection. Altogether, these results suggest an evolutionary conserved mechanism, similar to

the mammalian H2AK15 ubiquitin/acetylation switch, modulating yeast Rad9[53BP1] function on

the chromatin surrounding DNA breaks.

RESULTS

H2AS15ph function is distinct from H2AS129ph in response to DNA damage

In order to investigate if H2A-S15 has a role in DNA damage response, we mutated H2A-S15 to

alanine (non-phosphorylation mimicking) or glutamic acid (phosphorylation mimicking). In the

presence of DNA damaging agent methyl methanesulfonate (MMS), *H2A-S15E* mutant cells

show a slight growth defect, which is not seen for *H2A-S15A* (**Fig. 1B**). However, in combination

with the *H2A-S129A* mutant, both *H2A-S15A/E* show stronger growth defect compared to single

H2A-S129A mutant cells (**Fig. 1B**). Next, we tested if the *H2A-S15* mutants had any genetic

interaction with the other PIKK-dependent histone phosphorylation induced in response to

DSBs in yeast, H2BT129ph (10). *H2A-S15A/E H2B-T129A* double mutant cells do not show any

additional growth defect compared to *H2A-S15* single mutants in presence of MMS (**Fig. 1C**).

These results indicate that H2AS129 and H2AS15 phosphorylations play distinct roles in

response to DNA damage induced by MMS.

To decipher if the *H2A-S15* mutants affect H2AS129 phosphorylation *in vivo*, we

performed ChIP-qPCR at an inducible DSB. We used the yeast background in which a single

persistent DNA double strand break is created at the *MAT* locus by the homothallic

endonuclease (HO), inducible by galactose-containing media. The induced H2AS129ph signal

detected on each side of the DSB in wild type (WT) cells corresponds to what was previously

reported (10, 12). Both *H2A-S15A/E* mutants show similar pattern of H2AS129ph compared to

WT cells, not affecting the induction or spreading of H2A-S129ph beside a very small increase in

the *H2A-S15E* mutant (**Fig. 1D-E**). These results again support a role of H2AS15ph in DNA

damage response/repair that is distinct from H2AS129ph.

The lysine residues of H2A and H4 N-terminal tails are known targets of the NuA4

acetyltransferase complex (30). Studies have described histone modifications modulating the

occurrence of other modifications in chromatin, thus forming a network of histone crosstalk

(16). H4S1ph carried out by CK2 has been shown to inhibit acetylation of various lysine residues

on the N-terminal tail of H4 and play an active role in DSB repair as well as transcription (19,

20). This raised the question whether H2AS15ph could influence acetylation of lysine residues

on the H2A and H4 N-terminal tails. Western blot analysis of whole cell extracts from yeast

cells harboring the *H2A-S15* point mutants and grown in normal conditions reveals a small

decrease in both H2A and H4 acetylation, more noticeable for the *H2A-S15A* mutant compared

to wild type (**Fig. 1F**). We further investigated *in vivo* if the *H2A-S15* mutants affected the

acetylation of histones around DSBs. Both *H2A-S15A/E* mutants show a decrease in H4

acetylation signal but only close to the DNA break site (**Fig. 1G**). These results suggest that

H2AS15ph has a small but significant effect on other histone modifications in regions

surrounding the DSB.

H2AS15ph is induced over a large chromatin domain surrounding DNA double-strand breaks

To characterize the dynamics of H2AS15ph in response to DNA damage, we produced and

purified an antibody against the *S. cerevisiae* H2A N-terminal portion with Ser15 bearing the

phosphorylation mark. To demonstrate that H2AS15ph is indeed regulated by DNA damage, we

performed immunoblotting with purified yeast native chromatin. As shown in **figure 2A**,

H2AS15ph signal increases with MMS treatment, similarly to H2AS129ph. A control using

phosphatase treatment of the chromatin confirms that the signal detected by the antibody is

indeed phosphorylation-specific (**Fig. 2B**).

To determine whether H2AS15ph occurs at the DSB, we performed ChIP-qPCR at the

MAT locus before and after HO induction. The signal detected for H2AS15ph shows a clear

increase around the DSB site in WT background spanning over 10 kilobases on each side of the

break (**Fig. 2C**). As a control to check the specificity of the signal detected by the antibody, the

H2A-S15A mutant cells do not show such signal over a large domain after induction of the DSB

(**Fig. 2C**). However, close to the break site, *H2A-S15A* mutant cells do show a slight increase

which likely reflects some non-specific cross-reactivity of the antibody with another mark. To

characterize the dynamics of H2AS15ph appearance around the DSB, the ChIP-qPCRs were

performed at different time-points after induction of the HO endonuclease. Interestingly,

appearance of H2AS15ph seems delayed in comparison to H2AS129ph, reaching maximum

levels throughout the region after 3hrs, while H2AS129ph reaches its maximum levels at 3 and

7kb within an hour (**Fig. 2D-E**). Since H2AS129ph seems deposited before H2AS15ph, we tested

if it is required for H2AS15 phosphorylation to occur. ChIP-qPCR in wild type and H2AS129A

mutant cells does show a significant decrease of H2AS15ph signal around the HO-induced DSB,

but the mark is still present (**Fig. 2F**). These results indicate that γH2A/H2AS129ph can influence

phosphorylation of H2AS15 around the DSB but is not *per se* required for it to occur, in

agreement with the increase DNA damage sensitivity of the double mutants (**Fig. 1B**).

Altogether, these data clearly demonstrate that H2AS15ph is actively deposited over a large

region of chromatin surrounding DNA double-strand breaks.

Mec1^ATR is solely responsible for H2AS15 phosphorylation at DSBs

Tel1^{ATM} and Mec1^{ATR} signaling kinases phosphorylate histone H2AS129 to form γ-H2A (7, 11,

31). They also target various proteins to modulate DSB repair and checkpoint activation.

Recruitment of Tel1 to DSB requires Xrs2 (mammalian NBS1), subunit of the Mre11-Rad50-Xrs2

(MRX) complex (32). Deletion of *XRS2* affects Tel1-dependent checkpoint signaling and

telomere maintenance functions (33). Yeast Lcd1/Ddc2 (mammalian ATRIP) is responsible for

the recruitment of Mec1 to RPA-coated single-stranded DNA (ssDNA) produced during end

resection (34). Mec1 has been shown to be involved in spreading of γ-H2A in *trans* onto

adjacent undamaged chromosomes in close physical proximity (10, 35). Deletion of *LCD1* leads

to loss of Mec1 recruitment to RPA-coated ssDNA and, therefore, Mec1-dependent cell

signaling in response to damage (36, 37).

lcd1Δ and *lcd1Δxrs2Δ* mutant cells are respectively defective in the recruitment of Mec1

and both Mec1/Tel1 at the DSB. In *lcd1Δ* cells, where Tel1 recruitment and signaling is

unaffected, ChIP-qPCR experiments show that the signal for γ-H2A close to the break is similar

to wild type cells (**Fig. 3A**). However, farther from the break, at 7Kb and 10Kb, the signal is

significantly reduced, showing its relation to Mec1. In *lcd1Δxrs2Δ* cells, where both Mec1 and

Tel1 recruitments are abrogated, the γ-H2A signal is completely abolished throughout the

region (**Fig. 3A**). This is similar to what has been observed for *mec1Δ* and *mec1Δtel1Δ* mutant

cells in previous studies (10). Loss of *MEC1* affects γ-H2A spreading away from the break site,

whereas loss of *MEC1* and *TEL1* results in complete loss of γ-H2A.

A large scale phosphoproteomic study has detected phosphorylation of H2AS15 and

linked it to Mec1 (29). To investigate this, we carried out ChIP-qPCR using the H2AS15ph

antibody to study its spreading around the DSB in *lcd1Δ* and *lcd1Δxrs2Δ* mutant cells. As shown

above, wild type cells show enrichment for H2AS15ph over a large region on each side of the

DSB (**Fig. 3B**). However, *lcd1Δ* as well as *lcd1Δxrs2Δ* mutant cells do not show any H2AS15ph

signal. These results demonstrate that, unlike γ-H2A which is deposited by both the Mec1 and

Tel1 kinases, H2AS15 phosphorylation is carried out solely by Mec1, which links it to DNA end

resection after DSB formation.

H2AS15 is required to maintain fidelity of resection

Previous studies have shown that the *H2A-S129A* non-phosphorylation mutant leads to faster

resection from the HO-induced DSB compared to wild type cells (38, 39). This increased

resection may be attributed to the inability of Rad9 to bind H2AS129ph through its BRCT

domain, an essential interaction for its recruitment on chromatin (17, 27, 28, 40, 41). Budding

yeast Rad9, like its mammalian homolog 53BP1, functions in slowing down or keeping a check

on the rate of DNA end resection (42-45). To investigate whether H2AS15ph may affect

resection, we used an antibody against Rfa1, RPA largest subunit, as a proxy for the ssDNA

generated by resection, in ChIP-qPCR at the HO-induced DSB. Interestingly, *H2A-S15E*

(phosphorylation mimicking) mutant cells show increased RPA binding near the break, whereas

H2A-S15A mutant cells behave like wild type (**Fig. 4A**). The *H2A-S129A* cells also show increased

RPA signals, reflecting higher ssDNA/resection in this mutant. Double mutants *H2A-S15A/E*

H2A-S129A have similar increased RPA signals as *H2A-S129A* cells, maybe slightly less than

H2AS15E cells at some locations (**Fig. 4A**). To confirm that these RPA signals are the

consequence of increased DNA end resection, we directly measured by qPCR the DNA signal

5kb away from the DSB, compared to an unrelated genomic locus. In this assay, a very

significant and similar decrease of signal is detected in *H2A-S15E*, *H2A-S129A* and double

mutants, but not in *H2A-S15A* (**Fig. 4B**). These results indicate that H2AS15 phosphorylation

leads to increased resection at DSBs, similar to the effect of losing H2AS129 phosphorylation.

Furthermore, the effects of each mutant are non-additive, suggesting that they are

mechanistically related. An interesting possibility is whether H2AS15ph is also recognized by

the BRCT domain of Rad9, this domain being well characterized for its ability to bind

H2AS129ph (27). Surprisingly, *in vitro* peptide binding assays with recombinant Rad9 indicate a

clear binding preference for the histone H2A tail peptide, but in its non-phosphorylated form

(**Fig. 4B**). Importantly, this binding specificity is also confirmed using yeast whole cell extracts in

peptide pull-downs and detecting endogenous Rad9 by immunoblotting (**Fig. 4C**). In this case,

inducing DNA damage in the cells does not affect endogenous Rad9 ability to bind the H2A

peptide *in vitro*. This ability of H2AS15ph to block Rad9 binding to the H2A histone tail may in

part explain the increased end resection detected in *H2A-S15E* mutant cells but not in *H2A-*

S15A (**Fig. 4A-B**). As these results suggest that Rad9 may directly bind histone H2A tail in

chromatin *in vivo*, and that H2AS15 phosphorylation may regulate this interaction, we analyze

Rad9 binding near the HO-induced DSB by ChIP-qPCR. In wild-type conditions in this specific

yeast genetic background, we could only detect Rad9-myc signal by ectopic expression from a

plasmid. Furthermore, the specific signal based on the empty vector control could only be

detected very close (150bp) on each side of the HO break (**Fig. 4E**). At these locations, Rad9

binding is indeed abrogated in the *H2A-S129A* mutant, as expected, and validating the

specificity of the signal. In contrast, Rad9 binding does not seem affected by *H2A-S15A/E*

mutations. Since H2AS15 phosphorylation occurs more downstream away from the DNA break,

the Rad9 signal detected here close to the HO site may not reflect the population regulated by

H2AS15ph (see (42, 46) for models of two populations/functions of Rad9/53BP1 near DNA ends

versus larger domains). Another possibility is that H2AS15ph is not required for Rad9

recruitment but regulates the way it binds nucleosomes and influences end resection.

Altogether, these results implicate the DNA damage-induced H2AS15ph histone mark in the

regulation of end resection at DSBs.

DISCUSSION

Histone tails are sites of various post-translational modifications (PTMs) such as

phosphorylation, acetylation, methylation, ubiquitylation and SUMOylation. These

modifications are recognized by reader proteins containing specialized domains such as

ubiquitin binding domains, BRCT and FHA domains that bind phosphorylation, Chromodomains,

MBT domains, Bromodomains and PHD fingers that recognize methylation or acetylation.

Proteins and protein complexes use these interactions for their proper recruitment, retention

and/or orientation at the target sites to carry out their respective functions (47, 48).

In this study, we characterized yeast H2AS15ph as a novel histone modification in the

context of DNA damage. This modification is induced in the presence of DNA damaging agent

MMS, which is a known alkylating agent and produces DSBs as cells replicate their DNA in S-

phase. A large-scale proteomic study had identified H2AS15ph by mass spectrometry and

implicated Mec1^{ATR} for its detection (29). Using *in vivo* analysis, we show that indeed Mec1, and

not Tel1, is responsible for the induction of this mark around DNA breaks, unlike γ-H2A

(H2AS129ph) which is deposited by either Tel1 or Mec1 (**Fig. 3**). While the spreading of γ-H2A

at DSBs is not affected in *H2A-S15A/E* mutant cells, the *H2A-S129A* mutant does affect the

levels of H2AS15ph (**Fig. 1C-D, 2F**). Furthermore, a decrease of chromatin acetylation is

detected in *H2A-S15A/E* mutant cells near the DNA break (**Fig. 1F-G**). We know from previous

studies that acetylation of histone H4 and H2A by the NuA4 complex plays an important role for

the recruitment of various chromatin remodelers like INO80, RSC and SWI/ SNF required for

repair by HR (12, 49-51). Future work will be needed to determine if H2AS15 affects the

recruitment of these remodelers.

A most interesting finding reported here argues that yeast H2AS15ph corresponds to an

evolutionary conserved regulatory function to regulate 53BP1/Rad9 binding to the H2A tail and

DNA end resection. The yeast H2AS15 residue seems the functional homolog of mammalian

H2AK15 in modulating resection which in turn affects DNA repair pathway selection, albeit

through different PTMs. In mammals, when 53BP1 binds H2AK15ub in the vicinity of DSBs, it

inhibits resection and favors the NHEJ pathway. This binding inhibits TIP60-mediated

acetylation of H4 and H2A tails in chromatin, including H2AK15, which otherwise would block

ubiquitylation by RNF168 and 53BP1 binding, favoring resection and HR (25). *H2A-S15E* mutant

cells show increased resection, as seen with *H2A-S129A* mutant cells where Rad9[53BP1] is

incapable to bind and inhibit or slow down resection (**Fig. 4A-B**). *In vitro* data suggest that Rad9

binds the H2A N-terminal tail in its unmodified state and this binding is abolished when H2AS15

is phosphorylated (**Fig. 4C-D**). On the other hand, this regulated interaction does not seem to

affect Rad9 recruitment to the DSB (**Fig. 4E**). Since Rad9 is recruited to nucleosomes near DSBs

though bivalent interaction with H2AS129ph and H3K79me, our results suggest that Rad9

interaction with the H2A tail is not required for its recruitment but can regulate its function

linked to DNA end resection. 53BP1 also interacts with nucleosomes through three interfaces,

i.e. H2AK15ub-UDR, H4K20me-Tudor and H2A.XS139ph-BRCT. But, as in yeast, only two are

required for its binding to nucleosomes, γH2AX-BRCT being dispensable (24, 46, 52). We

proposed a model depicted in (Fig. 4F) where resection of DNA ends produces ssDNA which

gets coated by RPA and recruits Mec1[ATR] through its binding partner Lcd1/Ddc2[ATRIP] (34, 36).

Mec1 then phosphorylates H2AS15 which inhibits Rad9 function and therefore increases

resection. H2AS15ph may be a dynamic mark, switching between phosphorylation and non-

phosphorylation states to fine-tune the process of resection. This conserved function of the

H2A tail to regulate Rad9/53BP1 binding and resection is even further supported by a recent

report showing phosphorylation of the ubiquitin moiety of H2AK15ub (H2AK15pUbT12) to

inhibit 53BP1 binding and favor resection (53).

Future work will address whether specific proteins may bind H2AS15ph. While our

peptide pull-downs with yeast extracts followed by mass spectrometry did not yield clear

results (data not shown), we speculate that other DNA repair proteins such as Rtt107 and

Dpb11 which have BRCT domains could recognize H2AS15ph. Rtt107 was recently shown to

bind H4T80ph and γ-H2A to counteract Rad9-modulated checkpoint activity (54). Dpb11 has

been shown to act as a scaffold protein for various checkpoint proteins and is involved in Rad9

recruitment/retention at DSBs (55, 56). H2AS15 may also act redundantly with other DNA

damage induced histone phosphorylation marks and function as a scaffold for proper

orientation or retention of repair factors.

MATERIAL AND METHODS

Yeast strains

All yeast strains used in the current study are from FY406 (57) and JKM179 (58) (*hta1-*

htb1Δ::NatMX , *hta2-htb2Δ::KANMX*) backgrounds with deletion of both copies of H2A and H2B

genes. These strains carry an ectopic copy of *HTA1-HTB1* on low copy vector pRS413 (in FY406)

or pRS414 (in JKM179). Gene deletion and point mutations were obtained through standard

PCR techniques. Plasmid shuffling using 5-Fluoroorotic acid (5-FOA) was used to introduce the

plasmid with point mutations in respective backgrounds. Yeast background JKM139 (58) with

deletion of   *sml1Δ*::*NATMX* , *lcd1Δ*::*KANMX* , *xrs2Δ*::*TRP1* was used in this study. QY362

(JKM139 *RAD9-13MYC::KANMX*) was used for peptide pull down experiments while the Rad9-

myc ChIP-qPCR were done in the JKM179 background with *HTA1/2-HTB1/2* deletions covered

wild type or mutant H2A and ectopic Rad9-Myc expression from a pRS315 vector (40). General

lithium acetate method was used to transform the yeast cells. Cells were grown in YPD (1%

yeast extract, 2% peptone, 2% dextrose) at 30^{o}C until early log phase. For spot assay 1:10

dilution of the culture was spotted and grown at 30^{o}C for 3-4 days.

Chromatin Immunoprecipitation (ChIP)

ChIP-qPCR was performed as described previously (59). In brief, cells were grown overnight in

YP-Raffinose until OD_{600} of 0.5-1, at this point galactose (2%) was added to induced DSB at *MAT*

locus for 3 hours. This was followed by cross-linking of cells with formaldehyde and lysing with

bead-beater. Sonication was performed on Diagenode Bioruptor to obtain chromatin size of

~200-500bp. 100μg of chromatin was used to setup the immunoprecipitation reaction with

anti-H2A-S129ph (Upstate 07-745), anti-H2A-S15Ph (antibody was produced in rabbit using the

peptide sequence: Ac-GSAAKA(pS)QSRSAK-C. Affinity purification was performed to improve

the specificity of the antibody, with phosphorylated versus non-phosphorylated peptides.

Antibody was produced by Thermo Fisher Scientific), anti-Rfa1 (Agrisera AS07214), anti-H4ac

(Upstate 06-946), anti-H4 (Abcam 7311) and anti-Myc (9E10 Babco MMS150R) and reaction

was allowed to mix on a wheel overnight at 4^{o}C. Next morning Protein-A agarose beads

(protein-G magnetic beads for anti-Myc) were added followed by additional 3-4 hours on a

wheel at 4^{o}C. After multiple washing steps the DNA was eluted and incubated at 65^{o}C overnight

for de-cross linking followed by phenol-chloroform extraction of DNA. This DNA was

resuspended in NTE and used for qPCR. LC480 LightCycler (Roche) was used to quantify DNA

with the primer pairs as mentioned in the respective figures to calculate % of

immunoprecipitation (IP)/ input. The data represents mean of two independent biological

replicates. Error bars indicate the ranges between the two biological replicates. Fold

enrichment represents ratio of %IP/input at indicated loci around DSB normalized on signal at

negative-control locus IntergenicV. Levels of HO-induced DNA break was verified by qPCR

across the HO cut-site (normalized on control locus) and were similar in all experiments. Direct

measurement of end resection by qPCR was done using primers 4.8kb left of the HO cut-site

and presented as a ratio of signal over a control locus (*PRE1*, (60)). Primer sequences are

available upon request.

Purification of native yeast chromatin

Native yeast chromatin was prepared as described previously (61), cells were treated with MMS

(0.05%) or DMSO (control) for 2 hours to induce DNA damage. Lambda protein phosphatase

(λPP) (NEB) was used to dephosphorylate the purified chromatin. The reaction mixture was

incubated for 30 mins at 30^{o}C followed by booster of λPP and additional incubation for 30 mins

at 30^{o}C.

Recombinant protein purification

Recombinant Rad9 (C-terminal containing Tudor and BRCT domains) protein was purified as

described previously (62). In short, bacterial cells were grown overnight at 16^{o}C with IPTG

induction. Next day cells were lysed with lysozyme followed by sonication. The soluble portion

was incubated with glutathione sepharose (GE healthcare) beads for 3-4 hours at 4^{o}C and

eluted with glutathione. The eluted protein concentration was quantified by running on SDS-

PAGE with known BSA standards followed by coomassie staining.

Peptide-pull down

Peptide-pull down was performed with synthetic peptide corresponding to H2A N-terminal (aa

1-23) obtained from AnaSpec. 1μg of peptide was incubated with 1μg of GST-tagged protein

overnight on wheel at 4^{o}C in binding buffer (50 mM Tris–HCl pH 7.5, 50 mM NaCl, 0.1% NP-40, 1

mM PMSF and phosphatase inhibitors). Next morning streptavidin dynabeads were added to

the reaction mixture and kept at 4^{o}C for two additional hours followed by washing, running on

SDS-PAGE and western blotting with anti-GST (Sigma G1160).

Peptide-pull downs with yeast cell extracts were performed by growing cells in YPD and treating

with MMS (0.05%) or DMSO for 2 hours (final OD$_{600}$ ~1.0-1.5). Cells were centrifuged, the pellet

was washed with cold milliQ water, resuspended in sensitization buffer (100mM Tris-HCl pH

9.4, 10mM DTT) and incubated at 30^{o}C on a wheel for 15 mins. It was then centrifuged at 500g

for 5mins at 4^{o}C, resuspended in 30ml of spheroplasting buffer (1.2M Sorbitol 20mM Hepes pH

7.5), centrifuged again, resuspended in 40ml of spheroplasting buffer with lyticase and

incubated on wheel at 30^{o}C for 90 mins. After centrifugation at 500g for 5 mins, wash with

wash buffer (1.2M Sorbitol, 20mM Pipes pH 6.4, 1mM MgCl$_2$), it was resuspended in 25ml of

nuclear isolation buffer; NIB (250mM Sucrose, 60mM KCl, 1mM PMSF, 14mM NaCl, 0.8% Triton

X-100, 5mM MgCl$_2$, 1mM CaCl$_2$, 15mM MES), mixed well to remove clumps and kept on ice for

20 mins. After centrifugation and two washes with NIB (until white nuclear pellet was

obtained), the pellet was resuspended in pull-down buffer (400mM NaCl, 1mM PMSF, 50mM

Tris-HCl pH8, 0.5% NP-40, 5 mM β-glycerophosphate, 10 mM sodium butyrate, 5 mM NaF) to

get 300mM NaCl in the final volume, followed by incubation on rotating plate at 4^{o}C for 20

mins. After centrifugation at 14,000rpm at 4^{o}C for 30 mins, the supernatant was collected

(nuclear extract). Biotinylated peptides were bound to streptavidin dynabeads in peptide

binding buffer (150mM NaCl, 50mM Tris-HCl pH8.0, 0.1% NP-40, 5 mM β-glycerophosphate, 10

mM sodium butyrate, 5 mM NaF) by incubating at room temperature on wheel for 30 mins.

Peptide-dynabeads were washed with protein binding buffer (140mM NaCl, 50mM Tris-HCl pH

8.0, 0.1% NP-40, 0.5mM DTT, 10μM ZnCl$_2$, 5 mM β-glycerophosphate, 10 mM sodium butyrate,

5 mM NaF) followed by addition of 600μg of nuclear extract to each pull-down reaction and

incubated at 4^{o}C on wheel for 2 hours. Nuclear extract that was incubated with phospho

peptide was initially incubated with non-phospho peptide, on the other hand, nuclear extract

for non-phospho peptide was initially incubated with phospho peptide for 2 hours at 4^{o}C on

wheel. After 3-4 washed with final wash buffer (350mM NaCl, 50mM Tris-HCl pH8.0, 0.1% NP-

40, 0.5mM DTT, 10μM ZnCl$_2$, 5 mM β-glycerophosphate, 10 mM sodium butyrate, 5 mM NaF),

Laemmli buffer was added, boiled and resolved in SDS-PAGE followed by western blotting with

anti-Myc (9E10 Babco MMS150R).

Acknowledgments

We thank Xue Cheng for important advice. We are grateful to Jim Haber for sharing

reagents/strains and stimulating discussions. This work was supported by a grant from the

Canadian Institutes of Health Research (CIHR) to J.C. (FDN-143314). S.A. held a

Desjardins/Fondation du CHU de Québec studentship. J.C. holds the Canada Research Chair in

Chromatin Biology and Molecular Epigenetics.

REFERENCES

1.   Hanahan D, Weinberg RA. 2011. Hallmarks of cancer: the next generation. Cell 144:646-
74.
2.   Jackson SP, Bartek J. 2009. The DNA-damage response in human biology and disease.
Nature 461:1071-8.
3.   Chiruvella KK, Liang Z, Wilson TE. 2013. Repair of double-strand breaks by end joining.
Cold Spring Harb Perspect Biol 5:a012757.
4.   Symington LS. 2016. Mechanism and regulation of DNA end resection in eukaryotes. Crit
Rev Biochem Mol Biol 51:195-212.
5.   Abraham RT. 2001. Cell cycle checkpoint signaling through the ATM and ATR kinases.
Genes Dev 15:2177-96.
6.   Smith GC, Jackson SP. 1999. The DNA-dependent protein kinase. Genes & development
13:916-934.
7.   Downs JA, Lowndes NF, Jackson SP. 2000. A role for Saccharomyces cerevisiae histone
H2A in DNA repair. Nature 408:1001-1004.
8.   Paull TT, Rogakou EP, Yamazaki V, Kirchgessner CU, Gellert M, Bonner WM. 2000. A
critical role for histone H2AX in recruitment of repair factors to nuclear foci after DNA
damage. Current Biology 10:886-895.
9.   Rogakou EP, Boon C, Redon C, Bonner WM. 1999. Megabase Chromatin Domains
Involved in DNA Double-Strand Breaks in Vivo. The Journal of Cell Biology 146:905-916.
10.   Lee CS, Lee K, Legube G, Haber JE. 2014. Dynamics of yeast histone H2A and H2B
phosphorylation in response to a double-strand break. Nat Struct Mol Biol 21:103-9.
11.   Shroff R, Arbel-Eden A, Pilch D, Ira G, Bonner WM, Petrini JH, Haber JE, Lichten M. 2004.
Distribution and dynamics of chromatin modification induced by a defined DNA double-
strand break. Curr Biol 14:1703-11.

12.   Downs JA, Allard S, Jobin-Robitaille O, Javaheri A, Auger A, Bouchard N, Kron SJ, Jackson
412   SP, Côté J. 2004. Binding of Chromatin-Modifying Activities to Phosphorylated Histone
413   H2A at DNA Damage Sites. Molecular Cell 16:979-990.

414   13.   Morrison AJ, Highland J, Krogan NJ, Arbel-Eden A, Greenblatt JF, Haber JE, Shen X. 2004.
415   INO80 and gamma-H2AX interaction links ATP-dependent chromatin remodeling to DNA
416   damage repair. Cell 119:767-75.

417   14.   van Attikum H, Fritsch O, Gasser SM. 2007. Distinct roles for SWR1 and INO80 chromatin
418   remodeling complexes at chromosomal double-strand breaks. EMBO J 26:4113-25.

419   15.   van Attikum H, Fritsch O, Hohn B, Gasser SM. 2004. Recruitment of the INO80 complex
420   by H2A phosphorylation links ATP-dependent chromatin remodeling with DNA double-
421   strand break repair. Cell 119:777-88.

422   16.   Dantuma NP, van Attikum H. 2016. Spatiotemporal regulation of posttranslational
423   modifications in the DNA damage response. The EMBO Journal 35:6-23.

424   17.   Wysocki R, Javaheri A, Allard S, Sha F, Côté J, Kron SJ. 2005. Role of Dot1-Dependent
425   Histone H3 Methylation in G_1 and S Phase DNA Damage Checkpoint
426   Functions of Rad9. Molecular and Cellular Biology 25:8430-8443.

427   18.   Celeste A, Petersen S, Romanienko PJ, Fernandez-Capetillo O, Chen HT, Sedelnikova OA,
428   Reina-San-Martin B, Coppola V, Meffre E, Difilippantonio MJ, Redon C, Pilch DR, Olaru A,
429   Eckhaus M, Camerini-Otero RD, Tessarollo L, Livak F, Manova K, Bonner WM,
430   Nussenzweig MC, Nussenzweig A. 2002. Genomic Instability in Mice Lacking Histone
431   H2AX. Science 296:922-927.

432   19.   Cheung WL, Turner FB, Krishnamoorthy T, Wolner B, Ahn SH, Foley M, Dorsey JA,
433   Peterson CL, Berger SL, Allis CD. 2005. Phosphorylation of histone H4 serine 1 during
434   DNA damage requires casein kinase II in S. cerevisiae. Curr Biol 15:656-60.

435   20.   Utley RT, Lacoste N, Jobin-Robitaille O, Allard S, Cote J. 2005. Regulation of NuA4
436   histone acetyltransferase activity in transcription and DNA repair by phosphorylation of
437   histone H4. Mol Cell Biol 25:8179-90.

438   21.   Harvey AC, Jackson SP, Downs JA. 2005. Saccharomyces cerevisiae histone H2A Ser122
439   facilitates DNA repair. Genetics 170:543-53.

440   22.   Krishnamoorthy T, Chen X, Govin J, Cheung WL, Dorsey J, Schindler K, Winter E, Allis CD,
441   Guacci V, Khochbin S, Fuller MT, Berger SL. 2006. Phosphorylation of histone H4 Ser1
442   regulates sporulation in yeast and is conserved in fly and mouse spermatogenesis.
443   Genes & development 20:2580-2592.

444   23.   House NCM, Polleys EJ, Quasem I, De la Rosa Mejia M, Joyce CE, Takacsi-Nagy O, Krebs
445   JE, Fuchs SM, Freudenreich CH. 2019. Distinct roles for S. cerevisiae H2A copies in
446   recombination and repeat stability, with a role for H2A.1 threonine 126. eLife 8:e53362.

447   24.   Fradet-Turcotte A, Canny MD, Escribano-Díaz C, Orthwein A, Leung CC, Huang H, Landry
448   MC, Kitevski-LeBlanc J, Noordermeer SM, Sicheri F, Durocher D. 2013. 53BP1 is a reader
449   of the DNA-damage-induced H2A Lys 15 ubiquitin mark. Nature 499:50-4.

450   25.   Jacquet K, Fradet-Turcotte A, Avvakumov N, Lambert JP, Roques C, Pandita RK, Paquet E,
451   Herst P, Gingras AC, Pandita TK, Legube G, Doyon Y, Durocher D, Côté J. 2016. The TIP60
452   Complex Regulates Bivalent Chromatin Recognition by 53BP1 through Direct H4K20me
453   Binding and H2AK15 Acetylation. Mol Cell 62:409-421.

26. Grenon M, Costelloe T, Jimeno S, O'Shaughnessy A, FitzGerald J, Zgheib O, Degerth L, Lowndes NF. 2007. Docking onto chromatin via the Saccharomyces cerevisiae Rad9 Tudor domain. Yeast 24:105-119.

27. Hammet A, Magill C, Heierhorst J, Jackson SP. 2007. Rad9 BRCT domain interaction with phosphorylated H2AX regulates the G1 checkpoint in budding yeast. EMBO Rep 8:851-7.

28. Javaheri A, Wysocki R, Jobin-Robitaille O, Altaf M, Cote J, Kron SJ. 2006. Yeast G1 DNA damage checkpoint regulation by H2A phosphorylation is independent of chromatin remodeling. Proc Natl Acad Sci U S A 103:13771-6.

29. Bastos de Oliveira FM, Kim D, Cussiol JR, Das J, Jeong MC, Doerfler L, Schmidt KH, Yu H, Smolka MB. 2015. Phosphoproteomics reveals distinct modes of Mec1/ATR signaling during DNA replication. Mol Cell 57:1124-1132.

30. Allard S, Utley RT, Savard J, Clarke A, Grant P, Brandl CJ, Pillus L, Workman JL, Côté J. 1999. NuA4, an essential transcription adaptor/histone H4 acetyltransferase complex containing Esa1p and the ATM-related cofactor Tra1p. The EMBO Journal 18:5108-5119.

31. Redon C, Pilch DR, Rogakou EP, Orr AH, Lowndes NF, Bonner WM. 2003. Yeast histone 2A serine 129 is essential for the efficient repair of checkpoint-blind DNA damage. EMBO reports 4:678-684.

32. Nakada D, Matsumoto K, Sugimoto K. 2003. ATM-related Tel1 associates with double-strand breaks through an Xrs2-dependent mechanism. Genes Dev 17:1957-62.

33. Oh J, Al-Zain A, Cannavo E, Cejka P, Symington LS. 2016. Xrs2 Dependent and Independent Functions of the Mre11-Rad50 Complex. Mol Cell 64:405-415.

34. Zou L, Elledge SJ. 2003. Sensing DNA Damage Through ATRIP Recognition of RPA-ssDNA Complexes. Science 300:1542-1548.

35. Li K, Bronk G, Kondev J, Haber JE. 2020. Yeast ATM and ATR kinases use different mechanisms to spread histone H2A phosphorylation around a DNA double-strand break. Proceedings of the National Academy of Sciences 117:21354-21363.

36. Gobbini E, Cesena D, Galbiati A, Lockhart A, Longhese MP. 2013. Interplays between ATM/Tel1 and ATR/Mec1 in sensing and signaling DNA double-strand breaks. DNA Repair (Amst) 12:791-9.

37. Paciotti V, Clerici M, Lucchini G, Longhese MP. 2000. The checkpoint protein Ddc2, functionally related to S. pombe Rad26, interacts with Mec1 and is regulated by Mec1-dependent phosphorylation in budding yeast. Genes & Development 14:2046-2059.

38. Chen X, Cui D, Papusha A, Zhang X, Chu CD, Tang J, Chen K, Pan X, Ira G. 2012. The Fun30 nucleosome remodeller promotes resection of DNA double-strand break ends. Nature 489:576-80.

39. Eapen VV, Sugawara N, Tsabar M, Wu WH, Haber JE. 2012. The Saccharomyces cerevisiae chromatin remodeler Fun30 regulates DNA end resection and checkpoint deactivation. Mol Cell Biol 32:4727-40.

40. Nnakwe CC, Altaf M, Cote J, Kron SJ. 2009. Dissection of Rad9 BRCT domain function in the mitotic checkpoint response to telomere uncapping. DNA Repair (Amst) 8:1452-61.

41. Toh GW, O'Shaughnessy AM, Jimeno S, Dobbie IM, Grenon M, Maffini S, O'Rorke A, Lowndes NF. 2006. Histone H2A phosphorylation and H3 methylation are required for a novel Rad9 DSB repair function following checkpoint activation. DNA Repair (Amst) 5:693-703.

42.   Ferrari M, Dibitetto D, De Gregorio G, Eapen VV, Rawal CC, Lazzaro F, Tsabar M, Marini
F, Haber JE, Pellicioli A. 2015. Functional interplay between the 53BP1-ortholog Rad9
and the Mre11 complex regulates resection, end-tethering and repair of a double-strand
break. PLoS Genet 11:e1004928.
43.   Escribano-Díaz C, Orthwein A, Fradet-Turcotte A, Xing M, Young JT, Tkáč J, Cook MA,
Rosebrock AP, Munro M, Canny MD, Xu D, Durocher D. 2013. A cell cycle-dependent
regulatory circuit composed of 53BP1-RIF1 and BRCA1-CtIP controls DNA repair pathway
choice. Mol Cell 49:872-83.
44.   Noordermeer SM, Adam S, Setiaputra D, Barazas M, Pettitt SJ, Ling AK, Olivieri M,
Alvarez-Quilon A, Moatti N, Zimmermann M, Annunziato S, Krastev DB, Song F,
Brandsma I, Frankum J, Brough R, Sherker A, Landry S, Szilard RK, Munro MM, McEwan
A, Goullet de Rugy T, Lin ZY, Hart T, Moffat J, Gingras AC, Martin A, van Attikum H,
Jonkers J, Lord CJ, Rottenberg S, Durocher D. 2018. The shieldin complex mediates
53BP1-dependent DNA repair. Nature 560:117-121.
45.   Zimmermann M, Lottersberger F, Buonomo SB, Sfeir A, de Lange T. 2013. 53BP1
Regulates DSB Repair Using Rif1 to Control 5' End Resection. Science 339:700-704.
46.   Bothmer A, Robbiani DF, Di Virgilio M, Bunting SF, Klein IA, Feldhahn N, Barlow J, Chen
HT, Bosque D, Callen E, Nussenzweig A, Nussenzweig MC. 2011. Regulation of DNA end
joining, resection, and immunoglobulin class switch recombination by 53BP1. Mol Cell
42:319-29.
47.   Wilson MD, Durocher D. 2017. Reading chromatin signatures after DNA double-strand
breaks. Philos Trans R Soc Lond B Biol Sci 372.
48.   Rossetto D, Truman AW, Kron SJ, Cote J. 2010. Epigenetic modifications in double-strand
break DNA damage signaling and repair. Clin Cancer Res 16:4543-52.
49.   Bennett G, Peterson CL. 2015. SWI/SNF recruitment to a DNA double-strand break by
the NuA4 and Gcn5 histone acetyltransferases. DNA Repair 30:38-45.
50.   House NCM, Yang JH, Walsh SC, Moy JM, Freudenreich CH. 2014. NuA4 initiates dynamic
histone H4 acetylation to promote high-fidelity sister chromatid recombination at
postreplication gaps. Mol Cell 55:818-828.
51.   Cheng X, Cote V, Cote J. 2021. NuA4 and SAGA acetyltransferase complexes cooperate
for repair of DNA breaks by homologous recombination. PLoS Genet 17:e1009459.
52.   Zgheib O, Pataky K, Brugger J, Halazonetis TD. 2009. An oligomerized 53BP1 tudor
domain suffices for recognition of DNA double-strand breaks. Mol Cell Biol 29:1050-8.
53.   Walser F, Mulder MPC, Bragantini B, Burger S, Gubser T, Gatti M, Botuyan MV, Villa A,
Altmeyer M, Neri D, Ovaa H, Mer G, Penengo L. 2020. Ubiquitin Phosphorylation at
Thr12 Modulates the DNA Damage Response. Molecular Cell 80:423-436.e9.
54.   Millan-Zambrano G, Santos-Rosa H, Puddu F, Robson SC, Jackson SP, Kouzarides T. 2018.
Phosphorylation of Histone H4T80 Triggers DNA Damage Checkpoint Recovery. Mol Cell
72:625-635 e4.
55.   Liu Y, Cussiol JR, Dibitetto D, Sims JR, Twayana S, Weiss RS, Freire R, Marini F, Pellicioli A,
Smolka MB. 2017. TOPBP1(Dpb11) plays a conserved role in homologous recombination
DNA repair through the coordinated recruitment of 53BP1(Rad9). J Cell Biol 216:623-
639.

56. Granata M, Lazzaro F, Novarina D, Panigada D, Puddu F, Abreu CM, Kumar R, Grenon M,
Lowndes NF, Plevani P, Muzi-Falconi M. 2010. Dynamics of Rad9 chromatin binding and
checkpoint function are mediated by its dimerization and are cell cycle-regulated by
CDK1 activity. PLoS Genet 6:e1001047.
57. Hirschhorn JN, Bortvin AL, Ricupero-Hovasse SL, Winston F. 1995. A new class of histone
H2A mutations in Saccharomyces cerevisiae causes specific transcriptional defects in
vivo. Molecular and Cellular Biology 15:1999-2009.
58. Lee SE, Moore JK, Holmes A, Umezu K, Kolodner RD, Haber JE. 1998.
Saccharomyces Ku70, Mre11/Rad50, and RPA Proteins Regulate Adaptation
to G2/M Arrest after DNA Damage. Cell 94:399-409.
59. Rossetto D, Cramet M, Wang AY, Steunou AL, Lacoste N, Schulze JM, Cote V, Monnet-
Saksouk J, Piquet S, Nourani A, Kobor MS, Cote J. 2014. Eaf5/7/3 form a functionally
independent NuA4 submodule linked to RNA polymerase II-coupled nucleosome
recycling. EMBO J 33:1397-415.
60. Mojumdar A, Sorenson K, Hohl M, Toulouze M, Lees-Miller SP, Dubrana K, Petrini JHJ,
Cobb JA. 2019. Nej1 Interacts with Mre11 to Regulate Tethering and Dna2 Binding at
DNA Double-Strand Breaks. Cell Reports 28:1564-1573.e3.
61. Altaf M, Utley RT, Lacoste N, Tan S, Briggs SD, Cote J. 2007. Interplay of chromatin
modifiers on a short basic patch of histone H4 tail defines the boundary of telomeric
heterochromatin. Mol Cell 28:1002-14.
62. Setiaputra D, Ahmad S, Dalwadi U, Steunou A-L, Lu S, Ross JD, Dong M-Q, Côté J, Yip CK.
2018. Molecular Architecture of the Essential Yeast Histone Acetyltransferase Complex
NuA4 Redefines Its Multimodularity. Molecular and Cellular Biology 38:e00570-17.
564
565

FIGURE LEGENDS

Fig 1: *H2A-S15* mutants are sensitive to DNA damage and functions independently of γ-H2A.

(A) Sequence of H2A N-termini highlighting the SQ site at position 15 in *S. cerevisiae* and

corresponding lysine at position 15 in *H. sapiens*.

(B, C) Phenotypic analysis of yeast cells expressing wild type or the indicated H2A mutants. Ten-

fold serial dilutions of log phase cells were spotted on plates containing rich media alone or in

presence of DNA damaging agent methylmethane sulfonate (MMS). *H2A-S15E* showed a slight

growth defect in presence of the drug. Whereas, both *H2A-S15A/E* in background with *H2A-*

S129A showed stronger sensitivity than their respective single mutants. *H2B-T129A* did not

influence the growth of *H2A-S15A/E*.

(D, E) ChIP-qPCR showing γ-H2A at an inducible DSB is not affected by *H2A-S15A/E*. Cells were

grown in YP-raffinose overnight till log phase followed by addition of galactose to induce DSB

for 3 hours before cross-linking the chromatin. Primer pairs used flanked the DNA sequence on

both left and right side at 3Kb, 7Kb and 10Kb of the DSB at *MAT* locus. Anti-H4 signal was used

to normalize for histone occupancy. The data represents mean of two independent biological

replicates. Error bars indicate the range between the two biological replicates.

(F) *H2A-S15A/E* affects the acetylation of histone H4 and H2A. Western blot analysis of yeast

whole cell extracts (WCE) with anti-H4ac reveals a slight reduction in H4ac and H2Aac in *H2A-*

S15A and to a lesser extent in *H2A-S15E* compared to wild type. Histone H3 was used as loading

control.

(G) *H2A-S15A/ E* show reduced H4ac signal in regions close to the DNA break site. Signal for

H4ac was normalized for histone occupancy on total H4 signal. ChIP-qPCR was performed as

described in (D), with primers flanking 150bp, 1kb, 3Kb left and 0.5Kb, 1kb, 3Kb right of DSB.

Fig 2: H2AS15ph is locally induced upon DNA double strand break formation and spread over

several kilobases.

(A) H2AS15ph signal increased when yeast cells are treated with DNA damaging agent, MMS.

Western blot analysis was performed with 300ng of native chromatin purified from yeast cells

treated with DMSO (untreated control) or MMS. Anti-H2AS129ph was used as positive control

for the MMS treatment. Antibodies against histone H2B and H4 were used as loading control.

(B) Anti-H2AS15ph is specific for recognizing a phosphorylation mark on H2A. Western blot

analysis was performed with 300ng of native chromatin purified form MMS-treated yeast cells

after incubation with or without lambda protein phosphatase (λPP). Lane with λPP treatment

showed a significant decrease in signal for H2A-S15ph. Histone H4 was used as loading côntrol.

(C) H2AS15ph spreads on both sides of the HO endonuclease-induced single DSB at the *MAT*

locus. Signal for H2AS15ph showed enrichment after the induction of the DSB in wild type but

not in *H2A-S15A*. ChIP-qPCR was performed as described in Figure 1D. Fold enrichment was

calculated as ratio of %IP/ input normalized on total H4 for histone occupancy at indicated loci

around the DSB to the signal at the negative control locus intergenicV.

(D, E) Kinetics of H2S15ph and H2AS129ph accumulation after the induction of the HO DSB.

ChIP samples were analyzed after 0, 30min, 1hr and 3hrs of HO induction. Anti-H4 signal was

used to normalize for histone occupancy. Enrichments were measured as described in (C).

(F) Decreased H2AS15ph signals in cells lacking γH2A. ChIP-qPCR of H2AS15ph before and after

HO induction in wild type and *H2A-S129A* mutant cells. Enrichments were measured as

described in (C).

Fig 3: Mec1^ATR^ mediates phosphorylation of H2AS15 *in vivo*.

(A) Phosphorylation of H2AS129 carried out by Mec1^ATR^ and Tel1^ATM^. Signal for H2AS129ph at

DSB in *lcd1Δ* (loss of Mec1) is similar to wild type proximal to the break but much reduced distal

to the break. H2AS129ph signal is abolished in *lcd1Δxrs2Δ* (loss of Mec1 and Tel1). The line

representing the signal for GAL *lcd1Δxrs2Δ* is hidden behind since it has similar values as

uninduced/raffinose samples.

(B) Phosphorylation of H2AS15 is only mediated by Mec1^ATR^. Loss of H2AS15ph signal in *lcd1Δ*

and *lcd1Δxrs2Δ* compared to WT.

All yeast stains used in the above experiments have *sml1Δ* background. ChIP-qPCR was

performed as described in Figure 1D, primer pairs flanking 1Kb left and right of DSB were also

used.

Fig 4: H2AS15 regulates resection of DNA ends at DSB and Rad9^53BP1^ binding *in vitro*

(A) *H2A-S15E* mutant has increased DNA end resection compared to *H2A-S15A* and wild type

backgrounds, as shown by single-strand DNA binding RPA signal. *H2A-S129A* is used as a

positive control as it also leads to higher resection/RPA binding. ChIP-qPCR was performed as

described in Figure 1D with antibody against Rfa1 (largest RPA subunit) and primer pairs

flanking 1Kb, 3Kb and 7Kb left and right of DSB.

(B) Increased DNA end resection in the *H2A-S15E* mutant is also confirmed by directly

measuring DNA signal loss near the HO DSB. After 3hrs in inducing media, qPCR was used to

measure DNA at 4.8kb Left of the HO cut-site and at control locus in the genome (*PRE1*). Signal

ratios on the control locus show are decreased in *H2A-S15E* and *H2A-S129A* mutants, indicating

similar higher end resection in these cells.

(C) Peptide pull-down assay with GST-tagged recombinant Rad9 and indicated H2A N-terminal

peptides. rRad9 binds to unmodified peptide over phosphorylated peptide. Empty GST is used

as negative control.

(D) Peptide pull-down assay as in (B) but using extracts from yeast cells treated or not with

MMS and expressing myc-tagged Rad9.

(E) Binding of Rad9 close to the HO DSB is not affected by the H2AS15 mutants *in vivo*. ChIP-

qPCRs of Rad9-Myc ectopically expressed in the indicated backgrounds show significant binding

within 150bp of the break, which is lost in the *H2A-S129A* mutant but not in *H2A-S15E*. An

strain with an empty vector (E.V.) is used as control for background non-specific Myc signal.

(F) Model for the role H2AS15ph in DSB repair. Mec1 phosphorylates H2AS15 in the region

surrounding the DSB. This phosphorylation blocks Rad9 binding to the H2A tail in the bound

nucleosomes, which disrupts its function and results in increased resection. H2AS15ph may

function as a dynamic switch to modulate resection, phosphorylation enabling resection and

non-phosphorylation preventing hyper-resection.

(A)

```
             1              15
S.cerevisiae N-SGGKGGKAGSAAKASQSRSAK
H.sapiens    N-SGRGKQGGKARAKAKSRSSR
```

(B)

WT
H2A-S129A
H2A-S15A
H2A-S15E
H2A-S15A-S129A
H2A-S15E-S129A

YPD   MMS-0.03%

(C)

WT
H2A-S129A
H2B-T129A
H2A-S15A
H2A-S15E
H2A-S15A, H2B-T129A
H2A-S15E, H2B-T129A

YPD   MMS-0.035%

(D)

Raff WT
Raff *H2A-S15A*
GAL WT
GAL *H2A-S15A*

%IP H2AS129ph/H4

10 7 3 0 3 7 10 (kb)
Left   Right

(E)

Raff WT
Raff *H2A-S15E*
GAL WT
GAL *H2A-S15E*

%IP H2AS129ph/H4

10 7 3 0 3 7 10 (kb)
Left   Right

(F)

H2A
WT 15A 15E

Anti-H4ac
Anti-H2Aac
Anti-H3

(G)

WT
H2A-S15A
H2A-S15E

%IP H4ac/ H4

3 1 0.15 0 0.5 1 3 (kb)
Left   Right

FIGURE 2

(A)

(B)