Broca area homologue's asymmetry reflects gestural communication lateralisation in monkeys *(Papio anubis)*

Yannick Becker[1,2], Nicolas Claidière[1], Konstantina Margiotoudi[1], Damien Marie[1], Muriel Roth[2], Bruno Nazarian[2], Jean-Luc Anton[2], Olivier Coulon[3], & Adrien Meguerditchian[1*]

*Correspondence to: adrien.meguerditchian@univ-amu.fr.
[1]*Laboratoire de Psychologie Cognitive CNRS/Univ Aix-Marseille UMR7290*
[2]*Centre IRMf Institut de Neurosciences de la Timone CNRS/Univ Aix-Marseille*
[3] *Institut de Neurosciences de la Timone CNRS/Univ Aix-Marseille*
[4]*Station de Primatologie CNRS-CELPHEDIA UAR846*

Abstract:

Manual gestures and speech recruit a common neural network, involving Broca's area in the left hemisphere. Such speech-gesture integration gave rise to theories on the critical role of manual gesturing in the origin of language.

Within this evolutionary framework, research on gestural communication in our closer primate relatives has received renewed attention for investigating its potential language-like features.

Here, using in-vivo anatomical MRI in 50 baboons, we found that communicative gesturing is related to Broca homologue's marker in monkeys, namely the ventral portion of the Inferior Arcuate sulcus (*IA sulcus*). In fact, both direction and degree of gestural communication's handedness – but not handedness for object manipulation - are associated and correlated with contralateral depth asymmetry at this exact *IA sulcus* portion. In other words, baboons that prefer to communicate with their right hand have a deeper left-than-right *IA sulcus*, than those preferring to communicate with their left hand and vice versa. Interestingly, in contrast to handedness for object manipulation, gestural communication's lateralisation is not associated to the *Central sulcus* depth asymmetry, suggesting a double

dissociation of handedness' types between manipulative action and gestural

communication.

It is thus not excluded that this specific gestural lateralisation signature within the

baboons' frontal cortex might reflect a phylogenetical continuity with language-related

Broca lateralisation in humans.

Introduction:

Broca's area and its left hemispheric specialisation has historically been considered as the centre

of speech production (Friederici, 2017). Even if such a modular conception of language neural

bases was questioned by models of plastic and large distributed networks (Friederici, 2017;

Hickok & Poeppel, 2007), it is still well acknowledged that Broca's area in the left hemisphere

remains a key node for language specialisation within this neural distributed network.

Complementary work thereby highlighted Broca's area as an interface between speech and

multimodal motor integration including gesture and mouth mouvements (Gentilucci &Volta,

2008). Broca's area is also known for its involvement in motor planning, sequential and

hierarchical organization of behaviours, such as linguistic grammar or tool use and tool making

(Gentilucci & Volta, 2008; Koechlin & Jubault, 2006; Stout et al., 2015; Corballis, 2015; Wakita

et al., 2014). This body of work raises evolutionary questions about the role of the motor system

and gestural communication in language origins and its brain specialisation. Therefore, a

growing number of researchers proposed that language organization took some of its

phylogenetical roots into a gestural system across primate evolution (Gentilucci & Volta, 2008;

Corballis, 2015; Tomasello, 2008) Consequently, whereas comparative research has focused on

the potential continuities between language organisation and the vocal and auditory system in

nonhuman primates (eg. Boe et al., 2017; Jarvis, 2019; Wilson et al., 2017), the research on

gestural communication in apes and monkeys has recently received an increased interest within

this evolutionary framework.

A large body of non-human primate studies has documented some continuities of the

communicative gestural system with several key features of human language such as

intentionality, referentiality, learning flexibility and lateralisation (e.g., Tomasello, 2008;

Meguerditchian & Vauclair, 2014; Molesti et al., 2020). About manual lateralisation specifically,

studies in baboons and great apes have indeed showed that communicative manual gesturing

elicited stronger right-hand use in comparison to non-communicative manipulative actions

(reviewed in: Meguerditchain et al., 2013). In addition, at the individual level, a double

dissociation concerning the type of handedness has been documented between gestural

communication and object manipulation, showing that primates classified as right-handed for

communicative gesture are not especially classified as right-handed for object manipulation and

vice versa (Meguerditchian & Vauclair 2006, 2009). Those behavioural findings in different

primate species suggested a specific leftward lateralized system for communicative gestures,

which might be different from the one involves in handedness for object manipulation. This is

consistent with human literature showing that typical object-manipulation handedness measures

turned out to be rather a poor marker of language lateralisation (Fagard, 2013), as most left-

handers (78%) also show left-hemisphere dominance for language (Knecht et al., 2000; Mazoyer

et al., 2014), just like right-handed people. In both humans and nonhuman primates, direction of

handedness for object manipulation were found associated to contralateral asymmetries of the

motor hand area within the *Central sulcus* (e.g., humans: Amunts, 2000; Cykowski et al., 2008;

chimpanzees: Hopkins & Cantalupo, 2004; Dadda et al., 2008; Baboons: Margiotoudi et al.,

2019; Capuchin monkeys: Phillips & Sherwood, 2005; Squirrel monkeys: Nudo et al, 1992). In

fact, it has recently been demonstrated that the neural substrates of typical handedness measures

and language brain organisation might be not related but rather independent from each other

(Groen et al, 2013; Ocklenburg et al, 2014; Häberling et al., 2016).

Whether gestural communication's handedness in humans is a better predictor of language

lateralisation and is thus different than typical handedness measures remain unclear.

Nevertheless, several studies in human is supporting this hypothesis. In early human

development, the degree of right-handedness for preverbal gestures is more pronounced than

handedness for manipulation (Blake et al., 1994; Bonvillian et al., 1997; see also Fagard, 2013;

Cochet & Vauclair, 2010) and increases when the lexical spurt occurs in children contrary to

manipulation handedness (Cochet et al., 2011).

Given such potential lateralisation links between gesture and language in humans, it is thus not

excluded that the specific lateralisation's signature found for communicative gestures in

nonhuman primates might reflect evolutionary continuities with frontal hemispheric

specialisation for speech/gesture integration. This hypothesis might be relevant to investigate

given brain studies in nonhuman primates have shown human-like gross left-hemispheric

asymmetries of homologous language areas at a populational level: In particular Broca's

homologue in great apes (Cantalupo & Hopkins, 2001; Graïc et al., 2020) as well as the Planum

Temporale in great apes and even in baboons, an Old World monkey species, in both adult and

newborn (Gannon et al., 1998; Marie et al., 2018; Becker et al., 2021 a, Becker et al., 2021 b).

For Old World monkeys specifically, no study regarding structural asymmetry for Broca's

homologue has been investigated. One reason is that determining this area in monkeys is

particularly challenging in comparison to apes. In fact, the inferior precentral sulcus, the inferior

frontal sulcus and the fronto-orbital sulcus, which are common borders of Broca's homologue in

apes (Cantalupo & Hopkins, 2001), are absent in monkeys and thus delimitation is not trivial.

Nevertheless, all the detailed cytoarchitectonic studies addressing the Broca's homologue within

the frontal lobe in Old World monkeys (i.e. in mostly macaques but also in baboons) - and its

two components Area 44 and 45 - pointed towards the same sulcus of interest as the epicentre of

this region: the mid-ventral and ventral portion of the inferior arcuate sulcus (*IA sulcus*). The *IA*

sulcus is considered homologue to the ascending branch of the inferior precentral sulcus (Amiez

& Petrides, 2009) which delimits Broca's area posteriorly in humans and great apes. In monkeys,

Area 45 homologue sits in the anterior bank of the ventral *IA sulcus* (Petrides et al., 2005). In

contrast, Area 44 homologue might be located in the fundus and the posterior bank of the ventral

IA sulcus in monkeys (Petrides et al., 2005), which overlaps with F5 region related to the mirror

neuron system (Belmalih et al., 2009; Rizzolatti et al., 2019). Electric stimulation in the fundus

of the ventral *IA sulcus* elicited oro-facial and finger mouvements in macaques (Petrides et al.,

2005). Concerning baboons specifically, a cytoarchitectonic study (Watanabe☐Sawaguchi et al.,

1991) showed similarities to the macaque frontal lobe organisation given Area 45 anteriorly to

the *IA sulcus*, even if Area 44 was not described (Petrides et al., 2005; Belmalih et al., 2009;

Rizzolatti et al., 2019; Watanabe☐Sawaguchi et al., 1991). Therefore, in the absence of the usual

Broca's sulcal borders found in apes, the depth of the ventral part of the *IA sulcus* constitutes the

only critical neuroanatomical marker for delimiting the border and the surface of Broca's

homologue in monkeys.

In the present in-vivo MRI study conducted in 50 baboons (*Papio anubis*), we have (1) measured

the inter-hemispheric asymmetries of the *IA sulcus*' depth - from its dorsal to its most ventral

portion among subjects for which the *Central sulcus* depth measure was available from a

previous study (Margiotoudi et al., 2019) (2) as well as its potential links with direction and

degree of communicative gesture's handedness in comparison to handedness for manipulative

actions as measured with a bimanual tube task (see Hopkins, 1995).

To evaluate the hypothesis of an evolutionary continuity between the communicative gesture's

lateralised system in baboons and frontal hemispheric specialisation for language

(Meguerditchian et al., 2013) we tested specifically whether the depth asymmetry of the most

ventral Inferior Arcuate sulcus' portion (ventral *IA sulcus*, i.e., the Broca's homologue) - but not

the *Central sulcus* - was exclusively associated with communicative gestures' lateralisation.

Methods:

<u>Subjects:</u>

Inter-hemispheric asymmetries of the *IA sulcus*' depth were quantified from anatomical T1w

MRI images in 80 baboons *Papio anubis* born in captivity and free from developmental or

anatomical abnormalities or brain disorders (generation F1, 52 females, 28 males, age range = 7

to 32 years, mean age (years): M = 17,7 SE = 5,9). Out of this sample, 50 baboons (29 females

and 21 males, mean age (years): M = 12,3 SE = 5,8) had individual measures of handedness for

both communicative gestures (i.e. hand slapping gesture, Meguerditchian & Vauclair, 2006) and

manipulative actions (i.e. the bimanual tube task, Vauclair et al., 2005), as well as *Central sulcus*

depth measures as reported in Margiotoudi et al. (2019) in 63 individuals.

All baboons were housed in social groups at the Station de Primatologie CNRS (UPS 846,

Rousset, France; Agreement number for conducting experiments on vertebrate animals: D13-

087-7) and have free access to outdoor areas connected to indoor areas. Wooden and metallic,

ethologically approved, structures enrich the enclosures. Feeding times are held four times a day

with seeds, monkey pellets and fresh fruits and vegetables. Water is available ad libitum. The

study was approved by the "C2EA-71 Ethical Committee of Neurosciences" (INT Marseille)

under the number APAFIS#13553-201802151547729. The experimental procedure complied

with the current French laws and the European directive 86/609/CEE.

<u>Sulcal Parametrization:</u>

The *IA sulcus* and the *Central sulcus* were extracted from T1w images using the pipeline of the

free BrainVisa software (see Mangin et al., 2004 for details of the procedure). The sulcus

parametrization tool within the BrainVisa toolbox provides therefore sulcus-based morphometry

by subdividing the *sulci* of each hemisphere into 99 standardized positions from dorsal to ventral

sulcus extremities in order to quantify the variation of sulcal depth all across the *sulci's* 99

positions (Mangin et al., 2004). This automatic algorithm is free from observer's judgment. To

estimate asymmetries of the *sulci's* depth between the two hemispheres an asymmetry quotient

(AQ) for each of the 99 sulcal positions (AQ = (R – L) / [(R + L) × 0.5) was computed

(Margiotoudi et al., 2019). The AQ values vary between -2 and +2 with positive values

indicating right-hemispheric lateralisation and negative values indicating left-hemispheric

lateralisation.

<u>Behaviour Correlate:</u>

For further investigating its potential behavioural correlates, we tested whether the right- versus

left-handed groups classified for a given manual task (i.e., gestural communication versus

manipulative actions) differed in term of neurostructural depth asymmetries (AQ) within the *IA*

sulcus and the *Central sulcus*.

The two types of handedness measures were previously assessed (for communicative gesture:

Meguerditchian & Vauclair, 2006; Meguerditchian et al., 2011; and for manipulative actions:

Vauclair et al., 2005; Molesti et al., 2016) and were here included (1) when data of both manual

behaviours from the same subjects overlapped and (2) when sulci's depth extraction of the right

and left hemispheres were available in those subjects for both the *IA sulcus* and the *Central*

sulcus (i.e., Total: 50 baboons).

Communicative gesture was defined as a mouvement of the hand directed to a specific partner or

audience in order to affect its behaviour (Molesti et al., 2020). Similarly, then apes, some

communicative manual gestures in baboons have been found to share human-like intentional

control, referential properties, flexibility of acquisition and of use as well as similar specific

pattern of manual lateralisation (reviewed in Tomasello, 2008; Meguerditchian & Vauclair,

2014; Meguerditchain et al., 2013). The present study focused specifically on the "Hand

slapping" gesture which was previously found optimal for measuring gestural communication's

lateralisation in this species (Meguerditchian & Vauclair, 2006, 2009; Meguerditchian et al.,

2011). Indeed, the hand slapping behaviour – a probably innate gestures used by the baboon to

threat or intimidate the recipient - is the most common and frequent visual gesture of the

repertoire (Molesti et al., 2020) produced intentionally and unimanually in a lateralised manner

across similar agonistic contexts and similar emitter's postures (Meguerditchain et al., 2013).

Hand use was recorded in a baboon when slapping or rubbing quickly and repetitively the hand

on the ground in direction to a conspecific or a human observer at an out of reach distance.

Recorded events were taken from different bouts and not repeated measures from the same bout.

Handedness for manipulative actions was assessed using the well-documented bimanual

coordinated "Tube task" (Hopkins, 1995). Hand use was recorded when extracting food with a

finger out of a PVC tube hold by the other hand.

For both types of manual behaviours, in case a subject has been assessed in multiple sessions

within 2004-2015, the final classification as right or left-handed was selected based on the

session with the most observations.

The individual handedness index (HI) for a given manual behaviour, or degree of individual

manual asymmetry, was calculated based on the formula (#R-#L)/(#R+#L), with #R indicating

right hand responses and #L for left hand responses. The HI values vary between -1 and +1 with

positive values indicating right hand preference and negative values indicating left hand

preference.

Among the 50 baboons, for communicative gesture, 22 subjects were classified as left-handed,

28 as right-handed. For object-related manipulative actions (i.e., the bimanual tube task), 22

subjects were classified as left-handed, 28 as right-handed. Among the 50 baboons, 18 subjects

switched left-/right-handed categories of hand preference between communicative gesture and

manipulative actions (i.e., 9 from left-handed group for gestural communication to right-handed

group for manipulative actions, 9 from right-handed group for gestural communication to left-

handed group for manipulative actions).

<u>Statistical analysis:</u>

Statistical analysis was conducted using R 3.6.1 by Cluster Mass Permutation tests (Maris et al.,

2007).

First, an assembly of depth asymmetry measures was defined as a "cluster" when continuous

significant differences of the same sign across positions were found between groups (two-sided

t-tests, Welch corrected for inequality of variance, $p < .05$). Second, the sum of t-values within

each cluster was calculated (the "cluster mass").

Next, permutations were conducted for the between individual tests: For a given type of manual

behaviour, Left-handed individuals' AQ values versus Right-handed individuals' AQ values

were randomly redistributed between individuals and the maximum absolute cluster mass was

calculated for each randomly permuted set. This procedure was repeated 5000 times and the 99%

confidence interval (CI) of the maximum cluster mass was calculated. The clusters in the

observed data were considered significant at 1% level if their absolute cluster mass was above

the 99% CI of the distribution (i.e. $p < .01$).

We also performed a linear correlation between (1) the Handedness Index (HI) values for

communicative gesture calculated from the 50 individuals and (2) the Asymmetric Quotient

(AQ) values of those 50 baboons calculated from the respective left and right ventral *IA sulcus'*

depth sum of the continuous positions of the cluster for which a significant difference in AQ

score is detected by t-test comparison between the right- and left-handed groups. The same

procedure was followed for the HI values for non-communicative actions.

Data availability

The behavioural, neuro-anatomical and statistic code data that support the findings of this study

are available in "OSF Storage" with the identifier DOI 10.17605/OSF.IO/DPXS5.

https://osf.io/dpxs5/?view_only=f406ad972edd43e485e5e4076bae0f78

Results:

Between baboons communicating preferentially with the right hand versus the ones with the left hand, we found significant contralateral differences of depth asymmetries in the ventral portion of the *IA sulcus* (i.e., from the mid-ventral IA position to the most ventral *IA sulcus* portion, namely from contiguous positions 65 to 95 out of the 99 segmented positions of the entire *IA sulcus*) according to a cluster-based permutation test ($p < .01$, t-value clustermass= 76.16, for $p <$ .01 a clustermass of 65.28 was needed, see Maris et al., 2007). In other words, the 28 baboons using preferentially their right hand for communicative gestures showed more leftward *IA sulcus* depth asymmetry at this cluster than the 22 ones using preferentially their left-hand. In contrast, for non-communicative manipulative actions, we found no significant difference of depth sulcus asymmetries between the left- (N = 22) versus right-handed (N = 28) groups concerning any portion of the *IA sulcus,* according to a cluster-based permutation test ($p > .10$) (Fig. 1).

In addition, after calculating the AQ score per subject representing the sulcus depth asymmetry of the whole "Broca's cluster" (i.e., from the sum of the *IA sulcus* depths from positions 65 to 95 in the left hemisphere and the sum of *IA sulcus* depths from position 65 to 95 in the right hemisphere), we found a significant negative correlation between individual AQ depth values of the Broca's cluster (i.e., from positions 65 to 95) and individual handedness degree for communication (HI) : $r(48) = -.337$; $p < .05$ (i.e. stronger the hand preference is for one hand, deeper is the *IA sulcus* asymmetry from positions 65 to 95 in the contralateral hemisphere) (Fig. 2 A). In contrast, AQ depth values of the Broca's cluster did not show significant correlation with HI for non-communicative actions ($r(48) = -.037$; $p \approx 1$) (Fig. 2 B). Using the cocor package in R (Diedenhofen, B. & Musch, J., 2015), a comparison between these

two overlapping correlations based on dependent groups showed a significant difference between

the two correlations ($p < .05$).

When comparing with the control sulcus of interest, the *Central sulcus* related to the

primary motor cortex, opposite effect was found between handedness for manipulative actions

and hand preferences for communicative gesture. We found no significant difference of sulcus

depth asymmetries regarding communicative gestures. In contrast, Margiotoudi et al., (2019)

reported that the CS presented a contralateral asymmetry at continuous positions 56-60 (labelled

as the "Motor-hand area's cluster") for non-communicative manipulative actions, after

permutation tests for correction (Fig. 1).

Finally, we conducted a mixed-model analysis of variance with AQs depth values for the *IA*

sulcus "Broca's cluster" and for the *Central sulcus* "Motor hand area's cluster" (AQ derived

from continuous positions 56-60, see Margiotoudi et al., 2019) serving as the repeated measure

while communication handedness (left- *versus* right-handed) and action handedness (left- *versus*

right-handed) were between-group factors. The mixed-model analysis of variance demonstrated

a significant main effect on the AQ scores for communication handedness ($F1_{,46} = 14.08$, p

< 0.01) and for action handedness ($F_{1,46} = 4,1$, $p < 0.05$).

Figure 1: Effect of left-/right-hand direction of two handedness types (communication vs. action) on neuroanatomical sulcus depth asymmetries (*IA sulcus* vs. *Central sulcus*).

Left panel: Pictures of the two types of handedness measures in baboons. "Communication Handedness": a "Handslap" communicative gesture in a juvenile male "Action Handedness": the non-communicative bimanual coordinated "tube task" performed by an adult male.
Top panel: 3-D brain representation from BrainVisa software of the baboon's left hemisphere,

including the *IA sulcus* and the *Central sulcus* with the portion in purple where a significant
effect was found in Margiotoudi et al., 2019.
Graphs: Sulcus depth's asymmetry (AQ) comparison between right-handed group *versus* left-
handed group of baboons classified according to the type of manual tasks.Positive Mean
Asymmetry Quotient values (AQ) indicate rightward hemispheric asymmetry, negative Mean
Asymmetry Quotient values leftward hemispheric asymmetry. +/- SE indicated the Standard
Error.
(A) IA sulcus AQ between right-handed (N=28) *versus* left-handed (N=22) groups' classification
for communicative "Handslap" gesture. Significant contralateral AQ difference ($p < .01$)
between the two groups was found for a cluster including positions 65 to 95 (highlighted in
purple in the graph and the 3D representation of the *IA Sulcus*).
(B) Central Sulcus AQ between right-handed (N=28) *versus* left-handed (N=22) groups'
classification for non-communicative bimanual coordinated actions.
(C) IA sulcus AQ between right-handed (N=28) *versus* left-handed (N=22) groups' classification
for non-communicative bimanual coordinated actions.
(D) Initial graph *(Adapted from Margiotoudi et al., 2019)* of the *Central Sulcus* AQ showing the
significant contralateral AQ differences ($p < .05$) found between the left-handed (N=28) *versus*
right-handed (N=35) groups group for the non-communicative bimanual coordinated actions
(Action condition) for positions 56 to 61 (highlighted in purple in the graph and the 3D
representation of the *Central Sulcus*).

303 Figure 2: Correlation between handedness degree types and the Broca's cluster's asymmetry

305
A. Individual handedness degree (HI) for communicative gestures and AQ depth values of the
Broca's cluster (i.e., from positions 65 to 95) in dark blue dots.   Light blue line: Significant
negative correlation between HI and AQ.
B. Individual handedness degree (HI) for manipulative actions (HI) and AQ depth values of the
Broca's cluster (i.e., from positions 65 to 95) in dark green squares.   Light green line
(superposing on x axis): Non-significant correlation between HI and AQ.
312
313
Discussion:

The results of the study are straightforward. We showed that the IA sulcus left- or rightward

depth asymmetry at its mid-ventral and ventral portion (labelled as the "Broca cluster") is

associated exclusively with contralateral direction (left-/right-hand) of communicative manual

gestures' lateralisation in baboons but not handedness for non-communicative actions.

We also found a significant negative correlation between the Handedness Index (HI) values for

gestures and the Asymmetric Quotient (AQ) depth values of the *IA sulcus* "Broca cluster",

suggesting that the contralateral links between handedness for gestural communication and

depth asymmetries at the most ventral portion of the *IA sulcus* is evident not only at a qualitative

level but also at a quantitative level as well. In other words, individuals with a stronger degree of

manual lateralisation for communicative gestures have greater *IA   sulcus* depth asymmetries at

this ventral cluster in the hemisphere contralateral to their preferred hand for communication.

The ventral positions of such sulcal depth asymmetries are clearly at a crossroad of Broca-related

frontal regions including the fundus of the sulcus, Area 44 (Petrides et al., 2005), the anterior

bank, Area 45 (Petrides et al., 2005), the posterior bank and ventral F5 or granual frontal area

(GrF) (Belmalih et al., 2009; Rizzolatti et al., 2019). Since the sulcus depth might reflect a gyral

surface and its underlying grey matter volume, future work of delineating and quantifying grey

matter of the ventral *IA sulcus* would help determining which of those sub-regions of the Broca

homologue is driving the asymmetry specifically.

Whereas handedness for manipulative actions in baboons was previously found related to the

motor cortex asymmetry within the *Central sulcus* (Margiotoudi et al., 2019), our present

findings report the first evidence in monkeys that the neurostructural lateralisation's landmark of

communicative gesture is located in a frontal region, related to Broca homologue. Such a

contrast of results between manipulation and communication found at the cortical level is

consistent with what was found at the behavioural level in studies showing that communicative

gesture in baboons and chimpanzees elicited specific and independent patterns of manual

lateralisation in comparison to non-communicative manipulative actions (Meguerditchian &

Vauclair, 2009; Meguerditchian et al., 2010). Therefore, it provides additional support to the

hypothesis suggesting that gestural communication's lateralisation in nonhuman primates might

be, just as language brain organisation in human (see Häberling et al., 2016), related to a

different lateralised neural system than handedness for pure manipulative action. Its specific

correlates with Broca homologue's lateralisation is also consistent with what was found in our

closest relatives, the chimpanzee (Taglialatela et al., 2006; Meguerditchian et al., 2012).

Regarding Broca's area in humans, very recently, a functional segregation was proposed with

Broca's anterior part implicated in language syntax and its posterior part exclusively implicated

in motor actions (Zaccarella et al., 2021). The authors argued that action and language meet at

this interface. In this homologous area in macaques, neuronal recordings showed populations of

specific neurons activated for both volitional vocal and manual actions (Gavrilov & Nieder,

2021). In an evolutionary perspective we propose therefore that the intentional communicative

function of primate's communicative gesture might be a functional interface between actions and

language prerequisites, nested inside the monkeys' Broca's homologue (see also: Arbib 2006,

Rizzolatti 2017, Corballis, 2015)

It suggests that gestural communication may be a compelling embodied modality for one of the

evolutionary roots of the typical multimodal language system in humans and its left-hemispheric

specialisation.It is thus not excluded that language-related frontal lateralisation might be much

older than previously thought and inherited from a gestural communicative system dating back,

not to Hominid origins, but rather to the common ancestor of humans, great apes and Old World

monkeys, 25–35 million years ago.

363
Acknowledgments

General: We are very grateful to the Station de Primatologie CNRS, particularly the animal care

staff and technicians, Jean-Noël Benoit, Jean-Christophe Marin, Valérie Moulin, Fidji and

Richard Francioly, Laurence Boes, Célia Sarradin, Brigitte Rimbaud, Sebastien Guiol, Georges

Di Grandi for their critical involvement in this project; Leonard Samain-Aupic for sulci

labelling; Ivan Balansard, Sandrine Melot-Dusseau, Laura Desmis, Frederic Lombardo and

Colette Pourpe for additional assistance.

Funding: The project has received funding from the European Research Council under the

European Union's Horizon 2020 research and innovation program grant agreement No 716931 -

GESTIMAGE - ERC-2016-STG (P.I. Adrien Meguerditchian), from the French "Agence

Nationale de le Recherche" (ANR-12-PDOC-0014-01, LangPrimate Project, P.I. Adrien

Meguerditchian) as well as from grants ANR-16-CONV-0002 (ILCB) and the Excellence

Initiative of Aix-Marseille University (A*MIDEX). The Center IRM-INT (UMR 7289, AMU-

CNRS) is a platform member of France Life Imaging network (grant ANR-11-INBS-0006).

Author contributions: Y.B and A.M prepared the paper. Y.B performed the analyses. K.M.,
D.M. prepared the MRI images pre-processing and sulcus extraction pipeline. M.R designed
MRI sequences, B.N. designed the baboons' monitoring programs, R.L. designed the specific
procedures of welfare, anaesthesia and preparation of baboons in the MRI machine J.L.A
supervised and coordinated the MRI sessions. O.C. designed the sulcus parametrization tool. N.C
designed the statistic procedure. A.M. supervised the study and MRI acquisitions.

Competing interests: Authors declare no competing interests.

References and Notes

Amiez C, Petrides M. 2009. Anatomical organization of the eye fields in the human and non-
human primate frontal cortex. *Progress in Neurobiology* **89**:220–230.
doi:10.1016/j.pneurobio.2009.07.010

Amunts K, Jäncke L, Mohlberg H, Steinmetz H, Zilles K. 2000. Interhemispheric asymmetry of
the human motor cortex related to handedness and gender. *Neuropsychologia* **38**:304–312.
doi:10.1016/S0028-3932(99)00075-5

Arbib MA. 2005. From monkey-like action recognition to human language: An evolutionary
framework for neurolinguistics. *Behavioral and Brain Sciences* 63.

Becker Y, Phelipon R, Sein J, Velly L, Renaud L, Meguerditchian A. 2021a. Planum temporale
grey matter volume asymmetries in newborn monkeys (*Papio anubis*). *Brain Struct Funct*.
doi:10.1007/s00429-021-02278-9

Becker Y, Sein J, Velly L, Giacomino L, Renaud L, Lacoste R, Anton J-L, Nazarian B, Berne
C, Meguerditchian A. 2021b. Early Left-Planum Temporale Asymmetry in Newborn

Monkeys (*Papio anubis*): A longitudinal structural MRI study at two stages of development. *NeuroImage* 117575. doi:10.1016/j.neuroimage.2020.117575

Belmalih A, Borra E, Contini M, Gerbella M, Rozzi S, Luppino G. 2009. Multimodal architectonic subdivision of the rostral part (area F5) of the macaque ventral premotor cortex. *Journal of Comparative Neurology* **512**:183–217. doi:10.1002/cne.21892

Blake J, O'Rourke P, Borzellino G. 1994. Form and function in the development of pointing and reaching gestures. *Infant Behavior and Development* **17**:195–203. doi:10.1016/0163-6383(94)90055-8

Bonvillian JD, Richards HC, Dooley TT. 1997. Early Sign Language Acquisition and the Development of Hand Preference in Young Children. *Brain and Language* **58**:1–22. doi:10.1006/brln.1997.1754

Cantalupo C, Hopkins WD. 2001. Asymmetric Broca's area in great apes. *Nature* **414**:505. doi:10.1038/35107134

Cochet H, Jover M, Vauclair J. 2011. Hand preference for pointing gestures and bimanual manipulation around the vocabulary spurt period. *Journal of Experimental Child Psychology* **110**:393–407. doi:10.1016/j.jecp.2011.04.009

Cochet H, Vauclair J. 2010. Pointing gestures produced by toddlers from 15 to 30 months: Different functions, hand shapes and laterality patterns. *Infant Behavior and Development* **33**:431–441. doi:10.1016/j.infbeh.2010.04.009

Corballis MC. 2015. What's left in language? Beyond the classical model. *Annals of the New York Academy of Sciences* **1359**:14–29. doi:10.1111/nyas.12761

Cykowski MD, Coulon O, Kochunov PV, Amunts K, Lancaster JL, Laird AR, Glahn DC, Fox PT. 2008. The Central Sulcus: an Observer-Independent Characterization of Sulcal

Landmarks and Depth Asymmetry. *Cerebral Cortex* **18**:1999–2009.

doi:10.1093/cercor/bhm224

Dadda M, Cantalupo C, Hopkins WD. 2006. Further evidence of an association between

handedness and neuroanatomical asymmetries in the primary motor cortex of chimpanzees

(Pan troglodytes). *Neuropsychologia* **44**:2582–2586.

doi:10.1016/j.neuropsychologia.2006.03.037

Diedenhofen, B. & Musch, J. 2015. cocor: A Comprehensive Solution for the Statistical

Comparison of Correlations. *PLoS ONE*, 10(4): e0121945.

doi:10.1371/journal.pone.0121945

Fagard J. 2013. The nature and nurture of human infant hand preference. *Annals of the New York*

Academy of Sciences **1288**:114–123. doi:https://doi.org/10.1111/nyas.12051

Gannon PJ, Holloway RL, Broadfield DC, Braun AR. 1998. Asymmetry of Chimpanzee

Planum Temporale: Humanlike Pattern of Wernicke's Brain Language Area Homolog.

Science **279**:220–222. doi:10.1126/science.279.5348.220

Gavrilov N, Nieder A. 2021. Distinct neural networks for the volitional control of vocal and

manual actions in the monkey homologue of Broca's area. *eLife* **10**:e62797.

doi:10.7554/eLife.62797

Gentilucci M, Volta RD. 2008. Spoken language and arm gestures are controlled by the same

motor control system. *The Quarterly Journal of Experimental Psychology* **61**:944–957.

doi:10.1080/17470210701625683

Graïc J-M, Peruffo A, Corain L, Centelleghe C, Granato A, Zanellato E, Cozzi B. 2020.

Asymmetry in the Cytoarchitecture of the Area 44 Homolog of the Brain of the

Chimpanzee Pan troglodytes. *Front Neuroanat* **14**. doi:10.3389/fnana.2020.00055

Groen MA, Whitehouse AJO, Badcock NA, Bishop DVM. 2013. Associations between

Handedness and Cerebral Lateralisation for Language: A Comparison of Three Measures

in Children. *PLOS ONE* **8**:e64876. doi:10.1371/journal.pone.0064876

Häberling IS, Corballis PM, Corballis MC. 2016. Language, gesture, and handedness: Evidence

for independent lateralized networks. *Cortex* **82**:72–85. doi:10.1016/j.cortex.2016.06.003

Hopkins WD, Cantalupo C. 2004. Handedness in Chimpanzees (Pan troglodytes) Is Associated

With Asymmetries of the Primary Motor Cortex but Not With Homologous Language

Areas. *Behav Neurosci* **118**:1176–1183. doi:10.1037/0735-7044.118.6.1176

Hickok G, Poeppel D. 2007. The cortical organization of speech processing. *Nature Reviews*

Neuroscience **8**:393–402. doi:10.1038/nrn2113

Hopkins WD. 1995. Hand preferences for a coordinated bimanual task in 110 chimpanzees (Pan

troglodytes): Cross-sectional analysis. *Journal of Comparative Psychology* **109**:291–297.

doi:10.1037/0735-7036.109.3.291

Jarvis ED. 2019. Evolution of vocal learning and spoken language. *Science* **366**:50–54.

doi:10.1126/science.aax0287

Keller SS, Roberts N, García-Fiñana M, Mohammadi S, Ringelstein EB, Knecht S, Deppe M.

2010. Can the Language-dominant Hemisphere Be Predicted by Brain Anatomy? *Journal*

of Cognitive Neuroscience **23**:2013–2029. doi:10.1162/jocn.2010.21563

Koechlin E, Jubault T. 2006. Broca's area and the hierarchical organization of human behavior.

Neuron **50**:963–974. doi:10.1016/j.neuron.2006.05.017

Mangin J-F, Riviere D, Cachia A, Duchesnay E, Cointepas Y, Papadopoulos-Orfanos D, Collins

DL, Evans AC, Regis J. 2004. Object-based morphometry of the cerebral cortex. *IEEE*

Transactions on Medical Imaging **23**:968–982. doi:10.1109/TMI.2004.831204

Margiotoudi K, Marie D, Claidière N, Coulon O, Roth M, Nazarian B, Lacoste R, Hopkins WD,

Molesti S, Fresnais P, Anton J-L, Meguerditchian A. 2019. Handedness in monkeys

reflects hemispheric specialization within the central sulcus. An in vivo MRI study in

right- and left-handed olive baboons. *Cortex*. doi:10.1016/j.cortex.2019.01.001

Marie D, Roth M, Lacoste R, Nazarian B, Bertello A, Anton J-L, Hopkins WD, Margiotoudi K,

Love SA, Meguerditchian A. 2018. Left Brain Asymmetry of the Planum Temporale in a

Nonhominid Primate: Redefining the Origin of Brain Specialization for Language. *Cereb*

Cortex **28**:1808–1815. doi:10.1093/cercor/bhx096

Mazoyer B, Zago L, Jobard G, Crivello F, Joliot M, Perchey G, Mellet E, Petit L, Tzourio-

Mazoyer N. 2014. Gaussian Mixture Modeling of Hemispheric Lateralization for

Language in a Large Sample of Healthy Individuals Balanced for Handedness. *PLOS ONE*

9:e101165. doi:10.1371/journal.pone.0101165

Maris E, Oostenveld R. 2007. Nonparametric statistical testing of EEG- and MEG-data. *Journal*

of Neuroscience Methods **164**:177–190. doi:10.1016/j.jneumeth.2007.03.024

Meguerditchian A, Gardner MJ, Schapiro SJ, Hopkins WD. 2012. The sound of one-hand

clapping: handedness and perisylvian neural correlates of a communicative gesture in

chimpanzees. *Proc Biol Sci* **279**:1959–1966. doi:10.1098/rspb.2011.2485

Meguerditchian A, Molesti S, Vauclair J. 2011. Right-handedness predominance in 162

baboons for gestural communication : Consistency across time and groups. *Behavioral*

Neuroscience, **125**:653-660. doi: 10.1037/a0023823

Meguerditchian A, Vauclair J. 2014. Communicative Signaling, Lateralization and Brain

Substrate in Nonhuman Primates: Toward a Gestural or a Multimodal Origin of Language?

1 **7**:135–160.

Meguerditchian A, Vauclair J. 2009. Contrast of hand preferences between communicative

gestures and non communicative actions in baboons : implications for the origins of

hemispheric specialization for language. *Brain and Language* **108**:167–174. doi:

10.1016/j.bandl.2008.10.004.

Meguerditchian A, Vauclair J. 2006. Baboons communicate with their right hand. *Behavioral*

Brain Research, **171**:170-174. doi: 10.1016/j.bbr.2006.03.018

Meguerditchian A, Vauclair J, Hopkins WD. 2010. Captive chimpanzees use their right hand to

communicate with each other : Implications for the origins of hemispheric specialization

for language. *Cortex*, **46**:40-48. doi: 10.1016/j.cortex.2009.02.013

Meguerditchian A, Vauclair J, Hopkins WD. 2013. On the origins of human handedness and

language: A comparative review of hand preferences for bimanual coordinated actions and

gestural communication in nonhuman primates. *Developmental Psychobiology* **55**:637–

650. doi:10.1002/dev.21150

Molesti S, Meguerditchian A, Bourjade M. 2019. Gestural communication in olive baboons

(Papio anubis): repertoire and intentionality. *Animal Cognition*. doi:10.1007/s10071-019-

01312-y

Nudo RJ, Jenkins WM, Merzenich MM, Prejean T, Grenda R. 1992. Neurophysiological

correlates of hand preference in primary motor cortex of adult squirrel monkeys. *J*

Neurosci **12**:2918–2947. doi:10.1523/JNEUROSCI.12-08-02918.1992

Ocklenburg S, Beste C, Arning L, Peterburs J, Güntürkün O. 2014. The ontogenesis of language

lateralization and its relation to handedness. *Neuroscience & Biobehavioral Reviews*

43:191–198. doi:10.1016/j.neubiorev.2014.04.008

Petrides M, Cadoret G, Mackey S. 2005. Orofacial somatomotor responses in the macaque

monkey homologue of Broca's area. *Nature* **435**:1235–1238. doi:10.1038/nature03628

Phillips KA, Sherwood CC. 2005. Primary motor cortex asymmetry is correlated with

handedness in capuchin monkeys (cebus apella). *Behavioral Neuroscience* **119**:1701–1704.

doi:10.1037/0735-7044.119.6.1701

Rizzolatti G, Craighero L. 2004. The Mirror-Neuron System. *Annual Review of Neuroscience*

27:169–192. doi:10.1146/annurev.neuro.27.070203.144230

Stout D, Hecht E, Khreisheh N, Bradley B, Chaminade T. 2015. Cognitive Demands of Lower

Paleolithic Toolmaking. *PLOS ONE* **10**:e0121804. doi:10.1371/journal.pone.0121804

Taglialatela JP, Cantalupo C, Hopkins WD. 2006. Gesture handedness predicts asymmetry in

the chimpanzee inferior frontal gyrus: *NeuroReport* **17**:923–927.

doi:10.1097/01.wnr.0000221835.26093.5e

Tomasello M. 2008. Origins of human communication, The Jean Nicod lectures. Cambridge,

Mass: MIT Press.

Vauclair J, Meguerditchian A, Hopkins WD. 2005. Hand preferences for unimanual and

coordinated bimanual tasks in baboons (Papio anubis). *Cognitive Brain Research* **25**:210–

216. doi:10.1016/j.cogbrainres.2005.05.012

Wakita M. 2014. Broca's area processes the hierarchical organization of observed action. *Front*

Hum Neurosci **7**. doi:10.3389/fnhum.2013.00937

Watanabe☐Sawaguchi K, Kubota K, Arikuni T. 1991. Cytoarchitecture and intrafrontal

connections of the frontal cortex of the brain of the hamadryas baboon (Papio hamadryas).

Journal of Comparative Neurology **311**:108–133. doi:10.1002/cne.903110109

Wilson B, Marslen-Wilson WD, Petkov CI. 2017. Conserved Sequence Processing in Primate

Frontal Cortex. *Trends in Neurosciences* **40**:72–82. doi:10.1016/j.tins.2016.11.004

Zaccarella E, Papitto G, Friederici AD. 2021. Language and action in Broca's area:

Computational differentiation and cortical segregation. *Brain and Cognition* **147**:105651.

doi:10.1016/j.bandc.2020.105651

542