Effects of Perinatal Fluoxetine Exposure on Novelty-induced Social and Non-Social

Investigation Behaviors in a Seminatural Environment

Ole Christian Sylte[1,*], Jesper Solheim Johansen[1,*], Indrek Heinla[1], Danielle J Houwing[1,2],

Jocelien DA Olivier[2], Roy Heijkoop[1], and Eelke MS Snoeren[1,3]

[1] Department of Psychology, UiT the Arctic University of Norway, Norway

[2] Department of Neurobiology, Groningen Institute for Evolutionary Life Sciences, University

of Groningen, the Netherlands

[3] Regional Health Authority of North Norway

[*] These authors contributed equally.

22 Correspondence concerning this article should addressed to Eelke Snoeren, Department of

23 Psychology, UiT the Arctic University of Norway, 9037 Tromsø, Norway. E-mail address:

24 eelke.snoeren@uit.no

26   **Abstract**

Selective serotonin reuptake inhibitors (SSRIs) are increasingly prescribed as medication for

various affective disorders during pregnancy. SSRIs cross the placenta and affect serotonergic

neurotransmission in the fetus, but the neurobehavioral consequences for the offspring remain

largely unclear. Recent rodent research has linked perinatal SSRI exposure to alterations in

both social and non-social aspects of behavior. However, this research has mainly focused on

behavior within simplified environments. The current study investigates the effects of

perinatal SSRI exposure on social and non-social investigation behaviors of adult rat

offspring upon introduction to a novel seminatural environment with unknown conspecifics.

During the perinatal period (gestational day 1 until postnatal day 21), rat dams received daily

treatment with either an SSRI (fluoxetine, 10 mg/kg) or vehicle. Adult male and female

offspring were observed within the first hour after introduction to a seminatural environment.

The results showed that perinatal fluoxetine exposure altered aspects of non-social

investigation behaviors, while not altering social investigation behaviors. More specifically

both fluoxetine exposed males and females spent more total time on locomotor activity than

controls. Furthermore, fluoxetine exposed females spent less time exploring objects and

specific elements in the environment. The data suggest that perinatal exposure to SSRIs leads

to a quicker, less detailed investigation strategy in novel environments, and that the alteration

is mostly pronounced in females.

Keywords: antidepressants, fluoxetine, perinatal, behavior, social, rats, seminatural

environment, SSRI

1. Introduction

A considerable number of women experience depression or other mental disorders during pregnancy. Approximately 1 in 10 pregnant women fulfill the DSM-5 diagnostic criteria for major depressive disorder (Bennett et al. 2004; Woody et al. 2017). In treatment of maternal depression and anxiety, selective serotonin reuptake inhibitors (SSRIs) are the most frequently prescribed class of drugs, as it has been considered relatively safe for both mother and child. The prescription rate of SSRIs to pregnant women has increased tremendously in the last decades (Mitchell et al. 2011), and recent estimates suggest a worldwide prevalence of 3% (Molenaar et al. 2020) with significant geographically differences (Andrade et al. 2008; Charlton et al. 2015). Consequently, hundreds of thousands of babies exposed to SSRIs during early development are born every year. Despite the widespread use, we have limited knowledge on whether SSRI exposure during the early stages of brain development can lead to altered long-term behavioral outcomes, such as social and non-social behaviors.

Antidepressants, such as SSRIs, reach the fetus by crossing the placenta and are present in breast milk (Kristensen et al. 1999; Rampono et al. 2004). Thus, children can potentially be exposed to SSRIs during the entire perinatal period (Kim et al. 2006; Noorlander et al. 2008). SSRIs inhibit the function of the serotonin-reuptake transporter (SERT or 5-HTT), which leads to an accumulation of 5-HT in the synaptic cleft. This in turn increases the magnitude and duration of 5-HT activity at pre- and post-synaptic 5-HT receptors. In the adult brain, 5-HT acts mainly as a modulatory neurotransmitter, regulating emotion, cognition, sleep and stress responses (Olivier et al. 2011a). However, in the developing brain, 5-HT is widespread and acts as a neurotrophic factor regulating cell division, differentiation, migration, and synaptogenesis (Azmitia 2001; Gaspar et al. 2003). Consequently, developmental SSRI exposure is suggested to affect both neurodevelopment and later-life behaviors (Muller et al. 2016).

Previous studies in humans have shown associations between developmental SSRI exposure and impaired social behavior (Klinger et al. 2011), increased risk of speech and language disorders (Brown et al. 2016), and elevated levels of internalizing behavior, like anxiety and depression (Hermansen et al. 2016; Lupattelli et al. 2018; Malm et al. 2016), as well as increased risk of attention-deficit hyperactivity disorders (Man et al., 2018). While the existing literature has mainly examined the childhood years, little is known on whether these

associations persist into adulthood. In addition, outcomes such as depression may not emerge

before a certain age and could therefore remain undiscovered.

Epidemiological research on humans, like the above-mentioned studies, are

correlational in nature, and do not necessarily imply causation. A frequent problem with

human studies is the difficulty to isolate the effects of SSRI exposure from the effects of

maternal mental health. Women using SSRIs during pregnancy are likely suffering from

depression, which itself has been shown to have negative impact on the offspring (Dunkel

Schetter 2011; El Marroun et al. 2014; Goodman 2007). Animal research, on the other hand,

allows to control for potential interference from confounding factors, like maternal health,

drug dose and timing of exposure. As rodent and human serotonergic development is

remarkably similar (Glover and Clinton 2016), rodent studies can provide valuable

translational insight about how developmental SSRI exposure affects human offspring.

Rodent studies investigating the effects of developmental exposure to SSRIs have

reported alterations in different social and non-social behaviors in the offspring. In juvenile

male and female offspring, both pre- and post-natal SSRI exposure have been shown to

decrease social play behavior (Houwing et al. 2019b; Khatri et al. 2014; Olivier et al. 2011b;

Rodriguez-Porcel et al. 2011; Simpson et al. 2011). Similar tendencies were found in adult

rats with developmental SSRI exposure leading to less social interactions (Olivier et al.

2011b; Rodriguez-Porcel et al. 2011), or decreased interest to explore a novel conspecific

(Khatri et al. 2014; Rodriguez-Porcel et al. 2011; Simpson et al. 2011; Zimmerberg and

Germeyan 2015). SSRI exposure can also decrease (Houwing et al. 2020), or increase

(Gemmel et al. 2017; Kiryanova and Dyck 2014; Svirsky et al. 2016), aggressive-like social

behaviors. Furthermore, a recent meta-analysis revealed that developmental exposure to SSRI

was linked to reduced activity and explorative behaviors in adult rats and mice (Ramsteijn et

al. 2020).

Most rodent studies, however, have used simplified test set-ups which only

investigates a small fraction of all behaviors. Furthermore, these studies do not account for

the environmental and social complexity of real-world situations. To bypass these limitations,

recent studies from our research group have employed a seminatural environment enabling

rats to express many aspects of their natural behaviors (Hegstad et al. 2020; Heinla et al.

2020; Houwing et al. 2019a). These studies showed that perinatal SSRI fluoxetine (FLX)

exposure leads to various alterations in social and non-social behaviors in a naturalistic

setting. More specifically, perinatal fluoxetine exposure was associated with an increased

amount of passive social behaviors in both males and females, but a reduction of active social

behavior, general activity (Houwing et al. 2019a), and pro-social behaviors in females (Heinla

et al. 2020). Interestingly, these studies were performed in the seminatural environment

after the rats were familiarized to each other and the physical environment. It is currently

unknown how social and non-social behaviors manifest directly after introduction to a novel

environment with unfamiliar conspecifics. As perinatal SSRI exposure seem to alter stress-

coping behaviors (Houwing et al. 2019a), one could hypothesize that the stressor of a novel

environment with new conspecifics could lead to more pronounced changes in social and non-

social behaviors.

The aim of the current study was to investigate if perinatal SSRI exposure alters social

and non-social investigation behaviors in a novel environment with unknown conspecifics.

We define investigation as behaviors that provides the animal with information about a novel

stimulus. More specifically, social investigation refers to when the stimulus investigated is a

conspecific, such as when sniffing and grooming others, while non-social investigation refers

to investigation of inanimate objects and environmental locations. In line with previous

studies (Heinla et al. 2020; Houwing et al. 2019a), we expected perinatal fluoxetine exposure

to show a reduction in active social behavior in non-social investigation (exploratory)

behavior in the initial phase of the introduction to the seminatural environment. In addition, as

introduction to a new environment can be considered a stressful situation, we also expected to

observe an increase in self-grooming behavior in FLX-exposed animals.

2.   Material and Methods

The data was collected from video recordings obtained in a previously performed

experiment (Houwing et al. 2019a). The materials and methods are therefore similar to our

previous studies using data from the same experiment (Hegstad et al. 2020; Heinla et al. 2020;

Houwing et al. 2019a). However, the behavioral scoring scheme and the time window of

observation was unique for the current study.

2.1 Animals and dam housing

A total of 20 Wistar rats (10 males, 10 females), weighing 200-250 grams on arrival,

were obtained from Charles River (Sulzfeld, Germany) for breeding. After arrival, same-sex

pairs were housed in Makrolon IV cages (60 x 38 x 20 cm) on a reversed 12:12 hours

light/dark cycle, in which the lights were turned on at 23.00. Temperature in the room was 21

± 1°C, and the relative humidity was 55 ± 10 %. Standard rodent food pellets (standard chow,

Special Diets Services, Witham, Essex, UK), water and nesting material were available ad

libitum. Animal care and experimental procedures were conducted in agreement with

European Union council directive 2010/63/EU. The protocol was approved by the National

Animal Research Authority in Norway.

2.2 Breeding and antidepressant treatment

Daily, all females were checked for sexual receptivity by placing them together with a

male rat for 5 minutes. When lordosis behavior was observed, they were considered in

proestrus and thus ready for breeding. The female then got placed together with a male in an

isolated Makrolon IV cage for the next 24 hours (gestational day 0). Afterwards, they returned

to their initial same-sex pairs for the first two weeks of pregnancy. From gestational day 14,

the females were placed solitarily until delivery (gestational day 21/postnatal day 0).

During the 6-week period from conception (gestational day 0) to weaning (postnatal

day 21), females received either the SSRI fluoxetine 10 mg/kg (Apotekproduksjon, Oslo,

Norway) or vehicle (methylcellulose; Sigma, St. Louis, MO, USA) daily by oral gavage. The

offspring were thus exposed to perinatal fluoxetine via the treatment of the dams (in utero and

via breast feeding). The fluoxetine treatment was prepared with tablets for human usage that

were pulverized and dissolved in sterile water (2mg/mL) and injected at a volume of 5mL/kg.

Methylcellulose powder, the non-active filling of a fluoxetine tablet, was used as control

condition. The powder was dissolved in sterile water to create a 1% solution and administered

at a volume of 5mL/kg as well. Every third day, females were weighed to ensure correct

dosage of fluoxetine/vehicle. The chosen dosage of fluoxetine was decided upon comparison

of fluoxetine blood levels of humans and animals (Lundmark et al. 2001; Olivier et al.

2011b). When the rat dams got close to the end of pregnancy, they were checked two times a

day (09.00 and 15.00) for delivery.

2.3 Offspring housing

The offspring were housed together with their mothers until weaning (gestational day

21). After weaning, groups of two or three same-sex littermates were housed together in

Makrolon IV cages (see cage distribution in the supplemental materials, Table S1). They were

left undisturbed, except for the ovariectomy (see section for Procedure) and weekly cage

cleaning, until introduction to the seminatural environment at the age of 13-18 weeks. To

enable individual recognition, ears were punched. In Figure 1, a schematic overview shows all

experimental procedures from gestational day 0 to the end of the experiment.

Figure 1 Overview of experimental procedures

Figure 1. FLX = fluoxetine, CTR = control, P = postnatal day, G = gestational day. Created

with BioRender (https://biorender.com/).

2.4 Seminatural environment

The seminatural environment (SNE; 240 x 210 x 75 cm) consisted of two parts: an

open area and a burrow system (Figure 2; (Chu and Agmo 2014; Houwing et al. 2019a;

Snoeren et al. 2015)). Four openings (8 x 8 cm) connected the two areas. In the open area, two

partitions (40 x 75 cm) simulated natural obstacles. The burrow system consisted of

connected tunnels (width 7.6 cm, height 8 cm) and four nest boxes (20 x 20 x 20 cm).

Plexiglas covered the burrow at the height of 75 cm, while the open area remained open. A

curtain between the two parts allowed for different light settings. The burrow was left dark the

entire time. In the open area, on the other hand, light settings simulated a day-night cycle. A

lamp located 2,5 m above the floor, simulated daylight (180 lux) between 22.45 and 10.30.

From 10.30 to 11.00 the lights gradually decreased to 1 lux (simulating moonlight). The

darkness lasted until the light gradually increased from 1 to 180 lux between 22.15 and 22.45.

The whole ground of the SNE was covered with a layer (2 cm) of aspen wood chip

bedding (Tapvei, Harjumaa, Estonia). The nest boxes had 6 squares of nesting material in

each (non-woven hemp fibers, 5 x 5 fibers, 5 mm thickness, Datesend, Manchester, UK).

Three plastic shelters (15 x 16.5 x 8.5 cm, Datesend, Manchester, UK) were placed in the

open area. Additionally, 12 aspen wooden sticks (2 x 2 x 10 cm, Tapvei, Harjumaa, Estonia)

were randomly placed around in the SNE. A pile of food pellets (approx. 2 kg) and four

bottles of water were available at all time (see location in Figure 2A).

Two video cameras (Basler) were mounted on the ceiling, 2 m above the open area

(regular camera) and the burrow system (infrared camera) respectively. Media Recorder 2.5

was employed for video recordings. The data got immediately stored on an external hard

drive. The recording was manually stopped and restarted every 24 hours. The purpose was to

ensure that eventual errors only would affect one day of recorded data.

Figure 2 The Seminatural Environment

Figure 2. Illustration of the whole seminatural environment (A) and sectioning of the different

locations (B). 1 = open area close to burrow left, 2 = open area close to burrow right, 3 =

open area far away from burrow left, 4 = open area far away from burrow right, 5 = tunnels

far away from open a, 6 = tunnels close to OA, 7 = nestbox left, 8 = nestbox mid-left, 9 =

nestbox mid-right, 10 = nestbox right. Created with BioRender (https://biorender.com/). A

picture of the seminatural environment can be found in the Supplemental materials (Figure

S1).

2.5 Design of the study

Initially, five cohorts, each consisting of eight rat offspring, were placed one at the

time in the SNE. However, one day of video material was lost due to recording error, which

reduced the number of cohorts to four. A cohort consisted of 4 males and 4 females of which

each sex constituted 2 controls (CTR) and 2 fluoxetine (FLX) rats. Thus, data from this

experiment came from 8 CTR-males, 8 CTR-females, 8 FLX-males and 8 FLX-females (see

Table S2 for more details). Within a cohort, same sex rats came from different litters and were

thus unfamiliar to each other. Some rats had one sibling from the opposite sex in the same

cohort. However, these rats had been housed in different home cages since weaning. Analysis

of the data also revealed no differences in behavior towards males or females, suggesting that

the sibling factor did not influence the data.

2.6 Procedure

246   For the purpose of a previous study (Houwing et al. 2019a), the female offspring were

ovariectomized two weeks before entering the SNE in order to control their estrous cycle.

Although irrelevant for the objective of the current study, this procedure had the effect of

keeping the females in diestrus of the menstrual cycle during the observation period. Before

entering the SNE, the rats were shaved on the back and tail-marked under isoflurane

anesthesia for individual recognition (for more details, see (Houwing et al. 2019a)). All rats

were also weighed, confirming that there was no weight difference between CTR- and FLX-

rats.

254   Each cohort was placed in the SNE for 8 days. See Figure 1 for an overview of the

whole procedure. The cohorts were introduced to SNE on the first day (day 0) at 10.00 and

removed on day 8 at the same time. However, only data from the first hour was used for the

purpose of this study. All rats were again weighed after being removed from the SNE. No

difference in weight was observed between CTR- and FLX-rats. In order to remove olfactory

cues, the SNE was cleaned and bedding changed between cohorts.

2.7 Behavioral observations

262   The frequency and/or duration of several behaviors (see Table 1) were scored

manually using The Observer XT, version 12 (Noldus, Wageningen, The Netherlands). Two

observers, blinded for the animal treatment, independently scored either males or females

across all four cohorts. In addition to behavior, (1) location of the animal (see Figure 2B), (2)

whether the animal initiated the respective behavior or was respondent to it, (3) whether the

animal was in physical contact with another animal or not during the respective behavior, and

lastly, (4) ID of the interacting partner was scored. Since we were interested in observing how

the rats behaved in a novel environment with unfamiliar conspecifics, all rats were scored in

the first 60 minutes after entry to the SNE.

2.8 Data preparation and statistical analysis

275   As shown in Table 2, the recorded behaviors were combined into behavioral clusters.

For each rat, we calculated the total duration and the number of events for every behavior and

behavioral cluster. This data was later divided into six 10-minute time-bins in order to analyze

behavioral changes over time. Latencies to meet the other rats, and latencies to visit the

different locations of the SNE were also noted. This data was later divided and analyzed

cumulative over the first 1, 3, 5, 10, 20, 30, and 60 minutes. In this study, we operationalized

social investigation behaviors as the cluster "socially active behaviors" and the latencies to

meet all other rats, whereas non-social investigation behaviors were operationalized as the

cluster "general activity" and latencies to visit all the locations (See Figure 2B).

Table 1 Description of recorded behaviors

Behavior	Description
Walking/running	Walking or running through the environment
Chasing	Running forward in the direction of a conspecific
Non-social exploration	Exploring the environment by sniffing, usually when slowly walking or sitting still
Digging	Digging, pushing or carrying bedding/nesting/food material
Resting/immobile alone	Sitting or sleeping with minimal movement of the head without other rats in close vicinity
Resting/immobile socially	Sitting or sleeping with minimal movement of the head with at least 1 other rat on maximum 1 rat body length away
Hiding alone	Being in the shelter alone
Hiding socially	Being in the shelter with at least one other rat
Following	Walking or running in the same direction as another rat in front.
Allogrooming	Grooming any part of a conspecific's body, usually on the head or in the neck region
Sniffing anogenitally	Sniffing the anogenital region of the conspecific
Sniffing nose-to-nose	Sniffing the facial region of the conspecific
Sniffing body	Sniffing any part of the conspecifics body, except for the anogenital and facial region
Fighting	Kicking, pouncing, pushing, grapping, boxing or wrestling another rat
Nose-off	Facing another rat, usually in a tunnel, resulting in one rat moving forward and the other backing up
Self-grooming	Grooming itself
Freezing	Complete absence of movement in addition to a tense body posture
Rearing supported	Raising itself upright on its hind paws, facing a wall or an object

Rearing unsupported	Raising itself upright on its hind paws, not facing a wall or an object

288 **Table 2 Description of behavioral clusters**

Cluster	Behaviors within clusters
Socially active behaviors	Sniffing anogenitally, sniffing nose-to-nose, sniffing body, and allogrooming
General activity	Walking/running, non-social exploration
Non-socially passive behaviors	Resting alone, hiding alone
Socially passive behaviors	Hiding socially, resting socially
Conflict behaviors	Nose-off, fighting

290   Normality of data was determined with Shapiro-Wilks tests. Data with p < .05 was

analyzed non-parametrically. Simple group comparisons were performed with either a student

t-test or the non-parametric Mann-Whitney U test. Repeated measures ANOVA was used

when the behaviors were analyzed over time. In cases the Mauchly's test indicated violation

of sphericity from the ANOVA output, the degrees of freedom were corrected using

Greenhouse-Geisser estimates of sphericity. To correct for multiple comparisons, the

Benjamini-Hochberg procedure was performed on all significant results together with a

predetermined set of variables (sniffing, self-grooming, non-social exploration, conflict

behaviors). All tests reported were done 2-tailed.

299   Because male and female behavior were scored by two different observers, no

300 conclusions were drawn regarding potential differences between males and females.

302 **2.9 Statement Open Science Framework (OSF)**

303   The design of our study was preregistered on OSF on the 25th of March 2019

(https://osf.io/m87j5). There were no changes in analysis, except that we did not use the

originally planned additional control group. As stated at OSF, the planned control group was

not suitable, because it consisted of aged rats and had a different composition in number of

rats (7 versus 8). We therefore concluded that these differences would make it impossible to

compare the cohorts of the current study.

3 Results

From the behavioral scoring, we obtained a lot of data. A complete overview of all

behaviors can be found in Table S3 and S4. In this result section, we only mention the

relevant behavior to the purpose of the paper. A summary of the main finding are provided in

Table 3.

3.1 *Social investigation behaviors*

The data analysis revealed that CTR- and FLX-females did not differ in time spent on

($t = -1.04$, $p = .315$, $d = -0.52$, Figure 3A) or number of episodes ($t = -1.04$, $p = .318$, $d = -$

0.52) performing socially active behaviors. When looking separately at the different

behavioral components constituting the cluster (see Table 2), CTR- and FLX-rats did not

differ on any other behavioral components constituting the clusters relevant to social

behaviors (socially active behaviors, socially passive behaviors and conflict behaviors). No

difference was found between CTR- and FLX-males for socially active behaviors in total time

($t = 0.95$, $p = .356$, $d = 0.48$, Figure 3D) or on number of episodes ($t = 0.103$, $p = .919$, $d =$

0.05).

Although the treatment groups did not differ in the amount of socially active

behaviors, it could still be the case that the groups had different interest in meeting other rats.

To investigate this possibility, we first looked at the latencies to when the rats had met all

seven other cohort-members. The data analysis showed that there was no significant

difference in latency to meet all cohort-members between CTR- and FLX-rats for females ($t =$

0.84, $p = .418$, $d = 0.42$) or males ($U = 24.00$, $z = -0.84$, $p = .422$, $r = -.21$). We subsequently

measured how many cohort members the rats had met as a function of time. CTR- and FLX-

rats were compared on cumulative data measured at 1/3/5/10/20/30/60 minutes. For FLX-

females, there were no significant differences in the number of rats met (treatment effect:

$F(1,14) = 0.05$, $p = .821$) or in the pattern of rats met (timepoints x treatment: $F(1.73, 24.24)$

$= 0.28$, $p = .725$) over time compared to CTR-females (Figure 3G). Similarly, CTR- and

FLX-males did not differ in the number of rats met across all timepoints (treatment effect:

$F(1,14) = 0.49$, $p = .492$) or in the pattern of rats met over time (timepoints x treatment:

$F(2.05, 28.74) = 0.59$, $p = .563$, Figure 3H).

3.2 Other social behaviors

We also investigated some other social behaviors, such as socially passive behaviors

and conflict behaviors. No difference was found between CTR- and FLX-females in total time

($U = 33.00$, $z = 1.05$, $p = 1$, $r = .03$, Figure 3B) or number of episodes being socially passive (t

$= -0.28$, $p = .784$, $d = -0.14$). Furthermore, CTR- and FLX-females spent a similar amount of

time ($t = 0.03$, $p = .978$, $d = 0.01$, Figure 3C) and episodes ($t = -0.40$, $p = .692$, $d = -0.20$) in

conflict with other rats. Similarly, for males, no differences were found for time spent on

social passive behavior ($U = 41.00$, $z = 0.95$, $p = .382$, $r = .24$, Figure 3E), episodes of social

passive behavior ($t = 1.48$, $p = .161$, $d = 0.74$), time spent on conflict behavior ($t = -0.03$, $p =$

$.655$, $d = -0.02$, Figure 3F), or episodes in conflict behavior ($U = 42.00$, $z = 1.05$ $p = .786$, $r =$

$.26$).

Figure 3 Social behaviors in females and males

Figure 3. The data represent the time spent (s) on socially active behaviors (A, D), socially

passive behaviors (B, E), conflict behaviors (C, F), and the total number of rats met over time

(G, H). All graphs show comparisons between CTR-females (n = 8) and FLX-females (n = 8)

or between CTR-males (n = 8) and FLX-males (n = 8). Data are shown with individual data

points with bars representing the group means (A-F), or with squares and circles

representing respective group means (G-H). Error bars are representing SEM.

3.3 *Non-social investigation behaviors*

370   CTR- and FLX-females did not differ in time spent on ($t = -1.04$, $p = .311$, $d = 0.31$,

Figure 4A) or in the number of episodes of general activity ($t = -1.82$, $p = .090$, $d = -.0.91$).

However, FLX-females were found to spend significantly *more* time walking/running ($U =$

56.00, $z = 2.52$, $p = .025$, $r = .63$, Figure 4B) but *less* time on non-social exploration ($U =$

8.00, $z = -2.52$, $p = .025$, $r = -.63$, Figure 4C) compared to CTR-females. FLX-females were

also found to have more episodes of walking/running compared to CTR-females ($t = -4.29$, p

$= .005$, $d = -2.15$). CTR- and FLX-females did not differ in the number of non-social

exploration episodes ($t = -0.54$, $p = .693$, $d = -0.27$). Similar as for the females, no difference

in time spent on ($t = -1.69$, $p = .114$, $d = -0.85$, Figure 4D) or on number of episodes in

general activity ($t = -1.60$, $p = 0.131$, $d = -0.80$) were found between CTR- and FLX-males.

However, just as FLX-females, FLX-males spent more time walking/running than CTR-males

($t = -3.05$, $p = .045$, $d = -1.52$, Figure 4E), but there was no difference in time spent on non-

social exploration ($t = 0.06$, $p = .953$, $d = 0.03$, Figure 4F). FLX-males did not differ from

CTR-males in the number of episodes walking/running ($t = -1.61$, $p = .130$, $d = -0.80$) or non-

social exploration ($t = -0.73$, $p = .786$, $d = -0.36$).

385   We then investigated whether there were differences between CTR- and FLX-rats in

how long it took them to visit all the 10 predefined locations (see Figure 2B) of the

seminatural environment. Rats that did not visit all locations within the observation time were

given a score of 3600 (total observation time in seconds). The results indicated that FLX-rats,

both males and females, did not need significantly more or less time to visit all locations than

CTR-rats (females: $t = 1.33 = p = .212$, $d = 0.42$; males: $t = -1.15$, $p = .271$, $d = -0.57$). We

thereafter investigated how many locations the rats visited as a function of time

(1/3/5/10/20/30/60 minutes), measured on cumulative data. FLX-females were not

significantly faster at visiting the different locations compared to CTR-females (Figure 4G),

but when the different time-points were analyzed separately, they seem to have visited

significantly more locations within the first 3 minutes ($t = -2.46$, $p = .027$, $d = -1.23$)

compared to CTR-females. No difference in the number of locations visited (treatment effect:

$F(1,14) = 3.43$, $p = .085$) or in the pattern (time x treatment: $F(2.64, 36.97) = 0.39$, $p = .735$)

over time were found between the CTR- and FLX-males (Figure 4H).

3.4 Other non-social behaviors

We also looked at other relevant non-social behaviors, including non-socially passive

behaviors. The analysis revealed that there was no significant difference between CTR- and

FLX-females in time spent on ($U = 28.00$, $z = -0.42$, $p = .721$, $r = -0.11$) or in the number of

non-socially passive behaviors ($t = -0.12$, $p = .903$, $d = -0.06$). Similarly, for the male groups,

no significant difference was found for time spent on ($t = 1.62$, $p = .127$, $d = 0.81$) or in the

number of non-socially passive behaviors ($t = 0.62$, $p = .546$, $d = 0.31$)

Next, we investigated whether CTR- and FLX-rats showed different level of

anxiety/stress-related behaviors. The results revealed no significant difference between CTR-

and FLX-rats for time spent on (females: $t = 1.67$, $p = .195$, $d = 0.84$; males: $U = 37.00$, $z =$

0.53, $p = .806$, $r = .13$) or in the number of episodes (females: $t = 0.58$, $p = .693$, $d = 0.29$;

males: $t = -0.60$, $p = .860$, $d = -0.30$) self-grooming. When investigating the total time in the

open area, no significant difference was found between CTR- and FLX-rats (females: $t = -$

1.39, $p = .186$, $d = -0.70$; males: $t = -0.98$, $p = .345$, $d = -0.49$). Similarly, the treatment groups

did not differ on the total time spent in the burrow area (females: $t = 1.57$, $p = .138$, $d = 0.79$;

males: $t = 1.02$, $p = .323$, $d = 0.51$).

Figure 4 Non-social behaviors in females and males

Figure 4. The data represent the time spent (s) on general activity (A, D), walking/running (B, E), non-social exploration (C, F), and the total number of places in the SNE visited over time (G, H). All graphs show comparisons between CTR-females (n = 8) and FLX-females (n = 8) or between CTR-males (n = 8) and FLX-males (n = 8). Data are shown with individual data points with bars representing the group means (A-F), or with squares and circles representing respective group means (G-H). Error bars are representing SEM.

*p < 0.05

3.5 Behavioral adaption over time

Finally, we were interested to see whether the treatment groups adapted differently to the novel physical and social environment, and thus, whether the differences in behavior between the groups were stable over time. We therefore divided the dataset into six 10-minutes time-bins and assessed the differences between CTR- and FLX-rats on social and non-social behaviors over the course of the observation period.

The repeated measure analysis revealed that FLX-females and FLX-males did not show a significantly different pattern of time spent on socially active behaviors, compared to CTR-females (time-bin x treatment: $F(5,70) = 0.26$, $p = .932$, $\eta_p^2 = .02$, Figure 5A) or CTR-males (time-bin x treatment: $F(5,70) = 0.51$, $p = .765$, $\eta_p^2 = .04$, Figure 5B) respectively. Similarly, when looking at the frequency of socially active behaviors, no interaction between time-bin and treatment was found for female ($F(5,70) = 0.63$, $p = .675$, $\eta_p^2 = .04$) or male rats ($F(5,70) = 0.99$, $p = .431$, $\eta_p^2 = .07$).

For time spent on walking/running, no differences as a function of time were found between the CTR- and FLX-rats for females ($F(5,70) = 0.63$, $p = .679$, $\eta_p^2 = .04$) or males ($F(2.64. 36.92) = 0.69$, $p = .634$, $\eta_p^2 = .05$), meaning that the increase in walking/running was present during the whole course of the hour and was most pronounced during the first 10- ($t = -2.77$, $p = .015$, $d = -1.38$) and 30-minutes ($U = 59.00$, $z = 2.84$, $p = .003$, $r = .71$) in FLX-females, and during the first 40- ($t = -3.58$, $p = .003$, $d = -1.79$) and 50-minutes ($t = -2.56$, $p = .023$, $d = -1.28$) in FLX-males, compared to CTR-animals. Similar results were found when analyzing the frequency of walking/running (females: $F(5,70) = 0.88$, $p = .498$, $\eta_p^2 = .06$; males: $F(2.85, 39.88) = 0.82$, $p = .483$, $\eta_p^2 = .06$). In term of non-social exploration, neither FLX-females ($F(5,70) = 0.84$, $p = .529$, $\eta_p^2 = .06$) nor FLX-males ($F(2.87, 40.20) = 0.47$, $p = .697$, $\eta_p^2 = .03$) showed a significant different pattern of time spent on exploration compared to their control group. Thus, FLX-males did not differ from CTR-males throughout the different time points during the observation period. FLX-females other the other hand, scored lower than CTR-females during the whole hour, but most prominently in the first 10- ($t = 3.03$, $p = .009$, $d = 1.52$), 20- ($t = 4.38$, $p = < .001$, $d = 2.19$) and 30- minutes ($U = 12.00$, $z = -2.10$, $p = .038$, $r = -.53$). Similar results were revealed for the frequency of non-social exploration (females: $F(5,70) = 0.23$, $p = .948$, $\eta_p^2 = .02$; males: $F(5,70) = 1.76$, $p = .132$, $\eta_p^2 = .11$).

462 **Table 3. Summary of main findings**

No effects were found between CTR and FLX rats on:

- time spent on and number of social investigation behaviors
- time spent on and number of social passive behaviors
- time spent on and number of conflict behaviors
- latency to approach new conspecifics
- time spent generally active

FLX-female were found to (compared to CTR-females):

- spent more time and more often walking/running, mainly during the first 30 minutes
- spent more time on non-social exploration, mainly during the first half an hour
- visited more places within the first 3 minutes

FLX-males were found to (compared to CTR-males):

- spent more time walking/running, mainly during the last half an hour

Figure 5 Social and non-social investigation behaviors measured as a function of time.

Figure 5. The data represent the time spent (s) on different behaviors as a function of time

measured every 10 minutes. The graphs show socially active behaviors (A, B),

walking/running (C, D), and non-social exploration (E, F). All graphs show comparisons

between CTR-females (n = 8) and FLX-females (n = 8) or between CTR-males (n = 8) and

FLX-males (n = 8). Squares and circles represent respective group means, error bars

representing ± SEM.*p < 0.05, #p < 0.06

4. Discussion

In our study, we investigated how perinatal fluoxetine exposure affects adult social

and non-social investigation behaviors in a novel seminatural environment with unfamiliar

conspecifics. Our findings show that perinatal fluoxetine exposure does not induce alterations

on social investigation behaviors and strategies when introduced to a novel seminatural

environment and unknown conspecifics. However, perinatal fluoxetine exposure was found to

affect non-social investigation behaviors. More specifically, perinatal fluoxetine exposed

female and male rats showed increased locomotor activity (in terms of walking/running),

while perinatal fluoxetine exposed females showed decreased non-social exploration.

Furthermore, it was demonstrated that the observed differences maintained throughout the

whole observation.

4.1 Social behaviors

The first question we investigated was whether social investigation behaviors,

operationalized as active social behaviors (sniffing and grooming other rats) and latency to

meet all other cohort members, would be affected by perinatal SSRI exposure. The ability to

interact in line with social norms is crucial in everyday life, and deviant social behavior in the

initial phase of contact can make it difficult to establish social relationships. The results in

this study revealed no differences between CTR- and FLX-rats on the total time spent on, or

the number of, active social behaviors. Previous findings from the same experiment showed

that FLX-females, but not FLX-males, showed a tendency toward decreased active social

behaviors (Houwing et al. 2019a), which was not present after naturally occurring aggressive

encounters (Heinla et al. 2020). Nevertheless, in those studies, behaviors were observed after

the rats had already been housed together in the seminatural environment for several days,

and thus were familiar with each other. The effect of fluoxetine on social behaviors might

have different outcomes depending on whether the rats are interacting with familiar or

unfamiliar partners (Gemmel et al. 2019). In the present study, the rats were observed during

the first hour after introduction to the seminatural environment, allowing us to investigate

how the rats encounter the first social situations before knowing each other. It should be

noted, however, that the combination of a novel environment together with novel

conspecifics, might give different results than exposure to only one of the novelties. The use

of a novel environment may have masked differences in active social behaviors, since such

differences have been found in previous studies where animals were observed in a familiar

environment (Heinla et al. 2020; Houwing et al. 2019a).

509   We also measured how long it took the rats to meet the other cohort members after

being introduced to the novel environment. Such latency times could indicate whether the rats

have different interests in approaching other rats. Lack of social interest is a relevant trait to

examine since such symptoms commonly appear in various mental and neurodevelopmental

disorders (Barkus and Badcock 2019). However, the results did not reveal any differences in

latencies to meet conspecifics between CTR- and FLX-rats. From our findings, we conclude

that perinatal SSRI exposure does not affect social investigation behavior and strategies

during the first hour after introduction to a novel environment with unfamiliar conspecifics.

517   We further investigated whether SSRI exposure leads to behavioral alterations in other

aspects of social behaviors, such as social passive behaviors and conflict behaviors. The

results revealed no difference in passive social behavior between FLX-rats and CTR-rats.

Furthermore, neither FLX-females nor FLX-males differed from CTR-rats in terms of conflict

behavior. However, conflict behavior was not frequently occurring in our experiment. The

Wistar strain is generally known to exhibit little aggressive behavior compared to other strains

(Koolhaas et al. 2013). In addition, the experiment was not designed to trigger aggressive

behavior as competition for food, water or mating partners were not necessary.

4.2 Non-social behaviors

527   Next, we investigated whether perinatal SSRI exposure would affect non-social

investigation behaviors, operationalized as locomotor activity (walking/running), non-social

exploration and latency to visit all locations of the environment. We found that both FLX-

females and FLX-males spent more time on locomotor activity compared to control rats. The

display of subtle changes in motor development and in motor movement has also been found

in children exposed to SSRIs in (Casper et al. 2003). However, FLX-female rats also visited

more locations of the seminatural environment within the first 3 minutes after entrance

compared to CTR-females. This could indicate that perinatal SSRI exposure leads to an

increased interest to investigate paths and locations. Contrary to our findings, a recent meta-

analysis found evidence for *reduced* activity in developmentally SSRI exposed rats, as mostly

measured by total distance moved (Ramsteijn et al. 2020). Although we did not measure total

distance per se, it is reasonable to assume that total distance is related to total time spent

walking/running in the seminatural environment. Nevertheless, the meta-analysis is mainly

based on studies measuring activity in simplified open field boxes. Such set-ups allow the rats

to perceive the whole environment without necessarily having to move their bodies. We could

therefore assume that an increased interest to investigate locations and paths would only be

observable in situations where walking/running (movement) is needed to investigate the

environment. In addition, in the current environment, more rats were present leading to the

assumption that the odors and sounds from others may also elicit extra movement, making our

set-up more reliable to study the effects of perinatal fluoxetine exposure on a measure as

locomotor activity reflecting alterations in interest to investigate novel paths and locations.

With this in mind, we would expect the differences between FLX- and CTR-rats to disappear

(or diminish) when the animals get familiar with their surroundings and conspecifics.

Interestingly, previous studies from our research group that analyzed other relevant data from

the same experiment did indeed find no differences on locomotor activity between FLX- and

CTR- rats after the rats were already familiarized with the environment (Heinla et al. 2020;

Houwing et al. 2019a). This suggests that the current findings of increased locomotor activity

in FLX-rats are related to the introduction to a novel environment, and not the complexity of

the environment on itself.

We also found that FLX-females, but not FLX-males, spent less time on non-social

exploration than control rats, meaning they were sniffing less on objects (e.g. shelters,

wooden sticks) and specific elements in the environment (e.g. walls, the ground). This is in

line with previous findings from day 4 and day 7 in the same experiment (Houwing et al.

2019a), where reduced non-social exploration was found in FLX-females, but not in FLX-

males. However, since in the present study males and females were not compared directly,

any conclusion on the relative difference between the sexes can't be drawn. Other studies

have also reported reduced non-social explorative behaviors in SSRI exposed rats (Ansorge et

al. 2004; Karpova et al. 2009; Rebello et al. 2014; Sarkar et al. 2014; Simpson et al. 2011;

Zohar et al. 2016). Although we have shown that FLX-females seem to have increased

interest to explore paths and locations, shown by increased locomotor activity, our findings

also indicate that perinatal SSRI exposure in females leads to reduced interest to investigate

objects and other specific elements in the environment. Although the findings might seem

contradictive at first sight, locomotor activity and non-social exploration could possibly serve

different purposes. As locomotor activity could measure the interest to get quickly familiar

with the whole environment as a kind of screening behavior, non-social exploration reflects a

more detailed and accurate investigation of the environment. Therefore, we suggest that

perinatal SSRI exposure alters the strategy the animals use to investigate a novel environment

leading to a quicker, but less detailed investigation of novel environments. Previous studies

have found an association between perinatal SSRI exposure and diagnosis of attention-deficit

hyperactivity disorder (ADHD; Man et al, 2018). It could be speculated that the investigation

strategy observed in FLX-rats could reflect ADHD-related symptoms, such as often failing to

give close attention to details and disliking tasks requiring sustained mental effort (American

Psychiatric Association, 2013).

We further investigated other non-social behaviors such as anxiety/stress-related

behaviors. We did not find any difference between CTR- and FLX-rats on anxiety/stress-

related behaviors. Another subset of data from this experiment using the same cohorts of

animals found that white-noise exposure induced increased self-grooming in FLX-males

(Houwing et al. 2019a), which was explained as an altered stress-coping behavior. As

introduction to a new environment can be considered a stressful situation, we expected to

observe a similar increase in self-grooming behavior in FLX-males in the present study.

However, no differences were found between CTR- and FLX- rats on self-grooming behavior.

Moreover, no differences were found on the amount of time spent in the open area, as

measure for changes in anxiety-related behavior. Altogether, this makes us to conclude that

perinatal SSRI exposure does not affect anxiety/stress-related behavior during the first hour of

exposure to a novel environment with unfamiliar conspecifics.

4.3 Behavioral adaption over time

The last question we investigated was whether perinatal SSRI exposed rats adapt

differently to unfamiliarity (both environmental and socially) than their non-exposed

conspecifics. Therefore, we split the observational data into six 10-minute time-bins in order

to look at behavioral changes over time. As part of the familiarization process to a new

environment, we generally expected to see adjustments in behavior during the first hour, such

as decrease in general activity (Wilkinson et al. 2006). However, our main subject of interest

was whether perinatal SSRI exposed rats adjusted their behavior in a different manner than

controls.

Our results revealed that SSRI exposed animals adapted similarly to the novel

environment as control animals. As discussed, FLX-females spent less time exploring objects

and the physical environment, whereas both FLX-males and FLX-females spent more time on

locomotor activity compared to CTR-rats. Those differences remained relatively stable

throughout the first hour, meaning that FLX- and CTR- rats behaved differently, but adapted

similarly to the novel environment over time (the increased locomotor activity remained

higher during the full course of the observed hour). We conclude that perinatal fluoxetine

exposed rats do not adapt their behaviors differently than controls during the first hour after

introduction to the novel environment, instead the changes in non-social investigation

behavior remain stable over time.

5. Conclusion

In summary, our data showed that perinatal SSRI exposure alters aspects of non-social

investigation behaviors when introduced to a novel environment with unfamiliar conspecifics,

but did not alter social investigation behaviors. Both FLX-males and FLX-females showed a

higher amount of locomotor activity, while FLX-females visited more locations within the

first three minutes, and spent less time exploring objects and specific elements in the physical

environment. Perinatal fluoxetine exposure did not affect social behavior or how the animals

adapted to the unfamiliar seminatural environment over time. Altogether, we conclude that

perinatal SSRI exposure alters non-social investigation, to a quicker and less detailed strategy,

when exposed to a novel environment, and that the alteration is most pronounced in females.

Acknowledgements

Financial support was received from Helse Nord #PFP1295-16, Norway. We also

would like to thank Ragnhild Osnes, Carina Sørensen, Nina Løvhaug, Katrine Harjo, and

Remi Osnes for their excellent care of the animals.

References

American Psychiatric Association. (2013). Diagnostic and statistical manual of mental disorders (5th ed.). 10.1176/appi.books.9780890425596

Andrade SE, Raebel MA, Brown J, Lane K, Livingston J, Boudreau D, Rolnick SJ, Roblin D, Smith DH, Willy ME, Staffa JA, Platt R (2008) Use of antidepressant medications during pregnancy: a multisite study. Am J Obstet Gynecol 198.

Ansorge MS, Zhou M, Lira A, Hen R, Gingrich JA (2004) Early-life blockade of the 5-HT transporter alters emotional behavior in adult mice. Science 306: 879-81.

Azmitia EC (2001) Modern views on an ancient chemical: serotonin effects on cell proliferation, maturation, and apoptosis. Brain Res Bull 56: 413-24.

Barkus E, Badcock JC (2019) A Transdiagnostic Perspective on Social Anhedonia. Front Psychiatry 10: 216.

Bennett HA, Einarson A, Taddio A, Koren G, Einarson TR (2004) Prevalence of depression during pregnancy: systematic review. Obstetrics and gynecology 103: 698-709.

Brown AS, Gyllenberg D, Malm H, McKeague IW, Hinkka-Yli-Salomaki S, Artama M, Gissler M, Cheslack-Postava K, Weissman MM, Gingrich JA, Sourander A (2016) Association of Selective Serotonin Reuptake Inhibitor Exposure During Pregnancy With Speech, Scholastic, and Motor Disorders in Offspring. JAMA Psychiatry 73: 1163-1170.

Casper RC, Fleisher BE, Lee-Ancajas JC, Gilles A, Gaylor E, DeBattista A, Hoyme HE (2003) Follow-up of children of depressed mothers exposed or not exposed to antidepressant drugs during pregnancy. J Pediatr 142: 402-8.

Charlton RA, Jordan S, Pierini A, Garne E, Neville AJ, Hansen AV, Gini R, Thayer D, Tingay K, Puccini A, Bos HJ, Nybo Andersen AM, Sinclair M, Dolk H, de Jong-van den Berg LT (2015) Selective serotonin reuptake inhibitor prescribing before, during and after pregnancy: a population-based study in six European regions. BJOG : an international journal of obstetrics and gynaecology 122: 1010-20.

Chu X, Agmo A (2014) Sociosexual behaviours in cycling, intact female rats (Rattus norvegicus) housed in a seminatural environment. Behaviour 151: 1143-1184.

Dunkel Schetter C (2011) Psychological science on pregnancy: stress processes, biopsychosocial models, and emerging research issues. Annual review of psychology 62: 531-58.

El Marroun H, White T, Verhulst FC, Tiemeier H (2014) Maternal use of antidepressant or anxiolytic medication during pregnancy and childhood neurodevelopmental outcomes: a systematic review. Eur Child Adolesc Psychiatry 23: 973-92.

Gaspar P, Cases O, Maroteaux L (2003) The developmental role of serotonin: news from mouse molecular genetics. Nature reviews Neuroscience 4: 1002-12.

Gemmel M, De Lacalle S, Mort SC, Hill LA, Charlier TD, Pawluski JL (2019) Perinatal fluoxetine has enduring sexually differentiated effects on neurobehavioral outcomes related to social behaviors. Neuropharmacology 144: 70-81.

Gemmel M, Hazlett M, Bogi E, De Lacalle S, Hill LA, Kokras N, Hammond GL, Dalla C, Charlier TD, Pawluski JL (2017) Perinatal fluoxetine effects on social play, the HPA system, and hippocampal plasticity in pre-adolescent male and female rats: Interactions with pre-gestational maternal stress. Psychoneuroendocrinology 84: 159-171.

Glover ME, Clinton SM (2016) Of rodents and humans: A comparative review of the neurobehavioral effects of early life SSRI exposure in preclinical and clinical research. Int J Dev Neurosci 51: 50-72.

Goodman SH (2007) Depression in mothers. Annu Rev Clin Psychol 3: 107-35.

Hegstad J, Huijgens PT, Houwing DJ, Olivier JDA, Heijkoop R, Snoeren EMS (2020) Female rat sexual behavior is unaffected by perinatal fluoxetine exposure. Psychoneuroendocrinology 120: 104796.

Heinla I, Heijkoop R, Houwing DJ, Olivier JDA, Snoeren EMS (2020) Third-party prosocial behavior in adult female rats is impaired after perinatal fluoxetine exposure. Physiol Behav 222: 112899.

Hermansen TK, Roysamb E, Augusti EM, Melinder A (2016) Behavior and inhibitory control in children with prenatal exposure to antidepressants and medically untreated depression. Psychopharmacology (Berl) 233: 1523-35.

Houwing DJ, Esquivel-Franco DC, Ramsteijn AS, Schuttel K, Struik EL, Arling C, de Boer SF, Olivier JDA (2020) Perinatal fluoxetine treatment and dams' early life stress history have opposite effects on aggressive behavior while having little impact on sexual behavior of male rat offspring. Psychopharmacology (Berl) 237: 2589-2600.

Houwing DJ, Heijkoop R, Olivier JDA, Snoeren EMS (2019a) Perinatal fluoxetine exposure changes social and stress-coping behavior in adult rats housed in a seminatural environment. Neuropharmacology 151: 84-97.

Houwing DJ, Staal L, Swart JM, Ramsteijn AS, Wohr M, de Boer SF, Olivier JDA (2019b) Subjecting Dams to Early Life Stress and Perinatal Fluoxetine Treatment Differentially Alters Social Behavior in Young and Adult Rat Offspring. Frontiers in neuroscience 13: 229.

Karpova NN, Lindholm J, Pruunsild P, Timmusk T, Castren E (2009) Long-lasting behavioural and molecular alterations induced by early postnatal fluoxetine exposure are restored by chronic fluoxetine treatment in adult mice. Eur Neuropsychopharmacol 19: 97-108.

Khatri N, Simpson KL, Lin RC, Paul IA (2014) Lasting neurobehavioral abnormalities in rats after neonatal activation of serotonin 1A and 1B receptors: possible mechanisms for serotonin dysfunction in autistic spectrum disorders. Psychopharmacology (Berl) 231: 1191-200.

Kim J, Riggs KW, Misri S, Kent N, Oberlander TF, Grunau RE, Fitzgerald C, Rurak DW (2006) Stereoselective disposition of fluoxetine and norfluoxetine during pregnancy and breast-feeding. Br J Clin Pharmacol 61: 155-63.

Kiryanova V, Dyck RH (2014) Increased aggression, improved spatial memory, and reduced anxiety-like behaviour in adult male mice exposed to fluoxetine early in life. Developmental neuroscience 36: 396-408.

Klinger G, Frankenthal D, Merlob P, Diamond G, Sirota L, Levinson-Castiel R, Linder N, Stahl B, Inbar D (2011) Long-term outcome following selective serotonin reuptake inhibitor induced neonatal abstinence syndrome. Journal of perinatology : official journal of the California Perinatal Association 31: 615-20.

Koolhaas JM, Coppens CM, de Boer SF, Buwalda B, Meerlo P, Timmermans PJ (2013) The resident-intruder paradigm: a standardized test for aggression, violence and social stress. Journal of visualized experiments : JoVE: e4367.

Kristensen JH, Ilett KF, Hackett LP, Yapp P, Paech M, Begg EJ (1999) Distribution and excretion of fluoxetine and norfluoxetine in human milk. Br J Clin Pharmacol 48: 521-7.

Lundmark J, Reis M, Bengtsson F (2001) Serum concentrations of fluoxetine in the clinical treatment setting. Ther Drug Monit 23: 139-47.

Lupattelli A, Wood M, Ystrom E, Skurtveit S, Handal M, Nordeng H (2018) Effect of Time-Dependent Selective Serotonin Reuptake Inhibitor Antidepressants During Pregnancy on Behavioral, Emotional, and Social Development in Preschool-Aged Children. J Am Acad Child Adolesc Psychiatry 57: 200-208.

Malm H, Brown AS, Gissler M, Gyllenberg D, Hinkka-Yli-Salomaki S, McKeague IW, Weissman M, Wickramaratne P, Artama M, Gingrich JA, Sourander A (2016) Gestational Exposure to Selective Serotonin Reuptake Inhibitors and Offspring Psychiatric Disorders: A National Register-Based Study. J Am Acad Child Adolesc Psychiatry 55: 359-66.

Mitchell AA, Gilboa SM, Werler MM, Kelley KE, Louik C, Hernandez-Diaz S, Stud NBDP (2011) Medication use during pregnancy, with particular focus on prescription drugs: 1976-2008. Am J Obstet Gynecol 205.

Molenaar NM, Bais B, Lambregtse-van den Berg MP, Mulder CL, Howell EA, Fox NS, Rommel AS, Bergink V, Kamperman AM (2020) The international prevalence of antidepressant use before, during, and after pregnancy: A systematic review and meta-analysis of timing, type of prescriptions and geographical variability. J Affect Disorders 264: 82-89.

Muller CL, Anacker AMJ, Veenstra-VanderWeele J (2016) The serotonin system in autism spectrum disorder: From biomarker to animal models. Neuroscience 321: 24-41.

Noorlander CW, Ververs FF, Nikkels PG, van Echteld CJ, Visser GH, Smidt MP (2008) Modulation of serotonin transporter function during fetal development causes dilated heart cardiomyopathy and lifelong behavioral abnormalities. PLoS One 3: e2782.

Olivier JD, Blom T, Arentsen T, Homberg JR (2011a) The age-dependent effects of selective serotonin reuptake inhibitors in humans and rodents: A review. Progress in neuro-psychopharmacology & biological psychiatry 35: 1400-8.

Olivier JD, Valles A, van Heesch F, Afrasiab-Middelman A, Roelofs JJ, Jonkers M, Peeters EJ, Korte-Bouws GA, Dederen JP, Kiliaan AJ, Martens GJ, Schubert D, Homberg JR (2011b) Fluoxetine administration to pregnant rats increases anxiety-related behavior in the offspring. Psychopharmacology (Berl) 217: 419-32.

Rampono J, Proud S, Hackett LP, Kristensen JH, Ilett KF (2004) A pilot study of newer antidepressant concentrations in cord and maternal serum and possible effects in the neonate. Int J Neuropsychopharmacol 7: 329-34.

Ramsteijn AS, Van de Wijer L, Rando J, van Luijk J, Homberg JR, Olivier JDA (2020) Perinatal selective serotonin reuptake inhibitor exposure and behavioral outcomes: A systematic review and meta-analyses of animal studies. Neurosci Biobehav Rev 114: 53-69.

Rebello TJ, Yu Q, Goodfellow NM, Caffrey Cagliostro MK, Teissier A, Morelli E, Demireva EY, Chemiakine A, Rosoklija GB, Dwork AJ, Lambe EK, Gingrich JA, Ansorge MS (2014) Postnatal day 2 to 11 constitutes a 5-HT-sensitive period impacting adult mPFC function. J Neurosci 34: 12379-93.

Rodriguez-Porcel F, Green D, Khatri N, Harris SS, May WL, Lin RC, Paul IA (2011) Neonatal exposure of rats to antidepressants affects behavioral reactions to novelty and social interactions in a manner analogous to autistic spectrum disorders. Anatomical record 294: 1726-35.

Sarkar A, Chachra P, Vaidya VA (2014) Postnatal Fluoxetine-Evoked Anxiety Is Prevented by Concomitant 5-HT2A/C Receptor Blockade and Mimicked by Postnatal 5-HT2A/C Receptor Stimulation. Biological psychiatry 76: 858-868.

Simpson KL, Weaver KJ, de Villers-Sidani E, Lu JY, Cai Z, Pang Y, Rodriguez-Porcel F, Paul IA, Merzenich M, Lin RC (2011) Perinatal antidepressant exposure alters cortical network function in rodents. Proc Natl Acad Sci U S A 108: 18465-70.

Snoeren EM, Antonio-Cabrera E, Spiteri T, Musatov S, Ogawa S, Pfaff DW, Agmo A (2015) Role of Oestrogen alpha Receptors in Sociosexual Behaviour in Female Rats Housed in a Seminatural Environment. J Neuroendocrinol 27: 803-18.

Svirsky N, Levy S, Avitsur R (2016) Prenatal exposure to selective serotonin reuptake inhibitors (SSRI) increases aggression and modulates maternal behavior in offspring mice. Dev Psychobiol 58: 71-82.

Wilkinson JL, Herrman L, Palmatier MI, Bevins RA (2006) Rats' novel object interaction as a measure of environmental familiarity. Learn Motiv 37: 131-148.

Woody CA, Ferrari AJ, Siskind DJ, Whiteford HA, Harris MG (2017) A systematic review and meta-regression of the prevalence and incidence of perinatal depression. J Affect Disord 219: 86-92.

Zimmerberg B, Germeyan SC (2015) Effects of neonatal fluoxetine exposure on behavior across development in rats selectively bred for an infantile affective trait. Dev Psychobiol 57: 141-52.

Zohar I, Shoham S, Weinstock M (2016) Perinatal citalopram does not prevent the effect of prenatal stress on anxiety, depressive-like behaviour and serotonergic transmission in adult rat offspring. Eur J Neurosci 43: 590-600.