Effects of Perinatal Fluoxetine Exposure on Novelty-induced Social and Non-Social

Investigation Behaviors in a Seminatural Environment

Ole Christian Sylte[1,*], Jesper Solheim Johansen[1,*], Indrek Heinla[1], Danielle J Houwing[1,2],

Jocelien DA Olivier[2], Roy Heijkoop[1], and Eelke MS Snoeren[1,3]

[1] Department of Psychology, UiT the Arctic University of Norway, Norway

[2] Department of Neurobiology, Groningen Institute for Evolutionary Life Sciences, University

of Groningen, the Netherlands

[3] Regional Health Authority of North Norway

[*] These authors contributed equally.

22   Corresponding concerning this article should addressed to Eelke Snoeren, Department of

23   Psychology, UiT the Arctic University of Norway, 9037 Tromsø, Norway. E-mail address:

24   eelke.snoeren@uit.no

Abstract

Selective serotonin reuptake inhibitors (SSRIs) are increasingly prescribed as medication for

various affective disorders during pregnancy. SSRIs cross the placenta and affect serotonergic

neurotransmission in the fetus, but the neurobehavioral consequences for the offspring remain

largely unclear. Recent rodent research has linked perinatal SSRI exposure to alterations in

both social and non-social aspects of behavior. However, this research has mainly focused on

behavior within simplified environments. The current study investigates the effects of

perinatal SSRI exposure on social and non-social investigation behaviors of adult rat

offspring upon introduction to a novel seminatural environment with unknown conspecifics.

During the perinatal period (gestational day 1 until postnatal day 21), rat dams received daily

treatment with either an SSRI (fluoxetine, 10 mg/kg) or vehicle. Adult male and female

offspring were observed within the first hour after introduction to a seminatural environment.

The results showed that perinatal fluoxetine exposure altered aspects of non-social

investigation behaviors, while not altering social investigation behaviors. More specific, both

fluoxetine exposed males and females spent more total time on locomotor activity than

controls. Furthermore, fluoxetine exposed females spent less time exploring objects and

specific elements in the environment. The data suggest that perinatal exposure to SSRIs leads

to a quicker, less detailed investigation strategy in novel environments, and that the alteration

is mostly pronounced in females.

Keywords: antidepressants, fluoxetine, perinatal, behavior, social, rats, seminatural

environment, SSRI

1. Introduction

A considerable number of women experience depression or other mental disorders

during pregnancy. Approximately 1 in 10 pregnant women fulfill the DSM-5 diagnostic

criteria for major depressive disorder (Bennett et al., 2004; Woody et al., 2017). In treatment

of maternal depression and anxiety, selective serotonin reuptake inhibitors (SSRIs) are the

most frequently prescribed class of drugs, as it has been considered relatively safe for both

mother and child. The prescription rate of SSRIs to pregnant women has increased

tremendously in the last decades (Mitchell et al., 2011), and recent estimates suggest a

worldwide prevalence of 3% (Molenaar et al., 2020) with significant geographically

differences (Andrade et al., 2008; Charlton et al., 2015). Consequently, hundreds of thousands

of babies exposed to SSRIs during early development are born every year. Despite the

widespread use, we have limited knowledge on whether SSRI exposure during the early

stages of brain development can lead to altered long-term behavioral outcomes, such as social

and non-social behaviors.

Antidepressants, such as SSRIs, reach the fetus by crossing the placenta and are

present in breast milk (Kristensen et al., 1999; Rampono et al., 2004). Thus, children can

potentially be exposed to SSRIs during the entire perinatal period (Kim et al., 2006;

Noorlander et al., 2008). SSRIs inhibit the function of the serotonin-reuptake transporter

(SERT or 5-HTT), which leads to an accumulation of 5-HT in the synaptic cleft. This in turn

increases the magnitude and duration of 5-HT activity at pre- and post-synaptic 5-HT

receptors. In the adult brain, 5-HT acts mainly as a modulatory neurotransmitter, regulating

emotion, cognition, sleep and stress responses (Olivier et al., 2011a). However, in the

developing brain, 5-HT is widespread and acts as a neurotrophic factor regulating cell

division, differentiation, migration, and synaptogenesis (Azmitia, 2001; Gaspar et al., 2003).

Consequently, developmental SSRI exposure is suggested to affect both neurodevelopment

and later-life behaviors (Muller et al., 2016).

Previous studies in humans have shown associations between developmental SSRI

exposure and impaired social behavior (Klinger et al., 2011), increased risk of speech and

language disorders (Brown et al., 2016), and elevated levels of internalizing behavior, like

anxiety and depression (Hermansen et al., 2016; Lupattelli et al., 2018; Malm et al., 2016).

While the existing literature has mainly examined the childhood years, little is known on

whether these associations persist into adulthood. In addition, outcomes such as depression

may not emerge before a certain age and could therefore remain undiscovered.

Epidemiological research on humans, like the above-mentioned studies, are

correlational in nature, and do not necessarily imply causation. A frequent problem with

human studies is the difficulty to isolate the effects of SSRI exposure from the effects of

maternal mental health. Women using SSRIs during pregnancy are likely suffering from

depression, which itself has been shown to have negative impact on the offspring (Dunkel

Schetter, 2011; El Marroun et al., 2014; Goodman, 2007). Animal research, on the other hand,

allows to control for potential interference from confounding factors, like maternal health,

drug dose and timing of exposure. As rodent and human serotonergic development is

remarkably similar (Glover and Clinton, 2016), rodent studies can provide valuable

translational insight about how developmental SSRI exposure affects human offspring.

Rodent studies investigating the effects of developmental exposure to SSRIs have

reported alterations in different social and non-social behaviors in the offspring. In juvenile

male and female offspring, both pre- and post-natal SSRI exposure have been shown to

decrease social play behavior (Houwing et al., 2019b; Khatri et al., 2014; Olivier et al.,

2011b; Rodriguez-Porcel et al., 2011; Simpson et al., 2011). Similar tendencies have been

found in adult rats with developmental SSRI exposure leading to less social interactions

(Olivier et al., 2011b; Rodriguez-Porcel et al., 2011), or decreased interest to explore a novel

conspecific (Khatri et al., 2014; Rodriguez-Porcel et al., 2011; Simpson et al., 2011;

Zimmerberg and Germeyan, 2015). SSRI exposure can also decrease (Houwing et al., 2020),

or increase (Gemmel et al., 2017; Kiryanova and Dyck, 2014; Svirsky et al., 2016),

aggressive-like social behaviors. Furthermore, a recent meta-analysis revealed that

developmental exposure to SSRI was linked to reduced activity and explorative behaviors in

adult rats and mice (Ramsteijn et al., 2020).

Most of the animal studies, however, have used simplified rodent test set-ups which

only investigates a small fraction of all behaviors. Furthermore, these studies do not account

for the environmental and social complexity of real-world situations. To bypass these

limitations, recent studies from our research group have employed a seminatural environment

enabling rats to express all aspects of their natural behaviors (Hegstad et al., 2020; Heinla et

al., 2020; Houwing et al., 2019a). These studies showed that perinatal SSRI fluoxetine (FLX)

exposure leads to various alterations in social and non-social behaviors in a naturalistic

setting. More specifically, perinatal fluoxetine exposure was associated with an increased

amount of passive social behaviors in both males and females, but a reduction of active social

behavior, general activity (Houwing et al., 2019a), and pro-social behaviors in females

(Heinla et al., 2020). Interestingly, these studies were performed in the seminatural

environment *after* the rats were familiarized to each other and the physical environment. It is

currently unknown how social and non-social behaviors manifest directly after introduction to

a novel environment with unfamiliar conspecifics. As perinatal SSRI exposure seem to alter

stress-coping behaviors (Houwing et al., 2019a), one could hypothesize that the stressor of a

novel environment with new conspecifics could lead to more pronounced changes in social

and non-social behaviors.

The aim of the current study was to investigate if perinatal SSRI exposure alters social

and non-social investigation behaviors in a novel environment with unknown conspecifics.

We define investigation as behaviors that provides the animal with information about a novel

stimulus. More specifically, social investigation refers to when the stimulus investigated is a

conspecific, such as when sniffing and grooming others, while non-social investigation refers

to investigation of inanimate objects and environmental locations. In line with previous

studies (Heinla et al., 2020; Houwing et al., 2019a), we expected perinatal fluoxetine

exposure to show a reduction in active social behavior in in non-social investigation

(exploratory) behavior in the initial phase of the introduction to the seminatural environment.

In addition, as introduction to a new environment can be considered a stressful situation, we

also expected to observe an increase in self-grooming behavior in FLX-exposed animals.

2.   Material and Methods

The data was collected from video recordings obtained in a previously performed

experiment (Houwing et al., 2019a). The materials and methods are therefore similar to those

described previously (Hegstad et al., 2020; Heinla et al., 2020; Houwing et al., 2019a).

However, the behavioral scoring scheme was uniquely formed to the current study.

2.1 Animals and dam housing

A total of 20 Wistar rats (10 males, 10 females), weighing 200-250 grams on arrival,

were obtained from Charles River (Sulzfeld, Germany) for breeding. After arrival, same-sex

pairs were housed in Makrolon IV cages (60 x 38 x 20 cm) on a reversed 12:12 hours

light/dark cycle, in which the lights were turned on at 23.00. Temperature in the room was 21

± 1°C, and the relative humidity was 55 ± 10 %. Standard rodent food pellets (standard chow,

Special Diets Services, Witham, Essex, UK), water and nesting material were available ad

libitum. Animal care and experimental procedures were conducted in agreement with

European Union council directive 2010/63/EU. The protocol was approved by the National

Animal Research Authority in Norway.

2.2 Breeding and antidepressant treatment

Daily, all females were checked for sexual receptivity by placing them together with a

male rat for 5 minutes. When lordosis behavior was observed, they were considered in

proestrus and thus ready for breeding. The female then got placed together with a male in an

isolated Makrolon IV cage for the next 24 hours (gestational day 0). Afterwards, they returned

to their initial same-sex pairs for the first two weeks of pregnancy. From gestational day 14,

the females were placed solitarily until delivery (gestational day 21/postnatal day 0).

During the 6-week period from conception (gestational day 0) to weaning (postnatal

day 21), females received either the SSRI fluoxetine 10 mg/kg (Apotekproduksjon, Oslo,

Norway) or vehicle (methylcellulose; Sigma, St. Louis, MO, USA) daily by oral gavage. The

offspring were thus exposed to perinatal fluoxetine via the treatment of the dams (in utero and

via breast feeding). The fluoxetine treatment was prepared with tablets for human usage that

were pulverized and dissolved in sterile water (2mg/mL) and injected at a volume of 5mL/kg.

Methylcellulose powder, the non-active filling of a fluoxetine tablet, was used as control

condition. The powder was dissolved in sterile water to create a 1% solution and administered

at a volume of 5mL/kg as well. Every third day, females were weighed to ensure correct

dosage of fluoxetine/vehicle. The chosen dosage of fluoxetine was decided upon comparison

of fluoxetine blood levels of humans and animals (Lundmark et al., 2001; Olivier et al.,

2011b). When the rat dams got close to the end of pregnancy, they were checked two times a

day (09.00 and 15.00) for delivery.

2.3 Offspring housing

The offspring were housed together with their mothers until weaning (gestational day

21). After weaning, groups of two or three same-sex littermates were housed together in

Makrolon IV cages (see cage distribution in the supplemental materials). They were left

undisturbed, except for the ovariectomy (see section for Procedure) and weekly cage cleaning,

until introduction to the seminatural environment at the age of 13-18 weeks. To enable

individual recognition, ears were punched. In Figure 1, a schematic overview shows all

experimental procedures from gestational day 0 to the end of the experiment.

Figure 1 Overview of experimental procedures

Figure 1. FLX = fluoxetine, CTR = control, P = postnatal day, G = gestational day. Created

with BioRender (https://biorender.com/).

2.4 Seminatural environment

The seminatural environment (SNE; 240 x 210 x 75 cm) consisted of two parts: an

open area and a burrow system (Figure 2; (Chu and Agmo, 2014; Houwing et al., 2019a;

Snoeren et al., 2015)). Four openings (8 x 8 cm) connected the two areas. In the open area,

two partitions (40 x 75 cm) simulated natural obstacles. The burrow system consisted of

connected tunnels (width 7.6 cm, height 8 cm) and four nest boxes (20 x 20 x 20 cm).

Plexiglas covered the burrow at the height of 75 cm, while the open area remained open. A

curtain between the two parts allowed for different light settings. The burrow was left dark the

entire time. In the open area, on the other hand, light settings simulated a day-night cycle. A

lamp located 2,5 m above the floor, simulated daylight (180 lux) between 22.45 and 10.30.

From 10.30 to 11.00 the lights gradually decreased to 1 lux (simulating moonlight). The

darkness lasted until the light gradually increased from 1 to 180 lux between 22.15 and 22.45.

The whole ground of the SNE was covered with a layer (2 cm) of aspen wood chip

bedding (Tapvei, Harjumaa, Estonia). The nest boxes had 6 squares of nesting material in

each (non-woven hemp fibers, 5 x 5 fibers, 5 mm thickness, Datesend, Manchester, UK).

Three plastic shelters (15 x 16.5 x 8.5 cm, Datesend, Manchester, UK) were placed in the

open area. Additionally, 12 aspen wooden sticks (2 x 2 x 10 cm, Tapvei, Harjumaa, Estonia)

were randomly placed around in the SNE. A pile of food pellets (approx. 2 kg) and four

bottles of water were available at all time (see location in Figure 2A).

Two video cameras (Basler) were mounted on the ceiling, 2 m above the open area

(regular camera) and the burrow system (infrared camera) respectively. Media Recorder 2.5

was employed for video recordings. The data got immediately stored on an external hard

drive. The recording was manually stopped and restarted every 24 hours. The purpose was to

ensure that eventual errors only would affect one day of recorded data.

Figure 2 The Seminatural Environment

Figure 2. Illustration of the whole seminatural environment (A) and sectioning of the different

locations (B). 1 = open area close to burrow left, 2 = open area close to burrow right, 3 =

open area far away from burrow left, 4 = open area far away from burrow right, 5 = tunnels

far away from open a, 6 = tunnels close to OA, 7 = nestbox left, 8 = nestbox mid-left, 9 =

nestbox mid-right, 10 = nestbox right. Created with BioRender (https://biorender.com/).

2.5 Design of the study

Initially, five cohorts, each consisting of eight rat offspring, were placed one at the

time in the SNE. However, one day of video material was lost due to recording error, which

reduced the number of cohorts to four. A cohort consisted of 4 males and 4 females of which

each sex constituted 2 controls (CTR) and 2 fluoxetine (FLX) rats. Thus, data from this

experiment came from 8 CTR-males, 8 CTR-females, 8 FLX-males and 8 FLX-females (see

Table S2 for more details). Within a cohort, same sex rats came from different litters and were

thus unfamiliar to each other. Some rats had one sibling from the opposite sex in the same

cohort. However, these rats had been housed in different home cages since weaning.

2.6 Procedure

249   For the purpose of a previous study (Houwing et al., 2019a), the female offspring were

ovariectomized two weeks before entering the SNE in order to control their estrous cycle.

Although irrelevant for the objective of the current study, this procedure had the effect of

keeping the females in diestrus of the menstrual cycle during the observation period. Before

entering the SNE, the rats were shaved on the back and tail-marked under isoflurane

anesthesia for individual recognition (for more details, see (Houwing et al., 2019a)). All rats

were also weighed, confirming that there was no weight difference between CTR- and FLX-

rats.

257   Each cohort was placed in the SNE for 8 days. See Figure 1 for an overview of the

whole procedure. The cohorts were introduced to SNE on the first day (day 0) at 10.00 and

removed on day 8 at the same time. However, only data from the first hour was used for the

purpose of this study. All rats were again weighed after being removed from the SNE. No

difference in weight was observed between CTR- and FLX-rats. In order to remove olfactory

cues, the SNE was cleaned and bedding changed between cohorts.

2.7 Behavioral observations

265   The frequency and/or duration of several behaviors (see Table 1) were scored

manually using The Observer XT, version 12 (Noldus, Wageningen, The Netherlands). Two

observers, blinded for the animal treatment, independently scored either males or females

across all four cohorts. In addition to behavior, (1) location of the animal (see Figure 2B), (2)

whether the animal initiated the respective behavior or was respondent to it, (3) whether the

animal was in physical contact with another animal or not during the respective behavior, and

lastly, (4) ID of the interacting partner was scored. Since we were interested in observing how

the rats behaved in a novel environment with unfamiliar conspecifics, all rats were scored in

the first 60 minutes after entry to the SNE.

Table 1 Description of recorded behaviors

Behavior	Description
Walking/running	Walking or running through the environment
Chasing	Running forward in the direction of a conspecific
Non-social exploration	Exploring the environment by sniffing, usually when slowly walking or sitting still
Digging	Digging, pushing or carrying bedding/nesting/food material
Resting/immobile alone	Sitting or sleeping with minimal movement of the head without other rats in close vicinity
Resting/immobile socially	Sitting or sleeping with minimal movement of the head with at least 1 other rat on maximum 1 rat body length away
Hiding alone	Being in the shelter alone
Hiding socially	Being in the shelter with at least one other rat
Following	Walking or running in the same direction as another rat in front.
Allogrooming	Grooming any part of a conspecific's body, usually on the head or in the neck region
Sniffing anogenitally	Sniffing the anogenital region of the conspecific
Sniffing nose-to-nose	Sniffing the facial region of the conspecific
Sniffing body	Sniffing any part of the conspecifics body, except for the anogenital and facial region
Fighting	Kicking, pouncing, pushing, grapping, boxing or wrestling another rat
Nose-off	Facing another rat, usually in a tunnel, resulting in one rat moving forward and the other backing up
Self-grooming	Grooming itself
Freezing	Complete absence of movement in addition to a tense body posture
Rearing supported	Raising itself upright on its hind paws, facing a wall or an object
Rearing unsupported	Raising itself upright on its hind paws, not facing a wall or an object

277   **Table 2 Description of behavioral clusters**

Cluster	Behaviors within clusters
Socially active behaviors	Sniffing anogenitally, sniffing nose-to-nose, sniffing body, and allogrooming
General activity	Walking/running, non-social exploration
Non-socially passive behaviors	Resting alone, hiding alone
Socially passive behaviors	Hiding socially, resting socially
Conflict behaviors	Nose-off, fighting

279   **2.8 Data preparation and statistical analysis**

280   As shown in Table 2, the recorded behaviors were combined into behavioral clusters.

For each rat, we calculated the total duration and the number of events for every behavior and

behavioral cluster. This data was later divided into six 10-minute time-bins in order to analyze

behavioral changes over time. Latencies to meet the other rats, and latencies to visit the

different locations of the SNE were also noted. This data was later divided and analyzed

cumulative over the first 1, 3, 5, 10, 20, 30, and 60 minutes. In this study, we operationalized

social investigation behaviors as the cluster "socially active behaviors" and the latencies to

meet all other rats, whereas non-social investigation behaviors were operationalized as the

cluster "general activity" and latencies to visit all the locations (See Figure 2B).

289   Normality of data was determined with Shapiro-Wilks tests. Data with $p < .05$ was

analyzed non-parametrically. Simple group comparisons were performed with either a student

t-test or the non-parametric Mann-Whitney U test. Repeated measures ANOVA was used

when the behaviors were analyzed over time. In cases the Mauchly's test indicated violation

of sphericity from the ANOVA output, the degrees of freedom were corrected using

Greenhouse-Geisser estimates of sphericity. To correct for multiple comparisons, the

Benjamini-Hochberg procedure was performed on all significant results together with a

predetermined set of variables (sniffing, self-grooming, non-social exploration, conflict

behaviors). All tests reported were done 2-tailed.

2.9 Statement Open Science Framework (OSF)

The design of our study was preregistered on OSF on the 25th of March 2019

(https://osf.io/m87j5). There were no changes in analysis, except that we did not use the

originally planned additional control group. As stated at OSF, the planned control group was

not suitable, because it consisted of aged rats and had a different composition in number of

rats (7 versus 8). We therefore concluded that these differences would make it impossible to

compare the cohorts of the current study.

3 Results

From the behavioral scoring, we obtained a lot of data. A complete overview of all

behaviors can be found in Table S3 and S4.

3.1 Fluoxetine exposure does not influence social behaviors in a novel environment

We first investigated whether perinatal exposure to fluoxetine (FLX) affects social

behaviors when the animals are habituating to a novel environment. We therefore examined

social investigation behaviors, as in how the rats investigate unknown conspecifics. In

addition, we measured other relevant forms of social behaviors like passive and conflict

behaviors during the first period after introduction.

3.1.1 Social investigation behaviors

The data analysis revealed that CTR- and FLX-females did not differ in time spent on

($t = -1.04$, $p = .315$, $d = -0.52$, Figure 3A) or number of episodes ($t = -1.04$, $p = .318$, $d = -$

0.52) performing socially active behaviors. When looking separately at the different

behavioral components constituting the cluster (see Table 2), CTR- and FLX-rats did not

differ on any other behavioral components constituting the clusters relevant to social

behaviors (socially active behaviors, socially passive behaviors and conflict behaviors). No

difference was found between CTR- and FLX-males for socially active behaviors in total time

($t = 0.95$, $p = .356$, $d = 0.48$, Figure 3D) or on number of episodes ($t = 0.103$, $p = .919$, $d =$

0.05).

Although the treatment groups did not differ in the amount of socially active

behaviors, it could still be the case that the groups had different interest in meeting other rats.

To investigate this possibility, we first looked at the latencies to when the rats had met all

seven other cohort-members. The data analysis showed that there was no significant

difference in latency to meet all cohort-members between CTR- and FLX-rats for females ($t =$

0.84, $p = .418$, $d = 0.42$) or males ($U = 24.00$, $z = -0.84$, $p = .422$, $r = -.21$). We subsequently

measured how many cohort members the rats had met as a function of time. CTR- and FLX-

rats were compared on cumulative data measured at 1/3/5/10/20/30/60 minutes. For FLX-

females, there were no significant differences in the number of rats met (treatment effect:

$F(1,14) = 0.05$, $p = .821$) or in the pattern of rats met (timepoints x treatment: $F(1.73, 24.24)$

$= 0.28$, $p = .725$) over time compared to CTR-females (Figure 3G). Similarly, CTR- and

FLX-males did not differ in the number of rats met across all timepoints (treatment effect:

$F(1,14) = 0.49$, $p = .492$) or in the pattern of rats met over time (timepoints x treatment:

$F(2.05, 28.74) = 0.59$, $p = .563$, Figure 3H).

3.1.2 Other social behaviors

We also investigated some other social behaviors, such as socially passive behaviors

and conflict behaviors. No difference was found between CTR- and FLX-females in total time

($U = 33.00$, $z = 1.05$, $p = 1$, $r = .03$, Figure 3B) or number of episodes being socially passive (t

= -0.28, p = .784, d = -0.14). Furthermore, CTR- and FLX-females spent a similar amount of

time (t = 0.03, p = .978, d = 0.01, Figure 3C) and episodes (t = -0.40, p = .692, d = -0.20) in

conflict with other rats. Similarly, for males, no differences were found for time spent on

social passive behavior (U = 41.00, z = 0.95, p = .382, r = .24, Figure 3E), episodes of social

passive behavior (t = 1.48, p = .161, d = 0.74), time spent on conflict behavior (t = -0.03, p =

.655, d = -0.02, Figure 3F), or episodes in conflict behavior (U = 42.00, z = 1.05 p = .786, r =

.26).

Figure 3 Social behaviors in females and males

Figure 3. The data represent the time spent (s) on socially active behaviors (A, D), socially

passive behaviors (B, E), conflict behaviors (C, F), and the total number of rats met over time

(G, H). All graphs show comparisons between CTR-females (n = 8) and FLX-females (n = 8)

or between CTR-males (n = 8) and FLX-males (n = 8). Data are shown with individual data

points with bars representing the group means (A-F), or with squares and circles

representing respective group means (G-H). Error bars are representing SEM.

3.2 Fluoxetine exposure influence how the animals investigate a novel environment

We next investigated whether perinatal fluoxetine exposure alters non-social

investigation behaviors: how the animals investigate objects and the physical environment.

We also examined other relevant non-social behaviors, like passive and anxiety/stress-related

behavior, during the first hour after introduction to the environment.

3.2.1 Non-social investigation behaviors

CTR- and FLX-females did not differ in time spent on ($t = -1.04$, $p = .311$, $d = 0.31$,

Figure 4A) or in the number of episodes of general activity ($t = -1.82$, $p = .090$, $d = -.0.91$).

However, FLX-females were found to spend significantly *more* time walking/running ($U =$

56.00, $z = 2.52$, $p = .025$, $r = .63$, Figure 4B) but *less* time on non-social exploration ($U =$

8.00, $z = -2.52$, $p = .025$, $r = -.63$, Figure 4C) compared to CTR-females. FLX-females were

also found to have more episodes of walking/running compared to CTR-females ($t = -4.29$, p

$= .005$, $d = -2.15$). CTR- and FLX-females did not differ in the number of non-social

exploration episodes ($t = -0.54$, $p = .693$, $d = -0.27$). Similar as for the females, no difference

in time spent on ($t = -1.69$, $p = .114$, $d = -0.85$, Figure 4D) or on number of episodes in

general activity ($t = -1.60$, $p = 0.131$, $d = -0.80$) were found between CTR- and FLX-males.

However, just as FLX-females, FLX-males spent more time walking/running than CTR-males

($t = -3.05$, $p = .045$, $d = -1.52$, Figure 4E), but there was no difference in time spent on non-

social exploration ($t = 0.06$, $p = .953$, $d = 0.03$, Figure 4F). FLX-males did not differ from

CTR-males in the number of episodes walking/running ($t = -1.61$, $p = .130$, $d = -0.80$) or non-

social exploration ($t = -0.73$, $p = .786$, $d = -0.36$).

We then investigated whether there were differences between CTR- and FLX-rats in

how long it took them to visit all the 10 predefined locations (see Figure 2B) of the

seminatural environment. Rats that did not visit all locations within the observation time were

given a score of 3600 (total observation time in seconds). The results indicated that FLX-rats,

both males and females, did not need significantly more or less time to visit all locations than

CTR-rats (females: $t = 1.33 = p = .212$, $d = 0.42$; males: $t = -1.15$, $p = .271$, $d = -0.57$). We

thereafter investigated how many locations the rats visited as a function of time

(1/3/5/10/20/30/60 minutes), measured on cumulative data. FLX-females were not

significantly faster at visiting the different locations compared to CTR-females (Figure 4G),

but when the different time-points were analyzed separately, they seem to have visited

significantly more locations within the first 3 minutes ($t = -2.46$, $p = .027$, $d = -1.23$)

compared to CTR-females. No difference in the number of locations visited (treatment effect:

$F(1,14) = 3.43$, $p = .085$) or in the pattern (time x treatment: $F(2.64, 36.97) = 0.39$, $p = .735$)

over time were found between the CTR- and FLX-males (Figure 4H).

3.2.2 Other non-social behaviors

We also looked at other relevant non-social behaviors, including non-socially passive

behaviors. The analysis revealed that there was no significant difference between CTR- and

FLX-females in time spent on ($U = 28.00$, $z = -0.42$, $p = .721$, $r = -0.11$) or in the number of

non-socially passive behaviors ($t = -0.12$, $p = .903$, $d = -0.06$). Similarly, for the male groups,

no significant difference was found for time spent on ($t = 1.62$, $p = .127$, $d = 0.81$) or in the

number of non-socially passive behaviors ($t = 0.62$, $p = .546$, $d = 0.31$)

Next, we investigated whether CTR- and FLX-rats showed different level of

anxiety/stress-related behaviors. The results revealed no significant difference between CTR-

and FLX-rats for time spent on (females: $t = 1.67$, $p = .195$, $d = 0.84$; males: $U = 37.00$, $z =$

0.53, $p = .806$, $r = .13$) or in the number of episodes (females: $t = 0.58$, $p = .693$, $d = 0.29$;

males: $t = -0.60$, $p = .860$, $d = -0.30$) self-grooming. When investigating the total time in the

open area, no significant difference was found between CTR- and FLX-rats (females: $t = -$

1.39, $p = .186$, $d = -0.70$; males: $t = -0.98$, $p = .345$, $d = -0.49$). Similarly, the treatment groups

did not differ on the total time spent in the burrow area (females: $t = 1.57$, $p = .138$, $d = 0.79$;

males: $t = 1.02$, $p = .323$, $d = 0.51$).

Figure 4 Non-social behaviors in females and males

Figure 4. The data represent the time spent (s) on general activity (A, D), walking/running (B,

E), non-social exploration (C, F), and the total number of places in the SNE visited over time

(G, H). All graphs show comparisons between CTR-females (n = 8) and FLX-females (n = 8)

or between CTR-males (n = 8) and FLX-males (n = 8). Data are shown with individual data

points with bars representing the group means (A-F), or with squares and circles

representing respective group means (G-H). Error bars are representing SEM.

*p < 0.05

3.3 Fluoxetine exposure does not influence how the rats adapt to a novel environment

Finally, we were interested to see whether the treatment groups adapted differently to the novel physical and social environment, and thus, whether the differences in behavior between the groups were stable over time. We therefore divided the dataset into six 10-minutes time-bins and assessed the differences between CTR- and FLX-rats on social and non-social behaviors over the course of the observation period.

3.3.1 Social investigation behaviors

The repeated measure analysis revealed that FLX-females and FLX-males did not show a significantly different pattern of time spent on socially active behaviors, compared to CTR-females (time-bin x treatment: $F(5,70) = 0.26$, $p = .932$, $\eta_p^2 = .02$, Figure 5A) or CTR-males (time-bin x treatment: $F(5,70) = 0.51$, $p = .765$, $\eta_p^2 = .04$, Figure 5B) respectively. Similarly, when looking at the frequency of socially active behaviors, no interaction between time-bin and treatment was found for female ($F(5,70) = 0.63$, $p = .675$, $\eta_p^2 = .04$) or male rats ($F(5,70) = 0.99$, $p = .431$, $\eta_p^2 = .07$).

3.3.2 Non-social investigation behaviors

For time spent on walking/running, no differences as a function of time were found between the CTR- and FLX-rats for females ($F(5,70) = 0.63$, $p = .679$, $\eta_p^2 = .04$) or males ($F(2.64. 36.92) = 0.69$, $p = .634$, $\eta_p^2 = .05$), meaning that the increase in walking/running was present during the whole course of the hour and was most pronounced during the first 10- ($t = -2.77$, $p = .015$, $d = -1.38$) and 30-minutes ($U = 59.00$, $z = 2.84$, $p =.003$, $r = .71$) in FLX-females, and during the first 40- ($t = -3.58$, $p = .003$, $d = -1.79$) and 50-minutes ($t = -2.56$, $p = .023$, $d = -1.28$) in FLX-males, compared to CTR-animals. Similar results were found when

analyzing the frequency of walking/running (females: $F(5,70) = 0.88$, $p = .498$, $\eta_p^2 = .06$;

males: $F(2.85, 39.88) = 0.82$, $p = .483$, $\eta_p^2 = .06$). In term of non-social exploration, neither

FLX-females ($F(5,70) = 0.84$, $p = .529$, $\eta_p^2 = .06$) nor FLX-males ($F(2.87, 40.20) = 0.47$, $p =$

$.697$, $\eta_p^2 = .03$) showed a significant different pattern of time spent on exploration compared

to their control group. Thus, FLX-males did not differ from CTR-males throughout the

different time points during the observation period. FLX-females other the other hand, scored

lower than CTR-females during the whole hour, but most prominently in the first 10- ($t =$

3.03, $p = .009$, $d = 1.52$), 20- ($t = 4.38$, $p = < .001$, $d = 2.19$) and 30- minutes ($U = 12.00$, $z = -$

2.10, $p = .038$, $r = -.53$). Similar results were revealed for the frequency of non-social

exploration (females: $F(5,70) = 0.23$, $p = .948$, $\eta_p^2 = .02$; males: $F(5,70) = 1.76$, $p = .132$, η_p^2

$= .11$).

3.3.3 Other social and non-social behaviors

476   CTR- and FLX-rats showed no significant difference in pattern of time spent on

(females: time-bin x treatment: $F(1.21, 16.87) = 0.17$, $p = .729$, $\eta_p^2 = .01$; males: time-bin x

treatment: $F(1.89, 26.42) = 1.05$, $p = .361$, $\eta_p^2 = .07$) and number of (females: time-bin x

treatment: : $F(1.72, 24,12) = 0.41$, $p = .636$, $\eta_p^2 = .03$; males: time-bin x treatment: $F(2.04,$

$28.58) = 1.01$, $p = .378$, $\eta_p^2 = .07$) socially passive behaviors. Neither did they show any

significant difference in pattern of time spent on (females: time-bin x treatment: $F(5,70) =$

1.49, $p = .204$, $\eta_p^2 = .10$; males: time-bin x treatment: $F(1.99, 27.88) = 0.98$, $p = .388$, $\eta_p^2 =$

$.07$) or number of (females: time-bin x treatment: $F(5,70) = 0.63$, $p = .680$, $\eta_p^2 = .04$; males:

time-bin x treatment: $F(1.92, 26.84) = 0.77$, $p = .468$, $\eta_p^2 = .05$) conflict behaviors. However,

when analyzing the data non-linearly, a cubic interaction effect for the females group

indicated that when one group scored higher, the other scored lower ($F(1, 14) = 5.71$, $p =$

.031, $\eta_p^2 = .29$). For non-socially passive behaviors, no significant difference as a function of

time were found for females (duration: time-bin x treatment: $F(2.23, 31.14) = 0.26$, $p = .795$,

$\eta_p^2 = .02$; frequency: time-bin x treatment: $F(2.64, 36.99) = 0.18$, $p = .889$, $\eta_p^2 = .01$) and

males (duration: time-bin x treatment: $F(2.33, 32.62) = 0.58$, $p = .592$, $\eta_p^2 = .04$; frequency:

time-bin x treatment: $F(2.56, 35.90) = 0.51$, $p = .649$, $\eta_p^2 = .04$).

Figure 5 Social and non-social investigation behaviors measured as a function of time.

495
Figure 5. The data represent the time spent (s) on different behaviors as a function of time

measured every 10 minutes. The graphs show socially active behaviors (A, B),

walking/running (C, D), and non-social exploration (E, F). All graphs show comparisons

between CTR-females (n = 8) and FLX-females (n = 8) or between CTR-males (n = 8) and

FLX-males (n = 8). Squares and circles represent respective group means, error bars

*representing ± SEM.*p < 0.05, #p < 0.06*

4. Discussion

In our study, we investigated how perinatal fluoxetine exposure affects adult social

and non-social investigation behaviors in a novel seminatural environment with unfamiliar

conspecifics. Our findings show that perinatal fluoxetine exposure does not induce alterations

on social investigation behaviors and strategies when introduced to a novel seminatural

environment and unknown conspecifics. However, perinatal fluoxetine exposure was found to

affect non-social investigation behaviors. More specifically, perinatal fluoxetine exposed

female and male rats showed increased locomotor activity (in terms of walking/running),

while perinatal fluoxetine exposed females showed decreased non-social exploration.

Furthermore, it was demonstrated that the observed differences maintained throughout the

whole observation.

4.1 Social behaviors

The first question we investigated was whether social investigation behaviors,

operationalized as active social behaviors (sniffing and grooming other rats) and latency to

meet all other colony members, would be affected by perinatal SSRI exposure. The ability to

interact in line wth social norms is crucial in everyday life, and deviant social behavior in the

initial phase of contact can make it difficult to establish social relationships. The results in

this study revealed no differences between CTR- and FLX-rats on the total time spent on, or

the number of, active social behaviors. Previous findings from our research group showed that

FLX-females, but not FLX-males, showed a tendency toward decreased active social

behaviors (Houwing et al., 2019a), which was not present after naturally occurring aggressive

encounters. Nevertheless, in those studies, behaviors were observed after the rats had already

been housed together in the seminatural environment for several days, and thus were familiar

with each other. The effect of fluoxetine on social behaviors might have different outcomes

depending on whether the rats are interacting with familiar or unfamiliar partners (Gemmel et

al., 2019). In the present study, the rats were observed during the first hour after introduction

to the seminatural environment, allowing us to investigate how the rats encounter the first

social situations before knowing each other.

We also measured how long it took the rats to meet the other colony members after

being introduced to the novel environment. Such latency times could indicate whether the rats

have different interests in approaching other rats. Lack of social interest is a relevant trait to

examine since such symptoms commonly appear in various mental and neurodevelopmental

disorders (Barkus and Badcock, 2019). However, the results did not reveal any differences in

latencies to meet conspecifics between CTR- and FLX-rats. From our findings, we conclude

that perinatal SSRI exposure does not affect social investigation behavior and strategies

during the first hour after introduction to a novel environment with unfamiliar conspecifics.

Therefore, we suggest that the effect of perinatal SSRI exposure on social behaviors might be

dependent on the degree of familiarity between the rats. Although, the differences in results

could also have been caused by an altered interest in exploring the environment leading to

reduced social interactions.

We further investigated whether SSRI exposure leads to behavioral alterations in other

aspects of social behaviors, such as social passive behaviors and conflict behaviors. The

results revealed no difference in passive social behavior between FLX-rats and CTR-rats.

Furthermore, neither FLX-females nor FLX-males differed from CTR-rats in terms of conflict

behavior. However, conflict behavior was not frequently occurring in our experiment. The

Wistar strain is generally known to exhibit little aggressive behavior compared to other strains

(Koolhaas et al., 2013). In addition, the experiment was not designed to trigger aggressive

behavior as competition for food, water or mating partners were not necessary.

4.2 Non-social behaviors

Next, we investigated whether perinatal SSRI exposure would affect non-social

investigation behaviors, operationalized as locomotor activity (walking/running), non-social

exploration and latency to visit all locations of the environment. We found that both FLX-

females and FLX-males spent more time on locomotor activity compared to control rats.

FLX-females also had more episodes with locomotor activity compared to CTR-females. In

addition, FLX-females visited more locations of the seminatural environment within the first

3 minutes after entrance compared to CTR-females. Together, this could indicate that

perinatal SSRI exposure leads to an increased interest to investigate paths and locations.

Contrary to our findings, a recent meta-analysis found evidence for *reduced* activity in

developmentally SSRI exposed rats, as mostly measured by total distance moved (Ramsteijn

et al., 2020). Although we did not measure total distance per se, it is reasonable to assume that

total distance is related to total time spent walking/running in the seminatural environment.

Nevertheless, the meta-analysis is mainly based on studies measuring activity in simplified

open field boxes. Such set-ups allow the rats to perceive the whole environment without

necessarily having to move their bodies. We could therefore assume that an increased interest

to investigate locations and paths would only be observable in situations where

walking/running (movement) is needed to investigate the environment. In addition, in the

current environment, more rats were present leading to the assumption that the odors and

sounds from others may also elicit extra movement, making our set-up more reliable to study

the effects of perinatal fluoxetine exposure on a measure as locomotor activity reflecting

alterations in interest to investigate novel paths and locations. With this in mind, we would

expect the differences between FLX- and CTR-rats to disappear (or diminish) when the

animals get familiar with their surroundings and conspecifics. Interestingly, previous studies

from our research group did indeed find no differences on locomotor activity between FLX-

and CTR- rats after the rats were already familiarized with the environment (Heinla et al.,

2020; Houwing et al., 2019a). This suggests that the current findings of increased locomotor

activity in FLX-rats are related to the introduction to a novel environment, and not the

complexity of the environment on itself.

We also found that FLX-females, but not FLX-males, spent less time on non-social

exploration than control rats, meaning they were sniffing less on objects (e.g. shelters,

wooden sticks) and specific elements in the environment (e.g. walls, the ground). This is in

line with previous findings from day 4 and day 7 in the same experiment (Houwing et al.,

2019a), where reduced non-social exploration was found in FLX-females, but not in FLX-

males. Other studies have also reported reduced non-social explorative behaviors in SSRI

exposed rats (Ansorge et al., 2004; Karpova et al., 2009; Rebello et al., 2014; Sarkar et al.,

2014; Simpson et al., 2011; Zohar et al., 2016). Although we have shown that FLX-females

seem to have increased interest to explore paths and locations, shown by increased locomotor

activity, our findings also indicate that perinatal SSRI exposure in females leads to reduced

interest to investigate objects and other specific elements in the environment. Although the

findings might seem contradictive at first sight, locomotor activity and non-social exploration

could possibly serve different purposes. As locomotor activity could measure the interest to

get quickly familiar with the whole environment as a kind of screening behavior, non-social

exploration reflects a more detailed and accurate investigation of the environment. Therefore,

we suggest that perinatal SSRI exposure alters the strategy the animals use to investigate a

novel environment leading to a quicker, but less detailed investigation of novel environments.

598   We further investigated other non-social behaviors such as anxiety/stress-related

behaviors. We did not find any difference between CTR- and FLX-rats on anxiety/stress-

related behaviors. A previous study found that white-noise exposure induced increased self-

grooming in FLX-males (Houwing et al., 2019a), which was explained as an altered stress-

coping behavior. As introduction to a new environment can be considered a stressful

situation, we expected to observe a similar increase in self-grooming behavior in FLX-males

in the present study. However, no differences were found between CTR- and FLX- rats on

self-grooming behavior. Moreover, no differences were found on the amount of time spent in

the open area, as measure for changes in anxiety-related behavior. Altogether, this makes us

to conclude that perinatal SSRI exposure does not affect anxiety/stress-related behavior

during the first hour of exposure to a novel environment with unfamiliar conspecifics.

4.3 Behavioral adaption over time

611   The last question we investigated was whether perinatal SSRI exposed rats adapt

differently to unfamiliarity (both environmental and socially) than their non-exposed

conspecifics. Therefore, we split the observational data into six 10-minute time-bins in order

to look at behavioral changes over time. As part of the familiarization process to a new

environment, we generally expected to see adjustments in behavior during the first hour, such

as decrease in general activity (Wilkinson et al., 2006). However, our main subject of interest

was whether perinatal SSRI exposed rats adjusted their behavior in a different manner than

controls.

Our results revealed that SSRI exposed animals adapted similarly to the novel

environment as control animals. As discussed, FLX-females spent less time exploring objects

and the physical environment, whereas both FLX-males and FLX-females spent more time on

locomotor activity compared to CTR-rats. Those differences remained relatively stable

throughout the first hour, meaning that FLX- and CTR- rats behaved differently, but adapted

similarly to the novel environment over time (the increased locomotor activity remained

higher during the full course of the observed hour). Interestingly, the differences in locomotor

activity and non-social exploration in FLX-females was mostly pronounced during the first 30

minutes, while the FLX-males had more increased locomotor activity during the last 30

minutes. The reason for this remains unclear, but since our experiment employed a reversed

dark/light cycle in which the light conditions gradually decreased from daylight to moonlight

between 10.30 and 11.00 every morning, and the rats thus experienced the shift from light to

darkness during the first hour, this could have resulted in sex-specific alterations on

behavioral adjustment to light between male and female rats. However, future experiments are

needed before further conclusions can be drawn.

We conclude that perinatal fluoxetine exposed rats do not adapt their behaviors

differently than controls during the first hour after introduction to the novel environment,

instead the changes in non-social investigation behavior remain stable over time.

5. Conclusion

In summary, our data show that perinatal SSRI exposure alters aspects of non-social

investigation behaviors when introduced to a novel environment with unfamiliar conspecifics,

but did not alter social investigation behaviors. Both FLX-males and FLX-females showed a

higher amount of locomotor activity, while FLX-females visited more locations within the

first three minutes, and spent less time exploring objects and specific elements in the physical

environment. Perinatal fluoxetine exposure did not affect social behavior or how the animals

adapted to the unfamiliar seminatural environment over time. Altogether, we conclude that

perinatal SSRI exposure alters non-social investigation, to a quicker and less detailed strategy,

when exposed to a novel environment, and that the alteration is most pronounced in females.

Acknowledgements

Financial support was received from Helse Nord #PFP1295-16, Norway. We also

would like to thank Ragnhild Osnes, Carina Sørensen, Nina Løvhaug, Katrine Harjo, and

Remi Osnes for their excellent care of the animals.

References

Andrade, S.E., Raebel, M.A., Brown, J., Lane, K., Livingston, J., Boudreau, D., Rolnick, S.J., Roblin, D., Smith, D.H., Willy, M.E., Staffa, J.A., Platt, R., 2008. Use of antidepressant medications during pregnancy: A multisite study. American journal of obstetrics and gynecology 198.

Ansorge, M.S., Zhou, M., Lira, A., Hen, R., Gingrich, J.A., 2004. Early-life blockade of the 5-ht transporter alters emotional behavior in adult mice. Science 306, 879-881.

Azmitia, E.C., 2001. Modern views on an ancient chemical: Serotonin effects on cell proliferation, maturation, and apoptosis. Brain Res Bull 56, 413-424.

Barkus, E., Badcock, J.C., 2019. A transdiagnostic perspective on social anhedonia. Front Psychiatry 10, 216.

Bennett, H.A., Einarson, A., Taddio, A., Koren, G., Einarson, T.R., 2004. Prevalence of depression during pregnancy: Systematic review. Obstet Gynecol 103, 698-709.

Brown, A.S., Gyllenberg, D., Malm, H., McKeague, I.W., Hinkka-Yli-Salomaki, S., Artama, M., Gissler, M., Cheslack-Postava, K., Weissman, M.M., Gingrich, J.A., Sourander, A., 2016. Association of selective serotonin reuptake inhibitor exposure during pregnancy with speech, scholastic, and motor disorders in offspring. JAMA Psychiatry 73, 1163-1170.

Charlton, R.A., Jordan, S., Pierini, A., Garne, E., Neville, A.J., Hansen, A.V., Gini, R., Thayer, D., Tingay, K., Puccini, A., Bos, H.J., Nybo Andersen, A.M., Sinclair, M., Dolk, H., de Jong-van den Berg, L.T., 2015. Selective serotonin reuptake inhibitor prescribing before, during and after pregnancy: A population-based study in six european regions. BJOG 122, 1010-1020.

Chu, X., Agmo, A., 2014. Sociosexual behaviours in cycling, intact female rats (rattus norvegicus) housed in a seminatural environment. Behaviour 151, 1143-1184.

Dunkel Schetter, C., 2011. Psychological science on pregnancy: Stress processes, biopsychosocial models, and emerging research issues. Annu Rev Psychol 62, 531-558.

El Marroun, H., White, T., Verhulst, F.C., Tiemeier, H., 2014. Maternal use of antidepressant or anxiolytic medication during pregnancy and childhood neurodevelopmental outcomes: A systematic review. Eur Child Adolesc Psychiatry 23, 973-992.

Gaspar, P., Cases, O., Maroteaux, L., 2003. The developmental role of serotonin: News from mouse molecular genetics. Nat Rev Neurosci 4, 1002-1012.

Gemmel, M., De Lacalle, S., Mort, S.C., Hill, L.A., Charlier, T.D., Pawluski, J.L., 2019. Perinatal fluoxetine has enduring sexually differentiated effects on neurobehavioral outcomes related to social behaviors. Neuropharmacology 144, 70-81.

Gemmel, M., Hazlett, M., Bogi, E., De Lacalle, S., Hill, L.A., Kokras, N., Hammond, G.L., Dalla, C., Charlier, T.D., Pawluski, J.L., 2017. Perinatal fluoxetine effects on social play, the hpa system, and hippocampal plasticity in pre-adolescent male and female rats: Interactions with pre-gestational maternal stress. Psychoneuroendocrinology 84, 159-171.

Glover, M.E., Clinton, S.M., 2016. Of rodents and humans: A comparative review of the neurobehavioral effects of early life ssri exposure in preclinical and clinical research. Int J Dev Neurosci 51, 50-72.

Goodman, S.H., 2007. Depression in mothers. Annu Rev Clin Psychol 3, 107-135.

Hegstad, J., Huijgens, P.T., Houwing, D.J., Olivier, J.D.A., Heijkoop, R., Snoeren, E.M.S., 2020. Female rat sexual behavior is unaffected by perinatal fluoxetine exposure. Psychoneuroendocrinology 120, 104796.

Heinla, I., Heijkoop, R., Houwing, D.J., Olivier, J.D.A., Snoeren, E.M.S., 2020. Third-party prosocial behavior in adult female rats is impaired after perinatal fluoxetine exposure. Physiology & behavior 222, 112899.

Hermansen, T.K., Roysamb, E., Augusti, E.M., Melinder, A., 2016. Behavior and inhibitory control in children with prenatal exposure to antidepressants and medically untreated depression. Psychopharmacology 233, 1523-1535.

Houwing, D.J., Esquivel-Franco, D.C., Ramsteijn, A.S., Schuttel, K., Struik, E.L., Arling, C., de Boer, S.F., Olivier, J.D.A., 2020. Perinatal fluoxetine treatment and dams' early life stress history have opposite effects on aggressive behavior while having little impact on sexual behavior of male rat offspring. Psychopharmacology 237, 2589-2600.

Houwing, D.J., Heijkoop, R., Olivier, J.D.A., Snoeren, E.M.S., 2019a. Perinatal fluoxetine exposure changes social and stress-coping behavior in adult rats housed in a seminatural environment. Neuropharmacology 151, 84-97.

Houwing, D.J., Staal, L., Swart, J.M., Ramsteijn, A.S., Wohr, M., de Boer, S.F., Olivier, J.D.A., 2019b. Subjecting dams to early life stress and perinatal fluoxetine treatment differentially alters social behavior in young and adult rat offspring. Front Neurosci 13, 229.

Karpova, N.N., Lindholm, J., Pruunsild, P., Timmusk, T., Castren, E., 2009. Long-lasting behavioural and molecular alterations induced by early postnatal fluoxetine exposure are

restored by chronic fluoxetine treatment in adult mice. Eur Neuropsychopharmacol 19, 97-108.

Khatri, N., Simpson, K.L., Lin, R.C., Paul, I.A., 2014. Lasting neurobehavioral abnormalities in rats after neonatal activation of serotonin 1a and 1b receptors: Possible mechanisms for serotonin dysfunction in autistic spectrum disorders. Psychopharmacology 231, 1191-1200.

Kim, J., Riggs, K.W., Misri, S., Kent, N., Oberlander, T.F., Grunau, R.E., Fitzgerald, C., Rurak, D.W., 2006. Stereoselective disposition of fluoxetine and norfluoxetine during pregnancy and breast-feeding. Br J Clin Pharmacol 61, 155-163.

Kiryanova, V., Dyck, R.H., 2014. Increased aggression, improved spatial memory, and reduced anxiety-like behaviour in adult male mice exposed to fluoxetine early in life. Dev Neurosci 36, 396-408.

Klinger, G., Frankenthal, D., Merlob, P., Diamond, G., Sirota, L., Levinson-Castiel, R., Linder, N., Stahl, B., Inbar, D., 2011. Long-term outcome following selective serotonin reuptake inhibitor induced neonatal abstinence syndrome. J Perinatol 31, 615-620.

Koolhaas, J.M., Coppens, C.M., de Boer, S.F., Buwalda, B., Meerlo, P., Timmermans, P.J., 2013. The resident-intruder paradigm: A standardized test for aggression, violence and social stress. J Vis Exp, e4367.

Kristensen, J.H., Ilett, K.F., Hackett, L.P., Yapp, P., Paech, M., Begg, E.J., 1999. Distribution and excretion of fluoxetine and norfluoxetine in human milk. Br J Clin Pharmacol 48, 521-527.

Lundmark, J., Reis, M., Bengtsson, F., 2001. Serum concentrations of fluoxetine in the clinical treatment setting. Ther Drug Monit 23, 139-147.

Lupattelli, A., Wood, M., Ystrom, E., Skurtveit, S., Handal, M., Nordeng, H., 2018. Effect of time-dependent selective serotonin reuptake inhibitor antidepressants during pregnancy on behavioral, emotional, and social development in preschool-aged children. J Am Acad Child Adolesc Psychiatry 57, 200-208.

Malm, H., Brown, A.S., Gissler, M., Gyllenberg, D., Hinkka-Yli-Salomaki, S., McKeague, I.W., Weissman, M., Wickramaratne, P., Artama, M., Gingrich, J.A., Sourander, A., 2016. Gestational exposure to selective serotonin reuptake inhibitors and offspring psychiatric disorders: A national register-based study. J Am Acad Child Adolesc Psychiatry 55, 359-366.

Mitchell, A.A., Gilboa, S.M., Werler, M.M., Kelley, K.E., Louik, C., Hernandez-Diaz, S., Stud, N.B.D.P., 2011. Medication use during pregnancy, with particular focus on prescription drugs: 1976-2008. American journal of obstetrics and gynecology 205.

Molenaar, N.M., Bais, B., Lambregtse-van den Berg, M.P., Mulder, C.L., Howell, E.A., Fox, N.S., Rommel, A.S., Bergink, V., Kamperman, A.M., 2020. The international prevalence of antidepressant use before, during, and after pregnancy: A systematic review and meta-analysis of timing, type of prescriptions and geographical variability. Journal of Affective Disorders 264, 82-89.

Muller, C.L., Anacker, A.M.J., Veenstra-VanderWeele, J., 2016. The serotonin system in autism spectrum disorder: From biomarker to animal models. Neuroscience 321, 24-41.

Noorlander, C.W., Ververs, F.F., Nikkels, P.G., van Echteld, C.J., Visser, G.H., Smidt, M.P., 2008. Modulation of serotonin transporter function during fetal development causes dilated heart cardiomyopathy and lifelong behavioral abnormalities. PLoS One 3, e2782.

Olivier, J.D., Blom, T., Arentsen, T., Homberg, J.R., 2011a. The age-dependent effects of selective serotonin reuptake inhibitors in humans and rodents: A review. Prog Neuropsychopharmacol Biol Psychiatry 35, 1400-1408.

Olivier, J.D., Valles, A., van Heesch, F., Afrasiab-Middelman, A., Roelofs, J.J., Jonkers, M., Peeters, E.J., Korte-Bouws, G.A., Dederen, J.P., Kiliaan, A.J., Martens, G.J., Schubert, D., Homberg, J.R., 2011b. Fluoxetine administration to pregnant rats increases anxiety-related behavior in the offspring. Psychopharmacology 217, 419-432.

Rampono, J., Proud, S., Hackett, L.P., Kristensen, J.H., Ilett, K.F., 2004. A pilot study of newer antidepressant concentrations in cord and maternal serum and possible effects in the neonate. Int J Neuropsychopharmacol 7, 329-334.

Ramsteijn, A.S., Van de Wijer, L., Rando, J., van Luijk, J., Homberg, J.R., Olivier, J.D.A., 2020. Perinatal selective serotonin reuptake inhibitor exposure and behavioral outcomes: A systematic review and meta-analyses of animal studies. Neurosci Biobehav Rev 114, 53-69.

Rebello, T.J., Yu, Q., Goodfellow, N.M., Caffrey Cagliostro, M.K., Teissier, A., Morelli, E., Demireva, E.Y., Chemiakine, A., Rosoklija, G.B., Dwork, A.J., Lambe, E.K., Gingrich, J.A., Ansorge, M.S., 2014. Postnatal day 2 to 11 constitutes a 5-ht-sensitive period impacting adult mpfc function. J Neurosci 34, 12379-12393.

Rodriguez-Porcel, F., Green, D., Khatri, N., Harris, S.S., May, W.L., Lin, R.C., Paul, I.A., 2011. Neonatal exposure of rats to antidepressants affects behavioral reactions to novelty and

social interactions in a manner analogous to autistic spectrum disorders. Anat Rec (Hoboken) 294, 1726-1735.

Sarkar, A., Chachra, P., Vaidya, V.A., 2014. Postnatal fluoxetine-evoked anxiety is prevented by concomitant 5-ht2a/c receptor blockade and mimicked by postnatal 5-ht2a/c receptor stimulation. Biol Psychiatry 76, 858-868.

Simpson, K.L., Weaver, K.J., de Villers-Sidani, E., Lu, J.Y., Cai, Z., Pang, Y., Rodriguez-Porcel, F., Paul, I.A., Merzenich, M., Lin, R.C., 2011. Perinatal antidepressant exposure alters cortical network function in rodents. Proceedings of the National Academy of Sciences of the United States of America 108, 18465-18470.

Snoeren, E.M., Antonio-Cabrera, E., Spiteri, T., Musatov, S., Ogawa, S., Pfaff, D.W., Agmo, A., 2015. Role of oestrogen alpha receptors in sociosexual behaviour in female rats housed in a seminatural environment. J Neuroendocrinol 27, 803-818.

Svirsky, N., Levy, S., Avitsur, R., 2016. Prenatal exposure to selective serotonin reuptake inhibitors (ssri) increases aggression and modulates maternal behavior in offspring mice. Dev Psychobiol 58, 71-82.

Wilkinson, J.L., Herrman, L., Palmatier, M.I., Bevins, R.A., 2006. Rats' novel object interaction as a measure of environmental familiarity. Learning and Motivation 37, 131-148.

Woody, C.A., Ferrari, A.J., Siskind, D.J., Whiteford, H.A., Harris, M.G., 2017. A systematic review and meta-regression of the prevalence and incidence of perinatal depression. J Affect Disord 219, 86-92.

Zimmerberg, B., Germeyan, S.C., 2015. Effects of neonatal fluoxetine exposure on behavior across development in rats selectively bred for an infantile affective trait. Dev Psychobiol 57, 141-152.

Zohar, I., Shoham, S., Weinstock, M., 2016. Perinatal citalopram does not prevent the effect of prenatal stress on anxiety, depressive-like behaviour and serotonergic transmission in adult rat offspring. The European journal of neuroscience 43, 590-600.