Desert lizard diversity worldwide: effects of environment, time, and evolutionary rate

Running title: Desert lizard diversity

ABSTRACT

Aim: Biodiversity is not uniformly distributed across the Earth's surface, even among

physiographically comparable biomes in different biogeographic regions. For lizards, the

world's large desert regions are characterized by extreme heterogeneity in species richness,

spanning some of the most species-rich (arid Australia) and species-poor (central Asia)

biomes overall. Regional differences in species diversity may arise as a consequence of the

interplay of several factors (e.g., evolutionary time, diversification rate, environment), but

their relative importance for biogeographic patterns remains poorly known. Here we use

distributional and phylogenetic data to assess the evolutionary and ecological drivers of large-

scale variation in desert lizard diversity.

Location: Deserts worldwide.

Major taxa studied: Lizards (non-snake squamates).

Methods: We specifically test whether diversity patterns are best explained by differences in

the ages of arid-adapted lineages (evolutionary time hypothesis), by regional variation in

speciation rate, by geographic area of the arid systems, and by spatial variation related to

environment (climate, topography, and productivity).

Results: We found no effect of recent speciation rate and geographic area on differences in

desert lizard diversity. We demonstrate that the extreme species richness of the Australian

deserts cannot be explained by greater evolutionary time, because species began

accumulating more recently there than in more species-poor arid regions. We found limited

support for relationships between regional lizard richness and environmental variables, but

these effects were inconsistent across deserts, showing a differential role of the environment

in shaping the lizard diversity in different arid regions.

Main conclusions: Our results provide evidence against several classic hypotheses for

interregional variation in species richness, but also highlight the complexity of processes

underlying vertebrate community richness in the world's great arid systems.

Keywords: Arid regions; regional diversity; species richness; diversity anomalies;

evolutionary time; speciation rate.

INTRODUCTION

What causes the dramatic differences in species diversity that we observe among different

regions of the world? The latitudinal diversity gradient (LDG) is perhaps the most famous

spatial diversity pattern, but species richness can vary greatly even among physiographically

similar but geographically distinct biomes (Couvreur, 2015; Kelt et al., 1996; Pianka, 1986;

Ricklefs & Schluter, 1993; Valente & Vargas, 2013). There is overall agreement that multiple

ecological and evolutionary factors influence the assembly of regional species pools through

their effects on dispersal, speciation, and extinction (Cracraft, 1992; Rabosky, 2009; Ricklefs,

1987; Weber, Wagner, Best, Harmon, & Matthews, 2017; Wiens, 2011), but we still lack

general consensus on the relative importance of processes that shape regional differences in

species richness (Belmaker & Jetz, 2015; Mittelbach et al., 2007).

A key challenge with large-scale assessments of species richness is that climatic, historical, and geographic hypotheses for species diversity may covary, such that disentangling their relative contributions to species diversity may be difficult or impossible (Etienne et al., 2019; Rabosky & Hurlbert, 2015). For example, in the case of the LDG, the tropics are generally older, larger, warmer, and more productive than extratropical regions. These factors are generally predicted to affect species richness in the same direction, and it is therefore difficult to isolate the effects of any single process on the overall LDG. In contrast, regional differences in species richness among climatically-matched regions can facilitate tests of the role of historical and geographic factors on diversity, by controlling – at least in part – for broad-scale differences in climate between regions. Many previous studies have addressed the role of evolutionary processes in generating these "diversity anomalies" (Ricklefs & Latham, 1993) through intercontinental comparisons of tropical rainforests (Couvreur, 2015), temperate floras (Latham & Ricklefs, 1993; Qian & Ricklefs, 2000), Mediterranean floras (Valente & Vargas, 2013), mangrove ecosystems (Ricklefs & Latham, 1993), and others.

One of the most striking regional diversity anomalies is the exceptional species richness of Australian desert lizards relative to lizards from other deserts (Pianka, 1969, 1971, 1972). Arid Australia has long been known to have outstanding lizard diversity at the biogeographic scale (i.e., the total number of species in the arid zone) compared to other arid systems of the world (Pianka, 1985, 1986). In a recent synthesis of global reptile distributions, (Roll et al., 2017) found strong support for exceptional regional diversity (richness in one-degree grid cells) of lizards in arid Australia. This diversity holds not only in comparison to other arid systems, but even in comparison to lowland tropical rainforests (Roll et al., 2017). At the local scale, more than 70 species of lizards (non-snakes) can occur in regional sympatry in arid Australia, a number that far exceeds those for the world's second-

most species-rich lizard communities (e.g., Amazon rainforest, with 25-35 species;

Duellman, 2005; Rabosky et al., 2019). Numerous hypotheses have been proposed to explain

these dramatic differences in the number of species, including vegetative structure, habitat

specificity and heterogeneity, termite abundance, and other ecological, biogeographic, and

historical factors (Morton & James, 1988; Pianka, 1972, 1989).

The outstanding richness of Australian lizards at local, regional, and biogeographic

scales (Pianka, 1969, 1986), suggests the possibility that ecological and evolutionary factors

cascade upwards and/or downwards to influence species richness at larger (or smaller) spatial

scales. For instance, analysing range size and overlap for the diverse (~100 species)

Australian lizard genus *Ctenotus*, James & Shine (2000) concluded that the high local species

richness might be related to the large area of the arid zone, and the high regional diversity can

be explained by the wide distribution ranges favoured by climate homogeneity. Thus, local

richness may, at least in part, reflect processes that contribute to overall high species diversity

within the biogeographic and regional species pools (Ricklefs, 1987, 2004), such that

evolutionary and historical factors operating at regional-biogeographic scales also shape

species richness at local scales.

High levels of species diversity can be the outcome of different, but not mutually

exclusive, factors (Jablonski, Huang, Roy, & Valentine, 2017). For example, high diversity

might be simply due to the relatively ancient and undisturbed history of one region and the

lineages within (evolutionary time hypothesis; Fine & Ree, 2006; Fischer, 1960; Mittelbach

et al., 2007; Wallace, 1878). Net diversification rates, either through faster speciation or

lower extinction, might also contribute to elevated species richness (Schluter & Pennell,

2017). Other explanations include physical characteristics of the regions, which in turn might

affect species richness through their impact on evolutionary rates or through their impact on

the level at which diversity is regulated by diversity-dependent factors. Large geographic

areas may offer more opportunities for speciation than smaller regions (Rosenzweig &

Sandlin, 1997), and climate and topography, as well as environmental heterogeneity, can

further influence species diversity through impacts on primary productivity, habitat and

resource availability, and diversification rate (Antão et al., 2020; Badgley et al., 2017;

Cantalapiedra, Domingo, & Domingo, 2018; Garcia-Porta et al., 2019; Menéndez et al.,

2021; Stein et al., 2015; Stein, Gerstner, & Kreft, 2014; Velasco et al., 2018).

Here, we address the causes of the dramatic differences in lizard species richness

across desert regions, with an emphasis on the exceptional diversity in arid Australia,

focusing in particular on large (regional to biogeographic) spatial scales. We deliberately

omit snakes from our assessment, because geographic patterns of snake diversity are

markedly divergent from those for other (non-snake) squamate reptiles (e.g., "lizards"); snake

diversity is generally highest in humid tropical lowlands and is broadly congruent with

patterns in vascular plants, frogs, birds, and many other taxa (Roll et al., 2017). Through

targeted comparisons between the largest arid systems worldwide, we test a series of

hypotheses that could explain pronounced differences in lizard species richness at the

biogeographic and regional scales across these (broadly-comparable physiographic) regions.

Specifically, we test whether species diversity in these systems is related to: i) the age of the

arid-adapted lineages in each system (evolutionary time hypothesis), ii) within-region

speciation rates, iii) geographic area of the system, iv) environmental variables that could

also affect the carrying capacity of the system, as well as a measure of spatial heterogeneity

for each of them: topography, mean annual temperature, mean diurnal range, annual

precipitation, precipitation of the driest month, aridity, potential evapotranspiration, and

productivity.

After decades of interest in these questions, the increasing availability of data and the

development of new comparative phylogenetic and spatial methods allow us to finally make

broad comparisons between these regions from an evolutionary perspective (Schemske &

Mittelbach, 2017). Although our analyses cannot conclusively identify the factors that cause

geographic differences in species richness, they can reject potential explanations (Etienne et

al., 2019; Rabosky & Hurlbert, 2015).

MATERIAL AND METHODS

Delimitation of arid systems and species categorization

We analysed lizard faunas in the deserts of the world, understanding lizards to represent all

non-snake squamate reptiles. We first identified Earth's largest arid systems, focusing on

those delineated by Olson et al. (2001) in the Biome "Deserts and Xeric Shrublands" (Biome

13). Within this biome, we considered the major regions to be those with a substantial

component of arid and hyper-arid conditions, that is, with an Aridity Index (AI) < 0.2

(Parsons & Abrahams, 1994; Rocha, Godinho, Brito, & Nielsen, 2021). This delimitation set

includes 8 main arid systems (Figure 1): Atacama, Australian desert, Central Asian desert,

Gobi, Kalahari, North American desert, Persian desert, and Saharo-Arabian desert. These

'main arid systems' result from the aggregation of contiguous desert regions into single large

arid zones (e.g., the North American arid system includes the Sonoran, Mojave, Chihuahuan

and Great Basin arid subregions; the Saharo-Arabian system includes the Sahara desert and

several arid regions through the Arabian Peninsula and the Horn of Africa; etc.). Our

delimitation of contiguous arid systems (e.g., we merged the Sahara and the Arabian deserts,

and kept the Central Asian and the Persian deserts as separate entities) was based on a

regionalisation analysis through the interactive online tool for delimiting biogeographical

regions from species distribution Infomap Bioregions (Edler, Guedes, Zizka, Rosvall, &

Antonelli, 2017), which uses previously described clustering methods (Rosvall & Bergstrom,

2008; Vilhena & Antonelli, 2015). For this analysis, we used global reptile distribution data

(Roll et al., 2017; see below), setting the cluster cost to 1.0 and implementing 10 trials (Edler

et al., 2017).

We used lizard distribution data from a recently published global reptile dataset (Roll

et al., 2017) to categorize each lizard species into arid systems, following twofold criteria: a

species was considered as inhabiting a desert when at least 15% of its geographic range falls

within the desert, or when at least 15% of the total desert's geographic area is occupied by the

species. We tested the sensitivity of the categorization percentage across a range of

thresholds, and we found that the assigned species (and total diversities) for each desert were

similar across a broad range of numerical thresholds (Figure S1).

Evolutionary time hypothesis

The simplest hypothesis to explain geographic differences in species diversity relates to the

total evolutionary time over which species have accumulated in different regions. This idea

dates back to Wallace (1878) who proposed that the tropical diversity represents a 'more

ancient world' than the temperate zones to explain the latitudinal diversity gradient. In order

to test this hypothesis, which we refer to as the 'evolutionary time hypothesis', we estimated

the times when lineages started diversifying within the different arid systems of the world. To

do that, we reconstructed the ancestral presence of squamates in the deserts after assigning

extant species to arid systems. We used the consensus tree of squamates from Tonini et al.

(2016), excluding all the species without genetic or distribution data. The inclusion of species

whose phylogenetic position is imputed from taxonomic data alone leads to predictable biases

in ancestral state and evolutionary rate estimation and should be avoided (Rabosky, 2015).

The resulting tree contained 5,355 (out of a total of 11,183; Uetz, Freed, & Hošek, 2021)

squamate species (3,828 lizards and 1,526 snakes, plus the outgroup of all squamates, the

tuatara; Table S1 shows the percentage of squamates, lizards, and snakes of this pruned tree

in each desert relative to the phylogeny of all squamates). We assigned each one to the arid

systems or categorised as "non-arid". This classification resulted in 38 species that were

assigned to more than one desert, distributed in two or more contiguous systems: Gobi,

Central Asian, Persian, and Saharo-Arabian deserts (Table S2). We applied a model of

character evolution with each species coded as one out of 14 possible states: "non-arid",

"North American", "Central Asian", "Central Asian + Gobi", "Central Asian + Gobi +

Persian", "Central Asian + Persian", "Central Asian + Persian + Saharo-Arabian",

"Atacama", "Australian", "Gobi", "Kalahari", "Persian", "Persian + Saharo-Arabian",

"Saharo-Arabian". For the ancestral state estimation we used the function 'make.simmap'

from the R package 'phytools' (Revell, 2012), with an all-rates-different (ARD) model and

100 simulations. For phylogenetic data manipulation we used the R packages 'geiger'

(Harmon, Weir, Brock, Glor, & Challenger, 2008) and 'treeio' (Wang et al., 2020). This

approach allowed us to estimate the age at which lizard lineages started accumulating in each

arid system. Critically, we do not interpret character state changes to imply biogeographic

movements out of or into a particular desert. For example, an apparent transition from

"Australian" to "non-arid" could simply reflect an evolved shift in biome for a lineage. Such

a shift could occur even if species ranges are static through time: for example, a biome might

expand or contract through time, and species might simply adapt to the new climate regime

with little or no geographic movement (Donoghue & Edwards, 2014). This is particularly

relevant for our analysis of 'Desert colonisation dynamics' (see below).

We estimated the general trajectory of lineage accumulation for each arid zone by

summing the cumulative ancestral state probabilities for a given region through time (Mahler,

Revell, Glor, & Losos, 2010). This metric is clearly biased by the fact that only lineages with

extant descendants contribute to the lineage totals at each point in time, and thus cannot

detect earlier radiations within arid regions that failed to leave extant descendants. However,

in our case, we are testing hypotheses to explain the observed phylogenetic structure of arid-

zone lizard diversity, and thus it is the lineage accumulation curve for extant species that is

most relevant. If evolutionary time is the dominant factor influencing species richness, we

expect the lineage accumulation curve for the most species-rich deserts (e.g., Australia) to

predate the curves for other major arid regions. We performed correlation tests between the

total number of species in each desert and the estimated time by which each desert had

accumulated 10% or 30% of their total lineage diversity in the ancestral reconstruction, as a

way of testing if faunal build-up started earlier in high-diversity deserts. We note that a

positive correlation between age and richness is not diagnostic for identifying a contribution

of clade- or regional age on species diversity. Even if diversity in each region is strictly

equilibrial (and thus, not controlled by age), more diverse regions can appear "older", simply

because time to common ancestry for a regional community in equilibrium models should be

positively correlated with the equilibrium number of species (Rabosky & Hurlbert, 2015).

However, a negative correlation does reject the effect of age on richness. If a species-rich

biota is found to be "young", it is certainly possible that an unobserved history of

diversification predates the observed lineage accumulation curve (Louca & Pennell, 2020;

Quental & Marshall, 2010; Rabosky, 2016). However, it cannot then be the case that age

itself is the control on richness, because the observed "young" age would have still been fully

sufficient to account for all extant species richness in the species-rich region.

Desert colonisation dynamics

Colonisation (immigration) is a key process that affects regional biodiversity patterns, along

with evolutionary processes of speciation and extinction (Crouch, Capurucho, Hackett, &

Bates, 2019; Hua & Bromham, 2020; MacArthur & Wilson, 1967; Simberloff, 1974). We

explored the dynamics of colonisation of each desert, as well as the lizard interchange

between contiguous desert regions. With such purpose, we extracted the mean number of

biome shifts from external (non-arid) biomes to each arid system, as well as between arid

systems. This information resulted from the 100 simulations of the ancestral reconstruction of

desert occupancy that we implemented to test the evolutionary time hypothesis (see above).

Species richness, speciation rates and geographic area of the deserts

Another evolutionary mechanism that might explain richness patterns is variation in

diversification rates, e.g., areas and clades with higher diversification rates will presumably

have higher species diversity. In light of difficulties in estimating extinction rates from

molecular phylogenies (Rabosky, 2016), we tested this hypothesis using recent speciation

rates only (tip rates), which is likely the parameter that can be most reliably inferred from a

time-calibrated phylogeny (Louca & Pennell, 2020). All else being equal, elevated speciation

for a given biome should translate into elevated net diversification, unless extinction rates are

also sufficiently increased as to offset the effects of increased speciation rates. We estimated

recent speciation rates using the DR metric (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012).

Although sometimes presented as an estimate of net diversification rate, DR is a much better

estimate of speciation rate (Title & Rabosky, 2019). Using the Tonini et al. (2016)

pseudoposterior distribution of fully-sampled squamate phylogenies, we calculated the mean

DR metric for each of the 9,755 species of squamates (lizards and snakes) across 100 trees

from the posterior. The Tonini et al. (2016) phylogenies are complete at the species level but

include approximately 4,000 species whose position was imputed from taxonomic

information using a stochastic birth-death polytomy resolver (Thomas et al., 2013). DR

values computed from such trees are relatively robust to biases caused by incomplete taxon

sampling and afford greater power to infer variation in evolutionary rates relative to estimates

that use formal sampling fractions (Chang, Rabosky, & Alfaro, 2020). We then conducted

nonparametric Kruskal-Wallis and Wilcoxon post-hoc analyses to compare lizard speciation

rates among deserts to address how this metric relates with biogeographic species richness

(the total number of species within each desert). We performed a correlation test between the

log-transformed values of total number of species and mean DR values for each desert.

We also explored the relationship between species richness, speciation rates, and the

geographic area of the arid systems, since the size of an area may affect both the species

diversity and the speciation rates (Connor & McCoy, 1979; Rosenzweig, 1992). To do that,

we carried out correlation tests between the log-transformed values of species richness and

geographic area, and between mean tip speciation rates and geographic area among deserts.

This allowed us to look for potential patterns to explain richness levels (e.g., highly diverse

deserts with larger areas and/or faster speciation than less diverse ones). We implemented all

the analyses, as well as data visualization, in the R environment version 3.6.3 (R Core Team,

2019), using the packages within 'tidyverse' (Wickham et al., 2019).

Regional richness and environmental variables

We developed a desert lizard species richness grid overlaying the distribution data (Roll et

al., 2017) onto an equal-area 10 arcminutes (~18 x 18 Km; ~0.17 degrees) resolution grid in a

Behrmann equal-area projection. We believe that this resolution applied to the Roll et al.

(2017) data captures the regional species pool, due to the relatively coarse nature of the

geographic ranges themselves. The Roll et al. (2017) distributions are bounding curves that

describe the overall regional extent of taxa but which provide no information about the local

occurrence of taxa in the landscape. Hence, the community for any given point in space will

include all species whose geographic ranges overlap broadly with the focal point, even if the

point itself is localized to (for example) a habitat patch that is wholly unsuitable for many of

the species. Furthermore, it is clear from previous work that species turnover in some of these

systems would occur at a finer scale than the 18 x 18 km cells of our grid even if the Roll et

al. (2017) ranges tracked local occurrences; Australia for example is characterized by a

mosaic landscape with high habitat heterogeneity, and desert lizard communities show

appreciable beta diversity at the scale of just a few hundred meters to several kilometres

(Pianka, 1969; Rabosky, Cowan, Talaba, & Lovette, 2011). We investigated the potential

influence of three types of environmental descriptors on desert lizard diversity patterns:

topography, climate, and productivity. The topography data was obtained from the Shuttle

Radar Topography Mission (SRTM) elevation data with 30 arcsec spatial resolution as

processed by WorldClim v2.1 (Fick & Hijmans, 2017). We downloaded climate summary

data for the period between 1970-2000 at 30 arcsec resolution from WorldClim v2.1 (Fick &

Hijmans, 2017). We focused on climate variables that might better differentiate desert

environments due to limiting effects on biodiversity. These include Annual Precipitation,

Annual Temperature, Mean Diurnal Range and Precipitation of the Driest Month.

Additionally, we downloaded the WorldClim data derived Aridity and Potential

Evapotranspiration data from the CGIAR Consortium for Spatial Information (v.2; Trabucco

& Zomer, 2019; Zomer et al., 2008). The productivity data is based on the normalized

difference vegetation index (NDVI) over a period between 1981 and 2015 summarized on the

NDVI3g time series (v.1; (Pinzon & Tucker, 2014). We downloaded and processed the NDVI

data with the 'gimms' R package (Detsch, 2020) by obtaining the monthly data with 5 arcmin

spatial resolution and summarizing the yearly data with maximum NDVI found. The final

dataset is the average of the year maximum NDVI for the period between 1981 and 2015.

We spatially aggregated the environmental variables to match the target spatial

resolution of the richness grid (10 arcmin ~ 18 x 18 km). We used the mean and standard

deviation as aggregating function and derived three variables from each downloaded

environmental descriptor: 1) mean aggregated value describing the average value at each

pixel in the target resolution, 2) within-pixel standard deviation describing the variation

within the target pixel, which represents regional heterogeneity, and 3) neighbourhood

standard deviation describing the surrounding variation to the target pixel. We chose two

neighbours, equivalent to a 5x5 pixels search window at the target resolution, to produce the

last variable. This variable describes how the surrounding environment might contribute to

the species richness, as an additional estimation of environmental spatial heterogeneity. All

spatial processing was performed with the 'raster' R library (Hijmans, 2020).

Correlation between species richness and environmental descriptors

In order to quantify the relation between species richness and environmental descriptors, we

used a non-parametric Kendall correlation score under a randomized sub-sampling

framework. We conducted per-desert correlation tests between species richness and each of

the environmental variables. For each environmental predictor, we conducted three different

tests, corresponding to the three aggregation types (mean value, standard deviation within

pixel, and neighbourhood standard deviation; see *Regional richness and environmental*

variables). For each test, we used 250 randomly selected pixels from each desert to calculate

the correlation and repeated the process 1,000 times. The advantages of this framework are

two-fold: first, it allows us to compare same sample size, independently of the size of the

desert; second, by randomly selecting scattered pixels over the desert area we decrease the

effect of spatial autocorrelation of both species richness and the environmental descriptors on

the correlation score. We note that this approach does not eliminate spatial autocorrelation,

but it minimises the effect of non-independence of the neighbours resulting from it. From the

set with the correlation scores from each of the 1,000 replicates, we calculated the average

correlation and the respective 95% confidence interval. All analyses and data visualization were implemented with the R packages 'raster' (Hijmans, 2020), 'rgdal' (Bivand, Keitt, & Rowlingson, 2019), and the data science collection 'tidyverse' (Wickham et al., 2019).

RESULTS

Biogeographic-scale lizard diversity

Out of 6,303 global lizard species in the Roll et al. (2017) dataset, we identified 58 species from the Atacama Desert, 316 from the Australian deserts, 41 from the Central Asian desert, 26 from the Gobi, 207 from the Kalahari, 191 from the North American deserts, 133 from the Persian desert and 249 from the Saharo-Arabian desert (Figure 2a). As expected, arid Australia's lizard richness is the highest among the deserts of the world. Diversity in the Saharo-Arabian desert and in the Kalahari is also noteworthy, with more than 200 species each. The Gobi and the Central Asian deserts have the lowest diversity overall, while the North American and Persian deserts have intermediate richness (Figure 2a).

Speciation rate and geographic area

We found significant differences in tip speciation rates (DR metric) across desert regions (Kruskal-Wallis chi-squared = 186.12, df = 7, p-value < 0.001), with highest rates in the Atacama desert and lowest in the Kalahari (Figure 2b). The post-hoc Wilcoxon test shows the pairwise differences between deserts (Table S3). We found no significant relationship between species richness and mean tip speciation rate (cor = -0.22, p = 0.60), since levels of species richness are not reflected in rate values (Figure 2a, b, d). Lizards inhabiting the most diverse deserts (Australian, Saharo-Arabian, Kalahari, American) do not show particularly

high speciation rates. This decoupling is also true for systems with low lizard diversity:

Atacama desert, one of the least diverse, is inhabited by taxa with the highest tip speciation

rates.

We found no consistent or significant effect of geographic area on biogeographic

species richness (cor = 0.41, p = 0.31) or tip speciation rates (cor = -0.67 , p = 0.07; Figure

2). Regarding the richness-area relationship, the Kalahari desert, which is one of the most

diverse, is the smallest one after the Atacama. In the other extreme, the Saharo-Arabian

desert, of unparalleled extension, has lower diversity than Australia, and only few more

species than much smaller deserts such as the North American and the Kalahari (Figure 2a, c,

e). In respect of the area-speciation rate relationship, large deserts such as the Saharo-Arabian

and the Australian do not present higher tip speciation rates than smaller ones such as the

Atacama or the North American desert (Figure 2b, c, f).

Evolutionary time hypothesis

The ancestral state reconstruction for desert occupancy (Figure S2) allowed us to estimate the

age at which desert lizard diversity started accumulating and how the number of arid-adapted

lineages has increased through time according to the reconstructed node probabilities (Figure

3a shows the rise of desert occupancy cumulative probability over time in the four most

diverse deserts). Among the deserts with higher species diversity, the Saharo-Arabian and the

Kalahari are the ones in which lineages started accumulating earlier, around 80-70 million

years ago (Ma). Lizard diversity in the North American and the Australian deserts started

increasing around 60 Ma, with North American diversity remaining at low levels until around

40-30 Ma. In contrast, arid Australian diversity shows a steep rise beginning at roughly 40

Ma, with peak rates of extant lineage accumulation at approximately 20 Ma. Thus, the lizard

diversity in Australia according to our reconstructed cumulative probability through time plot

does not appear before than in less diverse deserts, and nonetheless reaches unparalleled

levels after such steep increase. In fact, our correlation tests yielded a negative relationship

between the total richness of the deserts and the age in which lizard communities began to

accumulate in each desert (e.g., the times when the ancestral state probability was a 10% or a

30% of the total cumulative probability in each desert; Figure 3b and 3c, respectively).

Desert colonisation dynamics

Ancestral state reconstruction analysis of desert occupancy implies contrasting dynamics of

biome shifts in high- and low-diversity regions present essentially different dynamics of

biome shifts (Figure 4; Table S4). Specifically, we observed that lineages from the most

species-rich deserts (Australian, Saharo-Arabian, Kalahari, North American, and Persian)

have undergone shifts to non-arid climate states much more frequently than they have

adapted to arid conditions. On the contrary, less diverse deserts show a more balanced pattern

of lineages shifting from non-arid to arid biomes and vice versa (e.g., Central Asian and Gobi

deserts), or even an inverse pattern where the evolution of the arid state has been the

prevailing direction of change (e.g., Atacama desert). Additionally, we found a generally

balanced interchange between contiguous desert regions (e.g., between the Saharo-Arabian

and the Persian, and between the Persian and the Central Asian).

Regional lizard richness and the effect of environmental variables

In our regional lizard diversity grid (10 arcmin, 18 x 18 km) we found the expected

outstanding levels in Australia, with most of the arid Australian grid cells containing between

60 and 80 species (Figure 5). Regional diversity levels are medium in the North American

and the Kalahari deserts, and they are low for the rest of the arid systems, especially the Gobi

and the Atacama deserts, with few or no grid cells having more than 20 species. This result

indicates that the moderate-to-high overall (biogeographic) species richness for the Saharo-

Arabian desert (Figure 2a) is largely a result of the vastly greater area of that region and

geographical turnover. After controlling for sampling area (Figure 5), it is clear that regional

assemblages in Saharo-Arabia contain far fewer species than arid Australia and the Kalahari.

We found no consistent effects of the environmental descriptors or aggregation types

on regional (grid-cell) assemblages within individual deserts (Figure 6; Figure S3; Table S5,

Appendix 1). The most notable relationships included a positive correlation of average

potential evapotranspiration with lizard richness in the Central Asian and North American

deserts (Kendall's Tau 0.67 ± 0.04 and 0.60 ± 0.06, respectively); a positive correlation of

average annual temperature in the Central Asian and the Gobi deserts (Kendall's Tau

0.64 ± 0.05 and 0.63 ± 0.05, respectively); and a negative correlation of average precipitation of

the driest month in the Central Asian desert (Kendall's Tau -0.61 ± 0.05) (Figure 6; Table S5;

Appendix 1). Overall, the most noticeable pattern is the high disparity of the effects of each

environmental variable in different deserts, which shows differential roles in each region.

DISCUSSION

We investigated long-standing hypotheses about the factors driving desert lizard diversity.

Even though an unknown number of species remain to be described and there are inevitable

differences in the knowledge of biodiversity among regions that could bias our results

(Hughes et al., 2021; Meiri, 2016), we included the most recent available data on reptile

distributions, finding the overall same diversity patterns that have been reported for decades.

Our results corroborate the uneven distribution of lizards across the deserts of the

world and further highlight the exceptional lizard richness reported previously for arid

Australia (Pianka, 1969, 1986; Roll et al., 2017), and illustrate how this pattern holds at both

regional (Figure 5) and biogeographic (Figure 2) scales. Australia represents one extreme of

the lizard diversity continuum among the deserts of the world, where we find some systems

with medium levels of biogeographic and regional diversity, such as the North American or

the Kalahari deserts, and very low levels in the other extreme, represented by deserts such as

the Gobi and the Atacama. There are also cases of contrasting richness levels at different

scales: the Saharo-Arabian region harbours high richness at the biogeographic scale but is

relatively depauperate at the regional (grid-cell) scale. However, our results do not

corroborate some current hypotheses explaining species diversity patterns in the deserts.

Specifically, we find no evidence that recent speciation rates are fastest in the most species-

rich deserts (Figure 2) and no evidence that the most species-rich deserts have accumulated

species for longer (evolutionary time hypothesis; Figure 3). Furthermore, we found little

evidence that the differences in species richness among deserts at biogeographic and regional

scales are the result of the many environmental factors we considered (Figure 6). Rather than

supporting universal explanations for which environmental variables determine patterns of

lizard richness in the deserts, our results suggest that some of these factors might have been

relevant to shape lizard diversity within the deserts, but with a differential role amongst the

arid systems.

Speciation is one of the main processes that directly influence the number of species

at large scales (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012; Ricklefs, 1987; Schluter &

Pennell, 2017). The idea of an immediate relationship between speciation rate and species

richness (i.e., high levels of species richness are primarily the result of high speciation rates)

is a long-standing hypothesis (MacArthur, 1969; Mittelbach et al., 2007) and has proven

useful to explain the latitudinal diversity gradient in mammals (Rolland, Condamine, Jiguet,

& Morlon, 2014), but it has been recently challenged in global analyses of other major groups

such as fishes (Rabosky et al., 2018), birds (Jetz, Thomas, Joy, Hartmann, & Mooers, 2012;

Kennedy et al., 2014; Quintero & Jetz, 2018; Rabosky, Title, & Huang, 2015), and plants

(Igea & Tanentzap, 2020). The "faster speciation" hypothesis was proposed by (Pianka,

1972) to explain Australia's extreme lizard diversity, but our study represents the first formal

test of this idea. We found no relationship between high species richness and high tip

speciation rates, suggesting that the exceptionally high diversity in arid Australia is not driven

by faster speciation, at least by recent levels of speciation. Likewise, deserts with low species

richness (e.g., Gobi and Central Asian) did not yield particularly low tip speciation rates. In

fact, recent speciation can sometimes be faster in the regions with lowest diversity (Rabosky

et al., 2018). Such decoupling might reflect the greater importance of extinction or

immigration relative to speciation (Gaston & Blackburn, 1996; López-Estrada, Sanmartín,

García-París, & Zaldívar-Riverón, 2019; Meseguer & Condamine, 2020), and/or the

importance of differential carrying capacities and diversity-dependence in mediating overall

diversity patterns (Marshall, 2007; Mittelbach et al., 2007; Rabosky, 2009).

It is important to note that our analyses concern present-day speciation rates only and

we did not attempt to estimate net diversification rates. Our results on lineage accumulation

might suggest effectively increased net diversification rates in the Australian deserts relative

to other less diverse deserts. However, net diversification rates require estimates of

extinction, which is notoriously difficult in the absence of fossils (Rabosky, 2016). Moreover,

we did not attempt to reconstruct historical speciation rates, which might have been faster in

the past; recent work has emphasized historical (deep-time) reconstructions of speciation rate

from phylogenetic trees may be unreliable regardless of the inference method used to

estimate them (Louca & Pennell, 2020). Nonetheless, analyses of recent speciation rates can

still provide great insight into the causes of diversity gradients. Processes underlying patterns

of species richness can be affected by mechanisms that are time- and diversity-independent.

For example, faster speciation could result from the kinetic effects of temperature on

processes that favour reproductive isolation (e.g., effects on metabolism; Allen, Brown, &

Gillooly, 2002; Rohde, 1992) and as such, low-latitude deserts should be characterized by

faster speciation rates than higher-latitude deserts. The shifting sandridge hypothesis of

(Pianka, 1972) proposes that the mobile dune sheets within interior Australia serve as a

continued source of geographic isolation for non-dune taxa — in effect, creating a continued

"species pump" that will result in overall high rates of speciation for the arid Australian biota.

Our finding of unexceptional recent speciation rates for the Australian deserts rejects this and

other species-pump models for this system. Note that this does not mean that such shifting

sandridges are unimportant for speciation, merely that speciation rates as influenced by such

geomorphic/landscape processes are unlikely to serve as rate-limiting controls on species

richness.

Geographic area is one of the main factors that may affect broad-scale patterns of

species richness (Connor & McCoy, 1979). The concept of the species-area relationship,

derived from early works (Gleason, 1922, 1925) and developed later from the island

biogeography theory (MacArthur & Wilson, 1967), predicts a positive effect of geographic

area on both species diversity and speciation rates (Rosenzweig, 1992; Rosenzweig &

Sandlin, 1997). This has been supported for island and non-island systems (Kisel &

Barraclough, 2010; Losos & Schluter, 2000; L. G. Marshall, Webb, Sepkoski, & Raup, 1982;

Sepkoski, 1976; Wagner, Harmon, & Seehausen, 2014), although patterns reported for some

systems have likely been confounded by assumptions that speciation rates are invariant

through time (Rabosky & Glor, 2010). Moreover, some authors have argued against

geographic area being the determinant factor on species diversity (Chown & Gaston, 2000;

Rohde, 1992, 1998). We found no significant effect of area on species richness, although the

correlation was positive (Figure 2e). It is possible that geographic area contributes to species

richness, but the relationship is incredibly noisy at best: the largest desert by far is the

500 Saharo-Arabian, which contains substantially lower diversity than Australia and roughly

501 equal diversity to the much smaller Kalahari. Conversely, the tiny Atacama desert – by far

502 the smallest in the dataset – has greater diversity than two much larger deserts.

503  Alternatively, evolutionary time may generate diversity patterns in which species

504 richness is decoupled from geographic area and other factors (Rohde, 1986; Stephens &

505 Wiens, 2003), although some evidence supports the combined effect of area and time (Fine &

506 Ree, 2006; Jetz & Fine, 2012). Age differences between geographic regions or biomes has

507 long been thought to be a potential contributor to differences in species richness (Wallace,

508 1878), for the simple reason that older, less-disturbed geographic regions have longer periods

509 of time for diversity to accumulate (Fischer, 1960; Mittelbach et al., 2007). Our results argue

510 against any simple role for time in explaining the observed heterogeneity in desert lizard

511 richness and even suggest a negative relationship between richness and age (Figure 3).

512 Remarkably, our results suggest that Australian lizard diversity began accumulating more

513 than ten million years after the onset of lineage accumulation in less-diverse deserts, like the

514 Saharo-Arabian and the Kalahari (Figure 3). Thus, greater time for diversification is unlikely

515 to account for the extreme lizard richness observed in arid Australia, and other factors are

516 likely responsible for the steep increase in lizard diversity observed in this region (e.g.,

517 historical diversification dynamics). Note that these results do not rely on accurately inferring

518 the true timing of squamate diversification in each region, but only on the phylogenetic age of

519 the extant biota. For example, now-extinct lineages of squamates may have undergone

520 extensive diversification in arid or semi-arid Australia many millions of years prior to our

521 reconstructions (Oliver & Hugall, 2017), but it is still true that the entire extant Australian

522 arid zone fauna was able to accumulate in a much briefer period of time.

523  It is worth noting that our estimation of the age of arid-adapted lineages does not aim

524 to serve as an absolute date for desert origination. The age of the deserts considered here is a

complex and contentious issue (Swezey, 2006). The onset of arid conditions might have

greatly preceded the formation of many modern-day arid landscapes (Pillans, 2018; Schuster

et al., 2006), and hence could potentially explain the presence of arid-adapted lineages long

before the estimated geological age of the desert (Byrne et al., 2008; Carranza, Arnold,

Geniez, Roca, & Mateo, 2008). Therefore, our estimation of the age of arid-adapted lineages

does not aim to serve as an absolute date for desert origination, but instead as a relative time

framework to compare the biotas inhabiting different desert systems.

Regarding colonisation dynamics, we found a consistent pattern distinguishing high-

and low-richness deserts (Figure 4). Species-rich deserts (e.g., Australian, Saharo-Arabian,

Kalahari, North American) are characterized by high levels of phylogenetic transitions to

non-arid conditions compared to shifts towards arid climate tolerance. This contrasts with

species-poor deserts, such as the Gobi or the Central Asian, where evolutionary shifts from

non-arid to arid conditions and vice versa are balanced. Further, in the case of the Atacama,

we observed a net excess of biome transitions into the region (Figure 4; Table S4). This

suggests essentially different processes of community assembly in regions with high and low

diversity. Based on these results, communities in high-diversity systems build through a few

colonisation events with high levels of subsequent intra-desert diversification, with many

lineages of those clades secondarily adapting to non-desert conditions. Less diverse deserts,

on the contrary, seem to be characterized by a relatively regular lineage interchange with

regions of lower aridity, without extensive radiations within the deserts. These differences

might be related to intrinsic system-specific characteristics that may offer more opportunities

for speciation in some regions, resulting in high diversity, while some other regions of low

diversity might serve as evolutionary 'dead-ends', with limited opportunities for

diversification within the desert.

Despite theoretical and empirical support linking climate and topography to species

richness in general (Currie, 1991; Francis & Currie, 2003; Jetz & Rahbek, 2002; Kreft & Jetz,

2007; Qian, 2010; Skeels, Esquerré, & Cardillo, 2020; Wright, 1983), we found ambiguous

support for the effects of environmental variables on desert lizard diversity at both regional

and biogeographic scales. Our per-desert environment-richness analyses show differential

effects of the environmental descriptors on lizard richness across deserts (Figure 6). We

found a considerable correlation of richness in some deserts with several variables, in

particular average evapotranspiration in Central Asia and North America, average annual

temperature in Central Asia and the Gobi, and (negative) average precipitation of the driest

month in Central Asia. Thus, the impact of the environment on the distribution of lizard

diversity in the Central Asian desert seems stronger than in other arid systems. It is

interesting to note the shared patterns in the Central Asian and the Gobi desert with respect to

most variables. Both deserts have similar levels of diversity (very low biogeographic and

regional diversity), and this might indicate a differential role of climate and other factors

between high- and low-diversity systems, as also suggested by our results on desert

colonisation dynamics (Figure 4). In the Atacama desert, environmental heterogeneity (e.g.,

surrounding heterogeneity in precipitation, productivity and aridity) appears to play a

significant role in the distribution of diversity. Average elevation is more related with the

higher diversity in Australia than any other variable, and its importance for Australian lizards

is remarkable in comparison with other deserts, while in the Kalahari and the Atacama

deserts the elevational heterogeneity seems more relevant. Additionally, in the Kalahari,

heterogeneity plays a rather constant role, although without strong correlations with richness.

In the Saharo-Arabian region, on the contrary, there are not remarkable effects of the

variables tested on lizard distribution. It should be noticed here that we only tested recent

environmental factors. It is likely that deep-time environmental dynamics (e.g., when most of

the desert diversity originated and assembled) had an important role in shaping lizard communities by affecting the stability of the arid systems through time (as well as historical diversification). In fact, the impact of biome fragmentation, expansion and contraction cycles might have been particularly relevant in the deserts (Barbolini et al., 2020; deMenocal, 1995; Hernández-Fernández & Vrba, 2005).

We did not find consistent effects of climate, productivity, and topography on species richness across deserts, but it is possible that other, unmeasured environmental variables and their interactions provide an important control on broad-scale richness. Our results, hence, are consistent with a scenario in which desert lizard diversity at regional-to-biogeographic scales depends mainly on system-specific ecological dynamics. Factors such as extinction, habitat heterogeneity, ecological preferences, and ecological interactions with other taxa, might be playing a relevant role in shaping the distribution of current lizard diversity in the deserts. Our results provide evidence against several general hypotheses proposed to explain desert lizard diversity patterns (age; evolutionary rate). However, we only studied part of the evolutionary rates hypothesis (recent speciation), and we speculate that the relationship between richness and environment may not be adequately captured with our simplified set of eight explanatory variables. Ultimately, our results emphasise the complexity of evolutionary and ecological processes that determine large-scale diversity patterns. Disentangling their underlying causes remains an essential task and one that is necessary in order to face the challenges imposed by the rapid pace of climatic and other change that is disrupting ecosystems worldwide.

DATA AVAILABILITY STATEMENT

All the data and scripts produced in this work will be made available at an online repository (e.g. GitHub), which will be updated and made public upon acceptance of the manuscript.

REFERENCES

Allen, A. P., Brown, J. H., & Gillooly, J. F. (2002). Global biodiversity, biochemical kinetics,

and the energetic-equivalence rule. *Science*, *297*, 1545–1548.

Antão, L. H., Bates, A. E., Blowes, S. A., Waldock, C., Supp, S. R., Magurran, A. E., …

Schipper, A. M. (2020). Temperature-related biodiversity change across temperate

marine and terrestrial systems. *Nature Ecology and Evolution*, *4*, 927–933.

Badgley, C., Smiley, T. M., Terry, R., Davis, E. B., DeSantis, L. R. G., Fox, D. L., …

Yanites, B. J. (2017). Biodiversity and Topographic Complexity: Modern and

Geohistorical Perspectives. *Trends in Ecology and Evolution*, *32*, 211–226.

Barbolini, N., Woutersen, A., Dupont-Nivet, G., Silvestro, D., Tardif, D., Coster, P. M. C., …

Hoorn, C. (2020). Cenozoic evolution of the steppe-desert biome in Central Asia.

Science Advances, *6*, eabb8227.

Belmaker, J., & Jetz, W. (2015). Relative roles of ecological and energetic constraints,

diversification rates and region history on global species richness gradients. *Ecology*

Letters, *18*, 563–571.

Bivand, R., Keitt, T., & Rowlingson, B. (2019). *rgdal: Bindings for the 'Geospatial' Data*

Abstraction Library. Retrieved from https://cran.r-project.org/package=rgdal

Byrne, M., Yeates, D. K., Joseph, L., Kearney, M., Bowler, J., Williams, M. A. J., …

Wyrwoll, K. H. (2008). Birth of a biome: Insights into the assembly and maintenance

of the Australian arid zone biota. *Molecular Ecology*, *17*, 4398–4417.

Cantalapiedra, J. L., Domingo, M. S., & Domingo, L. (2018). Multi-scale interplays of biotic

and abiotic drivers shape mammalian sub-continental diversity over millions of years.

Scientific Reports, *8*, 13413.

Carranza, S., Arnold, E. N., Geniez, P., Roca, J., & Mateo, J. A. (2008). Radiation, multiple

dispersal and parallelism in the skinks, *Chalcides* and *Sphenops* (Squamata:

Scincidae), with comments on *Scincus* and *Scincopus* and the age of the Sahara

Desert. *Molecular Phylogenetics and Evolution*, *46*, 1071–1094.

Chang, J., Rabosky, D. L., & Alfaro, M. E. (2020). Estimating Diversification Rates on

Incompletely Sampled Phylogenies: Theoretical Concerns and Practical Solutions.

Systematic Biology, *69*, 602–611.

Chown, S. L., & Gaston, K. J. (2000). Areas, cradles and museums: The latitudinal gradient

in species richness. *Trends in Ecology and Evolution*, *15*, 311–315.

Connor, E. F., & McCoy, E. D. (1979). The Statistics and Biology of the Species-Area

Relationship. *The American Naturalist*, *113*, 791–833.

Couvreur, T. L. P. (2015). Odd man out: Why are there fewer plant species in African rain

forests? *Plant Systematics and Evolution*, *301*, 1299–1313.

Cracraft, J. (1992). Explaining Patterns of Biological Diversity: Integrating Causation at

Different Spatial and Temporal Scales. In N. Eldredge (Ed.), *Systematics, Ecology*

and the Biodiversity Crisis (pp. 59–76). Columbia University Press.

Crouch, N. M. A., Capurucho, J. M. G., Hackett, S. J., & Bates, J. M. (2019). Evaluating the

contribution of dispersal to community structure in Neotropical passerine birds.

Ecography, *42*, 390–399.

Currie, D. J. (1991). Energy and large-scale patterns of animal and plant species richness. *The*

American Naturalist, *137*, 27–49.

deMenocal, P. B. (1995). Plio-Pleistocene African Climate. *Science*, *270*, 53–59.

Detsch, F. (2020). *gimms: Download and Process GIMMS NDVI3g Data*. Retrieved from

https://cran.r-project.org/package=gimms

Donoghue, M. J., & Edwards E.J. (2014). Biome shifts and niche evolution in plants. *Annual*

Review of Ecology and Systematics, *45*, 547–572.

Duellman, W. E. (2005). *Cusco Amazónico*. Comstock Pub. Associates.

Edler, D., Guedes, T., Zizka, A., Rosvall, M., & Antonelli, A. (2017). Infomap bioregions:

Interactive mapping of biogeographical regions from species distributions. *Systematic*

Biology, *66*, 197–204.

Etienne, R. S., Cabral, J. S., Hagen, O., Hartig, F., Hurlbert, A. H., Pellissier, L., … Storch,

D. (2019). A minimal model for the latitudinal diversity gradient suggests a dominant

role for ecological limits. *American Naturalist*, *194*, E122–E133.

Fick, S. E., & Hijmans, R. J. (2017). WorldClim 2: New 1-km spatial resolution climate

surfaces for global land areas. *International Journal of Climatology*, *37*, 4302–4315.

Fine, P. V. A., & Ree, R. H. (2006). Evidence for a time-integrated species-area effect on the

latitudinal gradient in tree diversity. *American Naturalist*, *168*, 796–804.

Fischer, A. G. (1960). Latitudinal variations in organic diversity. *Evolution*, *14*, 64–81.

Francis, A. P., & Currie, D. J. (2003). A globally consistent richness-climate relationship for

angiosperms. *The American Naturalist*, *161*, 523–536.

Garcia-Porta, J., Irisarri, I., Kirchner, M., Rodríguez, A., Kirchhof, S., Brown, J. L., …

Wollenberg Valero, K. C. (2019). Environmental temperatures shape thermal

physiology as well as diversification and genome-wide substitution rates in lizards.

Nature Communications, *10*, 4077.

Gaston, K. J., & Blackburn, T. M. (1996). The tropics as a museum of biological diversity:

An analysis of the New World avifauna. *Proceedings of the Royal Society B:*

Biological Sciences, *263*, 63–68.

Gleason, H. A. (1922). On the Relation Between Species and Area. *Ecology*, *3*, 158–162.

Gleason, H. A. (1925). Species and Area. *Ecology*, *6*, 66–74.

Harmon, L. J., Weir, J. T., Brock, C. D., Glor, R. E., & Challenger, W. (2008). GEIGER:

Investigating evolutionary radiations. *Bioinformatics*, *24*, 129–131.

Hernández-Fernández, M., & Vrba, E. S. (2005). Macroevolutionary processes and biomic

specialization: Testing the resource-use hypothesis. *Evolutionary Ecology*, *19*, 199–

219.

Hijmans, R. J. (2020). *raster: Geographic Data Analysis and Modeling*. Retrieved from

https://cran.r-project.org/package=raster

Hua, X., & Bromham, L. (2020). Modeling colonization rates over time: Generating null

models and testing model adequacy in phylogenetic analyses of species assemblages.

Evolution, *74*, 2605–2616.

Hughes, A., Orr, M., Ma, K., Costello, M., Waller, J., Provoost, P., … Qiao, H. (2021).

Sampling biases shape our view of the natural world. *Ecography*, *44*, 1–11.

Igea, J., & Tanentzap, A. J. (2020). Angiosperm speciation cools down in the tropics.

Ecology Letters, *23*, 692–700.

Jablonski, D., Huang, S., Roy, K., & Valentine, J. W. (2017). Shaping the Latitudinal

Diversity Gradient: New Perspectives from a Synthesis of Paleobiology and

Biogeography. *The American Naturalist*, *189*, 1–12.

James, C. D., & Shine, R. (2000). Why are there so many coexisting species of lizards in

Australian deserts? *Oecologia*, *125*, 127–141.

Jetz, W., & Fine, P. V. A. (2012). Global gradients in vertebrate diversity predicted by

historical area-productivity dynamics and contemporary environment. *PLoS Biology*,

10, e1001292.

Jetz, W., & Rahbek, C. (2002). Geographic range size and determinants of avian species

richness. *Science*, *297*, 1548–1551.

Jetz, W., Thomas, G. H., Joy, J. B., Hartmann, K., & Mooers, A. O. (2012). The global

diversity of birds in space and time. *Nature*, *491*, 444–448.

Kelt, D. A., Brown, J. H., Heske, E. J., Marquet, P. A., Morton, S. R., Reid, J. R. W., …

Shenbrot, G. (1996). Community Structure of Desert Small Mammals: Comparisons

Across Four Continents. *Ecology*, *77*, 746–761.

Kennedy, J. D., Wang, Z., Weir, J. T., Rahbek, C., Fjeldså, J., & Price, T. D. (2014). Into and

out of the tropics: The generation of the latitudinal gradient among New World

passerine birds. *Journal of Biogeography*, *41*, 1746–1757.

Kisel, Y., & Barraclough, T. G. (2010). Speciation has a spatial scale that depends on levels

of gene flow. *The American Naturalist*, *175*, 316–334.

Kreft, H., & Jetz, W. (2007). Global patterns and determinants of vascular plant diversity.

Proceedings of the National Academy of Sciences of the United States of America,

104, 5925–5930.

Latham, R. E., & Ricklefs, R. E. (1993). Continental comparisons of temperate-zone tree

species diversity. In R. E. Ricklefs & D. Schluter (Eds.), *Species Diversity in*

Ecological Communities: Historical and Geographical Perspectives (pp. 294–313).

University of Chicago Press.

López-Estrada, E. K., Sanmartín, I., García-París, M., & Zaldívar-Riverón, A. (2019). High

extinction rates and non-adaptive radiation explains patterns of low diversity and

extreme morphological disparity in North American blister beetles (Coleoptera,

Meloidae). *Molecular Phylogenetics and Evolution*, *130*, 156–168.

Losos, J. B., & Schluter, D. (2000). Analysis of an evolutionary species-area relationship.

Nature, *408*, 847–850.

Louca, S., & Pennell, M. W. (2020). Extant timetrees are consistent with a myriad of

diversification histories. *Nature*, *580*, 502–505.

721 MacArthur, R. H. (1969). Patterns of communities in the tropics. *Biological Journal of the*

722   *Linnean Society*, *1*, 19–30.

723 MacArthur, R. H., & Wilson, E. O. (1967). *The theory of island biogeography* (Vol. 1).

724   Princeton University Press.

725 Mahler, D. L., Revell, L. J., Glor, R. E., & Losos, J. B. (2010). Ecological opportunity and

726   the rate of morphological evolution in the diversification of Greater Antillean anoles.

727   *Evolution*, *64*, 2731–2745.

728 Marshall, C. R. (2007). Explaining Latitudinal Diversity Gradients. *Science*, *317*, 451–452.

729 Marshall, L. G., Webb, S. D., Sepkoski, J. J., & Raup, D. M. (1982). Mammalian evolution

730   and the Great American Interchange. *Science*, *215*, 1351–1357.

731 Meiri, S. (2016). Small, rare and trendy: Traits and biogeography of lizards described in the

732   21st century. *Journal of Zoology*, *299*, 251–261.

733 Menéndez, I., Gómez Cano, A. R., Cantalapiedra, J. L., Peláez-Campomanes, P., Álvarez-

734   Sierra, M. Á., & Hernández Fernández, M. (2021). A multi-layered approach to the

735   diversification of squirrels. *Mammal Review*, *51*, 66–81.

736 Meseguer, A. S., & Condamine, F. L. (2020). Ancient tropical extinctions at high latitudes

737   contributed to the latitudinal diversity gradient. *Evolution*, *74*, 1966–1987.

738 Mittelbach, G. G., Schemske, D. W., Cornell, H. V., Allen, A. P., Brown, J. M., Bush, M. B.,

739   … Turelli, M. (2007). Evolution and the latitudinal diversity gradient: Speciation,

740   extinction and biogeography. *Ecology Letters*, *10*, 315–331.

741 Morton, S. R., & James, C. D. (1988). The Diversity and Abundance of Lizards in Arid

742   Australia: A New Hypothesis. *The American Naturalist*, *132*, 237–256.

743 Oliver, P. M., & Hugall, A. F. (2017). Phylogenetic evidence for mid-Cenozoic turnover of a

744   diverse continental biota. *Nature Ecology and Evolution*, *1*, 1896–1902.

Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N.,

Underwood, E. C., … Kassem, K. R. (2001). Terrestrial Ecoregions of the World: A

New Map of Life on Earth. *BioScience*, *51*, 933–933.

Parsons, A. J., & Abrahams, A. D. (1994). Geomorphology of Desert Environments. In A. J.

Parsons & A. D. Abrahams (Eds.), *Geomorphology of Desert Environments* (pp. 3–8).

Chapman and Hall.

Pianka, E. R. (1969). Habitat specificity, speciation, and species density in Australian desert

lizards. *Ecology*, *50*, 498–502.

Pianka, E. R. (1971). Lizard species density in the Kalahari desert. *Ecology*, *52*, 1024–1029.

Pianka, E. R. (1972). Zoogeography and speciation of Australian desert lizards: An

ecological perspective. *Copeia*, *1*, 127–145.

Pianka, E. R. (1985). Some intercontinental comparisons of desert lizards. *National*

Geographic Research, *1*, 490–504.

Pianka, E. R. (1986). *Ecology and Natural History of Desert Lizards*. Princeton University

Press.

Pianka, E. R. (1989). Desert lizard diversity: Additional comments and some data. *The*

American Naturalist, *134*, 344–364.

Pillans, B. J. (2018). Seeing Red: Some Aspects of the Geological and Climatic History of the

Australian Arid Zone. In H. Lambers (Ed.), *On the Ecology of Australia's Arid Zone*

(pp. 5–43). Springer International Publishing.

Pinzon, J. E., & Tucker, C. J. (2014). A non-stationary 1981-2012 AVHRR NDVI3g time

series. *Remote Sensing*, *6*, 6929–6960.

Qian, H. (2010). Environment-richness relationships for mammals, birds, reptiles, and

amphibians at global and regional scales. *Ecological Research*, *25*, 629–637.

Qian, H., & Ricklefs, R. E. (2000). Large-scale processes and the Asian bias in species

diversity of temperate plants. *Nature*, *407*, 180–182.

Quental, T. B., & Marshall, C. R. (2010). Diversity dynamics: Molecular phylogenies need

the fossil record. *Trends in Ecology and Evolution*, *25*, 434–441.

Quintero, I., & Jetz, W. (2018). Global elevational diversity and diversification of birds.

Nature, *555*, 246–250.

R Core Team. (2019). *R: A Language and Environment for Statistical Computing*. Vienna,

Austria: R Foundation for Statistical Computing.

Rabosky, D. L., & Glor, R. E. (2010). Equilibrium speciation dynamics in a model adaptive

radiation of island lizards. *Proceedings of the National Academy of Sciences*, *107*,

22178–22183.

Rabosky, D. L. (2009). Ecological limits and diversification rate: Alternative paradigms to

explain the variation in species richness among clades and regions. *Ecology Letters*,

12, 735–743.

Rabosky, D. L. (2015). No substitute for real data: A cautionary note on the use of

phylogenies from birth-death polytomy resolvers for downstream comparative

analyses. *Evolution*, *69*, 3207–3216.

Rabosky, D. L. (2016). Challenges in the estimation of extinction from molecular

phylogenies: A response to Beaulieu and O'Meara. *Evolution*, *70*, 218–228.

Rabosky, D. L., Chang, J., Title, P. O., Cowman, P. F., Sallan, L., Friedman, M., … Alfaro,

M. E. (2018). An inverse latitudinal gradient in speciation rate for marine fishes.

Nature, *559*, 392–395.

Rabosky D. L., Cowan M. A., Talaba A. L., & Lovette I. J. (2011). Species interactions

mediate phylogenetic community structure in a hyperdiverse lizard assemblage from

arid Australia. *The American Naturalist*, *178*, 579–595.

Rabosky, D. L, & Hurlbert, A. H. (2015). Species Richness at Continental Scales Is

Dominated by Ecological Limits. *The American Naturalist*, *185*, 572–583.

Rabosky, D. L., Title, P. O., & Huang, H. (2015). Minimal effects of latitude on present-day

speciation rates in New World birds. *Proceedings of the Royal Society B: Biological*

Sciences, *282*, 20142889.

Rabosky, D. L., von May, R., Grundler, M. C., & Davis Rabosky, A. R. (2019). The Western

Amazonian richness gradient for squamate reptiles: Are There Really Fewer Snakes

and Lizards in Southwestern Amazonian Lowlands? *Diversity*, *11*, 199.

Revell, L. J. (2012). phytools: An R package for phylogenetic comparative biology (and other

things). *Methods in Ecology and Evolution*, *3*, 217–223.

Ricklefs, R. E., & Latham, R. E. (1993). Global patterns of diversity in mangrove floras. In

R. E. Ricklefs & D. Schluter (Eds.), *Species Diversity in Ecological Communities:*

Historical and Geographical Perspectives (pp. 215–229). University of Chicago

Press.

Ricklefs, R. E. (1987). Community Diversity: Relative roles of local and regional processes.

Science, *235*, 167–171.

Ricklefs, R. E. (2004). A comprehensive framework for global patterns in biodiversity.

Ecology Letters, *7*, 1–15.

Ricklefs, R. E., & Schluter, D. (1993). *Species diversity in ecological communities:*

Historical and geographical perspectives. University of Chicago Press Chicago.

Rocha, J. L., Godinho, R., Brito, J. C., & Nielsen, R. (2021). Life in Deserts: The Genetic

Basis of Mammalian Desert Adaptation. *Trends in Ecology and Evolution*, *36*, 637–

650.

Rohde, K. (1986). Differences in species diversity of Monogenea between the Pacific and

Atlantic Oceans. *Hydrobiologia*, *137*, 21–28.

Rohde, Klaus. (1992). Latitudinal Gradients in Species Diversity: The Search for the Primary

820   Cause. *Oikos*, *65*, 514–527.

821   Rohde, Klaus. (1998). Latitudinal Gradients in Species Diversity. Area Matters, but How

822   Much? *Oikos*, *82*, 184–184.

823   Roll, U., Feldman, A., Novosolov, M., Allison, A., Bauer, A. M., Bernard, R., … Meiri, S.

824   (2017). The global distribution of tetrapods reveals a need for targeted reptile

825   conservation. *Nature Ecology and Evolution*, *1*, 1677–1682.

826   Rolland, J., Condamine, F. L., Jiguet, F., & Morlon, H. (2014). Faster Speciation and

827   Reduced Extinction in the Tropics Contribute to the Mammalian Latitudinal Diversity

828   Gradient. *PLoS Biology*, *12*, e1001775.

829   Rosenzweig, M. L. (1992). Species Diversity Gradients: We Know More and Less Than We

830   Thought. *Journal of Mammalogy*, *73*, 715–730.

831   Rosenzweig, M. L., & Sandlin, E. A. (1997). Species Diversity and Latitudes: Listening to

832   Area's Signal. *Oikos*, *80*, 172–176.

833   Rosvall, M., & Bergstrom, C. T. (2008). Maps of random walks on complex networks reveal

834   community structure. *Proceedings of the National Academy of Sciences of the United

835   States of America*, *105*, 1118–1123.

836   Schemske, D. W., & Mittelbach, G. G. (2017). "Latitudinal Gradients in Species Diversity":

837   Reflections on Pianka's 1966 Article and a Look Forward. *American Naturalist*, *189*,

838   599–603.

839   Schluter, D., & Pennell, M. W. (2017). Speciation gradients and the distribution of

840   biodiversity. *Nature*, *546*, 48–55.

841   Schuster, M., Duringer, P., Ghienne, J. F., Vignaud, P., Mackaye, H. T., Likius, A., & Brunet,

842   M. (2006). The age of the Sahara Desert. *Science*, *311*, 821.

Sepkoski, J. J. (1976). Species diversity in the Phanerozoic: Species-area effects.

Paleobiology, *2*, 298–303.

Simberloff, D. S. (1974). Equilibrium Theory of Island Biogeography and Ecology. *Annual*

Review of Ecology and Systematics, *5*, 161–182.

Skeels, A., Esquerré, D., & Cardillo, M. (2020). Alternative pathways to diversity across

ecologically distinct lizard radiations. *Global Ecology and Biogeography*, *29*, 454–

469.

Stein, A., Beck, J., Meyer, C., Waldmann, E., Weigelt, P., & Kreft, H. (2015). Differential

effects of environmental heterogeneity on global mammal species richness. *Global*

Ecology and Biogeography, *24*, 1072–1083.

Stein, A., Gerstner, K., & Kreft, H. (2014). Environmental heterogeneity as a universal driver

of species richness across taxa, biomes and spatial scales. *Ecology Letters*, *17*, 866–

880.

Stephens, P. R., & Wiens, J. J. (2003). Explaining species richness from continents to

communities: The time-for-speciation effect in emydid turtles. *American Naturalist*,

161, 112–128.

Swezey, C. S. (2006). Revisiting the age of the Sahara Desert. *Science*, *312*, 1138–1139.

Thomas, G. H., Hartmann, K., Jetz, W., Joy, J. B., Mimoto, A., & Mooers, A. O. (2013).

PASTIS: An R package to facilitate phylogenetic assembly with soft taxonomic

inferences. *Methods in Ecology and Evolution*, *4*, 1011–1017.

Title, P. O., & Rabosky, D. L. (2019). Tip rates, phylogenies and diversification: What are we

estimating, and how good are the estimates? *Methods in Ecology and Evolution*, *10*,

821–834.

Tonini, J. F. R., Beard, K. H., Ferreira, R. B., Jetz, W., & Pyron, R. A. (2016). Fully-sampled

phylogenies of squamates reveal evolutionary patterns in threat status. *Biological*

Conservation, *204*, 23–31.

Trabucco, A., & Zomer, R. (2019). Global Aridity Index and Potential Evapotranspiration

(ET0)   Climate   Database   v2.   *CGIAR   Consort   Spat   Inf.*

https://doi.org/10.6084/m9.figshare.7504448.v3

Uetz, P., Freed, P., & Hošek, J. (2021). The Reptile Database. Retrieved from

http://www.reptile-database.org

Valente, L. M., & Vargas, P. (2013). Contrasting evolutionary hypotheses between two

mediterranean-climate floristic hotspots: The Cape of southern Africa and the

Mediterranean Basin. *Journal of Biogeography*, *40*, 2032–2046.

Velasco, J. A., Villalobos, F., Diniz-Filho, J. A. F., Algar, A. C., Flores-Villela, O., Köhler,

G., … Martinez-Meyer, E. (2018). Climatic and evolutionary factors shaping

geographical gradients of species richness in *Anolis<i/> lizards. Biological Journal of*

the Linnean Society, 123, 615–627.

Vilhena, D. A., & Antonelli, A. (2015). A network approach for identifying and delimiting

biogeographical regions. Nature Communications, 6, 6848.

Wagner, C. E., Harmon, L. J., & Seehausen, O. (2014). Cichlid species-area relationships

are shaped by adaptive radiations that scale with area. Ecology Letters, 17, 583–592.

Wallace, A. R. (1878). Tropical nature, and other essays. Macmillan and Company.

Wang, L. G., Lam, T. T. Y., Xu, S., Dai, Z., Zhou, L., Feng, T., … Yu, G. (2020). Treeio: An R

Package for Phylogenetic Tree Input and Output with Richly Annotated and

Associated Data. Molecular Biology and Evolution, 37, 599–603.

Weber, M. G., Wagner, C. E., Best, R. J., Harmon, L. J., & Matthews, B. (2017). Evolution in

a Community Context: On Integrating Ecological Interactions and Macroevolution.

Trends in Ecology and Evolution, 32, 291–304.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., ... Yutani, H.

(2019). Welcome to the Tidyverse. Journal of Open Source Software, 4, 1686.

Wiens, J. J. (2011). The causes of species richness patterns across space, time, and clades

and the role of 'ecological limits'. Quarterly Review of Biology, 86, 75–96.

Wright, D. H. (1983). Species-Energy Theory: An Extension of Species-Area Theory. Oikos,

41, 496–506.

Zomer, R. J., Trabucco, A., Bossio, D. A., & Verchot, L. V. (2008). Climate change

mitigation: A spatial analysis of global land suitability for clean development

mechanism afforestation and reforestation. Agriculture, Ecosystems and

Environment, 126, 67–80.

FIGURES

Figure 1. *Desert systems considered in this study. We focused on regions with aridity indices*

(AI) < 0.2, which describe systems that are formally "arid" or "hyper-arid" but which

exclude "semi-arid" (Parsons and Abrahams, 1994; Rocha et al., 2021). Map in an equal-

area Behrmann projection.

***Figure 2**. Species diversity, geographic area, and tip speciation rate across desert systems.*

The top row shows the characterization of arid systems: (a) biogeographic lizard richness,

(b) tip speciation rates, and (c) geographic area. The bottom row shows the correlation

between (d) speciation rates (mean DR) and lizard diversity in the deserts, (e) geographic

area and lizard diversity, and (f) geographic area and speciation rates. There is no clear

trend in the relationship between the three attributes: highly diverse deserts (e.g., Australian)

are not characterized by especially high speciation rates, while several smaller and less

diverse deserts (e.g., Atacama) show higher speciation rates than more diverse ones (e.g.,

Kalahari).

Figure 3. *Evolutionary time and species accumulation across major desert regions. a)*
Reconstructed lineage accumulation curves for the four most diverse arid systems:
Australian, Saharo-Arabian, Kalahari, and North American deserts. Lineage richness is
represented by the cumulative probability of the focal state (arid region) summed over all
nodes in the phylogeny that diverged prior to a given time point. This method necessarily
underestimates true richness at a given time as it cannot accommodate extinct (unobserved)
lineages, but nonetheless tracks the temporal accumulation of present-day lineages (extant)
through time within regions. Panels in the right show the correlation between richness and
the time in which each desert had (b) the 10% (ages10) and (c) the 30% (ages30) of their
total cumulative probability in the ancestral reconstruction. Remarkably, there is a negative
correlation between the age in which (extant) diversity started accumulating and the current
number of species in the deserts. Namely, the most-species rich region (Australia) is
associated with the most recent faunal buildup, thus rejecting the hypothesis that Australia's
exceptional diversity can be explained by greater time for diversity to accumulate.

Figure 4. Desert colonisation dynamics: frequency of biome shifts of lineages undergoing

new climate tolerances, from arid (desert systems) to non-arid ("OUT"), in each desert

region. Shifts in climate tolerance, as well as lineage interchange between deserts, are

depicted with directional arrows. Arrow thickness and colour intensity represent the number

of events. Circle size is proportional to the number of species within each desert. Information

of events into and out of the deserts are extracted from the ancestral reconstruction analysis

(Figure S2; Table S4).

Figure 5. Regional lizard species richness (10 arc-minute grid cell) for the deserts of the

world, with histograms of grid-cell richness for each system. Species richness of regional

lizard assemblages from Australia is far greater than in any other arid region, and the system

is thus exceptionally diverse at local (Pianka, 1969), regional, and biogeographic (Figure 2)

scales. Map in an equal-area Behrmann projection.

Figure 6. Correlation coefficient (mean and 95% confidence interval from 1,000 bootstrap replicates) between each environmental descriptor and regional lizard richness, for each arid region separately. "Average" column gives the correlations between species richness and the mean of the environmental variable per cell. "Heterogeneity within" and "Surrounding heterogeneity" give the correlation between species richness and two measures of variability in environmental conditions per cell: the standard deviation of the variable within the cell, and the standard deviation of the variable in the surrounding cells, respectively. There are no consistent effects of the environmental variables across deserts, and no clear relationships between environment and richness, although some variables seem relevant in the shaping of lizard diversity in certain deserts.

CAPTIONS FOR SUPPLEMENTARY FIGURES

Figure S1. Number of lizard species in each arid system under different threshold values for categorization of species into deserts: 10_10, 10_20, 15_15, etc. The number before the underscore indicates the percentage of the species distribution that needs to be within a desert to be categorized into that desert, and the number after the underscore indicates the percentage of a desert's extension that needs to be occupied by a species for the species to be categorized into that desert. If one of those criteria is met, a species is categorized into the arid system.

Figure S2. Ancestral reconstruction of desert occupancy on the Squamate phylogenetic tree from Tonini et al. (2016) excluding species without genetic and distribution data. Colors at the tips indicate the arid system(s) occupied by each species, and pie charts at nodes indicate the probability of occupying each desert.

Figure S3. Scatterplots showing the relationship between environmental variables and grid-

cell lizard richness, coloured by desert region.

CAPTIONS FOR SUPPLEMENTARY TABLES

Table S1. *Percentage of squamate, lizard, and snake species present in the ancestral*

reconstruction (i.e., with genetic and distribution data) for each desert, relative to the total

number of species in the phylogeny of all squamates (Tonini et al., 2016).

Table S2. *Categorisation of squamate species with genetic (Tonini et al., 2016) and*

distribution data (Roll et al., 2017) into the major arid regions of the world. This was the

dataset used for the ancestral reconstruction of desert occupancy (Figure S2).

1000

1001   ***Table S3***. *Significance values (p-values) of the post-hoc tests showing differences in tip*

1002   *speciation rates (DR metric) between desert regions.*

1003

Table S4. *Average number of biome transitions into and out of each desert region*

reconstructed in the 100 simulations of the evolution of desert occupancy (Figure S2). Rows

show the ancestral state, and columns show the descendant state. For example, the first row

shows the frequency of shifts from non-desert conditions ("Out") to each of the desert

regions. Likewise, the first column shows the frequency of shifts from each desert to non-arid

conditions ("Out").

1010

1011   ***Table S5***. *Results from the correlation tests between environmental variables and grid-cell*

1012   *lizard richness in each desert. For each correlation, 1,000 replicates were implemented, each*

1013   *one with a random sample of 200 grid cells. The table show the mean, the standard deviation*

1014   *(sd), and the 95% confidence interval (low.ci and up.ci) of the correlation coefficients from*

1015   *each analysis.*

1016

1017   ***CAPTIONS FOR SUPPLEMENTARY APPENDIX***

1018   ***Appendix 1****. Scatterplots showing the relationship between environmental variables and*

1019   *grid-cell lizard richness in each desert.*

1020