Stochastic effects on the establishment of mutants resistant to β-lactam antibiotics

Manja Saebelfeld[1,2,*], Suman G. Das[1,*], Arno Hagenbeek[2], Joachim Krug[1],
J. Arjan G.M. de Visser[2]

[1]*University of Cologne, Institute for Biological Physics, Köln, Germany*
[2]*Wageningen University, Laboratory of Genetics, Wageningen, Netherlands*

*These authors contributed equally to the work.

Corresponding authors: SG. Das sdas3@uni-koeln.de, M. Saebelfeld msaebelfeld@gmail.com

Abstract

For antibiotic resistance to arise, new resistant mutants must establish in a bacterial

population before they can spread via natural selection. Understanding the stochastic factors

influencing mutant establishment is crucial for a quantitative understanding of antibiotic

resistance emergence. Here, we quantify the single-cell establishment probability of four

Escherichia coli strains expressing β-lactamase alleles with different activity against the

antibiotic cefotaxime, as a function of concentration. Using a simple branching model, we

show that concentrations well below the minimal inhibitory concentration (MIC) can

substantially hamper establishment, particularly for highly resistant mutants, suggesting

cooperative effects due to antibiotic breakdown at high cell densities. We further show that

variation among cell lineages may explain the more gradual influence of increased antibiotic

concentrations on agar. Finally, we use the single-cell establishment probability to predict a

strain's MIC in the absence of social interactions, as a general reference to detect cooperative

resistance effects.

Introduction

A classical question in population genetics addresses the fate of a new adaptive allele in a population. Any new allele is initially present at a low frequency in the population and thus prone to extinction by genetic drift, even if the allele is beneficial (Patwa & Wahl, 2008). In 1927, Haldane stated that a novel beneficial allele has to overcome drift loss and *establish* in the population before it can be picked up by selection; the probability of which he estimated to be roughly two times its selective benefit (Haldane, 1927). Since then, Haldane's approximation has been generalised and extended by a great number of models, but none of them have been tested empirically until about a decade ago (Patwa & Wahl, 2008). With advances in genomic techniques, the issue of fixation probability has recently been studied in experimental populations of microorganisms (e.g. Lang, Botstein, & Desai, 2011; Frenkel, Good, & Desai, 2014; Good, McDonald, Barrick, Lenski, & Desai, 2017). These studies looked at the fate of beneficial alleles that are already established at a certain frequency in the population, finding that their fixation strongly depends on the competition with further beneficial alleles that appear via mutation over time. Only a small number of empirical studies has investigated the stochastic process of establishment of an initially rare beneficial allele arisen by *de novo* mutation in fungi (Gifford, De Visser, & Wahl, 2013), bacteria (Farrell, Gralka, Hallatschek, & Waclaw, 2017; Giometto, Nelson, & Murray, 2018) and nematodes (Chelo, Nédli, Gordo, & Teotónio, 2013). Overall, these studies found that genetic drift dominates the establishment process, and the probability for a beneficial allele to establish in a population depends on the alleles' initial frequency, its selective benefit, and the size of the population.

Whether a beneficial allele establishes successfully in a population is of particular relevance in cases of unwanted evolution, such as the evolution of antibiotic resistance. The worldwide growing concern about failing treatments of infections due to the emergence of resistant

bacterial pathogens has led to the awareness that this problem has to be tackled from

different angles, one of which is the better understanding of the emergence and evolution of

antibiotic resistance (Baym, Stone, & Kishony, 2016; Smith et al., 2016; Furusawa, Horinouchi,

& Maeda, 2018). Resistance alleles can be acquired via horizontal gene transfer or *de novo*

mutation. In the latter case, the resistant allele is initially present in a single individual within

a susceptible population. But how likely is it that this single genotype can establish in the

population and what factors influence this process? Using time-lapse microscopy, Coates et

al. (2018) followed the growth of single *Escherichia coli* cells on agar, supplemented with

different antibiotics. They found that the establishment of a single cell, i.e. the growth into a

visible macrocolony, was a highly stochastic process driven by fluctuations in cell death and

birth, leading to the extinction of microcolonies at various time points. This stochasticity was

more pronounced with increasing antibiotic concentration, resulting in higher extinction

rates. Similar results were found for streptomycin resistance in *Pseudomonas aeruginosa*

(Alexander & MacLean, 2020). By seeding one to a few cells of a resistant strain in liquid

medium with increasing antibiotic concentrations, it was found that at concentrations as low

as 1/8 of the minimum inhibitory concentration (MIC) of this strain, its establishment

probability was only 5% and that establishment of one cell was independent of the presence

of other cells in the inoculum. The authors speculated that the lack of social effects on mutant

establishment may depend on the mechanism of antibiotic resistance, and could be modified

when positive social interactions play a role, e.g. due to the expression of enzymes that break

down the antibiotic (Alexander & MacLean, 2020).

A common resistance mechanism potentially involving social effects involves β-lactamase

enzymes. β-lactamases hydrolyse the lactam ring of β-lactam antibiotics, which target

penicillin-binding proteins (PBPs) that are involved in cell wall synthesis of Gram-negative

bacteria, thereby leading to loss of cell wall integrity and eventual cell lysis (Samaha-Kfoury &

Araj, 2003; Bush, 2010). A well-studied and clinically relevant example is TEM-1 β-lactamase,

which has high activity toward first-generation penicillins, but low activity toward third-

generation β-lactams, including the cephalosporin cefotaxime (CTX). However, the TEM-1

allele can acquire mutations, rendering it more active to CTX (Salverda, de Visser, & Barlow,

2010; Schenk, Szendro, Krug, & de Visser, 2012; Van Dijk, Hwang, Krug, De Visser, & Zwart,

2017). As β-lactamases are expressed in the periplasmic space of gram-negative pathogens

like *E. coli*, this system has been recognized as potentially cooperative, since the enzymatic

breakdown in the periplasm reduces the antibiotic concentration in the environment (Brown,

West, Diggle, & Griffin, 2009), thereby cross-protecting susceptible strains (Medaney,

Dimitriu, Ellis, & Raymond, 2016). Such a cooperative behaviour can be expected to be more

pronounced on agar due to the local breakdown of the antibiotic, creating microenvironments

that allow for co-existence of strains with different levels of antibiotic resistance (Nicoloff &

Andersson, 2016; Frost et al., 2018; Geyrhofer & Brenner, 2020).

Here, we examined the single-cell establishment probability of antibiotic-degrading bacteria

under a range of antibiotic concentrations in a structured and an unstructured environment.

We used four *E. coli* strains expressing β-lactamase alleles with different activity, and hence

different levels of resistance, toward CTX. Each strain was tested separately at low cell

densities in both liquid and on agar medium with increasing CTX concentrations. To determine

the single-cell establishment probability from the data, we developed a simple branching

model, followed by a more detailed investigation of the cause of differences in stochasticity

between the two environments. We show that environmental structure and cooperative

behaviour play at most a minor role in the establishment of resistant cells at low initial

densities. Based on our results, we further introduce a new measure for resistance level that

takes inoculum size and stochasticity into account, in contrast to the classical MIC assay.

Results and Discussion

Single-cell establishment probability

To determine single cell establishment probabilities, we performed each one experiment in

liquid medium and on agar plates. In each experiment, four *E. coli* strains with increasing levels

of CTX resistance (referred to as Ancestor, Single mutant, Double mutant, Triple mutant; cf.

Table 1) were tested separately under exposure to a range of CTX concentrations. We use the

notation $p_e(x)$ to denote the probability that a single cell establishes a macroscopic

population (i.e. growth in liquid or colony formation on agar visible with the naked eye) at CTX

concentration x. When the antibiotic concentration is zero, a population should establish with

near certainty. Likewise, when the concentration is above the strains' minimum inhibitory

concentration (MIC), establishment is completely prevented. Stochastic outcomes can arise

only at intermediate concentrations.

In the liquid medium experiment, we grew bacteria in multi-well plates starting from a small

number of cells (1-3 on average) in each well and observed their establishment as a function

of CTX concentration. Applying aliquots of the cell cultures used for inoculating the well-plates

on agar, provided estimates of the viable cell numbers needed to evaluate the probabilities.

Using a bootstrapping algorithm and assuming a Poisson distribution of independently acting

cells in each inoculum (details in Supplement S4), we estimated the single-cell establishment

probability $p_e(x)$ for all strains and conditions, which are shown in Figure 1. As expected, the

mutants with higher resistance require higher antibiotic concentration to inhibit

establishment, and all strains show intermediate values of p_e for a range of subMIC CTX

concentrations.

Figure 1. Single-cell establishment probabilities, p_e, in liquid medium, estimated for the four
tested strains as a function of antibiotic concentration. For each strain, two inocula were used,
one of approximately 1 cell (blue circles) and another of 2-3 cells (red squares); average
inoculum size estimated from separate CFU counts are shown. The inferred single-cell
establishment probabilities are nearly identical for the two cases, implying that the
establishment probability of a lineage is not influenced by the presence of other lineages at
these low initial cell densities. The green diamonds are weighted averages over the two
inoculum sizes (see Supplement S3 for details on the method used to estimate p_e).

One complication in estimating the single-cell establishment probability is that multiple cells

may be present initially in some wells, and the lineages arising out of them may in principle

interact, for example through competition for resources or breakdown of antibiotic molecules.

In that case, the establishment probabilities of multiple lineages arising from different cells

are no longer independent, biasing our estimates of $p_e(x)$. To test whether lineages interact

from initially small inocula, we conducted the liquid experiment with two different mean

inoculum sizes and estimated $p_e(x)$ for both cases under the assumption that there is no

interaction (cf. Supplement S4). If interaction occurred, the inferred values should disagree.

However, Figure 1 shows that they agree closely, indicating that interaction between lineages

is insignificant at these low cell densities.

The absence of social interactions in liquid medium was also found in the study with *P.*

aeruginosa and streptomycin under similar conditions regarding culture volume and inoculum

size (Alexander & MacLean, 2020). In that study, social interactions were less expected

because streptomycin resistance does not involve degrading enzymes with the potential for

such interactions. Possibly social interactions play a role during the establishment of new β-

lactamase mutants in structured environments, where antibiotic breakdown may locally affect

establishment probability.

To determine the effect of environmental structure on establishment, we plated about 200

cells of each *E. coli* strain on CTX-containing agar, followed by counts of colony-forming units

(CFUs) after incubation. The establishment probability was estimated as the CFU count

relative to that at zero concentration (Figure 2). The patterns of p_e as a function of CTX

concentration are broadly similar to those in liquid medium. Overall, these results show that

the establishment process of single *E. coli* cells in our system can be described by simple

branching processes, assuming a uniform cell death probability within each lineage (see

Supplement S2), in both liquid and agar environments. However, when comparing the

estimates for liquid and agar medium more closely, one notices that the decline of p_e with

CTX concentration is less steep on agar than in liquid, even though the average establishment

probabilities are not systematically different. In the following section, we analyse the origin

and nature of this difference by modelling the function $p_e(x)$ and fitting it to our data.

Figure 2. Estimated single-cell establishment probabilities for the four strains as a function of

CTX concentration in liquid and on agar. For liquid, the values are obtained as weighted

averages over the two inoculum sizes (as shown in Figure 1). For agar, the inoculum size is

effectively 1, since each colony usually grows from a single cell. Green dots (with dashed lines)

show the per-cell division probability, p_d, (estimated using the branching model without

lineage variability), for two particular strains, one each for liquid and agar. The shape of the

p_d curve is similar for the other cases (not shown).

Sources of stochasticity and the branching model

When the establishment probability is different from 0 or 1, it means that the colony growth

process arising from genetically identical cells in the same environment can have different

outcomes. There can be two different sources of stochasticity in the system. One source we

call *inherent noise*, which arises from random thermal fluctuations in chemical reactions. Two

nascent cells that are identical in structure and composition may still have different

establishment fates due to inherent noise, potentially leading to different outcomes like cell

division for one and cell lysis for another. The other source we call *lineage variability*, which

we define as the systematic variation between founder cells of colonies, that can arise from

phenotypic differences already present between cells in the inoculum or can be caused by

small variations across the micro-environments in which the colonies grow. We aim to

differentiate between the two sources of stochastic effects in our data and understand the

relative contribution of each in the two tested environments with a different structure.

To do so, we model population growth as a branching process, where each cell has a cell

division probability, p_d, and consequently a cell death probability, $1 - p_d$. Inherent noise is

present whenever p_d is different from 0 and 1 and we assume all cells in a lineage to have the

same p_d. In a branching process, the simplest definition of establishment is that the population

continues to be of non-zero size indefinitely. For our branching model,

179
$$p_e = 2 - \frac{1}{p_d} \tag{1}$$

when $p_d \geq \frac{1}{2}$, and $p_e = 0$ when $p_d \leq \frac{1}{2}$ (see Supplement S2 for details). Considering p_d to be

identical for all cell lineages rules out lineage variation, and cell fate is purely a result of

inherent noise. To see if this is sufficient to explain the data, we estimate p_d from the observed

values of p_e, which are shown in Figure 2. The way p_d approaches $\frac{1}{2}$ asymptotically with

increasing concentrations is counter to the expectation that $p_d \rightarrow 0$ at very high concentration.

This indicates either that our definition of establishment is not biologically realistic, or that

inherent noise is inadequate for explaining the intermediate values of p_e. We can modify the

definition of establishment to make it more realistic by requiring that the process reaches a

certain threshold cell number in the time window of the experiment (Schenk et al., 2012). The

results (shown in Supplement S2) indicate that this is not sufficient to explain the data. We,

therefore, need to include lineage variability into our model.

<u>Model for $p_e(x)$</u>

To arrive at a model that incorporates both sources of stochasticity, inherent noise and lineage

variability, we start by modelling p_d as a Hill function of the antibiotic concentration, x, and

the antibiotic concentration where p_d is ½, x_0:

$$p_d(x, x_0) = \frac{1}{1 + \left(\frac{x}{x_0}\right)^n}. \qquad (2)$$

A mechanistic basis for this functional form is as follows: due to inactivation of penicillin-

binding proteins (PBPs, which are the main target of CTX; Rossolini, Arena, & Giani, 2017),

there is a shortage of PBP molecules, leading to the formation of cell wall pores, some of which

can eventually lead to bulging of the cell wall and lysis (Yao, Kahne, & Kishony, 2012). Let the

number of pores be Poisson distributed with mean λ, and let the probability that a pore does

not bulge be q. Then p_d is the probability that there are no bulging pores, which can be shown

to be equal to $e^{-\lambda(1-q)}$. The dependence of λ on the concentration is not known, but the

choice of a simple empirical form for λ produces the function in (2) (see Supplement S5 for

details). We emphasize that the choice of the Hill function is not unique; any inverted

sigmoidal function would suffice for the purpose. Note that in (2), x_0 is the concentration at

which p_d drops to $\frac{1}{2}$, and consequently p_e drops to 0; in other words, x_0 is the single-cell MIC

(Artemova, Gerardin, Dudley, Vega, & Gore, 2015). The value of x_0 is determined in part by the

level of β-lactamase activity, as a higher antibiotic degradation rate would produce a higher

x_0. The quantity $\frac{x}{x_0}$ can be interpreted as the effective (dimensionless) intracellular antibiotic

concentration. The Hill exponent n serves as a measure of the inherent noise. In the limit $n \rightarrow$

∞, $p_d(x)$ behaves like a step function that drops sharply from 1 to 0 at $x = x_0$, implying no

stochasticity since all cells behave identically (they all divide or all die). Thus high values of n

correspond to low stochasticity. For low n, the curve of $p_d(x)$ decays slowly, and therefore has

a larger window in which values of p_e are substantially different from 0 or 1, implying higher

stochasticity.

Lineage variability, on the other hand, is introduced via among-lineage variation in the

parameter x_0. Specifically, we model x_0 as a random variable drawn independently for each

lineage from the gamma distribution:

$$P(x_0) = \frac{1}{\Gamma(k)a^{k-1}} x_0^{k-1} e^{\frac{-x_0}{a}}.$$

The mean of x_0 is ka, and the variance is ka^2. The parameter a provides a scaling factor of

x_0. The coefficient of variation of x_0 is a measure of lineage variability, which in this case is

$k^{-\frac{1}{2}}$; k is therefore inversely related to the degree of lineage variability. The single-cell

establishment probability inferred from a large number of observations is then simply the

establishment probability (1) averaged over the distribution of x_0. The final model for

establishment probability is then:

$$p_e(x) = \int_x^\infty dx_0 \, P(x_0) \left(1 - \left(\frac{x}{x_0} \right)^n \right). \tag{3}$$

The parameter n for intrinsic noise is the same for all mutants, but may, in general, differ

between liquid and agar environments. The parameters for the distribution of x_0 can be

different for the tested strains and can depend on the environment as well. We performed

collective fits using the same value of n for both media, shown in Figure 3. It was estimated

that $n = 7.18 \pm 0.60$ and the coefficient of variation on agar was higher than in liquid for

most strains, consistent with the less steep curve for agar. In Figure 3, the x-values for each

strain have been scaled by the corresponding estimate for mean x_0. The curves for different

strains largely collapse, indicating small (but non-zero) lineage variability in x_0. The

coefficients of variation of x_0, which contain the effect of lineage variability, are larger for the

agar than for the liquid environment in most cases, as expected, but the difference is small.

We also performed model fits with separate estimates of n for liquid and agar. In this case,

the lower mean slope on agar was explained chiefly by a smaller value of n, as shown in the

Supplement (S3: Figure S3). The lineage variability was, in this case, similar for both media

except in the case of the Single mutant. The fit was equally good (see Supplement S6 for

details), making it hard to distinguish between the contribution of both sources of

stochasticity based on curve-fitting alone. However, the assumption of a universal value for n

is more parsimonious, and higher lineage variability on agar compared to liquid may also be

expected from its supposedly greater environmental heterogeneity. It thus seems likely that

a higher lineage variability is the main contributor for the lower slope of $p_e(x)$ on agar, but a

contribution from inherent noise cannot be ruled out.

Figure 3. Effect from intrinsic noise and lineage variability in x_0 on single-cell establishment
probability in liquid compared to agar. A joint fit of equation (3) with the same value for n for
both environments was used. **A**: $p_e(x)$ obtained from curve-fitting. The x-values for each
strain (Anc: TEM-1 Ancestor, Sm: Single mutant, Dm: Double mutant, Tm: Triple mutant) have
been scaled by their respective mean x_0, and the values for agar have been shifted along the
x-axis to the right by 0.5 to improve visibility only. **B**: The coefficient of variation of x_0 (mean
± S.D.), indicating higher lineage variability on agar than in liquid for most strains.

 <u>Minimum inhibitory concentration</u>

One of the main findings in the *P. aeruginosa* study mentioned above (Alexander & MacLean,

2020), was that the used mutant strain, when seeded alone at a few cells in a similar setup as

ours, showed a <5% establishment probability at a streptomycin concentration of 1/8 of the

strain's MIC. At the same MIC fraction, the establishment probabilities for our strains are

about 100, 100, 8 and 0% for the Ancestor, Single mutant, Double mutant and Triple mutant,

respectively. These results show that there is no general relationship between the MIC and

the single-cell establishment probability, neither across systems nor among genotypes of the

same species within the same system. The cause for this is likely rooted in how the MIC is

measured. In a standard MIC assay, 10^5 cells are inoculated into 200 μl medium in a series of

wells with 2-fold differences in antibiotic concentration. Thus, the "real" MIC value lies

between the smallest concentration where no growth is observed (defined as MIC) and half

this concentration. The concentration range of where the real MIC lies, is then smaller or

larger depending on the MIC of the respective strain. Another issue with the MIC is that even

the growth of an inoculum as large as 10^5 cells is a stochastic process under sufficiently high

antibiotic concentrations (Coates et al., 2018), which generally results in a reduced

reproducibility of MIC assays.

A conceptually cleaner approach to determine the MIC would be to recognise this

stochasticity. Here, we introduce a new measure, $MIC_q(N)$, defined as the concentration at

which an inoculum of size N has a probability q of not becoming established. In the absence

of interactions between lineages, $MIC_q(N)$ can be predicted from $p_e(x)$ (see Supplement S7

for details). Setting $q = \frac{1}{2}$ and $N = 10^5$, we obtain $MIC_q(N)$ values of 0.038, 0.23, 1.6, and

2.9 μg/ml CTX for the Ancestor, Single mutant, Double mutant and Triple mutant, respectively.

The values are smaller than the previously determined MICs (Table 1), and the differences

become increasingly pronounced with the level of resistance of the mutants. This suggests that positive interactions among cell lineages enhance their collective establishment due to the enzymatic breakdown of the antibiotic in the medium, which is particularly pronounced at high cell densities and for cells expressing enzymes with high activity against CTX.

The effect of inoculum size on the MIC through social interactions has been recognised before (Brook, 1989; Tan et al., 2012). Causes for such social effects have been suggested to include binding of the antibiotic to cellular components (Queenan, Foleno, Gownley, Wira, & Bush, 2004) or the expression of proteins that inactivate the antibiotic (Lenhard & Bulman, 2019). Our new measure includes a purely statistical inoculum effect, which leads to a logarithmic increase of $MIC_q(N)$ for large N (see Supplement S7 for details). Our results on the discrepancy between the MIC and the single-cell establishment probability are in line with the findings of other studies that criticised the MIC assay as not suitable for dosage determination for the treatment of antibiotic-resistant infections (Gould & MacKenzie, 1997; Artemova et al., 2015).

Conclusion

Our study contributes to the understanding of the establishment of adaptive mutants, a topic which has been under much theoretical scrutiny but is rarely investigated empirically. We show that establishment of populations starting from a single cell in the presence of an antibiotic can be well described by a simple branching model both in liquid and agar environments, where the cell division probability is dependent on the antibiotic concentration. We find that the establishment of single mutant cells is negatively affected by CTX concentrations well below each tested strains' MIC, consistent with previous findings(Alexander & MacLean, 2020).

One aim of our study was to test if cooperative behaviour (i.e. collective antibiotic breakdown)

among cells of the same strain would affect establishment probability and whether such an

effect differed between unstructured and structured environments. Our data show a

reasonable agreement between establishment in liquid and agar media, indicating that

environmental structure plays only a modest role, and that cooperative behaviour has a

negligible effect on the establishment probability of single resistant mutant cells. However,

the $p_e(x)$ curves for agar are slightly shallower than for the liquid environment, indicating

higher stochasticity. By evaluating the effect of two potential sources of stochasticity, random

intrinsic noise and systematic (e.g. heritable) variability among cell lineages, we found that

the simplest explanation for this observation is an enhanced lineage variability on agar,

possibly arising from small differences in local agar conditions.

Lastly, we show that social interaction effects on the establishment of antibiotic-resistant

mutants can be strong at high cell densities, making conventional MIC values highly sensitive

to the assay conditions. We suggest a more robust measure of resistance which includes both

inoculum size and probabilistic effects, $MIC_q(N)$, and can be compared easily with p_e to test

for the presence of interactions.

Material and Methods

<u>Strains and culture conditions</u>

Four strains were used for the experiments, derived from the *Escherichia coli* strain MG1655

DA28100. This strain had previously been modified with a YFP (yellow fluorescent protein)

marker cassette, containing a resistance gene for chloramphenicol (Gullberg, Albrecht,

Karlsson, Sandegren, & Andersson, 2014). For the current study, the chloramphenicol

resistance was removed, and each one of four TEM variants was inserted into the *galK* locus:

the ancestral allele TEM-1 (referred to as "Ancestor" strain), and three mutant alleles with

either 1, 2 or 3 point mutations in the TEM allele (referred to as "Single mutant", "Double

mutant" and "Triple mutant", respectively). While the TEM-1 ancestral allele confers only low

activity towards the β-lactam antibiotic cefotaxime (CTX), the three mutants show increasing

CTX resistance with each additional substitution (Table 1). All TEM loci are under the control

of the *lacI* repressor and are expressed by adding 50 μM Isopropyl β-D-1-

thiogalactopyranoside (IPTG) to the growth medium. The minimum inhibitory concentration

(MIC) of each strain for CTX had previously been determined in duplicates (Table 1), using 2-

fold increases in CTX concentration in microtiter plates filled with 200 μl M9 minimal medium

(containing 0.4% glucose, 0.2% casaminoacids, 2 μg/ml uracil and 1 μg/ml thiamine) and 50

μM IPTG, inoculated with 10^5 cells and incubated for 24h at 37°C.

For culturing, all strains were first recovered from glycerol stocks by streaking them out on

each one LB (Luria-Bertani) agar plate and incubated overnight at 37°C. From those plates,

each one colony was picked into 1 ml M9 medium (as above) and incubated overnight at 37°C,

250 rpm. The cultures were then serially diluted with phosphate-buffered saline (PBS) to the

density needed for the particular experiment (see below), assuming an initial density of

2.5×10^9 cells/ml.

Table 1: Overview and minimum inhibitory concentration (MIC) of the used strains.

Wildtype strain (without TEM allele)	TEM amino acid substitution	MIC for Cefotaxime (μg/ml)		Strain name (as used here)
		MIC*	$MIC_q(N)$**	
MG1655 DA28100 galK::YFPΔCAT-	TEM-1	0.08	0.038	Ancestor
	G238S	0.64	0.23	Single mutant
	E104K/G238S	10.24	1.6	Double mutant
	E104K/M182T/G238S	81.92	2.9	Triple mutant

*As determined prior to the study by standard MIC assays (see Material and Methods).

**As predicted from $p_e(x)$ in the current study (see the last section of Results and Discussion, and
Supplement S7).

<u>Liquid experiment</u>

Each of the four strains was tested separately in the absence of CTX and six CTX concentrations

(Supplement S1: Table S1), where the particular tested concentrations depended on the

strains' resistance level. The range of concentrations was chosen to display establishment

probabilities between 0 and 1, based on pilot experiments to find the right conditions (data

not shown).

Per strain and CTX concentration, on average either about 1 or 2 cells were seeded into 285

wells across three 96-well microtiter plates. For this, 190 µl M9 medium with the respective

cefotaxime concentration and 50 µM IPTG were pipetted into all wells. Overnight cultures of

the strains were diluted to 200 and 100 cells/ml with PBS. 10 µl of the dilutions were pipetted

into the wells. One well per plate served a medium control, mock-inoculated with 10 µl PBS.

The plates were incubated about 40 h at 37°C (static). After incubation, OD at 600 nm of all

wells was measured in a plate reader (Victor3™, PerkinElmer) without the plate lid. All medium

controls across the experiment showed no sign of growth; the OD ranged from 0.033 to 0.038.

Thus, a threshold of >0.05 was applied to determine whether a well successfully established

(i.e. showed growth).

To test for correct culture dilutions and Poisson distribution of seeded cell numbers, the same

culture dilutions used for seeding the well plates were dropped on LB agar plates (120 x 120

mm). Per dilution and strain, a total of 3x32 10 µl drops were applied to the plates, let dry for

about 10 to 20 minutes, incubated at 37°C until leaving, and then moved to 30°C overnight to

prevent overgrowth, followed by counting the number of colonies per droplet.

<u>Agar experiment</u>

As in the liquid experiment, all strains were tested separately and the CTX concentration range

depended on the strain. Five to eight concentrations were tested per strain (Supplement S1:

Table S1). The overnight cultures were diluted to 4000 cells/ml. 50 µl (~200 cells) of the

dilutions were spread onto 92 mm agar plates containing M9 medium (as above) with 1.5 %

agar, 50 µM IPTG and the respective cefotaxime concentration, using a bacterial spreader. Per

strain and cefotaxime treatment, eight replicate plates were used. The plates were incubated

at 37°C until colonies were big enough to count them unambiguously (20 to 48 hours). Each

colony is regarded as a successfully established single cell.

Acknowledgements

We would like to thank Helen Alexander for fruitful discussions of her work and advice on the

liquid experiment setup; and Tobias Bollenbach for advice on the p_e model.

Author contribution

All authors conceived the study. MS and AH conducted the experiments. SGD developed the

models with help from JK. MS and SGD wrote the manuscript with help from JK and JAGMdV.

All authors read the manuscript and approved the final version.

Competing Interests Statement

All authors declare that they have no significant competing financial, professional, or personal

interests that might have influenced the performance or presentation of the work described

in this manuscript.

References

Alexander, H. K., & MacLean, R. C. (2020). Stochastic bacterial population dynamics restrict the establishment of antibiotic resistance from single cells. *Proceedings of the National Academy of Sciences of the United States of America*, *117*(32), 19455–19464. https://doi.org/10.1073/pnas.1919672117

Artemova, T., Gerardin, Y., Dudley, C., Vega, N. M., & Gore, J. (2015). Isolated cell behavior drives the evolution of antibiotic resistance. *Molecular Systems Biology*, *11*(7), 822. https://doi.org/10.15252/msb.20145888

Baym, M., Stone, L. K., & Kishony, R. (2016). Multidrug evolutionary strategies to reverse antibiotic resistance. *Science*, *351*(6268). https://doi.org/10.1126/science.aad3292

Brook, I. (1989). Inoculum Effect. *Reviews of Infectious Diseases*, *11*(3), 361–368. https://doi.org/10.1093/clinids/11.3.361

Brown, S. P., West, S. A., Diggle, S. P., & Griffin, A. S. (2009). Social evolution in micro-organisms and a Trojan horse approach to medical intervention strategies. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *364*(1533), 3157–3168. https://doi.org/10.1098/rstb.2009.0055

Bush, K. (2010). Bench-to-bedside review: the role of β-lactamases in antibiotic-resistant Gram-negative infections. *Critical Care*, *14*(3), 224.

Chelo, I. M., Nédli, J., Gordo, I., & Teotónio, H. (2013). An experimental test on the probability of extinction of new genetic variants. *Nature Communications*, *4*, 1–8. https://doi.org/10.1038/ncomms3417

Coates, J., Park, B. R., Le, D., Şimşek, E., Chaudhry, W., & Kim, M. (2018). Antibiotic-induced population fluctuations and stochastic clearance of bacteria. *ELife*, *7*, 1–26. https://doi.org/10.7554/eLife.32976

Farrell, F. D., Gralka, M., Hallatschek, O., & Waclaw, B. (2017). Mechanical interactions in

bacterial colonies and the surfing probability of beneficial mutations. *Journal of the*

Royal Society Interface, *14*(131). https://doi.org/10.1098/rsif.2017.0073

Frenkel, E. M., Good, B. H., & Desai, M. M. (2014). The fates of mutant lineages and the

distribution of fitness effects of beneficial mutations in laboratory budding yeast

populations. *Genetics*, *196*(4), 1217–1226.

https://doi.org/10.1534/genetics.113.160069

Frost, I., Smith, W. P. J., Mitri, S., San Millan, A., Davit, Y., Osborne, J. M., … Foster, K. R.

(2018). Cooperation, competition and antibiotic resistance in bacterial colonies. *ISME*

Journal, *12*(6), 1582–1593. https://doi.org/10.1038/s41396-018-0090-4

Furusawa, C., Horinouchi, T., & Maeda, T. (2018). Toward prediction and control of

antibiotic-resistance evolution. *Current Opinion in Biotechnology*, *54*, 45–49.

https://doi.org/10.1016/j.copbio.2018.01.026

Geyrhofer, L., & Brenner, N. (2020). Coexistence and cooperation in structured habitats.

BMC Ecology, *20*(1), 1–15. https://doi.org/10.1186/s12898-020-00281-y

Gifford, D. R., De Visser, J. A. G. M., & Wahl, L. M. (2013). Model and test in a fungus of the

probability that beneficial mutations survive drift. *Biology Letters*, *9*(1).

https://doi.org/10.1098/rsbl.2012.0310

Giometto, A., Nelson, D. R., & Murray, A. W. (2018). Physical interactions reduce the power

of natural selection in growing yeast colonies. *Proceedings of the National Academy of*

Sciences of the United States of America, *115*(45), 11448–11453.

https://doi.org/10.1073/pnas.1809587115

Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E., & Desai, M. M. (2017). The dynamics

of molecular evolution over 60,000 generations. *Nature*, *551*(7678), 45–50.

https://doi.org/10.1038/nature24287

Gould, I. M., & MacKenzie, F. M. (1997). The response of *Enterobacteriaceae* to beta-lactam

antibiotics-'round forms, filaments and the root of all evil'. *Journal of Antimicrobial*

Chemotherapy, *40*(4), 495–499.

Gullberg, E., Albrecht, L. M., Karlsson, C., Sandegren, L., & Andersson, D. I. (2014). Selection

of a multidrug resistance plasmid by sublethal levels of antibiotics and heavy metals.

MBio, *5*(5), 19–23. https://doi.org/10.1128/mBio.01918-14.Editor

Haldane, J. B. S. (1927). A mathematical theory of natural and artificial selection, Part V:

selection and mutation. *Mathematical Proceedings of the Cambridge Philosophical*

Society, *23*(7), 838–844. https://doi.org/10.1017/S0305004100015644

Lang, G. I., Botstein, D., & Desai, M. M. (2011). Genetic variation and the fate of beneficial

mutations in asexual populations. *Genetics*, *188*(3), 647–661.

https://doi.org/10.1534/genetics.111.128942

Lenhard, J. R., & Bulman, Z. P. (2019). Inoculum effect of β-lactam antibiotics. *The Journal of*

Antimicrobial Chemotherapy, *74*(10), 2825–2843. https://doi.org/10.1093/jac/dkz226

Medaney, F., Dimitriu, T., Ellis, R. J., & Raymond, B. (2016). Live to cheat another day:

bacterial dormancy facilitates the social exploitation of β-lactamases. *ISME Journal*,

10(3), 778–787. https://doi.org/10.1038/ismej.2015.154

Nicoloff, H., & Andersson, D. I. (2016). Indirect resistance to several classes of antibiotics in

cocultures with resistant bacteria expressing antibiotic-modifying or -degrading

enzymes. *Journal of Antimicrobial Chemotherapy*, *71*(1), 100–110.

https://doi.org/10.1093/jac/dkv312

Patwa, Z., & Wahl, L. M. (2008). The fixation probability of beneficial mutations. *Journal of*

the Royal Society Interface, *5*(28), 1279–1289. https://doi.org/10.1098/rsif.2008.0248

Queenan, A. M., Foleno, B., Gownley, C., Wira, E., & Bush, K. (2004). Effects of inoculum and

β-lactamase activity in ampC- and extended-spectrum β-lactamase (ESBL)-producing

Escherichia coli and *Klebsiella pneumoniae* clinical isolates tested by using NCCLS ESBL

Methodology. *Journal of Clinical Microbiology*, *42*(1), 269–275.

https://doi.org/10.1128/JCM.42.1.269

Rossolini, G. M., Arena, F., & Giani, T. (2017). *Mechanisms of antibacterial resistance*.

Infectious Diseases (Fourth Edi). Elsevier Ltd. https://doi.org/10.1016/b978-0-7020-

6285-8.00138-6

Salverda, M. L. M., de Visser, J. A. G. M., & Barlow, M. (2010). Natural evolution of TEM-1 β-

lactamase: experimental reconstruction and clinical relevance. *FEMS Microbiology*

Reviews, *34*(6), 1015–1036. https://doi.org/10.1111/j.1574-6976.2010.00222.x

Samaha-Kfoury, J. N., & Araj, G. F. (2003). Recent developments in β lactamases and

extended spectrum β lactamases. *Bmj*, *327*(22 November 2003), 1209–1213.

https://doi.org/10.1136/bmj.327.7425.1209

Schenk, M. F., Szendro, I. G., Krug, J., & de Visser, J. A. G. M. (2012). Quantifying the adaptive

potential of an antibiotic resistance enzyme. *PLoS Genetics*, *8*(6), e1002783.

https://doi.org/10.1371/journal.pgen.1002783

Smith, E., Lichten, C. A., Taylor, J., MacLure, C., Lepetit, L., Harte, E., … Littmann, J. (2016).

Evaluation of the EC action plan against the rising threats from antimicrobial resistance.

Tan, C., Phillip Smith, R., Srimani, J. K., Riccione, K. A., Prasada, S., Kuehn, M., & You, L.

(2012). The inoculum effect and band-pass bacterial response to periodic antibiotic

treatment. *Molecular Systems Biology*, *8*(617), 1–11.

https://doi.org/10.1038/msb.2012.49

Van Dijk, T., Hwang, S., Krug, J., De Visser, J. A. G. M., & Zwart, M. P. (2017). Mutation supply

and the repeatability of selection for antibiotic resistance. *Physical Biology*, *14*(5).

https://doi.org/10.1088/1478-3975/aa7f36

Yao, Z., Kahne, D., & Kishony, R. (2012). Distinct single-cell morphological dynamics under

beta-lactam antibiotics. *Molecular Cell*, *48*(5), 705–712.

https://doi.org/10.1016/j.molcel.2012.09.016