DROP: Molecular voucher database for identification of *Drosophila* parasitoids

Resource article

Word count: 7850 excluding references

Authors

Chia-Hua Lue (C-HL), (0000-0002-5245-603X), chiachia926@gmail.com, corresponding

author

Matthew L. Buffington (MLB)

Sonja Scheffer (SS)

Matthew Lewis (ML)

Tyler A. Elliott (TAE)

Amelia R. I. Lindsey (AL)

Amy Driskell (AD), (0000-0001-8401-7923)

Anna Jandova (AJ)

Masahito T. Kimura (MTK)

Yves Carton (YC)

Robert R. Kula (RRK)

Todd A. Schlenke (TAS)

Mariana Mateos (MM), (0000-0001-5738-0145)

Shubha Govind (SG), (0000-0002-6436-639X)

Julien Varaldi (JV)

Emilio Guerrieri (EG), (0000-0002-0583-4667)

Massimo Giorgini (MG), (0000-0001-8670-0945)

Xingeng Wang (XW)

Kim Hoelmer (KH)

Kent M. Daane (KMD**)**

Paul K. Abram (PKA)

Nicholas A. Pardikes (NAP), (0000-0002-9175-4494)

Joel J. Brown (JJB), (0000-0002-3608-6745)

Melanie Thierry (MT)

Marylène Poirié (MP)

Paul Goldstein (PG), (0000-0002-1443-7030)

Scott E. Miller (SEM), (0000-0002-4138-1378)

W. Daniel Tracey (WDT)

Jeremy S. Davis (JSD), (0000-0002-5214-161X)

Francis M. Jiggins (FMJ)

Bregje Wertheim (BW)

Owen T. Lewis (OTL)

Jeff Leips (JL)

Phillip P. A. Staniczenko (PPAS)

Jan Hrcek (JH), (0000-0003-0711-6447), janhrcek@gmail.com

Affiliations

(C-HL, AJ, NAP, JJB, MT, JH) Biology Centre of the Czech Academy of Sciences, Institute

of Entomology, Branisovska 31, 37005 Ceske Budejovice, Czech Republic

(C-HL, PPAS) Department of Biology, Brooklyn College, City University of New York

(CUNY), 2900 Bedford Ave, Brooklyn, NY11210, USA

(MLB, SS, ML, RRK, PG) Systematic Entomology Laboratory, ARS/USDA c/o Smithsonian

Institution, National Museum of Natural History, 10th& Constitution Ave, NW,

Washington DC 20560, USA

(TAE) Centre for Biodiversity Genomics, 50 Stone Road East, University of Guelph,

Guelph, Ontario, N1G2W1, Canada

(AL) Department of Entomology, University of Minnesota, Saint Paul MN 55108.

(AD) Laboratories of Analytical Biology, Smithsonian Institution, National Museum of

Natural History, 10th & Constitution Ave, NW, Washington DC 20560, USA

(MTK) Hokkaido University Museum, Sapporo, Hokkaido 060-0810, Japan

(YC) "Évolution, Génomes, Comportement, Écologie", CNRS et Université Paris-Saclay

(TAS) Department of Entomology at the University of Arizona, Forbes 410, PO BOX

210036, Tucson, AZ 85721-0036.

(MM) Wildlife and Fisheries Sciences Department, Texas A&M University

(SG) The Graduate Center of the City University of New York, 365 Fifth Avenue, New

York, NY10016, USA

(JV) Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie

Evolutive UMR 5558, F-69622 Villeurbanne, France

(EG, MG) CNR- Institute for Sustainable Plant Protection (CNR-IPSP), National Research

Council of Italy, Piazzale E. Fermi 1, 80055 Portici, Italy.

(XW, KH) United States Department of Agriculture, Agricultural Research Services,

Beneficial Insects Introduction Research Unit. 501 S. Chapel St., Newark, DE 19713, USA

(KMD) Department of Environmental Science, Policy and Management. University of

California, Berkeley. Mulford Hall, 130 Hilgard Way, Berkeley, CA 94720.

(PKA) Agriculture and Agri-Food Canada, Agassiz Research and Development Centre,

6947 Hwy #7, Agassiz, V0M 1A0, British Columbia, Canada

(JJB, MT, JH) University of South Bohemia, Faculty of Science, Branisovska 31, 37005,

Czech Republic

(MP) Université "Côte d'Azur", INRAE, CNRS. and Evolution and Specificity of

Multitrophic Interactions (ESIM) Sophia Agrobiotech Institute, 400 Route des Chappes,

BP 167, 06903 Sophia Antipolis, France

(SEM) Smithsonian Institution, National Museum of Natural History, 10th & Constitution

Ave, NW, Washington DC 20560, USA

(WDT, JSD) Department of Biology, Indiana University Bloomington. 702 N. Walnut

Grove, Bloomington, IN47405

(WDT) Gill Center for Biomolecular Science, Indiana University Bloomington. 702 N.

Walnut Grove, Bloomington, IN47405

(JSD) Biology Department, University of Kentucky, 101 T. H. Morgan Building, Lexington,

KY, 40506

(FMJ) Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2

3EH UK

(BW) Groningen Institute for Evolutionary Life Sciences, University of Groningen,

Nijenborgh 9, 9747 AG Groningen, the Netherlands

(OTL) Department of Zoology, University of Oxford. 11a Mansfield Road, Oxford OX1

3SZ, UK.

(JL) Department of Biological Sciences, University of Maryland Baltimore County. 1000

Hilltop circle, Baltimore, MD, 21250

Abstract

Molecular identification is increasingly used to speed up biodiversity surveys and

laboratory experiments. However, many groups of organisms cannot be reliably

identified using standard databases such as GenBank or BOLD due to lack of sequenced

voucher specimens identified by experts. Sometimes a large number of sequences are

available, but with too many errors to allow identification. Here we address this

problem for parasitoids of *Drosophila* by introducing a curated open-access molecular

reference database, DROP (*Drosophila* parasitoids). Identifying *Drosophila* parasitoids is

challenging and poses a major impediment to realize the full potential of this model

system in studies ranging from molecular mechanisms to food webs, and in biological

control of *Drosophila suzukii*. In DROP (http://doi.org/10.5281/zenodo.4519656),

genetic data are linked to voucher specimens and, where possible, the voucher

specimens are identified by taxonomists and vetted through direct comparison with

primary type material. To initiate DROP, we curated 154 laboratory strains, 853

vouchers, 545 DNA sequences, 16 genomes, 11 transcriptomes, and 6 proteomes drawn

from a total of 183 operational taxonomic units (OTUs): 113 described *Drosophila*

parasitoid species and 70 provisional species. We found species richness of *Drosophila*

parasitoids to be acutely underestimated and provide an updated taxonomic catalogue

for the community. DROP offers accurate molecular identification and improves cross-

referencing between individual studies that we hope will catalyze research on this

diverse and fascinating model system. Our effort should also serve as an example for

researchers facing similar molecular identification problems in other groups of

organisms.

Key Words

Biodiversity, DNA sequences, Genomes, Integrative taxonomy, Molecular diagnostics,

Biological control

Introduction

Building a knowledge base that encompasses ecology, evolution, genetics, and biological control is contingent on reliable taxonomic identifications. Molecular identification is commonly used in groups of organisms with cryptic species that are difficult to identify morphologically (Fagan-Jeffries et al., 2018; Miller et al., 2016; Novotny & Miller, 2014), for the molecular detection of species interactions (Baker et al., 2016; Condon et al., 2014; Gariepy et al., 2019; Hrček & Godfray, 2015; Hrcek et al., 2011), and for identification of species from environmental DNA samples (Shokralla et al., 2012). The accuracy of molecular identification, however, depends on the accuracy of identifications associated with sequences databased in existing online depositories. The foundations of that accuracy are the voucher specimens which were sequenced and the collaboration of a taxonomic authority in the deposition of the sequence data.

GenBank serves as the most widely used sequence depository; however, deposition of sequences in GenBank, which is required by most peer-reviewed journals, does not require deposition of associated vouchers. The Barcode of Life Data System database (BOLD) (Ratnasingham & Hebert, 2007) explicitly aims to provide a framework for identifying specimens using single-locus DNA sequences (Hebert et al., 2003; Smith et al., 2005), and while these are associated with vouchers and metadata, the curation of these data is not consistently maintained by those submitting material. A recent study by Pentinsaari et al. (2020) showed misidentification in both databases caused by missteps in the protocols from query sequences to final determination.

Although the BOLD database function "BOLD-IDS" allows considerable database

curation (e.g., sequences are used for identification and/or flagging of

misidentified/contaminated records), it also automatically includes sequences from

GenBank, and may perpetuate the shortcomings previously mentioned since these

cannot be curated from within BOLD. As such, the quality of sequences and the

reliability of identifications obtained from BOLD-IDS can vary, and depends on the

curation by systematists focusing on individual taxa (Meiklejohn et al., 2019). BOLD-IDS

works well for taxa where qualified taxonomists have been involved with assuring data

quality; some insect examples include beetles (Hendrich et al., 2015), butterflies

(Escalante et al., 2010), geometrid moths (Hausmann et al., 2011, 2016; Miller et al.,

2016), true bugs (Raupach et al., 2014), and microgastrine wasps (Smith et al., 2013).

Unfortunately, this is not the case of parasitoids (Insecta: Hymenoptera) of

Drosophila flies (Insecta: Drosophilidae). There are vast numbers of *Drosophila*

parasitoid sequences readily available in GenBank and BOLD, as these parasitoids and

their hosts are important model organisms in biology. As of this writing, there are

88,666 nucleotide sequences deposited in GenBank for *Leptopilina heterotoma*

(Thomson) and *L. boulardi* (Barbotin, Carton & Kelner-Pillault) (Hymenoptera: Figitidae)

alone. However, less than 1 % of the identifications associated with these sequences

have been confirmed by taxonomists or are associated with voucher specimens

deposited in museum collections. With sequencing shifting from individual genes to

genomes we risk that the identification problems will soon apply to whole genomes.

There are around 4000 described species of Drosophilidae, and *Drosophila* contains

more than a third of the family's described species (O'Grady & DeSalle, 2018). By

contrast, although parasitic wasps are generally a species-rich group (Dolphin & Quicke,

2001; Quicke, 2015), the most recent catalogue of parasitoid species that attack

Drosophila lists only 50 described species (Carton et al., 1986). This disparity suggests

that the diversity of parasitic wasps attacking *Drosophila* is severely underestimated, an

assertion supported by the results presented here. This is largely a consequence of the

challenging nature of parasitoid taxonomy, in which morphological identification is

intractable for many species, and the fact that taxonomic specialists are greatly

outnumbered by the species they study.

Currently, only a few biological study systems have been characterized in

sufficient breadth and depth to allow researchers to connect various levels of biological

organization, from molecular mechanisms to food webs of interacting species.

Parasitoids of *Drosophila* represent one such system (Prévost, 2009). Moreover, the

practical feasibility of rearing parasitoids of *Drosophila* parasitoids under laboratory

conditions has led to a number of fundamental discoveries in ecology (Carton et al.,

1991; Terry et al. 2020; Thierry et al.,2021), evolution (Kraaijeveld & Godfray, 1997),

immunology (Kim-Jo et al., 2019; Nappi & Carton, 2001; Schlenke et al., 2007),

physiology (Melk & Govind, 1999), symbiosis (Xie et al., 2011, 2015), behavioral science

(Lefèvre et al., 2012) and other fields. In contrast to this large body of laboratory

studies, basic natural history of *Drosophila* parasitoids, especially their species richness

(Kimura & Mitsui, 2020; Lue et al., 2018), is little known. Addressing this knowledge gap

is especially pressing given current efforts to use parasitoids in biological control efforts,

such as those of the invasive pest spotted wing *Drosophila*, *Drosophila suzukii* (Abram et

al., 2020; Daane et al., 2016; Giorgini et al., 2019; Wang et al., 2020 a&b).

Properly executed molecular identification has the potential to be much more

efficient for the majority of researchers, and many laboratory strains are commonly

identified using DNA sequences alone. While it is practical for researchers to assign

species names based on a match to sequence records in genetic databases, this practice

often causes a cascade of inaccuracies. To illustrate the extent of the problem, we note

the example of *Ganaspis,* a genus of parasitoids commonly used in laboratories that

includes both superficially indistinguishable species with highly divergent sequences

that are often treated as conspecific, as well as specimens with identical sequences

identified under different names (Figure 1).

To address these issues, we introduce a newly curated molecular reference database

for <u>*Dro*</u>*sophila* <u>p</u>arasitoids —DROP— in which sequences are either linked to voucher

specimens identified by taxonomists or have a traceable provenance (Figure 2). The first

aim of DROP is to provide a reliable DNA sequence library for molecular identification of

Drosophila parasitoids that enables cross-referencing of original taxonomic concepts

with those of subsequent studies. We pay special attention to live parasitoid strains

which are available for future experiments. The second aim is to standardize and

expedite the linkage between specimens and available sequence data; we place a

premium on museum vouchers as they allow for repeatable scientific research. In DROP,

this goal is facilitated through a consolidated digital infrastructure of data associated

with laboratory strains, offering the opportunity for researchers to re-examine past

experimental results in a permanent context. The third aim is to provide an up-to-date

catalogue of the diversity of *Drosophila* parasitoids as a foundation for advancing the

understanding of their taxonomy. Finally, the fourth aim of DROP is for our collaborative

effort to serve as an inspiration to communities of researchers studying other groups of

organisms who are experiencing difficulties with the reliability of molecular reference

databases.

Materials and Methods

Drosophila species and their parasitoids

The phylogenetic and subgeneric structure within *Drosophila* and related genera is

not yet fully resolved (O'Grady & DeSalle, 2018). Various subgenera, including

Scaptomyza, *Zaprionus*, *Lordiphosa* and *Samoaia*, have been treated as both genera and

subgenera, and researchers have yet to achieve consensus on these various hypotheses

(O'Grady & DeSalle, 2018; Remsen & O'Grady, 2002; Yassin, 2013; Yassin & David,

2010). Species in *Drosophila* subgenera and genera closely related to *Drosophila*

commonly share niche space and natural histories and, as a result, are often attacked by

overlapping or identical groups of parasitoids. For instance, the invasive African fig fly,

Zaprionus indianus Gupta is attacked by *Pachycrepoideus vindemiae* (Rondani, 1875)

and *Leptopilina boulardi* (Pfeiffer et al., 2019; Santos et al., 2016), all of which have been

recorded from *Drosophila*. Therefore, we also include these groups within the contents

of DROP.

240   Parasitoids of *Drosophila* belong to four superfamilies of Hymenoptera

241 (Chalcidoidea, Cynipoidea, Ichneumonoidea, Diaprioidea) which evolved parasitism of

242 *Drosophila* flies independently (Carton et al., 1986; Prévost, 2009). All the parasitoids

243 known to attack *Drosophila* are solitary and attack either the larval or pupal stage; in

244 both cases, they emerge from the fly's puparium. The known *Drosophila* larval

245 parasitoids belong to two families, Braconidae (including the genera *Asobara,*

246 *Aphaereta, Phaenocarpa, Tanycarpa, Aspilota, Opius*) and Figitidae (*Leptopilina,*

247 *Ganaspis, Leptolamina, Kleidotoma*); all are koinobionts that allow the host to continue

248 development while the parasitoid grows within it. The known *Drosophila* pupal

249 parasitoids belong to three other families, Diapriidae (*Trichopria, Spilomicrus*),

250 Pteromalidae (*Pachycrepoideus, Spalangia, Trichomalopsis, Toxomorpha*) and Encytidae

251 (*Tachinaephagus*); they are all idiobionts that terminate host development immediately.

252 Host-specificity across the *Drosophila* parasitoids is poorly characterized—while some

253 can parasitize other families of Diptera (e.g., *Aphaereta aotea*) (Hughes & Woolcock,

254 1976), most are thought to be limited to *Drosophila* hosts.

256 ***Data sources***

257  To assemble the DROP database, we targeted 20 genera that potentially parasitize

258 frugivorous *Drosophila* species. We compiled DNA sequence and voucher data from four

259 sources: 1) museum collections, 2) publications, 3) molecular biodiversity inventories

260 publicly available in BOLD and GenBank, for which we managed to secure inspection of

the vouchers by taxonomists, and 4) conducted a sequencing and taxonomic inventory

of laboratory strains.

We first gathered species information into a catalogue of *Drosophila* parasitoid

species (Table 1) from 212 references (see DROP database reference table) and 36

institutes (Table S2). To ensure reliable names for nominal species (sequences identified

by a species name) in our database, we confirmed their taxonomic validity using the

Ichneumonoidea 2015 digital catalogue (Yu et al., 2016;

https://web.archive.org/web/20161022093945/http:/ichneumonoidea.name/global.ph

p) and Hymenoptera Online (HOL; http://hol.osu.edu/), both of which are curated by

taxonomic experts. To obtain reliable molecular identification data, we harvested 8,298

DNA sequences from GenBank and BOLD (all compiled into BOLD system: DS-DROPAR

dataset). These sequences represent 443 Barcode Index Numbers (BINs – a form of

provisionary taxa in BOLD) and 520 taxa, for a total of 963 operational taxonomic units

(OTUs). We use the term "OTU" as a general and neutral designation encompassing

described species, provisional species, undescribed species, cryptic species, and mis-

identified species.

The majority of the harvested sequences were Braconidae (6690), Diapriidae

(967), Figitidae (622), and Pteromalidae (19). Because of the concerns with generic

databases (noted above and in Figure 1 and Table S1), we assembled a list of sequences

with valid species names that could either be traceably linked to vouchers examined by

taxonomists or referred to directly in publications authored by a recognized expert in

the relevant taxon group. We then cross-checked species names with their

corresponding BINs in BOLD and flagged potential conflicts between species names and

BINs (Table S1).

A core goal of DROP besides that of a tool for biodiversity research is to function as a

platform that accommodates *Drosophila* parasitoids kept in culture (for experimental

work) or in quarantine (for biological control applications). So far, there has been a lack

of a coherent and reliable means of verifying species kept in laboratory settings, which

can be a serious problem. Since lab cultures are routinely contaminated by neighboring

cultures (e.g., through escapees), one species may be displaced by another even under a

vigilant eye.

For lab and quarantine lines in DROP, we deposited DNA extractions and vouchers in

the National Insect Collection, National Museum of Natural History, Smithsonian

Institution (USNM; Washington, DC, USA). During their initial assembly for DROP,

laboratory OTUs were designated by their strain name; most laboratory strains can be

associated with provisional species but some cannot yet be assigned. Three females and

three males of each strain were dry-mounted and individually assigned a USNMENT 'QR

code' specimen label as representative vouchers. For each molecular voucher, three legs

from a female wasp were removed for DNA extraction and sequencing (Supplementary

Methods for details), and the rest of the body was assigned a USNMENT specimen label

and preserved for morphological identification. Both DNA extraction and vouchers were

entered into the database and uploaded to BOLD (DROP project: DS-LABS) with an

associated GenBank ID [NOTE: the BOLD records will be pushed to GenBank at revision

stage; these data are not embargoed].

Where possible, we identified OTU strains using a combination of morphological and

sequence data, and characterized provisional species or species clusters using neighbor-

joining trees (Figure S1) based on the COI gene sequences (Supplemental material). For

establishing BIN limits in the context of DROP, we have adopted an initial percent cutoff

at 2%. As Ratnasingham & Hebert (2013) pointed out, this is a good starting point for

many taxa, but it also may be adjusted as more samples are acquired and compared.

Drosophila parasitoid database—DROP

To compile the above information, we built a simple Structured Query Language

(SQL) database in sqlite3 format using SQLiteStudio. There are eight linked tables in the

database—species, strain, voucher, sequence, genome, transcriptome, proteome and

reference—along with additional tables for linking these to reference table (Figure S2).

The database incorporates all sample fields used by BOLD for compatibility and includes

a number of new fields to accommodate a catalogue of *Drosophila* parasitoid species,

laboratory strain information, and links from the DROP database to BOLD and GenBank

records.

DROP is available on Zenodo (http://doi.org/10.5281/zenodo.4519656) for

permanent deposition and version control. In addition to the main database, the

Zenodo repository includes additional files to facilitate easy use of the database. These

files include: 1) the reference database in comma-separated text (.csv) and FASTA

format ready to be used for molecular identification; 2) a species catalogue with

taxonomic information; and 3) a list of laboratory strains with confirmed molecular

vouchers. DROP will be continued to be maintained by C-HL until further notice at the

Zenodo repository and sequences generated in the future will also be deposited in BOLD

(DROP project).

Species, provisional species, and OTU designations

In addition to the inherent value of a formal taxonomic name, a reliable provisional

taxon label can also be used for exchanging scientific information and conveying

experimental results among researchers (Schindel & Miller, 2010). Based on the amount

of sequence divergence between described species, we observed what appears to be a

significant number of provisional OTUs in the initial dataset we compiled. Furthermore,

among the data linked to a valid species name, some of these provisional OTUs are

actively being used in research and have sequences available to the public. We

therefore provide a list of potential new species with their molecular vouchers.

We use the following designation format for OTUs that refer to a provisional species:

"Drop_strainX_sp.1" or, when no other information is known, "DROP_sp.1". Where

possible, these OTUs are linked to BINs within BOLD and to a voucher USNM specimen

label number. If the genus of the OTU is known, the "Drop_Leptopilina_sp.1" format is

followed. These designations can facilitate species identification as well as discovery and

description of new species without compromising the existing taxonomy of the

described OTUs in question. As more complete species descriptions become available,

this provisional species framework can be updated while keeping the link to previous

provisional species name.

Results

Overview of DROP

We catalogued 182 OTUs in the DROP database with 113 described species of

Drosophila parasitoids and 69 provisional species (Table 1). In total, we documented 154

laboratory strains (Table S3), 853 vouchers from 36 institutions (Table S2). Among the

described species, 98 have voucher information, of which 61 are traceable to type

specimens, including 45 to holotypes (i.e., specimen used to root a name to the

taxonomic author's concept of the species). *Leptopilina* is represented by the highest

number of species with 45 OTUs, followed by *Asobara* with 26 OTUs. Within the 154

catalogued lab strains, 86 were actively being used in ongoing research (i.e., a live strain

being cultivated). These strains represent 39 OTUs: 11 described species and 28

provisional species (Table S3, Figure S1).

Molecular Vouchers

So far, DROP includes 545 DNA sequences and links to 16 genomes (Table 2.1), 11

transcriptomes (Table 2.2), and 6 proteomes (Table 2.3). From the total of 8298 DNA

sequences (dataset: DS-DROPAR) collected from public databases, only 322 sequences

(less than 4% of available sequences) satisfied the criteria for validity we imposed for

molecular vouchers (see material and methods) included in DROP. The DS-DROPAR

dataset initially referred to 520 taxa names, but only 52 names were valid, linked to

vouchers, or linked to a publication with evidence that the specimens had been

identified by taxonomists. The remaining 223 of 545 DROP DNA sequences were

generated by this project (dataset: DS-LABS, DS-AUSPTOID) and came from 121 OTUs

(12 provisional species and 101 lab strains).

The DROP database is largely made up of standard barcode COI sequences (340

sequences), which includes 77 OTUs: 43 described species and 33 provisional species.

We aimed to supplement COI with secondary markers (28SD2, 18S, ITS2) when possible,

resulting in an additional 120 sequences from 26 OTUs: 15 described species and 11

provisional species. There are currently 19 OTUs that have sequences from more than

one genetic marker.

Species Delimitation in Laboratory Strains

We used 298 COI sequences to resolve the identification of each laboratory

strain, and where possible, indicated potential species clusters for *Drosophila*

parasitoids (Fig. S1 and Table S3). Using a fixed 2% divergence cutoff, a total of 31 lab

strain OTUs were assignable to a valid species name, and the remaining 70 strain OTUs

were assigned to a provisional species. The taxonomic status of several of these

provisional species is also being investigated using an integrative taxonomic approach

involving morphological identification, genomic data, or other genetic data.

Discussion

In this paper, we introduce and describe a free and open-access database for the

reliable molecular identification of *Drosophila* parasitoids. The guiding principle of DROP

is data credibility, based on the prerequisite that genetic data be associated with explicit

criteria linking voucher specimens with taxonomic concepts of the original authors

(Troudet et al., 2018). When incorporating information from public genetic databases,

we include only sequences that have passed our filtering protocol. This protocol ensures

each entry is associated with a valid scientific name, provisional name, or consistently

applied OTU designation that can be used to integrate genetic and organismal data from

independent studies.

The following discussion expands on the utility of DROP and how we hope it will

benefit molecular species identification, connect research from various disciplines,

support biological control applications, and serve as a long-term molecular voucher

repository and clearinghouse for vetted data.

Molecular (mis-)identification

We observe that 17% of the described *Drosophila* parasitoid OTUs in BOLD and

GenBank (dataset: DS-DROPAR) are associated with more than one BIN; these are

examples of BIN-ID conflict. Roughly half of these OTUs are used as lab strains. This

latter observation is disturbing, because it demonstrates that the criteria used to

differentiate and reference species in active research programs are clouded. For

example, BIN-ID conflicts were observed in the *Drosophila* parasitoids *Ganaspis*

brasiliensis (Ihering) and *Asobara japonica* Belokobylskij (Table S1), both of which are in

active use in numerous research programs (e.g. Moreau et al., 2009; Nomano et al.,

2017; Reumer et al., 2012; Wang et al., 2020a & 2021) as well as in biological control

efforts against the invasive *D. suzukii* (e.g. Abram et al., 2020; Daane et al., 2016;

Giorgini et al., 2019). All the BINs from *G. brasiliensis* carry the name *G. xanthopoda*

(Figure 1). In such instances, assigning an identification by matching specimens to

barcode records in the genetic database is problematic, as two names are applied to the

same BIN. If sequences comprising the BIN are not linked to a voucher that can be

examined, teasing apart the two names and how they are applied is impossible.

Applying explicit, consistent criteria for species determination ensures that

experimental results can be reliably repeated, and that any potentially novel

observations will not be explained away as artifacts of identification. DROP addresses

these concerns by linking reliable reference sequences and vouchers for *G. brasiliensis*

(Figure 1) and from different studies: one with reference to the morphological

description (Buffington & Forshage, 2016) and the other with reference to the genome

(using voucher specimens from the morphological study; Blaimer et al., 2020).

We were not able to resolve all conflicts between BIN and species identity, for one

or more of the following three reasons: First, many records lack reliably identified

vouchers and have often been themselves used for molecular identification,

proliferating errors. Second, in some cases, it is not possible to verify whether the

genetic differences among BINs represent different species or simply intraspecific

genetic variation (Bergsten et al., 2012), because BINs themselves are not a species

concept. The only solution to this problem is to derive original sequence data from type

specimens (which is often either impractical or impossible for a number of technical

reasons), or from specimens whose conspecificity with the types has been corroborated.

Since species boundaries are always subject to testing, additional specimens from

multiple collecting events (e.g., representing different seasons and geographic regions)

may help provide the additional data to circumscribe a given species' limits. The third

difficulty in resolving BIN-ID conflict derives from the data themselves: Although the

mitochondrial COI gene is the locus most frequently chosen for identification of insects

and other animals, its effectiveness varies among insect groups (Brower & DeSalle,

2002; Gompert et al., 2008; Lin & Danforth, 2004). In part, this derives from gene-

tree/species-tree conflict as a function of mitochondrial DNA introgression (Gompert et

al., 2008; Klopfstein et al., 2016), parthenogenesis (Reumer et al., 2012), and/or

Wolbachia infection (Wachi et al., 2015; Xiao et al., 2012), any of which may lead to

complications in species delimitation using mitochondrial loci. Ideally, studies should

apply multiple loci, genomes, and comparative taxonomic data to clarify species

boundaries. As *Drosophila* parasitoids are often maintained in laboratory cultures, it is

also possible to use mating experiments to explore species boundaries under the

paradigm of the biological species concept (Seehausen et al., 2020).

DROP as a taxonomic tool

DROP offers an empirical platform for species discovery and a useful tool for

taxonomic research. The fact that the number of BINs reported here exceeds the

number of described species (Table S1, Figure S3) highlights the need for taxonomic

work. But such work cannot proceed on the basis of BINs or barcodes, but requires

integrative taxonomic approach employing a combination of molecular and

morphological data. Describing new species on the sole basis of a barcode or BIN,

without the benefit of independent character data, should, in general, be avoided. It

risks creating nomenclatural synonymy if it is later determined that a sequence can be

attributed to a specimen that bears a valid, available name. Moreover, BINs are based

on distance analyses which, by definition, are incompatible with diagnoses per se

(Ferguson, 2002; Prendini et al., 2002; Goldstein & DeSalle, 2011). Therefore, in

taxonomic treatments, it is critical to clarify the range of applicability of a given BIN and

it overlap with a taxonomic name (see example in Figure 1).

466   Public genetic databases have adopted a longstanding convention in treating

undetermined OTUs and sequences, referring to provisional species with numbers, as

for example "sp. 1", and these are rarely linked to vouchers. For OTUs designated as

provisional species, DROP enables cross-indexing of specimens, sequences and

references with studies and provides researchers with valuable tools for taxonomic

revisions, including the means of discovery, corroboration, and description of new

species. For example, "drop_Gan1_sp.1" refers to voucher USNMENT01557320

deposited in the USNM, Washington DC, COI sequence (DROP sequence_id 2), BOLD

process ID: DROP143-21, BIN number: XXXXXX (will update in the revision), 28D1

sequences (DROP sequence_id 289), and 28D2 sequences (DROP sequence_id 303). In

the future, when "drop_Gan1_sp.1" is described as a new species with a formal specific

epithet, DROP will update the species status and holotype information while keeping

this provisional species name as an informal "synonym." We recognize tracking these

informal 'tags' through time can be problematic; however, linking these tags in DROP to

a vouchered specimen and unique identifier will minimize confusion.

From molecular mechanisms to ecosystem structure

The use of molecular tools in insect biodiversity studies has gradually expanded from

barcoding single individuals to metabarcoding large environmental samples

representing entire food webs (Jeffs et al., 2020; Littlefair et al., 2016). *Drosophila* and

their parasitoids are among the few systems that currently allow us to explore

thoroughly the mechanisms of species interactions at scales ranging from the molecular

to the ecological. Here, we highlight two examples where information compiled in DROP

enables the study of the *Drosophila*-parasitoid system across multiple levels of biological

organization:

DROP includes a DNA reference library of Australian *Drosophila* parasitoids (DS-

AUSPTOID in BOLD) that connects laboratory experiments and field research. Molecular

vouchers of both hosts and parasitoids were collected along altitudinal gradients in the

rainforest of northern Queensland, Australia (Jeffs et al., 2020). With this DNA reference

library, researchers can detect interactions between *Drosophila* and their parasitoids

using PCR-based approaches and parasitized pupae (Hrcek & Godfray, 2015; Jeffs et al.,

2020). Surveying host-parasitoid interactions in this way will improve our understanding

of how environmental change alters the structure of host-parasitoid networks (Morris et

al., 2014; Staniczenko et al., 2017; Tylianakis et al., 2007) by accelerating data collection

in the field. In addition, JH established lab cultures of both hosts and their parasitoids

from the same Australian sampling sites with the aim of conducting laboratory

experiments (e.g. Thierry et al., 2021). Molecular vouchers of the lab strains were then

submitted to DROP as a reference database (DS-LABS in BOLD) to ensure that criteria for

species determination were applied consistently—and will continue to be applied

consistently—between the natural community studies and the laboratory experiments.

The presence of a foundational DNA reference library and species catalogue in

DROP will enable the process of exploring parasitoid biodiversity to become more

efficient. For example, DROP includes molecular vouchers from *Drosophila* parasitoids

that were collected across seasons and along latitudinal gradients in the eastern Unites

States (Lue et al., 2016, 2018). These data proved to be extremely useful for identifying

species in a more recent exploration of native parasitoid biodiversity across North

America (e.g., Abram et al., 2020). There are additional uses for DROP: curated

specimen collections may be used to document species distributions, phenology,

understand micro-evolutionary patterns, observe the effects of climate change, and

detect and track biological invasions (Funk, 2018; Schilthuizen et al., 2015; Tarli et al.,

2018).

Taxonomic accuracy for biocontrol studies

Unfortunately, the history of biological control includes many examples of

misidentifications that have resulted in failures to employ or establish the expected

control agent, thus hindering eventual success (Buffington et al., 2018; Rosen, 1986;

Huffaker et al. 1962). In the context of biological control research on *Drosophila* pest

species, a simple, reliable, and rapid identification tool for their natural enemies is

essential (Wang et al. 2020b). By anchoring the criteria for determining identities of

organisms being considered for biological control programs, DROP annotation enables

the direct examination of centers of origin for parasitoid species, their co-occurrence

with natural enemies, and the optimal timing for potential introductions of such

enemies (Abram et al., 2020; Daane et al., 2016; Girod et al., 2018a and b; Kimura, 2015;

Mitsui et al., 2007). Because most sequences from DROP are already vetted for

reliability, they can be used to identify biological control agents rapidly, before or after

being brought into quarantine facilities for safety and efficacy testing. This will decrease

the risk of non-target ecological impacts arising from misidentifications and facilitate

regulatory review for releases of effective and specific natural enemies.

In addition to species identification, reference sequences from DROP may be used to

create species-specific primers for the accurate identification of parasitoids, design

multiplex PCR assays that rapidly distinguish species in natural or agricultural

ecosystems (Ye et al., 2017), and apply high-throughput molecular identification

diagnostics (Fagan-Jeffries et al., 2018). Applications of such specific primers have been

used in bacteria, fungi, oomycetes and insect pests (Liu et al., 2017; Tedersoo et al.,

2019; Tsai et al., 2020).

Long-term molecular voucher preservation

During the curation of DROP, we found that holotype specimens were missing from

museums for several iconic *Drosophila* parasitoid species: *Asobara tabida* (Nees von

Esenbeck), *Leptopilina clavipes* (Hartig), and *Leptopilina longipes* (Hartig)*.* This is not

uncommon and impedes future taxonomic revisions regardless of whether molecular

data are used. To avoid contributing to this problem, DROP uses museums as

depositories for ensuring that sequenced vouchers of both described species and

provisional species are permanently stored. In order to stabilize nomenclature, we

further advocate the designation of neotypes (a replacement specimen for a missing

holotype) that have museum-vouchered DNA barcodes and additional genomic

extractions in storage.

Natural history museums are designed to maintain vouchers (including types) for

long-term preservation, and increasingly they implement institutionalized workflows

that link DNA sequences to specimens and specimen metadata (Prendini et al., 2002).

We strongly encourage the deposition of voucher specimens from field surveys and

experimental studies in museum collections, as has been urged by the Entomology

Collections Network (ECN) and required in many PhD programs. No matter how quickly

new molecular techniques are developed or refined, there is no substitute for a reliable

database of voucher specimens when it comes to ensuring the repeatability of biological

research (Funk et al., 2005; Lendemer et al., 2020).

Our results show that species richness of the parasitic wasps that attack *Drosophila*

is severely underestimated, and only a fraction of them have been described. In DROP,

38% of the OTUs are provisional species, and more than 46% of the named OTUs have

synonyms. Remarkably, *Leptopilina heterotoma*, one of the world's most studied

parasitoids, has more than 20 synonyms! As is generally the case, the rate of species

description and revision of *Drosophila* parasitoids lags far behind that with which

molecular sequence data are generated. Ensuring a consistent application of OTU

recognition is therefore essential. With DROP, researchers may ensure consistency is

their application of scientific names, and that those names are valid, making the

daunting process of describing *Drosophila* parasitoids more accurate and efficient.

In addition to the collection of physical museum resources, a central role

taxonomists play in DROP and its curation is that of fostering better integration of

taxonomy with experimental and biodiversity research. Our intention is to perpetuate

DROP beyond this introductory publication. We hope that experts in all areas of

Drosophila-parasitoid biology and related fields will join us in this effort.

Conclusion

Taxonomic confusion presents many obstacles in experimental and biodiversity

studies. One way of addressing this impediment is to provide a reliable DNA library with

traceable vouchers (Astrin et al., 2013). We developed DROP as a resource and platform

for gathering and sharing reliable genomic sequence data for *Drosophila* parasitoids. We

hope it will serve as a model for researchers working with organisms which present

similar difficulties. While compiling DROP, we found that the high number of provisional

versus named OTUs suggests that the diversity of parasitic wasps attacking *Drosophila* is

greatly underestimated. With this in mind, DROP represents the start of an important

knowledge base that will strengthen future studies of natural host-parasitoid

interactions, population dynamics, biocontrol, and the impact of climate change on

biodiversity and ecosystem services.

Acknowledgements

The DROP project was developed during the 2018 Entomology Society of

America conference, during the symposium "Drosophila parasitoids: from molecular to

ecosystem level". We thank Dr. Elijah Talamas for valuable comments on earlier drafts.

We also thank Chris Jeffs for providing Australian field samples. We are also thankful for

funding support from the Czech Science Foundation (17-27184Y). Additional fund for

sequencing was provided by MLB, OTL, and PPAS. Mention of trade names or

commercial products in this publication is solely for the purpose of providing specific

information and does not imply recommendation or endorsement by the USDA. USDA is

an equal opportunity provider and employer.

References

Abram, P. K., Mcpherson, A. E., Kula, R., Hueppelsheuser, T., Perlman, S. J., Curtis, C. I.,
Fraser, J. L., ... Buffington, M. (2020). New records of *Leptopilina*, *Ganaspis*, and
Asobara species associated with *Drosophila suzukii* in North America, including
detections of L. japonica and G. brasiliensis. *Journal of Hymenoptera Research*, 78,
1-17, https://doi.org/10.3897/jhr.78.55026
Astrin, J. J., Zhou, X., & Misof, B. (2013). The importance of biobanking in molecular
taxonomy, with proposed definitions for vouchers in a molecular context. *ZooKeys*,
365(SPEC.ISSUE), 67–70. https://doi.org/10.3897/zookeys.365.5875
Baker, C. C. M., Bittleston, L. S., Sanders, J. G., & Pierce, N. E. (2016). Dissecting host-
associated communities with DNA barcodes. *Philosophical Transactions of the
Royal Society B: Biological Sciences*, 371(1702).
https://doi.org/10.1098/rstb.2015.0328

Bergsten, J., Bilton, D. T., Fujisawa, T., Elliott, M., Monaghan, M. T., Balke, M.,
Hendrich, … Vogler, A. P. (2012). The effect of geographical scale of sampling on
DNA barcoding. *Systematic Biology,* 61 (5), 851-869.
https://doi.org/10.1093/sysbio/sys037
Blaimer, B. B., Gotzek, D., Brady, S. G., & Buffington, M. (2020). Comprehensive
phylogenomic analyses re-write the evolution of parasitism within cynipoid wasps.
BMC Ecology and Evolution, 20 (155). https://doi.org/10.1186/s12862-020-01716-2
Brower, A. V. Z., & DeSalle, R. (2002). Patterns of mitochondrial versus nuclear DNA
sequence divergence among nymphalid butterflies: The utility of wingless as a
source of characters for phylogenetic inference. *Insect Molecular Biology*, 7 (1), 73-
82. https://doi.org/10.1046/j.1365-2583.1998.71052.x
Buffington, M., &Forshage, M. (2016). Redescription of *Ganaspis brasiliensis* (Ihering,
1905), new combination, (Hymenoptera: Figitidae) a natural enemy of the Invasive
Drosophila suzukii (Matsumura, 1931) (Diptera: Drosophilidae). *Proceedings of the
Entomological Society of Washington*, 118(1), 1–13. https://doi.org/10.4289/0013-
8797.118.1.1
Buffington, M., Talamas, E. J., & Hoelmer, K. A. (2018). Team Trissolcus: Integrating
taxonomy and biological control to combat the brown marmorated stink bug.
American Entomologist, 64 (4), 224–232
Carton, Y., Boulétreau, M., van Alphen, J. J. M., & van Lenteren, J. C. (1986). The
Drosophila parasitic wasps. In Ashburner M, Carson HL, Thompson JN (Eds), *The
genetics and biology of Drosophila*, (3),348–394.
Carton, Y., Haouas, S., Marrakchi, M., & Hochberg, M. (1991). Two competing parasitoid
species coexist in sympatry. *Oikos*, 60, 222-230. https://doi.org/10.2307/3544869
Condon, M. A., Scheffer, S. J., Lewis, M. L., Wharton, R., Adams, D.C., & Forbes, A. A.
(2014). Lethal interactions between parasites and prey increase niche diversity in a
tropical community. *Science*, *343*(6176), pp.1240-1244.
Daane, K. M., Wang, X.-G., Biondi, A., Miller, B. E., Miller, J. C., Riedl, H., Shearer, P. W.,
… Walton, V. M. (2016). First exploration of parasitoids of *Drosophila suzukii* in
South Korea as potential classical biological agents. *Journal of Pest Science* 89, 823–
835, doi:10.1007/s10340-016-0740-0.
Dolphin, K., & Quicke, D. L. J. (2001). Estimating the global incompletely described
parasitoid wasps. *Biological Journal Of The Linnean Society*, 73 (3), 279-286,
https://doi.org/10.1006
Escalante, P., Ibarra-Vazquez, A., & Rosas-Escobar, P. (2010). Tropical montane
nymphalids in Mexico: DNA barcodes reveal greater diversity. *Mitochondrial DNA*,
21, 30-37, https://doi.org/10.3109/19401736.2010.535527
Fagan-Jeffries, E. P., Cooper, S. J. B., Bertozzi, T., Bradford, T. M., & Austin, A. D. (2018).
DNA barcoding of microgastrine parasitoid wasps (Hymenoptera: Braconidae) using
high-throughput methods more than doubles the number of species known for
Australia. *Molecular Ecology Resources*, 18(5), 1132–1143.
https://doi.org/10.1111/1755-0998.12904
Ferguson, J. W. H. (2002). On the use of genetic divergence for identifying species.
Biological Journal of the Linnean Society, 75, 509–16.

Forbes, A. A., Bagley, R. K., Beer, M. A., Hippee, A. C., & Widmayer, H. A. (2018).
Quantifying the unquantifiable: Why Hymenoptera, not Coleoptera, is the most
speciose animal order. *BMC Ecology*, 18(1), 1–11. https://doi.org/10.1186/s12898-
018-0176-x
Funk, V. A. (2018). Collections-based science in the 21st Century. *Journal of Systematics*
and Evolution, 56(3), 175–193. https://doi.org/10.1111/jse.12315
Funk, V. A., Hoch, P. C., Prather, L. A., & Wagner, W. L. (2005). The importance of
vouchers. *Taxon*, 54(1), 127–129. https://doi.org/10.2307/25065309
Gariepy, T. D., Bruin, A., Konopka, J., Scott-Dupree, C., Fraser, H., Bon, M. C., & Talamas,
E. (2019). A modified DNA barcode approach to define trophic interactions
between native and exotic pentatomids and their parasitoids. *Molecular Ecology*,
28(2), 456–470. https://doi.org/10.1111/mec.14868
Giorgini, M., Wang, X.-G., Wang, Y., Chen, F. S., Hougardy, E. , Zhang, H. M., Chen, Z.
Q., … Guerrieri, E. (2019). Exploration for native parasitoids of *Drosophila suzukii* in
China reveals a diversity of parasitoid species and narrow host range of the
dominant parasitoid. *Journal of Pest Science*, 92(2), 509–522.
https://doi.org/10.1007/s10340-018-01068-3
Girod, P., Borowiec, N., Buffington, M., Chen, G., Fang, Y., Kimura, M. T., Peris-Felipo, F.
J., … Kenis, M. (2018). The parasitoid complex of *D. suzukii* and other fruit feeding
Drosophila species in Asia. *Scientific Reports*, 8(1), e11839.
https://doi.org/10.1038/s41598-018-29555-8
Girod, P., Lierhmann, O., Urvois, T., Turlings, T. C. J., Kenis. M., & Haye, T. (2018). Host
specificity of Asian parasitoids for potential classical biological control of *Drosophila*
suzukii. Journal of Pest Science 91,1241–1250, https://doi. doi:10.1007/s10340-
018-1003-z
Goldstein, P. Z., & DeSalle, R. (2011). Integrating DNA barcode data and taxonomic
practice: Determination, discovery, and description. *BioEssays*, 33(2),135-147,
https://doi.org/10.1002/bies.201000036
Gompert, Z., Forister, M. L., Fordyce, J. A., & Nice, C. C. (2008). Widespread mito-nuclear
discordance with evidence for introgressive hybridization and selective sweeps in
Lycaeides. *Molecular Ecology*, 17(24), 5231-5244, https://doi.org/10.1111/j.1365-
294X.2008.03988.x
Grissell, E. (1999). Hymenopteran biodiversity: some alien notions. *American*
Entomologist, 45,236-244.
Hardy, I. C., van Alphen, J. J. M., & Godfray, H. C. J. (1994). Parasitoids: Behavioral and
evolutionary ecology. *The Journal of Animal Ecology*, 63(4), 1009-1010,
https://doi.org/10.2307/5282
Hausmann, A., Haszprunar, G., & Hebert, P. D. N. (2011). DNA barcoding the geometrid
fauna of bavaria (Lepidoptera): Successes, surprises, and questions. *PLoS ONE*, 6(2),
1–9. https://doi.org/10.1371/journal.pone.0017134
Hausmann, A., Miller, S. E., Holloway, J. D., Dewaard, J. R., Pollock, D., Prosser, S. W. J.,
& Hebert, P. D. N. (2016). Calibrating the taxonomy of a megadiverse insect family:
3000 DNA barcodes from geometrid type specimens (Lepidoptera, Geometridae).
Genome, 59(9), 671–684. https://doi.org/10.1139/gen-2015-0197

Hebert, P. D. N., Ratnasingham, S., & DeWaard, J. R. (2003). Barcoding animal life:
Cytochrome c oxidase subunit 1 divergences among closely related species.
Proceedings of the Royal Society B: Biological Sciences, 270 (Suppl.), 96-99,
https://doi.org/10.1098/rsbl.2003.0025
Hendrich, L., Morinière, J., Haszprunar, G., Hebert, P. D. N., Hausmann, A., Köhler, F., &
Balke, M. (2015). A comprehensive DNA barcode database for Central European
beetles with a focus on Germany: Adding more than 3500 identified species to
BOLD. *Molecular Ecology Resources*, 15(4), 795-818, https://doi.org/10.1111/1755-
0998.12354
Hrček, J., & Godfray, H. C. J. (2015). What do molecular methods bring to host-parasitoid
food webs? *Trends in Parasitology*, 31(1), 30–35.
https://doi.org/10.1016/j.pt.2014.10.008
Hrcek, J., Miller, S. E., Quicke, D. L. J., & Smith, M. A. (2011). Molecular detection of
trophic links in a complex insect host-parasitoid food web. *Molecular Ecology*
Resources, 11(5), 786–794. https://doi.org/10.1111/j.1755-0998.2011.03016.x
Huffaker, C. B., Kennett, C. E., Finney, G. L. (1962). Biological control of olive scale,
Pwrlatoria oleae (Cohree), in California by imported Aphytis maculicornis (Masi)
(Hymenoptera: Aphelinidae). *Hilgardia,* 32 (13): 541-636. DOI:
10.3733/hilg.v32n13p541
Hughes, R. D., Woolcock, L. T. (1976). Aphaereta aotea sp. N. (Hymenoptera:
Braconidae), an Alysiine parasite of dung breeding flies. J*ournal of Australian*
Entomological Society, 15, 191-196.
Jeffs, C. T., Terry, J. C. D., Higgie. M., Jandová, A., Konvičková. H., Brown. J. J., Lue. C.-H.,
… Lewis, O. T. (2020). Molecular analyses reveal consistent food web structure with
elevation in rainforest *Drosophila* - parasitoid communities. *Ecography*, 43, 1-11,
https://doi.org/10.1111/ecog.05390
Kim-Jo, C., Gatti, J. L., & Poirié, M. (2019). *Drosophila* cellular immunity against
parasitoid wasps: A complex and time-dependent process. *In Frontiers in*
Physiology, https://doi.org/10.3389/fphys.2019.00603
Kimura, M. T. (2015). Prevalence of exotic frugivorous *Drosophila* species, *D. simulans*
and *D. immigrans* (Diptera: Drosophilidae), and its effects on local parasitoids in
Sapporo, northern Japan. *Applied Entomology and Zoology*, 50(4), 509–515.
https://doi.org/10.1007/s13355-015-0361-8
Kimura, M. T., & Mitsui, H. (2020). *Drosophila* parasitoids (Hymenoptera) of Japan. I*n*
Entomological Science, 23(4), 359-368, https://doi.org/10.1111/ens.12432
Klopfstein, S., Kropf, C., & Baur, H. (2016). Wolbachia endosymbionts distort DNA
barcoding in the parasitoid wasp genus Diplazon (Hymenoptera: Ichneumonidae).
Zoological Journal of the Linnean Society, 177(3), 541–557.
https://doi.org/10.1111/zoj.12380
Kraaijeveld, A. R., & Godfray, H. C. J. (1997). Trade-off between parasitoid resistance and
larval competitive. *Nature*, 389, 278-280, https://doi.org/10.1038/38483
Lefèvre, T., De Roode, J. C., Kacsoh, B. Z., & Schlenke, T. A. (2012). Defence strategies
against a parasitoid wasp in Drosophila: Fight or flight? *Biology Letters*, 8(2), 230-
233, https://doi.org/10.1098/rsbl.2011.0725

Lendemer, J., Thiers, B., Monfils, A. K., Zaspel, J., Ellwood, E. R., Bentley, A., LeVan, K., …
Aime, M. C. (2020). The extended specimen network: A strategy to enhance US
biodiversity collections, promote research and education. *BioScience*, 70(1), 23-30,
https://doi.org/10.1093/biosci/biz140
Lin, C. P., & Danforth, B. N. (2004). How do insect nuclear and mitochondrial gene
substitution patterns differ? Insights from Bayesian analyses of combined datasets.
Molecular Phylogenetics and Evolution, 30(3), 686-702,
https://doi.org/10.1016/S1055-7903(03)00241-0
Littlefair, J. E., Clare, E. L., & Naaum, A. (2016). Barcoding the food chain: From Sanger to
high-throughput sequencing1. *Genome*, 59(11), 946–958.
https://doi.org/10.1139/gen-2016-0028
Liu, L. J., Pang, A. H., Feng, S. Q., Cui, B. Y., Zhao, Z. H., Kučerová, Z., Stejskal, V., … Li, Z.
H. (2017). Molecular Identification of ten species of stored-product psocids through
microarray method based on ITS2 rDNA. *Scientific Reports*, 7(1): 16694,
https://doi.org/10.1038/s41598-017-16888-z
Lue, C.-H., Borowy, D., Buffington, M. L., & Leips, J. (2018). Geographic and seasonal
variation in species diversity and community composition of frugivorous *Drosophila*
(Diptera: Drosophilidae) and their *Leptopilina* (Hymenoptera: Figitidae) parasitoids.
Environmental Entomology, 47(5): 1096-1106. https://doi.org/10.1093/ee/nvy114
Lue, C.-H., Driskell, A. C., Leips, J., & Buffington, M. L. (2016). Review of the genus
Leptopilina (Hymenoptera, Cynipoidea, Figitidae, Eucoilinae) from the Eastern
United States, including three newly described species. J*ournal of Hymenoptera
Research*, 53: 35-76. https://doi.org/10.3897/jhr.53.10369
Meiklejohn, K. A., Damaso, N., & Robertson, J. M. (2019). Assessment of BOLD and
GenBank – Their accuracy and reliability for the identification of biological
materials. *PLoS ONE,* 14(6): e0217084.
https://doi.org/10.1371/journal.pone.0217084
Melk, J. P., & Govind, S. (1999). Developmental analysis of *Ganaspis xanthopoda*, a larval
parasitoid of *Drosophila melanogaster*. *Journal of Experimental Biology*, 202, 1885-
1896
Miller, S. E., Hausmann, A., Hallwachs, W., & Janzen, D. H. (2016). Advancing taxonomy
and bioinventories with DNA barcodes. *Philosophical Transactions of the Royal
Society Biological Sciences*, 371(1702): 20150339. doi: 10.1098/rstb.2015.0339
Mitsui, H., van Achterberg, K., Nordlander, G., & Kimura, M. T. (2007). Geographical
distributions and host associations of larval parasitoids of frugivorous *Drosophilidae*
in Japan. *Journal of Natural History*, 41(25–28), 1731–1738.
https://doi.org/10.1080/00222930701504797
Moreau, S. J. M., Vinchon, S., Cherqui, A., & Prévost, G. (2009). Components of Asobara
venoms and their effects on hosts. *In Advances in Parasitology*, Prévost G (Ed). 70,
217-232, https://doi.org/10.1016/S0065-308X(09)70008-9
Morris, R. J., Gripenberg, S., Lewis, O. T., & Roslin, T. (2014). Antagonistic interaction
networks are structured independently of latitude and host guild. *Ecology Letters*,
17(3), 340-349, https://doi.org/10.1111/ele.12235
Nappi, A. J., & Carton, Y. (2001). Immunogenetic aspects of the cellular immune

response of *Drosophila* against parasitoids. *Immunogenetics*, 52(3–4), 157–164.
https://doi.org/10.1007/s002510000272
Nomano, F. Y., Kasuya, N., Matsuura, A., Suwito, A., Mitsui, H., Buffington ,M.L., &
Kimura, M. T. (2017). Genetic differentiation of *Ganaspis brasiliensis*
(Hymenoptera: Figitidae) from East and Southeast Asia. *Applied Entomology and*
Zoology, 52(3), 429–437. https://doi.org/10.1007/s13355-017-0493-0
Novotny, V., & Miller, S. E. (2014). Mapping and understanding the diversity of insects in
the tropics: Past achievements and future directions. *Austral Entomology*, 53(3),
259–267. https://doi.org/10.1111/aen.12111
O'Grady, P. M., & DeSalle, R. (2018). Phylogeny of the genus *Drosophila*. *Genetics*,
209(1), 1–25. https://doi.org/10.1534/genetics.117.300583
Pentinsaari, M., Ratnasingham, S., Miller, S. E., & Hebert, P. D. N. (2020). BOLD and
GenBank revisited – Do identification errors arise in the lab or in the sequence
libraries? *PLoS One,* 15(4): e0231814. https://doi.org/10.1371/journal.
pone.0231814
Pfeiffer, D. G., Shrader, M. E., Wahls, J. C. E., Willbrand, B. N., Sandum, I., van der Linde,
K., Laub, C. A., … Day, E. R. (2019). African Fig Fly (Diptera: Drosophilidae): Biology,
expansion of geographic range, and its potential status as a soft fruit pest. *Journal*
of Integrated Pest Management, 10(1), 1–8. https://doi.org/10.1093/jipm/pmz018
Prendini, L., Hanner, R., & DeSalle, R. (2002). Obtaining, storing and archiving specimens
and tissue samples for use in molecular studies. *In Techniques in Molecular*
Systematics and Evolution. https://doi.org/10.1007/978-3-0348-8125-8_11
Prévost, G. (2009). Parasitoids of *Drosophila*. *In Advances in parasitology*.
https://doi.org/10.1016/S0065-308X(09)70018-1
Quicke, D. L. J. (2015). The Braconid and Ichneumonid parasitoid wasps: Biology,
systematics, evolution and ecology. Wiley-Blackwell,
https://doi.org/10.1002/9781118907085
Ratnasingham, S., & Hebert, P. D. N. (2007). BARCODING: bold: The Barcode of Life Data
System (http://www.barcodinglife.org). *Molecular Ecology Notes*, 7(3), 355-364,
https://doi.org/10.1111/j.1471-8286.2007.01678.x
Ratnasingham, S., & Hebert, P. D. N. (2013). A DNA-based registry for all animal species:
the barcode index number (BIN) system. *PLoS ONE,* 8(7): e66213.
https://doi.org/10.1371/journal.pone.0066213
Raupach, M. J., Hendrich, L., Kuchler, S. M., Deister, F., Moriniere, J., & Gossner, M. M.
(2014). Building-Up of a DNA Barcode Library for true bugs (Insecta: Hemiptera:
Heteroptera) of Germany reveals taxonomic uncertainties and surprises. *PLoS ONE*,
9(9), 1–13. https://doi.org/10.1371/journal.pone.0106940
Remsen, J., & O'Grady, P. (2002). Phylogeny of Drosophilinae (Diptera: Drosophilidae),
with comments on combined analysis and character support. *Molecular*
Phylogenetics and Evolution, 24(2), 249-264, https://doi.org/10.1016/S1055-
7903(02)00226-9
Reumer, B, M., van Alphen, J. J. M., & Kraaijeveld, K. (2012). Occasional males in
parthenogenetic populations of Asobara japonica (Hymenoptera: Braconidae): Low
Wolbachia titer or incomplete coadaptation. *Heredity*, 108(3), 341-346,

https://doi.org/10.1038/hdy.2011.82
Rosen, D. (1986). The role of taxonomy in effective biological control programs.
Agriculture, Ecosystems & Environment, 15(2-3), 121-129.
https://doi.org/10.1016/0167-8809(86)90085-X
Santos, W. G. N., Fernandes, E. C., Souza, M. M., Guimarães, J. A., & Araujo, E. L. (2016).
First record of Eucoilinae (Hymenoptera: Figitidae), parasitoids of African fig fly
Zaprionus indianus Gupta (Diptera: Drosophilidae), in the Caatinga biome.
Semina:Ciencias Agrarias, 37(5), 3055–3058. https://doi.org/10.5433/1679-
0359.2016v37n5p3055
Schilthuizen, M., Vairappan, C. S., Slade, E. M., Mann, D. J., & Miller, J. A. (2015).
Specimens as primary data: Museums and "open science." *Trends in Ecology and
Evolution*, 30(5), 237–238. https://doi.org/10.1016/j.tree.2015.03.002
Schlenke, T. A., Morales, J., Govind, S., & Clark, A. G. (2007). Contrasting infection
strategies in generalist and specialist wasp parasitoids of *Drosophila melanogaster*.
PLoS Pathogens, 3(10):e158, https://doi.org/10.1371/journal.ppat.0030158
Schindel, D., & Miller, S. E. (2010). Provisional Nomenclature the on-ramp to taxonomic
names. In: Polaszek, A. (Ed), *Systema Nature*, 250: The Linnaean Ark. CRC, Boca
Raton, 109-115.
Seehausen, M. L., Ris, N., Driss, L., Racca, A., Girod, P., Warot, S., Borowiec, N., Tosevski,
I., & Kenis, M. (2020). Evidence for a cryptic parasitoid species reveals its suitability
as a biological control agent. *Scientific reports*, 10: 19096.
https://doi.org/10.1038/s41598-020-76180-5
Shokralla, S., Spall, J. L., Gibson, J. F., & Hajibabaei, M. (2012). Next-generation
sequencing technologies for environmental DNA research. *In Molecular Ecology*,
21(8), 1794-1805, https://doi.org/10.1111/j.1365-294X.2012.05538.x
Smith, M. A., Fisher, B. L., & Hebert, P. D. N. (2005). DNA barcoding for effective
biodiversity assessment of a hyperdiverse arthropod group: The ants of
Madagascar. *Philosophical Transactions of the Royal Society Biological Sciences*,
360(1462), 1825-1834, https://doi.org/10.1098/rstb.2005.1714
Smith, M. A., Fernandez-Triana, J. L., Eveleigh, E., Gomez, J., Guclu, C., Hallwachs, W.,
Hebert. P. D. N., … Zaldivar-Riveron, A. (2013). DNA barcoding and the taxonomy of
Microgastrinae wasps (Hymenoptera, Braconidae): impacts after 8 years and nearly
20000 sequences. *Molecular Ecology Resource*s, 13, 168-276,
https://doi.org/10.1111/1755-0988.12038
Staniczenko, P. P. A., Reed-Tsochas, F., Lewis, O. T., Tylianakis, J. M., Albrecht, M.,
Coudrain, V., & Klein, A. M. (2017). Predicting the effect of habitat modification on
networks of interacting species. *Nature Communications*, 8, 792,
https://doi.org/10.1038/s41467-017-00913-w
Tarli, V. D., Grandcolas, P., & Pellens, R. (2018). The informative value of museum
collections for ecology and conservation: A comparison with target sampling in the
Brazilian Atlantic forest. *PLoS ONE*, 13(11).
https://doi.org/10.1371/journal.pone.0205710
Tedersoo, L., Drenkhan, R., Anslan, S., Morales-Rodriguez, C., & Cleary, M. (2019). High-
throughput identification and diagnostics of pathogens and pests: Overview and

practical recommendations. *Molecular Ecology Resources*, 19(1), 47–76.
https://doi.org/10.1111/1755-0998.12959

Terry, J. C. D., Chen, J., & Lewis, O. T. (2020). The effect of natural enemies on the
coexistence of competing species - an empirical test using Bayesian modern
coexistence theory. *bioRxiv*: https://doi.org/10.1101/2020.08.27.270389

Thierry, M., Pardikes, N. A., Lue, C.-H., Lewis, O. L., & Hrcek, J. (2021). Experimental
warming influences species abundances in a *Drosophila* host community through
direct effects on species performance rather than altered competition and
parasitism. *PLOS ONE* (In Press).

Troudet, J., Vignes-Lebbe, R., Grandcolas, P., & Legendre, F. (2018). The increasing
disconnection of primary biodiversity data from specimens: How does it happen
and how to handle it? *Systematic Biology*, 67(6), 1110–1119.
https://doi.org/10.1093/sysbio/syy044

Tsai, C.-L., Chu, I.-H., Chou. M.-H., Chareonviriyaphap, T., Chiang, M.-Y., Lin, P.-A., Lu, K.-
H., & Yeh, W.-B. (2020). Rapid identification of the invasive fall armyworm
Spodoptera frugiperda (Lepidoptera, Noctuidae) using species-specific primers in
multiplex PCR. *Scientific Reports*, 10(1), 1–9. https://doi.org/10.1038/s41598-020-
73786-7

Tylianakis, J. M., Tscharntke, T., & Lewis, O. T. (2007). Habitat modification alters the
structure of tropical host-parasitoid food webs. *Nature*, 445(7124), 202–205.
https://doi.org/10.1038/nature05429

Wachi, N., Nomano, F. Y., Mitsui, H., Kasuya, N., & Kimura, M. T. (2015). Taxonomy and
evolution of putative thelytokous species of Leptopilina (Hymenoptera: Figitidae)
from Japan, with description of two new species. *Entomological Science*, 18(1), 41–
54. https://doi.org/10.1111/ens.12089

Wang, X.-G., Biondi, A., & Daane, K. M. (2020). Functional responses of three candidate
Asian larval parasitoids evaluated for classical biological control of *Drosophila*
suzukii. Journal of Economic Entomology, 113(1): 73–80. doi: 10.1093/jee/toz265

Wang, X.-G., Biondi, A., Nance. A. N., Zappalà, L., Hoelmer, K. A., & Daane, K. M. (2021).
Assessment of Asobara japonica as a potential biological control agent for the
spotted wing drosophila, Drosophila suzukii. *Entomologia Generalis (In Press)* doi:
10.1127/entomologia/2020/1100

Wang, X.-G., Lee, J., Daane, K.M., Buffington, M., & Hoelmer, K. A. (2020). Biological
control of *Drosophila suzukii*. *CAB Reviews* 54, 10.1079/PAVSNNR202015054

Xiao, J. H., Wang, N. X., Murphy, R. W., Cook, J., Jia, L. Y., & Huang, D. W. (2012).
Wolbachia infection and dramatic intraspecific mitochondrial DNA divergence in a
fig wasp. *Evolution*, 66, 1907-1916, https://doi.org/10.1111/j.1558-
5646.2011.01561.x

Xie. J., Tiner, B., Vilchez, I., & Mateos, M. (2011). Effect of the *Drosophila* endosymbiont
Spiroplasma on parasitoid wasp development and on the reproductive fitness of
wasp-attacked fly survivors. *Evolutionary Ecology*, 25, 1065-1079,
https://doi.org/10.1007/s10682-010-9453-7

Xie, J., Winter, C., Winter, L., & Mateos, M. (2015). Rapid spread of the defensive
endosymbiont *Spiroplasma* in *Drosophila hydei* under high parasitoid wasp

pressure. *FEMS Microbiology Ecology*, 91(2), 1-11,
https://doi.org/10.1093/femsec/iu017
Yassin, A. (2013). Phylogenetic classification of the Drosophilidae Rondani (Diptera): The
role of morphology in the postgenomic era. *Systematic Entomology*,
https://doi.org/10.1111/j.1365-3113.2012.00665.x
Yassin, A., & David, J. R. (2010). Revision of the Afrotropical species of *Zaprionus*
(Diptera, Drosophilidae), with descriptions of two new species and notes on
internal reproductive structures and immature stages. *ZooKeys,* 51, 33-72,
https://doi.org/10.3897/zookeys.51.380
Ye, Z., Vollhardt, I. M. G., Girtler, S., Wallinger, C., Tomanovic, Z., & Traugott, M. (2017).
An effective molecular approach for assessing cereal aphid-parasitoid-
endosymbiont networks. *Scientific Reports*, 7(1), 1–12.
https://doi.org/10.1038/s41598-017-02226-w

936
937

Data Accessibility

The DROP database is freely accessible at Zenodo depository

(http://doi.org/10.5281/zenodo.4519656). New sequences have been deposited in

BOLD in datasets DS-LABS, and DS-AUSPTOID. [NOTE: the doi records will be update at

revision stage]

Author Contributions

The initial project idea was originated by C-HL, MLB, JH, MM, TS, JV, SG, and

PPAS. Molecular work was conducted by C-HL, SS, ML, AJ, and AD. BOLD and GenBank

data was harvested by TAE and C-HL. Figures were made by AL and C-HL. Laboratory and

field sample preparations were conducted by MTK, YC, TS, MM, SG, JV, EG, MG, XW,

KM, KMD, PA, NAP, MT, JJB, MP, FMJ, WDT, JSD, BW, OTL, PPAS, JL and AL. Taxonomic

concepts and interpretations were conducted by RRK, MLB, CH-L, PG, and SEM. DROP

database was built by JH and C-HL. All authors contributed to review and final revisions

of the manuscript, which was written primarily by C-HL, MLB and JH.

Figures:

Figure 1: An example of difficulties of molecular identification demonstrated on
Ganaspis xanthopoda and *G. brasiliensis*. Only two sequences (in bold text) can be
reliably used for identification and are included in DROP database. To select the
sequences, we searched the BINs associated with the organism's name "*Ganaspis*
xanthopoda" (green) or *"Ganaspis brasiliensis"* (purple) in BOLD. From each BIN, two
sequences from each species were selected to build a neighbor-joining tree (bottom axis
indicated % genetic divergence). There was a total of 6 BINs (gray boxes) in this
sequence complex. Of these, 4 BINs contained both species names and without
examination of vouchers identification would be impossible. In DROP, vouchers from
two sequences, **MG755073** and **MG755072**, were deposited in CNR-IPSP (Table S2),
examined by taxonomists and identify as *G. brasiliensis*. These two COI sequences can
now be used to reliably identify *G. brasiliensis*. For *G. xanthopoda,* there were no
available vouchers or reliable sequences that passed DROP standards to use for
identification*.* Species delimitation between *G. brasiliensis* and *G. xanthopoda* is
convoluted (see discussion), varies according to arbitrary % genetic divergence (gray
vertical lines), and needs future an integrative taxonomic revision.

Figure 2: Concept of a centralized, vetted, integrated database for _Drosophila_
Parasitoids (DROP) we developed. First, we provide a species and provisional species
catalog with correct taxonomy. Second, to provide a reliable genetic reference library,
genetic data (DNA sequences, genomes, transcriptomes, proteomes) link to a voucher
connected to the species catalog. Third, we link the two primary sources of data (field
surveys and laboratory experiments) by requiring a permanent deposition of vouchers
and sequences in order to be included in DROP.

980
981
982 **Tables:**

983 **Table 1:** List of species and provisional species included in DROP. For additional
984 taxonomic details, see DROP.
985

Superfamily	Family	Genus	Species_Name	Author
Chalcidoidea	Encyrtidae		_drop_Cha2_sp12_	
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_drop_ IR1_sp41_	Kimura
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_drop_BG1_sp42_	Kimura
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_zealandicus_	Ashmead 1904
Chalcidoidea	Pteromalidae		_drop_Pte69_sp11_	
Chalcidoidea	Pteromalidae	_Pachycrepoideus_	_vindemmiae_	(Rondani, 1875)
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_IR1_sp38_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_NG1_sp39_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_SK1_sp40_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drosophilae_	Ashmead 1887
Chalcidoidea	Pteromalidae	_Spalangia_	_erythromera_	Foerster 1850
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_dubia_	(Ashmead, 1896)
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_microptera_	(Lindeman, 1887)
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_nigricola_	Boucek

Chalcidoidea	Pteromalidae	*Trichomalopsis*	*sarcophagae*	(Gahan, 1914)
Chalcidoidea	Pteromalidae	*Vrestovia*	*brevior*	Boucek 1993
Chalcidoidea	Pteromalidae	*Vrestovia*	*fidenas*	(Walker, 1848)
Chalcidoidea	Pteromalidae		*drop_ PacAtl_sp46*	
Chalcidoidea	Pteromalidae		*drop_ PachyPort_sp45*	
Chalcidoidea			*drop_ CH_sp64*	
Cynipoidea	Figitidae	*Ganaspis*	*brasiliensis*	(Ihering, 1905)
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp51*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp52*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp53*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp1_sp67*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp2_sp68*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp50_sp66*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ IR1_sp25*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*drop_ IR2_sp26*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*drop_Gan1_sp1*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_TK1_sp27*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*hookeri*	Craword 1913
Cynipoidea	Figitidae	*Ganaspis*	*mahensis*	Kieffer 1911
Cynipoidea	Figitidae	*Ganaspis*	*mellipes*	(Say, 1826)
Cynipoidea	Figitidae	*Ganaspis*	*mundata*	Forster 1869
Cynipoidea	Figitidae	*Ganaspis*	*seticornis*	(Hellen, 1960)
Cynipoidea	Figitidae	*Ganaspis*	*tenuicornis*	Kieffer 1904
Cynipoidea	Figitidae	*Ganaspis*	*xanthopoda*	(Ashmead, 1896)
Cynipoidea	Figitidae	*Kleidotoma*	*bicolor*	(Giraud, 1860)
Cynipoidea	Figitidae	*Kleidotoma*	*dolichocera*	Thomson 1877
Cynipoidea	Figitidae	*Kleidotoma*	*drop_TK1_sp28*	Kimura
Cynipoidea	Figitidae	*Kleidotoma*	*filicornis*	(Cameron, 1889)
Cynipoidea	Figitidae	*Kleidotoma*	*icarus*	(Quinlan, 1964)
Cynipoidea	Figitidae	*Kleidotoma*	*psiloides*	Westwood 1833
Cynipoidea	Figitidae	*Kleidotoma*	*tetratoma*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptolamina*	*drop_Fig64_sp5*	
Cynipoidea	Figitidae	*Leptolamina*	*drop_Lmn_sp6*	
Cynipoidea	Figitidae	*Leptolamina*	*drop_TK1_sp29*	Kimura
Cynipoidea	Figitidae	*Leptolamina*	*gressitti*	Yoshimoto & Yasumatsu 1965
Cynipoidea	Figitidae	*Leptolamina*	*papuensis*	Yoshimoto 1963
Cynipoidea	Figitidae	*Leptolamina*	*ponapensis*	Yoshimoto 1962
Cynipoidea	Figitidae	*Leptolamina*	*seychellensis*	(Kieffer, 1911)

Cynipoidea	Figitidae	*Leptopilina*	*atraticeps*	(Kieffer, 1911)
Cynipoidea	Figitidae	*Leptopilina*	*australis*	(Belizin, 1966)
Cynipoidea	Figitidae	*Leptopilina*	*boulardi*	(Barbotin, Carton & Kelner-Pillault, 1979)
Cynipoidea	Figitidae	*Leptopilina*	*clavipes*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptopilina*	*cupulifera*	(Kieffer, 1916)
Cynipoidea	Figitidae	*Leptopilina*	*decemflagella*	Lue & Buffington 2017
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp54*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp55*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp56*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp57*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp58*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp59*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp60*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp61*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp62*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_BG1_sp34*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig059_sp4*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig124_sp2*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig58_sp3*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_IR1_sp30*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_NG1_sp33*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_SK1_sp35*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_STL_sp7*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_TK2_sp31*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_TK3_sp32*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*fimbriata*	(Kieffer, 1901)
Cynipoidea	Figitidae	*Leptopilina*	*freyae*	Allemand & Nordlander 2002
Cynipoidea	Figitidae	*Leptopilina*	*guineaensis*	Allemand & Nordlander 2002
Cynipoidea	Figitidae	*Leptopilina*	*heterotoma*	(Thomson, 1862)
Cynipoidea	Figitidae	*Leptopilina*	*japonica japonica*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*lasallei*	Buffington & Guerrieri 2020
Cynipoidea	Figitidae	*Leptopilina*	*leipsi*	Lue & Buffington 2018
Cynipoidea	Figitidae	*Leptopilina*	*lonchaeae*	(Cameron, 1912)
Cynipoidea	Figitidae	*Leptopilina*	*longipes*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptopilina*	*mahensis*	(Kieffer, 1911)
Cynipoidea	Figitidae	*Leptopilina*	*maia*	Lue & Buffington 2016
Cynipoidea	Figitidae	*Leptopilina*	*maria*	(Girault, 1930)
Cynipoidea	Figitidae	*Leptopilina*	*orientalis*	Allemand & Nordlander 2002

Cynipoidea	Figitidae	*Leptopilina*	*pacifica*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*rufipes*	(Cameron, 1908)
Cynipoidea	Figitidae	*Leptopilina*	*rugipunctata*	(Yoshimoto, 1962)
Cynipoidea	Figitidae	*Leptopilina*	*ryukyuensis*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*tokioensis*	Wachi & Kimura 2015
Cynipoidea	Figitidae	*Leptopilina*	*tsushimaensis*	Wachi & Kimura 2015
Cynipoidea	Figitidae	*Leptopilina*	*victoriae*	Nordlander 1980
Cynipoidea	Figitidae	*Rhoptromeris*	*heptoma*	(Hartig, 1840)
Cynipoidea	Figitidae	*Rhoptromeris*	*nigriventris*	Nordlander 1978
Cynipoidea	Figitidae	*Rhoptromeris*	*rufiventris*	(Giraud, 1860)
Cynipoidea	Figitidae	*Rhoptromeris*	*villosa*	(Hartig, 1840)
Cynipoidea	Figitidae		*drop_Lg500_sp43*	
Ichneumonoidea	Braconidae	*Alysia*	*drop_SP1_sp24*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*aotea*	Hughes & Woolcock 1976
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_SP1_sp15*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_TK1_sp13*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_TM1_sp14*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*minuta*	(Nees, 1811)
Ichneumonoidea	Braconidae	*Aphaereta*	*pallipes*	(Say, 1829)
Ichneumonoidea	Braconidae	*Aphaereta*	*scaptomyzae*	Fischer 1966
Ichneumonoidea	Braconidae	*Areotetes*	*striatiferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Areotetes*	*carinuliferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Asobara*	*ajbelli*	Berry 2007
Ichneumonoidea	Braconidae	*Asobara*	*albiclava*	Berry 2007
Ichneumonoidea	Braconidae	*Asobara*	*antipoda*	(Ashmead, 1900)
Ichneumonoidea	Braconidae	*Asobara*	*bactrocerae*	(Gahan, 1952)
Ichneumonoidea	Braconidae	*Asobara*	*brevicauda*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*citri*	(Fischer, 1963)
Ichneumonoidea	Braconidae	*Asobara*	*drop_KG1_sp16*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_NG1_sp17*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_SK2_sp20*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_SP1_sp18*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_Sp2_sp19*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*elongata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*gahani*	(Papp, 1969)
Ichneumonoidea	Braconidae	*Asobara*	*japonica*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Asobara*	*kenyaensis*	Peris-Felipo 2014
Ichneumonoidea	Braconidae	*Asobara*	*leveri*	(Nixon, 1939)
Ichneumonoidea	Braconidae	*Asobara*	*mesocauda*	van Achterberg & Guerrieri 2016

Ichneumonoidea	Braconidae	*Asobara*	*orientalis*	Viereck 1913
Ichneumonoidea	Braconidae	*Asobara*	*persimilis*	(Prince, 1976)
Ichneumonoidea	Braconidae	*Asobara*	*pleuralis*	(Ashmead, 1905)
Ichneumonoidea	Braconidae	*Asobara*	*rossica*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Asobara*	*rufescens*	(Fˆrster, 1862)
Ichneumonoidea	Braconidae	*Asobara*	*tabida*	(Nees, 1834)
Ichneumonoidea	Braconidae	*Asobara*	*triangulata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*turneri*	Peris-Felipo 2014
Ichneumonoidea	Braconidae	*Asobara*	*unicolorata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Aspilota*	*albertica*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*andyaustini*	Wharton 2002
Ichneumonoidea	Braconidae	*Aspilota*	*angusta*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*concolor*	Nees 1812
Ichneumonoidea	Braconidae	*Aspilota*	*parecur*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*villosa*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*barrattae*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*longworthi*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*philipi*	Berry 2007
Ichneumonoidea	Braconidae		*drop_Aso_sp8*	
Ichneumonoidea	Braconidae	*Opiognathus*	*pactus*	(Haliday, 1837)
Ichneumonoidea	Braconidae	*Opius*	*bellus*	Gahan 1930
Ichneumonoidea	Braconidae	*Opius*	*cinerariae*	Fischer
Ichneumonoidea	Braconidae	*Opius*	*crenuliferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Opius*	*monilipalpis*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Opius*	*ocreatus*	(Papp)
Ichneumonoidea	Braconidae	*Opius*	*pallipes*	Wesmael 1835
Ichneumonoidea	Braconidae	*Opius*	*pteridiophilus*	Wharton & Austin 1990
Ichneumonoidea	Braconidae	*Opius*	*pterus*	Wharton & Austin 1990
Ichneumonoidea	Braconidae	*Opius*	*trimaculatus*	Spinola
Ichneumonoidea	Braconidae	*Opius*	*youi*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Phaenocarpa*	*conspurcator*	(Haliday, 1838)
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drop_ IR1_sp22*	Kimura
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drop_TK1_sp21*	Kimura
Ichneumonoidea	Braconidae	*Phaenocarpa*	*tacita*	Stelfox 1941
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drosophilae*	(Fischer 1975)
Ichneumonoidea	Braconidae	*Tanycarpa*	*bicolor*	(Nees, 1814)
Ichneumonoidea	Braconidae	*Tanycarpa*	*chors*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Tanycarpa*	*drop_NG1_sp23*	Kimura
Ichneumonoidea	Braconidae	*Tanycarpa*	*punctata*	van Achterberg 1976

Ichneumonoidea	Braconidae		*drop_ Aly_sp47*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp48*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp49*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp50*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp63*	
Ichneumonoidea	Braconidae		*drop_ Aso_sp69*	
Diaprioidea	Diapriidae	*Trichopria*	anastrephae	Costa Lima 1940
Diaprioidea	Diapriidae	*Trichopria*	*drop_ BG1_sp37*	Kimura
Diaprioidea	Diapriidae	*Trichopria*	*drop_ Dia70_sp65*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_ Tri_sp44*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_Bdia_sp10*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_Dia127_sp9*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_TK1_sp36*	Kimura
Diaprioidea	Diapriidae	*Trichopria*	drosophilae	(Kieffer, 1912)
Diaprioidea	Diapriidae	*Trichopria*	modesta	(Ratzeburg, 1848)

Table 2.1: *Drosophila* parasitoid whole-genome sequences included in DROP. For additional details, see DROP.

Genus	Species_Name	Species_id	Genome_id	Voucher_id	GenBank_id
Ganaspis	brasiliensis	19	8	868	GCA_009823575.1
Ganaspis	brasiliensis	19	16	872	SRX8882993
Ganaspis	brasiliensis	19	17	871	SRX8882992
Ganaspis	drop_ Gsp1_sp67	182	15	873	SRX8882994
Ganaspis	drop_ Gsp2_sp68	183	14	874	SRX8882995
Ganaspis	drop_ Gsp50_sp66	181	9	869	GCA_011057455.1
Leptolamina	ponapensis	48	13	875	SRX8883008
Leptopilina	boulardi	4	5	865	GCA_011634795.1
Leptopilina	boulardi	4	6	866	GCA_003121605.1
Leptopilina	boulardi	4	12	876	SRX8883009
Leptopilina	clavipes	5	7	867	GCA_001855655.1
Leptopilina	heterotoma	6	1	861	GCA_010016045.1
Leptopilina	heterotoma	6	2	862	GCA_009602685.1
Leptopilina	heterotoma	6	3	863	GCA_009026005.1
Leptopilina	heterotoma	6	4	864	GCA_009025955.1
Leptopilina	japonica japonica	13	11	877	SRX8883011

990
991
992

993 **Table 2.2:** *Drosophila* parasitoid transcriptome data included in DROP.
994

Genus	Species_Name	Strain_id	Transcriptome_id	Voucher_id	Genbank_id
Leptopilina	*boulardi*	126	2	858	2183568
Leptopilina	*boulardi*	127	3	859	2183567
Leptopilina	*boulardi*	151	8	882	15642271
Leptopilina	*boulardi*	151	9	883	15642270
Leptopilina	*heterotoma*	147	1	857	2183569
Leptopilina	*heterotoma*	152	5	884	2046288
Leptopilina	*heterotoma*	61	6	880	11581553
Leptopilina	*heterotoma*	60	7	881	11662592
Leptopilina	*boulardi*	11	10	908	GAJA00000000.1
Leptopilina	*heterotoma*	14	11	909	GAJC00000000.1
Ganaspis	*hookeri*	7	12	910	GAIW00000000.1

996 **Table 2.3:** *Drosophila* parasitoid proteomes data included in DROP
997

Genus	Species_Name	Strain_id	Proteomes_id	Voucher_id	Assession_id
Leptopilina	*heterotoma*	152	1	885	PRIDE: PXD005639
Leptopilina	*heterotoma*	61	2	886	PRIDE: PXD005632
Leptopilina	*boulardi*	27	3	911	Upon request to julien.varaldi@univ-lyon1.fr
Leptopilina	*boulardi*	11	4	912	PRIDE: PDX023836
Leptopilina	*heterotoma*	14	5	913	PRIDE: PDX023824
Ganaspis	*hookeri*	7	6	914	PRIDE: PDX023825