DROP: Molecular voucher database for identification of *Drosophila* parasitoids

Resource article

Word count: 8237 excluding references

Authors

Chia-Hua Lue (C-HL), (0000-0002-5245-603X), chiachia926@gmail.com, corresponding

author

Matthew L. Buffington (MLB)

Sonja Scheffer (SS)

Matthew Lewis (ML)

Tyler A. Elliott (TAE)

Amelia R. I. Lindsey (AL)

Amy Driskell (AD), (0000-0001-8401-7923)

Anna Jandova (AJ)

Masahito T. Kimura (MTK)

Yves Carton (YC)

Robert R. Kula (RRK)

Todd A. Schlenke (TAS)

Mariana Mateos (MM), (0000-0001-5738-0145)

Shubha Govind (SG), (0000-0002-6436-639X)

Julien Varaldi (JV)

Emilio Guerrieri (EG), (0000-0002-0583-4667)

Massimo Giorgini (MG), (0000-0001-8670-0945)

Xingeng Wang (XW)

Kim Hoelmer (KH)

Kent M. Daane (KMD**)**

Paul K. Abram (PKA)

Nicholas A. Pardikes (NAP), (0000-0002-9175-4494)

Joel J. Brown (JJB), (0000-0002-3608-6745)

Melanie Thierry (MT)

Marylène Poirié (MP)

Paul Goldstein (PG), (0000-0002-1443-7030)

Scott E. Miller (SEM), (0000-0002-4138-1378)

W. Daniel Tracey (WDT)

Jeremy S. Davis (JSD), (0000-0002-5214-161X)

Francis M. Jiggins (FMJ)

Bregje Wertheim (BW)

Owen T. Lewis (OTL)

Jeff Leips (JL)

Phillip P. A. Staniczenko (PPAS)

Jan Hrcek (JH), (0000-0003-0711-6447), janhrcek@gmail.com

Affiliations

(C-HL, AJ, NAP, JJB, MT, JH) Biology Centre of the Czech Academy of Sciences, Institute

of Entomology, Branisovska 31, 37005 Ceske Budejovice, Czech Republic

(C-HL, PPAS) Department of Biology, Brooklyn College, City University of New York

(CUNY), 2900 Bedford Ave, Brooklyn, NY11210, USA

(MLB, SS, ML, RRK, PG) Systematic Entomology Laboratory, ARS/USDA c/o Smithsonian

Institution, National Museum of Natural History, 10th& Constitution Ave, NW,

Washington DC 20560, USA

(TAE) Centre for Biodiversity Genomics, 50 Stone Road East, University of Guelph,

Guelph, Ontario, N1G2W1, Canada

(AL) Department of Entomology, University of Minnesota, Saint Paul MN 55108.

(AD) Laboratories of Analytical Biology, Smithsonian Institution, National Museum of

Natural History, 10th & Constitution Ave, NW, Washington DC 20560, USA

(MTK) Hokkaido University Museum, Sapporo, Hokkaido 060-0810, Japan

(YC) "Évolution, Génomes, Comportement, Écologie", CNRS et Université Paris-Saclay

(TAS) Department of Entomology at the University of Arizona, Forbes 410, PO BOX

210036, Tucson, AZ 85721-0036.

(MM) Wildlife and Fisheries Sciences Department, Texas A&M University

(SG) The Graduate Center of the City University of New York, 365 Fifth Avenue, New

York, NY10016, USA

(JV) Université de Lyon, Université Lyon 1, CNRS, Laboratoire de Biométrie et Biologie

Evolutive UMR 5558, F-69622 Villeurbanne, France

(EG, MG) CNR- Institute for Sustainable Plant Protection (CNR-IPSP), National Research

Council of Italy, Piazzale E. Fermi 1, 80055 Portici, Italy.

(XW, KH) United States Department of Agriculture, Agricultural Research Services,

Beneficial Insects Introduction Research Unit. 501 S. Chapel St., Newark, DE 19713, USA

(KMD) Department of Environmental Science, Policy and Management. University of

California, Berkeley. Mulford Hall, 130 Hilgard Way, Berkeley, CA 94720.

(PKA) Agriculture and Agri-Food Canada, Agassiz Research and Development Centre,

6947 Hwy #7, Agassiz, V0M 1A0, British Columbia, Canada

(JJB, MT, JH) University of South Bohemia, Faculty of Science, Branisovska 31, 37005,

Czech Republic

(MP) Université "Côte d'Azur", INRAE, CNRS. and Evolution and Specificity of

Multitrophic Interactions (ESIM) Sophia Agrobiotech Institute, 400 Route des Chappes,

BP 167, 06903 Sophia Antipolis, France

(SEM) Smithsonian Institution, National Museum of Natural History, 10th & Constitution

Ave, NW, Washington DC 20560, USA

(WDT, JSD) Department of Biology, Indiana University Bloomington. 702 N. Walnut

Grove, Bloomington, IN47405

(WDT) Gill Center for Biomolecular Science, Indiana University Bloomington. 702 N.

Walnut Grove, Bloomington, IN47405

(JSD) Biology Department, University of Kentucky, 101 T. H. Morgan Building, Lexington,

KY, 40506

(FMJ) Department of Genetics, University of Cambridge, Downing Street, Cambridge CB2

3EH UK

(BW) Groningen Institute for Evolutionary Life Sciences, University of Groningen,

Nijenborgh 9, 9747 AG Groningen, the Netherlands

(OTL) Department of Zoology, University of Oxford. 11a Mansfield Road, Oxford OX1

3SZ, UK.

(JL) Department of Biological Sciences, University of Maryland Baltimore County. 1000

Hilltop circle, Baltimore, MD, 21250

Abstract

Molecular identification is increasingly used to speed up biodiversity surveys and

laboratory experiments. However, many groups of organisms cannot be reliably

identified using standard databases such as GenBank or BOLD due to lack of sequenced

voucher specimens identified by experts. Sometimes a large number of sequences are

available, but with too many errors to allow identification. Here we address this

problem for parasitoids of *Drosophila* by introducing a curated open-access molecular

reference database, DROP (*Drosophila* parasitoids). Identifying *Drosophila* parasitoids is

challenging and poses a major impediment to realize the full potential of this model

system in studies ranging from molecular mechanisms to food webs, and in biological

control of *Drosophila suzukii*. In DROP (http://doi.org/10.5281/zenodo.4519656),

genetic data are linked to voucher specimens and, where possible, the voucher

specimens are identified by taxonomists and vetted through direct comparison with

primary type material. To initiate DROP, we curated 154 laboratory strains, 856

vouchers, 554 DNA sequences, 16 genomes, 14 transcriptomes, and 6 proteomes drawn

from a total of 183 operational taxonomic units (OTUs): 114 described *Drosophila*

parasitoid species and 69 provisional species. We found species richness of *Drosophila*

parasitoids to be heavily underestimated and provide an updated taxonomic catalogue

for the community. DROP offers accurate molecular identification and improves cross-

referencing between individual studies that we hope will catalyze research on this

diverse and fascinating model system. Our effort should also serve as an example for

researchers facing similar molecular identification problems in other groups of

organisms.

Key Words

Biodiversity, DNA sequences, Genomes, Integrative taxonomy, Molecular diagnostics,

Biological control

Introduction

Building a knowledge base that encompasses ecology, evolution, genetics, and biological control is contingent on reliable taxonomic identifications. Molecular identification is commonly used in groups of organisms with cryptic species that are difficult to identify morphologically (Fagan-Jeffries et al., 2018; Miller et al., 2016; Novotny & Miller, 2014), for the molecular detection of species interactions (Baker et al., 2016; Condon et al., 2014; Gariepy et al., 2019; Hrček & Godfray, 2015; Hrcek et al., 2011), and for identification of species from environmental DNA samples (Shokralla et al., 2012). The accuracy of molecular identification, however, depends on the accuracy of identifications associated with sequences databased in existing online depositories (Fontes et al., 2021). The foundations of that accuracy are the voucher specimens which were sequenced and the collaboration of a taxonomic authority in the deposition of the sequence data.

GenBank serves as the most widely used sequence depository; however, deposition of sequences in GenBank, which is required by most peer-reviewed journals, does not require deposition of associated vouchers. The Barcode of Life Data System database (BOLD) (Ratnasingham & Hebert, 2007) explicitly aims to provide a framework for identifying specimens using single-locus DNA sequences (Hebert et al., 2003; Smith et al., 2005), and while these are associated with vouchers and metadata, the curation of these data is not consistently maintained by those submitting material. A recent study by Pentinsaari et al. (2020) showed misidentification in both databases caused by missteps in the protocols from query sequences to final determination.

Although the BOLD database function "BOLD-IDS" allows considerable database

curation (e.g. flagging of misidentified/contaminated records), it also automatically

includes sequences from GenBank, and may perpetuate the shortcomings previously

mentioned since these cannot be curated from within BOLD. As such, the quality of

sequences and the reliability of identifications obtained from BOLD-IDS can vary, and

depends on the curation by taxonomists focusing on individual taxa (Meiklejohn et al.,

2019). BOLD-IDS works well for taxa where qualified taxonomists have been involved

with assuring data quality; some insect examples include beetles (Hendrich et al., 2015),

butterflies (Escalante et al., 2010), geometrid moths (Hausmann et al., 2011, 2016;

Miller et al., 2016), true bugs (Raupach et al., 2014), and microgastrine wasps (Smith et

al., 2013).

Unfortunately, this is not the case of parasitoids (Insecta: Hymenoptera) of

Drosophila flies (Insecta: Drosophilidae). There are vast numbers of *Drosophila*

parasitoid sequences readily available in GenBank and BOLD, as these parasitoids and

their hosts are important model organisms in biology. As of this writing, there are

88,666 nucleotide sequences deposited in GenBank for *Leptopilina heterotoma*

(Thomson) and *L. boulardi* (Barbotin, Carton & Kelner-Pillault) (Hymenoptera: Figitidae)

alone. However, less than 1 % of the identifications associated with these sequences

have been confirmed by taxonomists or are associated with voucher specimens

deposited in museum collections. With sequencing shifting from individual genes to

genomes we risk that the identification problems will soon apply to whole genomes.

Drosophila and their parasitoids

The phylogenetic and subgeneric structure within *Drosophila* and related genera is

not yet fully resolved (O'Grady & DeSalle, 2018). Various subgenera, including

Scaptomyza, *Zaprionus*, *Lordiphosa* and *Samoaia*, have been treated as both genera and

subgenera, and researchers have yet to achieve consensus on these various hypotheses

(O'Grady & DeSalle, 2018; Remsen & O'Grady, 2002; Yassin, 2013; Yassin & David,

2010). Species in *Drosophila* subgenera and genera closely related to *Drosophila*

commonly share niche space and natural histories and, as a result, are often attacked by

overlapping or identical groups of parasitoids. For instance, the invasive African fig fly,

Zaprionus indianus Gupta is attacked by *Pachycrepoideus vindemiae* (Rondani, 1875)

and *Leptopilina boulardi* (Pfeiffer et al., 2019; Santos et al., 2016), all of which have been

recorded from *Drosophila*. Therefore, we also include these groups within the contents

of DROP.

Parasitoids of *Drosophila* belong to four superfamilies of Hymenoptera

(Chalcidoidea, Cynipoidea, Ichneumonoidea, Diaprioidea) which evolved parasitism of

Drosophila flies independently. All the parasitoids known to attack *Drosophila* are

solitary and attack either the larval or pupal stage; in both cases, they emerge from the

fly's puparium (Carton et al., 1986; Prévost, 2009). The known *Drosophila* larval

parasitoids belong to two families (Table 1), Braconidae (including the genera *Asobara,*

Aphaereta, Phaenocarpa, Tanycarpa, Aspilota, Opius) and Figitidae (*Leptopilina,*

Ganaspis, Leptolamina, Kleidotoma); all are koinobionts that allow the host to continue

development while the parasitoid grows within it. The known *Drosophila* pupal

parasitoids belong to three other families (Table 1), Diapriidae (*Trichopria, Spilomicrus*),

Pteromalidae (*Pachycrepoideus, Spalangia, Trichomalopsis, Toxomorpha*) and Encytidae

(*Tachinaephagus*); they are all idiobionts that terminate host development immediately.

Host-specificity across the *Drosophila* parasitoids is poorly characterized—while some

can parasitize other families of Diptera (e.g., *Aphaereta aotea*) (Hughes & Woolcock,

1976), most are thought to be limited to *Drosophila* hosts.

There are around 4000 described species of Drosophilidae, and *Drosophila* contains

more than a third of the family's described species (O'Grady & DeSalle, 2018). By

contrast, although parasitic wasps are generally a species-rich group (Dolphin & Quicke,

2001; Quicke, 2015), the most recent catalogue of parasitoid species that attack

Drosophila lists only 50 described species (Carton et al., 1986). This disparity suggests

that the diversity of parasitic wasps attacking *Drosophila* is severely underestimated, an

assertion supported by the results presented here. This is largely a consequence of the

challenging nature of parasitoid taxonomy, in which morphological identification is

intractable for many species, and the fact that taxonomic specialists are greatly

outnumbered by the species they study.

Currently, only a few biological study systems have been characterized in

sufficient breadth and depth to allow researchers to connect various levels of biological

organization, from molecular mechanisms to food webs of interacting species.

Parasitoids of *Drosophila* represent one such system (Prévost, 2009). Moreover, the

practical feasibility of rearing parasitoids of *Drosophila* under laboratory conditions has

led to a number of fundamental discoveries in ecology (Carton et al., 1991; Terry et al.,

2021), evolution (Kraaijeveld & Godfray, 1997), immunology (Kim-Jo et al., 2019; Nappi

& Carton, 2001; Schlenke et al., 2007), physiology (Melk & Govind, 1999), symbiosis (Xie

et al., 2011, 2015), behavioral science (Lefèvre et al., 2012) and other fields. In contrast

to this large body of laboratory studies, basic natural history of *Drosophila* parasitoids,

especially their species richness is little known (Kimura & Mitsui, 2020; Lue et al., 2018).

Addressing this knowledge gap is especially pressing given current efforts to use

parasitoids in biological control efforts, such as those of the invasive pest spotted wing

Drosophila, *Drosophila suzukii* (Abram et al., 2020; Daane et al., 2016; Giorgini et al.,

2019; Wang et al., 2020 a&b).

Properly executed molecular identification has the potential to be much more

efficient for the majority of researchers, and many laboratory strains are commonly

identified using DNA sequences alone. While it is practical for researchers to assign

species names based on a match to sequence records in genetic databases, this practice

often causes a cascade of inaccuracies. To illustrate the extent of the problem, we

present the example of *Ganaspis,* a genus of parasitoids commonly used in laboratories

that includes both superficially indistinguishable species with highly divergent

sequences that are often treated as conspecific, as well as specimens with identical

sequences identified under different names (Figure 1).

Aims

To address these issues, we introduce a newly curated molecular reference database

for <u>Drosophila</u> parasitoids —DROP— in which sequences are either linked to voucher

specimens identified by taxonomists or have a traceable provenance (Figure 2). The first

aim of DROP is to provide a reliable DNA sequence library for molecular identification of

Drosophila parasitoids that enables cross-referencing of original taxonomic concepts

with those of subsequent studies. We pay special attention to live parasitoid strains

which are available for future experiments. The second aim is to standardize and

expedite the linkage between specimens and available sequence data; we place a

premium on museum vouchers as they allow for repeatable scientific research. In DROP,

this goal is facilitated through a consolidated digital infrastructure of data associated

with laboratory strains, offering the opportunity for researchers to re-examine past

experimental results in a permanent context. The third aim is to provide an up-to-date

catalogue of the diversity of *Drosophila* parasitoids as a foundation for advancing the

understanding of their taxonomy. Finally, the fourth aim of DROP is for our collaborative

effort to serve as an inspiration to communities of researchers studying other groups of

organisms who are experiencing difficulties with the reliability of molecular reference

databases.

Materials and Methods

Data sources

To assemble the DROP database, we targeted 20 wasp genera that potentially

parasitize frugivorous *Drosophila* species. We compiled DNA sequence and voucher data

from four sources: 1) museum collections, 2) publications, for which we selected the

reference with taxonomist or parasitoid biologists as coauthors to ensure reliable

species identity, 3) molecular biodiversity inventories publicly available in BOLD and

GenBank, for which we managed to secure inspection of the vouchers by taxonomists,

and 4) a sequencing and taxonomic inventory of laboratory strains we conducted.

We first gathered species information into a catalogue of *Drosophila* parasitoid

species (Table 1) from 216 references (see DROP database reference table) and 36

institutes (Table S2). To ensure reliable names for nominal species (sequences identified

by a species name) in our database, we confirmed their taxonomic validity using the

Ichneumonoidea 2015 digital catalogue (Yu et al., 2016) and Hymenoptera Online (HOL;

http://hol.osu.edu/), both of which are curated by taxonomic experts. To obtain reliable

molecular identification data, we harvested 8,298 DNA sequences from GenBank and

BOLD (all compiled in BOLD as DS-DROPAR dataset dx.doi.org/10.5883/DS-DROPAR). As

of writing, these sequences represented 445 Barcode Index Numbers (BINs – a form of

dynamic provisionary taxa in BOLD, more detail in Ratnasingham & Hebert 2013), and

211 named taxa.

The majority of the harvested sequences were Braconidae (6690), Diapriidae

(967), Figitidae (622), and Pteromalidae (19). Because of the concerns with generic

databases (noted above and in Figure 1 and Table S1), we assembled a list of sequences

with valid species names that could either be traced back to vouchers examined by

taxonomists or were referred to directly in publications authored by a recognized expert

in the relevant taxon group. We then cross-checked species names with their

corresponding BINs in BOLD and flagged potential conflicts between species names and

BINs (Table S1).

A core goal of DROP besides that of a tool for biodiversity research is to function as a

platform that accommodates *Drosophila* parasitoids kept in laboratory strains (for

experimental work) or cultures in quarantine facilities (for biological control

applications). So far, there has been a lack of a coherent and reliable means of verifying

identification of species kept in laboratory settings, which can be a serious problem.

Since lab cultures are routinely contaminated by neighboring cultures (e.g., through

escapees), one species may be displaced by another even under a vigilant eye.

For lab and quarantine lines in DROP, we deposited DNA extractions and vouchers in

the National Insect Collection, National Museum of Natural History, Smithsonian

Institution (USNM; Washington, DC, USA). During their initial assembly of DROP,

laboratory OTUs (operational taxonomic unit) were designated by their strain name;

most laboratory strains can be associated with provisional species, but some cannot yet

be assigned. Three females and three males of each strain were dry-mounted and

individually assigned a USNMENT 'QR code' specimen label as representative vouchers.

For each molecular voucher, three legs from a female wasp were removed for DNA

extraction and sequencing (Supplementary Methods for details), and the rest of the

body was assigned a USNMENT specimen label and preserved for morphological

identification. Both DNA extraction and vouchers were entered into the database and

uploaded to BOLD (DROP project: DS-LABS dx.doi.org/10.5883/DS-LABS) with an

associated GenBank ID.

Where possible, we identified OTU strains using a combination of morphological and

sequence data, and characterized provisional species or species clusters using neighbor-

joining trees (Figure S1) based on the COI gene sequences (Supplemental material). For

establishing BIN limits in the context of DROP, we have adopted an initial percent cutoff

at 2%. We acknowledge that 2% genetic diverge cutoffs (or BINs) are unlikely to work

well across range of widely distributed species (Lin et al., 2015). But as Ratnasingham &

Hebert (2013) pointed out, 2% is a good starting point for many taxa, also it may need to

be adjusted as more samples are acquired and compared. Note that we use the term

"OTU" as a general and neutral designation encompassing described species, provisional

species, undescribed species, and cryptic species.

Drosophila parasitoid database—DROP

To compile the above information, we built a simple Structured Query Language

(SQL) database in sqlite3 format using SQLiteStudio (step by step user instruction in

supplemental material). Sqlite3 is a cross-platform format which can be also be opened

using a number of other programs. There are eight linked tables in the database —

species, strain, voucher, sequence, genome, transcriptome, proteome and reference —

along with additional tables for linking these to reference table (Figure S2). The

database incorporates all sample fields used by BOLD for compatibility and includes a

number of new fields to accommodate a catalogue of *Drosophila* parasitoid species,

laboratory strain information, and links from the DROP database to BOLD and GenBank

records.

DROP is available on Zenodo (http://doi.org/10.5281/zenodo.4519656) for

permanent deposition and version control. In addition to the main database, the

Zenodo repository includes additional files to facilitate easy use of the database. These

files include: 1) the reference database in comma-separated text (.csv) and FASTA

format ready to be used for molecular identification; 2) a species catalogue with

taxonomic information; and 3) a list of laboratory strains with confirmed molecular

vouchers. DROP will be continued to be curated and maintained by C-HL at the Zenodo

repository and sequences generated in the future will also be deposited in BOLD (DROP

project). If the curator changes, this will be announced in the README.md file in Zenodo

repository. As the database relies on vouchers, we will aim for it to be continued to be

maintained by taxonomists with direct access to museums.

Species, provisional species, and OTU designations

In addition to the inherent value of a formal taxonomic name, a reliable provisional

taxon label can also be used for exchanging scientific information and conveying

experimental results among researchers (Schindel & Miller, 2010). Based on the amount

of sequence divergence between described species, we observed what appears to be a

significant number of provisional OTUs in the initial dataset we compiled. Furthermore,

among the data linked to a valid species name, some of these provisional OTUs are

actively being used in research and have sequences available to the public. We

therefore provide a list of provisional species (potential new species) with their

molecular vouchers.

We use the following designation format for OTUs that refer to a provisional species:

"Drop_strainX_sp.1" or, when no other information is known, "DROP_sp.1". Where

possible, these OTUs are linked to a voucher USNM specimen label number. If the genus

of the OTU is known, the "Drop_Leptopilina_sp.1" format is followed. These

designations can facilitate species identification as well as discovery and description of

new species without compromising the existing taxonomy of the described OTUs in

question. As more complete species descriptions become available, this provisional

species framework can be updated while keeping the link to previous provisional species

name through deposited vouchers.

Results

Overview of DROP

We catalogued 183 OTUs in the DROP database with 114 described species of

Drosophila parasitoids and 69 provisional species (Table 1). In total, we documented 154

laboratory strains (Table S3), and 853 vouchers from 36 institutions (Table S2). Among

the described species, 98 have voucher information, of which 61 are traceable to type

specimens, including 45 to holotypes (i.e., specimen used to root a name to the

taxonomic author's concept of the species). *Leptopilina* is represented by the highest

number of species with 45 OTUs, followed by *Asobara* with 26 OTUs. Within the 154

catalogued lab strains, 86 were actively being used in ongoing research (i.e*.,* a live strain

being cultivated). These strains represent 39 OTUs: 11 described species and 28

provisional species (Table S3, Figure S1).

Molecular Vouchers

So far, DROP includes 545 DNA sequences and links to 16 genomes (Table S4), 14

transcriptomes (Table S5), and 6 proteomes (Table S6). From the total of 8298 DNA

sequences (BOLD dataset: DS-DROPAR) collected from public databases, only 322

sequences (less than 4% of available sequences) satisfied the reliability criteria we

imposed for molecular vouchers to be included in DROP (see Materials and Methods).

The DS-DROPAR dataset dx.doi.org/10.5883/DS-DROPAR initially referred to 211 taxon

names, but only 52 names were valid, linked to vouchers, or linked to a publication with

evidence that the specimens had been identified by taxonomists. The remaining 223 of

545 DROP DNA sequences were generated by DROP project (datasets: DS-LABS

dx.doi.org/10.5883/DS-LABS and DS-AUSPTOID dx.doi.org/10.5883/DS-AUSPTOID) and

came from 121 OTUs (101 lab strains and 12 provisional species).

The DROP database is largely made up of standard barcode COI sequences (349

sequences), which includes 77 OTUs: 43 described species and 33 provisional species.

We aimed to supplement COI with secondary markers (28SD2, 18S, ITS2) when possible,

resulting in an additional 120 sequences from 26 OTUs: 15 described species and 11

provisional species. There are currently 19 OTUs that have sequences from more than

one genetic marker.

Species Delimitation in Laboratory Strains

We used 298 COI sequences to resolve the identification of each laboratory

strain, and where possible, indicated potential species clusters (Fig. S1 and Table S3).

Using a fixed 2% divergence cutoff, a total of 31 lab strain OTUs were assignable to a

valid species name, and the remaining 70 strain OTUs were assigned to a provisional

species. The taxonomic status of several of these provisional species is also being

investigated using an integrative taxonomic approach involving morphological

identification, genomic data, or other genetic data.

Discussion

In this paper, we introduce and describe a free and open-access database for the

reliable molecular identification of *Drosophila* parasitoids. The guiding principle of DROP

is data credibility, based on the prerequisite that genetic data are explicitly associated

with voucher specimens and taxonomic concepts of the original authors (Troudet et al.,

2018). When incorporating information from public genetic databases, we included only

sequences that have passed our filtering protocol. This protocol ensures each entry is

associated with a valid scientific name, provisional name, or consistently applied OTU

designation that can be used to integrate genetic and organismal data from

independent studies.

The following discussion expands on the utility of DROP and how we hope it will

benefit molecular species identification, connect research from various disciplines,

support biological control applications, and serve as a long-term molecular voucher

repository and clearinghouse for vetted data. We also provide specific guidance for

users how best to refer to DROP in their publications to allow cross-linking between

studies.

Molecular (mis-)identification

We observe that 17% of the described *Drosophila* parasitoid OTUs in BOLD and

GenBank (dataset: DS-DROPAR) are associated with more than one BIN; these are

examples of BIN-ID conflict. Roughly half of these OTUs are used as lab strains. This

latter observation is disturbing, because it demonstrates that the criteria used to

differentiate and reference species in active research programs are clouded. For

example, BIN-ID conflicts were observed in the *Drosophila* parasitoids *Ganaspis*

brasiliensis (Ihering) and *Asobara japonica* Belokobylskij (Table S1), both of which are in

active use in numerous research programs (e.g. Moreau et al., 2009; Nomano et al.,

2017; Reumer et al., 2012; Wang et al., 2020a & 2021) as well as in biological control

efforts against the invasive *D. suzukii* (e.g. Abram et al., 2020; Daane et al., 2016;

Giorgini et al., 2019). All the BINs from *G. brasiliensis* carry the name *G. xanthopoda*

(Figure 1). In such instances, assigning an identification by matching specimens to

barcode records in the genetic database is problematic, as two names are applied to the

same BIN. If sequences comprising the BIN are not linked to a voucher that can be

examined, teasing apart the two names and how they are applied is impossible.

Applying explicit, consistent criteria for species determination ensures that

experimental results can be reliably repeated, and that any potentially novel

observations will not be explained away as artifacts of identification. DROP addresses

these concerns by linking reliable reference sequences and vouchers for *G. brasiliensis*

(Figure 1) between different studies: one with reference to the morphological

description (Buffington & Forshage, 2016) and the other with reference to the genome

(using voucher specimens from the morphological study; Blaimer et al., 2020).

We were not able to resolve all conflicts between BIN and species identity, for one

or more of the following three reasons: First, many records lack reliably identified

vouchers and have often been themselves used for molecular identification,

proliferating errors. Second, in some cases, it is not possible to verify whether the

genetic differences among BINs represent different species or simply intraspecific

genetic variation (Bergsten et al., 2012), because BINs themselves are not a species

concept. The only solution to this problem is to derive original sequence data from type

specimens (which is often either impractical or impossible for a number of technical

reasons), or from specimens whose conspecificity with the types has been corroborated.

Since species boundaries are always subject to testing, additional specimens from

multiple collecting events (ideally representing different seasons and geographic

regions) may help provide the additional data to circumscribe a given species' limits. The

third difficulty in resolving BIN-ID conflict derives from the data themselves: Although

the mitochondrial COI gene is the locus most frequently chosen for identification of

insects and other animals, its effectiveness varies among insect groups (Brower &

DeSalle, 2002; Gompert et al., 2008; Lin & Danforth, 2004). In part, this derives from

gene-tree/species-tree conflict as a function of mitochondrial DNA introgression

(Gompert et al., 2008; Klopfstein et al., 2016), parthenogenesis (Reumer et al., 2012),

and/or *Wolbachia* infection (Ferrer-Suay et al., 2018; Wachi et al., 2015; Xiao et al.,

2012), any of which may lead to complications in species delimitation using

mitochondrial loci. Ideally, studies should apply multiple loci, genomes, and comparative

taxonomic data to clarify species boundaries. As *Drosophila* parasitoids are often

maintained in laboratory cultures, it is also possible to use mating experiments to

explore species boundaries under the paradigm of the biological species concept

(Seehausen et al., 2020).

DROP as a taxonomic tool

DROP offers an empirical platform for species discovery and a useful tool for

taxonomic research. The fact that the number of BINs reported here exceeds the

number of described species (Table S1, Figure S3) highlights the need for taxonomic

work. But such work cannot proceed on the basis of BINs or barcodes, but requires

integrative taxonomic approach employing a combination of molecular and

morphological data. Describing new species on the sole basis of a barcode or BIN,

without the benefit of independent character data, should, in general, be avoided

(Meier et al., 2021). It risks creating nomenclatural synonymy if it is later determined

that a sequence can be attributed to a specimen that bears a valid, available name.

Moreover, BINs are based on distance analyses which, by definition, are incompatible

with diagnoses per se (Ferguson, 2002; Prendini et al., 2002; Goldstein & DeSalle, 2011).

Therefore, in taxonomic treatments, it is critical to clarify the range of applicability of a

given BIN and its overlap with a taxonomic name (see example in Figure 1). DROP allows

cross-linking between studies and therefore provides researchers with valuable tools for

taxonomic revisions, including the means of discovery, corroboration, and description of

new species.

How to use DROP to ensure cross-linking between studies and reliable molecular

identification?

Public genetic databases have adopted a longstanding convention in treating

undetermined OTUs and sequences, referring to provisional species with numbers, as

for example "sp. 1", and these are rarely linked to vouchers. For OTUs designated as

provisional species, DROP enables cross-indexing of specimens, sequences and

references between any studies (ecological, taxonomic, evolutionary, genetic, etc). The

best way to ensure cross-linking is depositing a voucher in DROP, together with a

sequence or genome from the same individual (or individual from the same strain or

series). For example, one can write:

Provisional species "drop_Gan1_sp.1" refers to voucher USNMENT01557320

deposited in the USNM, Washington DC, COI sequence (DROP sequence_id: 2, BOLD

Process ID: DROP143-21), 28SD1 sequence (DROP sequence_id: 289), and 28SD2

sequence (DROP sequence_id: 303).

Similarly, laboratory strains can be reported in the same way, just adding the

DROP lab strain_id. It is important to periodically recheck identification of laboratory

strains as cultures are easily cross-contaminated, and deposit vouchers of laboratory

strains associated with experiments to DROP. In the future, when e.g.

"drop_Gan1_sp.1" is described as a new species with a formal specific epithet, DROP

curator will update the species status and holotype information while keeping this

provisional species name as an informal "synonym."

A weaker and thus much less preferred way of cross-linking is to state in the

study that the identification of organisms was performed based on molecular

identification match of a sequence to DROP sequences. This is the only available option

for environmental DNA studies. For example, one can write:

Provisional species "drop_Gan1_sp.1" was identified based on 99.9% blast match

of COI to DROP sequence_id: 2 (BOLD Process ID: DROP143-21).

DROP deposition in Zenodo allows referencing of DROP either through general

doi (the doi we use throughout this paper), which takes the user always to the latest

database version, or through a doi specific to DROP version. When referencing DROP

please primarily cite this paper, but for reproducibility it is also good practice to include

doi of the specific DROP version used.

There are two basic ways of molecular identification which should ideally be

used in combination: sequence matching (blast), and tree-building methods which

investigate membership to a cluster. Further, there are a number of decisions to be

made with each method, concerning locus (or loci) and thresholds. DROP leaves these

decisions up to the users, only provides raw sequences or links to them. Practically, the

choice of loci is currently mostly limited to COI, but in the future it is likely that

molecular identifications will be based on multiple loci or whole genomes. Over time we

will also get a better idea about what thresholds are more appropriate than a fixed 2%

cut off. For rarer parasitoid genera which attack also other hosts besides *Drosophila*

(e.g. *Opius*, or *Spalangia* wasps) we suggest caution in the identification using only

DROP sequences as DROP does not include all sequences from these genera, but just

from species which are already known to attack *Drosophila*.

From molecular mechanisms to ecosystem structure

The use of molecular tools in insect biodiversity studies has gradually expanded from

barcoding single individuals to metabarcoding large environmental samples

representing entire food webs (Jeffs et al., 2020; Littlefair et al., 2016). *Drosophila* and

their parasitoids are among the few systems that currently allow us to explore

thoroughly the mechanisms of species interactions at scales ranging from the molecular

to the ecological. Here, we highlight two examples where information compiled in DROP

enables the study of the *Drosophila*-parasitoid system across multiple levels of biological

organization:

DROP includes a DNA reference library of Australian *Drosophila* parasitoids (dataset:

DS-AUSPTOID dx.doi.org/10.5883/DS-AUSPTOID) that connects laboratory experiments

and field research. Molecular vouchers of both hosts and parasitoids were collected

along altitudinal gradients in the rainforest of northern Queensland, Australia (Jeffs et

al., 2021). With this DNA reference library, researchers can detect interactions between

Drosophila and their parasitoids using PCR-based approaches and parasitized pupae

(Hrcek & Godfray, 2015; Jeffs et al., 2020). Surveying host-parasitoid interactions in this

way will improve our understanding of how environmental change alters the structure

of host-parasitoid networks (Morris et al., 2014; Staniczenko et al., 2017; Tylianakis et

al., 2007) by accelerating data collection in the field. In addition, JH established lab

cultures of both hosts and their parasitoids from the same Australian sampling sites with

the aim of conducting laboratory experiments (e.g. Thierry et al., 2021). Molecular

vouchers of the lab strains were then submitted to DROP as a reference database

(datasets: DS-LABS dx.doi.org/10.5883/DS-LABS) to ensure that criteria for species

determination were applied consistently—and will continue to be applied consistently—

between the natural community studies and the laboratory experiments.

The presence of a foundational DNA reference library and species catalogue in

DROP will enable the process of exploring parasitoid biodiversity to become more

efficient. For example, DROP includes molecular vouchers from *Drosophila* parasitoids

that were collected across seasons and along latitudinal gradients in the eastern United

States (Lue et al., 2016, 2018). These data proved to be extremely useful for identifying

species in a more recent exploration of native parasitoid biodiversity across North

America (e.g., Abram et al., 2020). There are additional uses for DROP: curated

specimen collections may be used to document species distributions, phenology,

understand micro-evolutionary patterns, observe the effects of climate change, and

detect and track biological invasions (Funk, 2018; Schilthuizen et al., 2015; Tarli et al.,

2018).

Taxonomic accuracy for biocontrol studies

Unfortunately, the history of biological control includes many examples of

misidentifications that have resulted in failures to employ or establish the expected

control agent, thus hindering eventual success (Buffington et al., 2018; Rosen, 1986;

Huffaker et al. 1962). In the context of biological control research on *Drosophila* pest

species, a simple, reliable, and rapid identification tool for their natural enemies is

essential (Wang et al. 2020b). By anchoring the criteria for determining identities of

organisms being considered for biological control programs, DROP annotation enables

the direct examination of centers of origin for parasitoid species, their co-occurrence

with natural enemies, and the optimal timing for potential introductions of such

enemies (Abram et al., 2020; Daane et al., 2016; Girod et al., 2018a and b; Kimura, 2015;

Mitsui et al., 2007). Because most sequences from DROP are already vetted for

reliability, they can be used to identify biological control agents rapidly, before or after

being brought into quarantine facilities for safety and efficacy testing. This will decrease

the risk of non-target ecological impacts arising from misidentifications and facilitate

regulatory review for releases of effective and specific natural enemies.

In addition to species identification, reference sequences from DROP may be used as

a starting point to create species-specific primers for the accurate identification of

parasitoids, design multiplex PCR assays that rapidly distinguish species in natural or

agricultural ecosystems (Ye et al., 2017), and apply high-throughput molecular

identification diagnostics (Fagan-Jeffries et al., 2018).

Long-term molecular voucher preservation

During the curation of DROP, we found that holotype specimens were missing from

museums for several iconic *Drosophila* parasitoid species: *Asobara tabida* (Nees von

Esenbeck), *Leptopilina clavipes* (Hartig), and *Leptopilina longipes* (Hartig)*. This is not

uncommon and impedes future taxonomic revisions regardless of whether or not

molecular data are used. To avoid contributing to this problem, DROP uses museums as

depositories for ensuring that sequenced vouchers of both described species and

provisional species are permanently stored. In order to stabilize nomenclature, we

further advocate the designation of neotypes (a replacement specimen for a missing

holotype or type series) that have museum-vouchered DNA barcodes and additional

genomic extractions in storage.

Natural history museums are designed to maintain vouchers (including types) for

long-term preservation, and increasingly they implement institutionalized workflows

that link DNA sequences to specimens and specimen metadata (Prendini et al., 2002).

We strongly encourage the deposition of voucher specimens from field surveys and

experimental studies in museum collections, as has been urged by the Entomology

Collections Network (ECN) and required in many PhD programs. No matter how quickly

new molecular techniques are developed or refined, there is no substitute for a reliable

database of voucher specimens when it comes to ensuring the repeatability of biological

research (Funk et al., 2005; Lendemer et al., 2020).

Our results show that species richness of the parasitic wasps that attack *Drosophila*

is severely underestimated, and only a fraction of them have been described. In DROP,

38% of the OTUs are provisional species, and more than 46% of the named OTUs have

synonyms. Remarkably, *Leptopilina heterotoma*, one of the world's most studied

parasitoids, has more than 20 synonyms! As is generally the case, the rate of species

description and revision of *Drosophila* parasitoids lags far behind that with which

molecular sequence data are generated. Ensuring a consistent application of OTU

recognition is therefore essential. With DROP, researchers may ensure consistency in

their application of scientific names, and that those names are valid, making the

daunting process of describing *Drosophila* parasitoids more accurate and efficient. In

addition to the collection of physical museum resources, a central role taxonomists play

in DROP and its curation is that of fostering better integration of taxonomy with

experimental and biodiversity research. Our intention is to perpetuate DROP beyond

this introductory publication. We hope that experts in all areas of *Drosophila*-parasitoid

biology and related fields will join us in this effort.

Conclusion

Taxonomic confusion presents many obstacles in experimental and biodiversity

studies. One way of addressing this impediment is to provide a reliable DNA library with

traceable vouchers (Astrin et al., 2013). Compared to BOLD and GenBank, DROP is a

small database that provides some advantages over an immense genetic database. For

example, it is easier for the research community to have direct communication amongst

themselves, when there is a strong focus on a few specific taxa (Weigand et al., 2019). A

good database has to maintain good quality of molecular data, but even more

challenging is to maintain quality of identification from different sources (Fontes et al.,

2021). In a big database, setting up a universal standard that satisfied all the taxa and

researchers desires is particularly challenging. The curated nature of DROP will allow us

to make strong rules to govern this data and assure users of its fidelity. While GenBank

and BOLD each perform some amount of curation, it could be difficult to agree on

curators for the whole range of animal and plant species catalogued there. We

developed DROP as a resource and platform for gathering and sharing reliable genomic

sequence data for *Drosophila* parasitoids. We hope it will serve as a model for

researchers working with organisms which present similar difficulties. While compiling

DROP, we found that the high number of provisional *versus* named OTUs suggests that

the diversity of parasitic wasps attacking *Drosophila* is greatly underestimated. With this

in mind, DROP represents the start of an important knowledge base that will strengthen

future studies of natural host-parasitoid interactions, population dynamics, biocontrol,

and the impact of climate change on biodiversity and ecosystem services.

Acknowledgements

The idea of DROP project was developed during the 2018 Entomological Society

of America conference, during the symposium "*Drosophila* parasitoids: from molecular

to ecosystem level". We thank Dr. Elijah Talamas for valuable comments on earlier

drafts and Dr. Vid Bakovic for genomic consultation on this project. We also thank Chris

Jeffs for providing some Australian field samples. We are thankful for funding support

from the Czech Science Foundation (17-27184Y). Additional fund for sequencing was

provided by MLB, OTL, and PPAS. Mention of trade names or commercial products in

this publication is solely for the purpose of providing specific information and does not

imply recommendation or endorsement by the USDA. USDA is an equal opportunity

provider and employer.

References

Abram, P. K., Mcpherson, A. E., Kula, R., Hueppelsheuser, T., Perlman, S. J., Curtis, C. I.,
Fraser, J. L., … Buffington, M. (2020). New records of *Leptopilina*, *Ganaspis*, and
Asobara species associated with *Drosophila suzukii* in North America, including
detections of L. japonica and G. brasiliensis. *Journal of Hymenoptera Research*, 78,
1-17, https://doi.org/10.3897/jhr.78.55026
Astrin, J. J., Zhou, X., & Misof, B. (2013). The importance of biobanking in molecular
taxonomy, with proposed definitions for vouchers in a molecular context. *ZooKeys*,
365(SPEC.ISSUE), 67–70. https://doi.org/10.3897/zookeys.365.5875
Baker, C. C. M., Bittleston, L. S., Sanders, J. G., & Pierce, N. E. (2016). Dissecting host-
associated communities with DNA barcodes. *Philosophical Transactions of the*
Royal Society B: Biological Sciences, 371(1702).
https://doi.org/10.1098/rstb.2015.0328
Bergsten, J., Bilton, D. T., Fujisawa, T., Elliott, M., Monaghan, M. T., Balke, M.,
Hendrich, … Vogler, A. P. (2012). The effect of geographical scale of sampling on
DNA barcoding. *Systematic Biology,* 61 (5), 851-869.
https://doi.org/10.1093/sysbio/sys037
Blaimer, B. B., Gotzek, D., Brady, S. G., & Buffington, M. (2020). Comprehensive
phylogenomic analyses re-write the evolution of parasitism within cynipoid wasps.
BMC Ecology and Evolution, 20 (155). https://doi.org/10.1186/s12862-020-01716-2
Brower, A. V. Z., & DeSalle, R. (2002). Patterns of mitochondrial versus nuclear DNA
sequence divergence among nymphalid butterflies: The utility of wingless as a
source of characters for phylogenetic inference. *Insect Molecular Biology*, 7 (1), 73-
82. https://doi.org/10.1046/j.1365-2583.1998.71052.x
Buffington, M., &Forshage, M. (2016). Redescription of *Ganaspis brasiliensis* (Ihering,
1905), new combination, (Hymenoptera: Figitidae) a natural enemy of the Invasive
Drosophila suzukii (Matsumura, 1931) (Diptera: Drosophilidae). *Proceedings of the*
Entomological Society of Washington, 118(1), 1–13. https://doi.org/10.4289/0013-
8797.118.1.1

Buffington, M., Talamas, E. J., & Hoelmer, K. A. (2018). Team Trissolcus: Integrating
taxonomy and biological control to combat the brown marmorated stink bug.
American Entomologist, 64 (4), 224–232
Carton, Y., Boulétreau, M., van Alphen, J. J. M., & van Lenteren, J. C. (1986). The
Drosophila parasitic wasps. In Ashburner M, Carson HL, Thompson JN (Eds), *The*
genetics and biology of Drosophila, (3),348–394.
Carton, Y., Haouas, S., Marrakchi, M., & Hochberg, M. (1991). Two competing parasitoid
species coexist in sympatry. *Oikos*, 60, 222-230. https://doi.org/10.2307/3544869
Condon, M. A., Scheffer, S. J., Lewis, M. L., Wharton, R., Adams, D.C., & Forbes, A. A.
(2014). Lethal interactions between parasites and prey increase niche diversity in a
tropical community. *Science*, *343*(6176), pp.1240-1244.
Daane, K. M., Wang, X.-G., Biondi, A., Miller, B. E., Miller, J. C., Riedl, H., Shearer, P. W.,
… Walton, V. M. (2016). First exploration of parasitoids of *Drosophila suzukii* in
South Korea as potential classical biological agents. *Journal of Pest Science* 89, 823–
835, doi:10.1007/s10340-016-0740-0.
Dolphin, K., & Quicke, D. L. J. (2001). Estimating the global incompletely described
parasitoid wasps. *Biological Journal Of The Linnean Society*, 73 (3), 279-286,
https://doi.org/10.1006
Escalante, P., Ibarra-Vazquez, A., & Rosas-Escobar, P. (2010). Tropical montane
nymphalids in Mexico: DNA barcodes reveal greater diversity. *Mitochondrial DNA*,
21, 30-37, https://doi.org/10.3109/19401736.2010.535527
Ferrer-Suay, M., Staverløkk, A., Selfa, J., Pujade-Villar, J., Naik, S., & Ekrem., T. (2018)
Nuclear and mitochondrial markers suggest new species boundaries in *Alloxysta*
(Hymenoptera: Cynipoidea: Figitidae). Arthropod Systematics & Phylogeny, 76(3),
463-473, doi:10.5883/DSALLOXYST
Fagan-Jeffries, E. P., Cooper, S. J. B., Bertozzi, T., Bradford, T. M., & Austin, A. D. (2018).
DNA barcoding of microgastrine parasitoid wasps (Hymenoptera: Braconidae) using
high-throughput methods more than doubles the number of species known for
Australia. *Molecular Ecology Resources*, 18(5), 1132–1143.
https://doi.org/10.1111/1755-0998.12904
Ferguson, J. W. H. (2002). On the use of genetic divergence for identifying species.
Biological Journal of the Linnean Society, 75, 509–16.
Forbes, A. A., Bagley, R. K., Beer, M. A., Hippee, A. C., & Widmayer, H. A. (2018).
Quantifying the unquantifiable: Why Hymenoptera, not Coleoptera, is the most
speciose animal order. *BMC Ecology*, 18(1), 1–11. https://doi.org/10.1186/s12898-
018-0176-x
Fontes, J. T., Vieire, P. E., Ekrem, T., Soares, P., & Costa, F. O. (2021). BAGS: An
automated barcode, audit & grade system for DNA barcode reference libraries.
Molecular Ecology Resources, 21(2), 573-583. https://doi.org/10.1111/1755-
0998.13262
Funk, V. A. (2018). Collections-based science in the 21st Century. *Journal of Systematics*
and Evolution, 56(3), 175–193. https://doi.org/10.1111/jse.12315
Funk, V. A., Hoch, P. C., Prather, L. A., & Wagner, W. L. (2005). The importance of
vouchers. *Taxon*, 54(1), 127–129. https://doi.org/10.2307/25065309

Gariepy, T. D., Bruin, A., Konopka, J., Scott-Dupree, C., Fraser, H., Bon, M. C., & Talamas,
E. (2019). A modified DNA barcode approach to define trophic interactions
between native and exotic pentatomids and their parasitoids. *Molecular Ecology*,
28(2), 456–470. https://doi.org/10.1111/mec.14868
Giorgini, M., Wang, X.-G., Wang, Y., Chen, F. S., Hougardy, E. , Zhang, H. M., Chen, Z.
Q., … Guerrieri, E. (2019). Exploration for native parasitoids of *Drosophila suzukii* in
China reveals a diversity of parasitoid species and narrow host range of the
dominant parasitoid. *Journal of Pest Science*, 92(2), 509–522.
https://doi.org/10.1007/s10340-018-01068-3
Girod, P., Borowiec, N., Buffington, M., Chen, G., Fang, Y., Kimura, M. T., Peris-Felipo, F.
J., … Kenis, M. (2018). The parasitoid complex of *D. suzukii* and other fruit feeding
Drosophila species in Asia. *Scientific Reports*, 8(1), e11839.
https://doi.org/10.1038/s41598-018-29555-8
Girod, P., Lierhmann, O., Urvois, T., Turlings, T. C. J., Kenis. M., & Haye, T. (2018). Host
specificity of Asian parasitoids for potential classical biological control of *Drosophila*
suzukii. Journal of Pest Science 91,1241–1250, https://doi. doi:10.1007/s10340-
018-1003-z
Goldstein, P. Z., & DeSalle, R. (2011). Integrating DNA barcode data and taxonomic
practice: Determination, discovery, and description. *BioEssays*, 33(2),135-147,
https://doi.org/10.1002/bies.201000036
Gompert, Z., Forister, M. L., Fordyce, J. A., & Nice, C. C. (2008). Widespread mito-nuclear
discordance with evidence for introgressive hybridization and selective sweeps in
Lycaeides. *Molecular Ecology*, 17(24), 5231-5244, https://doi.org/10.1111/j.1365-
294X.2008.03988.x
Grissell, E. (1999). Hymenopteran biodiversity: some alien notions. *American*
Entomologist, 45,236-244.
Hardy, I. C., van Alphen, J. J. M., & Godfray, H. C. J. (1994). Parasitoids: Behavioral and
evolutionary ecology. *The Journal of Animal Ecology*, 63(4), 1009-1010,
https://doi.org/10.2307/5282
Hausmann, A., Haszprunar, G., & Hebert, P. D. N. (2011). DNA barcoding the geometrid
fauna of bavaria (Lepidoptera): Successes, surprises, and questions. *PLoS ONE*, 6(2),
1–9. https://doi.org/10.1371/journal.pone.0017134
Hausmann, A., Miller, S. E., Holloway, J. D., Dewaard, J. R., Pollock, D., Prosser, S. W. J.,
& Hebert, P. D. N. (2016). Calibrating the taxonomy of a megadiverse insect family:
3000 DNA barcodes from geometrid type specimens (Lepidoptera, Geometridae).
Genome, 59(9), 671–684. https://doi.org/10.1139/gen-2015-0197
Hebert, P. D. N., Ratnasingham, S., & DeWaard, J. R. (2003). Barcoding animal life:
Cytochrome c oxidase subunit 1 divergences among closely related species.
Proceedings of the Royal Society B: Biological Sciences, 270 (Suppl.), 96-99,
https://doi.org/10.1098/rsbl.2003.0025
Hendrich, L., Morinière, J., Haszprunar, G., Hebert, P. D. N., Hausmann, A., Köhler, F., &
Balke, M. (2015). A comprehensive DNA barcode database for Central European
beetles with a focus on Germany: Adding more than 3500 identified species to
BOLD. *Molecular Ecology Resources*, 15(4), 795-818, https://doi.org/10.1111/1755-

0998.12354

Hrček, J., & Godfray, H. C. J. (2015). What do molecular methods bring to host-parasitoid
food webs? *Trends in Parasitology*, 31(1), 30–35.
https://doi.org/10.1016/j.pt.2014.10.008

Hrcek, J., Miller, S. E., Quicke, D. L. J., & Smith, M. A. (2011). Molecular detection of
trophic links in a complex insect host-parasitoid food web. *Molecular Ecology*
Resources, 11(5), 786–794. https://doi.org/10.1111/j.1755-0998.2011.03016.x

Huffaker, C. B., Kennett, C. E., Finney, G. L. (1962). Biological control of olive scale,
Pwrlatoria oleae (Cohree), in California by imported Aphytis maculicornis (Masi)
(Hymenoptera: Aphelinidae). *Hilgardia,* 32 (13): 541-636. DOI:
10.3733/hilg.v32n13p541

Hughes, R. D., Woolcock, L. T. (1976). Aphaereta aotea sp. N. (Hymenoptera:
Braconidae), an Alysiine parasite of dung breeding flies. J*ournal of Australian*
Entomological Society, 15, 191-196.

Jeffs, C. T., Terry, J. C. D., Higgie. M., Jandová, A., Konvičková. H., Brown. J. J., Lue. C.-H.,
… Lewis, O. T. (2020). Molecular analyses reveal consistent food web structure with
elevation in rainforest *Drosophila* - parasitoid communities. *Ecography*, 43, 1-11,
https://doi.org/10.1111/ecog.05390

Kim-Jo, C., Gatti, J. L., & Poirié, M. (2019). *Drosophila* cellular immunity against
parasitoid wasps: A complex and time-dependent process. *In Frontiers in*
Physiology, https://doi.org/10.3389/fphys.2019.00603

Kimura, M. T. (2015). Prevalence of exotic frugivorous *Drosophila* species, *D. simulans*
and *D. immigrans* (Diptera: Drosophilidae), and its effects on local parasitoids in
Sapporo, northern Japan. *Applied Entomology and Zoology*, 50(4), 509–515.
https://doi.org/10.1007/s13355-015-0361-8

Kimura, M. T., & Mitsui, H. (2020). *Drosophila* parasitoids (Hymenoptera) of Japan. I*n*
Entomological Science, 23(4), 359-368, https://doi.org/10.1111/ens.12432

Klopfstein, S., Kropf, C., & Baur, H. (2016). Wolbachia endosymbionts distort DNA
barcoding in the parasitoid wasp genus Diplazon (Hymenoptera: Ichneumonidae).
Zoological Journal of the Linnean Society, 177(3), 541–557.
https://doi.org/10.1111/zoj.12380

Kraaijeveld, A. R., & Godfray, H. C. J. (1997). Trade-off between parasitoid resistance and
larval competitive. *Nature*, 389, 278-280, https://doi.org/10.1038/38483

Lefèvre, T., De Roode, J. C., Kacsoh, B. Z., & Schlenke, T. A. (2012). Defence strategies
against a parasitoid wasp in Drosophila: Fight or flight? *Biology Letters*, 8(2), 230-
233, https://doi.org/10.1098/rsbl.2011.0725

Lendemer, J., Thiers, B., Monfils, A. K., Zaspel, J., Ellwood, E. R., Bentley, A., LeVan, K., …
Aime, M. C. (2020). The extended specimen network: A strategy to enhance US
biodiversity collections, promote research and education. *BioScience*, 70(1), 23-30,
https://doi.org/10.1093/biosci/biz140

Lin, C. P., & Danforth, B. N. (2004). How do insect nuclear and mitochondrial gene
substitution patterns differ? Insights from Bayesian analyses of combined datasets.
Molecular Phylogenetics and Evolution, 30(3), 686-702,
https://doi.org/10.1016/S1055-7903(03)00241-0

Lin, X., Stur, E., & Ekrem, T. (2015). Exploring genetic divergence in a species-rich genus
using 2790 DNA barcodes. *PLoS ONE*, 10(9): e0138993.
https://doi.org/10.1371/journal.pone.0138993
Littlefair, J. E., Clare, E. L., & Naaum, A. (2016). Barcoding the food chain: From Sanger to
high-throughput sequencing1. *Genome*, 59(11), 946–958.
https://doi.org/10.1139/gen-2016-0028
Lue, C.-H., Borowy, D., Buffington, M. L., & Leips, J. (2018). Geographic and seasonal
variation in species diversity and community composition of frugivorous *Drosophila*
(Diptera: Drosophilidae) and their *Leptopilina* (Hymenoptera: Figitidae) parasitoids.
Environmental Entomology, 47(5): 1096-1106. https://doi.org/10.1093/ee/nvy114
Lue, C.-H., Driskell, A. C., Leips, J., & Buffington, M. L. (2016). Review of the genus
Leptopilina (Hymenoptera, Cynipoidea, Figitidae, Eucoilinae) from the Eastern
United States, including three newly described species. J*ournal of Hymenoptera*
Research, 53: 35-76. https://doi.org/10.3897/jhr.53.10369
Meiklejohn, K. A., Damaso, N., & Robertson, J. M. (2019). Assessment of BOLD and
GenBank – Their accuracy and reliability for the identification of biological
materials. *PLoS ONE,* 14(6): e0217084.
https://doi.org/10.1371/journal.pone.0217084
Meier, R., Blaimer, B., Buenaventura, E., Hartop, E., von Thomas, R., Srivathsan, A., &
Yeo, D. (2021) A re-analysis of the data in Sharkey et al.,'s (2021) minimalist
revision reveals that BINs do not deserve names, but BOLD Systems needs a
stronger commitment to open science. *bioRxiv*.
https://doi.org/10.1101/2021.04.28.441626
Melk, J. P., & Govind, S. (1999). Developmental analysis of *Ganaspis xanthopoda*, a larval
parasitoid of *Drosophila melanogaster*. *Journal of Experimental Biology*, 202, 1885-
1896
Miller, S. E., Hausmann, A., Hallwachs, W., & Janzen, D. H. (2016). Advancing taxonomy
and bioinventories with DNA barcodes. *Philosophical Transactions of the Royal*
Society Biological Sciences, 371(1702): 20150339. doi: 10.1098/rstb.2015.0339
Mitsui, H., van Achterberg, K., Nordlander, G., & Kimura, M. T. (2007). Geographical
distributions and host associations of larval parasitoids of frugivorous *Drosophilidae*
in Japan. *Journal of Natural History*, 41(25–28), 1731–1738.
https://doi.org/10.1080/00222930701504797
Moreau, S. J. M., Vinchon, S., Cherqui, A., & Prévost, G. (2009). Components of Asobara
venoms and their effects on hosts. *In Advances in Parasitology*, Prévost G (Ed). 70,
217-232, https://doi.org/10.1016/S0065-308X(09)70008-9
Morris, R. J., Gripenberg, S., Lewis, O. T., & Roslin, T. (2014). Antagonistic interaction
networks are structured independently of latitude and host guild. *Ecology Letters*,
17(3), 340-349, https://doi.org/10.1111/ele.12235
Nappi, A. J., & Carton, Y. (2001). Immunogenetic aspects of the cellular immune
response of *Drosophila* against parasitoids. *Immunogenetics*, 52(3–4), 157–164.
https://doi.org/10.1007/s002510000272
Nomano, F. Y., Kasuya, N., Matsuura, A., Suwito, A., Mitsui, H., Buffington ,M.L., &
Kimura, M. T. (2017). Genetic differentiation of *Ganaspis brasiliensis*

(Hymenoptera: Figitidae) from East and Southeast Asia. *Applied Entomology and*
Zoology, 52(3), 429–437. https://doi.org/10.1007/s13355-017-0493-0

Novotny, V., & Miller, S. E. (2014). Mapping and understanding the diversity of insects in
the tropics: Past achievements and future directions. *Austral Entomology*, 53(3),
259–267. https://doi.org/10.1111/aen.12111

O'Grady, P. M., & DeSalle, R. (2018). Phylogeny of the genus *Drosophila*. *Genetics*,
209(1), 1–25. https://doi.org/10.1534/genetics.117.300583

Pentinsaari, M., Ratnasingham, S., Miller, S. E., & Hebert, P. D. N. (2020). BOLD and
GenBank revisited – Do identification errors arise in the lab or in the sequence
libraries? *PLoS One,* 15(4): e0231814. https://doi.org/10.1371/journal.
pone.0231814

Pfeiffer, D. G., Shrader, M. E., Wahls, J. C. E., Willbrand, B. N., Sandum, I., van der Linde,
K., Laub, C. A., … Day, E. R. (2019). African Fig Fly (Diptera: Drosophilidae): Biology,
expansion of geographic range, and its potential status as a soft fruit pest. *Journal*
of Integrated Pest Management, 10(1), 1–8. https://doi.org/10.1093/jipm/pmz018

Prendini, L., Hanner, R., & DeSalle, R. (2002). Obtaining, storing and archiving specimens
and tissue samples for use in molecular studies. *In Techniques in Molecular*
Systematics and Evolution. https://doi.org/10.1007/978-3-0348-8125-8_11

Prévost, G. (2009). Parasitoids of *Drosophila*. *In Advances in parasitology*.
https://doi.org/10.1016/S0065-308X(09)70018-1

Quicke, D. L. J. (2015). The Braconid and Ichneumonid parasitoid wasps: Biology,
systematics, evolution and ecology. Wiley-Blackwell,
https://doi.org/10.1002/9781118907085

Ratnasingham, S., & Hebert, P. D. N. (2007). BARCODING: bold: The Barcode of Life Data
System (http://www.barcodinglife.org). *Molecular Ecology Notes*, 7(3), 355-364,
https://doi.org/10.1111/j.1471-8286.2007.01678.x

Ratnasingham, S., & Hebert, P. D. N. (2013). A DNA-based registry for all animal species:
the barcode index number (BIN) system. *PLoS ONE,* 8(7): e66213.
https://doi.org/10.1371/journal.pone.0066213

Raupach, M. J., Hendrich, L., Kuchler, S. M., Deister, F., Moriniere, J., & Gossner, M. M.
(2014). Building-Up of a DNA Barcode Library for true bugs (Insecta: Hemiptera:
Heteroptera) of Germany reveals taxonomic uncertainties and surprises. *PLoS ONE*,
9(9), 1–13. https://doi.org/10.1371/journal.pone.0106940

Remsen, J., & O'Grady, P. (2002). Phylogeny of Drosophilinae (Diptera: Drosophilidae),
with comments on combined analysis and character support. *Molecular*
Phylogenetics and Evolution, 24(2), 249-264, https://doi.org/10.1016/S1055-
7903(02)00226-9

Reumer, B, M., van Alphen, J. J. M., & Kraaijeveld, K. (2012). Occasional males in
parthenogenetic populations of Asobara japonica (Hymenoptera: Braconidae): Low
Wolbachia titer or incomplete coadaptation. *Heredity*, 108(3), 341-346,
https://doi.org/10.1038/hdy.2011.82

Rosen, D. (1986). The role of taxonomy in effective biological control programs.
Agriculture, Ecosystems & Environment, 15(2-3), 121-129.
https://doi.org/10.1016/0167-8809(86)90085-X

Santos, W. G. N., Fernandes, E. C., Souza, M. M., Guimarães, J. A., & Araujo, E. L. (2016).
First record of Eucoilinae (Hymenoptera: Figitidae), parasitoids of African fig fly
Zaprionus indianus Gupta (Diptera: Drosophilidae), in the Caatinga biome.
Semina:Ciencias Agrarias, 37(5), 3055–3058. https://doi.org/10.5433/1679-
0359.2016v37n5p3055
Schilthuizen, M., Vairappan, C. S., Slade, E. M., Mann, D. J., & Miller, J. A. (2015).
Specimens as primary data: Museums and "open science." *Trends in Ecology and*
Evolution, 30(5), 237–238. https://doi.org/10.1016/j.tree.2015.03.002
Schlenke, T. A., Morales, J., Govind, S., & Clark, A. G. (2007). Contrasting infection
strategies in generalist and specialist wasp parasitoids of *Drosophila melanogaster*.
PLoS Pathogens, 3(10):e158, https://doi.org/10.1371/journal.ppat.0030158
Schindel, D., & Miller, S. E. (2010). Provisional Nomenclature the on-ramp to taxonomic
names. In: Polaszek, A. (Ed), *Systema Nature*, 250: The Linnaean Ark. CRC, Boca
Raton, 109-115.
Seehausen, M. L., Ris, N., Driss, L., Racca, A., Girod, P., Warot, S., Borowiec, N., Tosevski,
I., & Kenis, M. (2020). Evidence for a cryptic parasitoid species reveals its suitability
as a biological control agent. *Scientific reports*, 10: 19096.
https://doi.org/10.1038/s41598-020-76180-5
Shokralla, S., Spall, J. L., Gibson, J. F., & Hajibabaei, M. (2012). Next-generation
sequencing technologies for environmental DNA research. *In Molecular Ecology*,
21(8), 1794-1805, https://doi.org/10.1111/j.1365-294X.2012.05538.x
Smith, M. A., Fisher, B. L., & Hebert, P. D. N. (2005). DNA barcoding for effective
biodiversity assessment of a hyperdiverse arthropod group: The ants of
Madagascar. *Philosophical Transactions of the Royal Society Biological Sciences*,
360(1462), 1825-1834, https://doi.org/10.1098/rstb.2005.1714
Smith, M. A., Fernandez-Triana, J. L., Eveleigh, E., Gomez, J., Guclu, C., Hallwachs, W.,
Hebert. P. D. N., … Zaldivar-Riveron, A. (2013). DNA barcoding and the taxonomy of
Microgastrinae wasps (Hymenoptera, Braconidae): impacts after 8 years and nearly
20000 sequences. *Molecular Ecology Resource*s, 13, 168-276,
https://doi.org/10.1111/1755-0988.12038
Staniczenko, P. P. A., Reed-Tsochas, F., Lewis, O. T., Tylianakis, J. M., Albrecht, M.,
Coudrain, V., & Klein, A. M. (2017). Predicting the effect of habitat modification on
networks of interacting species. *Nature Communications*, 8, 792,
https://doi.org/10.1038/s41467-017-00913-w
Tarli, V. D., Grandcolas, P., & Pellens, R. (2018). The informative value of museum
collections for ecology and conservation: A comparison with target sampling in the
Brazilian Atlantic forest. *PLoS ONE*, 13(11): e0205710.
https://doi.org/10.1371/journal.pone.0205710
Terry, J. C. D., Chen, J., & Lewis, O. T. (2021). The effect of natural enemies on the
coexistence of competing species - an empirical test using Bayesian modern
coexistence theory. *bioRxiv*: https://doi.org/10.1101/2020.08.27.270389
Thierry, M., Pardikes, N. A., Lue, C.-H., Lewis, O. L., & Hrcek, J. (2021). Experimental
warming influences species abundances in a *Drosophila* host community through
direct effects on species performance rather than altered competition and

parasitism. *PLoS ONE*, 16(2): e0245029.

https://doi.org/10.1371/journal.pone.0245029

Troudet, J., Vignes-Lebbe, R., Grandcolas, P., & Legendre, F. (2018). The increasing

disconnection of primary biodiversity data from specimens: How does it happen

and how to handle it? *Systematic Biology*, 67(6), 1110–1119.

https://doi.org/10.1093/sysbio/syy044

Tylianakis, J. M., Tscharntke, T., & Lewis, O. T. (2007). Habitat modification alters the

structure of tropical host-parasitoid food webs. *Nature*, 445(7124), 202–205.

https://doi.org/10.1038/nature05429

Wachi, N., Nomano, F. Y., Mitsui, H., Kasuya, N., & Kimura, M. T. (2015). Taxonomy and

evolution of putative thelytokous species of Leptopilina (Hymenoptera: Figitidae)

from Japan, with description of two new species. *Entomological Science*, 18(1), 41–

54. https://doi.org/10.1111/ens.12089

Wang, X.-G., Biondi, A., & Daane, K. M. (2020). Functional responses of three candidate

Asian larval parasitoids evaluated for classical biological control of *Drosophila*

suzukii. Journal of Economic Entomology, 113(1): 73–80. doi: 10.1093/jee/toz265

Wang, X.-G., Biondi, A., Nance. A. N., Zappalà, L., Hoelmer, K. A., & Daane, K. M. (2021).

Assessment of Asobara japonica as a potential biological control agent for the

spotted wing drosophila, Drosophila suzukii. *Entomologia Generalis (In Press)* doi:

10.1127/entomologia/2020/1100

Wang, X.-G., Lee, J., Daane, K.M., Buffington, M., & Hoelmer, K. A. (2020). Biological

control of *Drosophila suzukii*. *CAB Reviews* 54, 10.1079/PAVSNNR202015054

Weigand, H., Beermann, A. j., Čiampor, F., Costa, F. O., Csabai, Z., Duarte, S., Geiger, M.

F., … Ekrem, T. (2019). DNA barcode reference libraries for the monitoring of

aquatic biota in Europe: Gap-analysis and recommendations for future work.

Science of The Total Environment. 687(15), 499-254.

https://doi.org/10.1016/j.scitotenv.2019.04.247

Xiao, J. H., Wang, N. X., Murphy, R. W., Cook, J., Jia, L. Y., & Huang, D. W. (2012).

Wolbachia infection and dramatic intraspecific mitochondrial DNA divergence in a

fig wasp. *Evolution*, 66, 1907-1916, https://doi.org/10.1111/j.1558-

5646.2011.01561.x

Xie. J., Tiner, B., Vilchez, I., & Mateos, M. (2011). Effect of the *Drosophila* endosymbiont

Spiroplasma on parasitoid wasp development and on the reproductive fitness of

wasp-attacked fly survivors. *Evolutionary Ecology*, 25, 1065-1079,

https://doi.org/10.1007/s10682-010-9453-7

Xie, J., Winter, C., Winter, L., & Mateos, M. (2015). Rapid spread of the defensive

endosymbiont *Spiroplasma* in *Drosophila hydei* under high parasitoid wasp

pressure. *FEMS Microbiology Ecology*, 91(2), 1-11,

https://doi.org/10.1093/femsec/iu017

Yassin, A. (2013). Phylogenetic classification of the Drosophilidae Rondani (Diptera): The

role of morphology in the postgenomic era. *Systematic Entomology*,

https://doi.org/10.1111/j.1365-3113.2012.00665.x

Yassin, A., & David, J. R. (2010). Revision of the Afrotropical species of *Zaprionus*

(Diptera, Drosophilidae), with descriptions of two new species and notes on

internal reproductive structures and immature stages. *ZooKeys,* 51, 33-72,
https://doi.org/10.3897/zookeys.51.380
Ye, Z., Vollhardt, I. M. G., Girtler, S., Wallinger, C., Tomanovic, Z., & Traugott, M. (2017).
An effective molecular approach for assessing cereal aphid-parasitoid-
endosymbiont networks. *Scientific Reports*, 7(1), 1–12.
https://doi.org/10.1038/s41598-017-02226-w
Yu, D. S. K., 2016, Global index for Ichneumonoidea. Last updated October 22, 2016
https://web.archive.org/web/20161022093945/http:/ichneumonoidea.name/gl
obal.php
1008

Data Accessibility

The DROP database is freely accessible at Zenodo depository

(http://doi.org/10.5281/zenodo.4519656). Sequences from GenBank and BOLD, all

compiled in BOLD, DROP project, DS-DROPAR dataset dx.doi.org/10.5883/DS-DROPAR.

New sequences have been deposited in BOLD, DROP project (datasets: DS-LABS

dx.doi.org/10.5883/DS-LABS and DS-AUSPTOID dx.doi.org/10.5883/DS-AUSPTOID).

1015

Author Contributions

The initial project idea was originated by C-HL, MLB, JH, MM, TS, JV, SG, and

PPAS. Molecular work was conducted by C-HL, SS, ML, AJ, and AD. BOLD and GenBank

data was harvested by TAE and C-HL. Figures were made by AL and C-HL. Laboratory and

field sample preparations were conducted by MTK, YC, TS, MM, SG, JV, EG, MG, XW,

KM, KMD, PA, NAP, MT, JJB, MP, FMJ, WDT, JSD, BW, OTL, PPAS, JL and AL. Taxonomic

concepts and interpretations were conducted by RRK, MLB, CH-L, PG, and SEM. DROP

database was built by JH and C-HL. All authors contributed to review and final revisions

of the manuscript, which was written primarily by C-HL, MLB and JH.

1025

Figures:

1027

Figure 1: An example of difficulties of molecular identification demonstrated on
Ganaspis xanthopoda and *G. brasiliensis*. Only two sequences (in bold text) can be
reliably used for identification and are included in DROP database. To select the
sequences, we searched the BINs associated with the organism's name "*Ganaspis
xanthopoda*" (green) or "*Ganaspis brasiliensis*" (purple) in BOLD. From each BIN, two
sequences from each species were selected to build a neighbor-joining tree (bottom axis
indicated % genetic divergence). There was a total of 6 BINs (gray boxes) in this
sequence complex. Of these, 4 BINs contained both species names and without
examination of vouchers, identification would be impossible. In DROP, vouchers from
two sequences, **MG755073** and **MG755072**, were deposited in CNR-IPSP (Table S2),
examined by taxonomists and identified as *G. brasiliensis*. These two COI sequences can
now be used to reliably identify *G. brasiliensis*. For *G. xanthopoda,* there were no
available vouchers or reliable sequences that passed DROP standards to use for
identification. Species delimitation between *G. brasiliensis* and *G. xanthopoda* is
convoluted, varies according to arbitrary % genetic divergence (gray vertical lines), and
needs an integrative taxonomic revision.

1044

Figure 2: Concept of a centralized, vetted, curated database for _Drosophila_ Parasitoids
(DROP) we developed. First, we provide a species and provisional species catalog with
correct taxonomy. Second, to provide a reliable genetic reference library, we link
genetic data (DNA sequences, genomes, transcriptomes, proteomes) to a voucher
connected to the species catalog. Third, we link the two primary sources of data (field
surveys and laboratory experiments) by requiring a permanent deposition of vouchers
and sequences in order to be included in DROP.

Tables:

Table 1: List of species and provisional species included in DROP. For additional
taxonomic details, see DROP.

Superfamily	Family	Genus	Species_Name	Author
Chalcidoidea	Encyrtidae		_drop_Cha2_sp12_	
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_drop_ IR1_sp41_	Kimura
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_drop_BG1_sp42_	Kimura
Chalcidoidea	Encyrtidae	_Tachinaephagus_	_zealandicus_	Ashmead 1904
Chalcidoidea	Pteromalidae		_drop_Pte69_sp11_	
Chalcidoidea	Pteromalidae	_Pachycrepoideus_	_vindemmiae_	(Rondani, 1875)
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_IR1_sp38_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_NG1_sp39_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drop_SK1_sp40_	Kimura
Chalcidoidea	Pteromalidae	_Spalangia_	_drosophilae_	Ashmead 1887
Chalcidoidea	Pteromalidae	_Spalangia_	_erythromera_	Foerster 1850
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_dubia_	(Ashmead, 1896)
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_microptera_	(Lindeman, 1887)
Chalcidoidea	Pteromalidae	_Trichomalopsis_	_nigricola_	Boucek

Chalcidoidea	Pteromalidae	*Trichomalopsis*	*sarcophagae*	(Gahan, 1914)
Chalcidoidea	Pteromalidae	*Vrestovia*	*brevior*	Boucek 1993
Chalcidoidea	Pteromalidae	*Vrestovia*	*fidenas*	(Walker, 1848)
Chalcidoidea	Pteromalidae		*drop_ PacAtl_sp46*	
Chalcidoidea	Pteromalidae		*drop_ PachyPort_sp45*	
Chalcidoidea			*drop_ CH_sp64*	
Cynipoidea	Figitidae	*Ganaspis*	*brasiliensis*	(Ihering, 1905)
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp51*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp52*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gan_sp53*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp1_sp67*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp2_sp68*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ Gsp50_sp66*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_ IR1_sp25*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*drop_ IR2_sp26*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*drop_Gan1_sp1*	
Cynipoidea	Figitidae	*Ganaspis*	*drop_TK1_sp27*	Kimura
Cynipoidea	Figitidae	*Ganaspis*	*hookeri*	Craword 1913
Cynipoidea	Figitidae	*Ganaspis*	*mahensis*	Kieffer 1911
Cynipoidea	Figitidae	*Ganaspis*	*mellipes*	(Say, 1826)
Cynipoidea	Figitidae	*Ganaspis*	*mundata*	Forster 1869
Cynipoidea	Figitidae	*Ganaspis*	*seticornis*	(Hellen, 1960)
Cynipoidea	Figitidae	*Ganaspis*	*tenuicornis*	Kieffer 1904
Cynipoidea	Figitidae	*Ganaspis*	*xanthopoda*	(Ashmead, 1896)
Cynipoidea	Figitidae	*Kleidotoma*	*bicolor*	(Giraud, 1860)
Cynipoidea	Figitidae	*Kleidotoma*	*dolichocera*	Thomson 1877
Cynipoidea	Figitidae	*Kleidotoma*	*drop_TK1_sp28*	Kimura
Cynipoidea	Figitidae	*Kleidotoma*	*filicornis*	(Cameron, 1889)
Cynipoidea	Figitidae	*Kleidotoma*	*icarus*	(Quinlan, 1964)
Cynipoidea	Figitidae	*Kleidotoma*	*psiloides*	Westwood 1833
Cynipoidea	Figitidae	*Kleidotoma*	*tetratoma*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptolamina*	*drop_Fig64_sp5*	
Cynipoidea	Figitidae	*Leptolamina*	*drop_Lmn_sp6*	
Cynipoidea	Figitidae	*Leptolamina*	*drop_TK1_sp29*	Kimura
Cynipoidea	Figitidae	*Leptolamina*	*gressitti*	Yoshimoto & Yasumatsu 1965
Cynipoidea	Figitidae	*Leptolamina*	*papuensis*	Yoshimoto 1963
Cynipoidea	Figitidae	*Leptolamina*	*ponapensis*	Yoshimoto 1962
Cynipoidea	Figitidae	*Leptolamina*	*seychellensis*	(Kieffer, 1911)

Cynipoidea	Figitidae	*Leptopilina*	*atraticeps*	(Kieffer, 1911)
Cynipoidea	Figitidae	*Leptopilina*	*australis*	(Belizin, 1966)
Cynipoidea	Figitidae	*Leptopilina*	*boulardi*	(Barbotin, Carton & Kelner-Pillault, 1979)
Cynipoidea	Figitidae	*Leptopilina*	*clavipes*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptopilina*	*cupulifera*	(Kieffer, 1916)
Cynipoidea	Figitidae	*Leptopilina*	*decemflagella*	Lue & Buffington 2017
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp54*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp55*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp56*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp57*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp58*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp59*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp60*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp61*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_ Lep_sp62*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_BG1_sp34*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig059_sp4*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig124_sp2*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_Fig58_sp3*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_IR1_sp30*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_NG1_sp33*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_SK1_sp35*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_STL_sp7*	
Cynipoidea	Figitidae	*Leptopilina*	*drop_TK2_sp31*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*drop_TK3_sp32*	Kimura
Cynipoidea	Figitidae	*Leptopilina*	*fimbriata*	(Kieffer, 1901)
Cynipoidea	Figitidae	*Leptopilina*	*freyae*	Allemand & Nordlander 2002
Cynipoidea	Figitidae	*Leptopilina*	*guineaensis*	Allemand & Nordlander 2002
Cynipoidea	Figitidae	*Leptopilina*	*heterotoma*	(Thomson, 1862)
Cynipoidea	Figitidae	*Leptopilina*	*japonica japonica*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*lasallei*	Buffington & Guerrieri 2020
Cynipoidea	Figitidae	*Leptopilina*	*leipsi*	Lue & Buffington 2018
Cynipoidea	Figitidae	*Leptopilina*	*lonchaeae*	(Cameron, 1912)
Cynipoidea	Figitidae	*Leptopilina*	*longipes*	(Hartig, 1841)
Cynipoidea	Figitidae	*Leptopilina*	*mahensis*	(Kieffer, 1911)

Cynipoidea	Figitidae	*Leptopilina*	*maia*	Lue & Buffington 2016
Cynipoidea	Figitidae	*Leptopilina*	*maria*	(Girault, 1930)
Cynipoidea	Figitidae	*Leptopilina*	*orientalis*	Allemand & Nordlander 2002
Cynipoidea	Figitidae	*Leptopilina*	*pacifica*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*rufipes*	(Cameron, 1908)
Cynipoidea	Figitidae	*Leptopilina*	*rugipunctata*	(Yoshimoto, 1962)
Cynipoidea	Figitidae	*Leptopilina*	*ryukyuensis*	Novkovic & Kimura 2011
Cynipoidea	Figitidae	*Leptopilina*	*tokioensis*	Wachi & Kimura 2015
Cynipoidea	Figitidae	*Leptopilina*	*tsushimaensis*	Wachi & Kimura 2015
Cynipoidea	Figitidae	*Leptopilina*	*victoriae*	Nordlander 1980
Cynipoidea	Figitidae	*Rhoptromeris*	*heptoma*	(Hartig, 1840)
Cynipoidea	Figitidae	*Rhoptromeris*	*nigriventris*	Nordlander 1978
Cynipoidea	Figitidae	*Rhoptromeris*	*rufiventris*	(Giraud, 1860)
Cynipoidea	Figitidae	*Rhoptromeris*	*villosa*	(Hartig, 1840)
Cynipoidea	Figitidae		*drop_Lg500_sp43*	
Ichneumonoidea	Braconidae	*Alysia*	*drop_SP1_sp24*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*aotea*	Hughes & Woolcock 1976
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_SP1_sp15*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_TK1_sp13*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*drop_TM1_sp14*	Kimura
Ichneumonoidea	Braconidae	*Aphaereta*	*minuta*	(Nees, 1811)
Ichneumonoidea	Braconidae	*Aphaereta*	*pallipes*	(Say, 1829)
Ichneumonoidea	Braconidae	*Aphaereta*	*scaptomyzae*	Fischer 1966
Ichneumonoidea	Braconidae	*Areotetes*	*striatiferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Areotetes*	*carinuliferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Asobara*	*ajbelli*	Berry 2007
Ichneumonoidea	Braconidae	*Asobara*	*albiclava*	Berry 2007
Ichneumonoidea	Braconidae	*Asobara*	*antipoda*	(Ashmead, 1900)
Ichneumonoidea	Braconidae	*Asobara*	*bactrocerae*	(Gahan, 1952)
Ichneumonoidea	Braconidae	*Asobara*	*brevicauda*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*citri*	(Fischer, 1963)
Ichneumonoidea	Braconidae	*Asobara*	*drop_KG1_sp16*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_NG1_sp17*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_SK2_sp20*	Kimura

Ichneumonoidea	Braconidae	*Asobara*	*drop_SP1_sp18*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*drop_Sp2_sp19*	Kimura
Ichneumonoidea	Braconidae	*Asobara*	*elongata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*gahani*	(Papp, 1969)
Ichneumonoidea	Braconidae	*Asobara*	*japonica*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Asobara*	*kenyaensis*	Peris-Felipo 2014
Ichneumonoidea	Braconidae	*Asobara*	*leveri*	(Nixon, 1939)
Ichneumonoidea	Braconidae	*Asobara*	*mesocauda*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*orientalis*	Viereck 1913
Ichneumonoidea	Braconidae	*Asobara*	*persimilis*	(Prince, 1976)
Ichneumonoidea	Braconidae	*Asobara*	*pleuralis*	(Ashmead, 1905)
Ichneumonoidea	Braconidae	*Asobara*	*rossica*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Asobara*	*rufescens*	(Fˆrster, 1862)
Ichneumonoidea	Braconidae	*Asobara*	*tabida*	(Nees, 1834)
Ichneumonoidea	Braconidae	*Asobara*	*triangulata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Asobara*	*turneri*	Peris-Felipo 2014
Ichneumonoidea	Braconidae	*Asobara*	*unicolorata*	van Achterberg & Guerrieri 2016
Ichneumonoidea	Braconidae	*Aspilota*	*albertica*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*andyaustini*	Wharton 2002
Ichneumonoidea	Braconidae	*Aspilota*	*angusta*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*concolor*	Nees 1812
Ichneumonoidea	Braconidae	*Aspilota*	*parecur*	Berry 2007
Ichneumonoidea	Braconidae	*Aspilota*	*villosa*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*barrattae*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*longworthi*	Berry 2007
Ichneumonoidea	Braconidae	*Dinotrema*	*philipi*	Berry 2007
Ichneumonoidea	Braconidae		*drop_Aso_sp8*	
Ichneumonoidea	Braconidae	*Opiognathus*	*pactus*	(Haliday, 1837)
Ichneumonoidea	Braconidae	*Opius*	*bellus*	Gahan 1930
Ichneumonoidea	Braconidae	*Opius*	*cinerariae*	Fischer
Ichneumonoidea	Braconidae	*Opius*	*crenuliferus*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Opius*	*monilipalpis*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Opius*	*ocreatus*	(Papp)
Ichneumonoidea	Braconidae	*Opius*	*pallipes*	Wesmael 1835
Ichneumonoidea	Braconidae	*Opius*	*pteridiophilus*	Wharton & Austin 1990

Ichneumonoidea	Braconidae	*Opius*	*pterus*	Wharton & Austin 1990
Ichneumonoidea	Braconidae	*Opius*	*trimaculatus*	Spinola
Ichneumonoidea	Braconidae	*Opius*	*youi*	Li & van Achterberg 2013
Ichneumonoidea	Braconidae	*Phaenocarpa*	*conspurcator*	(Haliday, 1838)
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drop_IR1_sp22*	Kimura
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drop_TK1_sp21*	Kimura
Ichneumonoidea	Braconidae	*Phaenocarpa*	*tacita*	Stelfox 1941
Ichneumonoidea	Braconidae	*Phaenocarpa*	*drosophilae*	(Fischer 1975)
Ichneumonoidea	Braconidae	*Tanycarpa*	*bicolor*	(Nees, 1814)
Ichneumonoidea	Braconidae	*Tanycarpa*	*chors*	Belokobylskij 1998
Ichneumonoidea	Braconidae	*Tanycarpa*	*drop_NG1_sp23*	Kimura
Ichneumonoidea	Braconidae	*Tanycarpa*	*punctata*	van Achterberg 1976
Ichneumonoidea	Braconidae		*drop_ Aly_sp47*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp48*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp49*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp50*	
Ichneumonoidea	Braconidae		*drop_ Aly_sp63*	
Ichneumonoidea	Braconidae		*drop_ Aso_sp69*	
Diaprioidea	Diapriidae	*Trichopria*	*anastrephae*	Costa Lima 1940
Diaprioidea	Diapriidae	*Trichopria*	*drop_ BG1_sp37*	Kimura
Diaprioidea	Diapriidae	*Trichopria*	*drop_ Dia70_sp65*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_ Tri_sp44*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_Bdia_sp10*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_Dia127_sp9*	
Diaprioidea	Diapriidae	*Trichopria*	*drop_TK1_sp36*	Kimura
Diaprioidea	Diapriidae	*Trichopria*	*drosophilae*	(Kieffer, 1912)
Diaprioidea	Diapriidae	*Trichopria*	*modesta*	(Ratzeburg, 1848)

1058