1 SARS-CoV-2 infection models using *in vivo* and *in vitro* hACE2-lentivirus transduction
2
3

4 Daniel J. Rawle[1*], Thuy Le[1], Troy Dumenil[1], Kexin Yan[1], Bing Tang[1], Cameron Bishop[1], Andreas
5 Suhrbier[1,2].
6
7
8 [1] Immunology Department, QIMR Berghofer Medical Research Institute, Brisbane, Queensland. 4029,
9 Australia.
10 [2] GVN Center of Excellence, Australian Infectious Disease Research Centre, Brisbane, Queensland 4029
11 and 4072, Australia.
12
13
14 [*]Corresponding author. Email: Daniel.Rawle@qimrberghofer.edu.au

SUMMARY
SARS-CoV-2 uses the human ACE2 receptor (hACE2), with mouse ACE2 (mACE2) unable to
support infection. Herein we describe an ACE2-lentivirus system and illustrate its utility *in vitro*
and *in vivo*. Transduction of non-permissive cell lines with hACE2 imparted replication
competence, and transduction with mACE2 containing N30D, N31K, F83Y and H353K
mutations, to match hACE2, rescued SARS-CoV-2 replication. hACE2-lentivirus transduction
of C57BL/6J, IFNAR[-/-] and IL-28RA[-/-] mice lungs indicated type I and III IFN receptor knockout
had minimal effect on virus replication. However, RNA-Seq illustrated that they were required
for inflammatory responses in C57BL/6J mice, which were similar to those seen in COVID-19
patients. Finally, we illustrate the utility of hACE2 transduction for vaccine evaluation in
C57BL/6J mice. The hACE2-lentivirus system thus has broad application in SARS-CoV-2
research, in particular allowing access to GM mice, while also offering the inherent advantages
of lentiviral transduction such as stable genomic integration.

60
61
62
63 **KEYWORDS**
64 SARS-CoV-2, lentivirus, ACE2, GM mice, vaccination, mutations, interferon, inflammation, RNA-seq

INTRODUCTION

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) has spread rapidly into a global pandemic (Wu et al., 2020a). SARS-CoV-2 infection can be asymptomatic, but is also the etiological agent of coronavirus disease 2019 (COVID-19), with acute respiratory distress syndrome (ARDS) representing a common severe disease manifestation (Guan et al., 2020). The SARS-CoV-2 pandemic has sparked unprecedented global research to understanding mechanisms of virus replication and disease pathogenesis, and to generate vaccines and treatments. Key to these efforts has been the development of animal models of SARS-CoV-2 infection and COVID-19 disease (Muñoz-Fontela et al., 2020).

The receptor binding domain (RBD) of the spike glycoprotein of SARS-CoV-2 binds human Angiotensin-Converting Enzyme 2 (hACE2) as the primary receptor for cell attachment and entry (Zhao et al., 2020). Mice do not support productive virus replication because the SARS-CoV-2 spike binds poorly to mouse ACE2 (mACE2) (Zhou et al., 2020). Expression of hACE2 in mice often via a transgene allows SARS-CoV-2 infection and can recapitulate aspects of COVID-19, with hACE2 expressed under control of various promoters, including K18 (Golden et al., 2020; McCray et al., 2007; Moreau et al., 2020; Oladunni et al., 2020; Rathnasinghe et al., 2020; Winkler et al., 2020; Yinda et al., 2021), mACE2 (Bao et al., 2020; Sun et al., 2020b), HFH4 (Jiang et al., 2020b), or chicken β-actin (Tseng et al., 2007). These mouse models all differ in various aspects including level of virus replication, disease manifestations and tissue tropisms (Muñoz-Fontela et al., 2020), but do not provide a simple mechanism whereby genetically modified (GM) mice can be exploited for SARS-CoV-2/COVID-19 research. Two systems that do allow the latter, involve transient hACE2 expression in mouse lungs using adenovirus vectors (Ad5) or adeno-associated virus (AAV), which impart the capacity for productive SARS-CoV-2 replication in mouse lungs (Hassan et al., 2020; Israelow et al., 2020; Rathnasinghe et al., 2020). A third involves use of mouse adapted SARS-CoV-2 (Dinnon et al., 2020; Gu et al., 2020; Leist et al., 2020), although use of this mutated virus may complicate evaluation of interventions targeting human SARS-CoV-2.

Lentivirus-mediated gene expression in mice lung epithelium has been investigated as a treatment for cystic fibrosis and provides long term expression of the cystic fibrosis transmembrane conductance regulator (CFTR) (Alton et al., 2017b; Buckley et al., 2008; Copreni et al., 2010; Copreni et al., 2004; Kremer et al., 2007; Limberis et al., 2010; Wilson, 2004). These studies demonstrated that VSV-G pseudotyped lentivirus can transduce mouse airway epithelial cells and their progenitors resulting in long-term gene expression. Herein we describe an ACE2-lentiviral system that conveys SARS-CoV-2 replication competence *in vitro* and allows productive infection of mouse lungs *in vivo*. We illustrate that GM mice can thereby be accessed for SARS-CoV-2/COVID-19 research, with experiments using IFNAR$^{-/-}$ and IL-28RA$^{-/-}$ mice showing the importance of type I and III IFN responses for SARS-CoV-2-induced inflammation. We also illustrate the use of hACE2-lentiviral transduction of lungs of wild-type C57BL/6J mice for evaluation of vaccines.

RESULTS

hACE2-lentiviruses for evaluating the role of RBD binding residues

hACE2 coding sequence (optimized for mouse codon usage) was cloned into the dual promoter lentiviral vector pCDH-EF1α-MCS-BGH-PGK-GFP-T2A-Puro (herein referred to as pCDH) (Figure 1A), which had been modified to contain the full length EF1α promoter. Previous studies show the EF1α promoter effectively drives gene expression after intranasal VSV-G pseudotyped lentivirus transduction of mouse lungs (Alton et al., 2017a). The GFP marker and puromycin resistance gene (cleaved from GFP by T2A) is expressed from a separate 3-phosphoglycerate kinase (PGK) promoter (Fig 1A). VSV-G pseudotyped pCDH-hACE2 lentivirus was produced by co-transfection of the lentiviral vector with VSV-G and Gag-Pol-Rev plasmids in HEK293T cells, and was used to transduce HEK293T (human embryonic kidney)

cells followed by puromycin selection. HEK293T cells do not express hACE2 (Crawford et al., 2020)
and therefore do not support SARS-CoV-2 replication, whereas HEK293T cells transduced with hACE2
supported substantial virus replication (Figure 1A). A T92Q mutation in hACE2 was introduced to
determine if removing the N90 glycosylation motif increased SARS-CoV-2 replication, as computer-
based modelling predicted enhanced affinity for spike RBD with removal of this glycan (Chan et al.,
2020; Mehdipour and Hummer, 2020). We also introduced K31N/K353H and T27Y/L79Y/N330Y
mutations into hACE2 as these changes were predicted by computer modeling to reduce and enhance
affinity for spike RBD (Chan et al., 2020), respectively. HEK293T cells expressing these aforementioned
hACE2 mutations did not significantly affect SARS-CoV-2 replication (Fig 1B), suggesting that the N90
glycan, and K31/K353 residues in hACE2 are not alone key determinants of viral infectivity.
151
Characterizing residues responsible for the inability of mACE2 to support SARS-CoV-2 replication
The ACE2-lentivirus system was used to introduce amino acid substitutions into mACE2 to identify
amino acids that make mACE2-expressing cells unable to support SARS-CoV-2 replications. Analysis of
crystal structures and species susceptibilities has suggested residues that may be responsible for the
differences in binding of the RBD to hACE2 and mACE2 (Alexander et al., 2020; Damas et al., 2020;
Griffin et al., 2020; Rodrigues et al., 2020; Wang et al., 2020). Based on these *in silico* studies, we
identified seven residues potentially responsible for the human-mouse differences. mACE2-lentivirus
vectors were then generated that contained all seven substitutions
(N30D/N31K/T79L/S82M/F83Y/A329E/H353K), with two further vectors constructed to identify the
minimum requirements for SARS-CoV-2 replication. The latter contained a subset of the seven changes;
four amino acid changes (N30D/N31K/F83Y/H353K) and two amino acid changes (N31K/H353K).
As expected HEK293T cells expressing mACE2 did not support productive SARS-CoV-2
replication, and CPE was not observed. In contrast, all three mACE2 mutants rescued virus replication
and infection resulted in CPE (Figure 1C-D). However, virus replication was ≈1 log lower for cells
expressing mACE2 with only two substitution (N31K/H353K) (Figure 1C) and infection produced less
CPE (Figure 1D). Thus two amino acid changes (N31K and H353K) in mACE2 were sufficient to
support SARS-CoV-2 replication; however, at least four changes (N30D/N31K/F83Y/H353K) were
required for virus replication and CPE to be comparable to that seen in cells expressing hACE2.
Thus of all the potential interactions between the RBD and ACE2 previously identify by X ray
crystallography and of all the amino acid differences between human and mouse ACE2 (Damas et al.,
2020; Wang et al., 2020), it emerges 4 are key to determining infectivity differences for cells expressing
hACE2 or mACE2. In the aforementioned crystal structures, hACE2-D30 interacts with S-K417 via a salt
bridge, hACE2-K31 interacts with S-Q493, hACE2-Y83 interacts with S-N487, and hACE2-K353 forms
hydrogen bonds with S-G496 and S-G502 (Figure 1E) Our studies highlight the utility of the ACE2-
lentivirus system for studying the role of ACE2 residues in productive SARS-CoV-2 infections.
177
hACE2-lentivirus transduction of mouse cell lines imparts SARS-CoV-2 replication competence
To determine whether hACE2-lentivirus transduction could make mouse cell lines SARS-CoV-2
replication competent and thus available for SARS-CoV-2 research, 3T3 (mouse embryonic fibroblasts)
and AE17 (mouse lung mesothelioma cell line) cells were transduced with hACE2. Significant virus
replication was seen in transduced cells (Supplementary Figure 1), although somewhat lower than that
seen in transduced HEK293 cells. Overt CPE was not seen (Supplementary Figure 1). This illustrates that
the hACE2-lentivirus system can be used for mouse cell lines, but that the efficiency of viral replication
may be cell line or cell type dependent.
186
SARS-CoV-2 replication in hACE2-lentivirus transduced C57BL/6J, IFNAR[-/-] and IL-28RA[-/-] mice

lung
To illustrate the utility of the hACE2-lentivirus system for use in GM mice, we investigated the role of
type I and type III IFN receptor signaling in controlling SARS-CoV-2 replication. C57BL/6J, IFNAR$^{-/-}$
and IL-28RA$^{-/-}$ were inoculated i.n. with 1.2-2.2$\times10^4$ transduction units of hACE2-lentivirus, equivalent to
approximately 190-350 ng of p24 per mice (Figure 2A). An hour prior to administration of lentivirus,
mice were treated with 1% lysophosphatidylcholine (LPC) to enhance transduction efficiency (Kremer et
al., 2007; Limberis et al., 2002). One week later mice were challenged with SARS-CoV-2 (10^5 CCID$_{50}$
i.n. per mice), and lungs were collected at 2, 4 and 6 days post infection (Fig 2A). Mice did not display
overt clinical symptoms or weight loss (Figure 1B), consistent with some studies using Ad5-hACE2
(Rathnasinghe et al., 2020), but not others (Sun et al., 2020a). RT-qPCR analyses indicated similar levels
of hACE2 mRNA after transduction of mouse lungs in each of the 3 mouse strains, with expression
maintained over the 6 day course of the experiment (Figure 2C).
Lungs from infected C57BL/6J, IFNAR$^{-/-}$ and IL-28RA$^{-/-}$ were analysed for infectious virus tissue titers.
Untransduced lungs showed no virus titers (Figure 2D), whereas hACE2-lentivirus transduced lungs
showed significant viral titers ranging from 10^2 to 10^5 CCID50/g for all strains of mice on day 2, with
titers dropping thereafter (Figure 2D). Viral RNA levels were measured using RT-qPCR, with
untransduced lungs showing low and progressively decaying levels post-inoculation (Figure 2E). For all
hACE2-lentivirus transduced lungs, significantly elevated and persistent viral RNA levels were seen for
all strains and at all time points by RT-qPCR (Figure 2E). Reducing viral titers and lingering viral RNA
have been reported previously for K18-hACE2 mice (Winkler et al., 2020) and is suggested in human
infections (Jefferson et al., 2020). RNA-Seq also illustrated that virus replication in hACE2-lentivirus
transduced lungs, K18-hACE2 transgenic mice and Ad5-hACE2 transduced lungs (Sun et al., 2020a)
were comparable when normalized to hACE2 mRNA expression (Figure 2F), suggesting comparable
hACE2 translation efficiencies in these different expression systems.
Importantly, no significant differences in viral loads emerged between C57BL/6J, IFNAR$^{-/-}$ and
IL-28RA$^{-/-}$ mice (Figure 2D, E). This remained true even when viral RNA levels were normalized to
hACE2 mRNA levels (Supplementary Figure 2). Similar results were obtained by RNA-Seq, with viral
read counts not significantly different for the 3 mouse strains (Supplementary Figure 3A-D). Thus using
three different techniques, no significant effects on viral replication could be seen in mice with type I or
type III IFN receptor deficiencies.
218
SARS-CoV-2 replication in hACE2-lentivirus transduced mice lung induces interferon and cytokine
signatures by day 2 post-infection
To determine the innate response to SARS-CoV-2 replication in hACE2-transduced mouse lungs, gene
expression in lungs of infected hACE2-transduced mice was compared with that of inoculated
untransduced mice at day 2 post-infection using RNA-Seq. Differentially expressed genes (DEGs)
between these groups were identified using a false discovery rate (FDR, or q value) threshold of <0.05
(Figure 3A; Supplementary Table 1A-C). Of the 110 DEGs, 95 were upregulated, with ≈40% of these
classified as type I IFN-stimulated genes (ISGs) as determined using Interferome (using conservative
settings that only include genes validated *in vivo* for mice) (Figure 3B, full list in Supplementary Table
1D). Type III ISGs are poorly annotated in this database, and so this analysis likely under-estimates the
number of these genes.
Ingenuity Pathway Analysis (IPA) of the 110 DEGs produced a series of pro-inflammatory
Upstream Regulators (USRs) (Supplementary Table 1E-F) and included Th1, Th2 and Th17-associated
cytokine USRs, as well as type I IFN USRs (Figure 3C). SARS-CoV-2-specific T cells in humans were
reported to be predominantly Th1-assocaiated, but Th2-associated cytokines were also produced
(Weiskopf et al., 2020). A cytokine and type I IFN DEG list has been published for a RNA-Seq analysis

of SARS-CoV-2 infected K18-hACE2 transgenic mice (Winkler et al., 2020). These DEG lists were used
to interrogate the full pre-ranked (fold change) gene list (Figure 3A, Supplementary Table 1B) by Gene
Set Enrichment Analysis (GSEA). Significant enrichments were observed (Figure 3D), indicating similar
cytokine and interferon (IFN) responses after SARS-CoV-2 infection of K18-hACE2 transgenic and
hACE2-lentivirus transduced mice.
IPA *Diseases and Functions* analysis of the 110 DEGs revealed a series of annotations dominated
by cellular infiltration and innate responses signatures (Figure 3E, Supplementary Table G-H). Several of
the annotations were associated with macrophages (Figure 3E), consistent with a previous report showing
that inflammatory monocyte-macrophages (IMMs) were the major source of inflammatory cytokines in
mouse adapted SARS-CoV infected mice (Channappanavar et al., 2016). Innate immune signaling was
also the dominant signature in other analyses including GO Biological Processes and Cytoscape
(Supplementary Table 1I-J).
GSEA analysis using the >31,000 gene signatures available from the molecular signatures
database (MSigDB), indicated significant negative enrichment (negative NES scores) of gene sets
associated with translation and mitochondrial electron transport chain function (Supplementary Table
1K). Translation inhibition by SARS-CoV-2 Nsp1 blocking mRNA access to ribosomes has been
reported previously (Banerjee et al., 2020; Schubert et al., 2020). SARS-CoV-2 down-regulation of genes
associated with cellular respiration and mitochondrial function has also been reported previously (Miller
et al., 2021).
DEGs from human COVID-19 lung (Blanco-Melo et al., 2020) were also enriched in our hACE2-
lentivirus transduced mouse lungs, and analysis of the core enriched genes indicated the similarity was
due to cytokine, IFN, chemokine and inflammatory signaling genes (Figure 3F, Supplementary Table 1L-
M). A different human COVID-19 lung dataset was also used (Wu et al., 2020b) and confirmed
enrichment of immune-related DEGs in the hACE2-lentivirus transduced mice (Supplementary Table
1N). Overall this indicates that SARS-CoV-2 infection in hACE2-lentivirus transduced mice lungs
recapitulates responses seen in other murine models and in COVID-19 patients.
261
Day 6 post-infection is characterized by inflammation resolution and tissue repair signatures
To determine how response profiles progress in the hACE2-lentivirus model, RNA-Seq was used to
compare lungs on day 2 with lungs on day 6 post-infection, with day 6 broadly representing the time of
severe lung disease in the K18-hACE2 model (Winkler et al., 2020). As noted above (Figure 2D), the
virus titers had dropped in hACE2-lentivirus transduced mice lung over this time period by ≈2.8 logs, and
severe disease (typically measured by weight loss) was not evident in this hACE2-lentivirus model
(Figure 2B). In both hACE2-lentivirus transduced and untransduced C57BL/6J mice there was a clear
evolution of responses from day 2 to day 6 post-infection; however, DEGs obtained from the former were
largely distinct from DEGs obtained from the latter (Figure 4A). The RNA-Seq data thus illustrated that
the DEGs associated with virus infection were distinct from those associated with virus inoculation
(Figure 4A). RNA-Seq of infected lungs provided 551 DEGs, 401 up-regulated and 150 down-regulated
(Figure 4B, Supplementary Table 2A-C).
IPA USR analyses of the 551 DEGs showed a clear reduction in inflammatory cytokine
annotations, although the IL4 signature remained a dominant feature on day 6 (Figure 4C, Supplementary
Table 2D-E). A Th2 skew and IL-4 have both been associated with lung damage in COVID-19 patients
(Vaz de Paula et al., 2020). At day 6 there was an up-regulation of USRs primarily associated with tissue
repair (Figure 4D, Supplementary Table 2F); MAPK1 (also known as ERK) and MAPK9 (Lavoie et al.,
2020), BMP (Sountoulidis et al., 2012), TCF7L2 (Zhu et al., 2015), and KLF4 (Lin et al., 2017; Raslan
and Yoon, 2020). B and T cell-associated USRs were also seen (Figure 4D, green circles) (Duperray et
al., 1990; Nechanitzky et al., 2013; Wang et al., 2010), consistent with development of an adaptive

immune response. A relatively minor PARP1 signature was identified (Figure 4D), with PARP inhibitors being considered for COVID-19 therapy (Badawy, 2020; Curtin et al., 2020). A strong signature associated with estrogen receptor 1 (ESR1) was observed (Figure 4D), with estrogen having anti-inflammatory properties and believed to be associated with reduced COVID-19 severity in women (Jin et al., 2020; Peckham et al., 2020; Pinna, 2021), and reduced SARS-CoV disease severity in female mice (Channappanavar et al., 2016; Channappanavar et al., 2017). Follicle stimulating hormone (FSH) (Figure 4D) stimulates estrogen production, and inhibin subunit alpha (INHA) is involved in the negative feedback of FSH production (Nicks et al., 2010). IPA Diseases and Functions, Cytoscape, and GO Biological Processes analyses further supported the contention that on day 6 inflammation has abated and tissue repair is underway (Supplementary Table 2G-L).

  The >31,000 gene sets available in MSigDB were used to interrogate the complete gene list (pre-ranked by fold change) for day 6 vs day 2 (Supplementary Table 2B) by GSEA. A highly significant negative enrichment for inflammatory response, and a highly significant positive enrichment for cilium development was seen (Figure 4E). Cilia in mammals are found in the lining of the respiratory epithelium. These GSEAs thus provide further support for the conclusion above, together arguing that in the hACE2-lentivirus model, day 6 post infection is characterized by inflammation resolution and tissue repair.

Type I and III IFN signaling is required for SARS-CoV-2-induced lung inflammation
To determine what responses to SARS-CoV-2 infection are dependent on type I IFN signaling, the lentivirus system was used to analyze IFNAR$^{-/-}$ mice lung using RNA-Seq. IFNAR$^{-/-}$ and C57BL/6J mice were transduced with hACE2-lentivirus and were then infected with SARS-CoV-2 and lungs harvested for RNA-Seq on days 2 and 6 post-infection. The largest impact of type I IFN signaling deficiency was seen on day 2 post infection (Figure 5A), consistent with disease resolution on day 6 (Fig 4C-E). The total number of DEGs for IFNAR$^{-/-}$ vs C57BL/6J mouse lungs on day 2 was 192, with most of these down-regulated (Figure 5A, Supplementary Table 3A-C). Viral reads were not significantly different between these two mouse strains (Figure 5A), confirming that knocking out type I IFN signaling is insufficient to affect SARS-CoV-2 replication (Frieman et al., 2010).

  We showed in Figure 3 that RNA-Seq analysis revealed 110 DEGs associated with inflammatory responses when hACE2-transduced vs untransduced C57BL/6J mouse lungs on day 2 post infection were compared. When the same comparison was made for hACE2-lentivirus transduced vs untransduced lungs in SARS-CoV-2 infected IFNAR$^{-/-}$ mice, only 1 DEG (Kcnk5) was identified (Figure 5B). This clearly illustrated that the inflammatory responses on day 2 in C57BL/6J mice (Figure 3C) required intact type I IFN signaling.

  As a control, the number of DEGs for IFNAR$^{-/-}$ mice vs C57BL/6J in untransduced lungs was determined. Although virus replication (hACE2-lentivirus transduced lung) provided 192 DEGs, virus inoculation (untransduced lung) provided only 16 DEGs (Figure 5C). Thus again, virus inoculation had a limited effect on these analyses, with DEGs largely arising from virus infection.

  It is suggested that type III IFN acts as a first-line defense against infection at epithelial barriers, reserving the more potent and inflammatory type I IFN response in the event the type III IFN response is breached (Galani et al., 2017; Lazear et al., 2019). As type III IFN signaling has been implicated in reducing SARS-CoV-2 infection *in vitro* (Vanderheiden et al., 2020), the aforementioned analyses on day 2 was repeated in IL-28RA$^{-/-}$ mice. Viral reads were not significantly different between these two mouse strains (Figure 5D), confirming that knocking out type III IFN signaling was insufficient to affect SARS-CoV-2 replication (Boudewijns et al., 2020; Frieman et al., 2010). The DEG list for IL-28RA$^{-/-}$ vs C57BL/6J mice comprised 132 genes (Supplementary Table 4A-C), of which 84 were down-regulated (Figure 5D). Perhaps surprisingly, the DEG list was largely different from that seen in IFNAR$^{-/-}$ mice

(Figure 5D), arguing that the ISGs stimulated by these two pathways are distinct.

The majority (90%) of the 178 down-regulated DEGs in IFNAR$^{-/-}$ versus C57BL/6J mice at day 2
were ISGs according to the Interferome database (using *in vivo* and mouse only options) (Figure 5E, top
and Supplementary Table 3D). Of the 84 down-regulated genes in IL-28RA$^{-/-}$ lungs, only 27% were
identified as ISGs (Figure 5E, bottom and Supplementary Table 4D). However, there are no annotations
in the Interferome database for type III ISGs in mice, and changing the settings to include human data
(which is also under-annotated for type III IFN) did not increase the number of type III ISGs in our search
using down DEGs from IL-28RA$^{-/-}$ lungs.

To determine the pathways regulated by type I and III IFN in response to SARS-CoV-2 infection,
IPA USR analyses was performed on DEG lists from day 2 for hACE2-lentivirus transduced lungs for
IFNAR$^{-/-}$ vs C57BL/6J (192 DEGs) and IL-28RA$^{-/-}$ vs C57BL/6J (132 DEGs). IPA USR signatures
associated with pro-inflammatory cytokine signaling were significantly reduced in both IFNAR$^{-/-}$ and, to a
lesser extent, IL-28RA$^{-/-}$ mice when compared to C57BL/6J mice (Figure 5F, full lists in Supplementary
Table 3E and 4E). The same USR signatures (Figure 5F) were regulated by a largely distinct set of DEGs
(Figure 5D), possibly due to differences in signaling kinetics and cell-type responsiveness to type I vs
type III IFN (Lazear et al., 2019). Thus although there were not significant effects on viral replication,
lungs from type I and type III IFN receptor knockout mice both had a blunted pro-inflammatory response
to SARS-CoV-2 infection.

Immunization with infectious or UV-inactivated SARS-CoV-2 protects against virus infection
To evaluate the utility of the hACE2-lentivirus mouse model of SARS-CoV-2 replication for vaccination
studies, mice were immunized subcutaneously (s.c.) twice with either infectious or UV-inactivated SARS-
CoV-2 (Takasuka et al., 2004) formulated with adjuvant (Figure 6A). Serum SARS-CoV-2 neutralization
titers were determined 4-9 weeks post-boost, and revealed all mice had serum neutralization titers (Figure
6B), comparable with high end levels seen in serum samples from prior SARS-CoV-2 infected individuals
(Wajnberg et al., 2020). After challenge all mice were sacrificed on day 2 and all mice had detectable
hACE2 mRNA, with hACE2 levels higher in unvaccinated mice (Figure 6C), perhaps due to cell
infiltrates (see below) diluting hACE2 RNA levels. Mice vaccinated with both infectious and UV-
inactivated SARS-CoV-2 showed significant reductions in viral RNA levels by RT-qPCR (Figure 6D), in
tissue titrations (Figure 6E), and in RNA-Seq read counts (Figure 6F). These immunizations thus
protected mice against significant viral infections and demonstrated the utility of hACE2-lentivirus
transduced mice as a model for vaccine evaluation.

On day 2 the lungs were harvested and analyzed by RNA-Seq (all genes in Supplementary Table
5B), with 1376 DEGs identified for infectious virus immunization vs unimmunized (DEGs in
Supplementary Table 5C). The DEGs were analyzed by IPA using canonical pathways, USR, and diseases
and functions all showing infiltration signatures, dominated by T cell annotations (Supplementary Table
5D-F). This was confirmed by GSEA of human blood transcription modules (Li et al., 2014) against the
pre-ranked (by fold change) gene list for immunized vs unvaccinated mice (Supplementary Table 5G).
This showed the expected enrichment of T cell and B annotations in vaccinated mice (Figure 6F), and
illustrated that protection against virus replication in lungs of immunized mice was associated with a
significant infiltration of lymphocytes.

DISCUSSION

Here we describe a hACE2-lentivirus transduction system that allows the use of C57BL/6J and GM mice
for SARS-CoV-2/COVID-19 research. Virus titer in hACE2-lentivirus transduced mice lung were
comparable to other studies using mice transduced with Ad5 or AAV expressing hACE2 (Hassan et al.,

2020; Israelow et al., 2020; Rathnasinghe et al., 2020). These transduction methodologies have the
advantage of avoiding the brain infection seen in K18-hACE2 mice, where brain infection is the major
contributor to mortality (Carossino et al., 2021; Kumari et al., 2020). The key advantage of using
lentiviral over adenoviral systems is integration of the lentiviral payload into the host cell genome, thus
allowing long term stable expression. The development of the hACE2-lentivirus model of SARS-CoV-2
infection thus provides a new option which can be easily applied in laboratories with established lentiviral
vector systems. We provide a detailed characterization of the lentivirus-hACE2 model of SARS-CoV-2
infection in C57BL/6J mice and illustrate the use of this model to evaluate the role of type I and type III
IFNs in virus control and inflammation signatures, using IFNAR$^{-/-}$ and IL-28RA$^{-/-}$ mice. We also
illustrate the use of C57BL/6J mice transduced with lentivirus-hACE2 for SARS-CoV-2 vaccine
evaluation.
The lentivirus-hACE2 C57BL/6J mouse model of SARS-CoV-2 infection shares many cytokine
signatures with those seen in human COVID-19 patients; IL-2, IL-10, IL-6, TNFα, IL-4, IL-1β, IFNγ and
CSF3 signatures were found in SARS-CoV2-infected lentivirus-hACE2 transduced C57BL/6J mice on
day 2 post infection, and elevated levels of these cytokines are found in COVID-19 patients (Han et al.,
2020; Huang et al., 2020). Several of these cytokines have been implicated as mediators of COVID-19;
for instance, IL-6 (Herold et al., 2020; Masiá et al., 2020; Xia et al., 2020), IL-4 (Vaz de Paula et al.,
2020), IL-10 (Lu et al., 2021) and IFNγ (Gadotti et al., 2020). On day 6 post infection the lentivirus-
hACE2 C57BL/6J mouse model showed a series of signatures associated with tissue repair. Thus our
RNA-Seq analyses validate that this is a model of SARS-CoV-2 infection with acute inflammation and
subsequent resolution.
We used the hACE2-lentivirus model of SARS-CoV-2 infection in IFNAR$^{-/-}$ and IL-28RA$^{-/-}$ mice
to show that type I and III IFN signaling was involved in orchestrating the inflammatory response, while
virus replication was not altered. This is largely in line with other studies using different model systems,
which indicate that knocking out all IFN signaling (STAT1$^{-/-}$, STAT2$^{-/-}$, or combined IFNAR$^{-/-}$/IL-28RA$^{-/-}$
mice or hamsters) is required for increased virus replication (Boudewijns et al., 2020; Frieman et al.,
2010; Israelow et al., 2020; Mahlakõiv et al., 2012). This suggests that having either type I or III IFN
signaling alone is enough for partial restriction of virus replication in these mouse models. This is
supported by *in vitro* experiments that show SARS-CoV-2 is sensitive to type I and III IFN, although
infection doesn't stimulate particularly high levels of IFN (Felgenhauer et al., 2020; Vanderheiden et al.,
2020), with SARS-CoV-2 encoding proteins that inhibit IFN signaling including Orf9b (Jiang et al.,
2020a), Orf6 (Miorin et al., 2020), nsp6 and nsp13 (Xia et al., 2020). Thus while SARS-CoV-2 may be
susceptible to IFN, the role of type I and III IFN signaling in driving inflammation has implications for
proposed IFN therapies for COVID-19 patients (Monk et al., 2020).
Herein we also illustrate the use of the hACE2-lentivirus system to study the role of ACE2
residues on SARS-CoV-2 replication *in vitro*. We show that while the N31K and H353K substitutions in
mACE2 increased SARS-CoV-2 replication compared to WT mACE2, the reverse mutation in hACE2
(K31N/K353H) did not affect virus replication compared to WT hACE2. This suggests that these
residues alone are not true markers of species susceptibility or non-susceptibility, and the structure of the
RBD binding region of ACE2 as a whole likely determines SARS-CoV-2 tropism. It also suggests that
significant reductions in binding between spike RBD and ACE2 predicted *in silico* (Chan et al., 2020) do
not always translate to reduced virus replication. This suggests that even low affinity interactions may be
enough for productive virus replication, which has implications for design of hACE2 antagonists as
antivirals. Our results may also inform a different approach to 'humanized' ACE2 mice where as few as
two amino acid residues could be substituted in mACE2 to allow SARS-CoV-2 replication in mice.
In conclusion, we describe the development of a mouse model using intranasal hACE2-lentivirus
transduction to sensitize mice lung to SARS-CoV-2 infection, which recapitulated cytokine signatures

seen in other mouse models and human COVID-19. We demonstrate broad applicability of this new
mouse model in vaccine evaluation, GM mice infection, and *in vitro* evaluation of ACE2 mutants.

ACKNOWLEDGEMENTS
We thank Dr I Anraku for his assistance in managing the PC3 (BSL3) facility at QIMR Berghofer MRI.
We thank Dr Alyssa Pyke and Mr Fredrick Moore (Queensland Health, Brisbane) for providing the
SARS-CoV-2 isolate. We thank Dr. David Harrich for providing the lentivirus vectors. We thank Dr.
Simon Phipps for providing the IL-28RA$^{-/-}$ mice. We thank Monash Genome Modification Platform for
providing the plasmid containing the mouse-codon optimised human ACE2 gene. We thank Clive
Berghofer and Lyn Brazil (and many others) for their generous philanthropic donations to support SARS-
CoV-2 research at QIMR Berghofer MRI. A.S. holds an Investigator grant from the National Health and
Medical Research Council (NHMRC) of Australia (APP1173880).

434
AUTHOR CONTRIBUTIONS
Conceptualization, D.J.R.; Methodology, D.J.R. and A.S.; Formal analysis, D.J.R., A.S., and T.D.;
Investigation, D.J.R., T.L., K.Y., B.T.; Data curation, D.J.R., A.S., and T.D.; Writing – original draft,
D.J.R.; Writing – review and editing, A.S. and D.J.R.; Visualization, D.J.R., A.S., T.D., and C.B;
Supervision, D.J.R. and A.S., Project administration, D.J.R. and A.S.; Funding acquisition, A.S. and
D.J.R.

441
442 **DECLARATION OF INTERESTS**
443 The authors declare no competing interests.

444
FIGURE LEGENDS
Figure 1. Mutational analyses of human and mouse ACE2 using ACE2-lentiviruses *in vitro*. A)
Schematic of pCDH-EF1α-ACE2-BGH-PGK-GFP-T2A-Puro lentiviral vector. **B-C)** Growth kinetics of
SARS-CoV-2 over a three day time course in HEK293T cells transduced with WT or mutant hACE2 **(B)**
or mACE2 **(C)** infected at MOI=0.1. Data is the mean of 1 (mACE2-N31K/H353K) or 2 (all others)
independent experiments with 3 replicates in each and error bars represent SEM. **D)** Inverted light
microscopy images of HEK293T cells transduced with the indicated ACE2 lentivirus and infected with
SARS-CoV-2 at MOI=0.1. Images were taken at 72 hpi and were representative of triplicate wells. **E)**
Crystal structure of the Spike RBD:hACE2 complex (PBD: 6M0J) viewed in PyMOL and highlighting
key interactions (blue) between hACE2 (red) and Spike RBD (green).

455
Figure 2. SARS-CoV-2 infection of mice lung after hACE2-lentivirus intranasal transduction. A)
Timeline of lentivirus transduction of mice lung and SARS-CoV-2 infection. **B)** Percent weight loss for
mice in C-E. **C)** RT-qPCR of mice lung RNA using primers for hACE2 (introduced by lentivirus
transduction) normalized to mRPL13a levels. Data is individual mice and is expressed as RNA copy
number calculated against a standard curve for each gene. Horizontal line indicates cut-off for reliable
detection, with all untransduced mice falling below this line. **D)** Titer of SARS-CoV-2 in mice lung
determined using CCID$_{50}$ assay of lung homogenate. Horizontal line indicates the limit of detection of
1.57 $\log_{10}$ CCID50/g. **E)** RT-qPCR of mice lung RNA using primers for SARS-CoV-2 E gene
normalized to mRPL13a levels. Data is individual mice and is expressed as RNA copy number calculated
using a standard curve. See Supplementary Figure 2 for SARS-CoV-2 RNA copies normalised to hACE2
copies. **F)** SARS-CoV-2 read counts (also see Supplementary Figure 3) normalized to hACE2 read
counts from RNA-seq data for lentivirus-hACE2 transduced mice at day 2, K18-hACE2 transgenic mice
at day 4, and Ad5-hACE2 transduced mice at day 2 (Sun et al., 2020a).

Figure 3. SARS-CoV-2 infection induces cytokine responses at 2 days post-infection in hACE2-
lentivirus transduced mice. A) Volcano plot of gene expression from RNA-seq analyses of lung RNA at
day 2 comparing mice with and without hACE2-lentivirus transduction. Genes were considered DEGs if
they had a FDR value < 0.05 (above horizontal dotted line) (see Supplementary Table 1B-C for full gene
and DEG lists). **B)** Interferome analysis of 95 up DEGs from 'A'. 38 of 95 up DEGs (40%) were ISGs.
Red = type I IFN, blue = type II IFN, green = type III IFN (see Supplementary Table 1D for full
Interferome ISG list). **C)** Cytokine signatures identified by IPA USR analysis (see Supplementary Table
1E-F for full and cytokine only USR lists) of 110 DEGs identified in 'A'. Circle area reflects the number
of DEGs associated with each USR annotation. USRs were grouped into three categories; red = Th1
associated, green = Th2 associated, blue = Th17 associated, and black = pleiotropic/other. The vertical
dotted line indicates activation z-score of 0. **D)** GSEAs for cytokine-related DEGs or type I IFN-related
DEGs from Winkler et al. supplemental data (Winkler et al., 2020) against the pre-ranked all gene list
(Supplementary Table 1B). Normalised enrichment score (NES) and nominal p-value are shown. **E)** IPA
diseases and functions analysis (see Supplementary Table 1G-H for full lists) of 110 DEGs identified in
'A'. The 23 annotations with the most significant p-value and with an activation z-score of >2 were
plotted with dots/lines indicating activation z-score and bars indicating $-\log_{10}$ p-value. Annotations were
grouped into two categories; cellular infiltration (blue) and innate responses (red). **F)** GSEA for DEGs
from human COVID-19 lung vs healthy control from Blanco-Melo et al. (Blanco-Melo et al., 2020)
against the pre-ranked (by fold change) all gene list (Supplementary Table 1B). Normalised enrichment
score (NES) and nominal p-value are shown. Core enriched genes (see Supplementary Table 1L for full
core enriched gene list) determined by GSEA were entered into Enrichr and the top 10 GO Biological
Processes annotations sorted by p-value are shown (see Supplementary Table 1M for full GO processes
list).

Figure 4. Tissue repair signatures dominate at day 6 post-infection in hACE2-lentivirus transduced
mice. A) Venn diagram for DEGs comparing day 6 vs 2 post infection with (red) and without (green)
hACE2-lentivirus transduction. Number of DEGs that are FDR<0.05 are shown. Mean virus titer ($\log_{10}$
$CCID_{50}/g$) in lung tissue ± SEM (represents data shown in Figure 2D) is shown for hACE2 transduced
mice at day 2 or 6 post infection. **B)** Volcano plot of gene expression from RNA-seq analyses of hACE2-
lentivirus transduced lung RNA comparing day 6 with day 2 (see Supplementary Table 2B-C for gene and
DEG lists). Genes were considered DEGs if they had a FDR value < 0.05 (above horizontal dotted line).
C) Cytokine signatures identified by IPA USR analysis (see Supplementary Table 2D-E for full and
cytokine only lists) of 551 DEGs identified in 'B' (Supplementary Table 2C). Circle area reflects the
number of DEGs associated with each USR annotation. USRs were grouped into four categories; red =
Th1 associated, green = Th2 associated, blue = Th17 associated, and black = pleiotropic/other. The
vertical dotted line indicates activation z-score of 0. **D)** Upregulated USR signatures identified using IPA
analysis (see Supplementary Table 2F) of 551 DEGs identified in 'B'. Circle area reflects the number of
DEGs associated with each USR annotation. USRs (excluding chemicals) with an activation z-score of >2
are shown and were grouped into five categories; green = B cell function, blue = tissue repair – endocrine
signaling, orange = tissue repair – growth factor, red = tissue repair – kinase, purple = tissue repair – Wnt
signaling. **E)** Selected plots from GSEA of entire MSigDB database using pre-ranked (by fold change)
'all' gene list (Supplementary Table 2B). NES score and p-value is shown for the
"GO_inflammatory_response" and
"Wp_Genes_Related_To_Primary_Cilium_Development_Based_On_Crispr" annotations.

Figure 5. Type I and III IFN signaling is required for the cytokine response in C57BL/6J mice. A)

Venn diagram for number of DEGs (FDR<0.05) between IFNAR[-/-] vs C57BL/6J mice at day 2 or day 6 (green). Mean $\log_{10}$ SARS-CoV-2 read count in RNA-seq data ± SEM is shown (red text = day 2, green text = day 6). **B)** Venn diagram for number of DEGs (FDR<0.05) between plus vs minus hACE2-lentivirus transduction comparing C57BL/6J (red) and IFNAR[-/-] (green) mice at day 2. **C)** Venn diagram for IFNAR[-/-] vs C57BL/6J mice at day 2 comparing hACE2-lentivirus transduced (red) or untransduced (green) mice. **D)** Venn diagram comparing IFNAR[-/-] vs C57BL/6J (red) and IL-28RA[-/-] vs C57BL/6J (blue) (see Supplementary Table 3B-C and 4B-C for full gene and DEG lists) mice at day 2 (all mice had received hACE2-lentivirus transduction). Mean $\log_{10}$ SARS-CoV-2 read count in RNA-seq data ± SEM is shown (red text = IFNAR[-/-], black text = C57BL/6J, green text = IL-28RA[-/-]). **E)** Interferome analysis of down DEGs from 'D'. See supplementary Table 3D and 4D for full Interferome ISG list. **F)** Signatures identified using IPA USR analysis of DEGs identified in 'D' for IFNAR[-/-] vs C57BL/6J (red) and IL-28RA[-/-] vs C57BL/6J (blue). Circle area reflects the number of DEGs associated with each USR annotation. Signatures were selected focusing on data previously identified in 'Figure 3C' (entire list is in Supplementary Table 3E and 4E).

Figure 6. Immunization protects against SARS-CoV-2 replication in hACE2-transduced mice lung and induces T and B cell responses. A) Timeline of mice vaccination, lentivirus transduction and SARS-CoV-2 infection. **B)** Reciprocal 50% neutralization titers of naïve mice and infectious or UV SARS-CoV-2 vaccinated mice. Horizontal dotted line represents the limit of detection (1 in 10). **C)** RT-qPCR of mice lung RNA using primers for hACE2 introduced by lentivirus transduction normalized to mRPL13a levels. Data is individual mice and is expressed as RNA copy number calculated against a standard curve for each gene. Horizontal line indicates cut-off for reliable detection, with all hACE2-negative mice falling below this line. **D)** RT-qPCR of mice lung RNA using primers for SARS-CoV-2 E gene normalized to mRPL13a levels. Data is individual mice and is expressed as RNA copy number calculated against a standard curve for each gene. **E)** Titer of SARS-CoV-2 in mice lung determined using CCID50 assay of lung homogenate. Horizontal line indicates the limit of detection of 1.57 $\log_{10}$ CCID50/g. **F)** Mean $\log_{10}$ SARS-CoV-2 read count at day 2 post-infection in RNA-seq data ± SEM is shown (blue text = unvaccinated mice, red text = infectious SARS-CoV-2 immunized mice). GSEA for Blood Transcription Modules (BTMs) from Li et al. (Li et al., 2014) against the pre-ranked all gene list comparing unvaccinated vs infectious SARS-CoV-2 immunized (s.c) mice lung at day 2 (all mice had hACE2-lentivirus transduction). Selected BTM modules with a positive NES score and $p < 0.05$ are shown (full list in Supplementary Table 5). See Supplementary Table 5 for full gene and DEG lists and full downstream analyses lists.

STAR METHODS
Ethics statement
All mouse work was conducted in accordance with the "Australian code for the care and use of animals for scientific purposes" as defined by the National Health and Medical Research Council of Australia. Mouse work was approved by the QIMR Berghofer Medical Research Institute animal ethics committee (P3600, A2003-607). Mice were euthanized using CO_2 or cervical dislocation.

Cell lines and SARS-CoV-2 culture
Vero E6 (C1008, ECACC, Wiltshire, England; Sigma Aldridge, St. Louis, MO, USA), HEK293T, AE17, and NIH-3T3 cells were cultured in medium comprising RPMI1640 (Gibco) supplemented with 10% fetal calf serum (FCS), penicillin (100 IU/ml)/streptomycin (100 µg/ml) (Gibo/Life Technologies) and L-glutamine (2 mM) (Life Technologies). Cells were routinely checked for mycoplasma (MycoAlert Mycoplasma Detection Kit MycoAlert, Lonza) and FCS was assayed for endotoxin contamination before

purchase (Johnson et al., 2005). The SARS-CoV-2 isolate was kindly provided by Queensland Health
Forensic & Scientific Services, Queensland Department of Health, Brisbane, Australia. The virus (hCoV-
19/Australia/QLD02/2020) was isolated from a patient and sequence deposited at GISAID
(https://www.gisaid.org/; after registration and login, sequence can be downloaded from
https://www.epicov.org/epi3/frontend#1707af). Virus stock was generated by infection of Vero E6 cells
at multiplicity of infection (MOI)≈0.01, with supernatant collected after 2-3 days, cell debris removed by
centrifugation at 3000 x g for 15 min at 4°C, and virus aliquoted and stored at -80 °C. Virus titers were
determined using standard $CCID_{50}$ assays (see below). The virus was determined to be mycoplasma free
using co-culture with a non-permissive cell line (i.e. HeLa) and Hoechst staining as described (La Linn et
al., 1995).
573
$CCID_{50}$ assays
Vero E6 cells were plated into 96 well flat bottom plates at $2x10^4$ cells per well in 100 µl of medium. For
tissue titer, tissue was homogenized in tubes each containing 4 ceramic beads twice at 6000 x g for 15
seconds, followed by centrifugation twice at 21000 x g for 5 min before 5 fold serial dilutions in 100 µl
RPMI1640 supplemented with 2% FCS. For cell culture supernatant, 10 fold serial dilutions were
performed in 100 µl RPMI1640 supplemented with 2% FCS. 100 µl of serially diluted samples were
added to Vero E6 cells and the plates cultured for 5 days at 37°C and 5% CO_2. The virus titer was
determined by the method of Spearman and Karber (a convenient Excel $CCID_{50}$ calculator is available at
https://www.klinikum.uni-heidelberg.de/zentrum-fuer-infektiologie/molecular-
virology/welcome/downloads).
584
Lentivirus cloning
ACE2 genes were cloned into pCDH-EF1α-MCS-BGH-PGK-GFP-T2A-Puro Cloning and Expression
Lentivector (System Biosciences, catalogue number CD550A-1), where the EF1α promoter was replaced
with the full length EF1α using NheI and ClaI restriction enzymes (New England Biolabs). Human ACE2
coding sequence (codon optimized for mouse) was cloned into the pCDH lentivector using using PCR
fragments amplified using Q5® High-Fidelity 2X Master Mix (New England Biolabs) and the following
primers; vector backbone (pCDH amplified with Forward 5'- TAAATCGGATCCGCGG -3' and Reverse
5'- AATTCGAATTCGCTAGC) and hACE2 insert (Forward 5'-
GCTAGCGAATTCGAATTATGAGCAGCAGCTCTTGGC -3' and Reverse 5'-
CCGCGGATCGCATTTATCAGAAGCTTGTCTGCACGT -3'). The pCDH fragment was digested with
DpnI (New England Biolabs) and was purified using QIAquick Gel Extraction Kit (QIAGEN), as was the
hACE2 fragment. Fragments were recombined using NEBuilder HiFi DNA Assembly Cloning Kit as per
manufacturer instructions. This was transformed into NEB® 10-beta Competent E. coli (High Efficiency)
(New England Biolabs) as per manufacturer instructions and plated on ampicillin agar plates overnight.
Colony PCR was performed using Q5 High-Fidelity 2X Master Mix (New England Biolabs) with the
hACE2 insert primers to identify a positive colony, which was grown in LB broth with ampicillin and
plasmid was purified using NucleoBond Xtra Midi kit (Machery Nagel).
All ACE2 mutant coding sequences were ordered from Twist Bioscience/Decode Science containing
EcoRI upstream and BamHI downstream. Fragments were amplified using the using Q5® High-Fidelity
2X Master Mix (New England Biolabs) and the following primers; Forward 5'-
CCTGACCTTAGCGAATTCATG -3' and Reverse 5'- ACCTAGCCTCGCGGATC -3'. PCR fragments
were purified using Monarch DNA Gel Extraction Kit (New England Biolabs), and were digested with
EcoRI and BamHI (New England Biolabs), as was the pCDH vector before purification again with
Monarch® DNA Gel Extraction Kit (New England Biolabs). The ACE2 fragment was ligated with
pCDH vector using T4 DNA Ligase (New England Biolabs) as per manufacturer instructions and then

transformed into NEB® 10-beta Competent E. coli (High Efficiency) (New England Biolabs) as per
manufacturer instructions and plated on ampicillin agar plates overnight. Colonies were grown in LB
broth with ampicillin and plasmid was purified using NucleoBond Xtra Midi kit (Machery Nagel).
Plasmids were confirmed by PCR with the cloning primers above and fragments were gel purified and
were confirmed by Sanger sequencing with the forward primer.
615
Lentivirus production, titration and cell line transduction
ACE2 lentivirus was produced by co-transfection of HEK293T cells with the pCDH-ACE2 plasmid,
VSV-G and Gag-Pol using Lipofectamine 2000 Reagent (Thermo Fisher Scientific) or Xfect Transfection
Reagent (Takara Bio) as per manufacturer instructions. Supernatant was collected 2 days after
transfection and centrifuged at 500 x g for 10 min, and this was concentrated using Amicon Ultra-15
Centrifugal Filter Units with 100 kDa cutoff (Merck Millipore) as per manufacturer instructions.
Lentivirus was titrated by serial dilution of lentivirus and incubating with 5000 HEK293T cells in 96 well
plates with 8 µg/ml polybrene for 3 days followed by GFP detection by flow cytometry (BD Biosciences
LSRFortessa). Transduction units (TU) per ml was calculated by percentage of 5000 cells that are GFP
positive multiplied by the dilution. For example if 2 µl lentivirus gives 5% GFP positive cells, the TU/ml
is ((5/100)*5000) * (1000/2) = 125,000 TU/ml. The p24 equivalent (ng/ml) was measured using Lenti-X
GoStix Plus (Takara Bio) as per manufacturer instructions.
Cell lines were transduced by incubating with lentivirus and 8 µg/ml polybrene for 2-3 days followed by 5
µg/ml puromycin treatment until most resistant cells expressed GFP. Cells were then infected with
SARS-CoV-2 at MOI 0.1 for 1 hr at 37◦C, cells were washed with PBS and media replaced. Culture
supernatant was harvested at the indicated timepoints and titered by CCID50 assay as described.
632
UV-inactivated SARS-CoV-2 vaccine preparation and mice vaccination
120 ml of SARS-CoV-2 was inactivated with at least 7650 J/m^2 UVC in a UVC 500 Ultraviolet
Crosslinker (Hoefer). Virus was determined to be inactivated by incubation with Vero E6 cells for 4 days
without development of CPE. UV-inactive SARS-CoV-2 was then partially-purified using a 20% sucrose
cushion centrifuged at 175,000 x g for 3-4 hr using SW32Ti rotor (Beckman Coulter). The sample that
has passed through the sucrose was collected and concentrated using Amicon Ultra-15 Centrifugal Filter
Units with 100 kDa cutoff (Merck Millipore). For infectious SARS-CoV-2 immunization, virus was also
concentrated using Amicon Ultra-15 Centrifugal Filter Units with 100 kDa cutoff (Merck Millipore).
Mice were injected s.c. in the base of the tail with 100 ul of UV-inactive or infectious SARS-CoV-2 with
25 ug in 50 ul adjuvant made with the same ingredients as AS01 (Chlibek et al., 2013) as described
previously (Yan et al., 2020). A boost was given 5-6 weeks after prime, and mice were bled to measure
serum neutralizing titers 4-9 weeks after boost.
645
Neutralization assay
Mouse serum was heat inactivated at 56◦C for 30 min and incubated with 100 CCID50 SARS-CoV-2 for
2 hr at 37◦C before adding 10^5 Vero cells/well in a 96 well plate to 200 µl. After 4 days cells were fixed
and stained by adding 50 µl formaldehyde (15% w/v) and crystal violet (0.1% w/v) (Sigma-Aldrich)
overnight. The plates were washed in tap water, dried overnight and 100 µl/well of 100% methanol added
to dissolve the crystal violet and the OD was read at 595 nm using a 96-well plate reader (Biotek Synergy
H4). The 50% neutralizing titers were interpolated from optical density (OD) versus dilution plots.

Mice intranasal lentivirus transduction and SARS-CoV-2 infection
C57BL/6J, IFNAR$^{-/-}$ (Swann et al., 2007) (originally provided by P. Hertzog, Monash University,
Melbourne, VIC, Australia) and IL-28RA$^{-/-}$ (Ank et al., 2008; Galani et al., 2017) mice were bred in-house

at QIMRB. Female mice were 8 weeks to 1 year old (age matched between groups) at the start of the
experiment. Mice were anesthetized using isoflurane and 4 µl of 1% L-α-Lysophosphatidylcholine from
egg yolk (Sigma Aldrich) in water was administered intranasally. After 1 hour mice were inoculated
intranasally with approximately 2×10^4 TU of hACE2-pCDH lentivirus in 50 µl, and 1 week later
challenged with 10^5 CCID50 SARS-CoV-2 intranasally in 50 µl. Mice were sacrificed by cervical
dislocation at day 2, 4 or 6 and lungs were collected. Superior lobe, middle lobe and inferior lobe of the
right lung was immediately homogenized in tubes each containing 4 beads twice at 6000 x g for 15
seconds, and used in tissue titration as described above. Left lung was placed in RNAprotect Tissue
Reagent (QIAGEN) at 4∘C overnight then -80∘C. The post-caval lobe of the right lung was fixed in 10%
formalin for 4-5 days before transfer to 70% EtOh for histology.
K18-hACE2 mice (McCray et al., 2007) were purchased from The Jackson Laboratory and bred in-house
at QIMRB with C57BL/6J mice. Mice were genotyped using Extract-N-Amp Tissue PCR Kit (Sigma
Aldrich) according to manufacturer instructions with the following primers; Forward 5'-
CTTGGTGATATGTGGGGTAGA-3' and Reverse 5'-CGCTTCATCTCCCACCACTT-3'. hACE2
positive mice were infected with 5×10^4 SARS-CoV-2 i.n. as above and lung RNA was harvested at day 4
for RNA-seq.

RT-qPCR
Mice lung was transferred from RNAlater to TRIzol (Life Technologies) and was homogenized twice at
6000 x g for 15 sec. Homogenates were centrifuged at $14,000 \times g$ for 10 min and RNA was isolated as
per manufacturer's instructions. cDNA was synthesized using ProtoScript II First Strand cDNA Synthesis
Kit (New England Biolabs) and qPCR performed using iTaq Universal SYBR Green Supermix (Bio-Rad)
as per manufacturer instructions with the following primers; SARS-CoV-2 E Forward 5'-
ACAGGTACGTTAATAGTTAATAGCGT -3' and Reverse 5'- ATATTGCAGCAGTACGCACACA,
hACE2 Forward 5'- GATCACGATTCCCAGGACG -3' and Reverse 5'- TCCGGCTGAACGACAACTC
-3', mRPL13a (Schroder et al., 2010) Forward 5'- GAGGTCGGGTGGAAGTACCA -3' and Reverse 3'-
TGCATCTTGGCCTTTTCCTT -3'. PCR fragments of SARS-CoV-2 E, ACE2 and mRPL13a using the
same primers as above were gel purified and 10-fold serial dilutions of estimated copy numbers were used
as standards in qPCR to calculate copies in samples reactions. SARS-CoV-2 E and ACE2 copies were
normalized by mRPL13a copy number in each reaction. qPCR reactions were performed in duplicate and
averaged to determine the copy number in each reaction.

RNAseq
TRIzol extracted lung RNA was treated with DNase (RNAseq-Free DNAse Set (Qiagen)) followed by
purification using RNeasy MinElute Cleanup Kit (QIAGEN) as per manufacturer instructions. RNA
concentration and quality was measured using TapeStation D1K TapeScreen assay (Agilent). cDNA
libraries were prepared using the Illumina TruSeq Stranded mRNA library prep kit and the sequencing
performed on the Illumina Nextseq 550 platform generating 75bp paired end reads. Per base sequence
quality for >90% bases was above Q30 for all samples. The quality of raw sequencing reads was assessed
using FastQC (Simons, 2010) (v0.11.8) and trimmed using Cutadapt (Martin, 2011) (v2.3) to remove
adapter sequences and low-quality bases. Trimmed reads were aligned using STAR (Dobin et al., 2013)
(v2.7.1a) to a combined reference that included the mouse GRCm38 primary assembly and the
GENCODE M23 gene model (Harrow et al., 2012), SARS-CoV-2 isolate Wuhan-Hu-1 (NC_045512.2;
29903 bp) and the human ACE2 mouse codon optimized sequence (2418 bp). Mouse gene expression
was estimated using RSEM (Li and Dewey, 2011) (v1.3.0). Reads aligned to SARS-CoV-2 and hACE2
were counted using SAMtools (Li et al., 2009) (v1.9). Differential gene expression in the mouse was
analyzed using EdgeR (3.22.3) and modelled using the likelihood ratio test, glmLRT().

Analyses of K18-hACE2 and Ad5-hACE2 RNA-seq data

RNA-seq datasets generated from the Winkler et al. study (Winkler et al., 2020), and Sun et al. study (Sun et al., 2020a) were obtained from the Gene Expression Omnibus (GSE154104 and GSE150847 respectively) and trimmed using Cutadapt (v2.3). Trimmed reads were aligned using STAR (v2.7.1a) to a combined reference that included the mouse GRCm38 primary assembly and the GENCODE M23 gene model, SARS-CoV-2 isolate Wuhan-Hu-1 (NC_045512.2; 29903 bp) and the human ACE2 transcript variant 1 (NM_001371415.1; 3339 bp). Mouse gene expression was estimated using RSEM (v1.3.0). Reads aligned to SARS-CoV-2 and hACE2 were counted using SAMtools (v1.9). Differential gene expression in the mouse was analyzed using EdgeR (3.22.3) and modelled using the likelihood ratio test, glmLRT().

Pathway Analysis

Up-Stream Regulators (USR), Diseases and Functions and canonical pathways enriched in differentially expressed genes in direct and indirect interactions were investigated using Ingenuity Pathway Analysis (IPA) (QIAGEN).

Network Analysis

Protein interaction networks of differentially expressed gene lists were visualized in Cytoscape (v3.7.2) (Shannon et al., 2003). Enrichment for biological processes, molecular functions, KEGG pathways and other gene ontology categories in DEG lists was elucidated using the STRING database (Szklarczyk et al., 2019) and GO enrichment analysis (Mi et al., 2019).

Gene Set Enrichment Analysis

Preranked GSEA (Subramanian et al., 2005) was performed on a desktop application (GSEA v4.0.3) (http://www.broadinstitute.org/gsea/) using the "GSEAPreranked" module. Gene sets from the supplemental materials from the Winkler et al. (Winkler et al., 2020), Blanco-Melo et al. (Blanco-Melo et al., 2020) and Wu et al. (Wu et al., 2020b) (filename.GMT) studies were investigated for enrichment in our pre-ranked all gene list for SARS-CoV-2 infected day 2 mice lung comparing plus versus minus ACE2 (Supplementary Table 2B). Li et al. (Li et al., 2014) blood transcription modules (BTM_for_GSEA_20131008.gmt, n = 346) was also used in GSEA to determine enrichment in our 'vaccinated' versus unvaccinated mice lung at day 2 (Supplementary Table 5B). The complete Molecular Signatures Database (MSigDB) v7.2 gene set collection (31,120 gene sets) (msigdb.v7.2.symbols.gmt: https://www.gsea-msigdb.org/gsea/msigdb/download_file.jsp?filePath=/msigdb/release/7.2/msigdb.v7.2.symbols.gmt) was used to run GSEAs on pre-ranked gene list for plus versus minus ACE2 at day 2 (Supplementary Table 2B). Gene ontology (Ashburner et al., 2000; Consortium, 2021) and Enrichr (Chen et al., 2013; Kuleshov et al., 2016) web-tools were also consulted.

Interferome

The indicated DEG lists were entered into Interferome (www.interferome.org) (Rusinova et al., 2013) with the following parameters: *In Vivo*, Mus musculus, fold change 2 (up and down).

SUPPPLEMENTAL INFORMATION

Supplementary Figure 1. Replication of SARS-CoV-2 in hACE2-lentivirus transduced mouse cell lines. Growth kinetics of SARS-CoV-2 over a three day time course in untransduced or hACE2-lentivirus transduced 3T3 or AE17 cells infected at MOI=0.1. Data is the mean of triplicate wells and error bars

750 represent SEM. Images of cells were taken using an inverted light microscope at day 3 post-infection and
751 are representative of triplicate wells.
752
753 **Supplementary Figure 2. RT-qPCR of mice lung RNA using primers for SARS-CoV-2 normalized**
754 **to hACE2 introduced by lentivirus transduction.** Data is individual mice and is expressed as RNA
755 copy number calculated against a standard curve for each gene.
756
Supplementary Figure 3. SARS-CoV-2 and hACE2 read counts in RNA-Seq data. RNA-Seq was
performed on mice lung from the same mice as in 'Figure 2'. **A)** SARS-CoV-2 read counts normalized to
total read count. Data points below the horizontal dotted line had read counts of zero. **B)** hACE2 read
counts normalised to total read count. Data points below the horizontal dotted line had read counts of
zero. **C)** SARS-CoV-2 read count normalised to hACE2 read count. Circle with pink dotted outline had
hACE2 read count of 0, so the value was set to the SARS-CoV-2 read count (not normalised). **D)** SARS-
CoV-2 reads aligned to reference genome viewed in Integrative Genome Viewer (IGV). Mice with
hACE2 displayed reads mapped across the entire genome, with higher counts for structural gene sub-
genomic RNA as also evident in hACE2-adenoviral vector transduced cell lines (Blanco-Melo et al.,
2020).
767
768 **Supplementary Table 1. RNA-Seq gene lists and downstream bioinformatic analyses of responses**
769 **at day 2 post-infection.**
770
771 **Supplementary Table 2. RNA-Seq gene lists and downstream bioinformatic analyses of responses**
772 **at day 6 post-infection.**
773
774 **Supplementary Table 3. RNA-Seq gene lists and downstream bioinformatic analyses of responses**
775 **in IFNAR$^{-/-}$ vs C57BL/6J mice at day 2 post-infection.**
776
777 **Supplementary Table 4. RNA-Seq gene lists and downstream bioinformatic analyses of responses**
778 **in IL-28RA$^{-/-}$ vs C57BL/6J mice at day 2 post-infection.**
779
780 **Supplementary Table 5. RNA-Seq gene lists and downstream bioinformatic analyses of responses**
781 **in immunized vs unvaccinated mice at day 2 post-infection.**
782
783 **REFERENCES**
784

Alexander, M.R., Schoeder, C.T., Brown, J.A., Smart, C.D., Moth, C., Wikswo, J.P., Capra, J.A., Meiler, J.,
Chen, W., and Madhur, M.S. (2020). Predicting susceptibility to SARS-CoV-2 infection based on structural
differences in ACE2 across species. The FASEB Journal *34*, 15946-15960.
Alton, E.W., Beekman, J.M., Boyd, A.C., Brand, J., Carlon, M.S., Connolly, M.M., Chan, M., Conlon, S.,
Davidson, H.E., Davies, J.C.*, et al.* (2017a). Preparation for a first-in-man lentivirus trial in patients with
cystic fibrosis. Thorax *72*, 137-147.
Alton, E.W.F.W., Beekman, J.M., Boyd, A.C., Brand, J., Carlon, M.S., Connolly, M.M., Chan, M., Conlon, S.,
Davidson, H.E., Davies, J.C.*, et al.* (2017b). Preparation for a first-in-man lentivirus trial in patients with
cystic fibrosis. Thorax *72*, 137.
Ank, N., Iversen, M.B., Bartholdy, C., Staeheli, P., Hartmann, R., Jensen, U.B., Dagnaes-Hansen, F.,
Thomsen, A.R., Chen, Z., Haugen, H.*, et al.* (2008). An Important Role for Type III Interferon (IFN-λ/IL-28)
in TLR-Induced Antiviral Activity. The Journal of Immunology *180*, 2474-2485.

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight,
S.S., Eppig, J.T., *et al.* (2000). Gene ontology: tool for the unification of biology. The Gene Ontology
Consortium. Nature genetics *25*, 25-29.
Badawy, A.A.B. (2020). Immunotherapy of COVID-19 with poly (ADP-ribose) polymerase inhibitors:
starting with nicotinamide. Biosci Rep *40*, BSR20202856.
Banerjee, A.K., Blanco, M.R., Bruce, E.A., Honson, D.D., Chen, L.M., Chow, A., Bhat, P., Ollikainen, N.,
Quinodoz, S.A., Loney, C., *et al.* (2020). SARS-CoV-2 Disrupts Splicing, Translation, and Protein Trafficking
to Suppress Host Defenses. Cell *183*, 1325-1339.e1321.
Bao, L., Deng, W., Huang, B., Gao, H., Liu, J., Ren, L., Wei, Q., Yu, P., Xu, Y., Qi, F., *et al.* (2020). The
pathogenicity of SARS-CoV-2 in hACE2 transgenic mice. Nature *583*, 830-833.
Blanco-Melo, D., Nilsson-Payant, B.E., Liu, W.C., Uhl, S., Hoagland, D., Møller, R., Jordan, T.X., Oishi, K.,
Panis, M., Sachs, D., *et al.* (2020). Imbalanced Host Response to SARS-CoV-2 Drives Development of
COVID-19. Cell *181*, 1036-1045.e1039.
Boudewijns, R., Thibaut, H.J., Kaptein, S.J.F., Li, R., Vergote, V., Seldeslachts, L., Van Weyenbergh, J., De
Keyzer, C., Bervoets, L., Sharma, S., *et al.* (2020). STAT2 signaling restricts viral dissemination but drives
severe pneumonia in SARS-CoV-2 infected hamsters. Nature Communications *11*, 5838.
Buckley, S.M.K., Howe, S.J., Sheard, V., Ward, N.J., Coutelle, C., Thrasher, A.J., Waddington, S.N., and
McKay, T.R. (2008). Lentiviral transduction of the murine lung provides efficient pseudotype and
developmental stage-dependent cell-specific transgene expression. Gene Therapy *15*, 1167-1175.
Carossino, M., Montanaro, P., O'Connell, A., Kenney, D., Gertje, H., Grosz, K.A., Kurnick, S.A., Bosmann,
M., Saeed, M., Balasuriya, U.B.R., *et al.* (2021). Fatal neuroinvasion of SARS-CoV-2 in K18-hACE2 mice is
partially dependent on hACE2 expression. bioRxiv.
Chan, K.K., Dorosky, D., Sharma, P., Abbasi, S.A., Dye, J.M., Kranz, D.M., Herbert, A.S., and Procko, E.
(2020). Engineering human ACE2 to optimize binding to the spike protein of SARS coronavirus 2. Science
(New York, NY) *369*, 1261-1265.
Channappanavar, R., Fehr, Anthony R., Vijay, R., Mack, M., Zhao, J., Meyerholz, David K., and Perlman, S.
(2016). Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses Cause
Lethal Pneumonia in SARS-CoV-Infected Mice. Cell host & microbe *19*, 181-193.
Channappanavar, R., Fett, C., Mack, M., Ten Eyck, P.P., Meyerholz, D.K., and Perlman, S. (2017). Sex-
Based Differences in Susceptibility to Severe Acute Respiratory Syndrome Coronavirus Infection. J
Immunol *198*, 4046-4053.
Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and Ma'ayan, A. (2013).
Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. BMC Bioinformatics *14*,
128.
Chlibek, R., Bayas, J.M., Collins, H., de la Pinta, M.L.R., Ledent, E., Mols, J.F., and Heineman, T.C. (2013).
Safety and Immunogenicity of an AS01-adjuvanted Varicella-zoster Virus Subunit Candidate Vaccine
Against Herpes Zoster in Adults ≥50 Years of Age. The Journal of Infectious Diseases *208*, 1953-1961.
Consortium, G.O. (2021). The Gene Ontology resource: enriching a GOld mine. Nucleic Acids Res *49*,
D325-d334.
Copreni, E., Palmieri, L., Castellani, S., and Conese, M. (2010). A VSV-G Pseudotyped Last Generation
Lentiviral Vector Mediates High Level and Persistent Gene Transfer in Models of Airway Epithelium In
Vitro and In Vivo. Viruses *2*, 1577-1588.
Copreni, E., Penzo, M., Carrabino, S., and Conese, M. (2004). Lentivirus-mediated gene transfer to the
respiratory epithelium: a promising approach to gene therapy of cystic fibrosis. Gene Therapy *11*, S67-
S75.
Crawford, K.H.D., Eguia, R., Dingens, A.S., Loes, A.N., Malone, K.D., Wolf, C.R., Chu, H.Y., Tortorici, M.A.,
Veesler, D., Murphy, M., *et al.* (2020). Protocol and Reagents for Pseudotyping Lentiviral Particles with
SARS-CoV-2 Spike Protein for Neutralization Assays. Viruses *12*.

Curtin, N., Bányai, K., Thaventhiran, J., Le Quesne, J., Helyes, Z., and Bai, P. (2020). Repositioning PARP
inhibitors for SARS-CoV-2 infection(COVID-19); a new multi-pronged therapy for acute respiratory
distress syndrome? British Journal of Pharmacology *177*, 3635-3645.
Damas, J., Hughes, G.M., Keough, K.C., Painter, C.A., Persky, N.S., Corbo, M., Hiller, M., Koepfli, K.-P.,
Pfenning, A.R., Zhao, H.*, et al.* (2020). Broad host range of SARS-CoV-2 predicted by comparative and
structural analysis of ACE2 in vertebrates. Proceedings of the National Academy of Sciences *117*, 22311-
22322.
Dinnon, K.H., Leist, S.R., Schäfer, A., Edwards, C.E., Martinez, D.R., Montgomery, S.A., West, A., Yount,
B.L., Hou, Y.J., Adams, L.E.*, et al.* (2020). A mouse-adapted model of SARS-CoV-2 to test COVID-19
countermeasures. Nature *586*, 560-566.
Dobin, A., Davis, C.A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., and
Gingeras, T.R. (2013). STAR: ultrafast universal RNA-seq aligner. Bioinformatics *29*, 15-21.
Duperray, C., Boiron, J.M., Boucheix, C., Cantaloube, J.F., Lavabre-Bertrand, T., Attal, M., Brochier, J.,
Maraninchi, D., Bataille, R., and Klein, B. (1990). The CD24 antigen discriminates between pre-B and B
cells in human bone marrow. J Immunol *145*, 3678-3683.
Felgenhauer, U., Schoen, A., Gad, H.H., Hartmann, R., Schaubmar, A.R., Failing, K., Drosten, C., and
Weber, F. (2020). Inhibition of SARS-CoV-2 by type I and type III interferons. J Biol Chem *295*, 13958-
13964.
Frieman, M.B., Chen, J., Morrison, T.E., Whitmore, A., Funkhouser, W., Ward, J.M., Lamirande, E.W.,
Roberts, A., Heise, M., Subbarao, K.*, et al.* (2010). SARS-CoV Pathogenesis Is Regulated by a STAT1
Dependent but a Type I, II and III Interferon Receptor Independent Mechanism. PLOS Pathogens *6*,
e1000849.
Gadotti, A.C., de Castro Deus, M., Telles, J.P., Wind, R., Goes, M., Garcia Charello Ossoski, R., de Padua,
A.M., de Noronha, L., Moreno-Amaral, A., Baena, C.P.*, et al.* (2020). IFN-γ is an independent risk factor
associated with mortality in patients with moderate and severe COVID-19 infection. Virus Res *289*,
198171-198171.
Galani, I.E., Triantafyllia, V., Eleminiadou, E.E., Koltsida, O., Stavropoulos, A., Manioudaki, M., Thanos, D.,
Doyle, S.E., Kotenko, S.V., Thanopoulou, K.*, et al.* (2017). Interferon-λ Mediates Non-redundant Front-
Line Antiviral Protection against Influenza Virus Infection without Compromising Host Fitness. Immunity
46, 875-890.e876.
Golden, J.W., Cline, C.R., Zeng, X., Garrison, A.R., Carey, B.D., Mucker, E.M., White, L.E., Shamblin, J.D.,
Brocato, R.L., Liu, J.*, et al.* (2020). Human angiotensin-converting enzyme 2 transgenic mice infected with
SARS-CoV-2 develop severe and fatal respiratory disease. JCI insight.
Griffin, B.D., Chan, M., Tailor, N., Mendoza, E.J., Leung, A., Warner, B.M., Duggan, A.T., Moffat, E., He, S.,
Garnett, L.*, et al.* (2020). North American deer mice are susceptible to SARS-CoV-2. bioRxiv,
2020.2007.2025.221291.
Gu, H., Chen, Q., Yang, G., He, L., Fan, H., Deng, Y.-Q., Wang, Y., Teng, Y., Zhao, Z., Cui, Y.*, et al.* (2020).
Adaptation of SARS-CoV-2 in BALB/c mice for testing vaccine efficacy. Science (New York, NY) *369*, 1603-
1607.
Guan, W.-j., Ni, Z.-y., Hu, Y., Liang, W.-h., Ou, C.-q., He, J.-x., Liu, L., Shan, H., Lei, C.-l., Hui, D.S.C.*, et al.*
(2020). Clinical Characteristics of Coronavirus Disease 2019 in China. New England Journal of Medicine
382, 1708-1720.
Han, H., Ma, Q., Li, C., Liu, R., Zhao, L., Wang, W., Zhang, P., Liu, X., Gao, G., Liu, F.*, et al.* (2020). Profiling
serum cytokines in COVID-19 patients reveals IL-6 and IL-10 are disease severity predictors. Emerging
microbes & infections *9*, 1123-1130.
Harrow, J., Frankish, A., Gonzalez, J.M., Tapanari, E., Diekhans, M., Kokocinski, F., Aken, B.L., Barrell, D.,
Zadissa, A., Searle, S.*, et al.* (2012). GENCODE: the reference human genome annotation for The ENCODE
Project. Genome Res *22*, 1760-1774.

Hassan, A.O., Case, J.B., Winkler, E.S., Thackray, L.B., Kafai, N.M., Bailey, A.L., McCune, B.T., Fox, J.M.,
Chen, R.E., Alsoussi, W.B., *et al.* (2020). A SARS-CoV-2 Infection Model in Mice Demonstrates Protection
by Neutralizing Antibodies. Cell *182*, 744-753.e744.
Herold, T., Jurinovic, V., Arnreich, C., Lipworth, B.J., Hellmuth, J.C., von Bergwelt-Baildon, M., Klein, M.,
and Weinberger, T. (2020). Elevated levels of IL-6 and CRP predict the need for mechanical ventilation in
COVID-19. Journal of Allergy and Clinical Immunology *146*, 128-136.e124.
Huang, C., Wang, Y., Li, X., Ren, L., Zhao, J., Hu, Y., Zhang, L., Fan, G., Xu, J., Gu, X., *et al.* (2020). Clinical
features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet (London, England)
395, 497-506.
Israelow, B., Song, E., Mao, T., Lu, P., Meir, A., Liu, F., Alfajaro, M.M., Wei, J., Dong, H., Homer, R.J., *et al.*
(2020). Mouse model of SARS-CoV-2 reveals inflammatory role of type I interferon signaling. The Journal
of experimental medicine *217*.
Jefferson, T., Spencer, E., Brassey, J., and Heneghan, C. (2020). Viral cultures for COVID-19 infectivity
assessment. Systematic review. medRxiv, 2020.2008.2004.20167932.
Jiang, H.-w., Zhang, H.-n., Meng, Q.-f., Xie, J., Li, Y., Chen, H., Zheng, Y.-x., Wang, X.-n., Qi, H., Zhang, J., *et*
al. (2020a). SARS-CoV-2 Orf9b suppresses type I interferon responses by targeting TOM70. Cellular &
Molecular Immunology *17*, 998-1000.
Jiang, R.-D., Liu, M.-Q., Chen, Y., Shan, C., Zhou, Y.-W., Shen, X.-R., Li, Q., Zhang, L., Zhu, Y., Si, H.-R., *et al.*
(2020b). Pathogenesis of SARS-CoV-2 in Transgenic Mice Expressing Human Angiotensin-Converting
Enzyme 2. Cell *182*, 50-58.e58.
Jin, J.-M., Bai, P., He, W., Wu, F., Liu, X.-F., Han, D.-M., Liu, S., and Yang, J.-K. (2020). Gender Differences
in Patients With COVID-19: Focus on Severity and Mortality. Frontiers in Public Health *8*.
Johnson, B.J., Le, T.T., Dobbin, C.A., Banovic, T., Howard, C.B., Flores Fde, M., Vanags, D., Naylor, D.J.,
Hill, G.R., and Suhrbier, A. (2005). Heat shock protein 10 inhibits lipopolysaccharide-induced
inflammatory mediator production. J Biol Chem *280*, 4037-4047.
Kremer, K.L., Dunning, K.R., Parsons, D.W., and Anson, D.S. (2007). Gene delivery to airway epithelial
cells in vivo: a direct comparison of apical and basolateral transduction strategies using pseudotyped
lentivirus vectors. The journal of gene medicine *9*, 362-368.
Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L.,
Jagodnik, K.M., Lachmann, A., *et al.* (2016). Enrichr: a comprehensive gene set enrichment analysis web
server 2016 update. Nucleic Acids Res *44*, W90-97.
Kumari, P., Rothan, H.A., Natekar, J.P., Stone, S., Pathak, H., Strate, P.G., Arora, K., Brinton, M.A., and
Kumar, M. (2020). Neuroinvasion and encephalitis following intranasal inoculation of SARS-CoV-2 in K18-
hACE2 mice. bioRxiv, 2020.2012.2014.422714.
La Linn, M., Bellett, A.J., Parsons, P.G., and Suhrbier, A. (1995). Complete removal of mycoplasma from
viral preparations using solvent extraction. J Virol Methods *52*, 51-54.
Lavoie, H., Gagnon, J., and Therrien, M. (2020). ERK signalling: a master regulator of cell behaviour, life
and fate. Nature Reviews Molecular Cell Biology *21*, 607-632.
Lazear, H.M., Schoggins, J.W., and Diamond, M.S. (2019). Shared and Distinct Functions of Type I and
Type III Interferons. Immunity *50*, 907-923.
Leist, S.R., Dinnon, K.H., 3rd, Schäfer, A., Tse, L.V., Okuda, K., Hou, Y.J., West, A., Edwards, C.E., Sanders,
W., Fritch, E.J., *et al.* (2020). A Mouse-Adapted SARS-CoV-2 Induces Acute Lung Injury and Mortality in
Standard Laboratory Mice. Cell.
Li, B., and Dewey, C.N. (2011). RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. BMC Bioinformatics *12*, 323.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R.
(2009). The Sequence Alignment/Map format and SAMtools. Bioinformatics *25*, 2078-2079.

Li, S., Rouphael, N., Duraisingham, S., Romero-Steiner, S., Presnell, S., Davis, C., Schmidt, D.S., Johnson,
S.E., Milton, A., Rajam, G., *et al.* (2014). Molecular signatures of antibody responses derived from a
systems biology study of five human vaccines. Nature immunology *15*, 195-204.

Limberis, M., Anson, D.S., Fuller, M., and Parsons, D.W. (2002). Recovery of airway cystic fibrosis
transmembrane conductance regulator function in mice with cystic fibrosis after single-dose lentivirus-
mediated gene transfer. Human gene therapy *13*, 1961-1970.

Limberis, M.P., Bell, C.L., Heath, J., and Wilson, J.M. (2010). Activation of transgene-specific T cells
following lentivirus-mediated gene delivery to mouse lung. Mol Ther *18*, 143-150.

Lin, L., Han, Q., Xiong, Y., Li, T., Liu, Z., Xu, H., Wu, Y., Wang, N., and Liu, X. (2017). Krüpple-like-factor 4
Attenuates Lung Fibrosis via Inhibiting Epithelial-mesenchymal Transition. Scientific reports *7*, 15847-
15847.

Lu, L., Zhang, H., Dauphars, D.J., and He, Y.-W. (2021). A Potential Role of Interleukin 10 in COVID-19
Pathogenesis. Trends in Immunology *42*, 3-5.

Mahlaкõiv, T., Ritz, D., Mordstein, M., DeDiego, M.L., Enjuanes, L., Müller, M.A., Drosten, C., and
Staeheli, P. (2012). Combined action of type I and type III interferon restricts initial replication of severe
acute respiratory syndrome coronavirus in the lung but fails to inhibit systemic virus spread. The Journal
of general virology *93*, 2601-2605.

Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. 2011
17, 3.

Masiá, M., Fernández-González, M., Padilla, S., Ortega, P., García, J.A., Agulló, V., García-Abellán, J.,
Telenti, G., Guillén, L., and Gutiérrez, F. (2020). Impact of interleukin-6 blockade with tocilizumab on
SARS-CoV-2 viral kinetics and antibody responses in patients with COVID-19: A prospective cohort study.
EBioMedicine *60*.

McCray, P.B., Jr., Pewe, L., Wohlford-Lenane, C., Hickey, M., Manzel, L., Shi, L., Netland, J., Jia, H.P.,
Halabi, C., Sigmund, C.D., *et al.* (2007). Lethal infection of K18-hACE2 mice infected with severe acute
respiratory syndrome coronavirus. Journal of virology *81*, 813-821.

Mehdipour, A.R., and Hummer, G. (2020). Dual nature of human ACE2 glycosylation in binding to SARS-
CoV-2 spike. bioRxiv, 2020.2007.2009.193680.

Mi, H., Muruganujan, A., Ebert, D., Huang, X., and Thomas, P.D. (2019). PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic Acids Res *47*,
D419-d426.

Miller, B., Silverstein, A., Flores, M., Cao, K., Kumagai, H., Mehta, H.H., Yen, K., Kim, S.-J., and Cohen, P.
(2021). Host mitochondrial transcriptome response to SARS-CoV-2 in multiple cell models and clinical
samples. Scientific Reports *11*, 3.

Miorin, L., Kehrer, T., Sanchez-Aparicio, M.T., Zhang, K., Cohen, P., Patel, R.S., Cupic, A., Makio, T., Mei,
M., Moreno, E., *et al.* (2020). SARS-CoV-2 Orf6 hijacks Nup98 to block STAT nuclear import and
antagonize interferon signaling. Proceedings of the National Academy of Sciences *117*, 28344.

Monk, P.D., Marsden, R.J., Tear, V.J., Brookes, J., Batten, T.N., Mankowski, M., Gabbay, F.J., Davies, D.E.,
Holgate, S.T., Ho, L.-P., *et al.* (2020). Safety and efficacy of inhaled nebulised interferon beta-1a
(SNG001) for treatment of SARS-CoV-2 infection: a randomised, double-blind, placebo-controlled, phase
2 trial. The Lancet Respiratory Medicine.

Moreau, G.B., Burgess, S.L., Sturek, J.M., Donlan, A.N., Petri, W.A., and Mann, B.J. (2020). Evaluation of
K18-hACE2 Mice as a Model of SARS-CoV-2 Infection. The American journal of tropical medicine and
hygiene *103*, 1215-1219.

Muñoz-Fontela, C., Dowling, W.E., Funnell, S.G.P., Gsell, P.-S., Riveros-Balta, A.X., Albrecht, R.A.,
Andersen, H., Baric, R.S., Carroll, M.W., Cavaleri, M., *et al.* (2020). Animal models for COVID-19. Nature
586, 509-515.

Nechanitzky, R., Akbas, D., Scherer, S., Györy, I., Hoyler, T., Ramamoorthy, S., Diefenbach, A., and
Grosschedl, R. (2013). Transcription factor EBF1 is essential for the maintenance of B cell identity and
prevention of alternative fates in committed cells. Nature immunology *14*, 867-875.
Nicks, K.M., Fowler, T.W., Akel, N.S., Perrien, D.S., Suva, L.J., and Gaddy, D. (2010). Bone turnover across
the menopause transition : The role of gonadal inhibins. Ann N Y Acad Sci *1192*, 153-160.
Oladunni, F.S., Park, J.-G., Pino, P.A., Gonzalez, O., Akhter, A., Allué-Guardia, A., Olmo-Fontánez, A.,
Gautam, S., Garcia-Vilanova, A., Ye, C., *et al.* (2020). Lethality of SARS-CoV-2 infection in K18 human
angiotensin-converting enzyme 2 transgenic mice. Nature Communications *11*, 6122.
Peckham, H., de Gruijter, N.M., Raine, C., Radziszewska, A., Ciurtin, C., Wedderburn, L.R., Rosser, E.C.,
Webb, K., and Deakin, C.T. (2020). Male sex identified by global COVID-19 meta-analysis as a risk factor
for death and ITU admission. Nature Communications *11*, 6317.
Pinna, G. (2021). Sex and COVID-19: A Protective Role for Reproductive Steroids. Trends in
endocrinology and metabolism: TEM *32*, 3-6.
Raslan, A.A., and Yoon, J.K. (2020). WNT Signaling in Lung Repair and Regeneration. Mol Cells *43*, 774-
783.
Rathnasinghe, R., Strohmeier, S., Amanat, F., Gillespie, V.L., Krammer, F., García-Sastre, A., Coughlan, L.,
Schotsaert, M., and Uccellini, M. (2020). Comparison of Transgenic and Adenovirus hACE2 Mouse
Models for SARS-CoV-2 Infection. Emerging microbes & infections, 1-19.
Rodrigues, J.P.G.L.M., Barrera-Vilarmau, S., M. C. Teixeira, J., Sorokina, M., Seckel, E., Kastritis, P.L., and
Levitt, M. (2020). Insights on cross-species transmission of SARS-CoV-2 from structural modeling. PLOS
Computational Biology *16*, e1008449.
Rusinova, I., Forster, S., Yu, S., Kannan, A., Masse, M., Cumming, H., Chapman, R., and Hertzog, P.J.
(2013). INTERFEROME v2.0: an updated database of annotated interferon-regulated genes. Nucleic Acids
Research *41*, D1040-D1046.
Schroder, W.A., Le, T.T.T., Major, L., Street, S., Gardner, J., Lambley, E., Markey, K., MacDonald, K.P.,
Fish, R.J., Thomas, R., *et al.* (2010). A Physiological Function of Inflammation-Associated SerpinB2 Is
Regulation of Adaptive Immunity. The Journal of Immunology *184*, 2663.
Schubert, K., Karousis, E.D., Jomaa, A., Scaiola, A., Echeverria, B., Gurzeler, L.-A., Leibundgut, M., Thiel,
V., Mühlemann, O., and Ban, N. (2020). SARS-CoV-2 Nsp1 binds the ribosomal mRNA channel to inhibit
translation. Nature Structural & Molecular Biology *27*, 959-966.
Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N., Schwikowski, B., and
Ideker, T. (2003). Cytoscape: a software environment for integrated models of biomolecular interaction
networks. Genome Res *13*, 2498-2504.
Simons, A. (2010). A quality control tool for high throughput sequence data. Available online:
https://www.bioinformatics.babraham.ac.uk/projects/fastqc.
Sountoulidis, A., Stavropoulos, A., Giaglis, S., Apostolou, E., Monteiro, R., Chuva de Sousa Lopes, S.M.,
Chen, H., Stripp, B.R., Mummery, C., Andreakos, E., *et al.* (2012). Activation of the Canonical Bone
Morphogenetic Protein (BMP) Pathway during Lung Morphogenesis and Adult Lung Tissue Repair. PLOS
ONE *7*, e41460.
Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A.,
Pomeroy, S.L., Golub, T.R., Lander, E.S., *et al.* (2005). Gene set enrichment analysis: a knowledge-based
approach for interpreting genome-wide expression profiles. Proc Natl Acad Sci U S A *102*, 15545-15550.
Sun, J., Zhuang, Z., Zheng, J., Li, K., Wong, R.L.-Y., Liu, D., Huang, J., He, J., Zhu, A., Zhao, J., *et al.* (2020a).
Generation of a Broadly Useful Model for COVID-19 Pathogenesis, Vaccination, and Treatment. Cell *182*,
734-743.e735.
Sun, S.H., Chen, Q., Gu, H.J., Yang, G., Wang, Y.X., Huang, X.Y., Liu, S.S., Zhang, N.N., Li, X.F., Xiong, R., *et*
al. (2020b). A Mouse Model of SARS-CoV-2 Infection and Pathogenesis. Cell host & microbe *28*, 124-
133.e124.

Swann, J.B., Hayakawa, Y., Zerafa, N., Sheehan, K.C., Scott, B., Schreiber, R.D., Hertzog, P., and Smyth,
M.J. (2007). Type I IFN contributes to NK cell homeostasis, activation, and antitumor function. J Immunol
178, 7540-7549.
Szklarczyk, D., Gable, A.L., Lyon, D., Junge, A., Wyder, S., Huerta-Cepas, J., Simonovic, M., Doncheva,
N.T., Morris, J.H., Bork, P.*, et al.* (2019). STRING v11: protein-protein association networks with increased
coverage, supporting functional discovery in genome-wide experimental datasets. Nucleic Acids Res *47*,
D607-D613.
Takasuka, N., Fujii, H., Takahashi, Y., Kasai, M., Morikawa, S., Itamura, S., Ishii, K., Sakaguchi, M., Ohnishi,
K., Ohshima, M.*, et al.* (2004). A subcutaneously injected UV-inactivated SARS coronavirus vaccine elicits
systemic humoral immunity in mice. International immunology *16*, 1423-1430.
Tseng, C.T., Huang, C., Newman, P., Wang, N., Narayanan, K., Watts, D.M., Makino, S., Packard, M.M.,
Zaki, S.R., Chan, T.S.*, et al.* (2007). Severe acute respiratory syndrome coronavirus infection of mice
transgenic for the human Angiotensin-converting enzyme 2 virus receptor. Journal of virology *81*, 1162-
1173.
Vanderheiden, A., Ralfs, P., Chirkova, T., Upadhyay, A.A., Zimmerman, M.G., Bedoya, S., Aoued, H.,
Tharp, G.M., Pellegrini, K.L., Manfredi, C.*, et al.* (2020). Type I and Type III Interferons Restrict SARS-CoV-
2 Infection of Human Airway Epithelial Cultures. Journal of virology *94*, e00985-00920.
Vaz de Paula, C.B., de Azevedo, M.L.V., Nagashima, S., Martins, A.P.C., Malaquias, M.A.S., Miggiolaro,
A.F.R.d.S., da Silva Motta Júnior, J., Avelino, G., do Carmo, L.A.P., Carstens, L.B.*, et al.* (2020). IL-4/IL-13
remodeling pathway of COVID-19 lung injury. Scientific Reports *10*, 18689.
Wajnberg, A., Amanat, F., Firpo, A., Altman, D.R., Bailey, M.J., Mansour, M., McMahon, M., Meade, P.,
Mendu, D.R., Muellers, K.*, et al.* (2020). Robust neutralizing antibodies to SARS-CoV-2 infection persist
for months. Science (New York, NY) *370*, 1227.
Wang, H., Kadlecek, T.A., Au-Yeung, B.B., Goodfellow, H.E.S., Hsu, L.-Y., Freedman, T.S., and Weiss, A.
(2010). ZAP-70: an essential kinase in T-cell signaling. Cold Spring Harb Perspect Biol *2*, a002279-
a002279.
Wang, Q., Zhang, Y., Wu, L., Niu, S., Song, C., Zhang, Z., Lu, G., Qiao, C., Hu, Y., Yuen, K.-Y.*, et al.* (2020).
Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2. Cell *181*, 894-904.e899.
Weiskopf, D., Schmitz, K.S., Raadsen, M.P., Grifoni, A., Okba, N.M.A., Endeman, H., van den Akker, J.P.C.,
Molenkamp, R., Koopmans, M.P.G., van Gorp, E.C.M.*, et al.* (2020). Phenotype and kinetics of SARS-CoV-
2–specific T cells in COVID-19 patients with acute respiratory distress syndrome. Science Immunology *5*,
eabd2071.
Wilson, J.M. (2004). Adeno-associated virus and lentivirus pseudotypes for lung-directed gene therapy.
Proceedings of the American Thoracic Society *1*, 309-314.
Winkler, E.S., Bailey, A.L., Kafai, N.M., Nair, S., McCune, B.T., Yu, J., Fox, J.M., Chen, R.E., Earnest, J.T.,
Keeler, S.P.*, et al.* (2020). SARS-CoV-2 infection of human ACE2-transgenic mice causes severe lung
inflammation and impaired function. Nature immunology.
Wu, F., Zhao, S., Yu, B., Chen, Y.-M., Wang, W., Song, Z.-G., Hu, Y., Tao, Z.-W., Tian, J.-H., Pei, Y.-Y.*, et al.*
(2020a). A new coronavirus associated with human respiratory disease in China. Nature *579*, 265-269.
Wu, M., Chen, Y., Xia, H., Wang, C., Tan, C.Y., Cai, X., Liu, Y., Ji, F., Xiong, P., Liu, R.*, et al.* (2020b).
Transcriptional and proteomic insights into the host response in fatal COVID-19 cases. Proceedings of
the National Academy of Sciences *117*, 28336.
Xia, H., Cao, Z., Xie, X., Zhang, X., Chen, J.Y.-C., Wang, H., Menachery, V.D., Rajsbaum, R., and Shi, P.-Y.
(2020). Evasion of Type I Interferon by SARS-CoV-2. Cell Reports *33*, 108234.
Yan, K., Vet, L.J., Tang, B., Hobson-Peters, J., Rawle, D.J., Le, T.T., Larcher, T., Hall, R.A., and Suhrbier, A.
(2020). A Yellow Fever Virus 17D Infection and Disease Mouse Model Used to Evaluate a Chimeric
Binjari-Yellow Fever Virus Vaccine. Vaccines *8*.

1082 Yinda, C.K., Port, J.R., Bushmaker, T., Offei Owusu, I., Purushotham, J.N., Avanzato, V.A., Fischer, R.J.,
1083 Schulz, J.E., Holbrook, M.G., Hebner, M.J.*, et al.* (2021). K18-hACE2 mice develop respiratory disease
1084 resembling severe COVID-19. PLoS Pathog *17*, e1009195.
1085 Zhao, X., Chen, D., Szabla, R., Zheng, M., Li, G., Du, P., Zheng, S., Li, X., Song, C., Li, R.*, et al.* (2020). Broad
1086 and Differential Animal Angiotensin-Converting Enzyme 2 Receptor Usage by SARS-CoV-2. Journal of
1087 virology *94*.
1088 Zhou, P., Yang, X.-L., Wang, X.-G., Hu, B., Zhang, L., Zhang, W., Si, H.-R., Zhu, Y., Li, B., Huang, C.-L.*, et al.*
1089 (2020). A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature *579*,
1090 270-273.
1091 Zhu, Y., Wang, W., and Wang, X. (2015). Roles of transcriptional factor 7 in production of inflammatory
1092 factors for lung diseases. J Transl Med *13*, 273-273.

1093

Figure 1

A RSVp | 5' LTR | Ψ | Full length EF1α promoter | **ACE2** | BGH-poly A | PGKp | T2A GFP | Puro | WPRE | 3' LTR

B HEK293T cells hACE2 mutants

Virus titer, log$_{10}$ CCID50/ml ± SE

Days post-infection

- ● Untransduced
- ■ WT hACE2
- ▲ hACE2-T92Q
- ▼ hACE2-K31N/K353H
- ◆ hACE2-T27Y/L79Y/N330Y

C HEK293T cells mACE2 mutants

Virus titer, log$_{10}$ CCID50/ml ± SE

Days post-infection

- ● Untransduced
- ■ WT mACE2
- ▼ mACE2-N30D/N31K/T79L/S82M/F83Y/A329E/H353K
- ▲ **mACE2-N30D/N31K/F83Y/H353K**
- ◆ **mACE2-N31K/H353K**

D Untransduced

WT hACE2 | WT mACE2

mACE2 N30D/N31K/F83Y/H353K | mACE2 N31K/H353K

E Spike RBD

WT hACE2

N487 | K417 | Q493 | G496 | D30 | K31 | K353 | G502 | Y83

SARS-CoV-2 can infect cells via mACE2 with these changes

Figure 2

Figure 3

A

hACE2

Day 2 + -

Down DEGs
n=15

Up DEGs
n=95

Not significant

-Log$_{10}$ FDR

Log$_2$ Fold Change

B

95 up DEGs in
Interferome

Type I 9

Type II 0

Type III

D

Cytokine up DEGs from
K18-hACE2 mice vs pre-
ranked all genes from 'A'

NES = 1.627
p-value = <0.001

Type I IFN up DEGs from
K18-hACE2 mice vs pre-
ranked all genes from 'A'

NES = 1.48
p-value = 0.024

C

Upstream
regulators

IL4

○ Th1 associated
○ Th2 associated
○ Th17 associated
○ Pleiotropic/other

IL6   IL10

IFNG   IL13

CSF2

IFNA/B

CSF1

IFNA   TNF   IL2

TNFSF11

IL1B

CSF3

IL27   L33   IL5

IL3

IL21   IFNB1

-log$_{10}$ p-value

Activation z-score

E

Diseases and
functions

Activation z-score

Leukopoiesis
Response of antigen presenting cells
Hematopoiesis of mononuclear leukocytes
Immune response of antigen presenting cells
Engulfment of antigen presenting cells
Response of macrophages
Engulfment of phagocytes
Response of phagocytes
Engulfment by macrophages
Response of myeloid cells
Quantity of mononuclear leukocytes
Quantity of lymphocytes
Quantity of leukocytes
Engulfment of myeloid cells
Migration of cells
Phagocytosis of phagocytes
Phagocytosis of myeloid cells
Immune response of leukocytes
Quantity of blood cells
Phagocytosis
Immune response of phagocytes
Inflammatory response
Cell movement

■ Cellular infiltration
■ Innate responses

-log$_{10}$ p-value

F

Human COVID-19 lung
up DEGs vs pre-ranked
all genes from 'A'

NES = 1.41
p-value = 0.004

Core enriched genes analysed by GO Biological Processes
ontology (Enrichr). Top 10 sorted by p-value:

GO process	P-value
cytokine-mediated signaling pathway	1.39E-20
cellular response to type I interferon	8.70E-20
type I interferon signaling pathway	8.70E-20
cellular response to interferon-gamma	4.33E-13
chemokine-mediated signaling pathway	6.10E-09
response to interferon-gamma	4.14E-08
inflammatory response	4.36E-08
lymphocyte chemotaxis	7.20E-08
eosinophil migration	1.01E-07
eosinophil chemotaxis	1.01E-07

Figure 4

Figure 5

Figure 6

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3