**Single paternal Dexamethasone challenge programs offspring metabolism and reveals**

**circRNAs as novel candidates in RNA-mediated inheritance**

Katharina Gapp[#1,2,3,4], Guillermo Parada[1,2,5], Fridolin Gross[3], Alberto Corcoba[2], Evelyn Grau[2],

Martin Hemberg[1,2], Johannes Bohacek[3,4], Eric A. Miska*[1,2,5]

[1] Gurdon Institute, University of Cambridge, Tennis Court Rd, Cambridge, CB2 1QN, UK.

[2] Wellcome Sanger Institute, Wellcome Genome Campus, Hinxton, CB10 1SA, UK.

[3] Institute for Neuroscience, Department of Health Science Technology, ETH Zürich,

Winterthurerstr. 190, 8057 Zürich, Switzerland

[4] Neuroscience Center Zurich, ETH Zurich and University of Zurich, Switzerland

[5] Department of Genetics, University of Cambridge, Downing Street, Cambridge, CB2 3EH,

UK.

* corresponding

[#] co-corresponding author

**Summary**

Single traumatic events that elicit an exaggerated stress response can lead to the development of neuropsychiatric conditions. Studies in mice suggests germline RNA as a mediator of effects of chronic environmental exposures to the progeny. The effects of an acute paternal stress exposure on the germline and their potential consequences on offspring remain unknown. We find that acute administration of an agonist for the stress-sensitive Glucocorticoid receptor, using the common corticosteroid Dexamethasone, affects the RNA payload of post-meiotic mature sperm as soon as 3 hours post exposure, challenging the prevalent view that epididymal transit is required to alter sperm RNA. It further impacts early embryonic transcriptional trajectories, as determined by single embryo sequencing, and metabolism in the offspring.

Importantly, we show persistent regulation of tRNA fragments in sperm and the descendant 2-cell- embryos, suggesting actual transmission from sperm to embryo. Lastly, we unravel environmentally induced alterations in the previously underconsidered class of sperm circRNAs, and their targets in the early embryo, highlighting this class as a novel candidate in RNA-mediated inheritance.

**Introduction**

Acute stress elicits a complex but well-studied cascade of neuroendocrine responses regulated by the hypothalamic pituitary adrenal axis. It involves the release of neuropeptides in the brain that induce the secretion of corticosteroid hormones from the adrenals. These hormones in turn activate mainly two types of nuclear glucocorticoid receptors (GR) (de Kloet, Karst and Joëls, 2008) expressed throughout the body, which then regulate gene expression and enable physiological and behavioral adjustments (de Kloet, Joëls and Holsboer, 2005). In vulnerable individuals, this response is excessive and it can lead to long lasting maladaptations with consequences on psychological and metabolic health (Daskalakis *et al.*, 2012).

It is also known that parental experiences can compromise the health of their progeny both in humans (Pembrey *et al.*, 2006; Heijmans *et al.*, 2008; Bowers and Yehuda, 2016)  and in animal models  (Benyshek, Johnston and Martin, 2006; Roth *et al.*, 2009; Jimenez-Chillaron *et al.*, 2009; Carone *et al.*, 2010; Pentinat *et al.*, 2010; Shankar *et al.*, 2010; Franklin *et al.*, 2010, 2011; Morgan and Bale, 2011; Weiss *et al.*, 2011; Dietz *et al.*, 2011; Vassoler *et al.*, 2013; Fullston *et al.*, 2013; Martínez *et al.*, 2014; Gapp *et al.*, 2014; Marco *et al.*, 2014; Rodgers *et al.*, 2015; Sharma *et al.*, 2015; Wu *et al.*, 2016; Chen *et al.*, 2016; Zhang *et al.*, 2018). Research on the underlying mechanism of such transmission has found changes in germline epigenetic make-up, in particular DNA methylation, histone post translational modifications (PTMs), histone positioning and RNA (Gapp and Bohacek, 2017). These epigenetic regulators are responsive to the environment and have been implicated in a variety of environmentally induced diseases (Jirtle and Skinner, 2007). Altered modifications must circumvent epigenetic reprogramming events in zygote and, depending on the timing of exposure, during germline development (Bohacek and Mansuy, 2017; Gapp and Bohacek, 2017). In the male germline, RNA is excluded from reprogramming and therefore a promising candidate for transgenerational information delivery (Gapp and Bohacek, 2018; Bohacek and Rassoulzadegan, 2019). Several studies carried out in *D. melangolaster* and *C. elegans* reported on transgenerational inheritance of induced traits and provided firm evidence for the involvement of small RNAs in the mechanism of transmission (Ashe *et al.*, 2012; Grentzinger *et al.*, 2012; Shirayama *et al.*, 2012). In mammals, a causal implication in the transmission of environmentally induced effects across generations has also solely been demonstrated for sperm RNA (Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Sharma *et al.*, 2015; Chen *et al.*, 2016).  Such RNA differs substantially from somatic RNA since it mainly consists of small RNA, predominantly tRNA-derived small fragments (tsRNAs), but also miRNAs, piRNAs and circRNAs, among others (Bohacek and Rassoulzadegan, 2019)(Chen, Yan and Duan, 2016; Gapp and Bohacek, 2018). CircRNAs, comprise a very stable class of RNA that has recently been observed to be present in high amounts in testis but also to some extent in sperm (Barrett and Salzman, 2016). Some have been shown to act as miRNA sponges, thereby

competing with mRNA targets, while also regulating the expression of their host genes (Barrett and Salzman, 2016). Hence, circRNAs have a strong potential for amplifying an inherited signal, which makes circRNAs exceptionally interesting candidates for epigenetic germline inheritance. To date, the involvement of circRNAs in soma-to-germline signalling has not yet been investigated.

tsRNAs and miRNAs are crucial regulators of early embryonic development and players in non-genetic inheritance (Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Rodgers *et al.*, 2015; Sharma *et al.*, 2015; Chen *et al.*, 2016; Benito *et al.*, 2018; Tyebji, Hannan and Tonkin, 2020). They have been reported to be acquired through exosomal uptake during epididymal transfer from caput to cauda epididymis (16,44). This might explain their responsiveness to environmental perturbations, despite mature sperm's presumably transcriptionally silent state caused by tightly packed chromatin. Sperm RNA can indeed change in response to chronic stress or by chronic treatments that mimic stress exposure, such as repeated injection of GR agonists (Rodgers *et al.*, 2013; Gapp *et al.*, 2014; Short *et al.*, 2016; Wu *et al.*, 2016). Uptake of epididysomal miRNA was sufficient to replicate a chronic stress induced effect on stress response in offspring mice(Chan *et al.*, 2020) . Surprisingly, acute stress has also recently been shown to affect offspring weight and glucose metabolism in mice (Hoyer *et al.*, 2013) and some of these effects were germline dependent (Bohacek, von Werdt and Mansuy, 2016). Together these related lines of evidence led us to hypothesize that acute GR activation has an intergenerational effect on offspring phenotype and that the transmission potentially implicates changes in the germline. The male germline cells - including mature sperm (Kaufmann *et al.*, 1992; Haeussler and Claus, 2007) and their surrounding Sertoli cells (Hazra *et al.*, 2014) – as well es the epididymal epithelial cells (Silva *et al.*, 2010) express GRs that mediate the effects of glucocorticoids on transcription. Dexamethasone (Dex) a specific  GR agonist is known to directly activate GR in the rat epididymis (Silva *et al.*, 2014). It is unknown whether acute stress affects sperm RNA, and if so, whether uptake via epididymisomes is involved in establishing germline changes that are relevant for offspring phenotypic alterations.

Here we investigate the impact of acute GR agonist administration on the germline RNA payload including circRNAs, at various time points post administration and interrogate the fate of altered sperm RNA. We further test germ-line dependency of transmitted metabolic effects and dissect the underlying molecular trajectories during early embryonic development using single cell sequencing of in vitro fertilization (IVF) derived embryos. Identifying a readout of transgenerational risk load at the level of the paternal sperm epigenome could pave the way for future studies aiming at a prevention of the transmission of the effects of acute GR activation to the offspring.

**Results**

*Effects of acute Dex injection on the germline small RNA payload*

Two reports have suggested that a single foot shock could elicit effects on offspring phenotype (Hoyer *et al.*, 2013; Bohacek, von Werdt and Mansuy, 2016).To examine potential epigenetic mediators of such acute stressful impacts we investigated sperm RNA of males 2 weeks after a single activation of the GR (Figure 1A). This timeline was chosen to mimic the timing at which breeding occurred when effects on offspring had been observed in a previous study (Bohacek, von Werdt and Mansuy, 2016). We injected the specific GR agonist Dex once intraperitoneally into 8 adult males. This drug is in frequent clinical use, now also as an apparently effective treatment for patients suffering from lower respiratory tract infection as a consequence of Covid-19 virus (EU Clinical Trials Register, no date; 'Biggest COVID-19 trial tests repurposed drugs first', 2020; Horby *et al.*, 2020). A  sperm population was harvested from each animal and RNA was extracted for ultra-deep small RNA sequencing, resulting in 16 libraries representing one injected male each (8 vehicle and 8 Dex-injected). Purity of the sperm samples was confirmed by inspecting RNA size profiles generated on the bioanalyzer to be absent of ribosomal RNA peaks, that would indicate contamination by somatic cells (Supplementary Fig. 1A). Reaching an average of 55.4 Million sequencing reads while also using randomized adaptors for 3′ ligation put us in a position to reduce PCR biases (Dard-

Dascot *et al.*, 2018) and accurately quantify less abundant miRNAs that are by far outnumbered in sperm by other small RNAs e.g. tsRNAs(Peng *et al.*, 2012). Our data showed an average of 60% mappable reads across all libraries, including 34% of multimappers. We detected an expected dominant prevalence of reads mapping to tsRNAs and abundant miRNAs in all samples (Supplementary Fig. 2A). Differential gene expression analysis, using DEseq2 (Love, Huber and Anders, 2014), revealed that a single acute activation of GR receptors induced changes in tsRNAs and miRNAs collected 14 days post injection (Figure 1B,C, FDR q<0.05), as has been observed in response to chronic environmental stress previously (Rodgers *et al.*, 2013; Gapp *et al.*, 2014). Interestingly, tsRNA-Gly-GCC, a tsRNA previously associated with the effects of nutritional challenge (Sharma *et al.*, 2015), was among the most strongly altered tsRNAs.

Some recent publications have suggested that sperm miRNAs and tsRNAs are acquired during epididymal transit from caput to cauda (Sharma *et al.*, 2015, 2018; Conine *et al.*, 2018). Further, it was shown that chronic nutritional-challenge-induced change in sperm tsRNAs is acquired by uptake of distinct sets of tsRNAs (Sharma *et al.*, 2015). To investigate whether the changes in small RNA after acute stress are due to uptake during transit from caput to cauda epididymis we investigated the mature sperm small RNA payload at two time points, 3 hours and 7 days following injection (Figure 2A, Supplementary Fig. 1C). Cells collected from cauda 7 days after injection had already exited testis, but have had time to pass through the entire epididymal tract before collection. Cells collected 3 hours post injection will most likely not have passed through the corpus epididiymis, and already resided in cauda epididymis at the time of injection where sperm resides up to 5 days (Meistrich, 1975; Dadoune and Alfonsi, 1984). Importantly, spontaneous ejaculation regularly voids cauda epididymis of sperm, even in the absence of a mating partner(Huber and Bronson, 1980), excluding the retention of "old" mature sperm in cauda for prolonged periods of time. The cells collected 7 days after exposure therefore represent a mixture of cells that might have already resided in the cauda and those cells that indeed passed through the corpus

epididymis, yet the spontaneous ejaculation ensures that the sample predominantly contains

the latter.

The collected samples were confirmed for their purity (Supplementary Fig. 1) and again

processed separately to represent sperm from one animal per library. The resulting libraries

were analysed jointly as to test for (1) effects of Dex injection independent of sampling time

post injeciton (2) effects of sampling time post injection independent of Dex treatment and (3)

effects depending on both Dex injection and the sampling time post injection (interaction).

We report an average of 64% mappable reads including 46% of multimappers and observe

that tsRNAs were significantly affected by sampling time post injection independent of

treatment. This demonstrates the fluctuation of tsRNAs over time in response to external

signals such as injections, or potentially due to uncontrollable external inputs from the animal

husbandry (Figure 2C). Interaction between treatment and time was statistically significant

for 27 tsRNA mapping loci including Gly-GCC-6-1, showing all upregulation after 7 days but

either no regulation (26 tsRNAs) or downregulation (Thr-TGT1-1) after 3 hours

(Supplementary Table2 sheet 3, q<0.05). This finding is consistent with the dominating view

that tsRNAs are acquired during epididymal transit from caput to cauda epididymis. However,

most tsRNAs that showed a significant change in response to treatment not after 3 hours but

after 7 days (interaction between treatment and time post injection, Supplementary Table 2

sheet 3, q<0.05) were not persistently altered in the dataset of 14 days after injection

(Supplementary Fig. 3). This indicates that on the one hand changes in sperm RNA are

dynamic and many do not persist for prolonged time. On the other hand this suggests that

potentially relevant small RNA changes mostly require either sperm to reside in testis at

exposure time or rely on a prolonged residency in the exposed organism.  Interestingly,

tsRNA-Leu-CAA and tsRNA-Arg-CCT, (Figure 2B) were persistently affected 3 hours and 7

days post exposure, challenging the necessity of epididymal caput to cauda transit for

acquiring tsRNA changes. While the change in tsRNA-Leu-CAA was temporary and did not

persist, strikingly tsRNA-Arg-CCT-2 deregulation persisted until 14 days post injection

(Figure 1C). To additionally validate the Dex induced change of tsRNA-Arg-CCT-2

independent of epididymal transit from caput to cauda we replicated the effect observed in

mature sperm sampled from cauda epididymis 3 hours post injection using q-PCR (Figure

2D, Supplementary table 2). Additionally, we sampled caput sperm 3 hours post injection and

measured tsRNA-Arg-CCT-2 levels. Overall 2-way ANOVA revealed a significant interaction

between sperm sampling location (caput versus cauda) and treatment (vehicle versus Dex).

Post hoc tests confirm a significant increase in tsRNA-Arg-CCT-2 levels in response to Dex

in cauda but not in caput sperm and a significant increase in Arg-CCT-2 levels between

cauda and caput sperm independent of treatment.

The behaviour of miRNAs differed considerably from tsRNAs. As would be expected if

epididymal transit was required for miRNA changes to be implemented, we observe no group

effect of treatment on miRNAs (Supplementary Fig. 4A, Supplementary table 2) across 3

hours and 7 days post injection. Further, we detected no effect of time post injection on

sperm miRNA payload (Supplementary Fig. 4B, Supplementary table 2) confirming the

absence of an effect of injection on miRNAs per se. However, we neither detected an

interaction between Dex and time post injection (Supplementary Fig. 4C, Supplementary

table 2) in miRNAs 7 days and 3 hours post injection. Importantly, when inspecting those

miRNAs that were significantly altered 14 days after injection, no alterations were apparent 3

hours or 7 days post injection (Supplementary Fig. 4D), indicating that changes in miRNAs

occur more slowly or require sperm cells to reside in testis at the time of injection.

*Effects of acute GR activation on in vivo offspring metabolic phenotype*

Based on the two reports on effects of single foot shock on offspring weight and the impact of

a single GR activation on germline small RNA payload, we hypothesized that this acute

impact on the receptor is sufficient to elicit transgenerational effects. We thus injected Dex

once intraperitoneally, then harvested sperm 14 days post injection, and performed *IVF* using

naïve oocytes to generate offspring for phenotyping (Figure 3A). Dex treatment did not affect

sperm count, fertility rate or resulting litter-sizes (Supplemenatary Fig.10-12).

The weight and size of pups was measured every 2 to 4 weeks starting at weaning (3 weeks

of age) until adulthood (12 weeks of age) and body mass index (BMI) was calculated as a

ratio of weight and squared length. Overall ANOVA of the resulting offspring showed a

significant effect of treatment ((F1.71)= 76.55, p<0.0001), time post injection

((F2.087,144.7)=41.99, p<0.0001) and sex (F (1, 71) = 76.55, p<0.0001) on BMI, and a

significant interaction between time and sex (F (3, 208) = 33.75, p<0.0001) and time and

treatment (F (3, 208) = 5.834, p=0.0008) (Figure 3B, Supplementary Fig. 5C, Supplementary

table 3). These results show that while males had generally higher BMI, both male and

female offspring of Dex injected fathers had a higher BMI.

To further explore potential causes and consequences of altered BMI, adult animals were

additionally tested for their glucose tolerance following glucose injection. Overall ANOVA

analysis of blood glucose levels revealed a significant effect of sex (F (1, 44) = 54.80,

p<0.0001) and time post injection (F (2.593, 114.1) = 196.6, p<0.0001) and significant

interactions between sex and time post injection (F (4, 176) = 6.115, P=0.0001), and sex and

treatment (F (1, 44) = 15.62, P=0.0003) (Supplementary Fig. 5C). Follow up repeated

measurements ANOVA analysis separated by sex showed a significant effect of treatment,

time and interaction  in females (treatment: F (1, 22) = 12.35, p=0.0020; time: F (4, 88) =

110.1, p<0.0001; interaction: F (4, 88) = 2.835, p=0.0291) and significant effects of treatment

and time but no interaction in males (treatment: F (1, 22) = 6.019, p=0.0225; time: F (4, 88) =

96.36, p<0.0001; F (4, 88) = 0.5401, p=0.7067; Figure 3C). These data hence demonstrate a

sex-dependent effect of paternal Dex injection on glucose tolerance, with impaired tolerance

in females and decreased glucose levels in males in response to glucose challenge.

In addition, blood glucose levels were assessed during the insulin tolerance test. Overall

ANOVA analysis showed significant effects of sex (F (1, 37) = 162.6, P<0.0001)

and time (F (3.314, 122.6) = 23.85, P<0.0001) and revealed a significant interaction between

sex and time (F (4, 148) = 12.49, P<0.0001), time and treatment (F (4, 148) = 5.380,

P=0.0005) and time and treatment and sex (F (4, 148) = 5.392, P=0.0004) (Supplementary

Fig. 5C). Follow-up repeated measurements ANOVA separated by sex showed a significant

effect of time (F (2.982, 65.60) = 44.73, p<0.0001), yet no significant effect of treatment (F (1, 22) = 0.3465, p=0.5621) nor an interaction between time and treatment (F (4, 88) = 0.1373, p=0.9681) in females (Figure 3D). In males we observe no effect of treatment F (1, 15) = 1.467, p=0.2446 yet detected a significant effect of time post injection time: F (2.914, 43.71) = 4.538, p=0.0079, and a significant interaction between treatment and time post injection F (4, 60) = 7.003, p=0.0001, Figure 3D). These results indicate sex and time dependent effects of paternal Dex on insulin tolerance. They further show no change in insulin tolerance in female descendants of fathers injected with Dex, whereas impaired insulin tolerance in male progeny.

Lastly, we explored a potential reflection of altered BMI in tissue composition by necropsy and weighing the dissected organs and fat pads. Overall ANOVA of necropsy weights revealed a significant effect of sex (F (1, 140) = 28.27, P<0.0001), tissue (F (4, 140) = 232.7, P<0.0001) and a significant interaction between sex and tissue (F (4, 140) = 3.379, P=0.0113) yet no effect of treatment (F (1, 140) = 0.2587, P=0.6118), or interaction between treatment and sex (F (1, 140) = 0.0004794, P=0.9826) or treatment and tissue (F (4, 140) = 0.1635, P=0.9565) on tissue weight (Supplementary Fig. 5A,B,C). This confirms sex dependency, yet no effect of paternal Dex injection on tissue weight in both sexes.

*Effects of acute Dex on offspring early embryonic small RNA*

The small quantity of paternal RNAs in the zygote relative to the large pool of maternal RNAs poses serious obstacles to their accurate quantification (Chen, Yan and Duan, 2016). While initial reports on small RNA transmission relied on comparative sequencing or microarrays analyses of unfertilized oocytes and fertilized zygotes (Ostermeier *et al.*, 2004), today we are aware that such comparisons can be deceiving, as they rely heavily on both assessment method (eg. microarray restricted to a selective set versus unbiased genome-wide sequencing) and sequencing depth (Dard-Dascot *et al.*, 2018; Yeri *et al.*, 2018). An example are inconsistent results regarding miRNAs that are exclusively supplied from the sperm, such as miR-34c, -99a, -214 (Amanai, Brahmajosyula and Perry, 2006; Liu *et al.*, 2012).

Alternative approaches have used indirect measures, e.g. assessing mRNA targets of

paternally derived small RNAs (Amanai, Brahmajosyula and Perry, 2006; Tang *et al.*, 2007;

Krawetz *et al.*, 2011; Sharma *et al.*, 2015). We attempted to directly examine the relative

difference between the small RNA landscape in early embryos resulting from IVF of naïve

oocytes with sperm from either Dex or vehicle injected males (Figure 4A). We used small-

RNA sequencing to compare 2-cell embryos derived from Dex treated or control fathers. We

detected an average of 29 % mappable reads including 21% multimappers. While we only

detected subtle changes in miRNAs of Dex exposed progeny (Supplementary Fig.  6), we

observed downregulation of several tsRNAs from 6 different genomic locations (q<0.1)

(Figure 4B). Strikingly, two of the downregulated tsRNAs (Gly-GCC at several genomic loci

and Gly-CCC) were consistently downregulated in sperm 14 days post Dex injection. This

likely indicates a direct delivery of these tsRNAs in control conditions, yet absence or

reduced delivery of this sperm RNA cargo of Dex treated males to the oocytes they fertilize.

*Effects of acute Dex administration on offspring early embryonic transcriptome*

If sperm RNA was directly impacting the zygotic mRNA pool, or affecting early embryonic

gene expression we hypothesized this to be apparent in the 2 cell embryo's transcriptome

(Figure 5A). To examine the effect of paternal Dex on early embryonic RNA content we

subjected 2-cell embryos to the Smartseq single cell sequencing protocol (Supplementary

Fig. 7).  After performing quality control and filtering the sequenced 2 cell embryo data on

criteria such as minimal read count/embryo (Supplementary Fig. 7A), we carried out

unsupervised clustering based on their gene expression profiles using SC3 (Kiselev *et al.*,

2017). We identified two distinct clusters (C1 and C2), which were composed by a balanced

mixture of treated and control cells. (Figure 5B). Since the resolution of single cell

experiments allows characterizing distinctive transcriptomic profiles within early cell division

stages, we used scmap (Kiselev, Yiu and Hemberg, 2018),  to project each 2 cell embryo

gene expression profile onto a reference dataset of single cells from 2 cell embryo states

previously reported by Deng *et al.* (Deng *et al.*, 2014) (Figure 5C.) Most of the 2 cell embryos

belonging to cluster C1 projected to the late 2 cell stage, whereas embryos from C2

exclusively projected to cells from the mid 2 cell stage. This shows that the two clusters

identified through unsupervised clustering correspond to 2-cell embryos in the mid and late

2-cell stage respectively.

Principal component analysis (PCA) revealed a prominent separation between C1 and C2

along the PC1 axis, suggesting a correlation between PC1 and developmental transitions

between mid and late single cell embryos (Supplementary Fig. 7B.) Interestingly, 2 cell

embryo offspring of males injected with Dex exhibited a significant shift of the C1 cluster

across PC1 (two-sided Wilcox test p< 0.03), while the C2 clusters did not show significant

differences across PC1 between treatment and control groups (Figure 5D.) These results

suggest that the effect of paternal Dex treatment on the transcriptome only becomes

apparent at the late 2 cell embryo stage. To further explore this hypothesis, we calculated the

silhouette coefficient (Rousseeuw, 1987) on PC1, as a measure of distance between C1 and

C2 clusters, for the control and treatment group. We observed a significant increase of PC1

silhouette coefficient between treatment and controls for both C1 (one-sided Wilcoxon test p-

value <0.005) and C2 (one-sided Wilcoxon test p-value < $2x10^{-5}$.) This confirms that Dex

treatment affects embryonic gene expression, promoting altered late 2 cell embryo stages

since the divergence from mid 2 cell embryos is significantly bigger in Dex offspring

compared to control offspring (Figure 5E.)

Accordingly, differential gene expression analysis using Monocle2 (Qiu *et al.*, 2017) focused

on late 2-cell embryos (cluster C1), revealed significant gene expression changes between

offspring of males injected with Dex and controls across 38 genes, some of which were

already apparent to a less significant extent during mid-2 cell embryos (cluster C1; e.g. Tcl1;

Supplementary Fig. 8A,B.) In line with a potentially altered developmental trajectory

becoming apparent in cluster 1, the late 2-cell stage includes several affected genes that are

involved in early embryonic development. For example, Bcap31 (B-cell receptor-associated

protein 31) is an important element for endoplasmatic reticulum and Golgi apparatus

function, and Bcap31 mutations lead to developmental diseases with metabolic disturbances (Cacciagli *et al.*, 2013). This is reminiscent of the metabolic phenotype observed in the adult offspring of Dex injected fathers. Hprt (Hypoxanthine-guanine phosphoribosyltransferase) is crucial for cell cycle division, and Tcl1 (T-cell leukemia/lymphoma) regulates cell proliferation (Kang *et al.*, 2013; Miyazaki *et al.*, 2013). Hence, an upregulation of Hprt, and a concomitant down-regulation of Tcl1 might indicate that cell fate decisions later during development may be affected. Another differentially expressed gene is Rbbp7 (RB binding protein 7), which is part of many histone deacetylase complexes such as Nurd and PRC2/EED-EZH2, and thus plays an essential role in chromatin mediated gene regulation (Yu *et al.*, 2018). Interestingly, several forms of PRC mutations in humans lead to different kinds of overgrowth phenotypes (Deevy and Bracken, 2019), an abnormality reminiscent of the increased BMI observed in Dex-offspring (Figure 3B.)

A previous report associated changes in genes that are coregulated by the promoter of a specific retrotransposable element (MERVL) with misexpression of tsRNA Gly-GCC in sperm (Sharma *et al.*, 2015). Since this tsRNA is significantly altered in sperm of acutely stressed males 14 days post treatment we evaluated whether the differentially expressed genes showed any overlap with the MERVL targets defined by Macfarlan *et al* (Macfarlan *et al.*, 2012) and Peaston *et al* (Peaston *et al.*, 2004). We did not detect any intersection between MERVL targets and significantly affected genes in mid or late 2 cell embryos, arguing against a functional role of tsRNA-Gly-GCC reduction in eliciting pronounced gene expression regulation of MERVL targets in the early embryos in this model.

*Effects of Dex administration on a novel candidate for sperm RNA mediated inheritance*

Despite the observed changes in sperm tsRNAs following acute Dex injection, the inability to match these changes with the previously suggested mechanism linked to MERVL targets in 2-cell embryos prompted us to investigate whether other germline changes might be more crucial for the offspring *in vivo* alterations in our model. We previously showed that chronic

stress exposure also led to changes in sperm long RNAs that contributed functionally to the transmission of effects to the offspring (Gapp *et al.*, 2018), yet the fact that sperm RNA is stable through transmission and that the minute amounts of transmitted paternal RNA can elicit major changes in the embryo remains puzzling. Therefore, we evaluated the impact of Dex injection on the highly stable class of circRNAs in male sperm. CircRNAs were previously detected in swine(Gòdia *et al.*, 2020) and human sperm (Chioccarelli *et al.*, 2019) and suggested to have functional implications in epigenetic regulation. They have been attributed a critical role post transcriptional cessation in the male germline(Tang *et al.*, 2020). Using Circexplorer in combination with EdgeR, analysis of sperm long RNA sequencing of males treated with Dex and controls revealed significant upregulation of two circRNAs (Figure 6A, q<0.1, Supplementary Fig. 1B.) while we also observed several significant changes in the sperm long RNA protein coding transcripts following acute Dex treatment (Figure 6B,C, Supplementary Fig. 1B, Supplementary Fig. 2B, Supplementary table 5). Both circRNAs are hosted in genes relevant for immune function (Taspase 1: Tasp1 and DENN Domain Containing 1B Dennd1b), yet the host genes did not show differential abundance of the protein coding host- gene transcript (Supplementary table 5). CircAtlas(Wu, Ji and Zhao, 2020) revealed several potential miRNA sponge-targets to be captured by the altered circRNAs. Some of these miRNAs are common sponge-targets of both circRNAs such as mir3110-5p, mir706, mir1955 (Supplementary Fig. 9). Diamine acetyltransferase 1 (Sat1), one of 3110-5p's high confidence miRNA-targets, as predicted by TargetScan(Agarwal *et al.*, 2015), is indeed significantly upregulated in the embryos composing cluster 1 (later developmental stage). MiRNA-target-upregulation is expected if mir3110-5p was downregulated through circRNA-mediated-sponging and highlights a potential effective contribution of increased circRNA in sperm to embryonic pathway regulation. This is the first report of a change induced by environmental exposure in this compelling class of RNA in sperm.

**Discussion**

By generating offspring using assisted reproductive techniques (IVF), we circumvent potential confounding variables such as transmission via RNA contained in seminal exosomes (Vojtech *et al.*, 2014) and affected maternal care by altered mating behavior (Mashoodh *et al.*, 2012) and prove germline dependence (Bohacek and Mansuy, 2017). Consistent with the significant changes of miRNAs and tsRNAs in the germline 2 weeks post GR activation, previous studies including our own have observed regulation of mouse sperm small RNA in a variety of contexts (Rodgers *et al.*, 2013; Gapp *et al.*, 2014; Grandjean *et al.*, 2015; Sharma *et al.*, 2015; Chen *et al.*, 2016; Wu *et al.*, 2016; Short *et al.*, 2017; Benito *et al.*, 2018; Rompala *et al.*, 2018; Yeshurun and Hannan, 2018).

Especially relevant specifically for our analysis, sperm RNA sequencing after drinking water administration of corticosterone for 4 weeks followed by mating, led to strong downregulation of tsRNA-GluCTC and tsRNA CysGCA, two of our top down-regulated tsRNAs, indicating that these tsRNAs are responding similarly to acute and chronic insults. At the same time this chronic manipulation elicits changes of several miRNAs, e.g. 34c and 471 (Short *et al.*, 2016), albeit in the opposite direction of what we find in response to acute Dex treatment. These discrepancies may arise either by the Dex induced short-term suppression of internal corticosteroid (Barden, 1999) or due to adaptations in response to chronic administration.

While four (Petropoulos, Matthews and Szyf, 2014; Short *et al.*, 2016; Wu *et al.*, 2016; Cartier *et al.*, 2018) out of five (Bönisch *et al.*, 2016) previous studies did report phenotypic effects following chronic paternal Dex exposure, only two assessed sperm small RNAs to associate the alterations to the sperm RNA payload (Bönisch *et al.*, 2016; Cartier *et al.*, 2018) yielding conflicting outcomes.

These differences might be due to inconsistent life stages (adulthood versus gestational), sperm collection (swim up, somatic lysis, or no purification) and/or dosage of exposure. Depending on the dosage and timing, the complex autoregulation of the GR can lead to GR downregulation after prolonged activation (Gjerstad, Lightman and Spiga, 2018). Acute exposures have the advantage of avoiding such long-term feedback regulation, and hence

provide an elegant approach for studying the signaling pathways leading to germline changes.

Mature sperm tsRNAs and miRNAs have been shown to be acquired during epididymal transit (Sharma *et al.*, 2015, 2018) and miRNAs are necessary for early embryonic development (Liu *et al.*, 2012; Conine *et al.*, 2018), although these results have been disputed by the fact that caput derived mouse sperm cells can be used for intra cytoplasmatic sperm injection (Zhou *et al.*, 2019) and the creation of viable offspring of said miRNA KO mice (Yuan *et al.*, 2015). Furthermore, a recent publication suggests that chronic stress induced sperm miRNAs are taken up primarily from epididymisomes originating from the caput epididymis or the proximal epididymal tract (Chan *et al.*, 2020). Chronic nutritional manipulation with effects on offspring also report the necessity of epididymal transit to acquire tsRNA changes in sperm(Sharma *et al.,* 2015). Harvesting mature sperm 3 hours after exposure yields a population enriched for cells that had been exposed while already in the cauda epididymis, where spermatids reside for roughly 5 days (Meistrich, 1975). These cells have not traveled through the epididymis nor have they had a chance to potentially take up small RNA from caput-derived epididymisomes after Dex administration. As expected, we detect no changes in miRNAs in these samples. We do however detect changes in tsRNAs 3 hours post Dex, some of which even persist 14 days post injection, challenging the current view that all sperm tsRNAs are taken up during epididymal transit. The discrepancies between results in the context of chronic and our acute intervention might depend on the post treatment interval. Chronic interventions - based on their experimental design - did not assess changes in mature sperm soon after the first intervention. Their claims rely on a comparison of sperm harvested from different tracts of the epididymis and might require conceptual reevaluation. An additional option for sperm RNA alterations in transcriptionally inert sperms was suggested in a recent study that found mitochondrial tRNA cleavage in the T-loop in response to a one-week high sugar diet in humans(Nätt *et al.*, 2019). In line with this observation, Dex injection could trigger oxidative stress(Bera *et al.*, 2010) that provokes such cleavage to increase tsRNA levels(Thompson *et al.*, 2008). A role for oxidative stress in

sperm RNA dynamics is further supported by a recent study in boar sperm that found seasonal differences in sperm small and long RNA associated with changed abundance of transcripts mapping to oxidative stress-, DNA damage- and autophagy- related genes (Gòdia *et al.*, 2019).  Such potentially oxidative stress mediated mechanism does not explain though a rapid decrease of tsRNAs 3 hours post Dex injection.

Importantly, we show alterations in sperm tsRNAs that persist in the oocyte concomitant with changes in early embryonic gene expression and a metabolic phenotype in adulthood. tsRNAs and tRNA-Gly derived fragments in particular are known to induce chromatin structure mediated gene regulation and to regulate cell differentiation in various contexts (Li *et al.*, 2016; Guzzi and Bellodi, 2020). Hence, we propose that the transmitted reduction in key tsRNAs such as Gly-GCC and Gly-CCC explains in part the observed perturbations during late two cell embryo developmental stage. This might reflect an accelerated developmental transcriptional program in the preimplantation embryo of Dex injected males, ultimately resulting in aberrant BMI and glucose metabolism later in life.

Additionally, we have discovered alterations in circRNA abundance in mature sperm that might also impact the developmental program in the early embryo. CircRNAs have the potential to be translated into proteins via back-splicing(Legnini *et al.*, 2017). Accordingly, they are crucial contributors to spermiogenesis post transcriptional cessation, since they provide a stable alternative to linear mRNA templates for protein translation(Tang *et al.*, 2020). Transmitted sperm-circRNAs could likewise contribute to translation post fertilization, yet the unconventional lattice-state of ribosomes preventing normal rates of translation post transcription(Israel *et al.*, 2019) accompanied by a rapid increase in proteins of the ubiquitine/proteasome pathway(Wang *et al.*, 2010) make this unlikely.  Never the less a study on human sperm detected abundant levels of circRNA with predicted regulatory function of early developmental genes in sperm heads, suggesting transmission and function post fertilization(Chioccarelli *et al.*, 2019).  By sponging miRNAs that regulate early embryonic transcripts, circRNA could amplify minute signals from paternal environment, such

as might be the case for the gene transcript Sat1, that displays increased expression in Dex offspring in 2-cell embryos from cluster 1.

Besides altered RNA identity, nucleic acid modifications especially of RNA but also DNA methylation and chromatin accessibility might contribute further to the effects of Dex injections on offspring metabolism. While detection of changes in each player should be subject of further investigation and might reveal a glimpse of their potential implication, proof of the individual relative causal contribution is extremely challenging since they likely require tight interaction to unravel their orchestrated effects.

Finally, it might be useful to consider testing the translatability of our findings to humans. Here we investigated the effects of a single Dex administration soon after the injection in mice, mimicking a single GR activation such as elicited by treatment of an acute asthma exacerbation (Cross, Paul and Goldman, 2011). The recent report that Dex can reduce the number of deaths associated with the Covid-19 pandemic ('Biggest COVID-19 trial tests repurposed drugs first', 2020), further prompts the re-evaluation of the impact of prolonged Dex treatment on offspring phenotype. From a clinical perspective, additional consideration is warranted for consequences on offspring health when extended time has elapsed between treatment and time of conception. Such designs may pave the way for the extrapolation of our findings.

We conclude that acute Dex treatment can induce germline epimodifications in the form of small and long non-coding RNA, which likely are relevant in the transmission of the effects of single traumatic events on offspring well-being. Our data suggest that sperm small RNAs are not solely regulated through epididysomal uptake during transition from caput to cauda epididymis. This expands the interpretation from chronic dietary and stress exposures(Sharma *et al.*, 2015; Chan *et al.*, 2020), where uptake of tsRNAs and miRNAs via epididymisomes has been suggested to lead to differential sperm payload, yet required sperm to transit from caput to cauda to bring about the changes. A persistent detection of significant fold changes of the exact same sperm small RNA in the embryo suggests functional implication in the information transfer from father to offspring. Together with

potentially transmitted miRNA sponges in the form of circRNAs, this likely contributes to a slight developmental acceleration of gene expression programs in the early embryo and ultimately manifests in a metabolic phenotype. Future studies may aim at testing the causal contribution of specific sperm RNAs to the transmission of effects of acute impacts. Certainly, continuous methodological refinement will help dissect the relative implication and the interplay of the distinct germline modifications such as DNA-methylation, histone-PTMs and chromatin architecture in this highly complex process.

### *Acknowledgements*

We thank Wayo Matsushima and Pierre-Luc Germain for valuable advice on bulk sequencing analysis and the Sanger RSF for help with animal care, especially the teams of Michael Woods and Francesca Flack. We also thank Sergio Mompart Barrenechea and Jasmine Kaur for help with the q-RT-PCR.

KG was funded by the Swiss National Science Foundation early postdoc and advanced postdoc mobility a SPARK and Novartis foundation grant. Some of this work was supported by Cancer Research UK (C13474/A18583, C6946/A14492) and Wellcome (104640/Z/14/Z, 092096/Z/10/Z) to EAM. GP and MH were supported by a core grant from the Wellcome Trust. The lab of JB is currently funded by the ETH Zurich, SNSF Project Grant 310030_172889/1, ETH Research Grant ETH-20 19-1, Kurt und Senta Herrmann-Stiftung, and Botnar Research Center for Child Health. For the purpose of Open Access, the author has applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission.

### *Author´s contributions*

KG performed animal exposures, collected samples and prepared sequencing libraries. EG performed IVF and embryo culture. GP analysed single cell sequencing data. MH supervised the single-embryo sequencing analysis. AC and FG analyzed bulk sequencing data. KG, JB

and EAM designed the study, interpreted the results and wrote the manuscript with input

from the other authors.

Futher information and request for resources should be directed to and will be fulfilled by the

lead contact, Katharina Gapp katharina.gapp@hest.ethz.ch.

***Conflict of interests***

The authors declare no competing interests. E.A.M. is a founder and Director of STORM

Therapeutics Ltd. STORM Therapeutics had no role in the design of the study and collection,

analysis, and interpretation of data as well as in writing the manuscript.

**Figure legends**

**Figure 1** Effects of Dex on small RNA payload of sperm cells residing in testis at time of

administration. (A) Experimental design depicting a time window of two weeks between

injection of Dex and sperm collection for molecular analysis. (B) Volcano plot depicting fold

changes and significance level of miRNAs and tsRNAs (C) in mature sperm 14 days post

injection of Dex (n=8) versus vehicle (n=8) as assessed by small RNA sequencing.

**Figure 2** Effect of Dex on sperm cells at different time points post Dex administration. (A)

Experimental design showing the location of sperm at time of injection and timing of sperm

harvest. MA (log-intensity ratios (M-values) versus log-intensity averages (A-values) plots

depicting (B) effect of Dex (log2 fold changes control versus dexamethasone), (C) of time

post injection (log2 fold changes 7 days versus 3 hours) (7 days Dex n =4 and controls n =4,

3 hours Dex n =3 and controls n =4). TsRNAs are indicated by sequence identity for display

only, each dot represents one small RNA. MA plot depicts log2 fold changes on the y axis

and the expression level on the x axis (the higher the expression the further to the right).

Statistically significantly changed small RNAs are highlighted in red. (D) Relative expression

of ArgCCT-2 as obtained by q-RT-PCR (cauda: Dex n=4, controls n=4, caput: Dex n=4,

controls n=5; interaction F(1,13)=6.34, p=0.0257, treatment F(1,13)=5.97, p=0.0040, site of

collection (F(1,13)=12.15, p=0.0296; cauda control versus cauda Dex t(13)=3.42, p=0.0274,

cauda Dex verus caput Dex t(13)=4.137, p=0.007). Whiskers display minimum and

maximum.

**Figure 3** Effect of Dex on metabolic phenotype in the offspring (A) Experimental design

depicting timeline between injection, sperm harvest, in vitro fertilization and phenotyping. (B)

Impact of Dex on male and female adult offspring (B) Body mass index (males vehicle

offspring n=21, Dex offspring n=22, females vehicle offspring n=17, Dex offspring n=17) (C)

glucose tolerance (males vehicle offspring n=12, Dex offspring n=12, females vehicle

offspring n=12, Dex offspring n=12) and (D) insulin tolerance (males vehicle offspring n=9,

Dex offspring n=8, females vehicle offspring n=12, Dex offspring n=12). Error bars display

SEM. Detailed statistical results are depicted in Supplementary Fig. 5, raw data are provided

in supplementary table 3).

**Figure 4 Effect of paternal Dex injection on embryonic offspring small RNA** (A)

Experimental design depicting timeline between injection, sperm harvest, in vitro fertilization

and small RNA sequencing at 2-cell stage. (B) Volcano plot showing effect of paternal Dex

on small RNA tsRNAs (vehicle embryonic offspring n=5, Dex embryonic offspring n=4).

TsRNAs are grouped by sequence identity for display only.

**Figure 5** Effect of paternal Dex injection on embryonic offspring long RNA transcriptome (A)

Experimental design depicting timeline between injection, sperm harvest, in vitro fertilization

and Smartseq2 sequencing at 2-cell stage. (B) Consensus matrix representing the similarity

between cells as reported by SC3. Similarity 0 indicates that a given pair of embryos were

never assigned to the same cluster, whereas similarity 1 means that a pair of embryos were

always assigned to the same cluster.  (C) Sankey diagram showing projection of the

obtained clusters (C1 and C2) into clusters reported by Deng *et al.* for single cells obtained

from 2 cell embryos. (D) Principal component analysis of two-cell embryos gene expression.

The top panel indicates the density of 2 cell embryos along PC1 grouped by condition;

control (red) and Dex treatment (blue). The other two bottom panels show the distribution of

2 cell embryos across PC1 and PC2 for control (red) and treated (blue) groups. The cluster

membership of each embryo is denoted by the point shapes (C1 cycles; C2 triangles) and

centroids of each cluster is indicated with a black dot overlaid with an x. Wilcox tests were

performed to assess differences on PC1 values of C1 and C2 clusters between the treated

and control groups. NS denotes non-significant change for C2 cluster, while * indicates a

significant difference for C1 cluster (p-value< 0.05). (E) Silhouette coefficient comparison

between treatment and control, statistical significance was assessed with Wilcox test (** p-

value < 0.01; *** p-value < 0.005) (F) Selection of differentially expressed genes as

determined by Monocle within C1 corresponding to late 2-cell embryo stage (*** adjusted p-

value < 0.005).

**Figure 6** Effects of Dex on long RNA payload of sperm cells residing in testis at time of

administration. (A) Experimental design depicting a time window of two weeks between

injection of Dex and sperm collection for molecular analysis.

(B) Volcano plot depicting fold changes and significance level of long RNA in mature sperm

14 days post injection of Dex (n=4) versus vehicle (n=4) as assessed by small RNA

sequencing. (C) Heatmap showing significantly differentially expressed long RNA transcripts

of the same experiment (multiple comparison corrected, q<0.05).

**Materials and Methods**

*Animals*

C57Bl/6 mice were obtained from the Sanger Research support facility in-house-breeding

colony. They were housed in a temperature and humidity-controlled facility in individually

ventilated cages under a non-reversed light-dark cycle (Sanger Research support facility) or

a reversed-light-dark cycle (ETH EPIC). Standard chow (LabDiet(r) 5021-3 supplied by IPS)

and water were provided *ad libitum after weaning* unless stated otherwise (e.g. oocyte

donors). Breeding colony was provided SAFE R03-10 breeding diet, supplied by SAFE diets.

Experimental procedures were performed during the animals' inactive cycle at Sanger. Age

and weight matched (margin of one week) males were used in each experimental group

receiving Dex injections. Animals used for Dex injection followed by sperm sequencing were

all sexually mature (14 and 7 days or 3 hours post treatment were 13, 11 and 9 weeks of age

respectively) at the time of sperm collection.

C57Bl/6 males used for sperm sequencing 14 days post Dex injection and q-PCR

experiments/validation 3 hours post Dex were obtained from the ETH`s EPIC in house

breeding colony in Zürich and were 14-18 weeks old. These mice were fed chow #3734 by

Kliba/Granovit.

IVF oocyte donor females and embryo recipients were fed SAFE R03-10 breeding diet,

supplied by SAFE diets until 10 days post embryo transfer. Until this time embryo recipients

were housed in pairs after which they were split into single housing. IVF offspring was

weaned at PND21 and assigned to cages avoiding littermate cohousing. Offspring

phenotyping was carried out between 3.5 to 4 months and necropsy at 4.5 months of age in

balanced (offspring controls, offspring treatment) and age matched groups (all animals had

an age spread of 3 days). Animals were housed in groups of 4-5 mice/cage in the Sager

Institute barrier research support facility (all animals apart from animals for q-RT-PCR

experiments) and ETHZ`s EPIC facility (animals for q-RT-PCR).

All experiments were approved by the UK home office (project license P176396F2) and

Cantonal commission for animal experimentation Zürich (project license ZH222/19).

*Dex treatment and sample collection*

Age matched males with an age spread of 1 week were randomly assigned to control and

treatment groups. Males were injected with either 2mg/kg of Dex in 10 % DMSO, 0.9% saline

or vehicle (10%DMSO in 0.9% saline). Males used for sperm collection did not undergo any

metabolic testing. They were sacrificed 2 weeks, 7 days and 3 hours after Dex or vehicle

treatment.  Cauda epididymis and vas deference were dissected and placed in M2 medium.

After allowing sperm to diffuse into M2 medium, cells were pelleted by short centrifugation

and washed with PBS. For sperm RNA sequencing and q-PCR, mature sperm cells were

separated from potential somatic contamination by somatic lysis, followed by 2 washes with

PBS (Brykczynska *et al.*, 2010).  Sperm counts and fertilization rate appeared unaffected

post Dex injection (Supplementary Fig. 10&11).

*Superovulation, in vitro fertilization and embryo culture*

12 randomly selected, C57BL/6 females were superovulated at 26-31 days of age with Card

Hyperova (Cosmo Bio, KYD-010-EX-X5), followed by 7.5 IU human chorionic gonadotrophin

(HCG) 48 hours later.

Cumulus-oocyte complexes (COCs) were released from the ampulla of the oviduct 16-17

hours after HCG administration, and preincubated in high calcium HTF with Glutathione

medium for 30-60 minutes (in $CO_2$ incubator at 37 deg C, 5% $CO_2$ in air) before

insemination. Frozen sperm used for insemination was pooled from 2 males that had been

injected with Dex or vehicle 14 days prior to cryopreservation. Thawed sperm was

preincubated for 30 minutes in TYH (with Methyl-b-cyclodextrin, Sigma C4555) medium at 37

deg C, 5% $CO_2$ in air, before being added to the COC complexes for fertilisation. 4 hours

after insemination, the presumptive zygotes were washed through several drops of KSOM

(Millipore, MR-121-D) and incubated overnight in KSOM.

For in vivo offspring, 14-20 x 2 cell embryos from overnight culture in 6 individual IVF dishes

/group were implanted into 0.5 dpc pseudo-pregnant F1 females (6 females/group). Each

dish contained oocytes from one female with the exception of 2 dishes (out of 6) in the Dex

group that contained oocytes of the same female, since one female failed to super-ovulate.

For molecular (single) embryo gene expression analysis at the two cell stage, 2-cell-embryos

from overnight culture were frozen, and after thawing briefly cultured in preincubated KSOM

until/during plating into 96 well plates. The females used to generate these embryos were

superovulated with PMSG. The IVF Protocol is based on EMMA Harwell's protocol (adapted

from Takeo & Nakagata 2011(Nakagata, 2011)), and the Sperm Freeze Protocol is based on

Ostermeier G.C. et al (2008)(Ostermeier *et al.*, 2008). Resulting litter sizes did not differ

between Vehicle and Dex injected offspring (Supplementary Fig. 12)

*Sperm and embryo RNA extraction*

Total RNA was prepared from adult mouse sperm using Trizol (Thermo Scientific 15596026)

and Directzol (Zymo R2080). Total RNA was prepared from zygotes using the Trizol LS

protocol. Quantity and purity of RNA were determined by Agilent 2100 Bioanalyser (Agilent

Technologies) and Qubit fluorometer (Life Technologies). Absence of prominent ribosomal

peaks indicated absence of somatic cell contamination.

*Sperm RNA sequencing (RNAseq)*

Sequencing was done using an Illumina Genome Analyzer HiSeq 2500 (Illumina) in Rapid

run mode for long 100bp and small 50 bp RNA sequencing runs respectively.

Libraries for long RNA sequencing were prepared using the TruSeq Stranded Total RNA kit

according to the manufacturer's instructions with indices diluted at 1:3. 200 ng of total sperm

RNA was subjected to removal of rRNA using Ribozero gold kit. Approximately 100ng of

sperm RNA and total RNA of several 2-cell zygotes was subjected to TruSeq or Nextflex

(sperm 14 days post injection) small RNA library preparation following the manufacturer's

recommendations with the following modifications: adaptors were diluted 1:4 and PCR cycles

were augmented to 18 and 22 (Nextflex) PCR cycles respectively. When library preparation

of samples was split across days groups were balanced to circumvent batch effects.

*Single embryo seq.*

2 cell embryos were generated using the same conditions as indicated for in vivo offspring

yet followed by embryo cryopreservation until processing for library preparation. They were

thawed and those that appeared intact (34 controls and 37 Dex) pipetted into wells of 2 96

well culture plates containing lysis buffer and stored at -80°C before processing according to

the Smartseq 2 protocol and manufacturer's recommendations (Nextera). Libraries contained

a 1:19 Million dilution of External RNA Controls Consortium (ERCC) spike-ins (4456740 Ambion)

and were amplified for 18 PCR cycles. Sequencing was performed on a HiSeq V4 under

paired end 75bp mode.

*Insulin and Glucose tolerance test*

Animals were fasted 4 hours to establish a shared baseline glucose level. They received a

single injection of insulin (insulin: 1mU/g body weight) (Actrapid Novo Nordisk), glucose

(2mg/g body weight) or vehicle (saline) intraperitoneally. Blood samples were taken from

lateral tail vein in adult animals to assess blood glucose level using an Accuckeck aviva

device.

*Body mass index*

Animal lengths were measured using a standard ruler and weighed for assessing body

weight. Body mass index was calculated using the following formula: weight (g)/(length

(cm)^2).

*Necropsy*

Organs were dissected after sacrifice and weighed immediately on a scale using "g" as a unit

with an accuracy of 2 decimals (accurate down to 10 mg).

*q-RT-PCR*

5ng/sample RNA isolated from sperm was reverse transcribed (RT) using the miCURY LNA

RT kit (Qiagen #339340). Quantitative RT-PCR (qRT-PCR) was performed using SYBR

green based detection in a Biorad thermal cycler with MiRCURY LNA-based small RNA

probes designed against 5`end of tRNA ArgCCT-2

(5`GCCCCAGUGGCCUAAUGGAUAAGGCACUGGCC3`) with a polyA tail directed reverse

miRCURY primer (Qiagen # 339317). U6 was used as an internal control (Qiagen # 339306).

*Bioinformatic and statistical data analysis*

*Sperm RNA sequencing*

Each sequencing library represented sperm harvested from a single male.  Sequencing

quality was assessed with FastQC (Andrews, 2010) and MultiQC (Ewels *et al.*, 2016).

Adapters were removed from the 3' ends with cutadap t(Martin, 2011) (version 1.14) and

resulting sequences with 14 nucleotides of length or less were discarded. All other reads

were aligned end to end (no soft clipping) to the ENSEMBL *Mus musculus* genome (release

75) (Flicek *et al.*, 2014) with STAR (Dobin *et al.*, 2013). No mismatches were allowed.

Featurecounts was used to match the alignments against the miRbase(Kozomara and

Griffiths-Jones, 2011) annotation (version 21) and obtain a matrix of miRNA counts. We

applied fractional counts whenever alignment occurred at multiple genomic locations.

Differential expression was analyzed using DESeq2 (Love, Huber and Anders, 2014).

Quantification of tRNA fragments was performed as above, but all CCA-3☐ trinucleotides

were trimmed after adapter removal, sequences with 15 nucleotides or less were

subsequently discarded and GtRNAdb (Chan and Lowe, 2016) annotation (GRCm38/mm10)

was used to obtain the count matrix.

For the data set collected 14 days after Dex injection, library preparation included the

insertion of 2 random tetranucleotides between read and adapters. By including only unique

sequences in the analysis we removed duplicates due to PCR amplification.

Long RNAseq libraries were pre-processed with trimmomatic (Bolger, Lohse and Usadel, 2014)

to remove adapters. Reads were aligned to the genome using STAR (Dobin *et al.*, 2013) and

quantified using featurecounts (Liao, Smyth and Shi, 2014). Circular RNAs were quantified using

Circexplorer2 (Zhang *et al.*, 2016) based on junction reads as detected by STAR. Differential

expression analysis was performed on the combined set of counts for circular and non-circular RNAs

using edgeR (Anders *et al.*, 2013). Robust estimation of dispersion was used to avoid spurious

significance due to outliers.

*2-cell single embryo sequencing analysis*

Reads from 2 cell embryos were mapped to the mouse reference genome (mm10) and

ERCC spike-ins using STAR (Dobin *et al.*, 2013). Resultant alignments were processed to

quantify the expression of annotated genes by GENCODE (vM11) and ERCC spike-ins using

featureCounts (Liao, Smyth and Shi, 2014). To filter low-quality sequenced embryos we only

considered those which had a total read count of at least 0.5 million reads with less than 15%

and 10% their read counts mapping to mitochondrial genes and   ERCC spike-ins

respectively. After these filters were applied a total of 56 embryos (29 controls and 27

treated) remained. We clustered their gene expression profiles using SC3 (Kiselev *et al.*,

2017) obtaining two main clusters (C1 and C2). Using scmap (Kiselev, Yiu and Hemberg,

2018), we projected the gene expression profiles for the two cell embryos onto an index

containing  expression profiles from zygotic, early/mid/late 2 cell embryos and 4 cell embryo

cells reported by Deng *et al.* (Deng *et al.*, 2014). We performed PCA analyses using scater

(McCarthy *et al.*, 2017) (runPCA function) and we calculated the PC1 silhouette coefficient

using in-house R scripts. To perform differential gene expression analyses we normalized

the read counts of each embryo as FPKM and we used Census (Qiu *et al.*, 2017) algorithm to

convert these values into relative transcripts counts. We computed the obtained 'Census

counts' using Monocle (v 2.99.2), assuming a negative binomial distribution and a lower detection limit of 0.5. We performed differential gene expression analyses between the total treated and control embryos, and also between the treated and control embryos inside of C1 and C2 clusters.

*Remaining statistical analyses.*

Sample size for in vivo offspring phenotyping was estimated based on previous work on similar models (Hoyer *et al.*, 2013; Bohacek, von Werdt and Mansuy, 2016). 3-Way repeated measures ANOVA was used to assess statistical significance for BMI, GTT and ITT measurements. Necropsy data were analysed using 3-way ANOVA followed by multiple t-tests and corrected for multiple comparisons using the Benjamini-Hochberg method. Normality was assessed with the Kolmogorov Smirnov test and met in all necropsy data. Homogeneity of variances was assessed and met in all necropsy data unless gonadal WAT. All t-tests did not assume homogeneity of variances (applied Welchs correction). All statistics of behavioural, metabolic tests and q-RT-PCR were computed with Prism. Outliers were removed from q-PCR results using Prism^s inbuild ROUT method and are depicted in the supplementary table 3 containing raw data with a star. Q-RT-PCR results were anaylysed with a 2-Way Anova followed by posthoc tests to compare individual groups applying the Bonferroni correction for multiple comparisons. All reported replicates were biological replicates. Significance was set at $p < 0.05$ for all tests.

*Availability of Data.*

The datasets supporting the conclusions of this article are included within the article (supplementary tables). All sequencing data were deposited to Gene Omnibus (accession number: GSE162112)  and ENA (accession number: ERP105660). All code is available on Github (https://github.com/ETHZ-INS/Sperm-RNA-Dex)

**Supplementary Material**

This article contains supplementary Figures and tables. Supplementary figures are compiled

in one word document.

**Supplementary table legend**

*Supplementary Table 1*

Sheet 1: List of normalized miRNA read counts of sperm harvested 14 days post Dex and

vehicle injection.

Sheet 2: Deseq2 results of a comparison between miRNA from sperm harvested 14 days

post Dex and vehicle injection.

Sheet 3: List of normalized tsRNA read counts of sperm harvested 14 days post Dex and

vehicle injection.

Sheet 4: Deseq2 results of a comparison between tsRNAs from sperm harvested 14 days

post Dex and vehicle injection.

*Supplementary table 2*

Sheet 1: Deseq2 results of the comparison of tsRNAs in sperm harvested 3 hours and 7

days post Dex with tsRNAs in sperm harvested 3 hours and 7 days post vehicle injection.

Sheet 2: Deseq2 results of the comparison between tsRNAs from sperm harvested post Dex

and vehicle injection at 3 hours with tsRNAs in sperm harvested 7 days post Dex and vehicle

injection.

Sheet 3: Deseq2 results of the interaction between treatment and time for tsRNAs from

sperm harvested 3 hours and 7 days post Dex and vehicle injection.

Sheet 4: Deseq2 results of the comparison of miRNAs in sperm harvested 3 hours and 7

days post Dex with miRNAs in sperm harvested 3 hours and 7 days post vehicle injection.

Sheet 5: Deseq2 results of the comparison between miRNAs from sperm harvested post Dex

and vehicle injection at 3 hours with miRNAs in sperm harvested  7 days post Dex and

vehicle injection.

Sheet 6: Deseq2 results of the interaction between treatment and time for miRNAs from

sperm harvested 3 hours and 7 days post Dex and vehicle injection.

Sheet 7: Raw values of qRT-PCR analysis for tsRNA-Arg-CCT-2 in caput and cauda sperm

sampled 3 hours post Dex and vehicle injection.

*Supplementary table 3*

Raw data for BMI (sheet 1), GTT (sheet 2) , ITT (sheet 3) and necropsy weights (sheet 4) of

adult offspring animals resulting from IVF of wildtype oocytes and sperm harvested 14 days

post Dex and vehicle injection.

*Supplementary table 4*

Sheet 1: Monocle output list on significantly differentially regulated genes between 2-cell

embryos resulting from IVF of wildtype oocytes and sperm harvested 14 days post Dex and

vehicle injection.

*Supplementary table 5*

Sheet 1: List of normalized long RNA seq counts of sperm harvested 14 days post Dex and

vehicle injection.

Sheet 2: EdgeR results of a comparison between long RNA reads from sperm harvested 14

days post Dex and vehicle injection.

**References**

Agarwal, V. *et al.* (2015) 'Predicting effective microRNA target sites in mammalian mRNAs', *eLife*, 4.

doi: 10.7554/eLife.05005.

Amanai, M., Brahmajosyula, M. and Perry, A. C. F. (2006) 'A restricted role for sperm-borne

microRNAs in mammalian fertilization.', *Biology of reproduction*, 75(6), pp. 877–884. doi:

10.1095/biolreprod.106.056499.

Anders, S. *et al.* (2013) 'Count-based differential expression analysis of RNA sequencing data using R and Bioconductor', *Nature Protocols*, 8(9), pp. 1765–1786. doi: 10.1038/nprot.2013.099.

Andrews, S. (2010) 'FastQC: a quality control tool for high throughput sequence data.'

Ashe, A. *et al.* (2012) 'piRNAs can trigger a multigenerational epigenetic memory in the germline of C. elegans', *Cell*. 2012/06/29, 150(1), pp. 88–99. doi: 10.1016/j.cell.2012.06.018.

Barden, N. (1999) 'Regulation of corticosteroid receptor gene expression in depression and antidepressant action.', *Journal of psychiatry & neuroscience : JPN*. Canadian Medical Association, 24(1), pp. 25–39. Available at: http://www.ncbi.nlm.nih.gov/pubmed/9987205.

Barrett, S. P. and Salzman, J. (2016) 'Circular RNAs: analysis, expression and potential functions', *Development*. Oxford University Press for The Company of Biologists Limited, 143(11), pp. 1838–1847. doi: 10.1242/DEV.128074.

Benito, E. *et al.* (2018) 'RNA-Dependent Intergenerational Inheritance of Enhanced Synaptic Plasticity after Environmental Enrichment', *Cell Reports*. Cold Spring Harbor Laboratory, 23(2), pp. 546–554. doi: 10.1016/j.celrep.2018.03.059.

Benyshek, D. C., Johnston, C. S. and Martin, J. F. (2006) 'Glucose metabolism is altered in the adequately-nourished grand-offspring (F3 generation) of rats malnourished during gestation and perinatal life', *Diabetologia*. 2006/03/25, 49(5), pp. 1117–1119. doi: 10.1007/s00125-006-0196-5.

Bera, S. *et al.* (2010) 'Dexamethasone-induced oxidative stress enhances myeloma cell radiosensitization while sparing normal bone marrow hematopoiesis', *Neoplasia*. Neoplasia Press, Inc., 12(12), pp. 980–992. doi: 10.1593/neo.101146.

'Biggest COVID-19 trial tests repurposed drugs first' (2020) *Nature Biotechnology*, 38(5), p. 510. doi: 10.1038/s41587-020-0528-x.

Bohacek, J. and Mansuy, I. M. (2017) 'A guide to designing germline-dependent epigenetic inheritance experiments in mammals.', *Nature methods*. Nature Publishing Group, 14(3), pp. 243–249. doi: 10.1038/nmeth.4181.

Bohacek, J. and Rassoulzadegan, M. (2019) 'Sperm RNA: Quo vadis?', *Seminars in Cell & Developmental Biology*. Academic Press. doi: 10.1016/J.SEMCDB.2019.07.005.

Bohacek, J., von Werdt, S. and Mansuy, I. M. (2016) 'Probing the germline-dependence of epigenetic inheritance using artificial insemination in mice', *Environmental Epigenetics*, 2(1), p. dvv015. doi: 10.1093/eep/dvv015.

Bolger, A. M., Lohse, M. and Usadel, B. (2014) 'Trimmomatic: a flexible trimmer for Illumina sequence data.', *Bioinformatics (Oxford, England)*. Oxford University Press, 30(15), pp. 2114–2120. doi: 10.1093/bioinformatics/btu170.

Bönisch, C. *et al.* (2016) 'Dexamethasone treatment alters insulin, leptin, and adiponectin levels in male mice as observed in DIO but does not lead to alterations of metabolic phenotypes in the offspring', *Mammalian Genome*. Springer, 27(1–2), pp. 17–28. doi: 10.1007/s00335-015-9616-5.

Bowers, M. E. and Yehuda, R. (2016) 'Intergenerational transmission of stress in humans', *Neuropsychopharmacology*, pp. 232–244. doi: 10.1038/npp.2015.247.

Brykczynska, U. *et al.* (2010) 'Repressive and active histone methylation mark distinct promoters in human and mouse spermatozoa', *Nat Struct Mol Biol*. Nature Publishing Group, 17(6), pp. 679–687. doi: http://www.nature.com/nsmb/journal/v17/n6/suppinfo/nsmb.1821_S1.html.

Cacciagli, P. *et al.* (2013) 'Mutations in BCAP31 Cause a Severe X-Linked Phenotype with Deafness, Dystonia, and Central Hypomyelination and Disorganize the Golgi Apparatus', *The American Journal of Human Genetics*, 93(3), pp. 579–586. doi: 10.1016/j.ajhg.2013.07.023.

Carone, B. R. *et al.* (2010) 'Paternally induced transgenerational environmental reprogramming of metabolic gene expression in mammals.', *Cell*, 143(7), pp. 1084–1096. doi: 10.1016/j.cell.2010.12.008.

Cartier, J. *et al.* (2018) 'Investigation into the role of the germline epigenome in the transmission of glucocorticoid-programmed effects across generations.', *Genome biology*. BioMed Central, 19(1), p. 50. doi: 10.1186/s13059-018-1422-4.

Chan, J. C. *et al.* (2020) 'Reproductive tract extracellular vesicles are sufficient to transmit intergenerational stress and program neurodevelopment', *Nature Communications*. Nature Publishing Group, 11(1), p. 1499. doi: 10.1038/s41467-020-15305-w.

Chan, P. P. and Lowe, T. M. (2016) 'GtRNAdb 2.0: an expanded database of transfer RNA genes identified in complete and draft genomes.', *Nucleic acids research*. Oxford University Press, 44(D1), pp. D184--9. doi: 10.1093/nar/gkv1309.

Chen, Q. *et al.* (2016) 'Sperm tsRNAs contribute to intergenerational inheritance of an acquired metabolic disorder', *Science*, 351(6271), pp. 397–400. doi: 10.1126/science.aad7977.

Chen, Q., Yan, W. and Duan, E. (2016) 'Epigenetic inheritance of acquired traits through sperm RNAs and sperm RNA modifications.', *Nature reviews. Genetics*. Nature Publishing Group, 17(12), pp. 733–743. doi: 10.1038/nrg.2016.106.

Chioccarelli, T. *et al.* (2019) 'Expression patterns of circular RNAs in high quality and poor quality human spermatozoa', *Frontiers in Endocrinology*, 10(JUL), pp. 1–14. doi: 10.3389/fendo.2019.00435.

Conine, C. C. *et al.* (2018) 'Small RNAs Gained during Epididymal Transit of Sperm Are Essential for Embryonic Development in Mice.', *Developmental cell*. Elsevier, 46(4), pp. 470--480.e3. doi: 10.1016/j.devcel.2018.06.024.

Cross, K. P., Paul, R. I. and Goldman, R. D. (2011) 'Single-dose dexamethasone for mild-to-moderate asthma exacerbations: effective, easy, and acceptable.', *Canadian family physician Medecin de famille canadien*. College of Family Physicians of Canada, 57(10), pp. 1134–1136. Available at: http://www.ncbi.nlm.nih.gov/pubmed/21998227.

Dadoune, J. and Alfonsi, M. (1984) 'Autoradiographic investigation of sperm transit through the male mouse genital tract after tritiated thymidine incorporation', *Reproduction Nutrition and Development*, 24(6), pp. 927–935. doi: 10.1051/rnd:19840709.

Dard-Dascot, C. *et al.* (2018) 'Systematic comparison of small RNA library preparation protocols for next-generation sequencing', *BMC Genomics*. BioMed Central, 19(1), p. 118. doi: 10.1186/s12864-018-4491-6.

Daskalakis, N. P. *et al.* (2012) 'Testing the cumulative stress and mismatch hypotheses of psychopathology in a rat model of early-life adversity', *Physiol Behav*. 2012/02/07, 106(5), pp. 707–721. doi: 10.1016/j.physbeh.2012.01.015.

Deevy, O. and Bracken, A. P. (2019) 'PRC2 functions in development and congenital disorders', *Development*. Oxford University Press for The Company of Biologists Limited, 146(19). doi: 10.1242/DEV.181354.

Deng, Q. *et al.* (2014) 'Single-Cell RNA-Seq Reveals Dynamic, Random Monoallelic Gene Expression in Mammalian Cells', *Science*. American Association for the Advancement of Science, 343(6167), pp. 193–196. doi: 10.1126/SCIENCE.1245316.

Dietz, D. M. *et al.* (2011) 'Paternal transmission of stress-induced pathologies.', *Biological psychiatry*, 70(5), pp. 408–414. doi: 10.1016/j.biopsych.2011.05.005.

Dobin, A. *et al.* (2013) 'STAR: ultrafast universal RNA-seq aligner.', *Bioinformatics (Oxford, England)*. Oxford University Press, 29(1), pp. 15–21. doi: 10.1093/bioinformatics/bts635.

EU Clinical Trials Register (no date) 'Randomised Evaluation of COVID-19 Therapy (RECOVERY)'. Available at: https://www.clinicaltrialsregister.eu/ctr-search/trial/2020-001113-21/GB (Accessed: 7 July 2020).

Ewels, P. *et al.* (2016) 'MultiQC: summarize analysis results for multiple tools and samples in a single

report', *Bioinformatics*. Oxford University Press, 32(19), pp. 3047–3048. doi:

10.1093/bioinformatics/btw354.

Flicek, P. *et al.* (2014) 'Ensembl 2014.', *Nucleic acids research*. Oxford University Press, 42(Database

issue), pp. D749--55. doi: 10.1093/nar/gkt1196.

Franklin, T. B. *et al.* (2010) 'Epigenetic Transmission of the Impact of Early Stress Across

Generations', *Biological Psychiatry*, 68(5), pp. 408–415. doi: 10.1016/j.biopsych.2010.05.036.

Franklin, T. B. *et al.* (2011) 'Influence of early stress on social abilities and serotonergic functions

across generations in mice.', *PloS one*. Edited by C. Combs, 6(7), p. e21842. doi:

10.1371/journal.pone.0021842.

Fullston, T. *et al.* (2013) 'Paternal obesity initiates metabolic disturbances in two generations of mice

with incomplete penetrance to the F2 generation and alters the transcriptional profile of testis and

sperm microRNA content', *FASEB Journal*, 27(10), pp. 4226–4243. doi: 10.1096/fj.12-224048.

Gapp, K. *et al.* (2014) 'Implication of sperm RNAs in transgenerational inheritance of the effects of

early trauma in mice', *Nature neuroscience*. 2014/04/15. Nature Publishing Group, a division of

Macmillan Publishers Limited. All Rights Reserved., 17(5), pp. 667–669. doi: 10.1038/nn.3695.

Gapp, K. *et al.* (2018) 'Alterations in sperm long RNA contribute to the epigenetic inheritance of the

effects of postnatal trauma', *Molecular Psychiatry*. Nature Publishing Group, p. 1. doi:

10.1038/s41380-018-0271-6.

Gapp, K. and Bohacek, J. (2017) 'Epigenetic germline inheritance in mammals: Looking to the past to

understand the future', *Genes, Brain and Behavior*. doi: 10.1111/gbb.12407.

Gapp, K. and Bohacek, J. (2018) 'Epigenetic germline inheritance in mammals: looking to the past to

understand the future', *Genes, Brain and Behavior*, 17(3). doi: 10.1111/gbb.12407.

Gjerstad, J. K., Lightman, S. L. and Spiga, F. (2018) 'Role of glucocorticoid negative feedback in the

regulation of HPA axis pulsatility.', *Stress (Amsterdam, Netherlands)*. Taylor & Francis, 21(5), pp.

403–416. doi: 10.1080/10253890.2018.1470238.

Gòdia, M. *et al.* (2019) 'A RNA-seq analysis to describe the boar sperm transcriptome and its

seasonal changes', *Frontiers in Genetics*, 10(MAR), pp. 1–14. doi: 10.3389/fgene.2019.00299.

Gòdia, M. *et al.* (2020) 'Identification of circular RNAs in porcine sperm and evaluation of their relation

to sperm motility', *Scientific Reports*, 10(1), pp. 1–11. doi: 10.1038/s41598-020-64711-z.

Grandjean, V. *et al.* (2015) 'RNA-mediated paternal heredity of diet-induced obesity and metabolic

disorders', *Sci Rep*. 2015/12/15, 5, p. 18193. doi: 10.1038/srep18193.

Grentzinger, T. *et al.* (2012) 'piRNA-mediated transgenerational inheritance of an acquired trait.',

*Genome research*, 22(10), pp. 1877–1888. doi: 10.1101/gr.136614.111.

Guzzi, N. and Bellodi, C. (2020) 'RNA Biology Novel insights into the emerging roles of tRNA-derived

fragments in mammalian development'. doi: 10.1080/15476286.2020.1732694.

Haeussler, S. and Claus, R. (2007) 'Expression of the glucocorticoid receptor and 11$β$-

hydroxysteroid dehydrogenase 2 in pig testes cells along fetal development', *Reproduction, Fertility

and Development*, 19(5), p. 664. doi: 10.1071/RD07033.

Hazra, R. *et al.* (2014) 'In Vivo Actions of the Sertoli Cell Glucocorticoid Receptor', *Endocrinology*,

155(3), pp. 1120–1130. doi: 10.1210/en.2013-1940.

Heijmans, B. T. *et al.* (2008) 'Persistent epigenetic differences associated with prenatal exposure to

famine in humans', *Proc Natl Acad Sci U S A*. 2008/10/29, 105(44), pp. 17046–17049. doi:

10.1073/pnas.0806560105.

Horby, P. *et al.* (2020) 'Effect of Dexamethasone in Hospitalized Patients with COVID-19: Preliminary

Report', *medRxiv*. Cold Spring Harbor Laboratory Press, p. 2020.06.22.20137273. doi:

10.1101/2020.06.22.20137273.

Hoyer, C. *et al.* (2013) 'Preconceptional paternal exposure to a single traumatic event affects postnatal

growth of female but not male offspring', 24(15). doi: 10.1097/WNR.0000000000000007.

Huber, M. H. R. and Bronson, F. H. (1980) 'Social modulation of spontaneous ejaculation in the

mouse', *Behavioral and Neural Biology*, 29(3), pp. 390–393. doi: 10.1016/S0163-1047(80)90373-8.

Israel, S. *et al.* (2019) 'An integrated genome-wide multi-omics analysis of gene expression dynamics

in the preimplantation mouse embryo', *Scientific Reports*. Springer US, 9(1), pp. 1–15. doi:

10.1038/s41598-019-49817-3.

Jimenez-Chillaron, J. C. *et al.* (2009) 'Intergenerational transmission of glucose intolerance and

obesity by in utero undernutrition in mice', *Diabetes*. 2008/11/20, 58(2), pp. 460–468. doi:

10.2337/db08-0490.

Jirtle, R. L. and Skinner, M. K. (2007) 'Environmental epigenomics and disease susceptibility', *Nature

reviews. Genetics*. 2007/03/17, 8(4), pp. 253–262. doi: 10.1038/nrg2045.

Kang, T. H. *et al.* (2013) 'The Housekeeping Gene Hypoxanthine Guanine Phosphoribosyltransferase

(HPRT) Regulates Multiple Developmental and Metabolic Pathways of Murine Embryonic Stem Cell

Neuronal Differentiation', *PLoS ONE*. Edited by A. J. Cooney. Public Library of Science, 8(10), p.

e74967. doi: 10.1371/journal.pone.0074967.

Kaufmann, S. H. *et al.* (1992) 'Evidence that rodent epididymal sperm contain the Mr approximately

94,000 glucocorticoid receptor but lack the Mr approximately 90,000 heat shock protein.',

*Endocrinology*, 130(5), pp. 3074–3084. doi: 10.1210/endo.130.5.1572314.

Kiselev, V. Y. *et al.* (2017) 'SC3: consensus clustering of single-cell RNA-seq data', *Nature Methods*.

Nature Publishing Group, 14(5), pp. 483–486. doi: 10.1038/nmeth.4236.

Kiselev, V. Y., Yiu, A. and Hemberg, M. (2018) 'scmap: projection of single-cell RNA-seq data across

data sets', *Nature Methods*. Nature Publishing Group, 15(5), pp. 359–362. doi: 10.1038/nmeth.4644.

de Kloet, E. R., Joëls, M. and Holsboer, F. (2005) 'Stress and the brain: from adaptation to disease.',

*Nature reviews. Neuroscience*, 6(6), pp. 463–475. doi: 10.1038/nrn1683.

de Kloet, E. R., Karst, H. and Joëls, M. (2008) 'Corticosteroid hormones in the central stress response:

Quick-and-slow', *Frontiers in Neuroendocrinology*, 29(2), pp. 268–272. doi:

10.1016/j.yfrne.2007.10.002.

Kozomara, A. and Griffiths-Jones, S. (2011) 'miRBase: integrating microRNA annotation and deep-

sequencing data', *Nucleic Acids Research*. Oxford University Press, 39(Issue suppl_1), pp. D152--

D157. doi: 10.1093/nar/gkq1027.

Krawetz, S. A. *et al.* (2011) 'A survey of small RNAs in human sperm', *Hum Reprod*. 2011/10/13,

26(12), pp. 3401–3412. doi: der329 [pii] 10.1093/humrep/der329.

Legnini, I. *et al.* (2017) 'Circ-ZNF609 Is a Circular RNA that Can Be Translated and Functions in

Myogenesis', *Molecular Cell*. Elsevier Inc., 66(1), pp. 22-37.e9. doi: 10.1016/j.molcel.2017.02.017.

Li, Q. *et al.* (2016) 'tRNA-Derived Small Non-Coding RNAs in Response to Ischemia Inhibit

Angiogenesis', *Scientific Reports*. Nature Publishing Group, 6(1), p. 20850. doi: 10.1038/srep20850.

Liao, Y., Smyth, G. K. and Shi, W. (2014) 'featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features', *Bioinformatics*. Oxford University Press, 30(7), pp.

923–930. doi: 10.1093/bioinformatics/btt656.

Liu, W.-M. *et al.* (2012) 'Sperm-borne microRNA-34c is required for the first cleavage division in

mouse', *Proceedings of the National Academy of Sciences of the United States of America*. National

Academy of Sciences, 109(2), pp. 490–494. doi: 10.1073/pnas.1110368109.

Love, M. I., Huber, W. and Anders, S. (2014) 'Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2', *Genome Biology*, 15(12), p. 550. doi: 10.1186/s13059-014-0550-8.

Macfarlan, T. S. *et al.* (2012) 'Embryonic stem cell potency fluctuates with endogenous retrovirus

activity.', *Nature*. NIH Public Access, 487(7405), pp. 57–63. doi: 10.1038/nature11244.

Marco, A. *et al.* (2014) 'Overweight and CpG methylation of the Pomc promoter in offspring of high-fat-

diet-fed dams are not "reprogrammed" by regular chow diet in rats.', *FASEB journal : official*

*publication of the Federation of American Societies for Experimental Biology*. The Federation of

American Societies for Experimental Biology, 28(9), pp. 4148–4157. doi: 10.1096/fj.14-255620.

Martin, M. (2011) 'Cutadapt removes adapter sequences from high-throughput sequencing reads',

*EMBnet.journal*, 17(1), p. 10. doi: 10.14806/ej.17.1.200.

Martínez, D. *et al.* (2014) 'In Utero Undernutrition in Male Mice Programs Liver Lipid Metabolism in the

Second-Generation Offspring Involving Altered Lxra DNA Methylation.', *Cell metabolism*, pp. 1–11.

doi: 10.1016/j.cmet.2014.03.026.

Mashoodh, R. *et al.* (2012) 'Paternal social enrichment effects on maternal behavior and offspring

growth', *Proc Natl Acad Sci U S A*. 2012/10/10, 109 Suppl, pp. 17232–17238. doi:

10.1073/pnas.1121083109.

McCarthy, D. J. *et al.* (2017) 'Scater: pre-processing, quality control, normalization and visualization of

single-cell RNA-seq data in R', *Bioinformatics*, p. btw777. doi: 10.1093/bioinformatics/btw777.

Meistrich, M. L. (1975) 'Alteration of epididymal sperm transport and maturation in mice by oestrogen

and testosterone', *Nature*, (1).

Miyazaki, T. *et al.* (2013) 'Functional Analysis of Tcl1 Using Tcl1-Deficient Mouse Embryonic Stem

Cells', *PLoS ONE*. Public Library of Science, 8(8). doi: 10.1371/JOURNAL.PONE.0071645.

Morgan, C. P. and Bale, T. L. (2011) 'Early Prenatal Stress Epigenetically Programs

Dysmasculinization in Second-Generation Offspring via the Paternal Lineage', *The Journal of*

*Neuroscience*, 31(33), pp. 11748–11755. doi: 10.1523/jneurosci.1887-11.2011.

Nakagata, N. (2011) 'Cryopreservation of Mouse Spermatozoa and In Vitro Fertilization', in. Humana

Press, pp. 57–73. doi: 10.1007/978-1-60761-974-1_4.

Nätt, D. *et al.* (2019) 'Human sperm displays rapid responses to diet', *PLoS Biology*, 17(12), pp. 1–25.

doi: 10.1371/journal.pbio.3000559.

Ostermeier, G. C. *et al.* (2004) 'Reproductive biology: Delivering spermatozoan RNA to the oocyte',

*Nature*, 429(6988), p. 154. doi: 10.1038/429154a.

Ostermeier, G. C. *et al.* (2008) 'Conserving, Distributing and Managing Genetically Modified Mouse

Lines by Sperm Cryopreservation', *PLoS ONE*. Edited by H. A. El-Shemy. Public Library of Science,

3(7), p. e2792. doi: 10.1371/journal.pone.0002792.

Peaston, A. E. *et al.* (2004) 'Retrotransposons Regulate Host Genes in Mouse Oocytes and

Preimplantation Embryos', *Developmental Cell*. Elsevier, 7(4), pp. 597–606. doi:

10.1016/J.DEVCEL.2004.09.004.

Pembrey, M. E. *et al.* (2006) 'Sex-specific, male-line transgenerational responses in humans',

*European Journal of Human Genetics*, 14, pp. 159–166. doi: 10.1038/sj.ejhg.5201538.

Peng, H. *et al.* (2012) 'A novel class of tRNA-derived small RNAs extremely enriched in mature mouse

sperm', *Cell Research*, 22(11), pp. 1609–1612. doi: 10.1038/cr.2012.141.

Pentinat, T. *et al.* (2010) 'Transgenerational inheritance of glucose intolerance in a mouse model of

neonatal overnutrition', *Endocrinology*. 2010/10/15, 151(12), pp. 5617–5623. doi: 10.1210/en.2010-

0684.

Petropoulos, S., Matthews, S. G. and Szyf, M. (2014) 'Adult Glucocorticoid Exposure Leads to

Transcriptional and DNA Methylation Changes in Nuclear Steroid Receptors in the Hippocampus and

Kidney of Mouse Male Offspring1', *Biology of Reproduction*, 90(2). doi:

10.1095/biolreprod.113.115899.

Qiu, X. *et al.* (2017) 'Single-cell mRNA quantification and differential analysis with Census', *Nature

Methods*, 14(3), pp. 309–315. doi: 10.1038/nmeth.4150.

Rodgers, A. B. *et al.* (2013) 'Paternal Stress Exposure Alters Sperm MicroRNA Content and

Reprograms Offspring HPA Stress Axis Regulation', *Journal of Neuroscience*. 2013/05/24, 33(21), pp.

9003–9012. doi: 10.1523/JNEUROSCI.0914-13.2013.

Rodgers, A. B. *et al.* (2015) 'Transgenerational epigenetic programming via sperm microRNA

recapitulates effects of paternal stress', *Proceedings of the National Academy of Sciences*, 112(44), p.

201508347. doi: 10.1073/pnas.1508347112.

Rompala, G. R. *et al.* (2018) 'Heavy Chronic Intermittent Ethanol Exposure Alters Small Noncoding

RNAs in Mouse Sperm and Epididymosomes.', *Frontiers in genetics*. Frontiers Media SA, 9, p. 32. doi:

10.3389/fgene.2018.00032.

Roth, T. L. *et al.* (2009) 'Lasting Epigenetic Influence of Early-Life Adversity on the BDNF Gene',

*Biological Psychiatry*, 65(9), pp. 760–769. Available at:

http://www.sciencedirect.com/science/article/B6T4S-4VCNF2J-

1/2/f7007e8e39f1f645dd25d9dbf97487fe.

Rousseeuw, P. J. (1987) 'Silhouettes: A graphical aid to the interpretation and validation of cluster

analysis', *Journal of Computational and Applied Mathematics*. North-Holland, 20, pp. 53–65. doi:

10.1016/0377-0427(87)90125-7.

Shankar, K. *et al.* (2010) 'Maternal overweight programs insulin and adiponectin signaling in the

offspring.', *Endocrinology*, 151(6), pp. 2577–2589. doi: 10.1210/en.2010-0017.

Sharma, U. *et al.* (2015) 'Biogenesis and function of tRNA fragments during sperm maturation and

fertilization in mammals', *Science*. NIH Public Access, 351(6271), pp. 391–396. doi:

10.1126/science.aad6780.

Sharma, U. *et al.* (2018) 'Small RNAs Are Trafficked from the Epididymis to Developing Mammalian

Sperm', *Developmental Cell*. Cell Press, 46(4), pp. 481--494.e6. doi: 10.1016/J.DEVCEL.2018.06.023.

Shirayama, M. *et al.* (2012) 'piRNAs initiate an epigenetic memory of nonself RNA in the C. elegans

germline', *Cell*. 2012/06/29, 150(1), pp. 65–77. doi: 10.1016/j.cell.2012.06.015.

Short, A. K. *et al.* (2016) 'Elevated paternal glucocorticoid exposure alters the small noncoding RNA

profile in sperm and modifies anxiety and depressive phenotypes in the offspring', *Translational*

*Psychiatry*, 6(6), p. e837. doi: 10.1038/tp.2016.109.

Short, A. K. *et al.* (2017) 'Exercise alters mouse sperm small noncoding RNAs and induces a

transgenerational modification of male offspring conditioned fear and anxiety.', *Translational*

*psychiatry*. Nature Publishing Group, 7(5), p. e1114. doi: 10.1038/tp.2017.82.

Silva, E. J. R. *et al.* (2010) 'Glucocorticoid receptor in the rat epididymis: Expression, cellular

distribution and regulation by steroid hormones', *Molecular and Cellular Endocrinology*, 325(1–2), pp.

64–77. doi: 10.1016/j.mce.2010.05.013.

Silva, E. J. R. *et al.* (2014) 'Impact of adrenalectomy and dexamethasone treatment on testicular

morphology and sperm parameters in rats: insights into the adrenal control of male reproduction',

*Andrology*, 2(6), pp. 835–846. doi: 10.1111/j.2047-2927.2014.00228.x.

Tang, C. *et al.* (2020) 'm6A-dependent biogenesis of circular RNAs in male germ cells', *Cell Research*.

Springer US, 30(3), pp. 211–228. doi: 10.1038/s41422-020-0279-8.

Tang, F. *et al.* (2007) 'Maternal microRNAs are essential for mouse zygotic development.', *Genes &*

*development*. Cold Spring Harbor Laboratory Press, 21(6), pp. 644–648. doi: 10.1101/gad.418707.

Thompson, D. M. *et al.* (2008) 'tRNA cleavage is a conserved response to oxidative stress in

eukaryotes', *Rna*, 14(10), pp. 2095–2103. doi: 10.1261/rna.1232808.

Tyebji, S., Hannan, A. J. and Tonkin, C. J. (2020) 'Pathogenic Infection in Male Mice Changes Sperm

Small RNA Profiles and Transgenerationally Alters Offspring Behavior', *Cell Reports*. Cell Press,

31(4), p. 107573. doi: 10.1016/J.CELREP.2020.107573.

Vassoler, F. M. *et al.* (2013) 'Epigenetic inheritance of a cocaine-resistance phenotype', *Nat Neurosci*. 2012/12/18, 16(1), pp. 42–47. doi: 10.1038/nn.3280.

Vojtech, L. *et al.* (2014) 'Exosomes in human semen carry a distinctive repertoire of small non-coding RNAs with potential regulatory functions', *Nucleic Acids Research*, 42(11), pp. 7290–7304. doi: 10.1093/nar/gku347.

Wang, S. *et al.* (2010) 'Proteome of mouse oocytes at different developmental stages', *Proceedings of the National Academy of Sciences of the United States of America*, 107(41), pp. 17639–17644. doi: 10.1073/pnas.1013185107.

Weiss, I. C. *et al.* (2011) 'Inheritable effect of unpredictable maternal separation on behavioral responses in mice', *Frontiers in Behavioral Neuroscience*, 5, p. doi: 10.3389/fnbeh.2011.00003. doi: 10.3389/fnbeh.2011.00003.

Wu, L. *et al.* (2016) 'Paternal psychological stress reprograms hepatic gluconeogenesis in offspring', *Cell Metabolism*, 23(4), pp. 735–743. doi: 10.1016/j.cmet.2016.01.014.

Wu, W., Ji, P. and Zhao, F. (2020) 'CircAtlas: an integrated resource of one million highly accurate circular RNAs from 1070 vertebrate transcriptomes', *Genome Biology*. BioMed Central, 21(1), p. 101. doi: 10.1186/s13059-020-02018-y.

Yeri, A. *et al.* (2018) 'Evaluation of commercially available small RNASeq library preparation kits using low input RNA', *BMC Genomics*. BioMed Central, 19(1), p. 331. doi: 10.1186/s12864-018-4726-6.

Yeshurun, S. and Hannan, A. J. (2018) 'Transgenerational epigenetic influences of paternal environmental exposures on brain function and predisposition to psychiatric disorders', *Molecular Psychiatry*, pp. 1–13. Available at: http://www.ncbi.nlm.nih.gov/pubmed/29520039.

Yu, N. *et al.* (2018) 'RBBP7 is a prognostic biomarker in patients with esophageal squamous cell carcinoma', *Oncology Letters*. Spandidos Publications, 16(6), pp. 7204–7211. doi: 10.3892/ol.2018.9543.

Yuan, S. *et al.* (2015) 'miR-34b/c and miR-449a/b/c are required for spermatogenesis, but not for the first cleavage division in mice', *Biology Open*, 4(2), pp. 212–223. doi: 10.1242/bio.201410959.

Zhang, X.-O. *et al.* (2016) 'Diverse alternative back-splicing and alternative splicing landscape of circular RNAs.', *Genome research*. Cold Spring Harbor Laboratory Press, 26(9), pp. 1277–1287. doi: 10.1101/gr.202895.115.

Zhang, Y. Y. *et al.* (2018) 'Dnmt2 mediates intergenerational transmission of paternally acquired metabolic disorders through sperm small non-coding RNAs', *Nature Cell Biology*. Springer US,

20(May). doi: 10.1038/s41556-018-0087-2.

Zhou, D. *et al.* (2019) 'Caput Epididymidal Mouse Sperm Support Full Development', *Developmental Cell*, 50(1), pp. 5–6. doi: 10.1016/j.devcel.2019.05.012.

**Gapp et al. Figure 1**

**A**

14 days

Dexamethasone injection    Sperm collection    Small RNA sequencing

ATCTCCCG
CGGTATATC
AGTCTAAAT

**B**                                              **C**

**Gapp et al. Figure 2**

**A**

**B**

Effect of Dex on tRf payload

tRNA−Arg−CCT−2−1

**C**

Effect of time post injection on tRf payload

**D**

tRf ARG-CCT-2

**A**

**B**

Females · Males

**C**

Glucose Tolerance Test

**D**

Insulin Tolerance Test

# Gapp et al. Figure 4

## A

14 days

Dexamethasone injection

Cryopreservation of Sperm

In vitro fertilization

2-cell embryo small RNA sequencing

ATCTCCCG
CGGTATATC
AGTCTAAAT

## B

Treatment

tRNA−Val−AAC−1

tRNA−Val−CAC−2

tRNA−Gly−CCC−5

tRNA−Gly−GCC−3

tRNA−Gly−GCC−2

tRNA−Gly−GCC−4

Z−Score

2.5
2
1.5
1
0.5
0
−0.5

Treatment
CON
DEX

# Gapp et al. Figure 5

# Gapp et al. Figure 6

**A**

14 days

Dexamethasone injection  Sperm collection  Long RNA sequencing

ATCTCCCG
CGGTATATC
AGTCTAAAT

**B**

RNA_type
- regular
○ circular

Tasp1
Dennd1b

x-axis: logFC
y-axis: −log10(FDR)

**C**

Group
- Veh
- Dex

Sdf4
circ-Tasp1
Ebf1
Myh9
Cfap44
Hnrnpa2b1
Syncrip