Feed efficiency of lactating Holstein cows was not as repeatable across diets

as within diet over subsequent lactation stages

Amélie Fischer[1,2*], Philippe Gasnier[2] and Philippe Faverdin[2]

[1]Institut de l'élevage, 75595 PARIS, France

[2] PEGASE, INRAE, Institut Agro, 35590, SAINT-GILLES, France

*Corresponding author: Amélie Fischer – Amelie.fischer@idele.fr

ABSTRACT

Background: Improving feed efficiency has become a common target for dairy farmers to

meet the requirement of producing more milk with fewer resources. To improve feed

efficiency, a prerequisite is to ensure that the cows identified as most or least efficient will

remain as such, independently of diet composition. Therefore, the current research analysed

the ability of lactating dairy cows to maintain their feed efficiency while changing the energy

density of the diet by changing its concentration in starch and fibre. A total of 60 lactating

Holstein cows, including 33 primiparous cows, were first fed a high starch diet-low fibre (diet

S^+F^-), then switched over to a low starch diet-high fibre (diet S^-F^+). To know if diet affect feed

efficiency, we compared the ability of feed efficiency to be maintained within a diet over

subsequent lactation stages, known as repeatability of feed efficiency, with its ability to be

maintained across diets, known as reproducibility of feed efficiency. To do so we used two

indicators: the estimation of the error of repeatability/reproducibility, which is commonly

used in metrology, and the coefficient of correlation of concordance (CCC), which is used in

biology. The effect of diet change could also lead to a change in cows sorting behaviour

which could potentially affect feed efficiency if for example the most efficient cows select

more concentrate than the least efficient. We therefore analysed the relationship between the

differences in individual feed refusals composition and the differences in feed efficiency. To

do so, the composition of each feed refusal was described with its near infra-red (NIR)

spectroscopy and was performed on each individual feed ingredient, diet and refusals and

used as composition variable. The variability of the NIR spectra of the refusals was described

with its principal components thanks to a principal component analysis (PCA). The Pearson

correlation was estimated to check the relationship between feed efficiency and refusals

composition, i.e. sorting behaviour.

Results: The error of reproducibility of feed efficiency across diets was 2.95 MJ/d. This error

was significantly larger than the errors of repeatability estimated within diet, which were 2.01

MJ/d within diet S^-F^+ and 2.40 MJ/d within diet S^+F^-. The CCC was 0.64 between feed

efficiency estimated within diet S^+F^- and feed efficiency estimated within diet S^-F^+. This CCC

was smaller than the one observed for feed efficiency estimated within diet between two

subsequent lactation stages (CCC = 0.72 within diet S^+F^- and 0.85 within diet S^-F^+). Feed

efficiency was poorly correlated to the first two principal components, which explained 90%

of the total variability of the NIR spectra of the individual refusals. This suggests that feed

sorting behaviour did not explain differences in feed efficiency.

Conclusions: Feed efficiency was significantly less reproducible across diets than repeatable

within the same diet over subsequent lactation stages, but cow's ranking for feed efficiency

was not significantly affected by diet change. This loss in repeatability across diets could be

due to a more pronounced feed sorting subsequent to the change in diet composition.

However, the differences in sorting behaviour between cows were not associated to feed

efficiency differences in this trial neither with the S^+F^- diet nor with the S^-F^+ diet. Those

results have to be confirmed on diets having different forage to concentrate ratios to ensure

that the least and most efficient cows will not change.

BACKGROUND

To be more competitive, dairy farmers have to increase the efficiency of resources use while

reducing their environmental footprint. With an expected increase of world population

(United Nations, 2017), feeds for dairy cow may shift towards more non-human edible feeds.

The challenge for dairy farmers will be to improve feed efficiency, while facing more volatile

feed prices (HLPE, 2011) and lower availability of feeds, which are directly edible by human.

Dairy cows' diets will therefore become more variable in the future. Feed efficiency can be

improved either by selecting the most efficient cows, thanks to an index including feed

efficiency, or by improving feed efficiency of the least efficient cows with precision feeding.

If feed efficiency is included as a genomic selection trait, feed efficiency has to be

reproducible independently of diet and environment. The sensitivity of dairy cow's ranking

for feed efficiency to diet composition, also called interaction between genetic and

environment (Hill and Mackay, 2004), needs to be evaluated to know if some cows perform

better and some worse when changing diet composition.

Repeatability is defined as the capacity of a method to give the same results when using the

same sample and repeating measurements in the same experimental conditions (JCGM, 2012),

that is within diet when applied to efficiency. Reproducibility is a repeatability done while

changing one specific characteristic in the experimental conditions (temperature, diet,

operator; (JCGM, 2012)), that is for example by changing diet's composition. When

estimating reproducibility of an indicator under different environmental conditions, it is

therefore essential to compare it with its repeatability under similar conditions to isolate errors

associated with diet change from errors associated with the method. In literature, the

reproducibility of feed efficiency when changing the diet was lower compared to the

repeatability within a given diet, as estimated with the correlations of residual feed intake

(RFI) within and across diets (r = 0.33 vs 0.42 in steers, (Durunna et al., 2011); r = 0.44-0.64

vs 0.53-0.70 in dairy cows, (Potts et al., 2015)). Animals will thus not necessarily rank the

same for feed efficiency when changing diet composition. Repeatability estimation, as defined

by the proportion of genetic and permanent environment variances within and across

lactations in the total variance of RFI, is highly variable across studies and countries with

values between 0.47 and 0.90 (Connor et al., 2013; Tempelman et al., 2015). This variability

across studies is associated with differences in diet composition and in period length between

studies. Indeed, in lactating dairy cows, the correlation of short-term feed efficiency with full

lactation feed efficiency increases with later lactation stages and longer periods (Connor et al.,

2019). The comparison between repeatability and reproducibility needs therefore feed

efficiency to be estimated over a long enough period to get a robust estimation within diet.

Feed efficiency is generally estimated with diets offered ad libitum with a minimum amount

of refusals (in general $5 - 10\%$ of offered). Cows can therefore potentially perform feed

sorting, which may lead to a consumed diet that differs from the distributed diet both in

composition and in nutritive value. This difference between refusals composition and

distributed diet composition has therefore to be considered when analysing the change in feed

efficiency while changing diet's composition because feed sorting behaviour could affect feed

efficiency. For instance, Dykier et al. (2020) observed that RFI was negatively correlated to

the intake of long particles (r = - 0.30, $p < 0.05$) and positively correlated to intake of short

particles (r =0.22, $p < 0.1$) in beef steers fed ad libitum. This difference in sorting behaviour

resulted in a consumed diet which concentration in crude protein increased with RFI (r = 0.25,

$p < 0.05$). Differences in feed composition were characterized by particle size differences in

Dykier et al. (2020). However the method for particle size composition (Lammers et al., 1996;

Kononoff et al., 2003) is burdensome and time consuming. The advent of near-infrared (NIR)

spectroscopy opens new ways to determine diet or feed compositions at high throughput.

Indeed the NIR spectrum is sensitive to physical and chemical characteristics of the sample,

and has therefore been used to determine nutritive value of feed, but also to discriminate

samples according to their composition (Pérez-Marín et al., 2004; De la Roza-Delagado et al.,

2007; Li et al., 2007).

The main objective of the current study was therefore to check the ability of feed efficiency to

be maintained across different diets. To achieve those objectives a trial was set up with

lactating dairy cows that were fed with two diets. These diets differed in energy density, by

lowering the starch concentration and increasing the fibre concentration of the diet. The feed

efficiency was estimated within diet using the method developed in a previous paper (Fischer

et al., 2018). The novelty of the current paper is to estimate feed efficiency reproducibility

across diets by combining two methods: the commonly used CCC in biology and the

comparison of the error of reproducibility across diets with the error of repeatability within

diet, as commonly used in metrology (JCGM, 2012). Indeed, to estimate if FE is maintained

across diets, its reproducibility across diets has to be compared to its repeatability within diet.

If the reproducibility results are as good as the repeatability results within diet, then one can

conclude that FE is as repeatable across diets as it is within diet. Opposedly if the

reproducibility results are worse than repeatability within diet, then one can conclude that the

ability of FE to be maintained across diets is not as good as within diet. As highlighted in the

previous paragraph, a diet change could also lead to a change in cows sorting behaviour

which could potentially affect feed efficiency. A second objective was therefore to check that

the change in feed efficiency associated with diet change was not explained by differences in

sorting behaviour. We therefore checked that feed efficiency was not associated with feed

sorting behaviour by analysing the feed composition of each cow's diet refusals with NIR

spectroscopy.

 MATERIAL & METHODS

Experimental Design

The experimentation was performed at the INRAE-Institut Agro UMR PEGASE research

facility of Mejusseaume (Le Rheu, France). An initial group of 68 Holstein cows were housed

in a free-stall barn with free access to water. These cows were monitored for feed efficiency

from calving (grouped between end of August and middle of October 2017) to end of June

2018, near the end of the lactation. A sequential design was adopted to feed the cows with a

first diet that was high in starch and low in Fibre (diet S^+F^-) and a second diet that was low in

starch and high in fibre (diet S^-F^+). The protein content was adjusted in diet S^-F^+ to have a

ratio between energy and protein that was similar to the ratio in diet S^+F^-. The diet S^+F^- was

fed from calving to March 18^{th} 2018, then all cows switched over to diet S^-F^+ until end of

June 2018. To make the feed efficiency estimated over the two feeding periods comparable,

the same period length was chosen for both periods. Each diet was fed for at least 95 days.

The first 23 days subsequent to the change in diet from S^+F^- to diet S^-F^+ were considered as an

adaptation period to the new diet and were therefore removed from the dataset. Each period

included the last 72 days of data. 62 cows had data over both 72-day periods and were

therefore kept for further analysis. Each period has been split in two sub-periods of 36 days

(Figure 1) to be able to estimate a repeatability within diet and to reach a supposedly

correlation with full lactation RFI of at least 0.8 according to Connor et al. (2019).

Reproducibility was estimated using the last 36-days of the first period and the first 36-days

of the second period to be comparable and compared with repeatability estimation over the

two 36-days segments within diet (Figure 1).

*Segments = time unit used to average the variables NEI, NEMilk, metabolic
BW, and BW loss and gain to fit the models for RNEI.*

Figure 1 Diagram of the design used to characterize the repeatability (r) and reproducibility (R) of residual net energy intake (RNEI). The repeatability compares RNEI within diet; it is shown with the solid line (●—●). The reproducibility compares RNEI across diets; it is shown with the dashed line (●··●). The dairy cows started first with the diet high in starch and low in fibre (S^+F^-) and switched over to the diet low in starch and high in fibre (S^-F^+).

Table 1 Composition of both experimental diets (S^+F^- and S^-F^+) and their chemical analysis.

	Diet S^+F^-	Diet S^-F^+
Feed composition (% DM +/- SD[1])		
Maize silage	65.0 +/- 1.6	59.1 +/- 0.9
Soybean meal	17.8 +/- 0.9	13.3 +/- 0.3
Dehydrated lucerne	8.1 +/- 1.6	14.4 +/- 0.2
Energy concentrate[2] + Minerals and vitamin complement[3]	9.1 +/- 1	1.6 +/- 0.7
Wheat straw	0	11.6 +/- 0.3
Diet analysis (+/- SD[1])		
DM (%)	42.9 +/- 0.7	47.1 +/- 0.3
OM (%)	94.0 +/- 0.15	94.1 +/- 0.04
Crude protein (g/kg)	167 +/- 4.9	145 +/- 1.1
ADF (g/kg)	198 +/- 2.9	249 +/- 1.2
NDF (g/kg)	351 +/- 3.6	420 +/- 1.6
Starch (g/kg)	226 +/- 3	183 +/- 2.2
Net energy for milk (MJ/kg)	6.62 +/- 0.107	5.98 +/- 0.014
Metabolizable protein (g/kg)	105 +/- 1.8	93 +/- 0.4

[1]SD were calculated using the day-to-day change in offered diet composition on an individual cow basis. [2]The part of energy concentrate in the diet, as described here, includes the part of energy concentrates distributed at the Greenfeed® station. [3]Concentrates included 17.8% wheat, 17.8% maize, 17.8% sugar beet pulp, 17.8% barley, 13% wheat bran, 3% beet molasses, 0.9% oil, 0.9% salt, 11% minerals and vitamin complement (including 6% phosphorus, 24% calcium, 5% magnesium, and other minerals and vitamins).

DM = dry matter; OM = organic matter; ADF = acid detergent fibre; NDF = neutral detergent fibre

Phenotypic Measurements

Individual feed intake and feed nutrient analysis. Cows were fed individually twice a

day after each milking (7:00 and 16:00). Daily intake was estimated individually as the

difference between daily offered feed and next morning's refusals. The diet was offered ad

libitum to maintain an average of 10% refusals per cow. Each cow was fed in her own

manger, only accessible by one cow thanks to the identification chip on her neck collar. Each

feed has been sampled once a week for concentrates, and once a day for the forages to

estimate dry matter, which was used to estimate individual feed dry matter intake (DMI). A

bulk sample was taken for each ingredient and silo for nutrient value analysis. Diet S^+F^-

included maize silage, soybean meal, dehydrated alfalfa and a mix of energy concentrates,

minerals and vitamins (Table 1). Diet S^-F^+ was formulated to have a lower starch

concentration by replacing starch with fibre while keeping a similar ratio between

metabolizable protein and net energy for lactation than in diet S^+F^-. To do so, wheat straw was

added to the ingredients included in diet S^+F^- to replace the energy concentrate and part of

soybean meal and maize silage (Table 1). In addition to both diets, cows had access to a gas

emissions monitoring system, the Greenfeed® (see "Individual performance: milk, body

weight and body condition, methane and carbon dioxide emissions" section), which

distributes small drops of energy concentrates to maintain the cow in the gas recording

system. The amount of energy concentrates distributed per cow per day in the Greenfeed®

station was added to the daily intake at the manger. All feed samples were freeze-dried and

ground with a 3-blade knife mill through a 0.8-mm screen. The organic matter content was

determined by ashing for 5h at 550°C in a muffle Furnace (Association Française de

Normalisation, 1997). The concentrations of neutral and acid detergent fibre were measured

according to Van Soest et al. (1991) using a Fibersac extraction unit (Ankon Technology,

Fairport, NY, USA). Fat content was measured by ether extraction and starch analyses were

performed by polarimetry (LABOCEA, Ploufragan, France). Nitrogen concentration for all

samples was determined by the Dumas method (Association Française de Normalisation,

1997) with a LECO Nitrogen Determinator (Leco, St Joseph, MI, USA). Nutritive values of

feeds were calculated from their chemical composition according to equations in INRA

(2010). Diet S^+F^- had 226 g starch/kg DM, 351 g neutral detergent fibre/kg DM for a net

energy concentration of 6.62 MJ/kg DM and 105 g metabolizable protein/kg DM. Diet S^-F^+

was 19% lower in starch concentration and 16.4% higher in neutral detergent fibre

concentration than diet S^+F^- for a net energy concentration of 5.98 MJ/kg DM and 93 g

metabolizable protein/kg DM (Table 1).

Individual performance: milk, body weight and body condition, methane and carbon

dioxide emissions. Milk yield was recorded at each milking with milk meters (DeLaval,

Tumba, Sweden). Milk fat and protein concentrations were analysed by mid infrared

spectrometers (Lillab, Chateaugiron, France) from morning and afternoon milk samples of

two days per week. Milk fat and milk protein concentrations were calculated as weighted

averages relatively to the morning and afternoon milk production of the day of sampling.

Cows were weighed automatically after each milking (W-2000, DeLaval, Tumba, Sweden) to

get an empty udder body weight (**BW**). All cows were scored for body condition once a

month by 3 trained persons according to the scale developed by Bazin (1984), going from 0

for an emaciated cow to 5 for a fat cow with 0.25 unit increments.

Methane emissions were recorded with two Greenfeed® units (C-Lock Inc., Rapid City, SD,

USA). A Greenfeed® unit is designed as a dispenser of concentrates to measure methane and

carbon dioxide emissions each time a cow visits the feeder, therefore both methane and

carbon dioxide were also monitored. A maximum of 720 g (30g/drop) concentrates was

offered daily at the Greenfeed® to attract cows in the Greenfeed. Each Greenfeed® unit

records methane emissions for up to 23 cows. Given the barn configuration and the two

Greenfeed® units, only 42 cows could therefore be monitored for methane emissions during

the study. On average the cows visited the Greenfeed 2.2 /d/cow (+/- 0.9). Energy

concentrates distributed at the Greenfeed® units were included in the estimation of individual

daily feed intake.

Checking for feed sorting behaviour

Individual refusals were sampled by collecting about 0.5 to 1 kg of fresh weight refusals per

cow once a week for 6 weeks during each of the two experimental periods to check if the

change in diet's feed ingredients between diet S^+F^- and diet S^-F^+ modified cow's feed sorting

behaviour. The samples were dried in a forced-air oven at 60°C for 48 h. The refusals samples

of each cow were pooled within period to end up with one sample per cow and per period, and

ground through a 3-blade mill (0.8 mm screen). The same process was applied for each feed

ingredient and each diet to estimate the difference in composition between feeds and refusals.

Instead of determining each sample's feed composition, we used an indirect approach based

on near-infra red (NIR) spectroscopy. Indeed, we have seen in the background section of this

paper that NIR spectroscopy can be used to differentiate samples differing in ingredients

proportion. By definition, if the samples differ on a physical or chemical basis, their spectra

will also be different. In this study, the differences in refusals composition and diets

composition will be analysed through their NIR spectra, without estimating or analysing their

chemical or physical characteristics. A Fourier transform near-infrared (FT-NIR)

spectroscopy with MPA (Bruker Optik Gmbh, Ettligen, Germany) was used to characterize

the spectra (from 3595 to 12490 cm^{-1}, resolution 16 cm^{-1}) of each individual feed, diet and

refusal sample. A principal component analysis (PCA) was fitted on the spectra of the feeds,

both diets and all refusals to summarize the dataset into principal components. A second PCA

was performed on the spectra of the refusals only, with randomRNEI as a supplementary

variable, to summarize the spectra of the refusals into fewer variables. The relationship

between refusals composition and feed efficiency was estimated with the coefficient of

correlation between randomRNEI and each of the principal component of this second PCA.

Both PCA were performed with the FactomineR (Lê et al., 2008) and Factoshiny (Vaissie et

al., 2020) packages of R (R Core Team, 2018).

Outlier Detection

Among the 62 cows, 2 cows had issues with their manger and were therefore removed from

the dataset because their intake data were not reliable enough, to end up with a group of 60

cows, including 33 primiparous cows. For methane and carbon dioxide data, a least rectangles

regression of carbon dioxide emission against methane emission was fitted to detect methane

outliers using the least.rect function of package RVAideMemoire (Hervé, 2018) in R (R Core

Team, 2018). The data outside the range of three standard deviations of the residuals around

the regression line were considered as outliers, and were removed. On average 0.9% (SD =

1.3%) of the initial methane data were removed per cow for being outliers. Methane data were

then averaged per experimental period and per cow.

Variables Calculation to Estimate Feed Efficiency

Estimation of feed efficiency requires DMI data, as well as all energy outputs or energy inputs

to be considered for a lactating dairy cow. Energy outputs gather net energy in milk, energy

required for maintenance, energy gained as adipose tissue and energy required for gestation.

Other energy inputs include adipose tissue mobilization.

Net energy in milk was calculated according to the following equation (Faverdin et al., 2010):

$\text{NEMilk (MJ/d)} = 7.12 \times \text{MilkProd} \times \left(0.44 + 0.0055 \times (\text{MFC} - 40) + 0.0033 \times (\text{MPC} - 31)\right)$

Where MilkProd is the milk production in kg/d, MFC is the milk fat concentration in g/kg and

MPC is the milk protein concentration in g/kg.

Gestation requirement were estimated with the following equation defined by Faverdin et al.

(Faverdin et al., 2010):

$\text{Gestation (MJ / d)} = 7.12 \times 0.00072 \times \text{BWbirth} \times \exp^{0.116 \times \text{GestWeek}}$

Where BWbirth is calf's weight at birth and assumed to be 40 kg, GestWeek is the week of

gestation.

Morning BW data used were smoothed with a moving average of the 15% neighbouring data,

to better reflect change in maintenance and to be less sensitive to daily gutfill change.

Monthly body condition score (**BCS**) data were filled to get daily BCS using a cubic Spline

with the function smooth.spline in R (R Core Team, 2018) using each scoring day as a knot.

Maintenance requirements were estimated with the metabolic BW, using the smoothed BW

data, and calculated as $\text{BW}^{0.75}$. Energy gained and energy mobilized as body reserves were

estimated as the day-to-day change in smoothed BW. If the change was positive, it was

attributed to body reserves gain, and body reserves loss was null. Conversely, if the change

was negative, it was attributed to body reserves mobilization, and body reserves gain was null.

Both BW gain and BW loss were constructed to be positive variables. Both, BW gain and BW

loss, were multiplied by daily BCS to account for body reserves differences within a given

BW change, resulting in the variables **BWlossBCS** and **BWgainBCS**.

Estimation of feed efficiency

Feed efficiency was estimated as the residual feed intake with the method developed in a

previous paper (Fischer et al., 2018). Briefly, instead of being estimated as the residual of the

multiple linear regression that estimates the observed DMI with the main energy outputs and

inputs, RFI was defined with a mixed model as the repeatable animal effect. Applied to our

study, each sub-period of 36 days was subdivided in segments of 12 days to end up with three

repeated measures for each cow within each sub-period (Figure 1). The initial model

explained net energy intake (NEI) with the fixed effects of net energy in milk, metabolic BW,

BWlossBCS, BWgainBCS, gestation requirement, BCS, their interaction with parity and sub-

period, and the fixed effect of parity, sub-period, and 12-day segment nested within sub-

period. This model included the repeated effect of cow across the 12-day segments within

sub-period and the random effect of cow within sub-period, and were grouped within sub-

period. Only significant ($p \leq 0.05$) interactions and variables were kept in the model. As in

our previous paper (Fischer et al., 2018), feed efficiency was defined as the random part of the

intercept of the mixed model 1 below, that was performed using PROC MIXED in SAS

(Version 9.4 of the SAS System for Linux. 2017. SAS Institute Inc., Cary, NC, USA) with a

heterogeneous autoregressive variance covariance matrix for the repeated statement. The

variables were averaged per 12-day segment.

296
$$\begin{aligned}
\text{NEI (MJ / d)} = &\left(\mu + \mu_{cow}\right) + \text{NEMilk} + BW^{0.75} + \text{BWlossBCS} + \text{BWgainBCS} \\
&+ \text{gestation} + \text{parity} + \text{sub-period} + \text{parity} \times BW^{0.75} + \text{sub-period} \times \text{NEMilk} \quad \text{(model 1)} \\
&+ \text{sub-period} \times BW^{0.75} + \text{segment} \mid \text{sub-period} + \varepsilon
\end{aligned}$$

where NEMilk is the net energy in milk in MJ/d, $BW^{0.75}$ is the metabolic BW in $kg^{0.75}$,

gestation is the gestation requirement in MJ/d, parity is the fixed effect of parity, sub-period is

the fixed effect of sub-period, segment|sub-period is the fixed effect of 12-day segment nested

within sub-period, parity×$BW^{0.75}$ is the interaction between parity and $BW^{0.75}$, sub-

period×NEMilk and sub-period×$BW^{0.75}$ are the interactions between sub-period and NEMilk,

and sub-period and $BW^{0.75}$, μ is the fixed intercept and μ_{cow} is the random part of the intercept

and ε is the error. Feed efficiency was defined as μ_{cow} in model 1 and will be called random

residual net energy intake **(RandomRNEI)**.

Repeatability and Reproducibility of Feed Efficiency

Repeatability and reproducibility were estimated with 2 methods: the standard deviation of

repeatability and standard deviation of reproducibility as defined by ICAR (JCGM, 2012),

and Lin's coefficient of correlation of concordance (CCC) (Lin, 1989).

Estimating Error of Repeatability and Reproducibility. Repeatability was estimated

within diet with an analysis of variance. For repeatability the model 3 below of analysis of

variance was fitted once with the data of feed efficiency within diet S^+F^- to get the

repeatability within diet S^+F^-, and once with the data of diet S^-F^+ to get the repeatability

within diet S^-F^+. Reproducibility was estimated with an analysis of variance with the data of

the second sub-period when cows were fed diet S^+F^- and the data of the first sub-period when

cows were fed the diet S^-F^+, to be able to estimate the variance associated with diet change.

Both analysis of variance were performed using the Anova function of car package (Fox and

Weisberg, 2011) in R (R Core Team, 2018) as follows:

$NEI = \mu + Cow + \varepsilon$ (model 3)

Where Cow stands for the fixed effect of cow, μ is the intercept and ε is the error. Standard

deviations of repeatability and of reproducibility were defined as the standard deviation of ε in

model 3. An F-test was performed with the var.test function in R (R Core Team, 2018) to test

if the error of repeatability in diet S^+F^- was similar to the error of reproducibility across diets,

and similarly by comparing error of repeatability in diet S^-F^+ with the error of reproducibility

across diets.

Estimation of Repeatability and Reproducibility Correlations. The Lin's coefficient

of correlation of concordance was estimated between the feed efficiency estimated at two

different sub-periods within diet for repeatability, and between feed efficiency estimated

during the second sub-period in diet S^+F^- and first sub-period in diet S^-F^+ for reproducibility.

The CCC were estimated with the epi.ccc function of epiR package (Stevenson et al., 2020) in

R (R Core Team, 2018).

All statistical analysis were done with the significance level of 0.05 ($p \leq 0.05$).

RESULTS

Period Effect on Cows Performance

334
335
The decrease in dietary net energy for lactation and in metabolizable protein was confounded

with the increase of lactation stage as the experimentation was based on a sequential design.

Therefore when effect of diet is mentioned here, it is confounded with the effect of lactation

stage.

The diet change decreased net energy intake, without significantly changing dry matter intake

(p = 0.26). Indeed, Cows ate on average 23.4 kg DM/d over both periods. They ate 156 MJ/d

when they were fed with diet S^+F^-, which was 13.9 % (19 MJ/d) more net energy than when

they were fed with diet S^-F^+ (p < 0.01; Table 2).

Cows partitioned 92.1 MJ/d in milk on diet S^-F^+, which was 17.3% (19.3 MJ/d) less than with

diet S^+F^- (p < 0.01, Table 2). This difference in net energy exported in milk between diets was

also observed for milk production. Indeed, cows produced 29.9 kg milk/d with diet S^-F^+,

which was 17.9% (6.5 kg/d) less than with diet S^+F^- (p < 0.01, Table 2). Change in dietary net

energy and metabolizable protein concentrations did neither significantly affect milk protein

concentration (p = 0.88) nor milk fat concentration (p = 0.56), with average concentrations of

31.4 g protein/kg milk and 38.8 g fat/kg milk (Table 2). Given the steady milk content and a

decreasing milk production, the decrease in net energy and metabolizable protein in the diet

decreased milk protein production (p < 0.01) and milk fat production (p < 0.01).

Maintenance related variables, known as BW and BCS, were lower when cows were fed diet

S^+F^- (p = 0.03 for BW and p < 0.01 for BCS), which was also at earlier lactation stages, with

averages of 651 kg and 1.96 BCS with diet S^+F^-, and 680 kg and 2.08 BCS with diet S^-F^+

(Table 2).

Table 2 Dry matter intake, performance, methane emissions and efficiency of the 60 Holstein

cows fed S^+F^- diet first and then S^-F^+ diet three months later.

	Least squares means		Residual SD	p-value
	Diet S^+F^-	Diet S^-F^+		
n (primiparous)	60 (33)	60 (33)		
Intake				
DMI (kg/d)	23.7	23.1	2.94	0.26
NEI (MJ/d)	156	137	18.58	<0.01
Performance				
Milk production (kg/d)	36.4	29.9	4.93	<0.01
Milk Fat production (g/d)	1,387	1,154	194.0	<0.01
Milk Protein production (g/d)	1133	933	145.6	<0.01
Milk Protein concentration (g/kg)	31.4	31.4	2.44	0.88
Milk Fat concentration (g/kg)	38.5	39.0	4.27	0.56
NE in Milk (MJ/d)	111.4	92.1	14.17	<0.01
BW (kg/d)	651	679	68.2	0.03
BCS	1.96	2.08	0.23	<0.01
BW Gain (kg/d)	0.40	0.37	0.16	0.28
BW Loss (kg/d)	0.14	0.32	0.12	<0.01
Methane				
Methane production (g/d)	510	523	66.2	0.41
Methane yield (g/kg DMI)	21.4	22.6	2.39	0.03
Methane yield (g/kg Milk)	14.0	17.6	2.58	<0.01

DMI: dry matter intake; NEI: net energy intake estimated using equation in Faverdin et al. [12]; NE

in milk: net energy in milk estimated using equation in Faverdin et al. [12]; BW: body weight; BCS:

body condition score; CH_4: methane emission. Diet S^+F^-: high in starch and low in fibre; diet S^-F^+:

low in starch and high in fibre.

Variables associated with body reserves change, identified as BW loss and BW gain in table

2, were differently affected by diet. Cows mobilized more BW when they were fed with S^-F^+

diet, also known as period 2, with a loss of 0.32 kg/d, than when fed with S^+F^- diet, also

known as period 1, with a loss of 0.14 kg/d (p < 0.01, Table 2). Gain in BW did not

significantly differ between both diets, with an average gain of 0.39 kg/d.

Dietary decrease in starch, replaced with fibre, confounded with the effect of lactation stage,

did not significantly change methane emissions, as cows emitted on average 517 g methane/d

(Table 2). Nevertheless, this dietary change increased methane yield from 21.4 g methane/kg

DMI up to 22.6 g methane/kg DMI (p = 0.03) and from 14.0 g methane/kg milk up to

17.6 g methane/kg milk when switching from diet S^+F^- to diet S^-F^+ (p < 0.01, Table 2).

Effect of Diet change on Feed Efficiency

Feed efficiency was more variable when cows were fed the S^-F^+ diet than when fed the S^+F^-

diet. Feed efficiency, as estimated with randomRNEI, had a standard deviation of 4.49 MJ/d

in sub-period 1 and 4.61 MJ/d in sub-period 2 when cows were fed S^+F^- diet, and 5.18 MJ/d in

sub-period 1 and 5.21 MJ/d in sub-period 2 when cows were fed the S^-F^+ diet. The change in

randomRNEI induced by diet change (randomRNEI diet S^+F^- - randomRNEI diet S^-F^+) was

negatively correlated with the change in methane yield, as per kg DMI, (CH4/DMI diet S^+F^-

- CH4/DMI diet S^-F^+) with a Pearson correlation of - 0.31 (p = 0.05), but was neither

significantly correlated with the change in methane production per day (p = 0.12) nor with the

change in methane yield, as per kg milk (p = 0.98). This means that a cow that had a lower

randomRNEI (higher feed efficiency) in diet S^-F^+, also had a higher methane yield per kg

DMI in diet S^-F^+ than when fed with the S^+F^- diet, and conversely.

Feed Efficiency Reproducibility across Diets

Cows' feed efficiency was not as reproducible across diets than repeatable within diet (Table

3 and Figure 2). Errors of reproducibility, when comparing efficiency across diets, were larger

than the errors of repeatability within diet, regardless of diet (Table 3). Indeed, the

reproducibility error of randomRNEI that was 2.95 MJ/d was significantly larger (p < 0.05,

Table 3) than the repeatability errors for diet S^+F^- and for diet S^-F^+ that were 2.40 and

2.01 MJ/d, respectively (Table 3). This

lower reproducibility across diets

compared with repeatability within diet

tended to be observed with the CCC when

comparing reproducibility across diets with

repeatability in diet S^-F^+, but not when

compared with repeatability within diet

S^+F^- (Table 3). Indeed, the CCC between

randomRNEI estimated in diet S^+F^- and

randomRNEI estimated in diet S^-F^+ was

0.64, which was smaller compared with the

correlations of 0.72 (p = 0.55) within diet

S^+F^- and 0.85 (p = 0.055) within diet S^-F^+

(Table 3).

Figure 2 Relationship between feed

efficiency estimated within the same diet

for repeatability estimation within diet S^+F^-

(A), S^-F^+ (B) or across diets for

reproducibility estimation (C). The dashed

black line stands for the first bisector, and

the solid black line stands for the least

rectangles regression.

Table 3 Error of repeatability and reproducibility, and Lin's concordance correlation coefficient within diet for repeatability and across diets for reproducibility (Repro.) for feed efficiency.

	Repeatability[1]		Repro.[2]	P-value		
	Diet S^+F^-	Diet S^-F^+		S^-F^+ vs S^+F^-	S^-F^+ vs Repro.	S^+F^- vs Repro.
SD (MJ/d)	2.401	2.014	2.953	0.06	<0.01	0.02
CCC	0.72	0.85	0.64	0.19	0.05	0.55

All cows were first fed with the S^+F^- diet (S^+F^-: high in starch and low in fibre) and then with the S^-F^+

diet (S^-F^+: low in starch and high in fibre). CCC: Lin's coefficient of correlation of concordance; SD:

standard deviation of the residuals of the analysis of variance defined in model 3. [1]Repeatability was

estimated within diet using two repetitions of 36 days within diet. Repeatability was defined as the

standard deviation of the residuals of model 3. The lower the error of repeatability, the more

repeatable it is. [2]Reproducibility was estimated using two repetitions of 36 days (1 repetition/diet): the

last 36 days of diet S^+F^- and the first 36 days of diet S^-F^+. Reproducibility was defined as the standard

deviation of the residuals of model 3. The lower the error of reproducibility, the more reproducible it

is.

Effect on feed selection and feed efficiency

The change in diet composition was associated with a change in cows' feed sorting behaviour. Indeed, the first plan of the PCA (Figure 3) showed that the individual refusals were clustered around the origin of this first plan, close to the two diets samples, and slightly shifted to the more fibrous feed ingredients of the diets (leaves part of maize silage, and straw). This suggests that the cows may have left more fibrous ingredients than concentrates or grains in the refusals, and that individual refusals seem to be different to any particular feed ingredient in the diet. Despite the high number of spectra length waves used to describe the spectrum of each sample, the two first principal components of the PCA explained 94% of the total variability of the spectra of the refusals and the feed samples (Figure 3). When focusing only on the refusals samples (second PCA), the first plan explained 90% of the total variability of

Figure 3 Principal Component Analysis of NIR spectra of the feed ingredients (MS=Maize Silage, MS_leaves = leaves and cane of Maize Silage, MS_grain=Grain of maize silage, Conc_Ener=concentrate energy, SB Meal = soybean meal, Dehyd_Lucerne = dehydrated Lucerne, Mineral), diets (Diet 1 = diet S^+F^- and Diet 2 = diet S^-F^+) and refusals of diet S^+F^- (1) and diet S^-F^+ (2). The 2 first components explain 94% of the total variance.

Diet S^+F^- : diet high in starch and low in fibre; Diet S^-F^+: diet low in starch and high in fibre.

Figure 4 Principal Component Analysis of NIR spectra of the refusals of diet S^+F^- (1) and diet S^-F^+ (2). The 2 first components explain 90% of the total variance. The size of the points is proportional to the absolute value of the residual net energy intake and the color is red if randomRNEI is positive (inefficient cows) and blue if randomRNEI is negative (efficient cows).

Diet S^+F^- : diet high in starch and low in fibre; Diet S^-F^+: diet low in starch and high in fibre.

the refusals' spectra (Figure 4). This focus on refusals' spectra only showed that the refusals

were clustered in 2 groups identifiable as the 2 diets (Figure 4). This suggests that the refusals

reflect the composition difference between both diets. Within each diet, the refusals spectra

differed across cows, but this difference was not associated with feed efficiency differences.

Indeed, the randomRNEI was evenly distributed within diet with no clear clustering within

diet that was associated with randomRNEI. Moreover, the principal components were poorly

correlated with randomRNEI with a correlation of – 0.05 between randomRNEI and principal

component 1, and of 0.009 between randomRNEI and principal component 2. If some

variability exists in the composition of refusals due to feed sorting, this selection was not

associated to feed efficiency differences in this trial neither when cows were fed with the S^+F^-

diet nor when fed with the S^-F^+ diet.

DISCUSSION

Feed Efficiency was less reproducible across diets than within diet

Feed efficiency was less reproducible across diets than within diet when using the

method defined by ICAR (JCGM, 2012) based on the estimation of errors. However, as

shown with the CCC and the cow's ranking (see section "Availability of data and material"

for this supplementary material), the change in cow's ranking was similar when comparing

between the two diets than when comparing within diet over subsequent lactation stages. This

shows that the change in diet affected the absolute value of feed efficiency but not cow's

ranking.

This difference between the results observed with the error method and those observed

with the CCC shows that it is therefore important to use several indicators when the objective

is to characterize reproducibility and repeatability of a method, here of feed efficiency. In our

study, we showed that the error of reproducibility across diets was significantly higher than

the error of repeatability within diet, while the CCC were similar when comparing across diet

and within diet S^+F^-, but different when comparing across diets and within diet S^-F^+. With one

indicator we would have concluded that diet has a significant effect on feed efficiency

repeatability, while with the second we would have concluded that the effect of diet does not

seem significant. The two indicators are complimentary: the CCC will give the strength of the

relationship between the two diets feed efficiency without any information about the

dispersion of the residuals of this relationship, while the errors give the dispersion of the

relationship. Most of the studies characterizing the reproducibility of feed efficiency in cattle

used coefficient of correlations and percentages of cows which changed their efficiency group

(Durunna et al., 2011; Potts et al., 2015; Asher et al., 2018). Conclusions based on animals'

re-ranking are highly subjected to the definition of efficiency groups, which is variable across

studies. The conclusions about repeatability may even differ with different definitions of

efficiency group. We therefore preferred not to use re-ranking to characterize reproducibility

of feed efficiency.

Upon the objective of cow's selection based on feed efficiency, one will prefer one

indicator or the other. For example for selection purposes, one may especially be interested in

the rate of cows able to maintain their efficiency class, and therefore use the CCC indicator.

With this in mind, our results and the literature show (Durunna et al., 2011; Potts et al., 2015;

Asher et al., 2018) that the ranking is quite similar within diet over time and across diets, and

therefore that the risk to misidentify the most or least efficient cows is minimal. If the

objective is to improve efficiency on an existing herd, one may prefer to use the errors

indicator. With this in mind, our results show that a diet change affects the efficiency of the

herd because the errors are significantly higher when predicting feed efficiency after a diet

change than within the same diet. Indeed if efficiency would be reproducible when changing

diet's composition, all cows should behave the same way and their efficiency would be easily

predictable with their previous efficiency. This is not observed, because when randomRNEI

in first subperiod within diet S^-F^+ was predicted with the randomRNEI in second subperiod

within diet S^+F^-, the regression was able to explain 41 % of the total variability of

randomRNEI in first subperiod within diet S^-F^+. This low coefficient of determination is

mostly explained by the diversity of adaptation of each cow's DMI to the diet change (Figure

2). Indeed, for similar randomRNEI and similar DMI there were cows which decreased their

DMI and decreased their randomRNEI, while others increased both their DMI and

randomRNEI, and others maintained both their DMI and randomRNEI. This shows that cows

having similar feed efficiency and intake on a specific diet, will not necessarily end up with

similar efficiency and intake on a different diet.

To conclude about reproducibility, one should also estimate repeatability and compare

it to reproducibility. If the reproducibility is similar to repeatability, the method or estimation

is highly reproducible. If the reproducibility is significantly worse than repeatability, the

method is less reproducible than repeatable, and therefore the method or estimation is

sensitive to the environment. The estimation of repeatability is therefore essential when

characterizing the reproducibility of a method or estimation. The lack of confidence interval

or statistical test to compare the reproducibility and repeatability indicators in most of the

studies (Durunna et al., 2011; Potts et al., 2015; Asher et al., 2018) limits the possibility to

conclude about the reproducibility of feed efficiency.

Similar reproducibility results than in literature

The CCC comparing efficiencies before and after diet change was significantly lower

when compared with the CCC of repeatability within diet S^-F^+ for feed efficiency (p = 0.05),

but was not different when compared with diet S^+F^- CCC (p > 0.1). The lack of significance

observed when comparing the reproducibility CCC when diet changed with the repeatability

CCC estimated within diet S^+F^- can be explained by the lower repeatability of feed efficiency

observed when cows were fed with diet S^+F^-. The lower repeatability of randomRNEI within

diet S^+F^- can be explained by the lower repeatability observed for NEI within diet S^+F^- as its

error of repeatability was 4.60 MJ/d within diet S^-F^+ and 5.82 MJ/d within diet S^+F^-. The

observed decrease in correlation for feed efficiency under reproducibility conditions was also

found in literature with Pearson's correlations of 0.54-0.70 in heifers and 0.42 in steers within

diet, that were larger than those observed after diet changed with correlations of 0.40 for

heifers and 0.33 for steers (Durunna et al., 2011; Cassady et al., 2016). In dairy cows,

correlations averaged 0.65 within diet and 0.56 across diets (Potts et al., 2015). The

correlations they observed were lower and closer together compared to the correlations

observed in the current study. The higher correlations observed in our study can be explained

by the longer period used, that is 36 days, compared to Potts et al. (2015) who used 7 days per

sub-period to estimate repeatability correlations within diet. In fact, the longer and the closer

to middle of lactation the period was, the better the RFI approximates a RFI estimated over

the full lactation (Connor et al., 2019). It would be worth to look at the period length required

to reach the maximum repeatability within diet and within lactation for RFI, and then apply

this length within diet and estimate reproducibility of feed efficiency when changing diet.

Diet and Period Effects on Performance

Dietary starch and fibre concentration modified intake and performance in lactating dairy

cows. Diets high in starch and lower in fibre compared to diets low in starch and higher in

fibre do not necessarily increase intake (Boerman et al., 2015; Potts et al., 2015; Karlsson et

al., 2018). Similarly, in the current study, intake, expressed as DM, was not significantly

higher when cows were fed with S^+F^- diet than when they were fed with S^-F^+ diet (p = 0.26).

However, when expressed as net energy, cows ate more when fed with the S^+F^- diet than

when fed with the S^-F^+ diet (p < 0.01), such as observed in Karlsson et al. (2018). Decrease in

dietary starch and fibre concentrations reduced milk production (p < 0.01), as observed in

literature (Boerman et al., 2015; Potts et al., 2015; Karlsson et al., 2018), but in the current

study this effect was confounded with the effect of lactation stage. This decrease in milk

production subsequent to a decrease of dietary starch and increase of dietary fibre is usually

observed with an increased milk fat concentration (Boerman et al., 2015; Potts et al., 2015;

Karlsson et al., 2018). This was not observed in the current study because neither milk fat nor

milk protein concentrations were different between diet S^+F^- and diet S^-F^+ ($p = 0.56$ for milk

fat and 0.88 for milk protein). One could argue that our change in NDF between both diets, of

16%, was too low compared to the 18-38% across Boerman et al. (2015), Potts et al. (2015)

and Karlsson et al. (2018) to see significant changes in milk fat concentrations. However, in

Boerman et al. (2015) only the lowest increase in NDF (18%) had a significant change in milk

fat concentration; the other diets having a higher change in NDF did not significantly increase

the milk fat concentration. Therefore the increase in NDF may not systematically increase

milk fat concentration. This steadiness of milk solids concentrations, combined with the

decrease in milk production, resulted in lower net energy exported in milk ($p < 0.01$) and

lower milk fat and protein productions ($p < 0.01$) on diet S^-F^+. Having a diet higher in net

energy concentration should have led to more body reserves replenishment and less body

reserves mobilization, as observed in Boerman et al. (2015) and Potts et al. (2015). In the

current study, a higher dietary starch concentration was associated with lower BW loss

($p < 0.01$), and similar BW gain ($p = 0.28$). Those conclusions differences for change in milk

solids and BW gain between our study and literature results can be explained by the

difference in experimental design. Indeed, the current study was based on a sequential design

while the other studies were based on a crossover design (Boerman et al., 2015; Potts et al.,

2015; Karlsson et al., 2018). The main limit of a sequential design, as chosen in the current

study, is the confounding between the effect of stage of lactation and the effect of treatment,

which does not exist in a crossover design. However, crossover designs have limits too: it is

impossible to characterize the effect of interaction between treatment and time or to

characterize a possible remnant effect of the first treatment on the following treatments. It is

therefore difficult to conclude if diet change or just lactation stage explained the decrease in

NEI and in milk production in the current study. However, given that cows should replenish

their body reserves more and mobilize them less with increasing lactation stages, we can

hypothesize that the lack of change in BW gain and the increased BW loss is mostly due to

decreased dietary starch concentration. The observed increase in BW and BCS in the current

study may be explained by more advanced lactation stages, and less by diet change.

Reducing starch and increasing fibre concentrations of the diet increased methane yield, as

per kg DMI (p = 0.03) and as per kg milk produced (p < 0.01). This increase of methane per

kg milk produced is partly explained by a lower milk production with similar feed intake. The

increase of methane yield per kg of DM is more associated with the diet composition. By

decreasing starch concentration in the diet and increasing dietary fibre concentration with

increased wheat straw and lucerne, the production of volatile fatty acids in the rumen may

have shifted in favour of acetate or butyrate profile instead of propionate profile. This shift is

usually associated with a friendlier methanogenic environment (Knapp et al., 2014). The

expected higher methane yield with lower dietary starch concentration, as per kg DMI or per

kg milk, as observed in the current study, has also been observed in literature (Bougouin et

al., 2018) but was not consensual. For some research, methane yield per kg DMI did not differ

when changing dietary starch concentration (Hatew et al., 2015; Pirondini et al., 2015). A

decrease in dietary starch concentration has usually been associated with an increase in

methane emitted per day (Pirondini et al., 2015; Bougouin et al., 2018), which was not

significant in the current study (p = 0.41). These diverging results about effect of dietary

starch concentration on methane production may reflect differences in the effect of dietary

starch concentration on DMI, milk yield and diet digestibility. Higher methane emissions per

day were observed when the change in dietary starch concentration was associated with

significant differences in DMI (Bougouin et al., 2018) or significant differences in diet

digestibility (Pirondini et al., 2015; Bougouin et al., 2018). The significant change in methane

yield per kg DMI was observed when the change in dietary starch concentration was

associated with a change in DMI and a change in diet digestibility (Bougouin et al., 2018).

The conclusions on dietary starch concentration on methane emissions seem to depend on the

effect of diet composition on both, DMI and diet digestibility.

Limits of the study

Given that digestibility partly explains feed efficiency differences in lactating dairy

cows (Oliveira et al., 2016; Potts et al., 2017), it would be interesting to evaluate the cows'

ability to maintain their efficiency for diets with decreased DM digestibility or for diets with

greater physically effective fibre affecting rumen fill. For example, a future study could

compare a classic highly digestible fibre, as S^+F^- diet in current study, with a diet including

fresh grass or hay.

In the current study, the effect of diet was confounded with the effect of lactation stage

because all cows went through the same sequence at the same time. The change in

performance, as well as in intake, could therefore be attributed to the diet, but also to lactation

stage. This experimental design can also lead to remnant effects of the previous treatment to

subsequent periods: results could have been different if diet sequence would have been

reversed between the two periods. Repeatability errors in diet S^+F^- were similar to

repeatability errors in diet S^-F^+ ($p = 0.06$). Feed efficiency was therefore as repeatable in diet

S^+F^- as in diet S^-F^+ (Table 3), which suggests that there was no remnant effect of diet S^+F^- on

feed efficiency under diet S^-F^+ and that lactation stage did not affect feed efficiency

variability in our study. Another way to tackle confusion between lactation stage and

treatment would have been to adopt a crossover design. However, this design can possibly

lead to an interaction between time and treatment, which is not quantifiable, or leads to a

remnant effect of the first treatment. Moreover, a similar study was conducted by Fischer et

al. where a cross-over design (paper under review) was used. The results and conclusions

were similar to the current study. This supports the validity of the current paper.

A last limit to the study is the length of the adaptation period between diet S^+F^- and S^-

F^+. When comparing subperiod 2 of diet S^+F^- with subperiod 1 of diet S^-F^+, we compared two

periods which had different length of adaptation period. Indeed the first had at least subperiod

1 of diet S^+F^- (36 days) whereas the second had 23 days. As we commonly consider that 2 to

3 weeks are enough to ensure that the cows are fully adapted to a new diet, we considered that

the adaptation to the diets was achieved for the data used in the current study, and therefore

we considered that the difference in length of the adaptation period did not influence the

results.

CONCLUSIONS

The study estimated the reproducibility of feed efficiency in dairy cows when changing the

diet concentration in starch and fibre, by comparing the reproducibility across diets with its

repeatability within diet. The results showed that feed efficiency was significantly less

reproducible when changing diet's starch and fibre concentration than repeatable within diet

over subsequent lactation stage. However the change in feed efficiency ranking of dairy cows

was not significantly different when comparing the ranking when cows were fed with the two

different diets than when comparing the ranking when cows were fed with the same diet over

subsequent lactation stages. The diet change in starch and fibre concentration did not affect

the ranking of dairy cows more than does the lactation stage. The change in diet composition

affected the feed sorting behaviour of the dairy cows. However this feed sorting was neither

related to feed efficiency differences nor to the change in feed efficiency subsequent to diet

change.

LIST OF ABBREVIATIONS

659 BCS: body condition score; BW: body weight; CCC: Lin's coefficient of correlation of

660 concordance; CH4: methane; DM: dry matter; DMI: dry matter intake; S^+F^-: diet high in

661 starch and low in fibre; S^-F^+: diet low in starch and high in fibre; NEI: net energy intake;

662 randomRNEI: random residual net energy intake; SD: standard deviation.

DECLARATIONS

Ethics approval

665 The protocol has been approved by the ethical committee and the French Ministry of Higher

666 Education, Research and Innovation (Authorization of the French Ministry of Higher

667 Education, Research and Innovation reference APAFIS 3122-2015112718172611).

Consent for publication

669 Not applicable.

Availability of data and material

671 The datasets generated and/or analysed during the current study are available in the

672 data.INRAE repository: https://doi.org/10.15454/FHRTWJ. In this repository you will also

673 find the supplemental table with the cows' ranking per subperiod (see table "RFI").

Competing interests

675 The authors declare that they have no competing interests.

Funding

677 This study was funded by the ANR project Deffilait (ANR-15-CE20-0014) and by APIS-

678 GENE. They both funded data collection and salary of A. Fischer.

Author's contributions

PF designed the experiment and monitored it; AF gathered and cleaned the data; PG

contributed to define the methodology used to check the hypothesis of feed sorting, to analyse

and interpret NIRS analysis of feeds and refusals; AF and PF analysed the data and

interpreted the results; AF was the major contributor in writing the manuscript; AF and PF

read and approved the final manuscript.

Acknowledgements

The authors warmly thank technical staff and managers at the INRA Méjusseaume research

facility who helped managing this experimentation, did a great job in doing precision

monitoring of the data, extracting and preparing the data and for being very helpful for

biologically clarifying abnormal data.

<div align="center">**REFERENCES**</div>

Asher, A., A. Shabtay, M. Cohen-Zinder, Y. Aharoni, J. Miron, R. Agmon, I. Halachmi, A.

Orlov, A. Haim, L.O. Tedeschi, G.E. Carstens, K.A. Johnson, and A. Brosh. 2018.

Consistency of feed efficiency ranking and mechanisms associated with inter-animal

variation among growing calves. J. Anim. Sci. 96:990–1009.

doi:https://doi.org/10.1093/jas/skx045.

Association Française de Normalisation. 1997. Aliments des animaux - Dosage de l'azote -

Méthode par combustion (DUMAS) - NF V18-120; Dosage des cendres brutes - NF

V18-101. Association Française de Normalisation. Saint-Denis La Plaine, France.

Bazin, S. 1984. Grille de Notation de l'état d'engraissement Des Vaches Pie Noires. RNED

bovin. RNED Bovin. Institut technique de l'élevage bovin, Paris, France.

Boerman, J.P., S.B. Potts, M.J. VandeHaar, and A.L. Lock. 2015. Effects of partly replacing

dietary starch with fiber and fat on milk production and energy partitioning. J. Dairy

Sci. 98:7264–7276. doi:https://doi.org/10.3168/jds.2015-9467.

Bougouin, A., A. Ferlay, M. Doreau, and C. Martin. 2018. Effects of carbohydrate type or

bicarbonate addition to grass silage-based diets on enteric methane emissions and milk

fatty acid composition in dairy cows. J. Dairy Sci. 101:6085–6097.

doi:https://doi.org/10.3168/jds.2017-14041.

Cassady, C.J., T.L. Felix, J.E. Beever, D.W. Shike, and N.P. for G.I. of F.E. in B. Cattle.

2016. Effects of timing and duration of test period and diet type on intake and feed

efficiency of Charolais-sired cattle1. J. Anim. Sci. 94:4748–4758.

doi:https://doi.org/10.2527/jas.2016-0633.

Connor, E.E., J.L. Hutchison, H.D. Norman, K.M. Olson, C.P. Van Tassell, J.M. Leith, and

R.L. Baldwin. 2013. Use of residual feed intake in Holsteins during early lactation

shows potential to improve feed efficiency through genetic selection. J. Anim. Sci.

91:3978–3988. doi:https://doi.org/10.2527/jas.2012-5977.

Connor, E.E., J.L. Hutchison, C.P. Van Tassell, and J.B. Cole. 2019. Defining the optimal

period length and stage of growth or lactation to estimate residual feed intake in dairy

cows. J. Dairy Sci. 102:6131–6143. doi:https://doi.org/10.3168/jds.2018-15407.

De la Roza-Delgado , B., A., Soldado, A., Martínez-Fernández, F., Vicente, A., Garrido-Varo,

D., Pérez-Marín, M. J., De la Haba, and J. E. , Guerrero-Ginel. 2007. Application of

near-infrared microscopy (NIRM) for the detection of meat and bone meals in animal

feeds: A tool for food and feed safety. Food Chem. 105: 1164-1170.

doi:https://dx.doi.org/10.1016/j.foodchem.2007.02.041.

Durunna, O.N., F.D.N. Mujibi, L. Goonewardene, E.K. Okine, J.A. Basarab, Z. Wang, and

S.S. Moore. 2011. Feed efficiency differences and reranking in beef steers fed grower

and finisher diets. J. Anim. Sci. 89:158–167. doi:https://doi.org/10.2527/jas.2009-

2514.

Dykier, K.C., J.W. Oltjen, P.H. Robinson, and R.D. Sainz. 2020. Effects of finishing diet

sorting and digestibility on performance and feed efficiency in beef steers. animal

14:59–65. doi:https://doi.org/10.1017/S1751731119001988.

Faverdin, P., R. Delagarde, L. Delaby, and F. Meschy. 2010. Alimentation des vaches

laitières. INRA, ed. Quae, Versailles, France.

Fischer, A., N.C. Friggens, D.P. Berry, and P. Faverdin. 2018. Isolating the cow-specific part

of residual energy intake in lactating dairy cows using random regressions. Animal

12:1396–1404. doi:https://doi.org/10.1017/s1751731117003214.

Fox, J., and S. Weisberg. 2011. An {R} Companion to Applied Regression. Second. Sage,

Thousand Oaks, USA.

Hatew, B., S.C. Podesta, H. Van Laar, W.F. Pellikaan, J.L. Ellis, J. Dijkstra, and A. Bannink.

2015. Effects of dietary starch content and rate of fermentation on methane production

in lactating dairy cows. J. Dairy Sci. 98:486–499.

doi:https://doi.org/10.3168/jds.2014-8427.

Hervé, M. 2018. RVAideMemoire: Testing and PLotting Procedures for Biostatistics.

Hill, W.G., and T.F.C. Mackay. 2004. D. S. Falconer and Introduction to quantitative

genetics. Genetics 167:1529–1536.

HLPE. 2011. Price volatility and food security. A report by the High Level Panel of Experts

on Food Security and Nutrition of the Committee on World Food Security. Rome.

Institut National de la Recherche Agronomique (INRA). 2010. Alimentation Des Bovins,

Ovins et Caprins. Besoins Des Animaux - Valeurs Des Aliments: Tables Inra 2007

Mise à Jour 2010. Alimentation des bovins, ovins et caprins. Besoins des animaux -

Valeurs des aliments. Quae, Versailles, France.

JCGM. 2012. International vocabulary of metrology - basic and general concepts and

associated terms (VIM), 2008 version with minor corrections. Int. Vocab. Metrol.

Karlsson, J., R. Spörndly, M. Lindberg, and K. Holtenius. 2018. Replacing human-edible feed

ingredients with by-products increases net food production efficiency in dairy cows. J.

Dairy Sci. 101:7146–7155. doi:https://doi.org/10.3168/jds.2017-14209.

Knapp, J.R., G.L. Laur, P.A. Vadas, W.P. Weiss, and J.M. Tricarico. 2014. Invited review:

Enteric methane in dairy cattle production: Quantifying the opportunities and impact

of reducing emissions. J. Dairy Sci. 97:3231–3261.

doi:https://doi.org/10.3168/jds.2013-7234.

Kononoff, P.J., A.J. Heinrichs, and D.R. Buckmaster. 2003. Modification of the Penn State

forage and total mixed ration particle separator and the effects of moisture content on

its measurements. J. Dairy Sci. 86:1858–1863. doi:https://doi.org/10.3168/jds.S0022-

0302(03)73773-4.

Lammers, B.P., D.R. Buckmaster, and A.J. Heinrichs. 1996. A simple method for the analysis

of particle sizes of forage and total mixed rations. J. Dairy Sci. 79:922–928.

doi:https://dx.doi.org/10.3168/jds.S0022-0302(96)76442-1.

Lê, S., J. Josse, and F. Husson. 2008. FactoMineR: A Package for Multivariate

AnalysisAideMemoire: Testing and PLotting Procedures for Biostatistics. J. Stat.

Softw. 25:1–18. doi:https://doi.org/10.18637/jss.v025.i01.

Li, H., X., Lv, J., Wang, J., Li, H., Yang, and Y., Qin. 2007. Quantitative determination of

soybean meal content in compound feeds: comparison of near-infrared spectroscopy

and real-time PCR. Anal. Bioanal. Chem. 389:2313-2322.

doi:https://dx.doi.org/10.1007/s00216-007-1624-1.

Lin, L.I.-K. 1989. A Concordance Correlation Coefficient to Evaluate Reproducibility.

Biometrics 45:255–268. doi:https://doi.org/10.2307/2532051.

Oliveira, L.F., A.C. Ruggieri, R.H. Branco, O.L. Cota, R.C. Canesin, H.J.U. Costa, and

M.E.Z. Mercadante. 2016. Feed efficiency and enteric methane production of Nellore

cattle in the feedlot and on pasture. Anim. Prod. Sci. 58:886–893.

doi:https://dx.doi.org/10.1071/AN16303.

Pérez-Marín, D., A., Garrido-Varo, J. E., Guerrero-Ginel, and A., Gómez-Cabrera. 2004.

Near-infrared reflectance spectroscopy (NIRS) for the mandatory labelling of

compound feedingstuffs: chemical composition and open-declaration. Anim. Feed.

Sci. Tech. 116:333-349. doi:https://doi.org/10.1016/j.anifeedsci.2004.05.002.

Pirondini, M., S. Colombini, M. Mele, L. Malagutti, L. Rapetti, G. Galassi, and G.M.

Crovetto. 2015. Effect of dietary starch concentration and fish oil supplementation on

milk yield and composition, diet digestibility, and methane emissions in lactating dairy

cows. J. Dairy Sci. 98:357–372. doi:https://doi.org/10.3168/jds.2014-8092.

Potts, S.B., J.P. Boerman, A.L. Lock, M.S. Allen, and M.J. VandeHaar. 2015. Residual feed

intake is repeatable for lactating Holstein dairy cows fed high and low starch diets. J.

Dairy Sci. 98:4735–4747. doi:https://doi.org/10.3168/jds.2014-9019.

Potts, S.B., J.P. Boerman, A.L. Lock, M.S. Allen, and M.J. VandeHaar. 2017. Relationship

between residual feed intake and digestibility for lactating Holstein cows fed high and

low starch diets. J. Dairy Sci. 100:265–278. doi:https://doi.org/10.3168/jds.2016-

11079.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing. Vienna, Austria.

Stevenson, M., T. Nunes, C. Heuer, J. Marshall, J. Sanchez, R. Thornton, J. Reiczigel, J.

Robison-Cox, P. Solymos, K. Yoshida, G. Jones, S. Pirikahu, S. Firestone, R. Kyle,

M. Jay, and C. Reynard. 2020. EpiR: Tools for the Analysis of Epidemiological Data.

R package version 1.0-13.

Tempelman, R.J., D.M. Spurlock, M. Coffey, R.F. Veerkamp, L.E. Armentano, K.A. Weigel,

Y. de Haas, C.R. Staples, E.E. Connor, Y. Lu, and M.J. VandeHaar. 2015.

Heterogeneity in genetic and nongenetic variation and energy sink relationships for

residual feed intake across research stations and countries. J. Dairy Sci. 98:2013–2026.

doi:https://doi.org/10.3168/jds.2014.8510.

United Nations Department of Economic and Social Affairs Population Division (United

Nations). 2017. World Population Prospects: The 2017 Revision, Volume II:

Demographics Profiles. Accessed September 17, 2018.

https://population.un.org/wpp/Publications/Files/WPP2017_Volume-II-Demographic-

Profiles.pdf.

Vaissie, P., A. Monge, and F. Husson. 2020. Factoshiny: Perform Factorial Analysis from

"FactoMineR" with a Shiny Application.

Van Soest, P.J., J.B. Robertson, and B.A. Lewis. 1991. Methods for Dietary Fiber, Neutral

Detergent Fiber, and Nonstarch Polysaccharides in Relation to Animal Nutrition. J.

Dairy Sci. 74:3583–3597. doi:https://doi.org/10.3168/jds.S0022-0302(91)78551-2.