1 **TITLE:** **Serum and sputum MMP-9/TIMP-1 in winter sports athletes and**

2 **swimmers: relationships with airway function**

3
Subtitle: Serum and sputum MMP-9/TIMP-1 in athletes

Authors: Valérie Bougault[1,2], PhD, Julie Turmel[2], PhD, Louis-Philippe Boulet[2], MD,

FCCP, FRCPC

[1]Université Côte d'Azur, LAMHESS, Nice, France

[2]Centre de Recherche de l'Institut Universitaire de Cardiologie et de Pneumologie de Québec,

Québec, Canada

13
14 **Sources of support**: None. The authors have no conflict of interest directly relevant to the
15 contents of this study.

16
17
18

19 **Abstract**
20

Introduction: Competitive cross-country skiers and swimmers present characteristics of

airway inflammation and remodeling of the extracellular matrix similar to what is observed in

mild asthma. We aimed to compare serum and sputum MMP-9/TIMP-1, reflecting the

balance between airway fibrogenesis and inflammation process, in swimmers and winter sport

athletes, during the annual period of training resumption, and to observe its seasonal

variations in winter sports athletes.

Methods: Competitive winter sports athletes (n=41), swimmers (n=25) and healthy

nonathletes (n=8) had blood sampling, lung function measurement, skin prick tests, eucapnic

voluntary hyperpnea challenge, methacholine inhalation test (MIT), and induced sputum

analysis. Twelve winter sport athletes performed the test during both summer and winter.

Serum and sputum MMP-9 and TIMP-1 were measured by ELISA.

Results: No significant difference in baseline serum or sputum MMP-9/TIMP-1 ratio was

observed between the three groups, and no relationship was observed with airway function or

responsiveness. Serum MMP-9/TIMP-1 ratio was however significantly higher during the

summer in winter sport athletes compared with winter season (median [Interquartile range]:

1.27 [0.97-1.62] during the winter and 3.65 [2.47-4.03] during the summer, p=0.005). Sputum

MMP-9 correlated with Methacholine PC_{20} and serum CC16/SP-D ratio. No significant

correlation was observed between serum values and lung function or responsiveness.

Conclusion: Our results suggest that although MMP-9/TIMP-1 ratio in sputum or serum was

similar in our three groups of subjects, it may fluctuate according to the training season while

sputum MMP-9 correlated with Methacholine PC_{20} and serum CC16/SP-D ratio, suggesting a

link between airway responsiveness, epithelial damage and MMP-9 in the airways.

Key-words: Airways, Athletes, Biomarkers, Seasons, Fibrogenesis, Inflammation

Introduction

Competitive swimmers and winter sports athletes are particularly at risk to develop respiratory disease or symptoms during their career, due to the high levels of ventilation sustained during prolonged training periods in very specific environmental conditions (i.e. inhalation of cold dry air in winter sports athletes and chlorine by-products in swimmers) (1-3). On one hand, winter sports athletes frequently report exercise-induced respiratory symptoms, especially cough, due to cold dry air inhalation and disturbing their training or performance, but not necessarily related to exercise-induced bronchoconstriction (EIB) or airway hyperresponsiveness (AHR) (1,4-6). On the other hand, a majority of competitive swimmers present with AHR to direct or indirect challenge in a Lab, without complaining of respiratory symptoms or having EIB during swimming, except maybe when chlorine byproducts are elevated in the atmosphere of swimming pools (1,7-9). Both categories of competitive athletes present characteristics of airway remodeling of the extracellular matrix (ECM), such as fibrosis of the submucosa and reticular basement membrane thickening, whatever the AHR status to methacholine (3,10). The inflammatory profile in airway tissue looks, however, different in competitive cross-country skiers and swimmers, characterized by a marked increase in T-lymphocytes and macrophage cell counts in the former and mast cells infiltration in the latter (3,10). All these observations may indicate a different airway modification phenotype according to the training environment.

In some phenotypes of asthma, an imbalance between the expression of proteases and anti-proteases within the ECM, and particularly of matrix metalloproteinase-9 (MMP-9) and its inhibitor (TIMP-1), is believed to be a biomarker of tissue destruction or abnormal repair, and fibrosis (11-15). The activation of epithelial cells, and the recruitment of neutrophil and mast cells (16-20), as observed during exercise ventilation, may stimulate the secretion or release

of matrix metalloproteinase-9 (MMP-9) (21-24). In excess, MMP-9 and tissue-inhibitor

(TIMP-1) have been shown to lead to airway tissue damage and fibrosis in asthma (12-15,25).

In order to distinguish between both potential phenotypes, we compared the ratio of MMP-9

and its inhibitor (TIMP-1) during the annual beginning of the new training season in winter

sport athletes and swimmers, in relation with other classical biomarkers of inflammation.

Whereas a low MMP-9/TIMP-1 ratio would rather reflect a predominant fibrogenesis over

airway inflammation, an increased ratio would indicate a predominant inflammatory process,

which seems also to predict a better response to inhaled corticosteroids in asthma (26). The

secondary objective was to observe the variations of MMP-9/TIMP-1 ratio between winter

and summer, in winter sports athletes only, because if swimmers are exposed to the pool air

all year round, winter sport athletes are subjected to climatic variations related to their sport.

MATERIAL AND METHODS

Participants

Athletes were active competitors, 14 years of age or older, and trained at least 10 hours per

week in their sport during the last year. A group of non-athletes, non-asthmatic subjects with

no asthma-like symptoms and not involved in a competitive sport was also recruited. They

were paired for age (± 5 years) and sex with swimmers. All participants were non-smokers

and non-obese. All subjects gave their written informed consent and the study protocol was

approved by the IUCPQ Institutional Ethics Committee (CER 20159 and CER 20141). Data

from the lung function and responsiveness analysis between both populations were previously

published (1).

Study design

The visits were performed at the beginning of a new training season, mainly from the end of

August to early October in swimmers and non-athletes, and during the summer in winter sport

athletes. To check a potential change in serum biomarkers during the seasons, 12 winter sport

athletes also had a blood sample during the competitive season in winter. Each test was

performed at least 14h after the last low intensity training session for all participants. A

physical examination and allergy skin prick tests were done and a locally developed

standardized questionnaire was administered to all the subjects regarding respiratory health

and sport activities. Blood samples were taken before performing an eucapnic voluntary

hyperpnea challenge (EVH), followed by a methacholine inhalation test (MIT) approximately

45 min later. Both tests were performed to characterize airway responsiveness to indirect and

direct stimuli, respectively. A sputum induction was also carried out in all the subjects, about

one hour after the end of EVH test. The order of the tests was the same at each visit.

Atopy

Atopic status was determined by skin-prick tests with common airborne allergens, including animal danders, house-dust mite, mixed trees, mixed grasses, pollens, and molds. The presence of atopy was defined as at least one positive (>3 mm mean wheal diameter at 10 min) response to allergens in the presence of a negative saline control and a positive skin response to histamine.

Lung function

Spirometry was carried out according to the American Thoracic Society (ATS) specifications (27). Predicted spirometric values were defined according to Knudson et al. (28). Three reproducible measurements of forced expiratory volume in one second (FEV_1) and forced vital capacity (FVC) were obtained.

Airway responsiveness to Eucapnic Voluntary Hyperpnea and Methacholine challenge

Eucapnic voluntary hyperpnea challenge (EVH) and Methacholine inhalation test (MIT) were performed using the method described by Anderson and Brannan (29), and the 2-minute tidal breathing method (30), respectively. The methacholine PC_{20}, which is the concentration of methacholine inducing a 20% fall in FEV_1, and the post-EVH fall in FEV_1 were used for analysis. Airway hyperresponsiveness (AHR) to EVH was defined by a FEV_1 fall of at least 10% sustained for at least two consecutive time-points after EVH and to methacholine by a $PC_{20} < 16$ mg.mL^{-1}.

Induced sputum

Sputum was induced using the method described by Pizzichini et al. (31) and processed

within 2 h following induction. Briefly, mucus plugs were selected from saliva, weighed,

treated with four times their volume of dithiothreitol (DTT) (Cal-biochem Corp, La Jolla, CA,

USA) and rocked for 15 min. The reaction was stopped with an equal volume of Dulbecco's

phosphate buffered saline (D-PBS) (Life Technologies, Gibco BRL, Burlington, ON,

Canada), filtered and counted to determine total cell count and viability. The samples were

centrifuged and supernatants frozen at −80°C until further analysis.

Blood sampling

Blood was sampled into gel tube with a clotting accelerator. Just after sampling, the tube was

let in ice in a standing position for about 30 minutes to let the clot to form. Then, the blood

was centrifuged at 4°C, 1100g for 20 minutes, and micro-tubes were stored at -80°C until

analysis.

Serum and sputum analysis

Blood level of fibrinogen and C-reactive protein (CRP) were measured at the biochemistry

service of the hospital by colorimetry and turbidimetry, respectively. Serum club cell proteins

(CC16) and surfactant proteins D (SP-D) were measured by ELISA (Biovendor, LLC,

Candler, NC, USA), according to the manufacturer's instructions. Concentrations were

determined using a microplate reader (Molecular Devices, Microplate Reader, Sunnyvale,

CA, USA). The sensitivity of the assay was 500 pg.ml^{-1} for CC-16 and 2.2 ng.ml^{-1} for SP-D.

The ratio of CC16/SP-D was calculated and kept for analysis. Serum and sputum-supernatant

of MMP-9 and TIMP-1 measurements were also done by enzyme-linked immunosorbent

assay (Amersham, Oakville, ON, Canada), according to the manufacturer's instructions. The

sensitivity of the assay was 0.156 ng.mL^{-1} for MMP-9 and 0.08 ng.mL^{-1} for TIMP-1. The

ratios of MMP-9/TIMP-1 were then calculated.

Statistical analysis

Quantitative data were expressed as mean ± SD, or median [Interquartile range].

Methacholine PC_{20} is expressed as geometrical mean [Interquartile range]. Qualitative data

are expressed as the number of subjects and/or percentages. Univariate normality assumptions

were verified with the Kolmogorov-Smirnov test. To compare the groups, one-way ANOVA

was used for age, height, training (y/week), spirometric values, serum MMP-9, and Kruskal-

Wallis test was used to compare mass, training (h/week), percentage fall in FEV_1 post-EVH,

methacholine PC_{20}, serum fibrinogen, CRP, TIMP-1, CC16, SP-D, CC16/SP-D ratio, and

MMP-9/TIMP-1 ratio. To compare sputum values between swimmers and winter sport

athletes, a paired-t test was used for neutrophil, macrophage and lymphocyte cell counts,

MMP-9, MMP-9/TIMP-1 and a Mann-Whitney test for sputum total cell count, eosinophil

and bronchial cell counts, and TIMP-1. The Dunn's or Holm-Sidak's multiple comparisons

procedure were applied post hoc to the ANOVA. To compare data between winter and

summer, a paired t-test or non-parametric Wilcoxon test was used in winter sport athletes,

when normality or equality of variance failed. Correlations were evaluated using Pearson's

(r_p) or Spearman (r_s) test and the significance using a one-sample *t* test. The software used for

statistical analysis was Sigmastat (version 3.5, Systat Software GmbH, Erkrath, Germany)

with a limit of significance of $p < 0.05$.

Results

The characteristics of the participants are presented in Table 1. We were able to analyse

samples from 8 controls, 25 swimmers and 41 winter athletes.

Comparison of biomarkers in serum and sputum of swimmers, winter sports athletes and non-

athletes

No significant difference was observed in serum CRP, fibrinogen, CC16/SP-D ratio, MMP-9,

or MMP-9/TIMP-1 ratio between the three groups, during the same training season (Figure 1).

The level of serum TIMP-1 was significantly higher in swimmers compared with winter

sports athletes (median [interquartile range]: 208 ng.mL^{-1} [184-224] in swimmers *vs* 182

ng.mL^{-1} [162-195] in winter sports athletes, p=0.005), but no difference was observed

between both groups of athletes and controls (median [interquartile range] in controls: 188

ng.mL^{-1} [155-194]).

Sputum was obtained in 10 swimmers (40%), and 19 winter sports athletes (46%) and in only

2 controls. Only data from athletes were thus analyzed for sputum (Table 2). No significant

difference was observed in sputum MMP-9, TIMP-1 or MMP-9/TIMP-1 ratio between both

groups of athletes (Figure 2).

No significant correlation was observed between MMP-9, TIMP-1, MMP-9/TIMP-1 in serum

or sputum and training history, training hours /week, FEV$_1$, FVC, MIT PC$_{20}$, the maximum

fall in FEV$_1$ post-EVH, AHR status to MIT and EVH, serum CRP or fibrinogen, serum

CC16/SP-D and sputum inflammatory cells in the whole athletes group, in swimmers or in

winter sport athletes only. Sputum MMP-9 and TIMP-1 correlated positively in the whole

group of athletes (r$_s$=0.47, p=0.013), but no correlation was observed between serum MMP-9

and TIMP-1 values (p=0.96). Sputum MMP-9 in athletes (n=27) correlated to serum

CC16/SP-D ratio (r_s=0.45, p=0.019), and MIT PC_{20} (r_s=-0.47, p=0.013) (Figure 3).

Comparison of MMP-9 and TIMP-1 during winter and summer seasons in winter sports

athletes

Among the 12 winter sports athletes, 11 had atopy, including 9 seasonal allergy. In these

athletes, there was a significant increase from the winter to the summer in FEV_1 (4.67 ± 0.96

L *vs* 4.88 ± 0.95 L, respectively; p = 0.008) and in FVC (5.63 ± 1.13 L *vs* 5.92 ± 1.20 L,

respectively; p = 0.003). The difference in FEV_1 between both seasons was of 207 ± 222 mL

(4.29 ± 4.32 %) and of 285 ± 255 mL (4.60 ± 4.07 %) for FVC. Methacholine PC_{20} was not

significantly different from the winter to the summer (p=0.30). The FEV_1 fall in EVH was

significantly higher during the summer compared to winter (10.5% [7.2 - 14] *vs* 7.6% [4.8 –

11.3], respectively; p = 0.002). The minimum FEV_1 post-EVH was not significantly different

between both seasons (4.36 ± 0.93 L *vs* 4.30 ± 0.93 L, in winter and in summer, respectively;

p = 0.72).

Serum MMP-9 and MMP-9/TIMP-1 ratio were significantly greater, whereas serum TIMP-1

trend to be lower, during the summer (Figure 4). Serum fibrinogen was significantly lower

during the summer (2.84 ± 0.35 g.L^{-1} in winter *vs* 2.50 ± 0.35 g.L^{-1} in summer; p = 0.04), but

no significant difference was observed in serum CRP (p = 0.87). No significant correlation

was observed between changes from summer to the winter of serum MMP9, TIMP-1,

MMP9/TIMP-1 ratio and the changes of FEV_1, FVC, MIT PC_{20} or FEV_1 fall post-EVH.

Discussion

The aims of the study were to compare serum and sputum concentrations of MMP-9 and TIMP- in winter sport athletes and swimmers in relation with airway function at a similar period of their competition season, and to observe the variations from winter to summer in winter sport athletes. No significant difference in serum or sputum concentrations of MMP-9, MMP-9/TIMP-1 or sputum TIMP-1 was observed between swimmers, winter sport athletes and controls. Sputum MMP-9 correlated with Methacholine PC_{20} and serum CC16/SP-D ratio, suggesting a link between the degree of responsiveness, epithelial damage and MMP-9 in the airways, despite no correlation between CC16/SP-D and MIT PC_{20}. Finally, in winter sport athletes, the season of measurement had an impact on serum MMP-9 and TIMP-1, with a higher MMP-9/TIMP-1 ratio observed during the summer, not related to airway function or responsiveness.

The lack of difference in serum MMP-9 and TIMP-1 levels, and MMP-9/TIMP-1 ratio in both athletes' category and controls, indicates probably that these biomarkers, when measured in blood, are not sensitive nor specific enough in this population to reflect local changes in the airways, that occur after years of training although it does exclude the possibility that differences could only have been seen after an intense exercise (dynamic process). In that sense, there was no correlation between airway biomarkers of inflammation, responsiveness or function, and serum level of biomarkers. Serum TIMP-1 was lightly but significantly increased in swimmers compared with winter sport athletes, probably resulting from a slight different training recovery state, as TIMP-1 in induced sputum was similar in both groups of athletes. Some studies reported an acute increase in blood MMP-9 after intense exercise, but their origin remains to be determined as studies bring contradictory results (32-38). MMP-9 may originate from several tissues after exercising and participates to the cleaving of muscle-

specific proteins and extracellular matrix formation, remodeling and regeneration in skeletal

and cardiac muscle, as well as the surrounding vasculature (32,33,36,39).

Blood MMP-9 level may modulate the activation of growth factor cytokines and

angiogenesis, thus facilitating physiological adaptations to exercise training. Certain stimuli,

especially those producing an important mechanical stress, activate the local production of

MMP-9 in the skeletal muscle (32,35), and probably also in the airways. The slight increase in

serum TIMP-1 unlikely originates from the airways but may reflect an incomplete tissue

repair process in the swimmers, due to an insufficient time to recover from the last training

and/or chlorine exposure. We chose the same season to compare serum from swimmers and

winter sport athletes as it is their annual period of training resumption. However, the new

swimming training year, that was also an Olympic year for summer sports, began with a high

volume of training, compared with the beginning of the winter sport athletes' season. In the

summer, in winter sport athletes, the serum MMP-9/TIMP-1 ratio was increased whereas the

serum TIMP-1 and fibrinogen were decreased compared to the winter season. This suggests

an increased airway inflammatory process during the summer, and fibrogenesis in winter.

This is an interesting observation meaning that these biomarkers are sensitive to training but

also probably to the environment. As all, except 3, had seasonal atopy, we can not exclude a

role of allergies in the increase of serum MMP-9.

In acute inflammation due to ozone inhalation, MMP-9 may play a protective role against

inflammation in limiting neutrophils' recruitment and protecting from epithelial injuries and

increases in permeability (40). MMP-9 have also been shown to promote the migration of

repairing bronchial epithelial cells after injury (41). As Vermeer et al. (42) observed *in vitro* a

loss of bronchial epithelium integrity through the cleavage of one or more components of

tight junctions by MMP-9, the relationship between epithelial damage and MMP-9 remains to

be clarified. In our study, sputum MMP-9 correlated with serum CC16/SP-D ratio in all the

athletes, confirming the link between the loss of epithelial integrity and MMP-9 in the

airways, regardless of the sense of the relation between both.

Few studies have investigated the relationship between lung function and MMP-9 and results

are inconsistent (43,44). In general, in mild stable asthma, there is no link between baseline

values of airway MMP-9, that are often similar to the level observed in control subjects, and

lung function (44); but a link is observed between both variables when measured after an

allergen challenge (45) or in untreated mild asthmatics having a higher number of neutrophils

and macrophages in sputum compared with control subjects (43). Todorova et al. (46)

suggested that during a stable period of disease, the level of MMP-9 may de determined by

their production by structural cells, i.e. bronchial fibroblasts, epithelial cells and smooth

muscle cells, whereas during asthma exacerbation, their major production source are

inflammatory cells (46).

In their study, the authors reported a positive relationship between the activity of MMP-9 in

fibroblasts of stable mild to moderate asthma patients and the dose of methacholine inducing

a 20% fall in FEV_1 (r=0.86, p=0.01) (46). In our study, there is a slight correlation between

the level of MMP-9 in sputum and MIT PC_{20} (rs=0.45, p=0.019), but which is negative. The

activity of MMP-9 in the study from Todorova et al. (46) was confined to cultured fibroblasts

of asthmatics, whereas in our study they probably originate also from epithelial, and

inflammatory cells. Therefore, the level of MMP-9 may reflect an underlying inflammatory

cells activity, and a slight damage in epithelial integrity. In that sense, sputum MMP-9 also

correlated to serum CC16/SP-D ratio.

A limitation is that we had insufficient induced sputum to analyze in winter sport athletes,

and, therefore, we had no measurement of MMP-9 and TIMP-1 in sputum during winter and

summer. This may have help us to better understand the origin of MMP-9 during both

seasons. However, as induced sputum is sometimes difficult to obtain in athletes (47), we

thought serum biomarkers as an attractive alternative. However, we found no correlation

between serum and sputum values. As all, except 3, had seasonal atopy, we can not exclude a

role of allergies in the increase of serum MMP-9.

To conclude, we found no major differences in the level of MMP-9 and TIMP-1 in the serum

of winter sport athletes and swimmers, except for a slight increase in TIMP-1 in swimmers.

This was not correlated to airway responsiveness, lung function, or the level of airway

inflammation, indicating probably a more systemic origin of these markers, related to training

and environment. Sputum level of MMP-9 correlated to serum CC16/DP-D ratio and

Methacholine PC_{20}, suggesting a probably indirect link between the degree of responsiveness,

epithelial damage and MMP-9 in the airways. Finally, as serum MMP-9/TIMP-1 ratio was

increased in the summer compared with the winter in elite athletes, this may indicate an

important role of training and/or environment on these biomarkers, that may also affect the

airways. But this remains to be determined. Our study underlines the difficulty to assess

inflammation and remodeling mechanisms in athletes' airways, as serum biomarkers do not

reflect the airways' activity.

Perspectives

In a near future, priority should be given to search for easy tools and specific protocols to

better understand the development of airway structural and functional changes in athletes,

according to the training and environmental stimuli. This will make it possible to prevent and

better manage respiratory diseases that develop during sports careers, and to identify the

phenotypes resulting in identical symptoms.

Conflict of interest

The authors declare (1) they have no relationship with companies or manufacturers who will

benefit from the results of the present study, (2) the results of the present study do not

constitute endorsement by ACSM, (3) the results of the study are presented clearly, honestly,

and without fabrication, falsification, or inappropriate data manipulation.

References

1. Bougault V, Turmel J, Boulet LP. Bronchial challenges and respiratory symptoms in

elite swimmers and winter sport athletes: Airway hyperresponsiveness in asthma: its

measurement and clinical significance. *Chest*. 2010;138(2 Suppl):31S-37S

2. Mountjoy M, Fitch K, Boulet LP, Bougault V, van Mechelen W, Verhagen E.

Prevalence and characteristics of asthma in the aquatic disciplines. *J Allergy Clin*

Immunol. 2015;136(3):588-94

3. Karjalainen EM, Laitinen A, Sue-Chu M, Altraja A, Bjermer L, Laitinen LA.

Evidence of airway inflammation and remodeling in ski athletes with and without

bronchial hyperresponsiveness to methacholine. *Am J Respir Crit Care Med*.

2000;161(6):2086-91

4. Rundell KW, Anderson SD, Sue-Chu M, Bougault V, Boulet LP. Air quality and

temperature effects on exercise-induced bronchoconstriction. *Compr Physiol*.

2015;5(2):579-610

5. Stenfors N. Self-reported symptoms and bronchial hyperresponsiveness in elite cross-

country skiers. *Respir Med*. 2010;104(11):1760-3

6. Boulet LP, Turmel J, Irwin RS; CHEST Expert Cough Panel. Cough in the Athlete:

CHEST Guideline and Expert Panel Report. *Chest*. 2017;151(2):441-54

7. Potts JE Adverse respiratory health effects of competitive swimming: the prevalence

of symptoms, illness, and bronchial responsiveness to methacholine and exercise

[dissertation]. Vancouver (BC): University of British Columbia; 1994, 210 p.

8. Lévesque B, Duchesne JF, Gingras S, et al. The determinants of prevalence of health

complaints among young competitive swimmers. *Int Arch Occup Environ Health*.

2006;80(1):32-9

9. Couillard S, Bougault V, Turmel J, Boulet LP. Perception of bronchoconstriction
following methacholine and eucapnic voluntary hyperpnea challenges in elite athletes.
Chest. 2014;145(4):794-802

10. Bougault V, Loubaki L, Joubert P, et al. Airway remodeling and inflammation in
competitive swimmers training in indoor chlorinated swimming pools. *J Allergy Clin
Immunol.* 2012;129(2):351-8

11. Tanaka H, Miyazaki N, Oashi K, Tanaka S, Ohmichi M, Abe S. Sputum matrix
metalloproteinase-9: tissue inhibitor of metalloproteinase-1 ratio in acute asthma. *J
Allergy Clin Immunol*. 2000;105(5):900-5

12. Gueders MM, Foidart JM, Noel A, Cataldo DD. Matrix metalloproteinases (MMPs)
and tissue inhibitors of MMPs in the respiratory tract: potential implications in asthma
and other lung diseases. *Eur J Pharmacol*. 2006;533(1-3):133-44

13. Mautino G, Capony F, Bousquet J, Vignola AM. Balance in asthma between matrix
metalloproteinases and their inhibitors. *J Allergy Clin Immunol*. 1999;104(3 Pt 1):530-
3

14. Mautino G, Henriquet C, Jaffuel D, Bousquet J, Capony F. Tissue inhibitor of
metalloproteinase-1 levels in BAL fluid from asthmatics. *Am J Respir Crit Care Med.*
1999;60:324-30

15. Mautino G, Oliver N, Chanez P, Bousquet J, Capony, F. Increased release of matrix
metalloproteinase-9 in bronchoalveolar lavage fluid and by alveolar macrophages of
asthmatics. *Am J Respir Cell Mol Biol*. 1997;17:583–92.

16. Chimenti L, Morici G, Paternò A, et al. Bronchial epithelial damage after a half-
marathon in nonasthmatic amateur runners. *Am J Physiol Lung Cell Mol Physiol*.
2010:298:L857-62

17. Nanson CJ, Burgess JL, Robin M, Bernard AM. Exercise alters serum pneumoprotein

384   concentrations. *Respir Physiol*. 2001;127:259-65

385   18. Boulet LP, Turcotte H, Langdeau JB, Bernier MC. Lower airway inflammatory

386   responses to high-intensity training in athletes. *Clin Invest Med*. 2005;28(1):15-22

387   19. Kippelen P, Larsson J, Anderson SD, Brannan JD, Dahlén B, Dahlén SE. Effect of

388   sodium cromoglycate on mast cell mediators during hyperpnea in athletes. *Med Sci*

389   *Sports Exerc*. 2010;42(10):1853-60

390   20. Hallstrand TS, Kippelen P, Larsson J, et al. Where to from here for exercise-induced

391   bronchoconstriction: the unanswered questions. *Immunol Allergy Clin North Am*.

392   2013;33(3):423-42

393   21. Barbaro MP, Spanevello A, Palladino GP, Salerno FG, Lacedonia D, Carpagnano GE.

394   Exhaled matrix metalloproteinase-9 (MMP-9) in different biological phenotypes of

395   asthma. *Eur J Intern Med*. 2014;25(1):92-6

396   22. Cundall M, Sun Y, Miranda C, Trudeau JB, Barnes S, Wenzel SE. Neutrophil-derived

397   matrix metalloproteinase-9 is increased in severe asthma and poorly inhibited by

398   glucocorticoids. *J Allergy Clin Immunol*. 2003;112(6):1064-71

399   23. Mattos W, Lim S, Russell R, Jatakanon A, Chung KF, Barnes PJ. Matrix

400   metalloproteinase-9 expression in asthma: effect of asthma severity, allergen

401   challenge, and inhaled corticosteroids. *Chest*. 2002;122(5):1543-52

402   24. Ventura I, Vega A, Chacón P, et al. Neutrophils from allergic asthmatic patients

403   produce and release metalloproteinase-9 upon direct exposure to allergens. *Allergy*.

404   2014;69(7):898-905

405   25. Grzela K, Litwiniuk M, Zagorska W, Grzela T. Airway Remodeling in Chronic

406   Obstructive Pulmonary Disease and Asthma: the Role of Matrix Metalloproteinase-9.

407   *Arch Immunol Ther Exp (Warsz).* 2016;64(1):47-55

26. Bossé M, Chakir J, Rouabhia M, Boulet LP, Audette M, Laviolette M. Serum matrix metalloproteinase-9:Tissue inhibitor of metalloproteinase-1 ratio correlates with steroid responsiveness in moderate to severe asthma. *Am J Respir Crit Care Med*. 1999;159(2):596-602

27. Miller MR, Hankinson J, Brusasco V, Burgos F, Casaburi R, Coates A, et al. Standardisation of spirometry. *Eur Respir J*. 2005;26:319–38

28. Knudson RJ, Lebowitz MD, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow-volume curve with growth and aging. *Am Rev Respir Dis*. 1983;127(6):725-34

29. Anderson SD, Brannan JD. Methods for "indirect" challenge tests including exercise, eucapnic voluntary hyperpnea, and hypertonic aerosols. *Clin Rev Allergy Immunol*. 2003; 24:27–54.

30. Crapo RO, Casaburi R, Coates AL, Enright PL, Hankinson JL, et al. Guidelines for methacholine and exercise challenge testing-1999. This official statement of the American Thoracic Society was adopted by the ATS Board of Directors, July 1999. *Am J Respir Crit Care Med*. 2000; 161:309–29

31. Pizzichini MM, Popov TA, Efthimiadis A, et al. Spontaneous and induced sputum to measure indices of airway inflammation in asthma. *Am J Respir Crit Care Med*. 1996; 154: 866–9

32. Carmeli E, Moas M, Lennon S, Powers SK. High intensity exercise increases expression of matrix metalloproteinases in fast skeletal muscle fibres. *Exp Physiol*. 2005;90(4):613-9

33. Saenz AJ, Lee-Lewandrowski E, Wood MJ, et al. Measurement of a plasma stroke

biomarker panel and cardiac troponin T in marathon runners before and after the 2005

Boston marathon. *Am J Clin Pathol*. 2006;126(2):185-9

34. Reihmane D, Jurka A, Tretjakovs P. The Relationship Between Maximal Exercise-

Induced Increases in Serum IL-6, MPO and MMP-9 Concentrations. *Scand J*

Immunol. 2012;76(2):188-92

35. Rullman E, Rundqvist H, Wågsäter D, et al. A single bout of exercise activates matrix

metalloproteinase in human skeletal muscle. *J Appl Physiol (1985)*. 2007;102(6):2346-

51

36. Rullman E, Olsson K, Wågsäter D, Gustafsson T. Circulating MMP-9 during exercise

in humans. *Eur J Appl Physiol*. 2013;113(5):1249-55

37. Mazani M, Fard AS, Baghi AN, Nemati A, Mogadam RA. Effect of pomegranate

juice supplementation on matrix metalloproteinases 2 and 9 following exhaustive

exercise in young healthy males. *JPMA*. 2014;64:785-90

38. Nascimento Da C, Durigan R, Tibana RA, Durigan JL, Navalta JW, Prestes J. The

response of matrix metalloproteinase-9 and -2 to exercise. *Sports Med*.

2015;45(2):269-78

39. Koskinen SO, Wang W, Ahtikoski AM, et al. Acute exercise induced changes in rat

skeletal muscle mRNAs and proteins regulating type IV collagen content. *Am J*

Physiol Regul Integr Comp Physiol. 2001;280(5):R1292-300

40. Yoon HK, Cho HY, Kleeberger SR. Protective role of matrix metalloproteinase-9 in

ozone-induced airway inflammation. *Env Health Perspect*. 2007;115(11):1557-63

41. Legrand C, Gilles C, Zahm JM, et al. Airway epithelial cell migration dynamics.

MMP-9 role in cell-extracellular matrix remodeling. *J Cell Biol*. 1999;146(2):517-29

42. Vermeer PD, Denker J, Estin M, et al. MMP9 modulates tight junction integrity and
cell viability in human airway epithelia. *Am J Physiol Lung Cell Mol Physiol*.
2009;296(5):L751-62

43. Vignola AM, Riccobono L, Mirabella A, et al. Sputum metalloproteinase-9/tissue
inhibitor of metalloproteinase-1 ratio correlates with airflow obstruction in asthma and
chronic bronchitis. *Am J Respir Crit Care Med*. 1998;158(6):1945-50

44. Ko FW, Diba C, Roth M, et al. A comparison of airway and serum matrix
metalloproteinase-9 activity among normal subjects, asthmatic patients, and patients
with asthmatic mucus hypersecretion. *Chest*. 2005;127:1919-27

45. Cataldo DD, Bettiol J, Noël A, Bartsch P, Foidart JM, Louis R. Matrix
metalloproteinase-9, but not tissue inhibitor of matrix metalloproteinase-1, increases in
the sputum from allergic asthmatic patients after allergen challenge. *Chest*.
2002;122(5):1553-9

46. Todorova L, Bjermer L, Miller-Larsson A, Westergren-Thorsson G. Relationship
between matrix production by bronchial fibroblasts and lung function and AHR in
asthma. *Respir Med*. 2010;104:1799-808

47. Bougault V, Turmel J, St-Laurent J, Bertrand M, Boulet LP. Asthma, airway
inflammation and epithelial damage in swimmers and cold-air athletes. *Eur Respir J*.
2009;33(4):740-6

Table 1: Characteristics of the participants

	Controls	Swimmers	Winter sports athletes
Subjects (n)	8	25	41
Age (years)	23 ± 2	$19 \pm 3^{*}$	$21 \pm 2^{*}$
Sex (M/F)	6/2	$18/7^{*\dagger}$	$26/15^{*\S}$
BMI (kg.m^{2})	22.3 ± 2.8	21.3 ± 1.7	22.2 ± 1.3
Atopy n (%)	6 (75)	22 (88)	30 (73)
Training (years)	3 ± 3	$11 \pm 3^{*}$	$10 \pm 3^{*}$
Training (hours per week)	5 ± 3	$25 \pm 4^{*\dagger}$	$16 \pm 6^{*\S}$
FEV$_1$ (% pred.)	94 ± 10	$116 \pm 17^{*}$	$112 \pm 12^{*}$
FVC (% pred.)	100 ± 8	$128 \pm 15^{*\dagger}$	$118 \pm 13^{*\S}$
MIT PC$_{20}$ (mg/mL)	72.2 [41.7-103.2]	$7.7 [2.5-14.1]^{*\dagger}$	$42.0 [20.2-128]^{\S}$
MIT PC$_{20}$ < 16 mg.mL^{-1} n (%)	1 (13)	$19 (76)^{*\dagger}$	$9 (23)^{\S}$
EVH (% fall in FEV$_1$)	9.6 [8.3-14.1]	$13.1 [10.5-23.0]^{\dagger}$	$9.1 [7.6-12.2]^{\S}$
HIB to EVH n (%)	4 (50)	$17 (71)^{\dagger}$	$15 (38)^{\S}$

Data are expressed as mean $\pm$ SD, or as number of subjects (n, percentage). MIT PC$_{20}$ is

expressed as geometrical mean [interquartile range] and EVH (% fall in FEV$_1$) is expressed as

median [interquartile range]. M: male; F: female; BMI: Body mass index; FEV$_1$: forced

expiratory volume in one second; FVC: forced vital capacity; % pred: % of the predicted

value; MIT PC$_{20}$: concentration of methacholine inducing a 20% fall in FEV$_1$; EVH: eucapnic

voluntary hyperpnea challenge; HIB: Hyperventilation-induced bronchoconstriction defined

as a fall of at least 10% in FEV$_1$ after the 6-min EVH challenge. *p < 0.05 compared with

controls; §p < 0.05 compared with swimmers; †p < 0.05 compared with winter sport athletes.

Table 2: Sputum cell counts characteristics in participants.

	Swimmers	Winter sports athletes
Subjects (n)	10	18
Total cell count ($\times 10^6$ cells)	0.485 [0.203-1.745]	0.366 [0.143-0.986]
Neutrophils (%)	51 [46-61]	36 [24-49]
Eosinophils (%)	0.25 [0-1.75]	0 [0-0.3]
Macrophages (%)	42 [30-46]	55 [32-69]
Lymphocytes (%)	1.5 [0.3-2.0]	1.1 [0.8-2.0]
Bronchial epithelial cells (%)	2.5 [1.0-8.3]	2.1 [1.5-3.6]

Data are expressed as median [interquartile range]

[§]p < 0.05 compared with swimmers; [†]p < 0.05 compared with winter sport athletes.

Figures Legend

Figure 1 A to F: Serum values of inflammation/remodeling biomarkers in controls, swimmers and winter sports athletes.

(A) MMP-9: Metalloproteinase-9; (B) TIMP-1: Tissue inhibitor of metalloproteinase-1; (C) MMP-9/TIMP-1 ratio; (D) CRP: C-reactive protein; (E) Fibrinogen; (F) CC16: Club cell proteins; SP-D: surfactant protein-D. A value of CC16/SP-D ratio of a control subject was retired from the graph but not statistical analysis, as it was an extreme data. The horizontal bar represents the median for each group.

** Post-hoc P <0.01 between groups

Figure 2: Values of inflammatory cells in induced sputum of athletes.

MMP-9: Metalloproteinase-9; TIMP-1: Tissue inhibitor of metalloproteinase-1. The horizontal bar represents the median for each group.

Figure 3: Correlation between sputum MMP-9 and (A) Methacholine airway responsiveness, and (B) serum CC16/SP-D ratio.

rs : Spearman correlation

Figure 4: Serum values of inflammation/remodeling biomarkers in winter sport athletes during the winter and the summer.

MMP-9: Metalloproteinase-9; TIMP-1: Tissue inhibitor of metalloproteinase-1. The horizontal bar represents the median for each season.

CC16: Club cell proteins; MMP-9: Metalloproteinase-9; Methacholine PC_{20}: concentration of methacholine inducing a 20% fall in FEV_1; SP-D: surfactant protein-D; rs: Spearman correlation coefficient.

Figure 1

Figure 2

Figure 3

Figure 4