**Evidence of stock connectivity, hybridization and misidentification in white anglerfish support the need of a genetics-informed fisheries management framework**

**Running title: White anglerfish genetics and management**

Imanol Aguirre-Sarabia[1], Natalia Díaz-Arce[1], Iker Pereda-Agirre[1], Iñaki Mendibil[1], Agurtzane Urtizberea[1], Hans D. Gerritsen[2], Finlay Burns[3], Ian Holmes[4], Jorge Landa[5], Ilaria Coscia[6], Iñaki Quinconces[1], Marina Santurtún[1], Antonella Zanzi[7], Jann T. Martinsohn[7], Naiara Rodríguez-Ezpeleta[1,*]

[1] AZTI, Marine Research, Basque Research and Technology Alliance (BRTA). Txatxarramendi ugartea z/g, 48395 Sukarrieta – Bizkaia, Spain

[2] Fisheries Ecosystems Advisory Services, Marine Institute, Rinville, Oranmore Co. Galway, H91 R673 Ireland

[3] Marine Scotland Science, Marine Laboratory, PO Box 101, Victoria Road, Torry, Aberdeenshire, AB11 9DB, Scotland, UK

[4] Centre for Environment, Fisheries and Aquaculture Science, Lowestoft Laboratory Pakefield Road NR33 0HT Lowestoft Suffolk United Kingdom

[5] Instituto Español de Oceanografía (IEO), Centro Oceanográfico de Santander. Promontorio San Martín, s/n, 39004 Santander, Spain.

[6] University of Salford, School of Science, Engineering and Environment, The Crescent, Salford, M5 4WT, England, UK

[7] European Commission, Joint Research Centre (JRC), Via Enrico Fermi 2749, 21027 Ispra, Italy

[*] Correspondence: nrodriguez@azti.es

**Abstract**

Understanding population connectivity within a species as well as potential interactions with its close relatives is crucial to define management units and to derive efficient management actions. However, although genetics can reveal mismatches between biological and management units and other relevant but hidden information such as species misidentification or hybridization, the uptake of genetic methods by the fisheries management process is far from having been consolidated. Here, we have assessed the power of genetics to better understand the population connectivity of white angelfish (*Lophius piscatorius*) and its interaction with its sister species, the black anglerfish (*L. budegassa*). Our analyses, based on thousands of genome-wide single nucleotide polymorphisms, show three findings that are crucial for white anglerfish management. We found i) that white anglerfish is likely composed of a single panmictic population throughout the Northeast Atlantic, challenging the three-stock based management, ii) that a fraction of specimens classified as white anglerfish using morphological characteristics are genetically identified as black anglerfish (*L. budegassa*) and iii) that the two *Lophius* species naturally hybridize leading to a population of hybrids of up to 20% in certain areas. Our results set the basics for a genetics-informed white anglerfish assessment framework that accounts for stock connectivity, revises and establishes new diagnostic characters for *Lophius* species identification and evaluates the effect of hybrids in the current and future assessments of the white anglerfish. Furthermore, our study contributes to provide additional evidence of the potentially negative consequences of ignoring genetic data for assessing fisheries resources.

**Introduction**

Sustainable exploitation of fisheries resources relies on effective fisheries management

actions, which in turn rely on accurate fisheries assessment, that is, the process that

includes the synthesis of information on life history, fishery monitoring, and resource

surveys for estimating stock size and harvest rate relative to sustainable reference

points (Methot & Wetzel, 2013). The process of fisheries assessment is usually applied

independently to pre-established management units (so-called stocks), of which

parameters such as growth, recruitment, and natural and fishing mortality are

assumed to be intrinsic and not dependent on emigration or immigration rates (Cadrin,

2020). Genetic data has demonstrated ability to delineate populations within a species,

that is, to identify groups of sexually interbreeding individuals which possess a

common gene pool, but has also revealed hidden phenomena such as species

misidentification (Garcia-Vazquez, Machado-Schiaffino, Campo, & Juanes, 2012) or

hybridization (Pampoulie et al., 2020).

Fisheries stocks are often defined upon political and administrative considerations

(Stephenson, 2002); yet, failure to align stocks with natural populations results in

unfeasibility to establish an accurate relationship between productivity and harvest

rates and can result in local reduction of populations and, in extreme cases, to local

population collapse (Bonfil, 2005). Misidentification can be a common phenomenon

when morphologically similar species are caught simultaneously, and results in misled

exploitation rate estimations when those are based on reported catches (Marko et al.,

2004). Hybridization, the successful reproduction between different species (Arnold,

1997; Stronen & Paquet, 2013), has been reported in teleost fishes (Schwenke, 2013;

Yaakub, Bellwood, van Herwerden, & Walsh, 2006), but incidence and associated

consequences in commercial fisheries has been scarcely explored (Epifanio & Nielsen,

2000); yet, hybridization could play a key role in diversity loss and even parental

species extinction, which has important consequences for management and

conservation (Allendorf, Leary, Spruell, & Wenburg, 2001). An example of species

whose management could largely benefit from genetic-derived information is the

white anglerfish (*Lophius piscatorius*, Linnaeus, 1758), a bottom-living fish that moved

from being a bycatch species in the last century (Hislop et al., 2001) to become one of

the most valuable demersal species in southern and western Europe (ICES, 2010).

The white anglerfish inhabits the Northeast Atlantic and Mediterranean Sea, where it

is assessed by the International Council for the Exploration of the Sea (ICES) and the

General Fisheries Commission for the Mediterranean (GFCM), respectively. Within the

Northeast Atlantic, the white anglerfish is managed as three stocks (Figure 1): the

Northern Shelf stock (Skagerrak, Kattegat, North Sea, West of Scotland and Rockall),

the Northern stock (Celtic Seas and Northern Bay of Biscay) and the Southern Stock

(Atlantic and Iberian waters) (ICES, 2019a, 2019b); yet, the few studies assessing the

population structure, based on otolith shape analysis (Cañás, Stransky, Schlickeisen,

Sampedro, & Fariña, 2012), tagging surveys (J. Landa, Quincoces, Duarte, Fariña, &

Dupouy, 2008; Laurenson, Johnson, & Priede, 2005) and molecular markers, including

allozymes (Crozier, 1987), mitochondrial DNA (Charrier et al., 2006) and microsatellites

(Blanco, Borrell, Cagigas, Vázquez, & Prado, 2008), did not find differences between

stocks. However, this needs to be confirmed with the analysis of a large number of

genomic markers, which has been effective in  resolving the population structure of

marine fish when other markers failed (Leone, Álvarez, García, Saborido-Rey, &

Rodriguez-Ezpeleta, 2019; Rodríguez-Ezpeleta et al., 2016; Rodríguez-Ezpeleta et al.,

2019). The white anglerfish coexists with its sympatric sister species, the black bellied

anglerfish (*Lophius budegassa*), which has a more southern distribution (Relini, 1999;

Ungaro et al., 2002). The outer morphology of both species is similar which is why they

are often confused, although the color of the epithelium that covers the abdominal

cavity, called peritoneum (white in *L. piscatorius* and black in *L. budegassa*) is

considered a unequivocal diagnostic character for individuals larger than 15 cm

(Caruso, 1986). Yet, genetic analyses based on polymerase chain reaction amplification

of restriction fragment length polymorphisms (PCR-RFLP) used for species

identification have revealed mislabeling among the two species (Armani et al., 2012).

There is thus a need for resolving the population structure of white anglerfish within

the Northeast Atlantic, including its relationship with Mediterranean populations, and

for understanding its interaction with its sister species, the black anglerfish. For that

aim, we have used thousands of genome-wide distributed SNP markers discovered and

genotyped through restriction site associated DNA sequencing (RAD-seq) and found i)

that white anglerfish from the three stocks within the Northeast Atlantic are

genetically connected, ii) first evidence of hybridization between white and black

anglerfish and iii) records of black anglerfish misidentified as white anglerfish due to a

lack of black peritoneum. These findings have important implications for white

anglerfish management and conservation, while revealing the significant advantage of

including genomics in fisheries assessment in general.

**Material and methods**

*Tissue sampling, DNA extraction, and species identification*

*Lophius piscatorius* (n=693) and *Lophius budegassa* (n=31) samples were collected

from Northeast Atlantic Ocean and Mediterranean Sea locations using scientific

surveys and commercial fisheries (Table S1; Figure 1). Sampling of *L. piscatorius* was

carried out so as to cover a large part of the geographic range of this species within the

Atlantic, whereas samples of *L. budegassa* were collected opportunistically as they

were only used for comparison purposes. Each individual was assigned to either

species by the color of the peritoneum, white for *L. piscatorius* and black for *L.*

*budegassa* (Caruso, 1986). Maturity was assigned following the 5 stages key (ICES,

2007). From each individual, a ~1 cm$^3$ muscle tissue sample was excised and

immediately stored in 96% molecular grade ethanol at -20 °C until DNA extraction.

Genomic DNA was extracted from about 20 mg of muscle tissue using the Wizard®

Genomic DNA Purification kit (Promega, WI, USA) following the manufacturer's

instructions. Extracted DNA was suspended in Milli-Q water and concentration was

determined with the Quant-iT dsDNA HS assay kit using a Qubit® 2.0 Fluorometer (Life

Technologies). DNA integrity was assessed by electrophoresis, migrating about 100 ng

of Sybr™ Safe-stained DNA on an agarose 1.0% gel. A polymerase chain reaction

restriction fragment length polymorphism (PCR-RFLP) method based on mitochondrial

DNA (Armani et al., 2012) was used for authentication of all specimens collected. In

order to further validate the PCR-RFLP results, for 122 of the samples, a fragment of

the mitochondrial cytochrome b (cytb) gene was amplified with the primers GluFish-F

and THR-Fish-R (Sevilla et al., 2007) using the following amplification conditions:

denaturation at 95 °C for 3 min; 35 cycles at 95 °C for 30 s, 57 °C for 30 s, 72 °C for 60 s;

and final extension at 72 °C for 10 min. The PCR products were purified and sequenced

using Sanger.

*RAD-seq library preparation and sequencing*

Restriction-site-associated DNA libraries were prepared for 306 *L. piscatorius* and 27 *L.*

*budegassa* individuals (Table S1) following the methods of Etter, Bassham, Hohenlohe,

Johnson, and Cresko (2012). Five individuals were run by duplicate starting from the

tissue sample to check for replicability. Between 300 and 600 ng of starting DNA

(depending on integrity) was digested with the *SbfI* restriction enzyme and ligated to

modified Illumina P1 adapters containing 5bp sample-specific barcodes. Pools of 32

individuals were sheared using the Covaris® M220 Focused-ultrasonicator™ Instrument

(Life Technologies) and size selected to 300-500 bp with magnetic beads. Following the

Illumina P2 adaptor ligation, each library was amplified using 12 PCR cycles and

batches of three pools were paired-end sequenced (100 bp) on an Illumina HiSeq4000.

*SNP genotype table generation*

Generated RAD-tags were analyzed using *Stacks* version 2.4 (Catchen, Hohenlohe,

Bassham, Amores, & Cresko, 2013). Quality filtering and demultiplexing were per-

formed using the *Stacks* module *process_radtags* removing reads with ambiguous ba-

ses and with an average quality score lower than 20 in at least one stretch of 15 nucle-

otides and using a maximum of 1 mismatch when rescuing single-end barcodes. Only

reads whose forward and reverse pair passed quality filtering were kept and the mod-

ule *clone_filter* was applied to remove PCR duplicates. *De novo* and reference-based

assemblies were performed for consistency since each approach has disadvantages:

the genome used for the reference-based approach is highly fragmented (Dubin,

Jørgensen, Moum, Johansen, & Jakt, 2019) and represents only one of the species in-

cluded in this study, and the *de novo* approach could be affected by assembly parame-

ters choice. For the *de novo* assembly, the module *ustacks* was used to assemble fil-

tered reads into putative orthologous loci per individual, with a minimum coverage

depth required to create a stack (parameter -m) of 3, and a maximum nucleotide mis-

matches allowed between stacks (parameter -M) of 4. RAD-loci were then assembled

using the module *cstacks* with a maximum of 6 allowed mismatches between sample

loci when generating the catalogue (parameter -n). Matches to the catalogue for each

sample were searched using *sstacks* and transposed using *tsv2bam*. For the reference

mapped assembly, the filtered reads were mapped against the available draft white

anglerfish genome (Dubin et al., 2019) using the BWA-MEM algorithm (Li, 2013), and

the resulting SAM files converted to BAM format, sorted and indexed using SAMTOOLS

(Li et al., 2009). Mapped reads were filtered to include only primary alignments and

correctly mate mapped reads. The following steps were applied to both the *de novo*

and reference mapped catalogs including only samples with a minimum of 30,000

RAD-loci. Paired-end reads were assembled into contigs and SNPs derived were identi-

fied and genotyped using the module *gstacks.* In order to avoid ascertainment bias

(Rodríguez-Ezpeleta et al., 2016), SNP selection was performed separately on each

subset of individuals to be analyzed (Table S2). For that aim, the module *populations*

was used to select the tags present in RAD-loci found in at least 90% of the individuals,

and *PLINK* version 1.07 (Purcell et al., 2007) was used to select samples with a mini-

mum of 0.85 genotyping rate, and SNPs with a minimum of 0.95 genotyping rate and a

minimum allele frequency (MAF) bigger than 0.05. Tags on which the selected SNPs

were located were mapped against the complete mitochondrial genome of *Lophius*

*piscatorius* (NC_036988.1) with the BWA-MEM algorithm (Li, 2013) and it was con-

firmed that the final genotype dataset did not contain mitochondrial SNPs.

*Genetic diversity, population structure and hybridization analyses*

The following analyses were performed using only the first SNP per tag. Expected (He)

and observed (Ho) heterozygosity and average pairwise $F_{ST}$ values were computed with

PLINK (Purcell, 2009) and GENEPOP (Rousset, 2014) respectively. Principal component

analysis (PCA) was performed without any *a priori* assignment of individuals to

populations using the package *adegenet* (Jombart & Ahmed, 2011) in R version 3.5.0

(Team, 2018). The genetic ancestry of each individual was estimated using the model-

based clustering method implemented in ADMIXTURE (Alexander, Shringarpure,

Novembre, & Lange, 2015) assuming from 2 to 5 ancestral populations (K) and setting

1000 bootstrap runs. The value of K with the lowest associated error value was

identified using ADMIXTURE's cross-validation procedure assuming from 1 to 5

ancestral populations. Using *NewHybrids* (E. Anderson & Thompson, 2002), posterior

probabilities of each individual's membership as a pure parent, first (F1) or second (F2)

generation hybrids or backcrosses were calculated applying default parameters to  the

200 SNPs with less than 1% of missing data that presented the highest Fst between

genetically assigned *L. piscatorious* and *L. budegassa* samples.

**Results**

*Genetic species identification based on mitochondrial markers*

In all samples for which it was conducted, cytb gene sequencing confirmed PCR-RFLP

results (Table S1). Species authentication was positive for all 31 specimens identified as

*Lophius budegassa*. However, from the 693 specimens identified as *L. piscatorius*

based on the colour of their peritoneum, 76 were genetically identified as *L.*
*budegassa*. This incongruency between the colour of the peritoneum and the
mitochondrial based genetic identification (Figure 2) suggests that species
identification based on the colour of the peritoneum results in erroneous taxonomic
assignment or that mitochondrial markers are ambiguous for species identification,
which needs to be evaluated using nuclear markers.

*Population structure based on nuclear markers*

The genotype table including individuals from both species resulted in 323 and 326
individuals (of which 27 were *L. budegassa*) and 16712 or 23126 SNPs after filtering
when using a *de novo* or reference mapped assembly respectively (Table S2).
Replicates resulted in 0.9947% identical genotypes. The PCA and ADMIXTURE analyses
based on these datasets grouped most samples in three clearly distinguishable groups,
and results were virtually identical among the *de novo* and reference assembly
datasets (Figure 3; Figure S1) and subsequent analyses were based on the former. In
the PCA (Figure 3A), samples are disposed along the first principal component
(accounting for 80.82% of the total variance) into three main groups. The first group
includes individuals provided, and genetically confirmed, as *L. piscatorius*. The second
group (in the middle) includes 25 individuals initially identified as *L. piscatorius*,
although 22 of them are assigned to *L. budegassa* according to mitochondrial DNA. The
third group includes individuals initially identified as *L. piscatorius* and *L. budegassa*,
but all of them assigned to *L. budegassa* according to mitochondrial DNA. Additionally,
there are five individuals (all with *L. budegassa* mitochondrial DNA) located in between
the main three clusters. This sample grouping is coherent with the different types of

genetic admixture patterns observed according to the presence of two ancestral

populations, best K=2 (Figure 3B): samples included in the first and third groups in the

PCA are not admixed and belong to different ancestral populations, and samples

included in the middle group in the PCA are admixed with equal contribution from

both ancestral populations. Samples located between groups in the PCA show

consistent patterns in admixture analyses with about 75% or 25% of contribution from

the first and third groups, respectively.

*Misidentification and hybridization*

Together, these results indicate that some of the samples morphologically identified as

*L. piscatorius* are *L. budegassa* despite presenting a white peritoneum (Figure 3C).

Samples in the middle group and adjacent ones have a significantly higher observed

average heterozygosity (0.77) than samples in first and third groups (0.05 and 0.03,

respectively), which could indicate that these individuals are hybrids. This was con-

firmed by NewHybrids, which assigned all individuals in the middle group as new gen-

eration hybrids between the two species (F1), and the five remaining individuals as

backcrosses between F1 hybrids and white or black anglerfish (Figure 3C). Mitochon-

drial DNA from all hybrids except for 3 F1 was of *L. budegassa*. Considering the propor-

tion of hybrid individuals within the ones bearing *L. piscatorius* or *L. budegassa* mito-

chondrial DNA as estimated from RAD-sequencing data, we calculated the percentage

of hybrid and misidentified individuals among those provided as *L. piscatorius*. Overall,

we found about 4 and 10% misidentified and hybrid individuals, respectively. However,

these were not distributed homogeneously across stocks and areas within the same

stock (Figure 4). Misidentified individuals were more present in the most southern re-

gion of the Southern stock (Portuguese coast) and Mediterranean Sea, with 1/3 of the

individuals identified as *L. piscatorius* being *L. budegassa,* less frequent in the Northern

stock (2%), and absent in Northern Shelf stock. Hybrid individuals were absent in the

Southern stock and North Sea (belonging to the Northern Shelf stock) and were most

frequent in the northern Bay of Biscay and Celtic seas (13%).

*Connectivity of white anglerfish within the Atlantic*

The genotype table including only those individuals genetically identified as *L.*

*piscatorius* include 238 or 232 individuals and 6233 or 6246 SNPs when including or

not Mediterranean samples, respectively. PCA and ADMIXTURE analyses based on

those datasets (Figure 5) reveal strong differentiation between Mediterranean and

Atlantic samples ($F_{ST}$ = 0.057) and no genetic differentiation within the Atlantic ($F_{ST}$ =

0), as also suggested by ADMIXTURE (best K=1).

**Discussion**

*What´s in a white anglerfish sample?*

We found that several of the samples provided as white anglerfish (*Lophius*

*piscatorius*) by expert scientists involved in surveys targeting anglerfish were indeed

black anglerfish or hybrids. This implies that the commonly used diagnostic character

for species identification, the color of the peritoneum, is not discriminative as all the

black anglerfish and hybrid individuals provided as white anglerfish had white

peritoneum. Indeed, it has already been suggested that some young individuals cannot

be distinguished by the color of the peritoneum (*e.g.* some that show a white

peritoneum with small black dots can be assigned to either black or white anglerfish

when using another diagnostic character such as the number of dorsal fin rays (J.

Landa and A. Antolínez, unpublished data). Here, we found misidentified individuals

that are large (up to 75cm), suggesting that the reason for misidentification is not

related to the size of the specimens. In light of these findings, alternative

characteristics for species identification are needed (e.g. dorsal and anal fin ray counts

or length of the cephalic dorsal fin spines (Caruso, 1986)), so that misidentification

does not affect data collection. Additionally, a way of identifying hybrids would be

needed. Notably, in our dataset, the hybrid size distributions was significantly smaller

than that of the *L. piscatorius* individuals ($p<0.05$ in the Mann-Whitney test), although

the small sample size of hybrids and confounding factors (e.g. different proportion of

hybrids in the different stocks, age classes) prevent drawing conclusions from this fact

at this stage.

*Causes and consequences of interspecific hybridization*

We provide, to the best of our knowledge, the first evidence of natural hybridization

between *Lophius piscatorius* and *L. budegassa*, a phenomenon that can occur between

closely related species sharing morphological, ecological and reproductive

compatibilities (Montanari, Hobbs, Pratchett, Bay, & Van Herwerden, 2014). *Lophius*

*spp.* hybrids were predominant in Northern Bay of Biscay and Celtic seas, where both

species´ distribution range overlap and have been historically abundant (Figure S2).

There is also overlap in both species´ bathymetric distribution (20-1000m and 100-

500m depth respectively for white and black anglerfish)(Azevedo, 1995; Caruso, 1986;

García-Rodríguez, Pereda, Landa, & Esteban, 2005; Quincoces, Santurtún, & Lucio,

1998), and in spawning periods, both spawning in winter and spring (Ofstad, Angus,

Pedersen, & Steingrund, 2013; Quincoces, 2002.). This overlap in space, depth and

time could have even been increased by recent changes of geographical and

bathymetric distributions of both species, potentially as a consequence of climate

change (Maltby, Rutterford, Tinker, Genner, & Simpson, 2020). According to

mitochondrial DNA, which is maternally inherited in most animals (Moritz, Dowling, &

Brown, 1987), we found that most hybrids (27 out of 30) resulted from a black

anglerfish mother and a white anglerfish father. We found a few backcrosses (hybrids

crossed with parental species) and no crosses involving two hybrids. This could be due

the presence of a stable hybrid zone with hybrids being less able to produce offspring

(Hayden et al., 2010; Mirimin et al., 2014) or to recent changes that have induced

hybridization between both species so newly that they did not have time to produce a

hybrid-derived population. The presence of a stable hybrid zone between black and

white anglerfish does not forcedly imply changes in the evolution of the parental

species but could imply management uncertainties. However, if hybridization is recent,

we cannot discard a process of evolutionary novelty (Budd & Pandolfi, 2010), through

parental species acquiring new functions (T. M. Anderson et al., 2009) and even

producing new species (Verheyen, Salzburger, Snoeks, & Meyer, 2003), or of

biodiversity loss, through extinction of parental species (Seehausen, 2006). Thus,

monitoring the hybrid zone and adjacent areas is crucial to understand the role of

hybridization in management and conservation of white and black anglerfish.

*Management implications of stock connectivity, hybridization and misidentification*

Our results show that the Northeast Atlantic and Mediterranean white anglerfish

populations are genetically isolated, challenging previous findings based on genetic

data (Charrier et al., 2006), and that the white anglerfish within the Northeast Atlantic

Ocean constitutes a genetically homogeneous population, shedding light into previous

unconclusive results based on genetic and non-genetic data (Blanco et al., 2008; Cañás

et al., 2012; Crozier, 1987; Charrier et al., 2006; J. Landa et al., 2008; Laurenson et al.,

2005). These results point towards the necessity to harmonize current stock definitions

within the Northeast Atlantic. Morphometric characteristics such as length of maturity

(Duarte, Azevedo, Landa, & Pereda, 2001) or weight-length relationships (Jorge Landa

& Antolínez, 2018) differ between stocks. These differences are potentially due to

sampling biases, differences in scales and/or measurement methods or interannual

differences among studies, as well as the possible influence of different fishing

pressure or environmental conditions between areas. However, they could also be due

to the inclusion of different proportions of misidentified and hybrid individuals in each

stock.

Although in the Northern and Southern stocks the assessment is done separately for

white and black anglerfish, they are combined for management, being Total Allowable

Catches (TAC) for *Lophius spp*. In the Northern Shelf stock, the two species are

assessed and managed as one assuming that black anglerfish in this area are rare.

However, the obvious increase of black anglerfish in this stock (Figure S2) has created

concern and suggestions to separate species have been made (ICES, 2018). The

deliberate (in the North Shelf stock) or non-deliberate (in the Northern and Southern

stocks) inclusion of an unknown (and perhaps variable) proportion of black anglerfish

in the white anglerfish assessment will likely lead to some bias. Therefore, some

corrections could be applied to the total catches and the length composition data.

However, the proportions of black anglerfish included in these assessments seem

relatively small (<10%) and would likely not have a major impact on the assessments.

The implications of including hybrids in the stock assessment are more difficult to

anticipate as biological characteristics of hybrids are unknown. The main concern is

that inclusion of hybrids would lead to erroneous inferences of reproductive potential

because this feature is inferred from the biomass of mature fish and the reproductive

output of anglerfish hybrids is unknown. If, as in many other species (Mallet, 2007),

they have no or neglectable offspring, they should not be included in the spawning

biomass as otherwise they could lead to an overestimation of productivity, even to a

level below of required to support a sustainable fishery (Morgan et al., 2012). On the

other hand, if, as the presence of backcrossed individuals suggest, they are

reproductively viable, further studies are required to assess hybrid fitness compared to

pure individuals.

Despite supporting stock merging, the analyses presented here suggest a series of

concerns that should be considered for white anglerfish management and which affect

differently each of the stocks. The southernmost locations of the Southern Stock and

the Mediterranean Sea are more affected by misidentification, the northernmost

locations of the Northern stock are more affected by hybridization, and the North Shelf

stock does not seem to be affected by either. The over and under representation of

misidentified individuals (black anglerfish with white peritoneum) in the south and

north respectively could be simply explained by differences in black anglerfish

abundance. However, both species are moving northwards, as shown in the species

distribution maps (FigureS2), and so the proportion of misidentified individuals could

increase in northern areas as black anglerfish becomes more abundant. This northward

movement of both species could also contribute to enlarge the hybrid zone towards

northern areas. If the proportion of hybridization and mislabeling varies significantly

over time, then this will have implications for the robustness of the assessment for

these stocks. Thus, genetic studies monitoring white and black anglerfish populations

across the Northeast Atlantic and Mediterranean Sea through time are needed.

*Outlook*

Our study shows that genetic analyses can be used to confirm or reject existing

hypothesis about, for example, stock connectivity but, most importantly, that they can

reveal hidden phenomena that were not foreseen, such as the thus far unknown

hybridization between white and black anglerfish. Yet, despite the power of genetics

to provide fisheries assessment relevant information, there are still barriers for the

uptake of genetic data by fisheries management, which can be due to a variety of

factors (Bernatchez et al., 2017; Ovenden, Berry, Welch, Buckworth, & Dichmont,

2015), such as lack of clear communication of genetic concepts by geneticists to end-

users and reluctance to change or adapt established assessment and management

procedures. The white anglerfish is a highly valuable species (around 30K tons with a

corresponding value of around 142 million euro; data extracted from

https://stecf.jrc.ec.europa.eu/dd/fdi) with a well-established data collection, and

assessment and management frameworks. Thus, we anticipate that these results will

set the basics for an imminent genetics informed fisheries management for this

species. Indeed, a genetics informed fisheries management is essential to ensure the

basics of fisheries science, whereby maximum sustainable yield can only be reached by

efficient management of distinct populations.

**Acknowledgements**

We would like to thank Arkaitz Pedrajas and Arnaitz Mugerza (AZTI), Liz Clarke, Jim

Drewery and Ruadhan Gillespie-Mules (MSS, Scotland), Eoghan Kelly and David Tully

(MI, Ireland), Jose Carlos Fernández Franco, Ángela Cortina Burgueño and Oscar

Fernández Acevedo (OPPF-4, Spain), the fisheries surveys team (CEFAS), Izaskun

Preciado and Susana Ruiz (IEO), Corina Chaves (IPMA, Portugal), Anabel Colmenero

(ICM-CSIC, Spain), Chrysoula Gubili (FRI, Greece) and Nota Peristeraki (HCRM, Greece)

for sampling and Elisabete Bilbao for laboratory work. This work has been funded by

the Joint Research Centre (European Commission), through the project GECKA

(Contract JRC/IPR/2017/D.2/0016/NC), the Department of Environment, Planning,

Agriculture and Fisheries (Basque Government), through the project GENGES and a

predoctoral grant to Imanol Aguirre-Sarabia, and by the Department of Education

(Basque Government), through a predoctoral grant to Iker Pereda. This is contribution

1043 from the Marine Research Division of AZTI.

**Data availability statement**

Demultiplexed and quality-filtered RAD-tag reads are available at the NCBI Sequence

Read Archive (SRA) under accession number SUB9963797 and final datasets and scripts

utilized for this project are available at https://github.com/rodriguez-

ezpeleta/Lophius_PopGen

**Author Contributions**

NRE, JTM, AZ, MS, IQ and IC designed the study. HG, FB, IH and JL provided samples.

IM generated data. IAS, NDA, IPA and NRE analyzed data. IAS, NRE, AU, HDG, FB, IH, JL,

IQ interpreted data. IAS and NRE wrote the first draft of the manuscript and integrated

input from all authors.

**References**

Alexander, D. H., Shringarpure, S. S., Novembre, J., & Lange, K. (2015). Admixture 1.3 software
manual. *Los Angeles: UCLA Human Genetics Software Distribution*.
Allendorf, F., Leary, R. F., Spruell, P., & Wenburg, J. (2001). The problems with hybrids: setting
conservation guidelines. *Trends in Ecology and Evolution, 16*, 613-622.
Anderson, E., & Thompson, E. (2002). A model-based method for identifying species hybrids
using multilocus genetic data. *Genetics, 160*(3), 1217-1229.
Anderson, T. M., Candille, S. I., Musiani, M., Greco, C., Stahler, D. R., Smith, D. W., . . .
Bustamante, C. D. (2009). Molecular and evolutionary history of melanism in North
American gray wolves. *Science, 323*(5919), 1339-1343.
Armani, A., Castigliego, L., Tinacci, L., Gandini, G., Gianfaldoni, D., & Guidi, A. (2012). A rapid
PCR–RFLP method for the identification of Lophius species. *European Food Research*
*and Technology, 235*(2), 253-263. doi:10.1007/s00217-012-1754-3
Arnold, M. L. (1997). *Natural hybridization and evolution*: Oxford University Press on Demand.
Azevedo, M. (1995). A statistical analysis of black monkfish catch rates in ICES Division IXa.
*Fisheries Research, 24*(4), 281-289.
Bernatchez, L., Wellenreuther, M., Araneda, C., Ashton, D. T., Barth, J. M. I., Beacham, T. D., . . .
Withler, R. E. (2017). Harnessing the Power of Genomics to Secure the Future of
Seafood. *Trends in Ecology & Evolution, 32*(9), 665-680.
doi:https://doi.org/10.1016/j.tree.2017.06.010
Blanco, G., Borrell, Y., Cagigas, M., Vázquez, E., & Prado, J. S. (2008). Microsatellites-based
genetic analysis of the Lophiidae fish in Europe. *Marine and Freshwater Research,*
*59*(10), 865-875.
Bonfil, R. (2005). Management Techniques for Elasmobranch Fisheries. In R. B. John Musick
(Ed.), *FAO Fisheries Technical Paper 474* (pp. 6-14): FAO.
Budd, A. F., & Pandolfi, J. M. (2010). Evolutionary novelty is concentrated at the edge of coral
species distributions. *Science, 328*(5985), 1558-1561.
Cadrin, S. X. (2020). Defining spatial structure for fishery stock assessment. *Fisheries Research,*
*221*, 105397. doi:https://doi.org/10.1016/j.fishres.2019.105397
Cañás, L., Stransky, C., Schlickeisen, J., Sampedro, M. P., & Fariña, A. C. (2012). Use of the
otolith shape analysis in stock identification of anglerfish (Lophius piscatorius) in the
Northeast Atlantic. *ICES Journal of Marine Science, 69*(2), 250-256.
doi:10.1093/icesjms/fss006
Caruso, J. (1986). Lophiidae. *Fishes of the North-eastern Atlantic and the Mediterranean, 3*,
1362-1363.
Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an
analysis tool set for population genomics. *Molecular Ecology, 22*(11), 3124-3140.
Crozier, W. (1987). Biochemical genetic variation and population structure in angler-fish
Lophius piscatorius L. from the Irish Sea and west of Scotland. *Journal of Experimental*
*Marine Biology and Ecology, 106*(2), 125-136.
Charrier, G., Chenel, T., Durand, J. D., Girard, M., Quiniou, L., & Laroche, J. (2006).
Discrepancies in phylogeographical patterns of two European anglerfishes (Lophius
budegassa and Lophius piscatorius). *Molecular Phylogenetics and Evolution, 38*(3),
742-754. doi:10.1016/j.ympev.2005.08.002
Duarte, R., Azevedo, M., Landa, J., & Pereda, P. (2001). Reproduction of anglerfish (Lophius
budegassa Spinola and Lophius piscatorius Linnaeus) from the Atlantic Iberian coast.
*Fisheries Research, 51*(2), 349-361. doi:https://doi.org/10.1016/S0165-7836(01)00259-
4
Dubin, A., Jørgensen, T. E., Moum, T., Johansen, S. D., & Jakt, L. M. (2019). Complete loss of the
MHC II pathway in an anglerfish, Lophius piscatorius. *Biology Letters, 15*(10),
20190594.

Epifanio, J., & Nielsen, J. (2000). The Role of Hybridization in the Distribution, Conservation and
Management of Aquatic Species. *Reviews in Fish Biology and Fisheries, 10*(3), 245-251.
doi:10.1023/a:1016729132297
Etter, P. D., Bassham, S., Hohenlohe, P. A., Johnson, E. A., & Cresko, W. A. (2012). SNP
discovery and genotyping for evolutionary genetics using RAD sequencing. In
*Molecular methods for evolutionary genetics* (pp. 157-178): Springer.
García-Rodríguez, M., Pereda, P., Landa, J., & Esteban, A. (2005). On the biology and growth of
the anglerfish Lophius budegassa Spinola, 1807 in the Spanish Mediterranean: a
preliminary approach. *Fisheries Research, 71*(2), 197-208.
Garcia-Vazquez, E., Machado-Schiaffino, G., Campo, D., & Juanes, F. (2012). Species
misidentification in mixed hake fisheries may lead to overexploitation and population
bottlenecks.        *Fisheries        Research,        114,        52-55.*
doi:https://doi.org/10.1016/j.fishres.2011.05.012
Hayden, B., Pulcini, D., Kelly-Quinn, M., O'Grady, M., Caffrey, J., McGrath, A., & Mariani, S.
(2010). Hybridisation between two cyprinid fishes in a novel habitat: genetics,
morphology and life-history traits. *BMC Evolutionary Biology, 10*(1), 169.
Hislop, J. R., Gallego, A., Heath, M. R., Kennedy, F. M., Reeves, S. A., & Wright, P. J. (2001). A
synthesis of the early life history of the anglerfish, Lophius piscatorius (Linnaeus, 1758)
in northern British waters. *ICES Journal of Marine Science, 58*(1), 70-86.
ICES. (2007). *Report of the Workshop on Sexual Maturity Staging of Hake and Monk*
*(WKMSHM), 21–24 November 2007, Lisbon, Portugal. ICES CM 2007/ACFM:34.*
Retrieved from
ICES. (2010). *Report of the Working Group on the Assessment of Southern Shelf Stocks of Hake,*
*Monk and Megrim (WGHMM), 5 - 11 May 2010, Bilbao, Spain, ICES Document CM*
*2010/ACOM:11.* Retrieved from
ICES. (2018). *Report of the Benchmark Workshop on Anglerfish Stocks in the ICES Area*
*(WKANGLER), 12–16 February 2018, Copenhagen, Denmark. ICES CM 2018/ACOM:31.*
*177 pp.* Retrieved from
ICES. (2019a). *Working Group for the Bay of Biscay and the Iberian waters Ecoregion (WGBIE).*
*ICES Scientific Reports. 1:31. 692 pp.* http://doi.org/10.17895/ices.pub.5299. Retrieved
from http://doi.org/10.17895/ices.pub.5299
ICES. (2019b). Working Group for the Celtic Seas Ecoregion (WGCSE). ICES Scientific Reports.
1:29.        1078        pp.        http://doi.org/10.17895/ices.pub.4982.
doi:http://doi.org/10.17895/ices.pub.4982
Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide
SNP data. *Bioinformatics, 27*(21), 3070-3071. doi:10.1093/bioinformatics/btr521
Landa, J., & Antolínez, A. (2018). Weight–length relationships, weight conversion factors and
somatic indices from two stocks of black anglerfish (Lophius budegassa) and white
anglerfish (L. piscatorius) in north-eastern Atlantic waters. *Regional Studies in Marine*
*Science, 23,* 87-96. doi:https://doi.org/10.1016/j.rsma.2017.03.002
Landa, J., Quincoces, I., Duarte, R., Fariña, A. C., & Dupouy, H. (2008). Movements of black and
white anglerfish (Lophius budegassa and L. piscatorius) in the northeast Atlantic.
*Fisheries Research, 94*(1), 1-12. doi:https://doi.org/10.1016/j.fishres.2008.04.006
Laurenson, C. H., Johnson, A., & Priede, I. G. (2005). Movements and growth of monkfish
Lophius piscatorius tagged at the Shetland Islands, northeastern Atlantic. *Fisheries*
*Research, 71*(2), 185-195. doi:https://doi.org/10.1016/j.fishres.2004.08.020
Leone, A., Álvarez, P., García, D., Saborido-Rey, F., & Rodriguez-Ezpeleta, N. (2019). Genome-
wide SNP based population structure in European hake reveals the need for
harmonizing biological and management units. *ICES Journal of Marine Science.*
Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
*arXiv preprint arXiv:1303.3997.*

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Subgroup, G. P. D. P.
527           (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics, 25*(16),
528           2078-2079. doi:10.1093/bioinformatics/btp352

529    Maltby, K. M., Rutterford, L. A., Tinker, J., Genner, M. J., & Simpson, S. D. (2020). Projected
530           impacts of warming seas on commercially fished species at a biogeographic boundary
531           of the European continental shelf. *Journal of Applied Ecology, n/a*(n/a).
532           doi:10.1111/1365-2664.13724

533    Mallet, J. (2007). Hybrid speciation. *Nature, 446*(7133), 279-283. doi:10.1038/nature05706

534    Marko, P. B., Lee, S. C., Rice, A. M., Gramling, J. M., Fitzhenry, T. M., McAlister, J. S., . . . Moran,
535           A. L. (2004). Fisheries: mislabelling of a depleted reef fish. *Nature, 430*(6997), 309-310.
536           doi:10.1038/430309b

537    Methot, R. D., & Wetzel, C. R. (2013). Stock synthesis: A biological and statistical framework for
538           fish stock assessment and fishery management. *Fisheries Research, 142*, 86-99.
539           doi:https://doi.org/10.1016/j.fishres.2012.10.012

540    Mirimin, L., Kerwath, S., Macey, B., Bester-van der Merwe, A., Lamberth, S., Bloomer, P., &
541           Roodt-Wilding, R. (2014). Identification of naturally occurring hybrids between two
542           overexploited sciaenid species along the South African coast. *Molecular Phylogenetics
543           and Evolution, 76*, 30-33.

544    Montanari, S. R., Hobbs, J. P. A., Pratchett, M. S., Bay, L. K., & Van Herwerden, L. (2014). Does
545           genetic distance between parental species influence outcomes of hybridization among
546           coral reef butterflyfishes? *Molecular Ecology, 23*(11), 2757-2770.

547    Morgan, J. A. T., Harry, A. V., Welch, D. J., Street, R., White, J., Geraghty, P. T., . . . Ovenden, J.
548           R. (2012). Detection of interspecies hybridisation in Chondrichthyes: hybrids and
549           hybrid offspring between Australian (Carcharhinus tilstoni) and common (C. limbatus)
550           blacktip shark found in an Australian fishery. *Conservation Genetics, 13*(2), 455-463.
551           doi:10.1007/s10592-011-0298-6

552    Moritz, C., Dowling, T., & Brown, W. (1987). Evolution of animal mitochondrial DNA: relevance
553           for population biology and systematics. *Annual Review of Ecology and Systematics,
554           18*(1), 269-292.

555    Ofstad, L., Angus, C., Pedersen, T., & Steingrund, P. (2013). Age and growth of anglerfish
556           (Lophius piscatorius) in Faroese waters. *Fisheries Research, 139*, 51-60.

557    Ovenden, J. R., Berry, O., Welch, D. J., Buckworth, R. C., & Dichmont, C. M. (2015). Ocean's
558           eleven: a critical evaluation of the role of population, evolutionary and molecular
559           genetics in the management of wild fisheries. *Fish and Fisheries, 16*(1), 125-159.
560           doi:10.1111/faf.12052

561    Pampoulie, C., Gíslason, D., Ólafsdóttir, G., Chosson, V., Halldórsson, S. D., Mariani, S., . . .
562           Víkingsson, G. A. (2020). Evidence of unidirectional hybridization and second-
563           generation adult hybrid between the two largest animals on Earth, the fin and blue
564           whales. *Evolutionary Applications, n/a*(n/a). doi:https://doi.org/10.1111/eva.13091

565    Purcell, S. (2009). PLINK 1.07. In.

566    Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A., Bender, D., . . . Daly, M. J.
567           (2007). PLINK: a tool set for whole-genome association and population-based linkage
568           analyses. *The American Journal of Human Genetics, 81*(3), 559-575.

569    Quincoces, I. (2002.). *Crecimiento y reproducción de las especies" Lophius budegassa" Spinola
570           1807 y" Lophius piscatorius" Linneo 1758, del Golfo de Vizcaya.* (PhD), Departamento
571           de Zoología y Dinámica Celular Animal, Universidad del País Vasco (EHU/UPV), Spain.

572    Quincoces, I., Santurtún, M., & Lucio, P. (1998). Biological aspects of white anglerfish (Lophius
573           piscatorius) in the Bay of Biscay (ICES Division VIIIa, b, d) in 1996–1997. *ICES CM.*

574    Relini, G. (1999). Synthesis of the knowledge on bottom fishery resources in Central
575           Mediterranean (Italy and Corsica). *Biol Mar Medit, 6*(1), 276-291.

576    Rodríguez-Ezpeleta, N., Bradbury, I. R., Mendibil, I., Álvarez, P., Cotano, U., & Irigoien, X.
577           (2016). Population structure of Atlantic mackerel inferred from RAD-seq-derived SNP

markers: effects of sequence clustering parameters and hierarchical SNP selection. *Molecular Ecology Resources, 16*(4), 991-1001. doi:10.1111/1755-0998.12518

Rodríguez-Ezpeleta, N., Díaz-Arce, N., Walter Iii, J. F., Richardson, D. E., Rooker, J. R., Nøttestad, L., . . . Arrizabalaga, H. (2019). Determining natal origin for improved management of Atlantic bluefin tuna. *Frontiers in Ecology and the Environment, 17*(8), 439-444. doi:10.1002/fee.2090

Rousset, F. (2014). Genepop 4.3 for Windows/Linux/Mac OS X. *This documentation: July, 8*, 2014.

Schwenke, P. (2013). *History and extent of introgressive hybridization in Puget Sound rockfishes (Sebastes auriculatus, S. caurinus, and S. maliger).*

Seehausen, O. (2006). Conservation: losing biodiversity by reverse speciation. *Current Biology, 16*(9), R334-R337.

Sevilla, R. G., Diez, A., NorÉN, M., Mouchel, O., JÉRÔMe, M., Verrez-Bagnis, V., . . . Bautista, J. M. (2007). Primers and polymerase chain reaction conditions for DNA barcoding teleost fish based on the mitochondrial cytochrome b and nuclear rhodopsin genes. *Molecular Ecology Notes, 7*(5), 730-734. doi:10.1111/j.1471-8286.2007.01863.x

Stephenson, R. L. (2002). Stock structure and management structure: an ongoing challenge for ICES. *ICES marine science symposia, 215*, 305-314.

Stronen, A. V., & Paquet, P. C. (2013). Perspectives on the conservation of wild hybrids. *Biological Conservation, 167*, 390-395.

Team, R. C. (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. 2012. *URL http://www. R-project. org.*

Ungaro, N., Marano, G., Auteri, R., Voliani, A., Massutí, E., García-Rodríguez, M., & Osmani, K. (2002). Distribution, abundance and biological features of anglerfish (Lophius piscatoirus and Lophhius budegassa)(Osteichthyes: Lphiiformes) in the Mediterranean Sea. *Scientia Marina, 66*(S2), 55-63.

Verheyen, E., Salzburger, W., Snoeks, J., & Meyer, A. (2003). Origin of the superflock of cichlid fishes from Lake Victoria, East Africa. *Science, 300*(5617), 325-329.

Yaakub, S. M., Bellwood, D. R., van Herwerden, L., & Walsh, F. M. (2006). Hybridization in coral reef fishes: introgression and bi-directional gene exchange in Thalassoma (family Labridae). *Molecular Phylogenetics and Evolution, 40*(1), 84-100.

**Figure legends**

Figure 1. Samples included in this study, and in parentheses those genotyped with RAD-seq,
where black triangles indicate Lophius budegassa samples and coloured circles, Mediterranean
(red), Atlantic southern stock (purple), Atlantic northern stock (green) or Atlantic northern
shelf stock (blue) Lophius piscatorius samples.

Figure 2. A. Specimen of anglerfish captured in the northern coast of Scotland (65 cm and 4.4
kg) with white peritoneum, emerald green eyes, and stippled skin, genetically identified as
black anglerfish. B. Agarose gel electrophoresis of the resulting PCR-RFLP band for white (L.
pis) and black anglerfish (L. bud) showing that this specimen (Sample) is genetically identified
as black anglerfish.

Figure 3. A. Principal Component Analysis (PC1=80.82%; PC2=0.26%) where main
differentiated groups are circled. B. Individual ancestry proportions estimated by ADMIXTURE
when assuming two ancestral populations. C. For each group identified int the PCA, whether
individuals composing it are assigned to L. piscatorius (white fish or mitochondrion), L.
budegassa (black fish or mitochondrion) or both according to visual or mitochondrial DNA
based species identification, admixture pattern and assignment according to NewHybrids.

Figure 4. Proportion of hybrid (grey areas and numbers) and *L. budegassa* (black areas and
numbers) individuals among those morphologically identified as *L. piscatorius* for each stock
(large circles) and, within stocks, for smaller areas (small circles indicating number of sampled
individuals per area). Colours refer to stocks as in Figure 1.

Figure 5. A. Principal Component Analysis (PCA) of samples genetically identified as white
anglerfish when including (left) or not (right) Mediterranean samples. B. Individual ancestry
proportions of samples genetically identified as white anglerfish estimated by ADMIXTURE
when assuming two ancestral populations including (above) or not (below) Mediterranean
samples.

## A

|  | Gr1 | Gr2 | Gr3 |
|---|---|---|---|
| N. Plat | 82 | 2 | 0 |
| North | 144 | 25 | 4 |
| South | 20 | 0 | 2 |
| Med | 6 | 1 | 5 |
| Black | 0 | 0 | 27 |

## B

N. Shelf  North  South  Med  Black

## C

|  | Gr1 | Gr2 |  | Gr3 |
|---|---|---|---|---|
| Visual ID. | | | | |
| Mt DNA ID | | | | |
| ADMIXTURE | | | | |
| NewHybrids | *L. piscatorius* | Backcross L.pis | F1 hybrid | Backcross L.bud | *L. budegassa* |

A

PC1=1.04%, PC2=0.62%

PC1=0.64%, PC2=0.62%

B

Med

N. Shelf          North          South