Spinal premotor interneurons controlling antagonistic muscles are spatially intermingled

Remi Ronzano[*1], Sophie Skarlatou[*2], Bianca K. Barriga[*6,7], B. Anne Bannatyne[*3], Gardave S. Bhumbra[*4], Joshua D. Foster[4], Jeffrey D. Moore[8], Camille Lancelin[1], Amanda Pocratsky[1], Mustafa Görkem Özyurt[1], Calvin C. Smith[1], Andrew J. Todd[3], David J. Maxwell[3], Andrew J. Murray[5], Samuel L. Pfaff[6§], Robert M. Brownstone[1§], Niccolò Zampieri[2§], Marco Beato[3§]

[1] Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, University College London, London WC1N 3BG, UK

[2] Max Delbrück Center for Molecular Medicine (MDC), Roybert-Rössle-Str. 10, 13092 Berlin, Germany

[3] Institute of Neuroscience and Psychology, College of Medical, Veterinary and Life Sciences, University of Glasgow, West Medical Building, Glasgow G12 8QQ, UK

[4] Department of Neuroscience Physiology and Pharmacology (NPP), Gower Street, University College London, WC1E 6BT, UK

[5] Sainsbury Wellcome Centre for Neural Circuits and Behaviour, University College London, London W1T 4JG, UK

[6] The Salk Institute for Biological Studies, Gene Expression Laboratory, 10010 North Torrey Pines Road, La Jolla CA 92037, USA

[7] Biological Sciences Graduate Program, University of California, San Diego, 9500 Gilman Drive, La Jolla, CA 92037, USA.

[8] Howard Hughes Medical Institute and Department of Molecular and Cellular Biology, Center for Brain Science, Harvard University, Cambridge, MA 02138, USA

*equal contribution

§co-senior authors

Corresponding authors:
Niccolò Zampieri, Max Delbrück Center for Molecular Medicine (MDC), Robert-Rössle-Str. 10, 13092 Berlin, Germany
niccolo.zampieri@mdc-berlin.de

Samuel J. Pfaff, The Salk Institute for Biological Studies, Gene Expression Laboratory, 10010 North Torrey Pines Road, La Jolla CA 92037, USA
pfaff@salk.edu

Robert M. Brownstone, Department of Neuromuscular Diseases, UCL Queen Square Institute of Neurology, University College London, London WC1N 3BG, UK
r.brownstone@ucl.ac.uk

Marco Beato, Department of Neuroscience Physiology and Pharmacology (NPP), Gower Street, University College London, WC1E 6BT, UK
m.beato@ucl.ac.uk

Key words: spinal cord, premotor interneurons, flexor muscles, extensor muscles, rabies, viral tracing

1 Abstract

Elaborate behaviours are produced by tightly controlled flexor-extensor motor neuron activation patterns.
Motor neurons are regulated by a network of interneurons within the spinal cord, but the computational
processes involved in motor control are not fully understood. The neuroanatomical arrangement of motor
and premotor neurons into topographic patterns related to their controlled muscles is thought to facilitate
how information is processed by spinal circuits. Rabies retrograde monosynaptic tracing has been used
to label premotor interneurons innervating specific motor neuron pools, with previous studies reporting
topographic mediolateral positional biases in flexor and extensor premotor interneurons. To further
investigate the spatial organization of premotor interneurons, we used several different viral approaches
for retrograde transsynaptic tracing. In contradiction to previous reports, we consistently found that
premotor interneurons are not organized into topographic groups related to their innervation of flexor or
extensor motor neuron pools. We conclude that the premotor spinal neurons controlling different muscles
are spatially intermingled and operate in the absence of clear spatial patterns among the flexor-extensor
circuit components.

Introduction

Precise regulation in the timing and pattern of activation of muscle groups across a joint is at the basis of motor control. In limbed vertebrates, the activity of flexor and extensor muscles is directed by dedicated pools of motor neurons that receive inputs from different subtypes of excitatory and inhibitory interneurons. Several of these classes of interneurons have been described in electrophysiological, anatomical, and genetic studies (Hultborn et al., 1971; Jankowska, 2001; Goulding, 2009), however the incomplete knowledge of the composition of spinal circuitry that control the activity of flexor and extensor motor neurons limits progress towards a full understanding of motor circuits.

Viral trans-synaptic tracing techniques have been used for several decades to map motor circuits (Ugolini, 2020). Given that rabies virus (RabV) jumps across synapses in the retrograde direction to infect presynaptic neurons, its use via intramuscular injections led to the identification of polysynaptic pathways (Rathelot and Strick, 2009). But the identification of monosynaptic connectivity relied on the timing of transsynaptic jumps, thus leading to a degree of uncertainty about the number of synapses between a labelled neuron and the motoneurons innervating the injected muscle.

The introduction of RabV monosynaptic tracing provided a high-throughput method for mapping presynaptic connectivity of selected neuronal populations (Callaway and Luo, 2015). Monosynaptic restriction is achieved by using a mutant virus lacking the gene encoding the rabies glycoprotein (G; ΔG-RabV), which is necessary for transsynaptic transfer, combined with selective complementation of G expression in neurons of choice (Wickersham et al., 2007). Various methodologies have been used to restrict G expression to the target neuronal population, with these G-expressing neurons becoming "starter cells" from which infecting ΔG-RabV can jump only one synapse and selectively label presynaptic neurons (Wall et al., 2010; Callaway and Luo, 2015).

Shortly after its introduction, monosynaptic rabies tracing was applied to the study of premotor interneurons in the spinal cord. To obtain selective complementation of G and subsequent rabies monosynaptic transfer from a single motor neuron pool, an elegant approach based on intramuscular co-injection of an AAV expressing G (AAV-G) and ΔG-RabV, both of which can infect motor neurons retrogradely, was described (Stepien et al., 2010). Thus, starter cells are generated in one fell swoop by taking advantage of the stringent anatomical specificity of motor neuron to muscle connectivity (Figure 1A). When this method was applied to study the distribution of premotor interneurons controlling the activity of extensor and flexor muscles in the hindlimb, a prominent spatial segregation along the medio-lateral axis of the dorsal ipsilateral spinal cord was observed, with extensor premotor interneurons found in more medial positions than flexors (Tripodi et al., 2011). The authors proposed that this organisation led to

"private" disynaptic pathways from proprioceptive afferents to appropriate motoneurons, and that this might offer some circuit organisational advantages.

More recently, in order to address concerns that this method could also lead to rabies infection and transsynaptic transfer from the sensory route (Figure 1A; Zampieri et al., 2014), G expression was further restricted to motor neurons by combining the use of a mouse line expressing Cre recombinase under the control of choline acetyltransferase (ChAT) and intramuscular injection of an AAV driving expression of G in a conditional manner (AAV-flex-G; Figure 1B). Under these conditions, segregation of flexor and extensor premotor interneurons was also shown at forelimb level (Wang et al., 2017). Finally, a further modification to the original AAV complementation strategy was introduced: stereotactic injection of AAV-flex-G in the spinal cord of *ChAT::Cre* mice was used to target G expression to cholinergic neurons, with restriction of starter cells to a motor pool achieved by ΔG-RabV muscle injection. These experiments also showed medio-lateral segregation in the distribution of flexor and extensor premotor interneurons (Figure 1C; Takeoka and Arber, 2019). Thus, these different experiments that all used AAV for complementing G expression in motor neurons demonstrated similar segregation of extensor and flexor premotor interneurons.

Given the importance that neuronal position may play in circuit organization and function, we sought to identify premotor interneurons for further investigation. However, we elected to achieve G complementation by using a mouse genetic approach that takes advantage of a conditional mouse line that drives G expression under control of Cre recombinase (*RΦGT* mice; Figure 1D, 1E; Takatoh et al., 2013). This method has been previously used to trace premotor circuitry of the vibrissal and orofacial motor systems, in combination with *ChAT::Cre* mice (Takatoh et al., 2013; Stanek et al., 2014), as well as forelimb muscles in combination with *Olig2::Cre* mice (Skarlatou et al., 2020). We reasoned that using this approach, G should be available at high levels in all motor neurons, thereby leading to efficient monosynaptic transfer from all the cells infected by ΔG-RabV, which was supplied via intramuscular injection. In contrast with previous studies, we did not observe any difference in the distribution of flexor and extensor premotor interneurons. Thus, we decided to replicate the experiments using AAV complementation strategies: AAV-G injection in wild-type mice (Tripodi et al., 2011) and AAV-flex-G in *ChAT::Cre* mice (Wang et al., 2017). Surprisingly, we did not observe segregation in the spatial organization of flexor and extensor premotor interneurons. Finally, we resorted to a different viral tracing method and used timed infection with pseudorabies virus (PRV; Strack and Loewy, 1990; Jovanovic et al., 2010), which also resulted in overlapping distributions of flexor and extensor premotor neurons. Altogether, these experiments conducted in different laboratories and using most of the available methods described in the literature for viral transsynaptic tracing of premotor circuits do not show segregation of flexor and extensor premotor neuron distributions.

Results

Flexor and extensor premotor interneurons in ChAT::Cre +/-; RΦGT+/- mice

In order to determine the spatial distribution of premotor interneurons controlling flexion and extension of the ankle, we injected ΔG-RabV/mCherry and ΔG-RabV/eGFP in the TA (tibialis anterior; ankle flexor) and LG (lateral gastrocnemius; ankle extensor) muscles of postnatal day (P) 1-2 *ChAT::Cre +/-; RΦGT+/-* mice. Analysis of lumbar level (L) 2 and L5 sections 8-9 days after injection revealed two main clusters of premotor interneurons located in the dorsal ipsilateral and ventral contralateral spinal cord (Figure 2A and 2B). Next, we obtained coordinates for the labelled cells in each section of the lumbar spinal cord and mapped premotor interneuron positions in three dimensions. The projection of x-y coordinates along the rostro-caudal axis of the spinal cord showed no difference in medio-lateral and dorso-ventral positions of flexor and extensor premotor interneurons (Figure 2C, left panel). Convolved distributions fully overlapped for the two groups in all four quadrants. Similarly, projection along the coronal plane (Figure 2C, middle panel) or sagittal plane (Figure 2C, right panel) revealed no obvious differences in the rostro-caudal, dorso-ventral and medio-lateral distribution of flexor and extensor premotor interneurons.

To study in detail the positional organization of premotor interneurons controlling the activity of the ankle joint, we analysed 13 animals in which we had performed simultaneous ΔG-RabV-eGFP and ΔG-RabV-mCherry injections in 3 different pairs of antagonist and synergist muscles in P1-2 *ChAT::Cre +/-; RΦGT+/-* mice: TA and LG, LG and MG (medial gastrocnemius; ankle extensor and LG synergist) and TA and PL (peroneus longus; ankle flexor and TA synergist). We started by analysing the results from the antagonist pairs, LG (n=11) and TA (n=7), pooling experiments from single and double injections (Figures 3A-B). All LG and TA experiments are overlaid, with different shades of blue (LG) and orange (TA) representing different animals, showing the reproducibility of premotor interneurons distributions across single experiments (pooled distributions shown in Figure 3C, individual experiments are shown in Figure 12-figure supplements 1-7). The Hedges' G coefficients of the distributions in the ipsilateral dorsal quadrants for all pairs of experiments had a median of -0.06 (IQR -0.26, 0.14), showing homogeneity between experiments and no differences in the positional organization of flexor and extensor premotor interneurons. The reproducibility of the results is confirmed by analysis of the coordinates across all experiments, showing similar correlation values within or across muscles (Figure 3D; r≥0.78). The values of the medians of individual experiments for LG and TA injections were 321 and 306 μm respectively (Hedges' G=-0.06, Figures 3E-F).

Since it was previously reported that the medio-lateral segregation in the distribution of flexor and extensor premotor interneurons is more pronounced in spinal segments rostral to the infected motor nucleus

(Tripodi et al., 2011), we analysed the organization of premotor interneurons at different lumbar levels. Positional coordinates were pooled and divided into 800 μm rostro-caudal bins and distributions were plotted for each bin from L1 to L6 (Figure 3–figure supplement 1). No differences in the medio-lateral distributions of LG and TA premotor interneurons were observed in any segment analysed (median positions on the medio-lateral axis for L1, the segment with the largest visible medio-lateral segregation: LG = 309 μm and TA = 327 μm, hierarchical bootstrapped Hedges'G = -0.17 (IQR -0.12, -0.22)).

The identity of infected motor neurons

The identity of starter cells represents a critical element for the interpretation of rabies tracing experiments. For the rabies tracing approaches discussed here (Figure 1), it is difficult to determine unambiguously the identity of starter motor neurons since the complementation of G is not accompanied by the expression of a reporter gene. In addition, a precise estimation of the total number of infected motor neurons is complicated by rabies toxicity that kills many neurons shortly after infection (Reardon et al., 2016). Nevertheless, we took advantage of the topographic organization of motor neuron to muscle connectivity to evaluate the pool identity and number of infected motor neurons that survived until the end of the experiment (Romanes, 1964; McHanwell and Biscoe, 1981). As predicted by the known position of the TA and LG motor pools in the spinal cord, we found that the majority of infected motor neurons were localized in the dorsal part of the ventral horn (Figure 2C; Sürmeli et al., 2011). Surprisingly, we also found motor neurons in more ventral, "ectopic" positions (Figure 2C and supplement 1-7 to Figure 12), where pools that innervate muscles controlling the function of the knee and hip joints reside (Sürmeli et al., 2011). Motor neuron labelling occasionally extended outside the lower lumbar segments where most of the ankle flexors and extensor pools are located (Figure 12-figure supplement 1-7). Moreover, in double TA and LG injections we found instances of motor neurons infected with both viruses (Figures 2D, E). In 5 experiments, a total of 200 LG and 150 TA motor neurons were labelled, of which 13 were infected with both ΔG-RabV-eGFP and ΔG-RabV-mCherry (see Table 1).

The presence of ectopic and double labelled motor neurons could be explained by either secondary labelling due to rabies transsynaptic transfer from starter cells through recurrent connections onto other motor neurons or by unintended primary infection of motor neurons due to non-specific muscle injections. Careful post-hoc analysis of hindlimb muscles after ΔG-RabV injection did not reveal any evidence of non-specific muscle infection. In addition, it is unlikely that the relatively small volume of ΔG-RabV solution used in the experiments would lead to spread to other muscles, particularly those located in compartments on the other side of the bone or joint, where muscle targets of the ectopic motor neurons reside. We therefore suggest that ectopic motor neurons were mostly labelled by transsynaptic labelling and therefore represent second-order presynaptic neurons. This is not surprising, as motor neurons have been shown to form synapses with other motor neurons and their connections can extend to neighbouring spinal

segments (Bhumbra and Beato, 2018), suggesting that the ectopic motor neurons found in our experiments were most likely due to rabies transsynaptic transfer. Regardless of the underlying reasons for the observed ectopic motor neuron labelling, its presence raises the possibility that what we defined as flexor and extensor premotor networks, might, in fact, originate from a mixed population of starter cells containing not only motor neurons of a single pool identity but also a fraction of, "non-specific", motor neurons belonging to other pools, thereby potentially diluting any observable spatial difference between the premotor networks of flexor and extensor muscles. However, it is important to notice that in our experiments the number of presumed "non-specific" starter cells is low and therefore unlikely to confound the results (Table 1; see below and Discussion).

The number of infected motor neurons does not affect the distribution of premotor interneurons

Spinal or muscle injection of AAV to complement G expression is likely to result in infection of a subset of motor neurons within the targeted pool, whereas the genetic experiments will result in complementation in all motor neurons. Thus, it is possible that the difference in the results obtained with these two methods could be explained by the absolute number of motor neurons from which rabies synaptic transfer occurs. In order to test the effect of the number of starter cells in our experimental conditions, we reasoned that by reducing the viral titre of the rabies solution used for muscle injection, we would scale down the number of infected motor neurons. Therefore, we performed a series of muscle injections (7 LG and 6 TA, of which 3 double LG-TA) in *ChAT::Cre +/-; RΦGT+/-* mice (Figure 1D) as in the experiments described above, but with diluted rabies virus (titre $<10^9$ I.U./ml) to reduce its infection efficiency (see Table 1). In low titre experiments, we detected an average 4.7 infected motor neurons compared to an average of 35.2 in the high titre experiments (titre $>5\times10^9$ I.U./ml). Once again, we did not observe segregation in the medio-lateral distribution of LG and TA premotor interneurons (Figures 4A-C). The pairwise Hedges' G coefficients for all the experiments had median value of -0.05 (IQR -0.19,0.15, Figure 5D). While there was a higher degree of variability between experiments compared to high titre injections, as shown in the correlation matrix of individual experiments (Figure 4D, r>0.45 for all comparisons), the median value of the medio-lateral positions in each experiment were very similar (LG = 307 µm and TA = 317 µm, hierarchical bootstrapped Hedges' G=-0.02, IQR -0.09,0.03, Figures 4E, F, individual experiments shown in Figure 12-figure supplement 8-10). Next, we compared high and low titre experiments for each muscle injected (Figure 4-figure supplement 1). The distribution of premotor interneurons shows remarkable overlap for LG and TA injections (Figure 4-figure supplement 1A, E). The medio-lateral distributions were not different for LG (median Hedges' G = -0.05, IQR -0.21,0.11, with high correlation values across experiments, between 0.77 and 0.98, Figure 4-figure supplement 1B, and medians of 321 µm and 307 µm for high and low titre experiments respectively, Figure 4-figure supplement 1C-D). When comparing the high and low titre injections of TA, we found high correlation values across experiments (across experiment

median 0.83, IQR 0.74,0.90) and low Hedges' G (median -0.10, IQR -0.26, 0.13, Figure 4-figure supplement 1F) and no differences in the lateral direction for low titre injections into TA (medians: high = 306 µm and low = 317 µm, Figure 4-figure supplement 1G-H). Of note, in low titre experiments we never observed ectopic motor neurons outside the expected nucleus (individual experiments are shown in Figure 12- figure supplements 8-10). Together, these data indicate that neither the absolute number of starter motor neurons nor the infection of ectopic motor neurons observed in high titre experiments significantly affects the positional organization of premotor interneurons.

The distribution of premotor interneurons is similar across different pairs of ankle flexors or extensors

Since it has been proposed that medio-lateral segregation of premotor interneurons is a general feature of flexor and extensor muscles, we analysed premotor interneurons of two more muscles controlling the movement of the ankle joint, PL and MG using the same viral strategy employed for the LG-TA injections (Figure 1D). The distributions of premotor interneurons of LG (6 LG-MG injections, 8 LG-TA injections and 4 single LG injections) and MG (6 LG-MG injections and 4 MG injections) did not reveal any difference in spatial organization (Figure 5A, individual experiments are shown in Figure 12- figure supplements 3,4,6). The Hedges' G for the mediolateral positions in the ipsilateral dorsal quadrants were computed for each pair of experiment and had a median value of 0.01 (IQR -0.12,0.16). Throughout the cord, the positions of rabies-labelled neurons were highly correlated (Figure 4B, r≥0.74) and reproducible along the medio-lateral axis (Figure 5C), with median values of the medio-lateral position across experiments of 314 µm for LG and 315 µm for MG, (bootstrapped median Hedges' G= -0.06, IQR -0.03, -0.10, Figure 4D). The same result was observed for TA and PL premotor interneurons (Figure 5E; 2 TA-PL injections, 8 TA-LG injections, 3 TA single injections and 3 PL single injections, individual experiments are shown in Figure 12- figure supplements 5,7). The median of the Hedges' G coefficients for all the pairs of experiments was- 0.01 (IQR -0.20, 0.18), with high correlation values between experiments (Figure 5F; r≥0.66). The medians of the mediolateral pooled distributions were 308 µm for TA and 327 µm for PL, similar medio-lateral distributions (Figure 5G) and median values were observed (Figure 5H). Hierarchical bootstrap of the data from TA and PL injections resulted in a median Hedges' G of -0.09 (IQR -0.15, -0.03). Together, these data show that premotor interneuron maps obtained using ΔG-RabV muscle injection in *ChAT::Cre +/-; RΦGT+/-* mice (Figure 1D) do not reveal any difference in the positional organization of interneurons controlling the activity of any of the main flexor and extensor muscles of the ankle.

Spatial distribution of excitatory and inhibitory premotor interneurons

Next, we examined whether there are differences in the spatial organization of excitatory vs inhibitory premotor interneurons. We performed single LG or TA injections of ΔG-RabV/mCherry in *ChAT::Cre +/-; RΦGT+/-* mice carrying an allele expressing GFP under the control of the neuronal glycine transporter (GlyT2; Zeilhofer et al., 2005). Since a large overlap in the expression of GlyT2 and GABA has been shown in the spinal cord (Todd and Sullivan, 1990; Todd et al., 1996), the majority of GlyT2^{on} interneurons are inhibitory Glycinergic/GABAergic cells. We defined GlyT2^{off} neurons as excitatory, although a small number in the superficial dorsal horn could be purely GABAergic inhibitory interneurons.

We monitored the GlyT2 status of cells while performing transsynaptic labelling from LG and TA motor neurons. First, we compared the distribution of premotor excitatory interneurons (GlyT2^{off}; Figures 6A-C) and then examined the distribution of inhibitory (GlyT2^{on}; Figures 6D-F). The distribution of GlyT2^{off} premotor interneurons was the same for LG and TA motor neurons (4 LG and 3 TA single injections, Figure 6A). The medians of the medio-lateral position in the dorsal ipsilateral cord were 306 μm and 326 μm for LG and TA respectively (hierarchical bootstrapped Hedges'G = -0.03, IQR -0.08,0.01) and -112 μm and 74 μm (hierarchical bootstrapped Hedges'G = -0.14, IQR -0.24, -0.04) in the ventral spinal cord (Figures 6 B-C). Similarly, we did not observe segregation in the distribution of GlyT2^{on} LG and TA premotor interneurons (Figure 6D). The medians of the medio-lateral coordinates of the dorsal GlyT2^{on} interneurons were 302 μm for LG and 346 μm for TA (hierarchical bootstrapped Hedges'G = -0.30, IQR -0.35, -0.24), while for ventral interneurons were 395 μm for LG and 437 μm for TA (hierarchical bootstrapped Hedges'G = -0.33, IQR -0.40,0.27, Figures 6E-F). High correlation values (r>0.79). between all individual experiments underscored the conserved positional organization of LG and TA premotor interneurons. T~~hus, t~~hese data indicate that there is no significant difference in the distribution of putative excitatory or inhibitory premotor interneurons controlling the activity of flexor and extensor muscles.

Finally, we compared the distributions of excitatory and inhibitory premotor interneurons separately for each muscle, LG (Figures 6G-I) and TA (Figures 6J-L). No differences were observed for the medio-lateral distribution of inhibitory and excitatory dorsal premotor interneurons (medians for LG: GlyT2^{on} = 303 μm and GlyT2^{off} = 306 μm; hierarchical bootstrapped Hedges'G = -0.06, IQR 0.03,0.09. Medians for TA: GlyT2^{on} = 346 and GlyT2^{off} = 326 μm; hierarchical bootstrapped Hedges'G = 0.15, IQR 0.08,0.23). In contrast, ventral ipsilateral GlyT2^{on} were more abundant than GlyT2^{off} for both LG (Figure 6H-I) and TA (Figure 6K-L). Conversely, GlyT2^{off} premotor interneurons dominated the ventral contralateral side (Figure 6H,K; medians for LG: GlyT2^{off} = -112 μm and GlyT2^{on} = 395 μm; hierarchical bootstrapped Hedges'G = 0.72, IQR 0.61,0.84. Medians for TA: GlyT2^{off} = 73 μm and GlyT2^{on} = 437 μm; hierarchical bootstrapped Hedges'G = 0.91, IQR 0.82,0.99). Overall, the data show a clear segregation in the distributions of excitatory and inhibitory premotor interneurons in the ventral half of the spinal cord: GlyT2^{on} inhibitory interneurons are almost exclusively found in the ipsilateral side while GlyT2^{off} excitatory interneurons also

present a prominent peak in the contralateral side (Figures 6G-K). These observations are reflected in the overall low correlation value in the position of excitatory and inhibitory premotor interneurons for both LG and TA premotor interneurons (r≤0.4).

Taken together, these findings indicate that while we are able to detect significant differences in the positional organization of excitatory and inhibitory premotor interneurons, we found that the organization of flexor and extensor premotor circuits were always intermingled regardless of their neurotransmitter status.

Flexor and extensor premotor interneurons tracing in Olig2::Cre +/-; RΦGT+/- mice

An important consideration concerning the use of our genetic approach for G complementation is the expression specificity of the Cre driver: recombination in multiple neuronal subtypes can potentially result in loss of monosynaptic restriction and rabies transfer across multiple synapses (Figures 1D and 1E). In the spinal cord, *ChAT::Cre* is not only expressed in motor neurons but also cholinergic interneurons, including medial partition cells (V0c neurons) that have prominent projections to motor neurons (Zagoraiou et al., 2009). Therefore, given that under our experimental conditions, V0c neurons express G and are presynaptic to motor neurons, they could permit disynaptic rabies transfer: first from motor neurons to V0c neurons and second from V0c neurons to their presynaptic partners. However, it is important to note that V0c presynaptic partners have been previously characterised using rabies monosynaptic tracing and comprise mostly interneurons located in the dorsal laminae of the spinal cord (Zampieri et al., 2014), an area that is largely devoid of rabies labelling in our experiments.

In order to test whether disynaptic transfer from premotor interneurons is affecting our analysis, we performed a set of experiments (4 gastrocnemius, GS and 3 TA injections) using the *Olig2::Cre* (Dessaud et al., 2007) instead of the *ChAT::Cre* line (Figure 1E). This line would ensure recombination in motor neurons but not in V0c or other cholinergic interneurons. However, Olig2 is transiently expressed during embryonic development in subsets of p2 and p3 progenitors (Chen et al., 2011). We reasoned that if additional transsynaptic transfer from premotor interneurons is significantly affecting our results, using a different Cre line to drive G expression in a non-overlapping subset of premotor interneurons should result in different labeling patterns. We performed monosynaptic tracing experiments after single injections of ΔG-RabV/mCherry in either the TA or GS muscles of P4 *Olig2::Cre +/-; RΦGT+/-* mice. Six days following injection, we observed interneuron labelling with a pattern similar to that of those performed in *ChAT::Cre* mice. There was no difference in the positional organization of flexor and extensor premotor interneurons (Figures 7C-E individual experiments are shown in Figure 12- figure supplements 11-12), with median values along the mediolateral axis of 281 μm for GS and 301 μm for TA (bootstrapped Hedges' G = -0.02, IQR -0.07,0.03).Comparison of the premotor maps obtained from *ChAT::Cre* and *Olig2::Cre* experiments

showed that interneuron distributions were indistinguishable, as shown by the high correlation values across mouse lines and muscles (r>0.9) and the low values of hierarchical bootstraps of the Hedges' G coefficients across muscles and *Olig2-Cre* and *ChAT-Cre* injections (Figure 7F). In addition, the median interneuron positions along the medio-lateral axis for each experiment were similar between *ChAT::Cre* (314 µm for GS and 308 µm for TA, including high and low efficiency experiments) and *Olig2::Cre* (281 µm for GS and 301 µm for TA) animals injected in the same muscle (Figure 7G, Hedges' G =0.14 (IQR 0.09,0.18) for GS pairs and 0.13 (IQR 0.08,0.18) for TA pairs). Thus, these results indicate that under our experimental conditions the results of tracing experiments done in *ChAT::Cre +/-; RΦGT+/- and Olig2::Cre +/-; RΦGT+/-* mice are unlikely to be influenced by disynaptic rabies transfer from spinal premotor interneurons.

Flexor and extensor premotor interneurons tracing with AAV complementation methods

In contrast to previous findings using AAV-G (Tripodi et al., 2011), we found extensive intermingling of flexor and extensor premotor interneurons when using Cre-based methods to genetically-express G-protein in order to complement rabies replication (Figure 1D, E). Therefore, we sought to replicate the previous findings using AAV-G complementation strategies (Figure 1A and B). We first performed injections of 1 µl of a 3:1 AAV-G: ΔG-RabV mixture in GS and TA muscles of P2 wild-type mice (Figure 1A;(Tripodi et al., 2011). Two experiments were performed on GS only, with similar protocol, and GS single injections were pooled with the data resulting from the double injections. Ten days following injection, consistent labelling with both viruses was observed throughout the lumbar region (L1: Figure 8A, and L5: Figure 8B). Analysis of rabies-labelled interneurons did not reveal any apparent difference in the distributions of premotor circuits controlling the activity of antagonist muscles (Figure 8C, individual experiments are shown in Figure 12- figure supplements 13). The Hedges' G coefficients of premotor interneuron distributions in the ipsilateral dorsal quadrant for all the experiments had a median of -0.02 (IQR -0.29,0.28), indicating high uniformity across experiments, as also shown by the high correlation values of positional coordinates of flexor and extensor interneurons (Figure 8D). When the analysis was restricted to the medio-lateral positions in the ipsilateral dorsal quadrant, the median distances from the midline were 249 µm for GS and 295 µm for TA (Figure 8E), a negligible shift compared to the typical interneuron soma diameter. These findings obtained using AAV-G complementation are therefore consistent with the pattern of premotor interneuron mixing identified using genetic-delivery of G-protein.

In these AAV-G complementation experiments, we observed extensive labelling of superficial dorsal interneurons, a feature that was absent when using genetic complementation methods (compare Figures 8C with 2C and 7E). We therefore reasoned that contribution of transsynaptic transfer from sensory afferents, precluded by our genetic complementation approaches, may result in tracing of dorsal and

medial interneurons and thus affect analysis of premotor circuits (Zampieri et al., 2014). To test this idea, we restricted expression of G to motor neurons by injecting AAV-flex-G in GS and TA muscles of *ChAT::Cre* +/- mice (1 µl of a 3:1 AAV-flex-G: ΔG-RabV mix), a strategy that was previously used to show segregation of flexor and extensor premotor circuits (Figure 1B; Wang et al., 2017). In these experiments (4 GS and 3 TA injections), the distributions of GS and TA premotor interneurons were similar (Figure 9A-C, individual experiments are shown in Figure 12- figure supplements 14-15), with high correlation values across experiments (Figure 9D). Analysis of the medio-lateral positioning in the ipsilateral dorsal quadrant showed that the median position of TA related interneurons was shifted by only 30 µm with respect to that of the GS related interneurons, and that this was in a *medial* direction (375 µm for GS and 344 µm for TA; Hedges' G from hierarchical bootstrap = -0.15, IQR -0.24, -0.05, Figure 9E).

Anterograde transsynaptic jumps alter the distribution of premotor interneurons

Regardless of whether the expression of G was restricted to motoneurons (Figure 9) or not (Figure 8) we observed no differences in the mediolateral position of dorsal ipsilateral premotor interneurons related to flexor or extensor muscles. However, the use of AAV-G-Flex that, contrarily to AAV-G, prevents anterograde jumps by restricting G expression to motoneurons, led to important differences. Indeed, despite being performed and analysed by the same lab at similar timepoints for injection (P1-P2) and harvesting (7-8 days post-injection), experiments using AAV-G result in an extensive labelling of superficial dorsal horn neurons for both tested muscles (Figure 10A).

The median values along the dorsoventral axis were 249 and 375 µm for GS and 295 and 344 µm for TA following AAV-G or AAV-Flex-G injections respectively. The Hedges' G bootstrapped coefficients were -0.55 (IQR -0.64, -0.46) for the GS pairs and -0.14 (IQR -0.23, -0.08) for the TA pairs. In addition, there was an excess of medially located infected neurons, in both GS and TA experiments, when the expression of G was not restricted to motoneurons. The dorsoventral and mediolateral differences in the distributions observed in AAV-G and AAV-G-Flex experiments were substantial for both GS and TA, thus indicating that when G could be expressed by sensory neurons (AAV-G muscle injection), the average position of labelled cells extended more dorsally and medially compared to experiments in which G could only be expressed in motor neurons (AAV-G-Flex). The observed medio dorsal shift is compatible with a contribution of anterograde tracing from sensory neurons (Figure 10A-C), which is absent when G is restricted to motor neurons, either by using AAV-G-Flex, or by genetic restrictions. Nonetheless, regardless of the possible effects derived from anterograde tracing, the distributions of labelled cells observed in flexor and extensor experiments did not differ.

Flexor and extensor premotor interneurons organization using pseudorabies virus timed tracing experiments

Finally, we studied organization of premotor circuits using a different viral tracing method. Pseudorabies virus (PRV) is a neurotropic virus that travels transsynaptically in the retrograde direction. It has previously been used to resolve the connectivity order in polysynaptic circuits (Gu et al., 2017; Ugolini, 2020) using timed infections protocols. In particular, the Bartha strain has been shown to trace spinal premotor circuits around 40 hours (h) after injection (Jovanovic et al., 2010). We simultaneously injected P11 GS and TA muscles with 0.5 µl of PRV-Bartha (PRV-152 and PRV-614). Analysis of the spinal cords two days after injection showed extensive labelling of interneurons throughout the lumbar segments (Figures 11A-B). There were no differences in the distributions or median mediolateral positioning of flexor and extensor premotor interneurons (Figure 11C, individual experiments are shown in Figure 12- figure supplements 16), and correlation analysis showed high correlation coefficients between experiments and across muscles (Figure 11D) and low values of the Hedges' G for pairs of experiments (median 0.09, IQR 0.04,0.22). Accordingly, the medians of mediolateral positions of flexor and extensor premotor interneurons were similar (345 µm for GS and 362 µm for TA; Figure 11D).

We finally pooled together all the results obtained with different methods. In all cases, the correlations between flexor and extensor muscles labelled with the same methods were very high (Figure 12A). The lowest correlation coefficients (which were still high) were typically observed between experiments with AAV-G injection in wild type mice and all the other methods, regardless of the muscle injected, consistent with a substantial contamination from interneurons anterogradely labelled through sensory afferents. Overall, the medians of the mediolateral positions (Figure 12B) did not differ between muscles for any of the methods used and the range of bootstrapped Hedges' G coefficient was between -0.23 and 0.17 across pairs of antagonist muscles throughout five different experimental paradigms tested. The whole set of experiments is shown in Figure 12- figure supplements 1-16. The labelling of experiments keeps the original lab conventions and is also reported in Table 1. Taken together, our findings indicate that premotor interneurons innervating flexor and extensor motor neurons are not spatially segregated.

Discussion

Spinal circuits are responsible for integrating descending commands and sensory information to ensure precise control and coordination of movement. In order to understand how these circuits organise movement, it is necessary to first identify and then study the roles and contributions of spinal interneurons that control the activity of different muscles. Previous work (Tripodi et al., 2011) exploited rabies monosynaptic tracing to examine the organization of premotor circuits controlling the activity of selected muscles. These studies, using intramuscular injection of an AAV expressing the rabies glycoprotein G, identified clear segregation in the spatial organization of premotor interneurons directing the activity of flexor and extensor muscles (Tripodi et al., 2011; Wang et al., 2017; Takeoka and Arber, 2019). In contrast, our systematic study using either genetic or AAV complementation of G expression and PRV timed infections, demonstrates complete spatial overlap amongst flexor and extensor premotor interneurons.

Mouse genetic-based strategies for rabies monosynaptic tracing of premotor circuits

We opted for a mouse genetic strategy that was previously used to trace premotor circuits of vibrissal, orofacial, and forelimb muscles (Takatoh et al., 2013; Stanek et al., 2014; Skarlatou et al., 2020). Combining a conditional allele expressing G from the rosa locus (*RΦGT* mice; Takatoh et al., 2013) with either the *ChAT::Cre* or *Olig2::Cre* lines (Figure 1D-E) is predicted to result in high levels of G expression in all motor neurons at the time of rabies muscle injection and therefore in robust transsynaptic transfer. Indeed, under these conditions, several hundred premotor neurons can be reproducibly traced in each experiment (Table 1; (Skarlatou et al., 2020; Ronzano et al., 2021). On the other hand, this strategy suffers from the undesirable consequences of lineage tracing, namely G complementation in all Cre expressing cells in the spinal cord, including those that transiently activate the targeted promoter during development. This problem is in part shared with the AAV-based experiments using intraspinal injection of AAV-flex-G in *ChAT::Cre* mice (Figure 1C; Takeoka and Arber, 2019). Thus, it is unlikely that the differences in the results obtained using these two strategies was caused by disynaptic transfer through cholinergic interneurons. Indeed, work using rabies monosynaptic tracing to identify spinal neurons presynaptic to the most prominent population of premotor cholinergic interneurons, V0c neurons, found a high density of pre-V0c neurons located in superficial laminae of the dorsal horn (Zampieri et al., 2014), an area where no labelling was observed in our mouse genetic-based experiments.

Nonetheless, to further explore the possibility of disynaptic pathways via cholinergic spinal interneurons, we examined the organization of flexor and extensor premotor circuits in experiments conducted with the *Olig2::Cre +/-; RΦGT+/-* line. Olig2 is expressed in motor neurons and in subsets of p2 and p3 progenitors, thus potentially generating opportunities for disynaptic transfer through V2 and V3 premotor interneurons (Chen et al., 2011). We did not find any difference in the distribution of premotor interneurons obtained in

these mice compared to those in *ChAT::Cre+/-; RΦGT+/-* mice. Therefore, it appears that disynaptic transfer is not a frequent event in our experimental conditions; otherwise, neuronal labelling in *ChAT::Cre* experiments would reflect the contribution of cholinergic interneurons, and *Olig2::Cre* experiments would reflect jumps through V2 and V3 interneurons. It is also important to consider the timing of rabies transsynaptic transfer (Ugolini, 2011). The earliest expression of rabies in primary infected motor neurons is first observed 3-4 days after injection and monosynaptic transfer not earlier than 5 days after injection, with strong labelling observed around 7-8 days. Since in our experiments, mice were sacrificed between 6 to 9 days following RabV injections (8-9 days in *ChAT::Cre* and 6 days in *Olig2::Cre* mice), it is unlikely that many, if any, double jumps would have occurred in this time window. Together, these considerations indicate that contamination of premotor circuit maps with interneurons labelled from transsynaptic transfer events originating from G expressing spinal interneurons is not likely to be the source of the different results obtained in AAV and mouse genetic experiments.

AAV- based strategies for rabies monosynaptic tracing of premotor circuits

In previous studies in which segregation of flexor and extensor premotor interneurons have been observed, AAV was used to express G in motor neurons. In the first report, AAV-G and ΔG-RabV were co-injected intramuscularly in wild type mice (Tripodi et al., 2011). This approach has the advantage of complementing G only in motor neurons projecting to the targeted muscle, thus avoiding the problem of G expression in spinal interneurons that could lead to loss of monosynaptic restriction. However, since sensory neurons in the dorsal root ganglia also innervate muscles, such strategy could lead to anterograde transsynaptic spread to the spinal cord through the sensory route (Figure 1A, Zampieri et al., 2014). In order to avoid this problem, intramuscular co-injection of a conditional AAV vector (AAV-flex-G) with ΔG-RabV in *ChAT::Cre* mice was used (Figure 1B, Wang et al., 2017). In this more stringent condition, G would only be expressed in motor neurons. A more recent study used intraspinal injection of AAV-flex-G in *ChAT::Cre* mice (Figure 1C, Takeoka and Arber, 2019), which would also avoid transfer from sensory neurons. However, despite the fact that these AAV-based strategies have distinct advantages and disadvantages, they all resulted in labelling of flexor and extensor premotor interneurons with distributions that were medio-laterally segregated in the dorsal ipsilateral quadrant of the spinal cord in experiments performed on neonatal (Tripodi et al., 2011) and adult (Takeoka and Arber, 2019) mouse hindlimbs, as well as neonatal forelimbs (Wang et al., 2017).

In order to resolve the discrepancy with the results obtained with genetic complementation, we tried to replicate the previous findings by directly testing two of the AAV complementation strategies, namely AAV-G and ΔG-RabV co-injection in wild type mice (Figure 1A;(Tripodi et al., 2011), and AAV-flex-G and ΔG-RabV co-injection in *ChAT::Cre* mice (Figure 1B; Wang et al., 2017). Given the small size of the injected

muscles, we limited our injection volumes to 1 μl, as opposed to the 5 μl used in the original study (Tripodi et al., 2011) routinely checking for injection specificity by examination of the leg muscles. Surprisingly, despite our attempts at replication, we did not observe segregation of flexor and extensor premotor circuits. However, the contribution of anterograde transsynaptic spread to the spinal cord through the sensory route was clearly detected in AAV-G experiments (but, as expected, not in AAV-flex-G experiments), reflecting the contribution of muscle-innervating sensory afferents that have post-synaptic targets predominantly located in the dorsal and medial aspects of the spinal cord (Zampieri et al., 2014; Pimpinella et al, 2022). Notably, this observation does not explain why we failed to replicate segregation of flexor and extensor premotor circuits, as we detected overlapping distributions both in the presence (AAV-G) or absence (AAV-flex-G) of sensory contributions. It is nonetheless interesting to notice that in the original report using AAV-G and ΔG-RabV co-injection in wild type mice, the only condition where flexor-extensor segregation was not reported by the authors is upon elimination of proprioceptors with diphtheria toxin (Tripodi et al., 2011).

Given that we used the same serotype of AAV and strain of RabV, the one clear difference in the experiments is the total volume of fluid injected intramuscularly. We limited our volume to 1 μl as we measured the sizes of these muscles to be around 2 μl in the early post-natal period. Thus, it is possible that the larger volume injected in the original study might have skewed the results, for example by differential infection of proprioceptive afferents innervating the flexor vs extensor muscles, or by infection of cutaneous afferents supplying the overlying skin (Li et al., 2011).

The issue of starter motor neurons

The identity and number of starter cells are the main determinants of reproducibility in rabies tracing experiments and thus represent key parameters for comparing different approaches. For experiments using ΔG-RabV, starter cells are those that are both primarily infected with RabV and express G. In general, for both the AAV and mouse genetics methods discussed here, it is difficult to precisely determine these factors, as neither approach employs expression of a reporter gene to mark G-expressing cells. Moreover, RabV is known to be toxic to neurons and some primary infected motor neurons are likely to die before analysis (Reardon et al., 2016). Because of the well-known topographic organization of neuromuscular maps, muscle identity of infected motor neurons can be inferred by their stereotyped position in the spinal cord (Romanes, 1964; McHanwell and Biscoe, 1981). Thus, for all the methodologies discussed here, it is only possible to approximate the identity and number of starter motor neurons by surveying the position of RabV-infected motor neurons present at the end of the experiment.

Restriction of starter "status" to motor neurons connected to a single muscle is determined by two aspects: the specificity of rabies virus injection and the availability of sufficient levels of G protein in the same cells (Callaway and Luo, 2015). All the approaches discussed here used intramuscular injections of G-deleted

rabies virus (SAD-B19) to selectively infect a motor pool. In this step, sources of variability are represented by 1) specificity of muscle injection and 2) the titre of the rabies virus injected. Muscle injection specificity was routinely checked following injections of adjacent synergist muscles and for all the co-injections of AAV-G and RabV. Rabies leak from antagonist muscles (LG and TA) located in different anatomical compartments on opposite sides of the tibia and fibula would be very unlikely. The titre of the injected rabies virus can affect the efficiency of primary infection: the data presented here show that the RabV titre, while affecting the number of motor neurons and secondary neurons labelled, does not affect the overall distribution of premotor interneurons. The same data indicate that the presence of a small number of "ectopic" motor neurons (seen only following high titre injections) does not significantly contribute to the tracing results, as the premotor distributions in high and low titre experiments are not different. Furthermore, these "ectopic" motor neurons likely represent recurrently connected presynaptic motor neurons (Bhumbra and Beato, 2018); therefore any labelling originating from them would represent a much less frequent disynaptic transfer event.

In the short term, the introduction of a reporter system to label G-expressing neurons, as routinely done in many rabies experiments, combined with the use of non-toxic rabies variants that would prevent motor neuron death (Reardon et al., 2016; Ciabatti et al., 2017; Chatterjee et al., 2018) will help resolve potential confusion about the identity and number of starter cells. Such tools could be used in both the AAV and the mouse genetic approaches. In addition, the ability to precisely restrict the selection of starter motor neurons either by the introduction of more specific Cre lines (e.g. Koronfel et al., 2021) or through the use of novel intersectional genetic and viral strategies could improve premotor tracing experiments. Finally, tracing from single motor neurons using delivery of DNA for G and TVA expression via patch clamp is a precise way to generate specific starter cells (Marshel et al., 2010; Rancz et al., 2011). This approach, followed by intraspinal injection of EnvA-pseudotyped ΔG-RabV, would ensure infection and pre-synaptic tracing from only selected neurons. While technically challenging, this method would have the added value of directly showing whether functionally distinct motor neurons within a pool receive differentially distributed presynaptic input.

Functional implications

While it has been proposed that spatial segregation of premotor interneurons provides an anatomical substrate for labelled line inputs from proprioceptive afferents to motor neurons (Tripodi et al., 2011), our data do not support such a model. Proprioceptive afferents projecting to the intermediate and dorsal spinal cord relay many types of information (e.g., changes in muscle length, muscle length itself, force, joint position), and form synapses with both excitatory and inhibitory interneurons that process and convey these data to flexor and extensor motor neurons to precisely regulate patterns of contraction. It is clear that, at least in the case of motor neurons, position plays an important role in the specificity of afferent

inputs received (Sürmeli et al., 2011), and it is reasonable to think that it might also be the case for interneurons. Indeed, it has been shown for V1 inhibitory interneurons that cell body positioning constrains wiring from proprioceptive afferents(Bikoff et al., 2016). However, the location of V1 subtypes does not seem to influence their output connectivity to motor neurons, as indicated by the case of Sp8+ V1 interneurons that are located in a medial position in the intermediate spinal cord, in about the same location described for premotor interneuron serving extensor motor pools (Tripodi et al 2011) but send uniform synaptic output to both flexor and extensor motor pools (Bikoff et al., 2016). Our data are consistent with the idea that interneuron position plays an important role in controlling sensory input connectivity, but not their output connectivity to motor neurons. That is, positional organization has been shown to represent a developmental strategy that facilitates wiring onto spatially segregated populations of neurons, rather than determining output connectivity, where neuronal position does not seem to have a major role.

Conclusion

In conclusion, despite using five different methods, we have not been able to find evidence that spinal premotor interneurons innervating flexor vs extensor motoneurons are segregated. It is important to stress that none of the methods discussed here is completely exempt from potential problems (Table 2). However, full appreciation of the strengths and weaknesses of each approach can guide both the choice of method for mapping premotor circuits and the interpretation of the results obtained.

Methods

Experimental settings

The experiments were performed and analysed across 4 different laboratories. All the injections on *ChAT::Cre* mice were performed at UCL, with locally produced RabV, either in the Beato or in the Brownstone lab, by 5 different operators. Tissue preparation and image acquisition were performed either in the Brownstone lab (3 different operators) or at the University of Glasgow (UoG, Maxwell lab, 1 operator). The injections in *Olig2::Cre* mice were performed and analysed at the Max Delbrück Center (MDC), with locally produced RabV. The injections of AAV-G and AAV-G-Flex in wild type and *ChAT::Cre* mice respectively were performed and analysed at the Salk Institute. The AAV-G construct was produced from Applied Viromics, while the AAV-G-Flex was produced at the Salk GT3 Core and the RabV was obtained from Janelia Farm. Experiments with PRV-Bartha rabies strain were performed and analysed at the Salk Institute. PRV-Bartha was obtained from NIH CNNV.

Animal Experimentation Ethical Approval

All experiments at UCL were carried out according to the Animals (Scientific Procedures) Act UK (1986) and were approved by the UCL AWERB committee under project licence number 70/9098. All experiments performed at the MDC were carried out in compliance with the German Animal Welfare Act and approved by the Regional Office for Health and Social Affairs Berlin (LAGeSo). All experiments performed at the Salk Institute were conducted in accordance with IACUC and AAALAC guidelines of the Salk Institute for Biological Studies.

Mouse strains

Homozygous *ChAT::Cre* mice (Rossi et al., 2011, Jackson lab, stock #006410) or heterozygous *Olig2::Cre* mice (Dessaud et al., 2007) were crossed with homozygous *RΦGT* mice (Jackson Lab, stock #024708), to generate *ChAT::Cre +/-; RΦGT +/-* or *Olig2::Cre +/-; RΦGT +/-* mice (Takatoh et al., 2013; Skarlatou et al., 2020) that were used for rabies tracing experiments. For experiments aimed at distinguishing excitatory and inhibitory populations of premotor interneurons, we crossed homozygous *ChAT::Cre* mice with heterozygous *GlyT2-eGFP* mice (a gift from Prof. Zeilhofer, University of Zurich, Zeilhofer et al., 2005) and their double-positive offspring were mated with the homozygous *RΦGT* mice. AAV-G-Flex constructs were injected in heterozygous *ChAT::Cre* mice

To quantify possible "leak" of Cre expression in the spinal cord, we crossed *ChAT::Cre* and *RCL-tdTom* (Ai9, Jax stock #007909) mice (13 sections from 3 mice), and found tdTom expression in cholinergic motor neurons and interneurons as expected, as well as in some non-cholinergic neurons distributed in

intermediate (10%, 70/690) and dorsal (9%, 62/690) laminae, with the remaining 35 located ventrally
(Figure 13) indicating that ectopic expression of Cre in ChAT negative neurons is minimal but not nil, and
mostly confined to superficial dorsal laminae. The possible "leak" of G and the avian receptor protein (TVA)
expression in *RΦGT* mice was then tested by injecting EnvA-ΔG-Rab-eGFP, produced according to
standard protocols (Osakada and Callaway, 2013) to a titre of 1×10^9 IU/ml. Lack of contamination from
non-pseudotyped virus was confirmed by infecting HEK cells at high (up to 20) multiplicity of infection.
Three *RΦGT* heterozygous mice were injected in the lateral gastrocnemius muscle at P1 and fixed 9 days
post injections. The tissue was cut as described below, but along the sagittal plane in 60 μm sections, in
order to isolate the dorsal motor column. Following immunoreaction for eGFP, in each of the 3 cords, we
found a maximum of 3 labelled motor neurons (1, 1 and 3 motor neurons in n=3 animals) but no
interneurons labelled, indicating some leakage in the expression of the TVA-IRES-glycoprotein cassette
from the *RΦGT* mice, but insufficient G expression to support transsynaptic jumps (Figure 13, figure
supplement 1).

Virus production

Rabies virus used in experiments performed at UCL, was obtained from in house stocks of a variant of the
SAD-B19 rabies strain where the sequence coding for the glycoprotein was replaced by the sequence for
either eGFP or mCherry (Wickersham et al., 2007). Virus was produced at high concentration using the
protocol described in (Osakada and Callaway, 2013): baby hamster kidney cells stably expressing the
rabies glycoprotein (BHK-G, kindly provided by Dr. Tripodi (LMCB, Cambridge) were thawed and plated
in standard Dulbecco modified medium, supplemented with 10% fetal bovine serum (FBS), incubated at
37°C with 5% CO_2 and split until ~70% confluence was obtained in 5 dishes (10 ml medium each). The
cells were then inoculated at 0.2-0.3 multiplicity of infection with either the ΔG-RabV-eGFP or the ΔG-
RabV-mCherry (initial samples kindly provided by Prof. Arber and Dr. Tripodi). Cells were incubated for 6
hours at 35°C and 3% CO_2 and then split 1 to 4 into 20 dishes (10 ml) with 10% FBS medium and kept at
37°C and 5% CO_2 for 12-24 hours, until ~70% confluent and medium was then replaced with 2% FBS
medium and cells incubated at 35°C and 3% CO_2 for virus production. The supernatant was collected after
~3 days, new medium was added for another round of production and supernatant was filtered (0.45 μm
filter) and centrifuged for 2 hours at 19400 rpm (SW28 Beckman rotor). The pellets were suspended in
phosphate buffer saline (PBS), dispersed and collected in a single tube and further centrifuged for 4 hours
at 21000 rpm in a 20% sucrose gradient (SW55 Beckman rotor). The resulting pellet was suspended in
100 μl PBS and the virus was aliquoted (5-10 μl) and stored in a -80° freezer. The viral titre of each round
of production was measured by serial 10-fold dilution of three different aliquots using standard protocols
(Osakada and Callaway, 2013). For each injection, the virus titre is reported in Table 1. In a subset of
experiments, we diluted the virus 10-fold in order to limit the number of starter cells. Rabies virus used for

experiments performed at MDC was produced as previously described (Skarlatou et al., 2020). For the
premotor labelling experiments mediated through co-injection of AAV-Flex-G and delta-G-RabV at the Salk
Institute, pSADΔG-mCherry and pSADΔG-eGFP were obtained from Janelia Viral Tools facility. AAV6-
Ef1a-B19G (1x10^13 GC/mL) was generated at Applied Viriomics, while AAV6-CAG-Flex-oG was
generated at the GT3 core at the Salk Institute. For retrograde PRV-Bartha experiments, PRV-152 (GFP)
and PRV-614 (mRFP) were obtained from the Center for Neuroanatomy with Neurotropic Viruses (CNNV).

Intramuscular injection

Neonatal pups (P1-P2) were anaesthetized using isoflurane inhalation and an incision was made on the
skin to expose the belly of the targeted muscle, either lateral or medial gastrocnemius (LG, MG), tibialis
anterior (TA) or peroneus longus (PL). The virus was injected intramuscularly using a 5 µl Hamilton syringe
(model 7652-01) fixed to a manual Narishige micromanipulator (M-3333) and loaded with a bevelled glass
pipette of inner diameter 50-70 µm. The volume injected was 1 µl, compatible with the estimated volume
of the muscles at this age (~2 µl), in order to minimize the risk of leaks to adjacent muscles. Viral batches
of similar titres were slowly (> 1 minute) injected, the skin incision was sutured with Vicryl 8-0 (Ethicon,
USA) and the pups received a subcutaneous injection of carprofen (10%) for pain management. Mice were
closely monitored for the next 24 hours for signs of movement impairment and were perfused under
terminal general anaesthesia 8-9 days after injection.

In order to compare directly the distributions of flexor and extensor associated premotor interneurons and
avoid confounding factors in the coordinate representations of these interneurons across different spinal
cords, we performed double injections of ΔG-RabV in the same animal, using ΔG-Rab-eGFP and ΔG-
Rab-mCherry injected in pairs of antagonist muscles, lateral gastrocnemius and tibialis anterior. For
comparison, we also performed double injections in pairs of synergist muscles: LG and MG or TA and PL.
Due to the close proximity of these pairs of muscles, before cutting the spinal tissue for
immunohistochemistry, we dissected the injected leg and confirmed that there was no contamination of
virus across the injected muscles or in adjacent muscles below or above the knee. To exclude confounding
factors in our observed premotor interneuron distributions due to systematic viral interference (Ohara et
al., 2009b, 2009a), in a subset of experiments single injections of ΔG-Rab-mCherry (4 LG, 4 MG, 2 TA
and 4 PL) were performed in the progeny of either *ChAT:: Cre;RΦGT* or *GlyT2-eGFP;ChAT::Cre;RΦGT*
mice.

For experiments performed at MDC, intramuscular injections were done as previously described
(Skarlatou et al., 2020). Briefly, P4 animals were anesthetized with isoflurane and a small incision in the
skin was made to reveal either the gastrocnemius (GS, 4 experiments, no attempts were made at selective
targeting of the two heads of the GS muscle) or the tibialis anterior (3 experiments) muscles. A volume of

1.5 µl of ΔG-RabV-mCherry was injected *in Olig2::Cre +/-; RΦGT +/-* mice using a glass capillary. Animals

were perfused at P10, 6 days after injection in order to minimize the chance of disynaptic transfer.

For wild type ΔG-RabV premotor labelling experiments performed at the Salk Institute, P2 pups were

anesthetised on ice and 1 µl of a 3:1 mixture of AAV-Ef1a-B19G and ΔG-Rab-GFP or ΔG-Rab-mCherry

was injected through the skin into the GS or TA muscles of the same animal, using a 10 µl Hamilton syringe

with 30-gauge metal needle. Animals were fixed by perfusion under terminal general anaesthesia, as

described below, 10 days post injection. For *ChAT::Cre* experiments, a 3:1 mixture of AAV-CAG-Flex-oG

(Addgene #74292) and deltaG-Rab-GFP was injected into either the GS or TA muscles of P1-P2 pups.

Anaesthesia, injection volume, and syringe used were the same as above. Animals were fixed by perfusion

under terminal general anaesthesia, as described below, 7 days post-injection. Hindlimbs were visualized

under a stereoscope to confirm correct targeting of muscles.

For PRV experiments, intramuscular injections of PRV were performed in P11 pups under 2-3% isoflurane

anaesthesia. Briefly, a small incision was made in the skin to expose the TA and GS and 0.5 µl of PRV-

152 or PRV-614 was delivered to either muscle, in the same animal, using a glass needle and picospritzer.

Animals were perfused 48 hours post injection. Hindlimbs were visualized under a stereoscope and

sectioned to confirmed correct targeting of muscles.

Tissue collection and immunohistochemistry

Under ketamine/xylazine terminal anaesthesia (i.p. 80 mg/kg and 10 mg/kg respectively), mice were

intracardially perfused with phosphate buffer solution (0.1 M PBS), followed by 4% paraformaldehyde in

PBS. The spinal cords were dissected and post-fixed for 2 hours at 4ºC, cryoprotected overnight at 4ºC in

30% PBS sucrose solution and embedded in OCT (optimal cutting temperature, Tissue-Tek, #4583)

compound.

For UCL experiments, injections were all conducted at UCL, whereas the immunohistochemistry and

imaging were conducted on different animals independently in two different laboratories (Maxwell at

Glasgow University and Beato/Brownstone at UCL). For tissue processing performed at UCL, lumbar

spinal cords were cut (30 µm thickness) in series in the transverse plane with a cryostat (Bright

Instruments, UK) mounted onto charged glass slides (VWR, #631-0108), and stored at -20ºC. Sections

were incubated for 36 hours at 4˚C with primary antibodies and overnight at 4˚C with secondary antibodies

in PBS containing 0.3M NaCl, 0.2% Triton 100-X (Sigma, T9284-500ml), 7% donkey normal serum (Sigma,

D9663-10ml). The primary antibodies used were: guinea pig anti-Isl1 (1:7500, from Dr. T Jessell, Columbia

University, New York), goat anti choline acetyl transferase (ChAT, 1:100, Millipore, AB144P), rabbit anti-

GFP (1:2500, Abcam. Ab290), chicken anti-mCherry (1:2500, Abcam, Ab205402). The secondary

antibodies were: donkey anti-guinea pig Alexa 647 (1:700, Millipore, AP193SA6), donkey anti-goat preabsorbed Alexa 405 (1:200, Abcam, ab175665), donkey anti-rabbit Alexa 488 (1:1000, Thermofisher, A21206), and donkey anti-chicken Cy3 (1:1000, Jackson Immunoresearch, #703-165-155). The slides were mounted in Mowiol (Sigma, 81381-250G) and cover-slipped (VWR, #631-0147) for imaging.

At Glasgow University, the spinal cords were sectioned using a Leica VT1000 vibratome (thickness 60 μm) and incubated in 50% ethanol for 30 minutes. Primary antibodies used were: chicken anti-GFP (1:1000, Abcam, Ab13970), rabbit anti-mCherry (1:2000, Abcam, Ab167453) and goat anti-ChAT (1:100, Millipore, AB254118). The secondary antibodies were: donkey anti-chicken A488 (1:500, Jackson Immunoresearch, 703-545-155), donkey anti-rabbit Rhodamine red (1:100, Jackson Immunoresearch, 711-295-152) and either Pacific Blue (1:200, prepared on site using unconjugated donkey anti-rabbit IgG, Jackson Immunoresearch, 711-005-152 and Pacific Blue kit, Invitrogen, P30012) or Alexa 647 (1:500, Jackson Immunoresearch, 705-605-003). Sections were mounted in Vectashield (Vector Laboratories, Peterborough, UK) and coverslipped.

For the experiments performed at MDC, spinal cords were processed as previously described (Skarlatou et al., 2020). Briefly, animals were anesthetized by intraperitoneal injection of ketamine /xylazine mix and transcardially perfused with ice-cold PBS (until the liver was cleared of blood), followed by freshly made ice-cold 4% PFA. Spinal cords were dissected and post-fixed for 90 minutes with 4% PFA on ice. Consecutive 40 μm spinal cord cryosections including the caudal thoracic and lumbar spinal regions were obtained using a Leica cryostat and incubated overnight at 4°C with rabbit anti-ChAT 1:16000 (Sürmeli et al., 2011 RRID:AB_2750952) followed by 1 hour incubation at room temperature with secondary antibody (Alexa-Fluor 488, 1:1000). Slides were mounted in Vectashield.

For PRV and ΔG-RabV experiments performed at the Salk Institute, animals were transcardially perfused with ice cold 1x PBS followed by ice cold 4% PFA (Electron Microscopy Sciences, 100504-858). Spinal cords were dissected and post-fixed in 4% PFA for 2 hours at 4°C. After the post-fixation, samples were washed with 1x PBS for 10 minutes followed by a 1-3 day incubation in 30% sucrose for cryoprotection (4°C). The lumbar spinal cord was then embedded in Tissue-Tek OCT (Sakura 4583) for cryosectioning onto glass slides (Fisherbrand Superfrost slides 12-550-15) in the transverse plane. The section thickness was 60 μm for delta-G-RabV experiments and 30 μm for PRV experiments. Immunohistochemistry was performed by incubating tissue with primary antibodies overnight at 4°C and secondary antibodies for 2 hours at room temperature, in a 0.3% Triton-X and 20% donkey serum blocking buffer. After staining slides were mounted with a glass coverslip and Fluoromount-G (00-49-58-02). The antibodies used for this portion of the study were: goat anti-GFP (1:1000, Rockland 600-101-215), rabbit anti RFP (1:1000, Rockland 600-401-379), donkey anti-goat Alexa 488 (1:1000, Invitrogen A11055), donkey anti-rabbit-Alexa 555 (1:1000, Invitrogen A32794).

Confocal imaging and analysis

For UCL experiments, confocal images were acquired using a Zeiss LSM800 confocal microscope with a 20X (0.8 NA) air objective and tile advanced set up function of ZEN Blue 2.3 software for imaging of the entire slice. The tiles were stitched using ZEN Blue software and cell detection was performed using Imaris (version up to 9.1, Bitplane) software. Cell counts were manually performed on every other section, in order to minimize the risk of counting the same cell twice in two consecutive sections.

For experiments performed at Glasgow University, the images were acquired using a Zeiss LSM710, with a x20 (0.8 NA) air objective and cells were counted manually using Neurolucida. Only a subset of sections was analysed (1 in every 8 consecutive sections), thus accounting for approximately 2 sections for every spinal segment.

For experiments performed at the MDC, confocal images were acquired using a Zeiss LSM800 confocal microscope. Regions of interest corresponding to each section and consisting of 8 tiles were imaged with a 10x (0.3 NA) air objective. The tiles were subsequently stitched using ZEN 2.3 Software. Acquisition and processing were performed immediately after immunohistochemistry where possible, to obtain the best possible signal. The resulting images were used for three-dimensional positional analysis as previously described (Skarlatou et al., 2020).

For experiments performed at the Salk Institute, fluorescent images were acquire using an Olympus VS-120 virtual slide scanner microscope with a 10x objective (0.3 NA). For delta-G-RabV experiments cells counts were performed on every section. For PRV experiments cell counts were performed on every 8th section. Both samples were analyzed in MATLAB using a custom script for cell detection.

A consistent system of coordinates was established using the central canal as origin of the x-y plane. The y-axis was defined as parallel to the dorso-ventral axis, with positive values towards the dorsal side and the x-axis was determined by the mediolateral direction, with positive values on the side of injection. For UCL and University of Glasgow experiments the L4-L5 border in the z direction was determined during the slicing procedure and its location was confirmed post-processing by identifying the slices with the widest mediolateral width. For MDC and Salk Institute experiments, the border between T13 and L1 was chosen as the starting point for slicing. The z coordinates obtained in different labs were subsequently aligned using the widest section as a point of reference for the border between L4 and L5 segments. For both Neurolucida and Imaris data files, in order to account for the different shapes of sections throughout the lumbar cord and deformation of individual sections, normalization of coordinates was performed independently for each quadrant using as reference points those indicated in Figure 13, figure supplement

2: the x dimension was normalized to the outer edge of the white matter at the level of the central canal, while the y dimension was normalized for each quadrant using the outermost points of the white matter for both dorsal and ventral horns. The resulting cylindrical reconstruction of the spinal cord was then scaled to the idealized spinal cord size (1700 μm in the mediolateral direction and 900 μm in the dorsoventral direction) for illustrational purposes. All coordinate transformations were performed using a custom script in MATLAB, adapted to read both Neurolucida and Imaris file formats. For experiments performed at UCL and MDC, infected motor neurons were identified by co-localization of either Isl1 or ChAT, and the presence of the reporter fluorescent protein (eGFP or mCherry) expressed after rabies infection. For experiments analysed at the Salk Institute, no attempt was made to identify infected motoneurons, therefore, the resulting spatial distributions do not distinguish between primary and secondary infected cells. Since infected motoneurons are located in the ventral horn, the inclusion of motoneurons in the overall distributions does not affect our analysis of the mediolateral segregation in the dorsal horn. Distributions of infected interneurons were calculated using a Gaussian convolution with a kernel size evaluated from the standard deviation of the original data (Bhumbra and Dyball, 2010).

Gaussian convolutions were calculated splitting the transverse, normalized, spinal cord profile into ipsi- and contra-lateral, and dorsal and ventral halves, with the corresponding distributions shown surrounding the transverse spinal cord maps. Areas under the top-bottom or left-right distributions of each label sum to 1. Correlations across individual experiments were calculated from the x-y coordinates projected along the rostrocaudal axis by computing a density matrix $\rho_n(x_i, y_i)$ for each experiment n and evaluating the correlation coefficient r_{nm} between experiments n and m using the formula

$$r_{nm} = \frac{\sum_i \sum_j (\rho_n(x_i, x_j) - median(\rho_n))(\rho_m(x_i, x_j) - median(\rho_m))}{\sqrt{(\sum_i \sum_j (\rho_n(x_i, x_j) - median(\rho_n))^2)(\sum_i \sum_j (\rho_m(x_i, x_j) - median(\rho_m))^2)}}$$

Correlations across groups of experiments were calculated using the same method, after pooling all the experiments corresponding to one muscle and one injection technique. We decided not to base our analysis on p-values standard hypothesis testing for two reasons: first, the p-value obtained for any statistical test, be it parametric or not, depends strongly on the sample size (Gómez-de-Mariscal et al., 2021). Given our large sample size (coordinates for several thousands of neurons), this would lead to an unacceptable rate of false positive. Indeed, within our dataset, we obtain very low ($<10^{-4}$) p-values for distributions whose medians where only a few μm apart. Since the spatial resolution of our cell location procedure is no higher than the diameter of the infected cells, such results cannot be considered meaningful. Second, our data have an intrinsically nested structure, therefore, hypothesis testing on pooled experiments would lead to an error caused by ignoring the hierarchical nature of our data. Therefore, when comparing different methods of injections where distributions from multiple animals were

analysed, we first performed a hierarchical bootstrap procedure (described in detail in Saravanan et al., 2020): briefly if *n* animals were injected with a given technique, we resampled (with replacement) the animals (first level bootstrap) and then within each animal, we resampled (with replacement) the coordinates distribution (second level bootstrap). This procedure was repeated 5000 times, giving rise to 5000 resampled experiments. In order to compare such experiments, we calculated the Hedges' G coefficient (Hedges, 1981)across pairs randomly selected within the 5000 replicas and obtained a distribution of Hedges' G. The medians of such distribution and inter quartile range (IQR) are reported for each pair of techniques or muscles we compared. When individual experiments are compared, the Hedges' G is reported for each pair of individual experiments. Rather than providing a dichotomous decision, absolute values of effect sizes can be classified as no effect (0-0.19), small (0.2-0.49), medium (0.5-0.79) and large (0.8 and above, (Hedges, 1981). Sample sizes and median values are reported individually for each experiment in Table 1. All data processing was performed in MATLAB, using custom written software (available on request from corresponding authors).

Acknowledgements

This work was supported by a BBSRC grant to MB, AJT and DJM (grant number BB/L001454), a Leverhulme Trust grant to MB (grant number RPG-2013-176) and a Wellcome Trust Investigator Award to RMB (110193). JDM was supported by a fellowship from the Jane Coffins Child Memorial Fund for Medical Research and 5K99HD096512 from NICHD. BKB was supported by UCSD Cell and Molecular Genetics Training Program (T32 GM007240), Timken-Sturgis Foundation, Salk Women in Science and Marguerite Vogt Award. RMB is supported by Brain Research UK. SS and NZ were supported by the DFG (ZA 885/1-1 and EXC 257 NeuroCure). SLP is supported as the Benjamin H. Lewis chair in neuroscience and Sol Goldman Charitable Trust, and National Institutes of Health (1 U19 NS112959-01 and 1 R01 NS123160-01). We thank Dr. Ariel Levine for participation in some early experiments and helpful discussions on the data. We are grateful to Drs. Silvia Arber and Marco Tripodi for constructive discussions and helpful comments during the preparation of this manuscript.

Figure legends

Figure 1: Schematic diagram of the currently available techniques for labelling premotor interneurons. A: simultaneous injection of RabV and AAV-G into muscles. Rabies transfer pathways that could potentially contaminate the distribution of premotor interneurons are labelled by a question mark and indicated by dashed lines. A: Afferent labelling could lead to anterograde labelling of sensory related interneurons. B: simultaneous muscle injection of RabV and a Cre dependent AAV-G into mice expressing Cre in motor neurons eliminates the risk of anterograde transfer from afferents. C: intraspinal injection of a flexed AAV-G in mice expressing Cre in motor neurons is followed by intramuscular rabies injection. D and E: RabV muscle injection is performed on mice selectively expressing the rabies glycoprotein in cholinergic neurons (D) or neuron expressing the Olig2 transcription factor (E). F: PRV Bartha is injected in muscles and retrogradely spreads through synapses. Restriction to first order interneurons can be achieved by extracting the tissue early (~48 hours) after injection.

Figure 2: Double injections of flexor and extensor muscles shows no segregation of premotor interneurons. A: Maximum intensity projection of a single 30 μm section taken from the L2 segment of a P10 cord infected with ΔG-Rab-mCherry in the LG and ΔG-Rab-EGFP in the TA in ChAT-*RΦGT* mice. B: Same experiment as A, showing a L5 section. C: Projections along the transverse (left), sagittal (middle) and coronal (right) axes throughout the lumbar region. Dots denote individual premotor interneurons, triangles denote infected motor neurons. Convolved density along each axe are shown to the sides of the raw data (top-bottom and left-right distributions in all panels sum to 1). D: Half section of a cord on the side of a double injection of LG and TA in the L4 segment. Some isl1+ motor neurons are labelled in the dorsal nuclei and one (indicated by arrowhead and enlarged in E) is labelled by both fluorescent proteins, indicating a potential transsynaptic jump between antagonist motor neurons (scale bars, 250 μm and 50 μm in D and E respectively).

Figure 3: Consistent distribution of flexor and extensor premotor interneurons across all individual experiments. A, B: Distribution of premotor interneurons of LG (A) and TA (B) for all the injections. Distributions for each individual experiment are represented with different shades of blue and orange. C: All experiments (single or double ΔG-RabV injections) pooled, showing an overlap of the flexor and extensor related distribution in all quadrants of the spinal cord, with individual dots replaced by contours. D: Correlation (top right) and absolute value of Hedges' G coefficient (bottom left) across all pairs of experiments, indicating a high degree of consistency and small effect sizes across all experiments, independent of the muscle injected. E: Box and whisker plot of the mediolateral position of dorsal ipsilateral premotor interneurons in each experiment. F: Values of the median for each LG and TA experiment.

Figure 4: Injection with diluted RabV do not reveal any segregation between flexor and extensor premotor interneurons. A and B show the distribution of LG and TA premotor interneurons on the transverse plane for individual experiments, represented with different colour shades. C: LG and TA distributions overlap and the premotor interneuron distributions are highly correlated across experiments and absolute value of Hedges' G coefficients are small (D). E: Box and whisker plot of the mediolateral position of dorsal premotor interneurons in each experiment. F: Values of the median for each LG and TA experiment.

Figure 5: Pairs of flexor or extensor muscles show similar distributions of premotor interneurons. A, comparison of pooled data from extensor muscles LG and MG injections. B: correlation and absolute value of Hedges' G coefficients across all experiments. C: box and whisker plots of the mediolateral position of dorsal ipsilateral premotor interneurons for each experiment and distribution of median values (D). E: similar plot as A, showing the distribution of premotor interneurons following injections of the flexor muscles TA and PL. Correlations and absolute value of Hedges' G coefficients across each experiment are shown in F. G and H shows the mediolateral distribution and the position of the median for each experiment, respectively.

Figure 6: GlyT2^{on} and GlyT2^{off} premotor interneurons are distributed similarly for TA and LG. A, B: Distribution of GlyT2^{off} (A) and GlyT2^{on} (D) premotor interneurons following LG and TA injections pooled from 4 LG and 3 TA experiments in *GlyT2-EGFP::ChAT-Cre* mice crossed with RΦGT mice, indicating that neither class of premotor interneurons is segregated across muscles. Boxplots and violin plots (B for GlyT2^{off} and E for GlyT2^{on}) show uniformity of distribution across experiments in both the dorsal (top) and ventral (bottom) halves of the cord. C (GlyT2^{off}) and F (GlyT2^{on}) show boxplots and individual values for the medians of the mediolateral distributions restricted to dorsal (top) or ventral (bottom) part of the cord. Ventral premotor GlyT2^{off} and GlyT2^{on} interneurons are differentially distributed. Comparison of excitatory and inhibitory premotor interneurons in LG (G) and TA (J) muscles are similar in the dorsal cord, but differ in the ventral cord, where most ipsilateral premotor interneurons are GlyT2^{on}, and the majority of contralateral premotor interneurons are GlyT2^{off}. Boxplots and violin plots of individual experiments are shown in H for LG and K for TA, highlighting the mediolateral differences in the ventral cord. The medians of the ventral and dorsal distributions are shown in I for LG and L for TA.

Figure 7: The distribution of flexor and extensor premotor INs is similar in *Olig2::Cre; RΦGT*. A, B: Single lumbar sections form animals injected in the GS (A) or TA (B) muscles. C - E: Overlay of individual GS (C) and TA (D) experiments and pooled experiments (E). F: Correlation coefficients and absolute value of hierarchical bootstrapped Hedges' G effect sizes between injections of different muscles and using a

different driver for Cre expression. G: Box and whisker plots of median values of all the medio-lateral distributions.

Figure 8: Injection of AAV-G and RabV into GS and TA muscles does not reveal segregation of premotor interneurons. Example of double infected upper (A and lower (B) lumbar sections. Transverse and longitudinal pooled distributions of premotor interneurons from two experiments are overlapping in all quadrants (C) and individual experiments are highly correlated and have low absolute value of Hedges' G effect sizes (D) and with similar medians in the ipsilateral dorsal quadrant. Scale bars 300 μm.

Figure 9: Injection of AAV-FLEX-G in ChAT::Cre mice gives rise to overlapping distributions of flexor and extensor related premotor interneurons. Extensive labelling is observed in two lower lumbar sections of two GS (A) and TA (B) injected mice. The pooled distributions of flexor and extensor related infected neurons are similar on the transverse and longitudinal planes (C). There is strong correlation and low effect sizes across individual experiments (D) and the medians are not different (E). Scale bars 400 μm.

Figure 10: Injection of AAV-FLEX-G prevents anterograde spreading of RabV from afferent terminals. A: side by side comparison of pooled transverse distribution of infected interneurons from GS (left) and TA (right) with AAV-G or AAV-FLEX-G and RabV. The distributions obtained with AAV-G show a marked mediolateral shift and extensive labelling of superficial dorsal horn neurons, a feature that is absent when anterograde transfer is prevented by restricting the expression of the G protein to motoneurons only. The distributions obtained are very similar across the different muscles, but the mediolateral shift is reflected in the lower correlation values and high hierarchical bootstrapped effect size (B) and difference in medians (C) between AAV-G and AAV-FLEX-G injections. Scale bars 400 μm.

Figure 11: Injections of PRV-Bartha in GS and TA muscles give rise to similar distribution of premotor interneurons. Example of a lower lumbar section of an animal injected in GS(A) and TA (B). The distribution of premotor interneurons are similar along all axis (C). Individual experiments are highly correlated, with small effect size (D) and their median values along the mediolateral axis are identical for flexor and extensor injections. Scale bars 300 μm.

Figure 12: Regardless of the tracing methods, flexor and extensor related premotor interneurons distributions are always overlapping. The correlation and absolute value of hierarchical bootstrapped Hedges' G coefficient matrix across pooled experiments obtained with different injection strategies exhibits high values of correlation and low effect sizes across different techniques and across the two different

880   muscles (A). The median values of the mediolateral position of premotor interneurons are similar for all
881   conditions (B).
882
Figure 13: Analysis of ectopic Cre expression in *ChAT-IRES-Cre* and *RCL-tdTom* mice. A: representative
lumbar section stained with antibodies against ChAT (green) and tdTom (red) and (B) map of neurons
labelled with both or one of the two antibodies in all the 13 analysed sections from 3 mice, showing that
some of the tdTom positive neurons do not express ChAT, indicating either a developmental
downregulation of ChAT expression or a modest leak in the Cre expression. C: Venn diagram showing the
overall number of mapped neurons.
889
890

Supplementary figure legends

Figure 3 figure supplement 1: The distribution of premotor interneurons is similar throughout the lumbar
spinal cord: data pooled from 18 experiments (11 LG and 7 TA injections) show that within each lumbar
segment, from L1 to L6, the distributions of LG and TA premotor interneurons are overlapping.

Figure 4 figure supplement 1: High and low efficiency infections give rise to the same premotor
interneurons distributions. Comparison of high and low titre injections are shown in A and E for LG and TA
respectively. Correlations between high and low titre experiments are high and effect sizes are low for both
LG and TA muscles (B and F) The distributions are similar across experiments for both muscles (B and E)
and the median values of the distributions in the ipsilateral dorsal quadrant are not different for high and
low efficiency of infection (C and F).

Figure 12-figure supplement 1: Representation of individual experiments (LG and TA double injections in
ChAT::Cre; RΦGT mice) obtained at UCL. Same experiment code as Table 1. Individual dots denote
premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 2: Representation of individual experiments (LG and TA double injections in
ChAT::Cre; RΦGT mice) obtained at University of Glasgow. Same experiment code as in Table 1.
Individual dots denote premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 3: Representation of individual experiments (LG and MG, double injections
in *ChAT::Cre; RΦGT* mice) obtained at UCL. Same experiment code as in Table 1. Individual dots denote
premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 4: Representation of individual experiments (LG and MG, double injections
in *ChAT::Cre; RΦGT* mice) obtained at University of Glasgow. Same experiment code as in Table 1.
Individual dots denote premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 5: Representation of individual experiments (TA and PL, double injections in
ChAT::Cre; R□GT mice) obtained at UCL. Same experiment code as in Table 1. Individual dots denote
premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 6: Representation of individual experiments (MG single injections in
ChAT::Cre; R☐GT mice) obtained at University of Glasgow. Same experiment code as in Table 1.
Individual dots denote premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 7: Representation of individual experiments (PL single injections in
ChAT::Cre; R☐GT mice) obtained at University of Glasgow. Same experiment code as in Table 1.
Individual dots denote premotor interneurons, triangles denote infected motor neurons.

Figure 12-figure supplement 8: Representation of individual experiments (LG and TA double injections in
ChAT::Cre; R☐GT mice) obtained at University of Glasgow using a diluted RabV (titre <10^9 I.U./ml). Same
experiment code as in Table 1. Individual dots denote premotor interneurons, triangles denote infected
motor neurons.

Figure 12-figure supplement 9: Representation of individual experiments (LG single injections in
ChAT::Cre; R☐GT mice) obtained at University of Glasgow using a diluted RabV (titre <10^9 I.U./ml). Same
experiment code as in Table 1. Individual dots denote premotor interneurons, triangles denote infected
motor neurons.

Figure 12-figure supplement 10: Representation of individual experiments (TA single injections in
ChAT::Cre; R☐GT mice) obtained at University of Glasgow using a diluted RabV (titre <10^9 I.U./ml).. Same
experiment code as in Table 1. Individual dots denote premotor interneurons, triangles denote infected
motor neurons.

Figure 12-figure supplement 11: Representation of individual experiments (GS, single injections in
Olig2::Cre; RΦGT mice) obtained at Max Delbrück Center. Same experiment code as in Table 1. Individual
dots denote premotor interneurons, triangles denote infected motor neurons.

951
952 Figure 12-figure supplement 12: Representation of individual experiments (TA, single injections in
953 *Olig2::Cre; RΦGT* mice) obtained at Max Delbrück Center. Same experiment code as in Table 1. Individual
954 dots denote premotor interneurons, triangles denote infected motor neurons.

955
Figure 12-figure supplement 13: Representation of 2 double injections (GS and TA) and 2 single injections
(GS) of AAV-G and RabV in wt mice) obtained at the Salk Institute. Individual dots denote infected cells,
with no distinction between interneurons and motoneurons

Figure 12-figure supplement 14: Representation of 4 individual experiments (GS single injections of AAV-
FLEX-G and RabV in ChAT::CRe mice) obtained at the Salk Institute. Individual dots denote infected cells,
with no distinction between interneurons and motoneurons

963
964 Figure 12-figure supplement 15: Representation of 3 individual experiments (TA single injections of AAV-
965 FLEX-G and RabV in ChAT::CRe mice) obtained at the Salk Institute. Individual dots denote infected cells,
966 with no distinction between interneurons and motoneurons

967
968 Figure 12-figure supplement 16: Representation of 4 individual experiments (GS and TA double injections
969 of PRV-Bartha) obtained at the Salk Institute. Individual dots denote infected cells, with no distinction
970 between interneurons and motoneurons.

971
972 Figure 13 figure supplement 1: Two examples of a longitudinal section of two different spinal cords from a
973 heterozygous *RΦGT* mouse injected in the LG with EnvA-ΔG-Rab-EGFP, showing a small number of
974 infected motor neurons, but no evidence of transsynaptic jumps, indicating ectopic expression of the TVA
975 receptor, but not of the rabies glycoprotein.

976
Figure 13 figure supplement 2 Schematic of a section of the spinal cord, indicating the reference points
used for normalization. Each section was translated to have the origin of a Cartesian set of axes centered
on the central canal (CC). A line passing through the central canal and perpendicular to the dorso-ventral
axis was used to identify the edge of the white matter in the mediolateral direction (ml) and the ml-CC
distance was used to normalize the x-coordinates. Along the dorso-ventral axes, the 4 edges of the white
matter (ne, nw, se, sw) were identified and their distance from the horizontal line passing through the
central canal were used to normalize the y-coordinates independently in each of the 4 quadrants.
984

Table 1

Code	Lab	Injection	Perfusion	Muscle	Titre I.U.	MNs	Double labelled MNs	Ipsi dorsal median (µm)	Ipsi premotor INs	Contra premotor INs	Total premotor INs	premotor INs/MNs ratio	Section sampling
170427n2	UCL	P2	P10	LG	1×10^{10}	46	3	285	741	116	857	19	1/2 (30 µm)
				TA	5×10^{9}	41		298	912	88	1000	24	
170427n3	UCL	P2	P10	LG	1×10^{10}	32	2	267	620	87	707	22	1/2 (30 µm)
				TA	5×10^{9}	6		276	386	34	420	70	
170503n6	UCL	P2	P10	LG	1×10^{10}	83	1	294	1935	639	2574	31	1/2 (30 µm)
				TA	5×10^{9}	55		315	1887	315	2202	40	
170125n3	UCL	P2	P10	LG	5×10^{9}	39	0	365	670	107	777	20	1/2 (30 µm)
				MG	5×10^{9}	39		353	819	307	1126	29	
170508n7	UCL	P2	P10	LG	1×10^{10}	110	3	352	1955	382	2337	21	1/2 (30 µm)
				MG	5×10^{9}	67		322	1497	429	1926	29	
170125n7	UCL	P2	P10	TA	5×10^{9}	47	0	363	907	308	1215	26	1/2 (30 µm)
				PL	5×10^{9}	39		380	1044	195	1239	32	
170125n8	UCL	P2	P10	TA	5×10^{9}	22	2	343	920	157	1077	49	1/2 (30 µm)
				PL	5×10^{9}	22		330	741	83	824	37	
1570	UoG	P1	P10	LG	2×10^{8}	11	-	322	1111	404	1515	138	1/8 (60 µm)
1571	UoG	P1	P10	LG	2×10^{8}	12	-	340	760	196	956	80	1/8 (60 µm)
1573	UoG	P1	P10	TA	5×10^{8}	10	-	332	447	68	515	52	1/8 (60 µm)
1574	UoG	P1	P10	TA	5×10^{8}	14	-	365	297	26	323	23	1/8 (60 µm)
1577	UoG	P2	P10	LG	2×10^{9}	18	2	329	313	43	356	20	1/8 (60 µm)
				TA	5×10^{9}	26		312	688	105	793	31	
1578	UoG	P2	P10	LG	2×10^{9}	21	5	330	292	34	326	16	1/8 (60 µm)
				TA	5×10^{9}	22		346	790	130	920	42	

1579	UoG	P2	P10	LG	2×10^9	30	1	322	1023	194	1217	41	1/8 (60 µm)
				MG	5×10^8	7		306	169	19	188	27	
1580	UoG	P2	P10	LG	2×10^9	14	0	316	414	48	462	33	1/8 (60 µm)
				MG	5×10^8	8		348	470	87	557	70	
1605	UoG	P1	P10	MG	1×10^8	6	-	340	412	110	522	87	1/8 (60 µm)
1611	UoG	P1	P10	PL	1×10^8	2	-	328	167	24	191	96	1/8 (60 µm)
1613	UoG	P2	P10	PL	1×10^8	1	-	340	164	16	180	180	1/8 (60 µm)
1639	UoG	P2	P10	TA	2×10^8	15	-	341	591	94	685	46	1/8 (60 µm)
1640	UoG	P2	P10	PL	2×10^8	20	-	322	629	122	751	38	1/8 (60 µm)
1644	UoG	P2	P10	LG	1×10^8	1	-	344	142	32	174	174	1/8 (60 µm)
				TA	2×10^8	-		296	57	11	68	-	
1646	UoG	P2	P10	LG	1×10^8	1	-	261	90	16	106	106	1/8 (60 µm)
				TA	2×10^8	3		305	76	13	89	30	
1653	UoG	P2	P10	LG	1×10^8	2	-	307	60	6	66	33	1/8 (60 µm)
				TA	2×10^8	2		312	58	8	66	33	
1656	UoG	P2	P10	LG	1×10^8	-	-	311	563	145	708	-	1/8 (60 µm)
1657	UoG	P2	P10	LG	1×10^8	1	-	321	323	51	374	374	1/8 (60 µm)
1660	UoG	P2	P10	MG	2×10^8	7	-	324	509	3	512	73	1/8 (60 µm)
1661	UoG	P2	P10	MG	2×10^8	10	-	338	175	63	238	24	1/8 (60 µm)
1662	UoG	P2	P10	MG	2×10^8	10	-	313	375	230	605	61	1/8 (60 µm)
1701	UoG	P2	P10	LG	2×10^9	8	2	351	169	26	195	24	1/8 (60 µm)
				MG	5×10^9	34		329	594	190	784	23	
1702	UoG	P2	P10	LG	2×10^9	14	2	331	561	107	668	48	1/8 (60 µm)
				MG	5×10^9	2		322	76	11	87	44	
353	MDC	P4	P10	GS	1×10^9	31	-	283	1542	431	1973	64	All (40 µm)
399	MDC	P4	P10	GS	1×10^9	41	-	286	569	77	646	16	All (40 µm)

1332	MDC	P4	P10	GS	1×10^9	18	-	317	1605	323	1928	107	All (40 µm)
1349	MDC	P4	P10	GS	1×10^9	18	-	305	1416	459	1875	104	All (40 µm)
700	MDC	P4	P10	TA	1×10^9	47	-	318	1723	122	1845	39	All (40 µm)
721	MDC	P4	P10	TA	1×10^9	22	-	310	1934	465	2399	109	All (40 µm)
1324	MDC	P4	P10	TA	1×10^9	17	-	292	2041	301	2342	138	All (40 µm)
1	Salk	P2	P10	GS	1×10^{11}	N/A	N/A	328	9185	2735	11920	N/A	All (60 µm)
				TA	1×10^{11}	N/A	N/A	349	3330	731	4061	N/A	
2	Salk	P2	P10	GS	1×10^{11}	N/A	N/A	303	8827	3867	12694	N/A	All (60 µm)
				TA	1×10^{11}	N/A	N/A	294	3198	1132	4330	N/A	
a	Salk	P1	P8	GS	1×10^{10}	N/A	N/A	248	334	42	376	N/A	1/9 (30 µm)
b	Salk	P1	P8	GS	1×10^{10}	N/A	N/A	237	275	30	305	N/A	1/9 (30 µm)
22a_4	Salk	P2	P9	GS	3×10^{11}	N/A	N/A	403	464	58	522	N/A	All (60 µm)
26a_1	Salk	P2	P9	GS	3×10^{11}	N/A	N/A	383	941	91	1032	N/A	All (60 µm)
26a_2	Salk	P2	P9	GS	3×10^{11}	N/A	N/A	351	1910	401	2311	N/A	All (60 µm)
26a_4	Salk	P2	P9	GS	3×10^{11}	N/A	N/A	382	1923	392	2315	N/A	All (60 µm)
26_1	Salk	P2	P9	TA	3×10^{11}	N/A	N/A	348	3236	263	3499	N/A	All (60 µm)
26_3	Salk	P2	P9	TA	3×10^{11}	N/A	N/A	367	2078	465	2543	N/A	All (60 µm)
26_4	Salk	P2	P9	TA	3×10^{11}	N/A	N/A	350	2494	597	3091	N/A	All (60 µm)
1_1 PRV	Salk	P11	P13	GS	1×10^9	N/A	N/A	318	430	54	484	N/A	1/4 (60 µm)
				TA	1×10^9	N/A	N/A					N/A	
1_4 PRV	Salk	P11	P13	GS	1×10^9	N/A	N/A	349	238	23	261	N/A	1/4 (60 µm)
				TA	1×10^9	N/A	N/A					N/A	
2_2 PRV	Salk	P11	P13	GS	1×10^9	N/A	N/A	357	515	82	597	N/A	1/4 (60 µm)
				TA	1×10^9	N/A	N/A					N/A	
3_3 PRV	Salk	P11	P13	GS	1×10^9	N/A	N/A	377	1005	53	1058	N/A	1/4 (60 µm)
				TA	1×10^9	N/A	N/A					N/A	

Table 1: details of individual experiments performed in the four different laboratories (University College London, UCL, University of Glasgow, UoG, Max Delbrück
Center for Molecular Medicine, MDC, Salk Institute), with individual cell count and virus concentration. Same experimental code as in Figure 3D, 5D, 6C-G, 7F,
9D, 10D, 12D and figure supplements. Experiments performed at the Salk Institute (except the PRV-Bartha experiments) were co-injections of either AAV6-
B19G (3×10^{12} titre) or of AAV6 - CAG-Flex (1×10^{12} titre). AAV and RV were injected in a 3:1 ratio. For all labs and muscles the total volume injected was 1
µl, with the exception of PRV-Bartha experiments, where 0.5 µl were injected. In the Salk Institute experiments, no attempt was made to identify primarily
infected motoneurons, therefore the cell count refers to both motoneurons and interneurons.
992

Method	Pros	Cons	Outcome	Reference
Muscle injection of AAV-G (serotype 2.6) + RabV (Figure 1A)	Avoids the possibility of retrograde disynaptic transfer from second order motor neurons due to restriction of G expression to targeted motor neurons Avoids the possibility of retrograde disynaptic transfer from premotor spinal interneurons.	The labelled premotor population could be contaminated by anterogradely labelled neurons from primary sensory neurons.	Flexor- extensor segregation No flexor- extensor segregation	(Tripodi et al., 2011) Present study
Muscle injection of AAV-flex-G (serotype 2.6) + RabV in *ChAT::Cre* mice (Figure 1B)	Avoids the possibility of retrograde disynaptic transfer from second order motor neurons due to restriction of G expression to targeted motor neurons Avoids the possibility of retrograde disynaptic transfer from premotor spinal interneurons. Avoids potential anterograde sensory contamination.	Conditional expression of G may be inefficient	Flexor- extensor segregation No flexor- extensor segregation	(Wang et al., 2017) Present study
Central injection of AAV-flex-G (serotype 2.9) in *ChAT::Cre* mice followed by muscle injection of RabV, in adults (Figure 1C)	Limits the issue of potential disynaptic transfer from cholinergic interneurons Avoids potential anterograde tracing from sensory neurons	Potential for disynaptic transfer from cholinergic premotor interneurons, transsynaptically labelled motor neurons and mis-targeted primary motor neurons	Flexor- extensor segregation	(Takeoka and Arber, 2019)
	Avoids potential anterograde tracing from sensory neurons	Potential for disynaptic transfer from premotor spinal interneurons, transsynaptically labelled	No flexor- extensor segregation	Present study

Genetically driven expression of G in *ChAT::Cre or Olig2::Cre* mice + muscle RabV injection in neonates (Figure 1D and E)	Ensures homogenous expression of G in all motor neurons	motor neurons and mis-targeted primary motor neurons.		
Muscle injection of PRV-Bartha with strictly timed fixation of tissue (Figure 1F)	High efficiency in transsynaptic transmission. Not reliant on viral recombination.	Timed fixation does not guarantee that transsynaptic jumps occur only up to the second order	No flexor-extensor segregation	Present study

Table 2: Summary of pros and cons of each described method

References

Bacskai T, Rusznak Z, Paxinos G, Watson C (2013) Musculotopic organization of the motor
neurons supplying the mouse hindlimb muscles: a quantitative study using Fluoro-
Gold retrograde tracing. Brain Struct Funct Available at:
http://www.ncbi.nlm.nih.gov/pubmed/23288256.

Bhumbra GS, Beato M (2018) Recurrent excitation between motoneurones propagates
across segments and is purely glutamatergic. PLoS Biology 16.

Bhumbra GS, Dyball RE (2010) Reading between the spikes of the hypothalamic neural
code. J Neuroendocrinol 22:1239–1250.

Bikoff JB, Gabitto MI, Rivard AF, Drobac E, MacHado TA, Miri A, Brenner-Morton S,
Famojure E, Diaz C, Alvarez FJ, Mentis GZ, Jessell TM (2016) Spinal Inhibitory
Interneuron Diversity Delineates Variant Motor Microcircuits. Cell 165.

Callaway EM, Luo L (2015) Monosynaptic Circuit Tracing with Glycoprotein-Deleted Rabies
Viruses. J Neurosci 35:8979–8985.

Chatterjee S et al. (2018) Nontoxic, double-deletion-mutant rabies viral vectors for
retrograde targeting of projection neurons. Nat Neurosci 21:638–646.

Chen J-A, Huang Y-P, Mazzoni EO, Tan GC, Zavadil J, Wichterle H (2011) Mir-17-3p
Controls Spinal Neural Progenitor Patterning by Regulating Olig2/Irx3 Cross-
Repressive Loop. Neuron 69:721–735.

Ciabatti E, González-Rueda A, Mariotti L, Morgese F, Tripodi M (2017) Life-Long Genetic
and Functional Access to Neural Circuits Using Self-Inactivating Rabies Virus. Cell
170:382-392.e14.

Dessaud E, Yang LL, Hill K, Cox B, Ulloa F, Ribeiro A, Mynett A, Novitch BG, Briscoe J
(2007) Interpretation of the sonic hedgehog morphogen gradient by a temporal
adaptation mechanism. Nature 450:717–720.

Gómez-de-Mariscal E, Guerrero V, Sneider A, Jayatilaka H, Phillip JM, Wirtz D, Muñoz-
Barrutia A (2021) Use of the p-values as a size-dependent function to address
practical differences when analyzing large datasets. Sci Rep 11:20942.

Gu Z, Serradj N, Ueno M, Liang M, Li J, Baccei ML, Martin JH, Yoshida Y (2017) Skilled
Movements Require Non-apoptotic Bax/Bak Pathway-Mediated Corticospinal Circuit
Reorganization. Neuron.

Hedges LV (1981) Distribution Theory for Glass's Estimator of Effect Size and Related
Estimators. Journal of Educational Statistics 6:107–128.

Hultborn H, Jankowska E, Lindstrom S (1971) Recurrent inhibition of interneurones
monosynaptically activated from group Ia afferents. The Journal of Physiology Online
215:613–636.

Jankowska E (2001) Spinal interneuronal systems: identification, multifunctional character
and reconfigurations in mammals. J Physiol 533:31–40.

Jovanovic K, Pastor AM, O'Donovan MJ (2010) The use of PRV-Bartha to define premotor
inputs to lumbar motoneurons in the neonatal spinal cord of the mouse. PLoS ONE
5:e11743–e11743.

Koronfel LM, Kanning KC, Alcos A, Henderson CE, Brownstone RM (2021) Elimination of
glutamatergic transmission from Hb9 interneurons does not impact treadmill
locomotion. Sci Rep 11:16008.

Li L, Rutlin M, Abraira VE, Cassidy C, Kus L, Gong S, Jankowski MP, Luo W, Heintz N,
Koerber HR, Woodbury CJ, Ginty DD (2011) The Functional Organization of
Cutaneous Low-Threshold Mechanosensory Neurons. Cell 147:1615–1627.

Marshel JH, Mori T, Nielsen KJ, Callaway EM (2010) Targeting single neuronal networks for
gene expression and cell labeling in vivo. Neuron 67:562–574.

McHanwell S, Biscoe TJ (1981) The localization of motoneurons supplying the hindlimb
muscles of the mouse. Philos Trans R Soc Lond B Biol Sci 293:477–508.

Ohara S, Inoue K, Witter MP, Iijima T (2009a) Untangling neural networks with dual
retrograde transsynaptic viral infection. Front Neurosci 3 Available at:
https://www.frontiersin.org/articles/10.3389/neuro.01.032.2009/full [Accessed
November 18, 2020].

Ohara S, Inoue K, Yamada M, Yamawaki T, Koganezawa N, Tsutsui K-I, Witter MP, Iijima T
(2009b) Dual transneuronal tracing in the rat entorhinal-hippocampal circuit by
intracerebral injection of recombinant rabies virus vectors. Front Neuroanat 3
Available at: https://www.frontiersin.org/articles/10.3389/neuro.05.001.2009/full
[Accessed November 18, 2020].

Osakada F, Callaway EM (2013) Design and generation of recombinant rabies virus vectors.
Nature Protocols 8:1583–1601.

Pimpinella S, Zampieri N (2021) Rabies anterograde monosynaptic tracing reveals
organization of spinal sensory circuits. Available at:
https://www.biorxiv.org/content/10.1101/2021.02.11.429920v1 [Accessed September
3, 2021].

Rancz EA, Franks KM, Schwarz MK, Pichler B, Schaefer AT, Margrie TW (2011) Transfection
via whole-cell recording in vivo: bridging single-cell physiology, genetics and
connectomics. Nat Neurosci 14:527–532.

Rathelot J-A, Strick PL (2009) Subdivisions of primary motor cortex based on cortico-
motoneuronal cells. Proceedings of the National Academy of Sciences 106:918–923.

Reardon TR, Murray AJ, Turi GF, Wirblich C, Croce KR, Schnell MJ, Jessell TM, Losonczy A
(2016) Rabies Virus CVS-N2c Strain Enhances Retrograde Synaptic Transfer and
Neuronal Viability. Neuron Available at:
http://www.ncbi.nlm.nih.gov/pubmed/26804990.

Romanes GJ (1964) THE MOTOR POOLS OF THE SPINAL CORD. Prog Brain Res 11:93–
119.

Ronzano R, Lancelin C, Bhumbra GS, Brownstone RM, Beato M (2021) Proximal and distal
spinal neurons innervating multiple synergist and antagonist motor pools Thoby-
Brisson M, Calabrese RL, Gosgnach S, eds. eLife 10:e70858.

Rossi J, Balthasar N, Olson D, Scott M, Berglund E, Lee CE, Choi MJ, Lauzon D, Lowell BB,
Elmquist JK (2011) Melanocortin-4 Receptors Expressed by Cholinergic Neurons
Regulate Energy Balance and Glucose Homeostasis. Cell Metabolism 13:195–204.

Saravanan V, Berman GJ, Sober SJ (2020) Application of the hierarchical bootstrap to multi-
level data in neuroscience. Neuron Behav Data Anal Theory
3:https://nbdt.scholasticahq.com/article/13927-application-of-the-hierarchical-
bootstrap-to-multi-level-data-in-neuroscience.

Skarlatou S, Hérent C, Toscano E, Mendes CS, Bouvier J, Zampieri N (2020) Afadin
Signaling at the Spinal Neuroepithelium Regulates Central Canal Formation and Gait
Selection. Cell Rep 31:107741.

Stanek E IV, Cheng S, Takatoh J, Han B-X, Wang F (2014) Monosynaptic premotor circuit
tracing reveals neural substrates for oro-motor coordination Mason P, ed. eLife
3:e02511.

Stepien AE, Tripodi M, Arber S (2010) Monosynaptic rabies virus reveals premotor network
organization and synaptic specificity of cholinergic partition cells. Neuron 68:456–
472.

Strack AM, Loewy AD (1990) Pseudorabies virus: a highly specific transneuronal cell body
marker in the sympathetic nervous system. J Neurosci 10:2139–2147.

Sürmeli G, Akay T, Ippolito GC, Tucker PW, Jessell TM (2011) Patterns of Spinal Sensory-
Motor Connectivity Prescribed by a Dorsoventral Positional Template. Cell 147:653–
665.

Takatoh J, Nelson A, Zhou X, Bolton MM, Ehlers MD, Arenkiel BR, Mooney R, Wang F
(2013) New modules are added to vibrissal premotor circuitry with the emergence of
exploratory whisking. Neuron 77:346–360.

Takeoka A, Arber S (2019) Functional Local Proprioceptive Feedback Circuits Initiate and
Maintain Locomotor Recovery after Spinal Cord Injury. Cell Reports 27:71-85.e3.

Todd AJ, Sullivan AC (1990) Light microscope study of the coexistence of GABA-like and
glycine-like immunoreactivities in the spinal cord of the rat. Journal of Comparative
Neurology 296:496–505.

Todd AJ, Watt C, Spike RC, Sieghart W (1996) Colocalization of GABA, glycine, and their
receptors at synapses in the rat spinal cord. J Neurosci 16:974–982.

Tripodi M, Stepien AE, Arber S (2011) Motor antagonism exposed by spatial segregation and
timing of neurogenesis. Nature 479:61–66.

Ugolini G (2011) Rabies Virus as a Transneuronal Tracer of Neuronal Connections.

Ugolini G (2020) Viruses in connectomics: Viral transneuronal tracers and genetically
modified recombinants as neuroscience research tools. Journal of Neuroscience
Methods 346:108917.

Wall NR, Wickersham IR, Cetin A, De La Parra M, Callaway EM (2010) Monosynaptic circuit
tracing in vivo through Cre-dependent targeting and complementation of modified
rabies virus. Proc Natl Acad Sci U S A 107:21848–21853.

Wang X, Liu Y, Li X, Zhang Z, Yang H, Zhang Y, Williams PR, Alwahab NSA, Kapur K, Yu B,
Zhang Y, Chen M, Ding H, Gerfen CR, Wang KH, He Z (2017) Deconstruction of
Corticospinal Circuits for Goal-Directed Motor Skills. Cell 171:440-455.e14.

Wickersham IR, Lyon DC, Barnard RJ, Mori T, Finke S, Conzelmann KK, Young JA,
Callaway EM (2007) Monosynaptic restriction of transsynaptic tracing from single,
genetically targeted neurons. Neuron 53:639–647.

Zagoraiou L, Akay T, Martin JF, Brownstone RM, Jessell TM, Miles GB (2009) A Cluster of
Cholinergic Premotor Interneurons Modulates Mouse Locomotor Activity. Neuron
64:645–662.

Zampieri N, Jessell TM, Murray AJ (2014) Mapping sensory circuits by anterograde
transsynaptic transfer of recombinant rabies virus. Neuron 81:766–778.

Zeilhofer HU, Studler B, Arabadzisz D, Schweizer C, Ahmadi S, Layh B, Bosl MR, Fritschy
JM (2005) Glycinergic neurons expressing enhanced green fluorescent protein in
bacterial artificial chromosome transgenic mice. J Comp Neurol 482:123–141.

1130

Figure 1

Figure 1: Schematic diagram of the currently available techniques for labelling premotor interneurons. A: simultaneous injection of RabV and AAV-G into muscles. Rabies transfer pathways that could potentially contaminate the distribution of premotor interneurons are labelled by a question mark and indicated by dashed lines. A: Afferent labelling could lead to anterograde labelling of sensory related interneurons. B: simultaneous muscle injection of RabV and a Cre dependent AAV-G into mice expressing Cre in motor neurons eliminates the risk of anterograde transfer from afferents. C: intraspinal injection of a flexed AAV-G in mice expressing Cre in motor neurons is followed by intramuscular rabies injection. D and E: RabV muscle injection is performed on mice selectively expressing the rabies glycoprotein in cholinergic neurons (D) or neuron expressing the Olig2 transcription factor (E). F: PRV Bartha is injected in muscles and retrogradely spreads through synapses. Restriction to first order interneurons can be achieved by extracting the tissue early (~48 hours) after injection.

1145

Figure 2: Double injections of flexor and extensor muscles shows no segregation of premotor interneurons.
A: Maximum intensity projection of a single 30 μm section taken from the L2 segment of a P10 cord infected
with ΔG-Rab-mCherry in the LG and ΔG-Rab-EGFP in the TA in ChAT-*RΦGT* mice. B: Same experiment as
A, showing a L5 section. C: Projections along the transverse (left), sagittal (middle) and coronal (right) axes
throughout the lumbar region. Dots denote individual premotor interneurons, triangles denote infected motor
neurons. Convolved density along each axe are shown to the sides of the raw data (top-bottom and left-right
distributions in all panels sum to 1). D: Half section of a cord on the side of a double injection of LG and TA
in the L4 segment. Some isl1+ motor neurons are labelled in the dorsal nuclei and one (indicated by
arrowhead and enlarged in E) is labelled by both fluorescent proteins, indicating a potential transsynaptic
jump between antagonist motor neurons (scale bars, 250 μm and 50 μm in D and E respectively).

Figure 3: Consistent distribution of flexor and extensor premotor interneurons across all individual experiments. A, B: Distribution of premotor interneurons of LG (A) and TA (B) for all the injections. Distributions for each individual experiment are represented with different shades of blue and orange. C: All experiments (single or double ΔG-RabV injections) pooled, showing an overlap of the flexor and extensor related distribution in all quadrants of the spinal cord, with individual dots replaced by contours. D: Correlation (top right) and absolute value of Hedges' G coefficient (bottom left) across all pairs of experiments, indicating a high degree of consistency and small effect sizes across all experiments, independent of the muscle injected. E: Box and whisker plot of the mediolateral position of dorsal ipsilateral premotor interneurons in each experiment. F: Values of the median for each LG and TA experiment.

1166

Figure 4: Injection with diluted RabV do not reveal any segregation between flexor and extensor premotor interneurons. A and B show the distribution of LG and TA premotor interneurons on the transverse plane for individual experiments, represented with different colour shades. C: LG and TA distributions overlap and the premotor interneuron distributions are highly correlated across experiments and absolute value of Hedges' G coefficients are small (D). E: Box and whisker plot of the mediolateral position of dorsal premotor interneurons in each experiment. F: Values of the median for each LG and TA experiment.

Figure 5: Pairs of flexor or extensor muscles show similar distributions of premotor interneurons. A, comparison of pooled data from extensor muscles LG and MG injections. B: correlation and absolute value of Hedges' G coefficients across all experiments. C: box and whisker plots of the mediolateral position of dorsal ipsilateral premotor interneurons for each experiment and distribution of median values (D). E: similar plot as A, showing the distribution of premotor interneurons following injections of the flexor muscles TA and PL. Correlations and absolute value of Hedges' G coefficients across each experiment are shown in F. G and H shows the mediolateral distribution and the position of the median for each experiment, respectively.

1181

Figure 6: GlyT2^{on} and GlyT2^{off} premotor interneurons are distributed similarly for TA and LG. A, B: Distribution
of GlyT2^{off} (A) and GlyT2^{on} (D) premotor interneurons following LG and TA injections pooled from 4 LG and
3 TA experiments in *GlyT2-EGFP::ChAT-Cre* mice crossed with RΦGT mice, indicating that neither class of

premotor interneurons is segregated across muscles. Boxplots and violin plots (B for GlyT2^{off} and E for
GlyT2^{on}) show uniformity of distribution across experiments in both the dorsal (top) and ventral (bottom)
halves of the cord. C (GlyT2^{off}) and F (GlyT2^{on}) show boxplots and individual values for the medians of the
mediolateral distributions restricted to dorsal (top) or ventral (bottom) part of the cord. Ventral premotor
GlyT2^{off} and GlyT2^{on} interneurons are differentially distributed. Comparison of excitatory and inhibitory
premotor interneurons in LG (G) and TA (J) muscles are similar in the dorsal cord, but differ in the ventral
cord, where most ipsilateral premotor interneurons are GlyT2^{on}, and the majority of contralateral premotor
interneurons are GlyT2^{off}. Boxplots and violin plots of individual experiments are shown in H for LG and K for
TA, highlighting the mediolateral differences in the ventral cord. The medians of the ventral and dorsal
distributions are shown in I for LG and L for TA.
1195

Figure 7: The distribution of flexor and extensor premotor INs is similar in *Olig2::Cre; RΦGT*. A, B: Single lumbar sections form animals injected in the GS (A) or TA (B) muscles. C - E: Overlay of individual GS (C) and TA (D) experiments and pooled experiments (E). F: Correlation coefficients and absolute value of hierarchical bootstrapped Hedges' G effect sizes between injections of different muscles and using a different driver for Cre expression. G: Box and whisker plots of median values of all the medio-lateral distributions.

Figure 8: Injection of AAV-G and RabV into GS and TA muscles does not reveal segregation of premotor interneurons. Example of double infected upper (A and lower (B) lumbar sections. Transverse and longitudinal pooled distributions of premotor interneurons from two experiments are overlapping in all quadrants (C) and individual experiments are highly correlated and have low absolute value of Hedges' G effect sizes (D) and with similar medians in the ipsilateral dorsal quadrant. Scale bars 300 μm.

Figure 9: Injection of AAV-FLEX-G in ChAT::Cre mice gives rise to overlapping distributions of flexor and extensor related premotor interneurons. Extensive labelling is observed in two lower lumbar sections of two GS (A) and TA (B) injected mice. The pooled distributions of flexor and extensor related infected neurons are similar on the transverse and longitudinal planes (C). There is strong correlation and low effect sizes across individual experiments (D) and the medians are not different (E). Scale bars 400 μm.

Figure 10

1216

Figure 10: Injection of AAV-FLEX-G prevents anterograde spreading of RabV from afferent terminals. A: side
by side comparison of pooled transverse distribution of infected interneurons from GS (left) and TA (right)
with AAV-G or AAV-FLEX-G and RabV. The distributions obtained with AAV-G show a marked mediolateral
shift and extensive labelling of superficial dorsal horn neurons, a feature that is absent when anterograde
transfer is prevented by restricting the expression of the G protein to motoneurons only. The distributions
obtained are very similar across the different muscles, but the mediolateral shift is reflected in the lower
correlation values and high hierarchical bootstrapped effect size (B) and difference in medians (C) between
AAV-G and AAV-FLEX-G injections. Scale bars 400 µm.
1225

Figure 11: Injections of PRV-Bartha in GS and TA muscles give rise to similar distribution of premotor interneurons. Example of a lower lumbar section of an animal injected in GS(A) and TA (B). The distribution of premotor interneurons are similar along all axis (C). Individual experiments are highly correlated, with small effect size (D) and their median values along the mediolateral axis are identical for flexor and extensor injections. Scale bars 300 μm.

Figure 12: Regardless of the tracing methods, flexor and extensor related premotor interneurons distributions are always overlapping. The correlation and absolute value of hierarchical bootstrapped Hedges' G coefficient matrix across pooled experiments obtained with different injection strategies exhibits high values of correlation and low effect sizes across different techniques and across the two different muscles (A). The median values of the mediolateral position of premotor interneurons are similar for all conditions (B).

1233

Figure 13

Figure 13: Analysis of ectopic Cre expression in *ChAT-IRES-Cre* and *RCL-tdTom* mice. A: representative lumbar section stained with antibodies against ChAT (green) and tdTom (red) and (B) map of neurons labelled with both or one of the two antibodies in all the 13 analysed sections from 3 mice, showing that some of the tdTom positive neurons do not express ChAT, indicating either a developmental downregulation of ChAT expression or a modest leak in the Cre expression. C: Venn diagram showing the overall number of mapped neurons.

1