Prenatal environmental conditions underlie alternative reproductive tactics that drive the formation of a mixed-kin cooperative society

Shailee S. Shah[1][*] and Dustin R. Rubenstein[1,2]

[1]Department of Ecology, Evolution, and Environmental Biology, Columbia University, 1200 Amsterdam Avenue, New York, NY 10027 USA
[2]Center for Integrative Animal Behavior, Columbia University, New York, NY 10027 USA

*Correspondence to: Shailee S. Shah

Email: shailee.shah@columbia.edu

Author contributions: Both authors designed the study, collected and analyzed the data, and wrote the paper.

Competing Interest Statement: The authors declare no competing interests.

Keywords: alternative reproductive tactics; dispersal; inclusive fitness; cooperative breeding; mixed-kin society; direct benefits

This PDF file includes:
Abstract
Main Text
Figures 1 to 4
Tables 1 to 3

Abstract

Although animal societies often evolve due to limited natal dispersal that results in kin clustering and

facilitates cooperation among relatives, many species form cooperative groups with low and variable kin

structure. Groups in such mixed-kin societies often comprise residents and immigrants of the same sex

that compete for breeding opportunities. To understand how such mixed-kin societies form despite their

potential for social conflict, we investigated the environmental causes and subsequent fitness

consequences of dispersal decisions in male cooperatively breeding superb starlings (*Lamprotornis*

superbus) that live in a climatically unpredictable environment. We show that the two alternative

reproductive tactics—natal dispersal or philopatry—exhibit reproductive tradeoffs resulting in equal

lifetime inclusive fitness. The tactic an individual adopts is governed by the environment its parents

experience prior to laying rather than the environment it experiences during its juvenile stage. When

individuals adopt the tactic not predicted by early life environmental conditions, their fitness is reduced.

Ultimately, climate-driven oscillating selection may help stabilize mixed-kin animal societies, despite

their reduced kin structure and potential for social conflict.

Main

Dispersal decisions that enable individuals to escape unfavorable environmental conditions or reduce

inbreeding risk also influence ecological and evolutionary processes like range expansion, gene flow, and

the formation of animal societies [1,2]. Limited natal dispersal—the permanent movement of an individual

from its natal to breeding site—plays a crucial role in social evolution by creating spatial clustering of kin

that facilitates cooperative care of young by relatives [3–5]. Although most animal societies are therefore

characterized by groups with high kin structure, a surprising number of species form social groups with

low kin structure [6] that consist of same-sex residents and immigrants, which often compete for breeding

opportunities [7–10]. Since indirect benefits alone cannot explain the evolution of these mixed-kin societies

[11–13], direct reproductive or survival benefits may underlie their formation [7,9,14]. Understanding how

mixed-kin cooperative societies form and remain stable despite their reduced kin structure and potential

for high social conflict will require examining the causes and lifetime fitness consequences of individual

dispersal decisions to determine why some individuals remain in their natal groups with kin while others

do not.

Dispersal decisions are influenced not only by aspects of the social or ecological environment that

individuals experience during their lifetime after reaching nutritional independence [15–17], but also by how

these factors affect their parents prior to birth and impact their development early-in-life (termed prenatal

factors) [18,19]. The influence of prenatal factors typically manifests via parental effects, modifications

provided by the mother or father to the offspring during development that alter offspring phenotype [20].

Parental effects may be particularly important in cooperatively breeding species [21], where parents can

limit offspring dispersal by influencing helping behavior or promote dispersal to reduce kin competition

[19,22,23]. Although dispersal in most social species tends to be sex-biased [24], in many cooperative breeders,

individuals of one sex can adopt either of two alternative reproductive tactics: natal dispersal or

philopatry [8,15]. Such sex-specific behavioral polyphenisms can exist either because (i) individuals face

conditional constraints in adopting the tactic with higher fitness and must instead make "the-best-of-a-

bad-job" by choosing the tactic with lower fitness [25,26], or (ii) the relative fitness payoffs are

environmentally-dependent such that both tactics can persist within a population when conditions

fluctuate [27,28]. Consistent with the "best-of-a-bad-job" hypothesis, the fitness benefits of alternative

reproductive tactics in cooperative breeders tend to be unequal in the short-term [29–32]. Yet, few studies

have compared the lifetime inclusive fitness outcomes of alternative reproductive tactics, especially in

mixed-kin societies where they are most common [8]. Thus, the evolutionary mechanism underlying the

formation of mixed-kin societies remains largely unknown and can only be revealed by considering the

lifetime inclusive fitness consequences of alternative reproductive tactics.

Here, we leveraged a 16-year longitudinal dataset from Kenya to examine how reproductive tactics in

male cooperatively breeding superb starlings (*Lamprotornis superbus*) are influenced by ecological and

social factors in both the prenatal and juvenile periods, and ultimately how these alternative tactics impact

lifetime inclusive fitness. Superb starlings are plural cooperative breeders in which both sexes help [33,34],

and although males tend to be more philopatric than females, these mixed-kin social groups have low but

variable kin structure overall as well as within both sexes [33,35]. While resident females never breed in their

natal groups, both resident and immigrant males can acquire breeding status [33]. These savanna-dwelling

birds inhabit one of the most variable and unpredictable environments on earth [36], where annual

fluctuation in seasonal rainfall governs insect prey availability [37]. Rainfall is most variable during the dry

season [38], and this variation has been shown to influence many aspects of superb starling social behavior

and life-history. In particular, rainfall experienced by the parents (prenatal rainfall) influences offspring

sex ratio [34], epigenetic modification of genes related to the avian stress response [39], and male reproductive

decisions [39]. Dry season rainfall experienced by individuals of both sexes also influences access to

breeding opportunities [40] and helping behavior [41]. In addition, the social environment (group size) also

impacts superb starling fitness, both survival [42] and reproductive success [38].

To examine the environmental causes of alternative reproductive tactics in superb starlings, we first

investigated how ecological and social factors during both the prenatal and juvenile stage influence male

dispersal decisions. We predicted that under harsh prenatal environmental conditions (low rainfall),

particularly for small groups with female-skewed sex ratios, parental effects may indirectly limit dispersal

of male offspring by promoting helping behavior. Additionally, males that remain in their natal group

would be more likely to provide alloparental care in their first year of life than males that ultimately

disperse. Similarly, male superb starlings that experience harsh environmental conditions as juveniles, the

life history stage when dispersal decisions are likely made [33], would be more likely to remain in their natal

group in order avoid the costs of dispersal or reap the benefits of philopatry. Furthermore, a social

environment conducive to higher survival (i.e., when in larger groups) and with greater access to

reproductive opportunities (groups with female-skewed sex ratios) may also promote philopatry. Next, to

understand the fitness consequences of these alternative reproductive tactics, we compared access to

reproductive opportunities, reproductive success, lifetime inclusive fitness, and survivorship of resident

and immigrant males. According to the best-of-a-bad-job hypothesis, we predicted that the two alternative

tactics would result in unequal lifetime inclusive fitness. Alternatively, if both dispersal and philopatry

maximize individual fitness, then the two tactics would have equal lifetime inclusive fitness but exhibit

reproductive tradeoffs such that individuals would have lower lifetime fitness when they adopt the

reproductive tactic not predicted by the prevailing environmental conditions. Ultimately, by linking the

environmental causes and fitness consequences of alternative reproductive tactics in a cooperative breeder

we will be able to determine how two fundamentally different reproductive phenotypes can persist and

potentially give rise to mixed-kin, plural breeding societies.

Results

Patterns of male dispersal

Consistent with previous reports in this species [33], we found that males were the primary philopatric sex;

only 59 of 198 males (30%) in our marked social groups for whom we had data on dispersal status (i.e.,

excluding individuals already present in the groups at the beginning of 2001 (dispersal status unknown)

and individuals not seen past one year of age (presumed dispersed)) were immigrants (compared to 121 of

248 (49%) of females). However, among all males for whom we had complete life history data (N = 162),

more than half the breeders (30 of 59; 51%) were immigrants.

Ecological and social predictors of male dispersal

Next, we examined the role of ecological and social factors experienced by males during their juvenile

stage (i.e., the period when dispersal occurs) as well as by their parents (i.e., prenatal) in influencing male

dispersal decisions. We found that harsh prenatal ecological conditions were associated with limited

dispersal such that males were less likely to disperse when they were born in years following low prenatal

rainfall (Z = 2.03, P = 0.04, CI = 0.02-0.74; Fig. 1, Table 1). Since prenatal ecological conditions did not

affect hatchling mass prior to fledging (see *SI*), this pattern was likely driven by dispersal, and not

mortality. In contrast, the social environment as well as the ecological conditions experienced during the

juvenile period had no significant impact on the likelihood of males dispersing (Table 1). Together, these

results suggest that male dispersal is influenced more by (i) conditions experienced by their parents than

the conditions they experience as juveniles and (ii) ecological rather than social factors. Finally, we also

found that males that subsequently dispersed were less likely to act as alloparents in their first year of life

than males that subsequently remained in their natal group ($\chi^2_1 = 7.63$, P = 0.006).

Fitness consequences of male dispersal decisions

We then explored the fitness consequences of the different male dispersal decisions. More than half the

males in the population (85 of 152 males; 56%) failed to accrue any inclusive fitness in their lifetimes

(Fig. S3) [33]. Residents unsurprisingly accrued higher indirect fitness than immigrants (Fig. 2C, U = 2103,

P = 0.02), who were unlikely to accrue meaningful indirect fitness in their natal group prior to dispersal

(see *SI*). In contrast, resident and immigrant males had similar direct and inclusive lifetime fitness (Fig.

2A-B; inclusive: U = 2337, P = 0.22; direct: U = 2701, P = 0.59). While this analysis does not quantify

indirect fitness accrued by immigrants in their natal group prior to dispersing, natal males that dispersed

accrued significantly lower fitness in their first year than residents, suggesting that immigrants into our

study population also did not likely accrue considerable fitness prior to dispersing (*see SI*). Nonetheless,

to account for the possibility that immigrant males accrued at least some indirect fitness in their natal

group prior to dispersal, we repeated our analysis after excluding fitness accrued by residents in their first

year of life. We found qualitatively similar results, with residents and immigrants still having similar

direct and inclusive fitness, as well as similar indirect fitness (see *SI*).

Although the survival likelihood of immigrants after successful dispersal is not significantly different

from that of residents (Z = 0.06, P = 0.95, CI = -0.36-0.38) (Fig. S4), our analysis did not account for any

potential mortality of immigrants during dispersal. While we expect the mortality of immigrants to be low

because there are no floaters in this system[33], to account for potential immigrant mortality during

dispersal, we ran a sensitivity analysis that added immigrants with zero fitness in 10% increments to the

analyses of direct, indirect, and inclusive lifetime fitness. Our sensitivity analysis found that the mortality

threshold for immigrants at which direct fitness would be higher in residents than immigrants was 70%

(Table S1). Similarly, the mortality threshold for immigrants at which inclusive fitness would be higher in

residents than immigrants ranged from 20% to 50%, depending on whether first-year resident males were

included in the analysis (Table S1). Together, these results indicate that resident and immigrant males

have similar lifetime inclusive fitness, even when accounting for the potential of mortality during

dispersal.

Despite similar lifetime inclusive fitness, immigrants were more likely than residents to breed at least

once in their lifetimes ($\chi^2_1 = 7.39$, P = 0.006), had higher lifetime breeding effort (the number of breeding

attempts as a proportion of an individual's adult lifespan) (U = 3869, P = 0.006), and began breeding at a

younger age (U = 265.5, P = 0.01). In contrast, residents that did breed had higher nest success than

immigrant breeders (Z = 2.46, P = 0.01, CI = 0.27-2.34). In addition, both lifetime breeding effort and

dispersal status affected a male's likelihood of accruing some nonzero inclusive fitness during his lifetime

(Fig. 3A, Table 2). Among males with some lifetime inclusive fitness, residents accrued higher inclusive

fitness than immigrants with the same amount of lifetime breeding effort (Fig. 3B, Table 2).

Finally, we assessed whether there were reproductive tradeoffs for individuals that adopted the tactic not

predicted by prenatal ecological conditions. Since we did not have precise rainfall data from the

immigrants' natal sites (though rainfall is highly spatially correlated and temporally synchronous within

the dispersal radius of our study site; see *SI*), we used a categorical measure of prenatal rainfall based on

the mean long-term pre-breeding rainfall at our study site (see *Methods* and *SI*). Consistent with the idea

of reproductive tradeoffs and the hypothesis that the fitness consequences of alternative reproductive

tactics depend on prenatal environmental conditions, we found that immigrant males were more likely to

accrue nonzero lifetime inclusive fitness when born following periods of high prenatal rainfall, but that

residents were more likely to accrue nonzero lifetime inclusive fitness when born following periods of

low prenatal rainfall (Fig. 4, Table 3).

Discussion

Although mixed-kin cooperative groups characterized by low relatedness are surprisingly common (45%

of all cooperatively breeding birds)[6], it remains unclear how these societies arise and are maintained given

the potential for high social conflict among unrelated group members competing for reproductive

opportunities. We examined the environmental causes and inclusive fitness consequences of dispersal

decisions in male superb starlings living in a climatically variable and unpredictable savanna environment

[36,42], for whom philopatry and dispersal can both be pathways to reproductive opportunities [33]. We found

that dispersal in males is influenced by their parents' prenatal environment rather than the environment

that the males experience as juveniles, suggesting that prenatal parental effects play an important role in

governing dispersal decisions. Ultimately, natal dispersal and philopatry had equal lifetime inclusive

fitness outcomes, meaning that they are two equal, alternative male reproductive tactics. Yet, the two

tactics showed reproductive tradeoffs—immigrants had greater access to reproductive opportunities via

the acquisition of dominant breeding positions, whereas residents that bred had higher reproductive

success. When males adopted a tactic not predicted by prevailing prenatal environmental conditions, their

lifetime inclusive fitness was reduced. Together, these results suggest that environmental variability

experienced early in life may help maintain a reproductive polyphenism with equal lifetime inclusive

fitness, which ultimately enables the formation and persistence of a mixed-kin cooperative society in an

unpredictable environment.

Unlike in other cooperatively breeding species where conditions experienced during the juvenile stage

shape dispersal decisions [15,16,43], male dispersal in superb starlings was only governed by prenatal

ecological conditions. Although we were unable to directly distinguish between dispersal and morality,

several lines of evidence—including the modal age at which natal males disappear from and modal age of

immigrants into our study population, the age at which males reach sexual maturity, the close match in

the proportions of dispersers and immigrants, the increased likelihood of dispersal following benign

ecological conditions, no effect of prenatal ecological conditions on hatchling mass prior to fledging, and

the high likelihood of post-fledgling survival from year-to-year [42]—strongly suggest that the pattern we

observed is driven by dispersal, and not mortality (see *Methods and SI* for further details). The effect of

prenatal conditions on dispersal suggests that parental effects influence the dispersal phenotype of male

superb starlings, a result that has been suggested previously for male reproductive opportunities in this

species [39]. Parental effects, which can result from physiological or molecular modifications to offspring

during development [20], may be particularly important in cooperatively breeding species [21] and species

inhabiting unpredictable environments [44]. While the specific physiological mechanisms of prenatal

parental effects on dispersal remain unknown for this species, previous studies have demonstrated an

effect of maternal condition on offspring sex allocation, maternal investment in eggs, and DNA

methylation in superb starlings [34,39] and other vertebrates [23,45,46]. Studies of two other avian species,

western bluebirds (*Sialia mexicana*) and great tits (*Parus major*), found an effect of maternal androgen

deposition on offspring dispersal in response to ecological conditions [18,19]. Viviparous lizards (*Lacerta*

vivipara) show a similar effect [46], and maternal hormones have also been shown to increase helping

behavior in subordinate female meerkats (*Suricata suricatta*) [22]. Alternatively, conditions experienced by

parents may affect their behavior towards their offspring after birth, such that dispersal is driven purely by

behavioral interactions prior to offspring becoming self-sufficient [47,48]. Further work in this and other

cooperatively breeding species should examine the mechanism by which environmentally-mediated

parental effects might influence future reproductive tactics adopted by offspring.

Parental effects can be selfish, to manipulate offspring phenotype in order to increase parental fitness, or

anticipatory, to maximize offspring fitness [21,49]. We know that offspring sex ratio in superb starlings is

male-biased when mothers are in poorer body condition following harsher prenatal conditions, a pattern

that suggests that females maximize their inclusive fitness in an unpredictable environment by investing

in the sex with lower fitness variance [34]. However, our results suggest that mothers could also be

influencing cooperative behavior in their offspring because males that participated in alloparental care in

their first year of life were subsequently more likely to remain in their natal group. Male superb starlings

are more likely to become alloparents and help more than females [38], and alloparents buffer the

detrimental effects of harsh environmental conditions on reproductive success in this species [41]. Thus,

prenatal parental effects promoting alloparental behavior following harsher prenatal conditions might

serve to selfishly increase the parents' short-term future reproductive success and indirectly limit

dispersal of male offspring [22]. Although males born following benign prenatal conditions could,

conversely, be more likely to disperse so as to reduce kin competition, this seems less probable because

kin competition has actually been shown to decrease under benign environmental conditions in other

cooperatively breeding species [50,51]. Additionally, our results are consistent with the hypothesis that

prenatal parental effects on dispersal in males are anticipatory because when males adopted the tactic

predicted by the prevailing prenatal ecological conditions, they were more likely to accrue some inclusive

fitness in their lifetimes. Since more than half of superb starling males fail to accrue any fitness in their

lifetimes, largely due to high nest predation pressure [33], securing any amount of nonzero lifetime fitness is

crucial in this species, an idea consistent with the bet-hedging hypothesis proposed for this [38] and other

social species [52]. Males born following harsh prenatal conditions thus appear to benefit from avoiding the

costs of dispersal and remaining in their natal group [17,53], whereas males born following benign conditions

might be better able to cope with the costs of dispersal and maximize their fitness by immigrating into

another social group [31].

In contrast to other studies in kin-only cooperatively breeding societies [29,30,54–57], we found that the two

reproductive tactics adopted by males—natal dispersal and philopatry—have equal lifetime inclusive

fitness in a mixed-kin cooperative society. Although resident males had significantly higher lifetime

indirect reproductive success than immigrants, an unsurprising result given the kin structuring among

males in superb starling social groups [33], this was more than balanced out by the relatively larger

contribution of lifetime direct fitness, even when accounting for potential mortality during dispersal.

While it was not logistically possible to compare the fitness of resident and dispersing males born to the

same social group, our approach of comparing resident to immigrant males has proven informative in

other avian studies [58]. While we do not have an estimate of dispersal mortality in this species, we have

never observed floaters in our study area, suggesting that individuals move from their natal group to their

non-natal group over a relatively short time period and likely do not incur significant mortality in transit

[33]. Following successful dispersal, our results show that the survival likelihood of immigrants is similar to

that of resident males. Furthermore, natal males that subsequently remain in their natal group accrue

significantly higher indirect fitness in their first year of life, suggesting that immigrants are unlikely to

accrue considerable indirect fitness in their natal group prior to dispersing. Yet, when even accounting for

this possibility, immigrant and resident males still had similar lifetime inclusive fitness. Similar variation

in alloparental effort and indirect fitness in relation to dispersal tactics has been shown in other

cooperatively breeding species [58,59].

Finally, we also show that the equal lifetime inclusive fitness outcomes of the two dispersal tactics are

due to a reproductive tradeoff, an idea that has been proposed theoretically [28,60] but rarely tested

empirically. Although immigrant superb starlings had greater access to reproductive opportunities via the

acquisition of dominant breeding positions with the group, residents that acquired breeding positions were

more likely to successfully fledge young. Residents can therefore afford to have lower access to breeding

opportunities over their lifetimes since they have higher nest success. This higher nest success is likely

due to greater alloparental care at nests of resident males who have more kin in the group to act as helpers

than do immigrants [61–63]. Thus, for superb starling males, philopatry leads to lower reproductive quantity

but higher quality, whereas dispersal results in higher reproductive quantity but lower quality. Since

immigrants and residents are equally likely to survive from one breeding season to the next, these

differences in reproductive quality and quantity are unrelated to longevity [29]. Together these results

suggest that the two alternative dispersal tactics are equivalent in terms of lifetime inclusive fitness and

refute the hypothesis that remaining in the natal group—the tactic favored following harsh prenatal

conditions—is simply making "the-best-of-a-bad-job" [25,26]. The coexistence of resident and immigrant

males in cooperative social groups has been shown to be similarly facilitated by equal reproductive rates

in spotted hyenas (*Crocuta crocuta*) [8] and inclusive fitness at different age stages in dwarf mongooses

(*Helogale parvula*) [64], both species that live in mixed-kin societies. However, to the best of our

knowledge, ours is the first study in a cooperatively breeding vertebrate to demonstrate equal lifetime

inclusive fitness outcomes of alternative dispersal tactics.

This behavioral polyphenism in reproductive tactics is likely maintained by a flexible response to high

and unpredictable environmental variability [27]. A temporally variable and unpredictable environment has

been shown both theoretically [28] and empirically [65] to generate conditional strategies that result in a

developmental switch between alternative tactics. If alternative reproductive tactics have a strong genetic

(or perhaps epigenetic) basis [66,67], environmental variability can reverse the selective differential between

the two tactics from one year to the next, resulting in alternative tactics with equal fitness outcomes [28,68].

The savanna habitat inhabited by superb starlings has high temporal ecological variability [36,38] that may

allow both dispersal tactics to persist, resulting in the formation of mixed-kin cooperative groups.

Interestingly, spotted hyenas [69] and dwarf mongooses [64] experience the same unpredictable African

savanna environments as superb starlings, suggesting that environmental uncertainty—and perhaps

oscillating selection pressures more generally—maintains alternative dispersal tactics and leads to the

formation of and/or stabilizes societies with low kin structure. Oscillating selection may be particularly

important in arid and semi-arid environments where climatic variability is high [68], and may help explain

the evolution of both mixed-kin societies and plural breeding more generally. Indeed, many other plural

cooperatively breeding birds across the globe live in harsh arid and semi-arid environments characterized

by unpredictable variation in rainfall and food resources [13,70–77]. While we do not yet know whether

dispersal has a genetic basis in superb starlings—though there are indications of a potential epigenetic

basis [39]—our results suggest that the two tactics face oscillating selection since the relative lifetime

inclusive fitness of the two tactics fluctuates in response to prenatal environmental conditions. Ultimately,

identifying the mechanism by which prenatal parental effects lead to developmental differences that

determine male reproductive tactics will be important for understanding the role that environmentally-

driven selection pressures play in shaping behavioral polyphenisms in this and other social species.

In summary, we have shown that natal dispersal and philopatry in male cooperatively breeding superb

starlings represent two alternative reproductive tactics with equal lifetime inclusive fitness. The tactics are

likely mediated by parental effects during development and are maintained by oscillating selection

pressures characteristic of their variable and unpredictable savanna environment. Our study suggests a

direct link between environmental uncertainty, behavioral polyphenism, and the evolution of mixed-kin

animal societies that cannot be explained by indirect fitness benefits alone. Our work also underscores the

importance of prenatal environmental conditions and parental effects in determining offspring phenotype,

especially in cooperative societies where early life conditions have direct implications on the future

fitness of both parents and offspring. Moreover, the direct benefits derived from environmental selective

pressures appear to play a significant role in the evolution and maintenance of cooperative societies,

alongside or in the absence of kin selection. Ultimately, understanding how fluctuation in early life

environmental conditions helps mediate reproductive tradeoffs and lifetime fitness is critical in an era of

rapid anthropogenic climate change because climatic uncertainty is only likely to increase across much of

the globe for the foreseeable future.

Materials and Methods

Data collection

Nine superb starling social groups have been monitored continuously since 2001 at the Mpala Research

Centre, Kenya (0° 17'N 37° 52'E) [33,78]. The study population is distributed over a 9 km extent from North

to South. Groups (mean size ± SD = 22 ± 12) defend stable territories year-round and consist of breeding

(mean ± SD = 2.70 ± 1.49 pairs per group) and non-breeding individuals, some of whom act as

alloparents that guard and/or provision young [33,41]. Birds breed twice a year during the long (March-June)

and short rains (October-November) [33]. We used data from the beginning of the 2001 short rains breeding

season through the end of the 2017 long rains breeding season (N = 33 breeding seasons over 16 years),

since not all birds in the study population were banded in the first breeding season. Limited focal

observation data for nests in 2001 short rains was accounted for in our analyses (see *Data Analysis*).

Birds were banded with a unique combination of colored leg bands and a numbered metal ring.

Hatchlings were banded in the nest; all other individuals were captured in baited pull-string traps or mist

nets and banded after fledging from the nest [78]. If banded post-fledgling, age was assessed via eye color as

fledgling (black eyes), sub-adult (less than one year of age; cloudy eyes), or adult (one year of age or

older; white eyes) [42]. Dispersal status (i.e., resident or immigrant) was assigned based on age at banding

and genetic parentage. Residents were either banded as hatchlings in the nest or as juveniles whose

parents were genetically identified as members of the same group using 15 microsatellite markers [79].

Immigrants were defined as those banded as juveniles or older whose parents were not genetically

identified as belonging to the same group. We also used the same microsatellite markers to estimate (i)

parentage and direct reproductive success in Cervus v3.0 [80] using methods described previously for this

species [40,81–83] and (ii) pairwise relatedness [84] between all individuals with the R package *related* [85]. Sex

was determined genetically [86] as previously described for this species [78].

We performed daily nest searches throughout the breeding season. Active nests were observed with a

spotting scope for 60-120 min per observation period (total observation time per nest: mean ± SD =

314.33 ± 248.82 min). All superb starlings within 30 m of the nest were identified, and their times of

arrival and departure were recorded [83]. Parents are the primary nest builders, and only the mother

incubates the eggs [83]. All other members of the social group seen visiting or guarding the nest were

categorized as alloparents [41].

Census data were used to estimate group size and sex ratio (calculated by dividing the number of males

by the sum of number of males and immigrant females in the group to estimate potential mate

competition. Natal females were excluded since they never breed and are thus not viable mates for males

in the group). Groups were opportunistically censused year-round and each individual marked as either

present or absent in its social group twice a year in six-month increments. Individuals not seen for five or

more breeding seasons (i.e. 2.5 years) were assumed to be dead [35,42]. We inferred that the dispersal

window for males is between fledging and around one year of age (age of sexual maturity) using four

lines of evidence: (i) the modal age at which males disappear from their natal group; (ii) the categorical

age of immigrant males dispersing into the study population; (iii) the likelihood of males being detected

in the census in their first year of life; and (iv) the minimum age of first breeding by resident males (see

SI for details). If a male was observed in its natal group after one year of age, we classified it as a

resident; if not, we classified it as having dispersed. Although we could not distinguish between dispersal

and mortality, the positive relationship between the likelihood of dispersal and benign ecological

conditions (see *Results*) suggests that these males did indeed disperse since we would have expected a

negative relationship if mortality were higher following harsh ecological conditions. In addition, we

found no variation in hatchling mass before fledging (measured at 10 days of age) with variation in

ecological conditions, further suggesting that this pattern is driven by dispersal not mortality (see *SI*).

Following the same lines of evidence, immigrant males were assumed to have entered the group at about

one year of age (see *SI*).

We considered the period between end of the breeding season when a male is born to the beginning of the

breeding season one year later (i.e., its juvenile stage) as its dispersal window (see *SI* for more details).

Ecological and social conditions experienced by juveniles in this period were termed juvenile stage

conditions. In contrast, ecological and social conditions experienced by parents in the pre-breeding period

preceding the breeding season of an individual's birth were termed prenatal conditions. A breeding season

was defined as the period between two weeks before the first nest and two weeks after the last nest of the

season. The period between breeding seasons was termed the pre-breeding period. Daily rainfall was

measured using an automated Hydrological Services TB3 Tipping Bucket Rain Gauge at Mpala Research

Centre [87], supplemented by a manual gauge at the same location when the automated gauge failed [38]. To

calculate prenatal and juvenile period rainfall, we summed the rainfall within the prenatal and juvenile

stages of an individual's lifetime, as defined above.

Data analysis

Our dataset comprises individuals that are (i) known to have been born in the groups ("natal"), (ii) known

to have immigrated into the group ("immigrant"), and (iii) were already part of the group in 2001 when

the study population was banded. All individuals are either known to have disappeared (classified as

either "dead" or "dispersed" depending on age at disappearance) or were still present in the study

population in 2017. Natal individuals that did not disperse were termed "residents". Thus, individuals

with "known complete life histories" are natal and immigrant individuals who were classified as dead.

Using data from individuals with known complete life histories as well as natal individuals that are still

alive and present in the study population past one year of age (N_M =198, N_F = 248), we first examined

patterns of male dispersal by quantifying the proportion of immigrant adult males and females in the

study population. We then determined how breeding opportunities were shared between immigrants and

residents by calculating the proportion of immigrant breeders of both sexes in the study population using

data from breeders with known complete life histories (N_M = 59, N_F = 121).

Next, using data from all natal males—including those classified as dispersed or dead as well as those still

present in the study population past the age of 1 year in 2017—we investigated the effect of ecological

and social conditions during the prenatal and juvenile stages on the likelihood of a male dispersing. We

built a generalized linear mixed model (GLMM) with a binomial error structure and "logit" link function.

We used the binomial response of reproductive tactic (1 = dispersed, 0 = remained) as the dependent

variable ($N_{dispersed}$ = 52, $N_{remained}$ = 133). Fixed effects, which were standardized using z-scores [88], included

prenatal and juvenile stage rainfall, prenatal group size, and prenatal sex ratio. Since group size and sex

ratio were highly correlated within a year (group size: Pearson's r = 0.95; sex ratio: Pearson's r = 0.80),

we only included prenatal group size and sex ratio in the final model. All two-way interactions were

included in the model, but later removed if their effect was not significant [89]. Breeding season of birth,

season (long vs. short rains), social group, mother ID, and father ID were included as random effects. All

random effects other than breeding season of birth had variance components equal to zero and were thus

removed from the final model to facilitate the computation of 95% confidence intervals. This had no

effect on the estimates of fixed effects[90]. Furthermore, to investigate the relationship between alloparental

care provided by males prior to reaching one year of age and their subsequent reproductive tactic, we used

a Chi-square test to compare alloparental status (ever alloparent/never alloparent) and reproductive tactic

(dispersed/remained) for all natal males (N = 185).

Finally, we sought to understand the fitness consequences of the alternative reproductive tactics using

data from males with known complete life histories. Although it was not possible to compare the fitness

consequences of the alternative reproductive tactics adopted by males born from same social group (since

dispersers rarely remain within the study population; only 3 males in 16 years), we compared access to

reproductive opportunities, reproductive success, lifetime inclusive fitness, and survivorship of resident

and immigrant males. Previous studies of cooperatively breeding birds have used the same approach [58].

Access to reproductive opportunities was quantified as: (i) the likelihood of a male ever breeding in its

lifetime (Chi-squared test) (N_{Imm} = 59, N_{Res} = 103); (ii) lifetime breeding effort, or the number of breeding

attempts as a proportion of an individual's adult lifespan (Mann-Whitney test with continuity correction)

(N_{Imm} = 59, N_{Res} = 103); and (iii) for males that bred at least once (N_{Imm} = 30, N_{Res} = 29), the age at first

breeding (Mann-Whitney test with continuity correction). To evaluate the likelihood of a male obtaining

any reproductive success in their lifetimes, we modeled the effect of dispersal status on nest success (1 =

succeeded, 0 = failed) using a GLMM with a binomial error structure and "logit" link function ($N_{Succeeded}$

= 49, N_{Failed} = 229). Since some males nested multiple times in their lifetimes, we included individual ID

as a random effect in the model along with social group and breeding season.

Following Green and Hatchwell [58], we calculated lifetime inclusive fitness (I) as the sum of lifetime direct

and indirect fitness according to the equation

$$I = \left[\sum_{j=1}^{J} M_j ER - \frac{\sum_{j=1}^{J}\sum_{n=1}^{N_j}\sum_{m=1}^{M_j} e_{nj}r_{mnj}}{2}\right]_{direct\ fitness} + \left[\sum_{o=1}^{O}\sum_{m=1}^{M_o} e_o\ r_{mo}\right]_{indirect\ fitness} \quad (1),$$

where direct fitness was calculated as the product of the number of fledglings per nest (M_j), paternal care

effort (E = 0.5, the other half being attributed to the mother), and the mean offspring-father relatedness (R

= 0.5) summed over all successful nests (J), minus half of the indirect fitness attributed to alloparents (the

other half being subtracted from the mother's direct fitness). Indirect fitness attributed to alloparents at

each nest was calculated as the product of the mean alloparental effort at a nest per social group per

breeding season (e_{nj}) and the alloparent's relatedness to the fledglings (r_{mnj}), summed over all fledglings

(M_j) and alloparents at the nest (N_j) for all successful nests (J). Alloparents with $r_{mnj} \leq 0$ received no

indirect fitness. Indirect fitness of the focal male was similarly calculated as the product of the mean

alloparental effort (e_o) and its relatedness to the fledglings (r_{mo}), summed over all of the nests he visited in

his lifetime (O). We used population means instead of individual measures of alloparental effort because

total observation time for nests varied. Alloparental effort was calculated as the proportion of time an

individual spent attending a nest (both guarding and bringing food) relative to the length of the

observation period [41].

We compared the lifetime inclusive, direct, and indirect fitness of resident and immigrant males using a

Mann-Whitney test with continuity correction (N_{Imm} = 53, N_{Res} = 99). Since males accrue indirect fitness

as juveniles, we excluded males born in 2001 short rains, the earliest breeding season in our dataset,

during which limited focal observations were conducted at nests. To account for indirect fitness accrued

by immigrant males in their natal group prior to dispersing, we additionally (i) compared fitness of

residents and immigrants after excluding fitness accrued by residents during the first year of their life (see

SI), and (ii) compared the fitness accrued in the first year of life by natal males that remained in their natal

group to that accrued by natal males that subsequently dispersed (see *SI*). To account for mortality during

dispersal, we conducted a sensitivity analysis by adding incremental proportions of immigrants with zero

lifetime fitness to the dataset (see *SI*).

To determine whether differences in reproductive access and success between resident and immigrant

males affected their lifetime inclusive fitness, we used GLMMs to model the effect of lifetime breeding

effort and dispersal status on lifetime inclusive fitness of males. The first model had a binomial response

(zero/nonzero lifetime inclusive fitness) and a "logit" link function ($N_{Imm} = 53$, $N_{Res} = 99$). The second

model had a continuous dependent variable of all nonzero inclusive fitness observations ($N_{Imm} = 20$, N_{Res}

$= 47$) with an "identity" link function and a gamma error distribution. The fixed effects for both models

were lifetime breeding effort and dispersal status. Social group was included as a random effect, but later

removed from the second model, where its variance component equaled zero, to facilitate the computation

of 95% confidence intervals [90].

We also built a GLMM to investigate the lifetime fitness consequences of adopting the tactic not

predicted by the prevailing environmental conditions, with a binomial response variable (zero/nonzero

lifetime inclusive fitness) and prenatal rainfall and dispersal status as fixed effects ($N_{Imm} = 53$, $N_{Res} = 99$).

Although we did not have precise rainfall data from the immigrants' natal sites, we used monthly rainfall

data from three sites within the dispersal radius (~30 km) (Shah and Rubenstein, unpubl. data) of our

study population (Mpala Research Centre, MRC)—UHURU Central (12 km from MRC) [91], UHURU

North (20 km from MRC) [91], and Nanyuki (38 km from MRC) (East African Livestock Early Warning

System)—to demonstrate high spatiotemporal correlation of prenatal rainfall for immigrant and resident

males (Table S1). Nonetheless, to take a more conservative approach, we used a categorical measure

based on mean long-term pre-breeding rainfall at Mpala Research Centre (low prenatal rainfall: rainfall $\leq$

long-term mean; high prenatal rainfall: rainfall > long-term mean). Rainfall within 10 mm of the long-

term mean was categorized as "low" to account for the accuracy of the rainfall gauge [87].

Finally, we used a time-varying Cox proportional hazard model to determine whether dispersal status

affected male survival (N_{Imm} = 59, N_{Res} = 103) [42]. The model was built using the R package *survival* [92].

We checked that our dataset did not violate the proportional hazard assumption using the "cox.zph"

function. Social group was included as a random factor. We performed all data analysis in R [93]. VIF < 2

for all fixed effects in GLMMs, excluding interaction terms [94].

Data Availability. Data and R code will be made available on Dryad.

Acknowledgements. We thank B. Heidinger, M. Uriarte, M. VanAcker, J. Jensen, P. Downing, S. Caro,

S. Siller, Y. Cheng, A. Earl, J. Guan, S. Guindre-Parker, Y. Haba, E. Greig, R. Gloag, M. Paquet, and an

anonymous reviewer for statistical advice and comments on the manuscript. We are grateful to W.

Watetu, G. Manyaas and J. Mosiany for their assistance in the field, as well as all other volunteers,

students and field assistants that have contributed to this project since its inception in 2001.

References

1.   Matthysen, E. Multicausality of dispersal : a review. in *Dispersal Ecology and Evolution* (Oxford

488   University Press, 2012).

489   2.   Taylor, P. D. Altruism in viscous populations - an inclusive fitness model. *Evol. Ecol.* **6**, 352–356

490   (1992).

491   3.   Emlen, S. T. Benefits, constraints and the evolution of the family. *Trends Ecol. Evol.* **9**, 282–285

492   (1994).

493   4.   Hamilton, W. D. The Genetical Evolution of Social Behavior II. *J. Theor. Biol.* **7**, 1–52 (1964).

494   5.   Hamilton, W. D. The genetical evolution of social behaviour. I. *J. Theor. Biol.* **7**, 1–16 (1964).

495   6.   Riehl, C. P. Evolutionary routes to non-kin cooperative breeding in birds. *Proc. R. Soc. London B*

496   *Biol. Sci.* **280**, 20132245 (2013).

497   7.   Baglione, V., Marcos, J. M., Canestrari, D. & Ekman, J. Direct fitness benefits of group living in a

498   complex cooperative society of carrion crows, Corvus corone corone. *Anim. Behav.* **64**, 887–893

499   (2002).

500   8.   Davidian, E., Courtiol, A., Wachter, B., Hofer, H. & Höner, O. P. Why do some males choose to

501   breed at home when most other males disperse? *Sci. Adv.* **2**, e1501236 (2016).

502   9.   Kaiser, S. A., Martin, T. E., Oteyza, J. C., Armstad, C. & Fleischer, R. C. Direct fitness benefits

503   and kinship of social foraging groups in an Old World tropical babbler. *Behav. Ecol.* **29**, 468–478

504   (2018).

505   10.   Reyer, H. U. Flexible helper structure as an ecological adaptation in the pied kingfisher (Ceryle

506   rudis rudis L.). *Behav. Ecol. Sociobiol.* **6**, 219–227 (1980).

507   11.   Rubenstein, D. R. & Abbot, P. The evolution of social evolution. in *Comparative Social Evolution*

508   (eds. Rubenstein, D. R. & Abbot, P.) 1–18 (Cambridge University Press, 2017).

509   doi:10.1017/9781107338319.002.

510   12.   Liu, M. *et al.* The genetic relatedness in groups of joint-nesting Taiwan yuhinas: Low genetic

511   relatedness with preferences for male kin. *PLoS One* **10**, 1–15 (2015).

512 13. Riehl, C. P. & Jara, L. Natural history and reproductive biology of the communally breeding

513   Greater Ani (Crotophaga major) at Gatún Lake, Panama. *Wilson J. Ornithol.* **121**, 679–687 (2009).

514 14. Clutton-Brock, T. H. Breeding together: kin selection and mutualism in cooperative vertebrates.

515   *Science (80-.).* **296**, 69–72 (2002).

516 15. Nelson-Flower, M. J., Wiley, E. M., Flower, T. P. & Ridley, A. R. Individual dispersal delays in a

517   cooperative breeder: Ecological constraints, the benefits of philopatry and the social queue for

518   dominance. *J. Anim. Ecol.* (2018) doi:10.1111/1365-2656.12814.

519 16. Baglione, V., Canestrari, D., Marcos, J. M. & Ekman, J. Experimentally increased food resources

520   in the natal territory promote offspring philopatry and helping in cooperatively breeding carrion

521   crows. *Proc. R. Soc. B Biol. Sci.* **273**, 1529–1535 (2006).

522 17. Eikenaar, C., Brouwer, L., Komdeur, J. & Richardson, D. S. Sex biased natal dispersal is not a

523   fixed trait in a stable population of Seychelles warblers. *Behaviour* **147**, 1577–1590 (2010).

524 18. Tschirren, B., Fitze, P. S. & Richner, H. Maternal modulation of natal dispersal in a passerine bird:

525   An adaptive strategy to cope with parasitism? *Am. Nat.* **169**, 87–93 (2007).

526 19. Duckworth, R. A. Maternal effects and range expansion: A key factor in a dynamic process?

527   *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 1075–1086 (2009).

528 20. Mousseau, T. A. & Fox, C. W. The adaptive significance of maternal effects. *Trends Ecol. Evol.*

529   **13**, 403–407 (1998).

530 21. Russell, A. F. & Lummaa, V. Maternal effects in cooperative breeders: From hymenopterans to

531   humans. *Philos. Trans. R. Soc. B Biol. Sci.* **364**, 1143–1167 (2009).

532 22. Dantzer, B. *et al.* The development of individual differences in cooperative behaviour: Maternal

533   glucocorticoid hormones alter helping behaviour of offspring in wild meerkats. *Philos. Trans. R.*

534   *Soc. B Biol. Sci.* **374**, (2019).

535 23. Silk, J. B. & Brown, G. R. Local resource competition and local resource enhancement shape

536   primate birth sex ratios. *Proc. R. Soc. B Biol. Sci.* **275**, 1761–1765 (2008).

537 24. Greenwood, P. J. & Harvey, P. H. The natal and breeding dispersal of birds. *Annu. Rev. Ecol. Syst.*

1–21 (1982).

25.   Koprowski, J. L. Alternative reproductive tactics in male eastern gray squirrels: 'making the best

of a bad job'. *Behav. Ecol.* **4**, 165–171 (1993).

26.   Dawkins, R. Good strategy or evolutionarily stable strategy. in *Sociobiology: Beyond*

nature/nurture (eds. Barlow, G. W. & Silverberg J.) 331–367 (Westview Press, 1980).

27.   West Eberhard, M. J. Phenotypic Plasticity and the Origins of Diversity. *Annu. Rev. Ecol. Syst.* **20**,

249–278 (1989).

28.   Moran, N. A. The evolutionary maintenance of alternative phenotypes. *Am. Nat.* **139**, 971–989

(1992).

29.   Walters, J. R., Doerr, P. D. & Carter, J. H. Delayed dispersal and reproduction as a life-history

tactic in cooperative breeders: fitness calculations from red-cockaded woodpeckers. *Am. Nat.* **139**,

623–643 (1992).

30.   Ekman, J., Eggers, S., Griesser, M. & Tegelström, H. Queuing for preferred territories: Delayed

dispersal of Siberian jays. *J. Anim. Ecol.* **70**, 317–324 (2001).

31.   Bergmüller, R., Heg, D. & Taborsky, M. Helpers in a cooperatively breeding cichlid stay and pay

or disperse and breed, depending on ecological constraints. *Proc. R. Soc. B Biol. Sci.* **272**, 325–331

(2005).

32.   Kingma, S. A., Bebbington, K., Hammers, M., Richardson, D. S. & Komdeur, J. Delayed dispersal

and the cost and benefits of different routes to independent breeding in a cooperative breeding

bird. *Evolution (N. Y).* **70**, 2595–2610 (2016).

33.   Rubenstein, D. R. Superb starlings: cooperation and conflict in an unpredictable environment. in

Cooperative Breeding in Vertebrates: Studies of Ecology, Evolution, and Behavior (eds. Koenig

Walter, D. & Dickinson, J. L.) 181–196 (Cambridge University Press, 2016).

34.   Rubenstein, D. R. Temporal but Not Spatial Environmental Variation Drives Adaptive Offspring

Sex Allocation in a Plural Cooperative Breeder. *Am. Nat.* **170**, 155 (2007).

35.   Pollack, L. & Rubenstein, D. R. The fitness consequences of kin-biased dispersal in a

cooperatively breeding bird. *Biol. Lett.* **11**, 20150336 (2015).

36.   Rubenstein, D. R. & Lovette, I. J. Temporal Environmental Variability Drives the Evolution of

Cooperative Breeding in Birds. *Curr. Biol.* **17**, 1414–1419 (2007).

37.   Rubenstein, D. R. Territory quality drives intraspecific patterns of extrapair paternity. *Behav. Ecol.*

18, 1058–1064 (2007).

38.   Rubenstein, D. R. Spatiotemporal environmental variation, risk aversion, and the evolution of

cooperative breeding as a bet-hedging strategy. *Proc. Natl. Acad. Sci.* **108**, 10816–10822 (2011).

39.   Rubenstein, D. R., Skolnik, H., Berrio, A. & Frances, A. Sex-specific fitness effects of

unpredictable early life conditions are associated with DNA methylation in the avian

glucocorticoid receptor. *Mol. Ecol.* **25**, 1714–1728 (2016).

40.   Rubenstein, D. R. Stress hormones and sociality: integrating social and environmental stressors.

Proc. R. Soc. B Biol. Sci. **274**, 967–975 (2007).

41.   Guindre-Parker, S. & Rubenstein, D. R. Multiple benefits of alloparental care in a fluctuating

environment. *R. Soc. Open Sci.* **5**, 172406 (2018).

42.   Guindre-Parker, S. & Rubenstein, D. R. Survival benefits of group living in a fluctuating

environment. *Am. Nat.* **195**, 1027–1036 (2020).

43.   Suh, Y. H., Pesendorfer, M. B., Tringali, A., Bowman, R. & Fitzpatrick, J. W. Investigating social

and environmental predictors of natal dispersal in a cooperative breeding bird. *Behav. Ecol.* **31**,

692–701 (2020).

44.   Meylan, S., Miles, D. B. & Clobert, J. Hormonally mediated maternal effects, individual strategy

and global change. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 1647–1664 (2012).

45.   Van Petegem, K. H. P. *et al.* Empirically simulated spatial sorting points at fast epigenetic changes

in dispersal behaviour. *Evol. Ecol.* **29**, 299–310 (2015).

46.   Meylan, S., Belliure, J., Clobert, J. & De Fraipont, M. Stress and body condition as prenatal and

postnatal determinants of dispersal in the common lizard (Lacerta vivipara). *Horm. Behav.* **42**,

319–326 (2002).

47.   Cant, M. A., Otali, E. & Mwanguhya, F. Eviction and dispersal in co-operatively breeding banded

591   mongooses (Mungos mungo). *J. Zool.* **254**, 155–162 (2001).

592   48.   Nelson-Flower, M. J. & Ridley, A. R. Nepotism and subordinate tenure in a cooperative breeder.

593   *Biol. Lett.* **12**, (2016).

594   49.   Marshall, D. J. & Uller, T. Why is a maternal effect adaptive? *Oikos* **116**, 1957–1963 (2007).

595   50.   Korb, J. & Schmidinger, S. Help or disperse? Cooperation in termites influenced by food

596   conditions. *Behav. Ecol. Sociobiol.* **56**, 89–95 (2004).

597   51.   Nichols, H. J., Bell, M. B. V., Hodge, S. J. & Cant, M. A. Resource limitation moderates the

598   adaptive suppression of subordinate breeding in a cooperatively breeding mongoose. *Behav. Ecol.*

599   **23**, 635–642 (2012).

600   52.   Kennedy, P., Higginson, A. D., Radford, A. N. & Sumner, S. Altruism in a volatile world. *Nature*

601   **555**, 359–362 (2018).

602   53.   Ligon, J. D., Stacey, P. B. & Ligon, J. D. The Benefits-of-Philopatry Hypothesis for the Evolution

603   of Cooperative Breeding: Variation in Territory Quality and Group Size Effects. *Am. Nat.* **137**,

604   831–846 (1991).

605   54.   Ligon, J. D. & Ligon, S. H. The cooperative breeding behavior of the green woodhoopoe. *Sci. Am.*

606   **247**, 126–134 (1982).

607   55.   Sparkman, A. M., Adams, J. R., Steury, T. D., Waits, L. P. & Murray, D. L. Direct fitness benefits

608   of delayed dispersal in the cooperatively breeding red wolf (Canis rufus). *Behav. Ecol.* **22**, 199–

609   205 (2011).

610   56.   MacColl, A. D. C. & Hatchwell, B. J. Determinants of lifetime fitness in a cooperative breeder, the

611   long-tailed tit Aegithalos caudatus. *J. Anim. Ecol.* **73**, 1137–1148 (2004).

612   57.   Antunes, D. F. & Taborsky, B. Early social and ecological experience triggers divergent

613   reproductive investment strategies in a cooperative breeder. *Sci. Rep.* **10**, 1–8 (2020).

614   58.   Green, J. P. & Hatchwell, B. J. Inclusive fitness consequences of dispersal decisions in a

615   cooperatively breeding bird, the long-tailed tit (Aegithalos caudatus). *Proc. Natl. Acad. Sci.*

201815873 (2018) doi:10.1073/PNAS.1815873115.

59.   Clutton-Brock, T. H. *et al.* Evolution and development of sex differences in cooperative behavior

in meerkats. *Science (80-.).* **297**, 253–256 (2002).

60.   Stearns, S. C. Trade-Offs in Life-History Evolution. *Funct. Ecol.* **3**, 259–268 (1989).

61.   Wright, J., McDonald, P. G., Te Marvelde, L., Kazem, A. J. N. & Bishop, C. M. Helping effort

increases with relatedness in bell miners, but 'unrelated' helpers of both sexes still provide

substantial care. *Proc. R. Soc. B Biol. Sci.* **277**, 437–445 (2010).

62.   Emlen, S. T. & Wrege, P. H. The role of kinship in helping decisions among white-fronted bee-

eaters. *Behav. Ecol. Sociobiol.* **23**, 305–315 (1988).

63.   Komdeur, J. The effect of kinship on helping in the cooperative breeding Seychelles warbler

(Acrocephalus sechellensis). *Proc. R. Soc. B Biol. Sci.* **256**, 47–52 (1994).

64.   Creel, S. R. & Rabenold, K. N. Inclusive fitness and reproductive strategies in dwarf mongooses.

Behav. Ecol. **5**, 339–348 (1994).

65.   Lampert, K. P. & Linsenmair, K. E. Alternative life cycle strategies in the West African reed frog

Hyperolius nitidulus: The answer to an unpredictable environment? *Oecologia* **130**, 364–372

(2002).

66.   Beldade, P., Mateus, A. R. A. & Keller, R. A. Evolution and molecular mechanisms of adaptive

developmental plasticity. *Mol. Ecol.* **20**, 1347–1363 (2011).

67.   Saastamoinen, M. *et al.* Genetics of dispersal. *Biol. Rev.* **93**, 574–599 (2018).

68.   Gibbs, H. L. & Grant, P. R. Oscillating selection on Darwin's finches. *Nature* **327**, 511–513

(1987).

69.   Holekamp, K. E. & Smale, L. Dispersal status influences hormones and behavior in the male

spotted hyena. *Horm. Behav.* **33**, 205–216 (1998).

70.   Clarke, M. F. Co-operative breeding by the Australian Bell Miner Manorina melanophrys Latham:

A test of kin selection theory. *Behav. Ecol. Sociobiol.* **14**, 137–146 (1984).

71.   Russell, A. F., Portelli, D. J., Russell, D. J. F. & Barclay, H. Breeding ecology of the Chestnut-

crowned Babbler: A cooperative breeder in the desert. *Emu* **110**, 324–331 (2010).

72.   Brown, J. L., Brown, E. R., Sedransk, J. & Ritter, S. Dominance, age, and reproductive success in

a complex society: A long-term study of the Mexican Jay. *Auk* **114**, 279–286 (1997).

73.   Hegner, R. E., Emlen, S. T. & Demong, N. J. Spatial organization of the white-fronted bee•eater.

264–266 (1982).

74.   Rowley, I., Russell, E., Payne, R. B. & Payne, L. L. Plural breeding in the Splendid Fairy-wren,

Malurus splendens (Aves: Maluridae), a cooperative breeder. *Ethology* **83**, 229–247 (1989).

75.   Curry, R. L. & Grant, P. R. Galapagos mockingbirds: territorial cooperative breeding in a

climatically variable environment. in *Cooperative breeding in birds* (eds. Stacey, P. B. & Koenig,

W. D.) 291–331 (Cambridge University Press, 1990).

76.   Rowley, I. Communal Activities Among White-Winged Choughs Corcorax Melanorhamphus. *Ibis*

(Lond. 1859). **120**, 178–197 (1978).

77.   Veltman, C. J. Flock, pair and group living lifestyles without cooperative breeding by Australian

Magpies Gymnorhina tibicen. *Ibis (Lond. 1859).* **131**, 601–608 (2008).

78.   Rubenstein, D. R. The evolution of the social and mating systems of the plural cooperatively

breeding superb starling, Lamprotornis Superbus. (Cornell University, 2006).

79.   Rubenstein, D. R. Isolation and characterization of polymorphic microsatellite loci in the plural

cooperatively breeding superb starling, Lamprotornis superbus. *Mol. Ecol. Notes* **5**, 739–744

(2005).

80.   Kalinowski, S. T., Taper, M. L. & Marshall, T. C. Revising how the computer program CERVUS

accommodates genotyping error increases success in paternity assignment. *Mol. Ecol.* **16**, 1099–

1106 (2007).

81.   Weinman, L. R., Solomon, J. W. & Rubenstein, D. R. A comparison of single nucleotide

polymorphism and microsatellite markers for analysis of parentage and kinship in a cooperatively

breeding bird. *Mol. Ecol. Resour.* **15**, 502–511 (2015).

82.   Apakupakul, K. & Rubenstein, D. R. Bateman's principle is reversed in a cooperatively breeding

bird. *Biol. Lett.* **11**, (2015).

83.   Rubenstein, D. R. Female extrapair mate choice in a cooperative breeder: trading sex for help and

increasing offspring heterozygosity. *Proc. R. Soc. B Biol. Sci.* **274**, 1895–1903 (2007).

84.   Queller, D. C. & Goodnight, K. F. Estimating Relatedness Using Genetic Markers. *Evolution (N.*

Y). **43**, 258 (1989).

85.   Pew, J., Wang, J., Muir, P. & Frasier, T. related: an R package for analyzing pairwise relatedness

data based on codominant molecular markers. (2015).

86.   Griffiths, R., Double, M. C., Orr, K. & Dawson, R. J. G. A DNA test to sex most birds. *Mol. Ecol.*

7, 1071–1075 (1998).

87.   Caylor, K. K., Gitonga, J. & Martins, D. J. Mpala Research Centre Meteorological and

Hydrological Dataset [Datafile]. (2017).

88.   Schielzeth, H. Simple means to improve the interpretability of regression coefficients. *Methods*

Ecol. Evol. **1**, 103–113 (2010).

89.   Engqvist, L. The mistreatment of covariate interaction terms in linear model analyses of

behavioural and evolutionary ecology studies. *Anim. Behav.* **70**, 967–971 (2005).

90.   Pasch, B., Bolker, B. M. & Phelps, S. M. Interspecific dominance via vocal interactions mediates

altitudinal zonation in neotropical singing mice. *Am. Nat.* **182**, (2013).

91.   Goheen, J. R. *et al.* Piecewise Disassembly of a Large-Herbivore Community across a Rainfall

Gradient: The UHURU Experiment. *PLoS One* **8**, e55192 (2013).

92.   Therneau, T. A Package for Survival Analysis in R. (2020).

93.   R Core Team. R: A language and environment for statistical computing. (2019).

94.   Fox, J. & Monette, G. Generalized collinearity diagnostics. *J. Am. Stat. Assoc.* **87**, 178–183

(1992).

95.   Bourne, A. R., Cunningham, S. J., Spottiswoode, C. N. & Ridley, A. R. High temperatures drive

offspring mortality in a cooperatively breeding bird. *Proc. R. Soc. B Biol. Sci.* **287**, 20201140

(2020).

96.   Ridley, A. R., Wiley, E. M., Bourne, A. R., Cunningham, S. J. & Nelson-Flower, M. J.

Understanding the potential impact of climate change on the behavior and demography of social

species: The pied babbler (Turdoides bicolor) as a case study. in *Advances in the Study of*

Behavior 1–42 (Elsevier Inc., 2021). doi:10.1016/bs.asb.2021.03.005.

 Figures

Figure 1. Effect of prenatal rainfall on the probability of male superb starlings dispersing from

their natal groups. Males were more likely to disperse when born following periods of higher prenatal

rainfall (N = 185, P = 0.04). Model estimate (solid line) is bounded by the 95% confidence interval

(shaded areas). Circles indicate raw data (blue = dispersed, orange = remained).

Figure 2. Mean lifetime fitness of immigrant and resident male superb starlings. Immigrant and

resident males (N_{Imm} = 59, N_{Res} = 99) have equal lifetime (A) inclusive (P = 0.22) and (B) direct fitness (P

= 0.59), though residents have higher (C) indirect fitness (P = 0.02). Black dots represent means and error

bars denote standard deviation. Shaded areas are kernel probability densities illustrating the distribution of

the data. Asterisks indicate significance (ns: P > 0.05; *: P < 0.05).

Figure 3. Effect of reproductive tradeoffs on lifetime inclusive fitness of male superb starlings. The

effect of lifetime breeding effort and dispersal status (immigrants = blue, triangles; residents = orange,

circles) on (A) the probability of accruing any nonzero lifetime inclusive fitness (N = 152), and (B) the

value of nonzero lifetime inclusive fitness (N = 67). For visualization purposes, we excluded one

individual with lifetime breeding effort greater than three standard deviations above the mean; this

individual was included in the statistical analysis, though excluding it did not alter the results. Lifetime

breeding effort (i.e., the number of breeding attempts as a proportion of an individual's adult lifespan) (P

= 0.001) and dispersal status (P = 0.04) affected a male's likelihood of accruing some nonzero inclusive

fitness during his lifetime. Among males with some lifetime inclusive fitness, residents accrued higher

inclusive fitness than immigrants for the same amount of lifetime breeding effort (P = 0.04). Model

estimates (solid lines) are bound by 95% confidence intervals (shaded areas). Points indicate raw data.

Dashed vertical lines indicate mean lifetime breeding effort.

Figure 4. Fitness consequences of male superb starlings adopting the tactic not predicted by the prevailing prenatal environmental conditions. The effect of prenatal rainfall and dispersal status (immigrants = blue, residents = orange) on the probability of accruing any nonzero lifetime inclusive fitness (fill = nonzero fitness, no fill = zero fitness) (N = 152). Immigrants were more likely to accrue nonzero lifetime inclusive fitness when born following periods of high prenatal rainfall, whereas residents were more likely to accrue nonzero lifetime inclusive fitness when born following periods of low prenatal rainfall (P = 0.02).

Table 1. Factors affecting dispersal decisions of natal male superb starlings. Results of a generalized

linear mixed model with a binomial response (dispersed vs. remained) for natal males (N = 185). Since

sex ratio and group size are highly correlated within a year, we only included the former in the final

model. Breeding season was included as a random effect. Random effects of social group, father ID,

mother ID, and season (short vs. long rains) had variance components equal to zero, and were thus

excluded from the final model to facilitate the computation of 95% confidence intervals.

Effect	Estimate	Std. Error	Z value	95% CI		p-value
Intercept	-0.97	0.18	-5.45	-1.36	-0.60	<0.001
Prenatal rainfall	0.36	0.18	2.03	0.02	0.74	0.04
Prenatal sex ratio	-0.13	0.20	-0.68	-0.54	0.24	0.50
Prenatal group size	0.05	0.19	0.27	-0.34	0.44	0.78
Juvenile rainfall	-0.12	0.17	-0.69	-0.48	0.24	0.49

Table 2. Effects of lifetime breeding effort and dispersal status on lifetime inclusive fitness of male

superb starlings. Results of generalized linear mixed models with a (A) binomial (zero vs. nonzero) (N =

152) and (B) continuous, positive (N = 67) response variable for lifetime inclusive fitness of males. Social

group was included as a random effect in the first model, but since it explained none of the variance and

hindered the calculation of 95% confidence intervals for the fixed effects, we removed it from the second

model.

(A) Response variable: binomial (zero/nonzero) lifetime inclusive fitness

Effect	Estimate	Std. Error	Z value	95% CI		p-value
Intercept	-1.10	0.35	-3.14	-1.86	-0.44	0.002
Lifetime breeding effort	1.03	0.31	3.28	0.46	1.72	0.001
Dispersal status	0.78	0.39	2.02	0.01	1.56	0.04

(B) Response variable: continuous, positive lifetime inclusive fitness

Effect	Estimate	Std. Error	t value	95% CI		p-value
Intercept	0.02	0.01	2.04	0.01	0.06	0.04
Lifetime breeding effort	0.55	0.15	3.73	0.33	0.95	<0.001
Dispersal status	0.04	0.02	2.27	-0.00	0.08	0.03

754 **Table 3. Effects of male superb starlings adopting the tactic not predicted by the prevailing**

755 **prenatal ecological conditions on the likelihood of gaining nonzero inclusive fitness during their**

756 **lifetimes**. Results of a generalized linear mixed model with a binomial response (zero vs. nonzero lifetime

757 inclusive fitness) variable for males (N = 152). Social group was included as a random effect.

Effect	Estimate	Std. Error	Z value	95% CI		p-value
Intercept	-0.10	0.46	-0.23	-1.03	0.80	0.82
Categorical prenatal rainfall	-0.63	0.59	-1.08	-0.80	0.52	0.28
Dispersal status	-0.84	0.64	-1.31	-2.13	0.40	0.19
Categorical prenatal rainfall x Dispersal status	1.74	0.77	2.25	0.24	3.29	0.02