Postpartum corticosterone and fluoxetine shift the tryptophan-kynurenine pathway in dams

Wansu Qiu[1], Kimberly A. Go[2], Yvonne Lamers[3], Liisa A. M. Galea[1,2,4*]

[1]Graduate Program in Neuroscience, [2]Department of Psychology, [3]Food Nutrition and Health Program, Faculty of Land and Food Systems, and [4]Djavad Mowafaghian Centre for Brain Health

University of British Columbia

*Corresponding Author:

2215 Wesbrook Mall

Vancouver, BC, Canada

V6T 1Z3

liisa.galea@ubc.ca

Highlights

- Tryptophan-kynurenine pathway (TKP) is altered by postpartum corticosterone (CORT)

- Postpartum CORT increased neurotoxic metabolites (3-HK, 3-HAA)

- Postpartum fluoxetine (FLX) increased xanthurenic acid concentrations

- Postpartum CORT and FLX together shifted the TKP balance towards neurotoxicity

Abstract

Perinatal depression (PND) affects 15% of mothers. Selective serotonin reuptake inhibitors

(SSRIs) are currently the first-line of treatment for PND, but are not always efficacious.

Previously, we found significant reductions in plasma tryptophan concentrations and higher

hippocampal proinflammatory cytokine, IL-1β levels, due to maternal SSRI treatment. Both

inflammation and tryptophan-kynurenine metabolic pathway (TKP) are associated with SSRI

efficacy in individuals with major depressive disorder (MDD). TKP is divided into

neuroprotective and neurotoxic pathways. Higher metabolite concentrations of the neurotoxic

pathway are associated with depression onset and implicated in SSRI efficacy. Metabolites in

TKP were investigated in a rodent model of *de novo* postpartum depression (PPD) given

treatment with the SSRI, fluoxetine (FLX). Dams were administered corticosterone (CORT)

(40mg/kg, s.c.), and treated with the SSRI, fluoxetine (FLX) (10mg/kg, s.c.), during the

postpartum for 22 days after parturition. Plasma TKP metabolite concentrations were quantified

on the last day of treatment. Maternal postpartum CORT increased neurotoxic metabolites and

co-enzyme/cofactors in dams (3-hydroxykynurenine, 3-hydroxyanthranilic acid, vitamin B2,

flavin adenine dinucleotide). The combination of both CORT and FLX shifted the

neuroprotective-to-neurotoxic ratio towards neurotoxicity. Postpartum FLX decreased plasma

xanthurenic acid concentrations. Together, our data indicate higher neurotoxic TKP expression

due to maternal postpartum CORT treatment, similar to clinical presentation of MDD. Moreover,

maternal FLX treatment showed limited efficacy to influence TKP metabolites, which may

correspond to its limited efficacy to treat depressive-like endophenotypes. Overall suggesting

changes in TKP may be used as a biomarker of *de novo* PPD and antidepressant efficacy and

targeting this pathway may serve as a potential therapeutic target.

Keywords: Postpartum depression, antidepressants, fluoxetine, tryptophan-kynurenine pathway

1. Introduction

Perinatal depression (PND) is a heterogeneous disease, in which symptoms and remission vary depending on depression onset timing and depression history (reviewed in Qiu et al., 2020). Early postpartum is a period of high susceptibility for first time psychiatric disease in females but not in males (Munk-Olsen et al., 2006). Postpartum depression (PPD) affects 15% of females with approximately 40% of these cases being individuals experiencing depressive symptoms for the first time (Wisner et al., 2013). Maternal postpartum corticosterone (CORT) is used to model *de novo* PPD in rodents (reviewed in Qiu et al., 2020).

Inflammation and altered metabolism in the tryptophan-kynurenine pathway (TKP) are associated with major depressive disorder (MDD) and PND (Corwin et al., 2008; Maes et al., 2011; Haapakoski et al., 2015). The immune system and TKP are related as proinflammatory cytokines, such as interferon (IFN)-γ and interleukin (IL)-1β, can increase enzymatic conversion of tryptophan into its downstream metabolite, kynurenine (reviewed by Maes et al., 2011). Higher expression of proinflammatory cytokines such as IL-6, tumor necrosis factor (TNF)-α, IFN-γ, and IL-1β are associated with both MDD and PND (Corwin et al., 2008; Haapakoski et al., 2015). TKP can be divided into either a "neurotoxic" pathway or a "neuroprotective" pathway and higher activation in the neurotoxic branch is seen in MDD and PND (see Figure 1A; reviewed in Maes et al., 2011). Previously, we found reduced cytokine (TNF-α and IFN-γ) expression and elevated concentrations of a TKP enzyme cofactor (vitamin B6) due to postpartum CORT (Qiu et al., 2020a), indicating an altered immune system and TKP in this model of *de novo* PPD.

Selective serotonin reuptake inhibitors (SSRIs) are the first line treatment for PND

although efficacy of SSRIs with PND is equivocal (reviewed in Qiu et al., 2020). Both

inflammation and TKP are implicated in antidepressant efficacy both in humans (Syed et al.,

2018; Sun et al., 2020) and in rodents (Qiu et al., 2020a). Individuals with MDD who are non-

responsive to SSRI treatment show either no change or an elevation of circulating cytokine levels

(Syed et al., 2018) and display higher baseline tryptophan metabolism defined by kynurenine to

tryptophan ratio (Sun et al., 2020). Previously, limited efficacy of the SSRI, fluoxetine (FLX),

was commensurate with increased levels of hippocampal IL-1β and decreased circulating

tryptophan concentrations (Qiu et al., 2020a). Thus, tryptophan catabolism via TKP may be

disrupted by CORT and could be a potential biomarker of SSRI efficacy, and the present study

sought to understand this relationship. Here, using a model of *de novo* PPD, we investigated the

effects of maternal postpartum CORT, to induce depressive-like endophenotypes, and concurrent

FLX to understand how these treatments may affect metabolites in the TKP. We hypothesized

that maternal postpartum CORT treatment will alter TKP metabolite concentrations and that

postpartum FLX will have limited efficacy to inhibit these effects.

2. Methods

Forty adult female Sprague-Dawley rats and six male rats (2.5 months old) were

purchased from Charles River (Montreal, QC, Canada). Males were used as breeders, and co-

housed with two females until pregnancy (determined via sperm in daily vaginal lavage

samples). All pregnant rats were undisturbed until parturition. Thirty-seven females were

included in the study. Protocols were in accordance with ethical guidelines set by the Canadian

Council for Animal Care and were approved by the University of British Columbia Animal Care

Committee.

All methods are previously described in Qiu et al. (2020a). Briefly, all dams received

daily injections postpartum, starting from two days after birth, for 22 days. All injections

occurred between 08:00-10:00. Dams received CORT (40mg/kg, s.c.; Sigma-Aldrich, St. Louis,

MO, USA) or its vehicle (OIL; sesame oil + 10% EtOH, 1ml/kg, s.c.) and either FLX (10mg/kg,

s.c.; Sequoia Research Products, Pangbourne, UK) or its vehicle, dextrose (DXT; 5% dextrose in

sterile water, 1ml/kg, s.c.). Dams were randomly assigned to one out of four treatment groups

(OIL+DXT, n = 9; OIL+FLX, n = 10; CORT+DXT, n = 9; and CORT+FLX, n = 9).

At least 2 hours after the last injection, all animals were euthanized by rapid decapitation,

and trunk blood was collected within three minutes of touching the cage. Plasma was collected in

cold EDTA coated tubes and centrifuged 4h later for 10min at 4°C. Data for 5 samples were not

collected due to human error.

We quantified kynurenine (KYN), kynurenic acid (KYNA), xanthurenic acid (XA),

anthranilic acid (AA), 3-hydroxykynurenine (3-HK), 3-hydroxyanthranilic acid (3-HAA),

riboflavin (vitamin B2) and its cofactor, flavin adenine dinucleotide (FAD), using isotope

dilution liquid chromatography coupled with tandem mass spectrometry based on a modified

method by Midttun et al. (2005). The intra- and inter-assay CVs for the quantitation of these

analytes were all $< 7\%$. A neuroprotective-to-neurotoxic z score ratio was calculated to account

for multiple neuroprotective metabolites or neurotoxic metabolites/cofactors: (zKYNA + zAA) /

(z3-HK + z3-HAA + zB2). All metabolite and metabolite ratio data were analyzed using two-

way general linear model of analysis of covariance (ANCOVA) with CORT (CORT or OIL) and

FLX (FLX or DXT) as between-subject factors and estrous cycle stage as covariate, unless

otherwise specified. Effect sizes were reported as η_p^2 or Cohen's *d*. Post hoc comparisons used

Newman-Keuls. *A priori* comparisons were subjected to Bonferroni correction. The covariate of

estrous cycle stage did not significantly affect any analyses (all *p*'s $\geq$ 0.097). Outliers were

removed when two standard deviations away from the mean, and this happened in two analyses

(one outlier in FAD and two outliers removed for the neuroprotective-to-neurotoxic ratio).

3. Results

Maternal FLX treatment decreased XA compared to vehicle DXT treatment (main effect

of FLX: $F_{(1, 29)} = 6.009$, $p = 0.020$, $\eta_p^2 = 0.172$; Figure 1B), with no other significant effects

(all *p*'s $\geq$ 0.296).

Maternal CORT increased AA (main effect of CORT: $F_{(1, 29)} = 44.506$, $p < 0.001$, $\eta_p^2 =$

0.605), 3-HK ($F_{(1, 29)} = 9.282$, $p = 0.005$, $\eta_p^2 = 0.242$), and 3-HAA ($F_{(1, 29)} = 33.693$, $p <$

0.001, $\eta_p^2 = 0.537$) concentrations compared to vehicle-treated animals (Figure 1C-E), with no

other significant effects (all *p*'s $\geq$ 0.253). Maternal CORT also increased plasma KYNA ($F_{(1,}$

$_{29)} = 17.731$, $p < 0.001$; $\eta_p^2 = 0.379$; Figure 1F) with no other significant effects (all *p*'s $\geq$

0.747). CORT+FLX significantly lowered the neuroprotective-to-neurotoxic z score ratio

compared to all other treatment groups, indicating a shift towards the neurotoxic pathway (*a*

priori all *p*'s $\leq$ 0.007; Cohen's $d = 1.785$ compared to OIL+DXT, Cohen's $d = 1.372$ compared

to OIL+FLX, Cohen's $d = 1.963$ compared to CORT+DXT; interaction between CORT and

FLX, $F_{(1, 27)} = 3.906$, $p = 0.058$, $\eta_p^2 = 0.126$; main effect of CORT, $F_{(1, 27)} = 4.265$, $p =$

0.049, $\eta_p^2 = 0.136$; main effect of FLX, $F_{(1, 27)} = 10.055$, $p = 0.004$, $\eta_p^2 = 0.271$, Figure 1G).

There were no other significant effects on KYN or other metabolite concentrations ($p = 0.410$)

Figure 1

Figure 1. Effects of maternal corticosterone (CORT) and fluoxetine (FLX) on plasma
tryptophan-kynurenine pathway metabolites. A. Scheme of the tryptophan-kynurenine
pathway (TKP). **B.** FLX decreased plasma xanthurenic acid concentrations. **C.-F.** CORT
treatment significanly increased plasma anthranilic acid concentrations **(C.)**, plasma 3-
hydroxykynurenine concentrations **(D.)**, plasma 3-hydroxyanthranilic acid concentrations **(E.)**,
and plasma kynurenic acid concentrations **(F.)**. **G.** CORT+FLX significantly decreased the
neuroprotive-to-neurotoxic z score ratio compared to all other groups. *$p < 0.05$. Data
represented in means + standard error of the mean, overlaid with individual data points, n's = 7-
10 per group.

Vitamin B2 and its cofactor, FAD, are critical for the enzymatic conversion of KYN to

the neurotoxic metabolite 3-HK, playing an important role in the neurotoxic pathway of TKP.

Maternal CORT increased plasma vitamin B2 (main effect of CORT: $F_{(1, 29)} = 5.070$, $p =$

0.032, $\eta_p^2 = 0.149$), with no other significant effects (all p's ≥ 0.254, Figure 2A). CORT+FLX

had significantly higher FAD than all other groups (all p's ≤ 0.003; Cohen's $d = 1.718$ compared

to OIL+DXT, Cohen's $d = 2.166$ compared to OIL+FLX, Cohen's $d = 2.332$ compared to

CORT+DXT; interaction between CORT and FLX, $F_{(1, 28)} = 6.862$, $p = 0.014$, $\eta_p^2 = 0.197$;

main effect of CORT, $F_{(1, 28)} = 11.969$, $p = 0.002$, $\eta_p^2 = 0.299$; main effect of FLX, $F_{(1, 28)} =$

4.335, $p = 0.047$, $\eta_p^2 = 0.134$, Figure 2B).

Figure 2

Figure 2. Effects of maternal corticosterone (CORT) and fluoxetine (FLX) treatment on
tryptophan-kynurenine pathway coenzyme and cofactor. A. CORT significantly increased
plasma vitamin B2 concentrations. **B.** CORT+FLX significantly increased plasma flavin adenine
dinucleotide (FAD) concentrations compared to all other groups. *$p < 0.05$. Data represented in
means + standard error of the mean, overlaid with individual data points, n's = 7-10 per group.
163

4. Discussion

Maternal postpartum CORT increased two neurotoxic TKP metabolites, 3-HK and 3-

HAA, and one coenzyme of the neurotoxic pathway, vitamin B2. However, postpartum CORT

also increased two neuroprotective metabolites, KYNA and AA, without significantly affecting

KYN concentrations. Further, maternal postpartum CORT with postpartum FLX shifted the

neuroprotective-to-neurotoxic balance towards neurotoxicity and increased FAD, a coenzyme to

drive conversion to the neurotoxic pathway. Maternal postpartum FLX treatment alone decreased

plasma XA concentration. Together, these data indicate maternal postpartum CORT upregulated

TKP towards the neurotoxic pathway, and this effect together with FLX increased neurotoxic

signalling in the TKP.

Maternal postpartum CORT increased plasma concentrations of neurotoxic metabolites,

3-HK and 3-HAA, which is consistent with findings using inflammation-induced depression

models (Parrott et al., 2016; Tao et al., 2020). Inflammation-induced depression, via

lipopolysaccharide, results in higher concentrations of 3-HK in the frontal cortex and

hippocampus of male mice (Tao et al., 2020). Lipopolysaccharide-induced depressive

endophenotypes are dependent on the activation of the neurotoxic pathway of TKP, via 3-HK, at

least in male mice (Parrot et al., 2016). Maternal CORT, also increased the neuroprotective

metabolite AA, which is consistent with findings showing elevated peripheral concentrations in

both male and female individuals with high risk for MDD and in an animal model of depression

(Sakurai et al., 2020). Together these results suggest that maternal postpartum CORT activates

the neurotoxic pathway of TKP, which shares similarities to other models of depression as well

as in individuals with MDD, and therefore may prove to be an effective therapeutic target in the

future.

Maternal postpartum CORT increased depressive-like endophenotypes (Qiu et al., 2020a)

and in the present study increased plasma KYNA, a neuroprotective metabolite, which is

consistent with another study showing higher gestational KYNA concentrations in women with

the onset of postpartum depressive symptoms compared to healthy controls (Teshigawara et al.,

2019). It is possible the increases in both neuroprotective and neurotoxic metabolites by maternal

postpartum CORT reflects higher TKP activation, which is upregulated by glucocorticoids in

both rodents and humans (reviewed in Maes et al., 2011). Regardless, postpartum CORT

increased activation of the neurotoxic pathway via elevated concentrations of two neurotoxic

metabolites (3-HK and 3-HAA), and the neurotoxic branch coenzyme vitamin B2, and its

cofactor FAD. Thus, higher activation of the neurotoxic branch may have contributed towards

the CORT-induced depressive-like endophenotypes reported previously in Qiu et al. (2020a) and

future studies could use this as therapeutic target and/or increase the efficacy of FLX by

targeting this pathway.

Previously, we reported a significant decrease in plasma tryptophan concentration and

higher levels of the proinflammatory cytokine, IL-1β, in the maternal hippocampus as well as

higher peripheral CXCL1 levels due to FLX treatment in the same animals (Qiu et al., 2020a).

Higher levels of IL-1β can also upregulate TKP (reviewed in Maes et al., 2011), so this may

have contributed to the current findings. Maternal postpartum FLX decreased plasma XA

concentration, which is consistent with data in individuals with MDD (Colle et al., 2020). XA

has not been specifically described as either neuroprotective or neurotoxic, but its downstream

effects on the glutaminergic system have been implicated in psychiatric diseases (reviewed Fazio

et al., 2018). Moreover, maternal postpartum FLX with CORT increased FAD, a cofactor in the

neurotoxic branch, and shifted the balance of the neuroprotective-to-neurotoxic ratio towards the

neurotoxic branch. Previously, FLX failed to reverse depressive-like endophenotypes in this

model (Qiu et al., 2020a). The present findings coupled with the results of higher IL-1β and

reduced TRP concentrations with maternal postpartum FLX (Qiu et al., 2020a) may have

contributed towards the limited efficacy of FLX previously reported in this model of *de novo*

PPD.

Conclusion

Overall, maternal postpartum CORT upregulated metabolites in the neurotoxic pathway

of the TKP, consistent with the profiles of TKP in MDD. Maternal postpartum CORT also led to

higher concentrations of neuroprotective metabolites, KYNA and AA, both of which have been

associated with MDD and PPD. This suggests that the association between TKP metabolites and

depression may be an important therapeutic target. Maternal postpartum FLX commensurate

with maternal CORT increased TKP activation by increasing FAD concentrations and shifting

the neuroprotection-to-neurotoxic balance towards the neurotoxic pathway of TKP. These results

indicate that higher activation of metabolites of the neurotoxic TKP pathway may be a biomarker

of depression and targeting this pathway may lead to novel therapeutic targets for PND and

MDD.

Declaration of conflict

None.

Acknowledgements

The authors would like to thank all the wonderful staff members at the Centre for Disease

Modelling animal facility at the University of British Columbia for the continuous support and

help throughout the project. Financial support for this research was provided by an operating

grant from the Canadian Institutes for Health Research (MOP142308) to LAMG. WQ is

supported by a 4-year Fellowship granted by the University of British Columbia, Faculty of

Medicine, Graduate Program in Neuroscience, and the University of British Columbia Institute

of Mental Health Marshalls Scholars Program in Mental Health, Department of Psychiatry.

References

Colle, R., Masson, P., Verstuyft, C., Fève, B., Werner, E., Boursier-Neyret, C., Walther, B., David, D.J., Boniface, B., Falissard, B., Chanson, P., Corruble, E., Becquemont, L., 2020. Peripheral tryptophan, serotonin, kynurenine, and their metabolites in major depression: A case–control study. Psychiatry and Clinical Neurosciences 74, 112–117. https://doi.org/10.1111/pcn.12944

Corwin, E.J., Johnston, N., Pugh, L., 2008. Symptoms of Postpartum Depression Associated With Elevated Levels of Interleukin-1 Beta During the First Month Postpartum. Biological Research For Nursing 10, 128–133. https://doi.org/10.1177/1099800408323220

Fazio, F., Ulivieri, M., Volpi, C., Gargaro, M., Fallarino, F., 2018. Targeting metabotropic glutamate receptors for the treatment of neuroinflammation. Current Opinion in Pharmacology, Neurosciences 38, 16–23. https://doi.org/10.1016/j.coph.2018.01.010

Haapakoski, R., Mathieu, J., Ebmeier, K.P., Alenius, H., Kivimäki, M., 2015. Cumulative meta-analysis of interleukins 6 and 1β, tumour necrosis factor α and C-reactive protein in patients with major depressive disorder. Brain, Behavior, and Immunity 49, 206–215. https://doi.org/10.1016/j.bbi.2015.06.001

Maes, M., Leonard, B.E., Myint, A.M., Kubera, M., Verkerk, R., 2011. The new '5-HT' hypothesis of depression: Cell-mediated immune activation induces indoleamine 2,3-dioxygenase, which leads to lower plasma tryptophan and an increased synthesis of detrimental tryptophan catabolites (TRYCATs), both of which contribute to the onset of depression. Progress in Neuro-Psychopharmacology and Biological Psychiatry, The Neuro-inflammatory and Neuroprogressive Pathways in Depression 35, 702–721. https://doi.org/10.1016/j.pnpbp.2010.12.017

Midttun, O., Hustad, S., Solheim, E., Schneede, J., Ueland, P.M., 2005. Multianalyte
quantification of vitamin B6 and B2 species in the nanomolar range in human plasma by
liquid chromatography-tandem mass spectrometry. Clin. Chem. 51, 1206–1216.
https://doi.org/10.1373/clinchem.2005.051169
Munk-Olsen, T., Laursen, T.M., Pedersen, C.B., Mors, O., Mortensen, P.B., 2006. New parents
and mental disorders: a population-based register study. JAMA 296, 2582–2589.
https://doi.org/10.1001/jama.296.21.2582
Parrott, J.M., Redus, L., Santana-Coelho, D., Morales, J., Gao, X., O'Connor, J.C., 2016.
Neurotoxic kynurenine metabolism is increased in the dorsal hippocampus and drives
distinct depressive behaviors during inflammation. Transl Psychiatry 6, e918.
https://doi.org/10.1038/tp.2016.200
Qiu, W., Hodges, T.E., Clark, E.L., Blankers, S.A., Galea, L.A.M., 2020. Perinatal depression:
Heterogeneity of disease and in animal models. Frontiers in Neuroendocrinology 59,
100854. https://doi.org/10.1016/j.yfrne.2020.100854
Sakurai, M., Yamamoto, Y., Kanayama, N., Hasegawa, M., Mouri, A., Takemura, M.,
Matsunami, H., Miyauchi, T., Tokura, T., Kimura, H., Ito, M., Umemura, E., (Boku),
A.S., Nagashima, W., Tonoike, T., Kurita, K., Ozaki, N., Nabeshima, T., Saito, K., 2020.
Serum Metabolic Profiles of the Tryptophan-Kynurenine Pathway in the high risk
subjects of major depressive disorder. Scientific Reports 10, 1961.
https://doi.org/10.1038/s41598-020-58806-w
Sun, Y., Drevets, W., Turecki, G., Li, Q.S., 2020. The Relationship between Plasma Serotonin
and Kynurenine Pathway Metabolite Levels and the Treatment Response to Escitalopram
and Desvenlafaxine. Brain, Behavior, and Immunity.
https://doi.org/10.1016/j.bbi.2020.01.011
Syed, S.A., Beurel, E., Loewenstein, D.A., Lowell, J.A., Craighead, W.E., Dunlop, B.W.,
Mayberg, H.S., Dhabhar, F., Dietrich, W.D., Keane, R.W., de Rivero Vaccari, J.P.,
Nemeroff, C.B., 2018. Defective Inflammatory Pathways in Never-Treated Depressed
Patients Are Associated with Poor Treatment Response. Neuron 99, 914-924.e3.
https://doi.org/10.1016/j.neuron.2018.08.001
Tao, X., Yan, M., Wang, L., Zhou, Y., Wang, Z., Xia, T., Liu, X., Pan, R., Chang, Q., 2020.
Homeostasis Imbalance of Microglia and Astrocytes Leads to Alteration in the
Metabolites of the Kynurenine Pathway in LPS-Induced Depressive-Like Mice.
International Journal of Molecular Sciences 21, 1460.
https://doi.org/10.3390/ijms21041460
Teshigawara, T., Mouri, A., Kubo, H., Nakamura, Y., Shiino, T., Okada, T., Morikawa, M.,
Nabeshima, T., Ozaki, N., Yamamoto, Y., Saito, K., 2019. Changes in tryptophan
metabolism during pregnancy and postpartum periods: Potential involvement in
postpartum depressive symptoms. Journal of Affective Disorders 255, 168–176.
https://doi.org/10.1016/j.jad.2019.05.028
Wisner, K.L., Sit, D.K.Y., McShea, M.C., Rizzo, D.M., Zoretich, R.A., Hughes, C.L., Eng, H.F.,
Luther, J.F., Wisniewski, S.R., Costantino, M.L., Confer, A.L., Moses-Kolko, E.L.,
Famy, C.S., Hanusa, B.H., 2013. Onset Timing, Thoughts of Self-harm, and Diagnoses in
Postpartum Women With Screen-Positive Depression Findings. JAMA Psychiatry 70,
490–498. https://doi.org/10.1001/jamapsychiatry.2013.87