CTP regulates membrane-binding activity of the nucleoid occlusion protein Noc

Adam S. B. Jalal[1], Ngat T. Tran[1], Ling J. Wu[2], Karunakaran Ramakrishnan[1], Martin Rejzek[3], Clare E. M. Stevenson[4], David M. Lawson[4], Jeff Errington[2], and Tung B. K. Le[1]

[1]Department of Molecular Microbiology
John Innes Centre, Norwich, NR4 7UH, United Kingdom
[2]Centre for Bacterial Cell Biology, Biosciences Institute, Medical School
Newcastle University, Newcastle Upon Tyne, NE2 4AX, United Kingdom
[3]Chemistry Platform
John Innes Centre, Norwich, NR4 7UH, United Kingdom
[4]Department of Biological Chemistry
John Innes Centre, Norwich, NR4 7UH, United Kingdom

*Correspondence: tung.le@jic.ac.uk

ABSTRACT
ATP and GTP-dependent molecular switches are extensively used to control functions of proteins in a wide range of biological processes. However, CTP switches are rarely reported. Here, we report a nucleoid occlusion protein Noc as the first example of a CTPase enzyme whose membrane-binding activity is directly regulated by a CTP switch. In *Bacillus subtilis*, Noc nucleates on 16-bp *NBS* sites before associating with neighboring non-specific DNA to form large membrane-associated nucleoprotein complexes to physically occlude assembly of the cell division machinery. By *in vitro* reconstitution, we show that (i) CTP is required for Noc to form the *NBS*-dependent nucleoprotein complex, and (ii) CTP binding, but not hydrolysis, switches Noc to a membrane-active state. Overall, we suggest that CTP couples membrane-binding activity of Noc to nucleoprotein complex formation to ensure productive recruitment of DNA to the bacterial cell membrane for nucleoid occlusion activity.

INTRODUCTION

While ATP and GTP switches are ubiquitous in biology, CTP switches have rarely been identified but may be more widespread than previously appreciated. A recent discovery showed that ParB, a crucial protein for bacterial chromosome segregation, is the founding member of a new class of CTP-dependent molecular switches[1,2]. ParB nucleates on a *parS* DNA sequence and associates with neighboring DNA, a process known as spreading, to enable faithful chromosome segregation[3–8]. CTP induces ParB self-dimerization to create a clamp-like molecule[2]. The ParB clamp self-loads at *parS*, then spreads by sliding to neighboring DNA while still entrapping DNA[2,9]. Essentially, CTP serves to switch ParB from a *parS*-nucleating open clamp to a DNA-sliding closed clamp state[2,9]. The end result is the formation of a higher-order nucleoprotein complex with multiple ParB-CTP clamps entrapped in the vicinity of the *parS* locus. The higher-order nucleoprotein complex stimulates the ATPase activity of ParA, a partner of ParB, driving the segregation of replicated chromosomes to daughter cells[3,10–14].

In Firmicutes, the nucleoid occlusion protein Noc is a paralog of ParB[15–17]; however, the role of Noc is different from that of a canonical ParB[18–20]. Noc helps to direct the assembly of the cell division machinery towards the middle of a dividing cell where the concentration of chromosomal DNA (the nucleoid) is the least, thus ensuring a binary cell division[18,21–23]. Noc does so by nucleating on 16-bp *NBS* (Noc-binding site) sites scattered around the chromosome before spreading to neighboring DNA to form large Noc-DNA nucleoprotein complexes[24]. Unusually, Noc is also a peripheral membrane protein that directly associates with the cell membrane via an N-terminal amphipathic helix[21] (Fig. 1A). The recruitment of the chromosomal DNA to the membrane is crucial for preventing the assembly of the division machinery over the chromosome; indeed a Noc variant lacking the amphipathic helix is impaired in nucleoid occlusion activity[21]. In *Bacillus subtilis*, Noc was observed to associate with the cell membrane in a transient manner *in vivo*[24]. It was thought that a strong membrane-binding activity of Noc might have been selected against since a stable association with the membrane might hamper chromosome replication and segregation[21], yet it is unclear

how the membrane-binding activity of Noc is modulated. Furthermore, Noc must bring the chromosomal DNA to the membrane to physically inhibit the assembly of the division machinery[21]; an unregulated membrane-binding activity would likely confine apo-Noc permanently to the cell membrane, thus unfavorably limiting the recruitment of DNA to the membrane[21]. Again, it remains unclear whether the membrane-binding activity of Noc is regulated, and if so how.

To investigate further, we have biochemically reconstituted *NBS*-dependent Noc spreading and membrane association events using purified *B. subtilis* Noc protein and phospholipid vesicles. We show that, similar to a canonical ParB, Noc is a CTPase enzyme that binds CTP to form a protein clamp that can slide and entrap DNA. Importantly, CTP binding, but not hydrolysis, is required to switch Noc-DNA from a membrane inactive to an active state, thus locking Noc into a pathway in which it has to spread before associating with the cell membrane. We solve an X-ray crystal structure of a C-terminal domain truncated apo-Noc from *Geobacillus thermoleovorans*, in which its membrane-targeting amphipathic helix adopts an autoinhibitory conformation, restricted from interacting with the membrane. We suggest that CTP binding might liberate the amphipathic helix, thereby switching Noc to a membrane-active state. Altogether, we demonstrate that CTP directly regulates the membrane-binding activity of the nucleoid occlusion protein Noc, further expanding the role of CTP switches in biology.

RESULTS

NBS DNA increases the CTP binding and hydrolysis rate of Noc

Given the shared ancestry between ParB and Noc, we wondered if *B. subtilis* Noc also binds and hydrolyses CTP. To investigate, we employed a membrane-spotting assay (DRaCALA) and showed that *B. subtilis* Noc binds radiolabeled CTP but only in the presence of a cognate 22-bp *NBS* DNA (Fig. 1B). An excess of unlabeled CTP, but no other NTP, outcompeted radiolabeled CTP for binding to Noc, suggesting that *B. subtilis* Noc binds CTP specifically (Fig. 1C). Similarly, an N-terminally truncated Noc variant lacking the 10-AA membrane targeting amphipathic helix (NocNΔ10) also bound radiolabeled CTP in the

presence of *NBS* DNA (Fig. 1D). However, the Noc (R89A) and Noc (N121S) variants, whose equivalent substitutions in ParB have been shown to impair spreading and CTP binding[1,2,9] (Fig. S1A), did not bind radiolabeled CTP at the tested concentration (Fig. 1D). Next, we performed isothermal titration calorimetry (ITC) to obtain quantitative binding parameters between a non-hydrolyzable CTP analog (CTPɣS) and *B. subtilis* Noc. We found that *B. subtilis* Noc binds CTPɣS with a moderate affinity (K_d = 67.5 ± 23 µM), while Noc (N121S) bound CTPɣS weaker at K_d = 231.5 ± 66 µM, and Noc (R89A) did not detectably bind nucleotide (Fig. S1B). Consistent with a previous report[2], *B. subtilis* Noc also showed CTP hydrolysis activity, albeit at a low rate of ~1 CTP per Noc per hour when only the purified protein and CTP were included (Fig. 1E)[2]. The addition of a 22-bp *NBS* DNA, but not a non-cognate 22-bp *parS* DNA, increased the CTP hydrolysis rate ninefold to ~9 CTP per Noc per hour (Fig. 1E)[2]. The Noc (R89A) and Noc (N121S) variants did not show noticeable CTP-hydrolyzing activity (Fig. 1E). Lastly, despite binding CTP equally or more strongly than the WT (Fig. 1D), NocNΔ10 hydrolyzed CTP at a reduced rate of ~5 CTP per Noc per hour (Fig. 1E) (see Discussion). Altogether, our data suggest that *B. subtilis* Noc is a CTPase enzyme that binds and hydrolyses CTP in the presence of cognate DNA.

CTP and *NBS* DNA stimulate the engagement of the N-terminal domain of Noc *in vitro*

In the presence of CTP, ParB self-engages at the N-terminal domain (NTD) to create a clamp-like molecule[2,9]. To investigate whether CTP elicits a similar response in Noc, we employed site-specific crosslinking of a purified *B. subtilis* Noc (E29C) variant by a sulfhydryl-to-sulfhydryl crosslinker bismaleimidoethane (BMOE). Residue E29 at the NTD were substituted by cysteine on an otherwise cysteine-less Noc (WT) background (Fig. S2A) to create a variant in which symmetry-related cysteines become covalently linked together if they are within 8 Å of each other. The cross-linked form was detectable as a protein form with reduced mobility on SDS-PAGE (labelled X in Figure 2). In the absence of CTP, Noc (E29C) crosslinked minimally (~10% crosslinked fraction, lanes 2-4 of Fig. 2A). The crosslinking efficiency increased threefold (~30%) in the presence of CTP (lane 8 of Fig. 2A), but not CDP or ATP (Fig. 2A and Fig. S2B). The crosslinking efficiency further

increased fivefold (~55%) when both CTP and a 22-bp *NBS* were included (lane 10 of Fig 2A, and Fig. S2D). However, the addition of a non-cognate 22-bp *parS* DNA did not result in the same high level of crosslinking even when CTP was present (lane 9 of Fig. 2A). Noticeably, a non-hydrolyzable CTPɣS readily promoted crosslinking (~45% crosslinked fraction) regardless of the presence or absence of *NBS* DNA (lanes 8-10 of Fig. 2B, and Fig. S2E). Therefore, our data suggest that CTP binding, but not hydrolysis, is required for the NTD engagement of *B. subtilis* Noc. Consistent with the requirement of CTP binding for NTD engagement, the Noc (E29C R89A) variant, in which the R89A substitution incapacitated CTP binding, did not crosslink beyond the background level in any tested condition (Fig. 2C). The Noc (E29C N121S) variant, which binds CTP (albeit at a reduced affinity) but cannot hydrolyze CTP, crosslinked similarly to Noc (E29C) (~30% crosslinked fraction) at the saturating concentration of 1 mM CTP, although the *NBS*-stimulated crosslinking was abolished (Fig. 2D). Lastly, the NocNΔ10 (E29C) variant, which lacks the N-terminal amphipathic helix, crosslinked similarly to Noc (E29C) in the presence of CTP and *NBS* DNA (Fig. S2C).

A clamp-like Noc associates with a closed *NBS* DNA substrate in a CTP-dependent manner

To further investigate if the NTD engagement produces a clamp-like Noc that can entrap DNA, we performed crosslinking reactions on a dual-cysteine Noc (E29C S253C) variant (Fig. 2E). Residues E29 and S253 at the NTD and the C-terminal domain (CTD) of Noc, respectively, were substituted by cysteine (Fig. 2E). While Noc (E29C) crosslinked in a CTP and *NBS*-dependent manner (Fig. 2A), Noc (S253C) crosslinked efficiently (~80% crosslinked fraction) but independently of CTP and/or *NBS* (Fig. S2F), consistent with the known role of the CTD as a primary dimerization domain[21,25]. We observed that purified Noc (E29C S253) formed a high molecular weight (HMW) species near the top of the polyacrylamide gel in the presence of CTP, a circular *NBS*-harboring plasmid, and the crosslinker BMOE (lane 7 of Fig. 2E-F and Fig. S2G). The HMW smear contained both protein and DNA, as apparent from dual staining with a protein stain (Coomassie) and a DNA stain (Sybr green) (Fig. 2F). The slowly-migrating DNA-stained bands were also observed when the

crosslinking products were resolved on an agarose gel (Fig. 2G). Therefore, the HMW smear most likely contained protein-DNA catenates between a circular *NBS* plasmid and a denatured but otherwise circularly crosslinked Noc (E29C S253C) polypeptide. To verify the nature of the catenates, crosslinking products were treated with Benzonase, a non-specific DNase, before SDS-PAGE analysis (lane 8 of Fig. 2E-F and Fig. S2G). Here, the use of Benzonase eliminated the HMW smear, likely by unlinking the protein-DNA catenates. Lastly, the HMW smear was not observed when an empty plasmid was employed or when CTP was omitted from the reaction (lanes 4-5 and 9-10 of Fig. 2E-F and Fig. S2G). Altogether, our results indicate that *B. subtilis* Noc entraps DNA in an *NBS*- and CTP-dependent manner.

To follow the DNA entrapment and spreading of Noc in real-time, we employed a 170-bp dual biotin-labeled *NBS* DNA that had been tethered at both ends onto a streptavidin-coated probe to form a closed DNA and measured bio-layer interferometry (BLI) signal (Fig. 3)[9]. BLI monitors wavelength shifts resulting from changes in the optical thickness of the probe during the association/dissociation of Noc from the closed *NBS* DNA substrate. In the absence of CTP, we observed only the nucleation event on *NBS* DNA with 1 µM purified Noc (Fig. 3A). Premixing Noc with ATP, GTP, or UTP did not change the sensorgram markedly, however the addition of 1 mM CTP increased the BLI response by ~sixfold (Fig. 3A), consistent with Noc-CTP spreading from *NBS* to accumulate more on the 170-bp closed DNA substrate than by nucleation alone. We did not observe a noticeable BLI response when a 170-bp closed *parS* DNA substrate was employed instead (Fig. 3A), confirming that nucleation and spreading by *B. subtilis* Noc is strictly dependent on the *NBS*. We further observed by BLI assay that the Noc (R89A) and Noc (N121S) variants could nucleate but could not spread to accumulate on the closed *NBS* substrate even in the presence of 1 mM CTP (Fig. 3A), consistent with the requirement of NTD engagement for DNA entrapment (see also Fig. 2C-D).

Next, we investigated whether a DNA substrate with a free end (an open DNA) can also support Noc accumulation in our BLI setup. The 170-bp dual biotin-labeled was designed with an unique BamHI recognition site flanking the *NBS* site

(Fig. 3B). To generate a free end on the DNA, the DNA-coated probe was immersed in a buffer containing *Bam*HI restriction enzyme. Before *Bam*HI digestion, Noc showed an enhanced association on a closed DNA substrate in the presence of CTP (Fig. 3A). However, after *Bam*HI digestion, the addition of CTP did not affect the BLI response beyond the nucleation of Noc at the *NBS* (Fig. 3B). We reasoned that, similar to the canonical ParB[9], Noc spreads but quickly escapes by sliding off a free DNA end. Overall, our BLI analysis is consistent with the site-specific crosslinking analysis and suggests a clamp-like Noc-CTP that can spread and accumulate on a closed DNA substrate by entrapping DNA.

Noc binds liposomes in the presence of CTP
It has been shown previously *in vivo* that Noc possesses membrane-binding activity and it must bring chromosomal DNA to the cell membrane to prevent the formation of cell division machinery[21]. Puzzlingly, however, we could not observe noticeable association between purified *B. subtilis* Noc and liposomes by a co-sedimentation assay (lanes 1-4 of Fig. 4A). We wondered if CTP might be the missing co-factor that activates the membrane-binding activity of Noc. To test this possibility, purified Noc was incubated with liposomes with or without CTP and 22-bp *NBS* DNA, and ultracentrifuged (Fig. 4A and Fig. S3A). The pellet contained sedimented liposomes and associating protein while the supernatant contained unbound protein; protein and DNA species from both fractions were analyzed by polyacrylamide gel electrophoresis. We did not observe a significant increase in the amount of Noc in the pellet when CTP alone, CTP and a non-cognate 22-bp *parS* DNA (lanes 5-8 of Fig. 4A), or other nucleotides were employed (Fig. S3B). However, in the presence of both CTP and a 22-bp *NBS* DNA, ~45% of the total amount of Noc was detected robustly in the pellet (lane 9-12 of Fig. 4A), suggesting that the *in vitro* membrane-binding activity of Noc is CTP- and *NBS*-dependent. *NBS* DNA is most likely required to stimulate CTP binding, hence the membrane-binding activity of Noc, rather than to concentrate a large amount of weakly binding Noc in the vicinity of *NBS* to collectively enhance its affinity for the cell membrane. Supporting this proposition, the short length of a 22-bp *NBS* DNA duplex means that only a dimer of Noc or Noc-CTP complex can occupy the DNA at a

time. Furthermore, nearly all of the 22-bp *NBS* DNA was present in the supernatant (instead of in the pellet) after centrifugation (Fig. 4A), most likely because Noc-CTP clamps rapidly escaped the open linear *NBS* DNA, further arguing that individual Noc-CTP possesses a substantial membrane-binding capability.

Next, we observed that the addition of CTPγS alone caused ~35% of Noc to associate with the pelleted vesicles (lanes 5-6, Fig. 4B). The vesicle-bound fraction further increased to ~45% when both CTPγS and a 22-bp *NBS* DNA were present (lanes 7-8, Fig. 4B). We infer that CTP binding, but not hydrolysis, is required for the *in vitro* Noc-liposomes interaction. Consistent with the requirement of CTP binding for membrane-binding activity, the Noc (R89A) and Noc (N121S) variants which do not bind CTP or bind CTP weakly failed to co-sediment with liposomes even when CTP and *NBS* were included (lanes 3-4 and 5-6, Fig. 4C). The N-terminal amphipathic helix was previously shown *in vivo* to be the membrane-targeting determinant of *B. subtilis* Noc[21]. Here, we also confirmed that a purified NocNΔ10 lacking this amphipathic helix was unable to co-sediment with liposomes *in vitro* regardless of the presence or absence of CTP or *NBS* DNA (lanes 7-8, Fig. 4C). Lastly, in another control experiment, *C. crescentus* ParB, which binds CTP but not the cell membrane[9,10,26], did not co-sediment with liposomes in the presence of CTP ± *parS* or *NBS* DNA (Fig. S3C).

Noc recruits *NBS* plasmid to liposomes in the presence of CTP

The recruitment of chromosomal DNA to the membrane is essential for Noc to exert the nucleoid occlusion activity *in vivo*[21]. Indeed, ectopic expression of neither *noc (R89A)*, *noc (N121S)*, nor *nocNΔ10* could rescue the synthetic cell division defect of a *B. subtilis* ΔnocΔminCD double mutant at elevated temperature (Fig. 4D)[21]. Epi-fluorescence microscopy of *B. subtilis* cells harboring *yfp*-tagged *noc* mutant alleles also confirmed that Noc (R89A), Noc (N121S), and NocNΔ10 failed to form punctate foci near the cell periphery i.e. were defective in the formation of large membrane-associated nucleoprotein complexes[21] (Fig. 4E). We wondered if the Noc-dependent recruitment of DNA to the membrane can be biochemically reconstituted. To this end, we assembled a reaction containing purified Noc, liposomes, CTP, and a ~5-kb circular *NBS*-

harboring plasmid before ultracentrifugation (Fig. 5). Unlike the 22-bp *NBS* DNA, the *NBS* plasmid is topologically closed and therefore should not be able to escape entrapment by Noc-CTP clamps. Unfortunately, owing to its high molecular weight, ~45 to 55% of the circular plasmid sedimented independently of the liposomes (lanes 5-6 of Fig. 5A, and lanes 1-4 of Fig. 5B). Nevertheless, in the presence of liposomes and CTP, the *NBS* plasmid completely co-sedimented with Noc (lanes 11-12 of Fig. 5A), demonstrating that Noc can recruit plasmid DNA to liposomes in the presence of CTP. The pellet/ supernatant distribution of a control plasmid with no *NBS* ("empty") was unaffected by the presence of Noc and CTP, and only ~2% of Noc was found in the pellet (lanes 9-10 of Fig. 5A). Next, in an attempt to minimize the sedimentation of a plasmid by itself, we performed a vesicle flotation assay in which liposomes and associating protein/DNA migrate up a sucrose gradient to the topmost faction rather than down into the pellet (Fig. S4A). Despite the basal level of ~20% total *NBS* plasmid in the top fraction even when liposomes were omitted (lane 3 of Fig. S4B), we again observed ~80% of total *NBS* plasmid being recruited to the liposomes when purified Noc and CTP were also present (lane 12 of Fig. S4C).

In light of the above results, we wondered if the role of *NBS* was to stimulate the membrane-binding activity of Noc-CTP. To test this possibility, we first assembled a co-sedimentation reaction as described above. Subsequently, Benzonase was added to eliminate *NBS* plasmid before ultracentrifugation (lanes 7-8 of Fig. 5B). The nuclease treatment eliminated intact *NBS* plasmid from both the supernatant and the pellet fractions, however ~65% of the total amount of Noc still remained co-sedimented to the pellet in comparison to ~80% when nuclease was omitted (lanes 7-8 vs. lanes 5-6 of Fig. 5A). In parallel, we tested spreading-defective Noc (R89A) and Noc (N121S), and a membrane-binding-defective NocNΔ10 for their ability to recruit *NBS* plasmid to liposomes in a co-sedimentation assay (Fig. 5C) as well as in a flotation assay (Fig. S4D). Consistent with previous and the above *in vivo* data[21], these mutants could not recruit DNA to the pellet fraction beyond the basal level (Fig. 5C and Fig. S4D). Altogether, these results suggest that the *NBS* specifically activates the membrane-binding activity of Noc in the

presence of CTP, thereby recruiting Noc-DNA complexes to the membrane.

The association of Noc-*NBS* DNA with liposomes is reversible

Once the membrane-associated Noc-DNA nucleoprotein complexes form, can this process be reversed? To investigate, we employed a buffer supplemented with EDTA to sequester coordinated Mg^{2+} cation to disrupt CTP binding from the preformed liposome-bound Noc-DNA complexes (Fig. 5D). In this experiment, the pellet containing preformed liposomes-bound Noc-DNA complexes (lanes 2 or 4 of Fig. 5D) was either resuspended in EDTA-minus or EDTA-plus buffer before being ultracentrifuged again. After the second round of centrifugation, both the pellet and the supernatant fractions were analyzed for the presence of protein and DNA (lanes 5-8 of Fig. 5D). We observed that while nearly all *NBS* plasmid remained in the pellet when an EDTA-minus buffer was used (lane 6 of Fig. 5D), ~38% of total plasmid returned to the supernatant in the presence of EDTA (lane 7 of Fig. 5D). These results demonstrate that the membrane-binding activity of Noc can be reversed and suggest a possible inhibitory mechanism that keeps apo-Noc in the membrane inactive mode in the absence of CTP.

Crystal structure of *Geobacillus thermoleovorans* NocΔCTD shows the membrane-targeting amphipathic helix in an autoinhibitory conformation

To gain further insights into the membrane inactive state, we sought to solve a crystal structure of Noc. We could not obtain high quality crystals of *B. subtilis* Noc either in full-length or truncated forms despite extensive efforts. However, we could grow and collect diffraction data for a C-terminal domain truncated apo-Noc (NocΔCTD) from a thermophilic bacterium, *Geobacillus thermoleovorans*, to 2.5 Å resolution. *B. subtilis* Noc and *G. thermoleovorans* Noc share 72% sequence identity (Fig. S5A). The structure was solved by iodide SAD phasing since no other Noc protein family structure was available as a template for molecular replacement. The asymmetric unit contains two similar copies of monomeric apo-Noc (RMSD = 0.9 Å) (Fig. S5B), hence we used the more complete subunit for all further analysis.

Each NocΔCTD subunit contains an N-terminal domain (NTD) (helices α1-α6) and an *NBS*-specific DNA-binding domain (DBD) (helices α7-α12) (Fig. 6A)[17,24]. The primary dimerization domain at the C-terminal side of Noc was truncated in NocΔCTD, hence is not present in this structure. Most notably, electron density for five of the 10 amino acids (AA) comprising the membrane-targeting helix α1 are visible and adopt a 3_{10} helical conformation (Fig. 6A-B). From the structure, it is apparent that the membrane-targeting sequence of Noc is indeed amphipathic, with distinct polar and hydrophobic faces (Fig. 6B). The amphipathic helix α1 is immediately followed by helix α2 and subsequently by an 8-AA α2-β1 loop that precedes the main NTD (Fig. 6A). By sequence comparison with a canonical ParB[2], the main NTD (β1-α6) of Noc contains the CTP-binding motifs, while the amphipathic α1, α2, and the α2-β1 loop are specific to the Noc protein family (Fig. S5A). We noted that the hydrophobic face of the amphipathic α1 helix is buried towards α5 and α7 at the core of Noc (Fig. 6A and 6C), thus is unexposed and unlikely to be available for membrane interaction. Specifically, the side chain of S6 hydrogen bonds with the side chain of R150, and the side chain of R7 hydrogen bonds with the main chain oxygen of N104 (Fig. 6C). Additionally, the main chain oxygen of S10 hydrogen bonds with the side chain of Q16, and lastly, the main chain oxygen of F11 hydrogen bonds with the side chain of Q120 (Fig. 6C). Sidechains of F9 and F11 also interact hydrophobically with side chains of I116 and I81, respectively (Fig. 6C). These interactions thus bury α1 in a potential autoinhibitory conformation. We further noted that α2, which does not target the membrane *per se*, but is conserved in Noc homologs (Fig. S5A)[21], also contributes to holding α1 in the repressed state (Fig. 6D). Specifically, the side chains of both E13 and Q16 form water-mediated contacts with the side chains of I65, R86, and K103, while the side chain of E20 forms a hydrogen bond with Q66 (Fig. 6D). Overall, our apo-NocΔCTD structure suggests a repressed state that might keep Noc in the membrane inactive state in the absence of CTP.

Crystal structure of the *G. thermoleovorans* NocNΔ26ΔCTD variant is incompatible with an autoinhibitory conformation of the amphipathic helix

Next, we attempted to obtain a co-crystal structure of Noc in complex with nucleotides. In the presence of CTPγS, we were able to grow and collect 2.95 Å diffraction data for a crystal of a *G. thermoleovorans* NocNΔ26ΔCTD variant, which lacks both the N-terminal membrane-targeting helix and the C-terminal domain. After solving its structure, it is apparent that NocNΔ26ΔCTD has adopted an alternative conformation to that of NocΔCTD (Fig. 7A vs. Fig. 7B-C). This alternative conformation of NocΔN26ΔCTD is compatible with homodimer formation, giving an interfacial area of ~2700 Å^2 (as evaluated with jsPISA[27]), which resembles that observed for a co-crystal structure of *B. subtilis* ParBΔCTD with bound cytidine diphosphate (CDP) (RMSD = 2.17 Å) (Fig. S6A)[2]. However, there was no clear electron density for a bound nucleotide in our structure, instead a sulfate anion from the crystallization solution occupies a position equivalent to that of the β-phosphate of the nucleotide in the *B. subtilis* ParB-CDP structure (Fig. S6B)[2]. We further observed that helix α5 in the NocΔN26ΔCTD structure swings outwards by 104 degrees and no longer forms a bundle with helix α6 from the same subunit (Fig. 7A vs. Fig. 7B), and that this movement might drive the self-dimerization at the N-terminal domain of Noc i.e. the NTD engagement (Fig. 7C). By superimposing the NTDs of NocΔCTD and NocNΔ26ΔCTD (RMSD = 1.49 Å), we detected severe clashes between α1, α2, and the opposite subunit of NocNΔ26ΔCTD (Fig. 7D and Fig. S6C-D). Therefore, it is clear that the autoinhibitory state of α1 and α2 (as observed in NocΔCTD) is not compatible with the alternative conformation in the NocNΔ26ΔCTD structure. We speculate that the amphipathic helix α1 and α2 might be liberated from its autoinhibitory conformation to be compatible with the NTD-engagement conformation in the NocNΔ26ΔCTD structure.

DISCUSSION
Assembly of the membrane-associated Noc-DNA nucleoprotein complex: roles of CTP
The nucleoid occlusion protein Noc increases cell division efficiency[22] by directing the division machinery towards mid-cell either by inhibiting FtsZ formation over the nucleoid[18,21] and/or by concentrating FtsZ in the vicinity of a pre-existing mid-cell Z-ring[23]. The extensive Noc-mediated DNA-membrane interaction is at the heart of both models for nucleoid occlusion[21,23]. In this study, we show that CTP regulates the nucleoid occlusion activity of Noc. We provide evidence that (i) CTP is required for Noc to form the *NBS*-dependent nucleoprotein complex, and (ii) CTP binding switches Noc from a membrane-inactive auto-repressed state to a membrane-active state. We propose that the dual dependency of Noc's membrane-binding activity on *NBS* and CTP might ensure productive recruitment of DNA to the bacterial cell membrane to exert the nucleoid occlusion activity (Fig. 8). It has been estimated that the intracellular concentrations of Noc and CTP are ~5 μM and ~1 mM, respectively[24,28]. At these concentrations, if the membrane-binding activity of Noc was solely dependent on CTP, most intracellular DNA-unbound Noc would be in the CTP-bound state and confined to the cell membrane, thus potentially limiting the recruitment of chromosomal DNA to the cell membrane. We therefore reason that the *NBS*-stimulated Noc-CTP interaction provides a mechanism to commit Noc into a pathway in which only DNA-entrapped Noc molecules are able to associate with the cell membrane (Fig. 8B). Another consequence of coupling membrane activity to *NBS* binding is that membrane-associated nucleoprotein complexes are spatially confined near the vicinity of *NBS* sites. This spatial confinement is important for directing division machinery formation towards mid-cell where the concentration of *NBS* sites, hence the nucleoid occlusion activity, is lowest[24].

The lack of a bound nucleotide in the NocΔN26ΔCTD structure prevents us from drawing a conclusion on the role of CTP and the possible conformational liberation of the amphipathic helix α1. However, several lines of evidence suggest that CTP might favor the NTD-engagement conformation as observed in our NocNΔ26ΔCTD structure: (i) CTP/CTPγS promoted the crosslinking between symmetrical E29C residues, a readout for NTD-engagement (Fig. 2A), (ii) the conformation of NocNΔ26ΔCTD is highly similar to that of a CDP-bound *B. subtilis* ParB (Fig. S6A)[2], and (iii) CTP/CTPγS enables Noc (WT) to co-sediment with liposomes (Fig. 4). Altogether, it is not unreasonable to speculate that the CTP-induced NTD engagement might liberate the amphipathic helix from its autoinhibitory state to interact with the cell membrane. The 8-AA loop that connects the amphipathic helix to the rest of Noc might offer the flexibility to orient the amphipathic helix

parallel to the membrane plane for binding (Fig. 8).

This study also provides evidence that *B. subtilis* Noc possesses CTPase activity but CTP hydrolysis is neither required for the clamp formation nor membrane association. What might possibly the role of CTP hydrolysis be? Similar to the counterpart ParB[1,2,9], CTP hydrolysis and/or the subsequent release of hydrolytic products might disengage the NTD to open the clamp to release entrapped DNA (Fig. 8B). Concomitantly, Noc might revert to the membrane-inactive state, enabling its extraction from the cell membrane. Our experiment that employed EDTA to sequester coordinated Mg^{2+} to artificially promote the dissociation of CTP from Noc supports the proposal that membrane association can be reversible (Fig. 5D). Furthermore, *B. subtilis* Noc foci associate with the cell membrane in a transient manner, perhaps suggesting a weak and fast on/off membrane interaction *in vivo*[21]. The transient association with the membrane, possibly endowed by a CTP hydrolysis event that facilitates the release of Noc, might be advantageous for the cells because a strong and permanent mode of binding might hamper chromosome replication or damage DNA. Supporting this view, fusing NocNΔ10 to a synthetic transmembrane helix led to broken, bisected chromosomes and eventually chromosome segregation defects in many *B. subtilis* cells[21,29]. Lastly, we noted that NocNΔ10 binds CTP equally/slightly stronger than Noc (WT) (Fig. 1D) yet its CTPase rate is reduced by a half (Fig. 1E). It is still unclear why this is the case; however, we speculate that the autoinhibitory conformation of this N-terminal most region might play a role in CTP binding/hydrolysis. Lending support to our speculation, this N-terminal-most region interacts with residues Q66 and R86 (apo-NocΔCTD structure, Fig. 6D) whose equivalent residues in *B. subtilis* ParB are known to be critical for CTP binding[2].

The expanding roles of CTP switches in biology

ATP and GTP switches that control membrane activity are widespread in all domains of life. For example, ATP binding promotes the dimerization of MinD (role in bacterial cell division site selection) and concomitantly increases its affinity for the cell membrane[30–33]. MinE, a partner of MinD, stimulates the ATPase activity of MinD, promoting MinD dissociation from the membrane[31,33–35]. In eukaryotes, both the Ras-related protein (Sar) and ADP-ribosylation factor (Arf) (role in vesicle trafficking) function as GTP-dependent switches, cycling between the active GTP-bound form and the inactive GDP-bound form[36–43]. In the GDP-bound form, the amphipathic helix of Sar/Arf1 adopts a repressed conformation by burying itself into a hydrophobic pocket[36,44]. The exchange of GDP for GTP induces conformational changes that push the myristoylated amphipathic helix out of the hydrophobic pocket, enabling membrane association[36,44]. Our study provides the first evidence for a CTP switch that controls membrane activity (Fig. 8), suggesting that CTP switches are likely to control more diverse functions in biology than previously appreciated.

FINAL PERSPECTIVES

Our work unveils a nucleotide-dependent regulatory layer, in addition to the previously described DNA-dependent regulation[21], in the activity of the nucleoid occlusion protein Noc. The dual dependency on nucleotide and DNA guarantees a productive formation of the tripartite DNA-protein-membrane super-complex for nucleoid occlusion. In this work, we also provide evidence for a CTP switch that controls membrane-binding activity, adding the control of membrane association in Noc to the role of ParB-CTP in bacterial chromosome segregation. It is likely that CTP switches are pervasive in biology but have so far been underappreciated[1,2,45]. Finally, evolution is replete with examples of functional domains being adapted to diversify functions of proteins. Gene encoding Noc resulted from gene duplication and neo-functionalization from *parB*[15–17], our work furthers the understanding of how a CTP switch has been adapted to a new function, and hence might have implications in understanding biological innovations by evolution.

ACKNOWLEDGEMENTS

This study was funded by the Royal Society University Research Fellowship Renewal (URF\R\201020 to T.B.K.L), and a Wellcome Investigator grant (209500) to Jeff Errington that supported L.J.W. A.S.B.J's PhD studentship was funded by the Royal Society (RG150448), and N.T.T was funded by the BBSRC grant-in-add (BBS/E/J/000C0683 to the John Innes Centre).

We thank Diamond Light Source for access to beamlines I04 and I04-1 under proposals MX18565 and MX25108 with support from the European Community's Seventh Framework Program (FP7/2007–2013) under Grant Agreement 283570 (BioStruct-X).

MATERIALS AND METHODS
Strains, media, and growth conditions
Escherichia coli and *B. subtilis* were grown in LB and CH medium[46], respectively. Carbenicillin (100 µg/mL), chloramphenicol (50 µg/mL), and tetracycline (12 µg/mL) were used for selection in *E. coli*, as required. Kanamycin (5 µg/mL), spectinomycin (50 µg/mL), and tetracycline (10 µg/mL) were used for selection in *B. subtilis*, as required. Xylose were added as needed at the concentration indicated.

Plasmid and strain construction
All plasmids and primers used in strain and plasmid construction are listed in Supplementary Table S1. All strains used in this study are listed in Supplementary Table S2.

Construction of pET21b::noc-his$_6$
A double-stranded DNA (dsDNA) fragment containing a codon-optimized *B. subtilis noc* gene was chemically synthesized (gBlocks, IDT). The pET21b plasmid backbone was generated via a double digestion of pET21b::*Caulobacter crescentus parB-(his)$_6$* with NdeI and HindIII[10]. The resulting backbone was subsequently gel-purified and assembled with the *noc* gBlocks fragment using a 2x Gibson master mix. Briefly, 2.5 µL of the gBlocks fragment and 2.5 µL of NdeI-HindIII-cut pET21b at equimolar concentration were added to 5 µL of a 2x Gibson master mix (NEB). The mixture was incubated at 50°C for 60 min. Subsequently, 5 µL was used to transform chemically competent *E. coli* DH5α cells. Gibson assembly was possible owing to a 23-bp sequence shared between the NdeI-HindIII-cut pET21b backbone and the gBlocks fragment. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pET21b::nocN∆10-his$_6$
The coding sequence of a 10-amino-acid N-terminally truncated Noc (NocN∆10) was amplified by PCR using primers AJ65 and AJ66, and pET21b::*noc-his$_6$* as template. The resulting PCR product was gel-purified and assembled into an NdeI-HindIII-cut pET21b using a 2x Gibson master mix. Gibson assembly was possible owing to a 23-bp sequence shared between the NdeI-and-HindIII cut pET21b backbone and the PCR amplified fragment. The 23-bp homologous region was introduced during the synthesis of primers AJ65 and AJ66. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pET21b::noc (R89A)-his$_6$
To introduce the R89A mutation into the coding sequence of Noc, primers P3296 and AJ73, and primers P3297 and AJ74 were used in PCR reactions to amplify the left half and the right half of *noc (R89A)*, respectively, from the pET21b::*noc-his$_6$* template. A 15-bp overlapping region between the two PCR fragments contained the point mutation and also enabled their assembly by a Gibson master mix. Briefly, 1.7 µL of each PCR-amplified DNA fragment and 1.6 µL of a gel purified NdeI-HindIII-cut pET21b at equimolar concentration were added to 5 µL of a 2x Gibson master mix. The mixture was incubated at 50°C for 60 min. Subsequently, 5 µL was used to transform chemically competent *E. coli* DH5α cells. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pET21b::noc (N121S)-his$_6$
The same procedure as above was used to introduce the N121S mutation into the coding sequence of Noc, except that primers P3296 and AJ87, and primers P3297 and AJ86 were used to amplify the left half and the right half of *noc (N121S)*, respectively, from the *pET21b::noc-his$_6$* template.

Construction of pET21b::noc (E29C)-his$_6$
The same procedure as above was used to introduce the E29C mutation into the coding sequence of Noc, except that primers P3296 and AJ85, and primers P3297 and AJ84 were used to amplify the left half and the right half of *noc (E29C)*, respectively, from the *pET21b::noc-his$_6$* template.

Construction of pET21b::noc (S253C)-his$_6$
The same procedure as above was used to introduce the S253C mutation into the coding sequence of Noc, except that primers P3296 and AJ89, and primers P3297 and AJ88 were used to amplify the left half and the right half of *noc (S253C)*, respectively, from the *pET21b::noc-his$_6$* template.

Construction of pET21b::noc (E29C R89A)-his$_6$
The same procedure as above was used to introduce the R89A mutation into the coding sequence of Noc (E29C), except that primers P3296 and AJ73, and primers P3297 and AJ74 were used to amplify the left half and the right half of *noc (E29C R89A)*, respectively, from the *pET21b::noc (E29C)-his$_6$* template.

Construction of pET21b::Noc (E29C N121S)-his$_6$
The same procedure as above was used to introduce the N121S mutation into the coding sequence of Noc (E29C), except that primers P3296 and AJ86, and primers P3297 and AJ87 were used to amplify the left half and the right half of *noc (E29C N121S)*, respectively, from the *pET21b::noc (E29C)-his$_6$* template.

Construction of pET21b::noc (E29C S253C)-his$_6$
The same procedure as above was used to introduce the E29C mutation into the coding sequence of Noc (S253C), except that primers P3296 and AJ85, and primers P3297 and AJ84 were used to amplify the left half and the right half of *noc (E29C S253C)*, respectively, from the *pET21b::noc (S253C)-his$_6$* template.

Construction of pET21b:: Geobacillus thermoleovorans NocΔCTD-his$_6$
A dsDNA fragment containing the coding sequence of a 42-amino-acid C-terminally truncated *G. thermoleovorans* Noc was chemically synthesized (gBlocks, IDT). The gBlocks fragment was assembled into an NdeI-HindIII-cut pET21b using a 2x Gibson master mix. Gibson assembly was possible owing to a 23-bp sequence shared between the NdeI-HindIII-cut pET21b backbone and the gBlocks fragment. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pET21b:: Geobacillus thermoleovorans NocNΔ26ΔCTD-his$_6$
The coding sequence of a 26-amino-acid N-terminally truncated and 42-amino-acid C-terminally truncated *G. thermoleovorans* Noc was amplified by PCR using primers AJ76 and AJ81, and pET21b:: *Geobacillus thermoleovo*rans NocΔCTD-his$_6$ as template. The resulting PCR product was gel-purified and assembled into an NdeI-HindIII-cut pET21b using a 2x Gibson master mix. Gibson assembly was possible owing to a 23-bp sequence shared

between the NdeI-and-HindIII cut pET21b backbone and the PCR amplified fragment. The 23-bp homologous region was introduced during the synthesis of primers AJ76 and AJ81. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pMCS5-4xNBS
A dsDNA fragment containing four *NBS* sites were chemically synthesized (gBlocks, IDT). The gBlocks fragment was assembled into an EcoRI-cut pMCS5[47] using a 2x Gibson master mix. Gibson assembly was possible owing to a 23-bp sequence shared between the EcoRI-cut pMCS5 backbone and the gBlocks fragment. The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pUC19-4xNBS
A 260-bp DNA region containing four *NBS* sites were amplified by PCR using primers 4xNBS-F and 4xNBS-R, and pMCS5-4x*NBS* as template. The resulting PCR product was gel-purified, 5'-phosphorylated using T4 PNK enzyme (NEB), and subsequently ligated into a SmaI-cut pUC19 backbone using T4 ligase (NEB). The resulting plasmid was verified by Sanger sequencing (Eurofins, Germany).

Construction of pSG4926N121S and B. subtilis strains harboring the noc(N121S)-yfp fusion
The N121S substitution was introduced into the *noc(WT)-myfp* plasmid pSG4926 by site-directed mutagenesis using PFU Turbo DNA polymerase (Agilent Technologies), and primers Noc(N121S)-F and Noc(N121S)-R. After verification of the mutation by DNA sequencing, the mutagenized plasmid (pSG4926N121S) was introduced into the *B. subtilis* Δnoc mutant (DWA117) by transformation[48] to generate strain 4746. To construct strain 4747, the *ΔminCminD* deletion was introduced into strain 4746 by transforming strain 4746 with the chromosomal DNA of strain DWA564, with selection for kanamycin resistance.

DNA preparation for *in vitro* assays
A 22-bp palindromic single-stranded DNA (ssDNA) oligomer (*NBS*: GGA<u>TATTTCCCGGGAAATA</u>TCC or *parS*: GGA<u>TGTTTCACGTGAAACA</u>TCC) [dissolved to 100 µM in buffer containing 1 mM Tris-HCl pH 8.0 and 5 mM NaCl] was heated at 98°C for 5 min before being left to cool down to room temperature overnight to form 50 µM *parS* or

NBS DNA duplex. The core sequences of *NBS* and *parS* are underlined.

Construction of biotinylated DNA substrates for bio-layer interferometry (BLI) analysis

DNA constructs were chemically synthesized (gBlocks dsDNA fragments, IDT) with M13F and M13R homologous regions at each end. To generate a dual biotin-labeled DNA substrate, PCR reactions were performed using a 2x GoTaq PCR master mix (Promega), biotin-labeled M13F and biotin-labeled M13R primers, and gBlocks fragments as template. PCR products were gel purified, and subsequently used in BLI assays.

Protein overexpression and purification

Plasmids pET21b::*noc-his$_6$* (WT or mutants) were introduced into *E. coli* Rosetta (DE3) competent cells by heat-shock transformation. A 10-mL overnight culture was used to inoculate 1 L of LB medium + carbenicillin + chloramphenicol. Cells were grown at 37°C with shaking at 250 rpm to an OD_{600} of ~0.4. The culture was then left to cool down to 4°C before isopropyl-β-D-thiogalactopyranoside (IPTG) was added to a final concentration of 1 mM. The culture was shaken for an additional 3 hrs at 28°C before the cells were pelleted by centrifugation. Pelleted cells were resuspended in a buffer containing 100 mM Tris-HCl pH 8.0, 250 mM NaCl, 10 mM imidazole, 5% (v/v) glycerol, 10 mg of lysozyme, and an EDTA-free protease inhibitor tablet (Merck). Cells were lyzed by sonication (10 cycles of 15 s with 10 s resting on ice in between each cycle). The cell debris was removed through centrifugation at 28,000 g for 30 min and the supernatant was filtered through a 0.45 µm filter (Sartorius). The lysate was then loaded into a 1-mL HisTrap column (GE Healthcare) that had been equilibrated with buffer A [100 mM Tris-HCl pH 8.0, 250 mM NaCl, 10 mM imidazole, and 5% (v/v) glycerol]. Protein was eluted from the column using an increasing gradient of imidazole (10 mM to 500 mM) in the same buffer. Noc-containing fractions were pooled together and diluted to a conductivity of 16 mS/cm before being loaded onto a 1-mL Heparin HP column (GE Healthcare) that had been equilibrated with 100 mM Tris-HCl pH 8.0, 25 mM NaCl, and 5% (v/v) glycerol. Protein was eluted from the Heparin column using an increasing gradient of NaCl (25 mM to 1 M) in the same buffer. Lastly, proteins were polished using a gel filtration column. To do so, Noc-containing fractions were concentrated using an Amicon Ultra-15 10 kDa cut-off spin filter (Merck) before being loaded onto a Superdex-75 gel filtration column (GE Healthcare). The gel filtration column was pre-equilibrated with buffer containing 10 mM Tris-HCl pH 8.0 and 250 mM NaCl. Eluted protein fractions were analyzed for purity by SDS-PAGE.

Noc variants that were used in crosslinking experiments were purified using a one-step Ni-affinity column, and all buffers were adjusted to pH 7.4 which was optimal for crosslinking reactions. Purified proteins were desalted using a PD-10 column (Merck), concentrated using an Amicon Ultra-4 10 kDa cut-off spin column (Merck), and stored at -80°C in a storage buffer [100 mM Tris-HCl pH 7.4, 250 mM NaCl, 10% (v/v) glycerol, and 1 mM TCEP].

Measurement of NTPase activity by EnzChek phosphate assay

NTP hydrolysis was monitored using an EnzChek Phosphate Assay Kit (Thermo Fisher). Samples (100 µL) containing a reaction buffer ± 0 to 1 mM of CTP ± 1 µM *NBS*/*parS* dsDNA + 1 µM of purified Noc (WT or mutants) were assayed in a Biotek EON plate reader for 15 hrs with readings every minute. The reaction buffer (1 mL) typically contained 640 µL ultrapure water, 100 µL 10x customized reaction buffer [100 mM Tris pH 8.0, 2 M NaCl, and 20 mM $MgCl_2$], 200 µL MESG substrate solution, and 10 µL purine nucleoside phosphorylase (1 unit). Reactions with buffer only, buffer + protein only or buffer + NTP only were also included as controls. The plates were shaken at 280 rpm continuously for 15 hrs at 25°C. The inorganic phosphate standard curve was also constructed according to the manual. Each assay was triplicated. The NTPase rates were calculated using a linear regression fitting in Excel.

In vitro crosslinking using a sulfhydryl-to-sulfhydryl crosslinker bismaleimidoethane (BMOE)

A 50 µL mixture of 10 µM Noc (WT/mutants) ± 1 mM NTP ± 1 µM 22-bp *NBS*/*parS* dsDNA was assembled in a reaction buffer [10 mM Tris-HCl pH 7.4, 200 mM NaCl, and 1 mM $MgCl_2$] and was incubated for 10 min at 22°C or for 1, 5, 10, 15, and 30 min at 4°C. Subsequently, BMOE was added to the final concentration of 1 mM, and the reaction was quickly mixed by three

pulses of vortexing. SDS-PAGE sample buffer containing 23 mM β-mercaptoethanol was then added immediately to quench the crosslinking reaction. Samples were heated to 50°C for 10 min before being loaded on 12% WedgeWell Tris-Glycine polyacrylamide gels (Thermo Fisher). Each experiment was triplicated. Polyacrylamide gels were stained in an InstantBlue Coomassie solution (Abcam) and band intensity was quantified using Image Studio Lite (LI-COR Biosciences). The crosslinked fractions were averaged, and their standard errors were calculated in Excel.

DNA loading assays were performed similarly to the *in vitro* BMOE crosslinking experiments above. A 50 µL mixture of 10 µM Noc (E29C S253C) ± 1 mM CTP ± 100 nM *NBS*-harboring/*NBS*-scrambled plasmid was assembled in a reaction buffer [10 mM Tris-HCl pH 7.4, 200 mM NaCl, and 1 mM MgCl$_2$] and was incubated for 10 min at 22°C. Subsequently, BMOE was added to the final concentration of 1 mM, and the reaction was quickly mixed by three pulses of vortexing. SDS-PAGE sample buffer containing 23 mM β-mercaptoethanol was then added immediately to quench the crosslinking reaction. Samples were heated to 50°C for 10 min before being loaded on 4-12% WedgeWell Tris-Glycine polyacrylamide gels.

For nuclease treatment, crosslinking reactions were performed as described above, however the reactions were quenched using a quenching buffer [10 mM Tris-HCl pH 7.4, 200 mM NaCl, 1 mM MgCl$_2$, and 2.3 mM β-mercaptoethanol] instead. Subsequently, 0.5 µL of a non-specific DNA nuclease (Benzonase, 250 units/µL, Merck) was added, the mixture was incubated at 22°C for 10 min before SDS-PAGE sample buffer was added. Samples were heated to 50°C for 10 min before being loaded on 4-12% WedgeWell Tris-Glycine polyacrylamide gels. Gels were either stained in an InstantBlue Coomassie solution (to detect protein bands) or in a Sybr Green solution (to detect DNA bands).

CTPγS was custom synthesized either in-house[49] or by Jena Biosciences.

Measurement of protein-DNA interactions by bio-layer interferometry (BLI)
Bio-layer interferometry (BLI) experiments were conducted using a BLItz system equipped with Dip-and-Read Streptavidin (SA) Biosensors

(Molecular Devices). BLItz monitors wavelength shifts (nm) resulting from changes in the optical thickness of the sensor surface during association or dissociation of the analyte. All BLI experiments were performed at 22°C. Briefly, the streptavidin biosensor was hydrated in a low-salt binding buffer [100 mM Tris-HCl pH 8.0, 150 mM NaCl, 1 mM MgCl$_2$, and 0.005% Tween 20] for at least 10 min before each experiment. Biotinylated double-stranded DNA (dsDNA) was immobilized onto the surface of the SA sensor through a cycle of Baseline (30 s), Association (120 s), and Dissociation (120 s). Briefly, the tip of the biosensor was dipped into a binding buffer for 30 s to establish the baseline, then to 1 µM biotinylated dsDNA for 120 s, and finally to a low salt binding buffer for 120 s to allow for dissociation. After the immobilization of DNA on the sensor, association reactions were monitored at 1 µM dimer concentration of Noc (WT/mutants) (with or without 1mM NTP) for 600 s. At the end of each binding step, the sensor was transferred into a protein-free binding buffer to follow the dissociation kinetics for 600 s. The sensor can be recycled by dipping in a high-salt buffer [100 mM Tris-HCl pH 8.0, 2 M NaCl, 1mM EDTA, and 0.005% Tween 20] for 20 min to remove bound Noc.

For experiments where a closed DNA substrate was cleaved to generate a free DNA end, DNA-coated sensors were dipped into 300 µL of restriction solution [266 µL of water, 30 µL of 10x CutSmart buffer (NEB), and 4 µL of *Eco*RI-HF or *Bam*HI-HF restriction enzyme (20,000 units/mL)] for 30 min at 37°C.

Measurement of Noc-CTPγS interaction by isothermal titration calorimetry (ITC)
All ITC experiments were recorded using a MicroCal PEAQ ITC instrument (Malvern Panalytical, UK). Experiments were performed at 4°C and both protein and ligand were in the buffer 100 mM Tris-HCl pH 8.0, 100 mM NaCl, and 5 mM CaCl$_2$. The calorimetric cell was filled with 100 µM monomer concentration of either *B. subtilis* Noc (WT), Noc (R89A), or Noc (N121S), and was titrated with 3 mM CTPγS. For each ITC run, a single injection of 0.5 µL of 3 mM CTPγS was performed first, following by 19 injections of 2 µL each. Injections were made at 120 s intervals with a stirring speed of 750 rpm. Each experiment was run in duplicate. The raw titration data were integrated and fitted to a one-site binding model using the built-in software of

the MicroCal PEAQ ITC. Controls (CTPγS into buffer and buffer into protein) were performed and no signal was observed.

Liposomes preparation

E. coli total lipid extract (25 mg/mL in chloroform, Avanti) was used to generate model liposomes. Briefly, an argon stream was used to evaporate chloroform from the lipids, and the resulting lipid cake was further dried under vacuum for 2 hrs. The lipids were subsequently re-suspended in a buffer containing 100 mM Tris-HCl pH 7.4 and 200 mM NaCl. The mixture was incubated at 30°C for 30 min with vigorous vortexing every 5 min. The final concentration of the resuspended lipids was 50 mg/mL. The resuspended lipids were then extruded to ~100 nm single unilamellar vesicles (SUV) using a mini-extruder (Avanti) equipped with polycarbonate membranes (0.1 μm pore size). The size of the resulting SUVs was confirmed by dynamic light scattering.

Liposome sedimentation assays

A 500 μL mixture of 0.75 μM Noc (WT/mutants) ± 1.0 mg/mL liposomes ± 1 mM NTP ± DNA (either 1 μM of 22-bp *NBS*/*parS* dsDNA or 100 nM of *NBS*-harboring/empty plasmid) was assembled in a binding buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, and 1 mM MgCl$_2$]. The mixture was incubated at 22°C for 20 min before being centrifuged at 50,000 rpm for 20 min at 22°C (TLA120.2 rotor, Optima Max-E Benchtop Ultracentrifuge). After centrifugation, the supernatant was transferred to a new 1.5-mL Eppendorf tube. The pellet was resuspended in 500 μL of binding buffer before being transferred to another 1.5-mL Eppendorf tube. SDS-PAGE sample buffer was then added, and the samples were heated at 70°C for 5 min before being loaded onto either 12% WedgeWell Tris-Glycine polyacrylamide gels, Novex 20% TBE polyacrylamide gels, or 1% agarose gels. Gels were subsequently stained in an InstantBlue Coomassie solution (to detect protein bands) or in a Sybr Green solution (to detect DNA bands). Each assay was triplicated. Protein/DNA band intensity was quantified using Image Studio Lite (LI-COR Biosciences). The protein/DNA fractions were averaged, and their standard errors were calculated in Excel.

For nuclease treatment, a 500 μL mixture of 0.75 μM Noc (WT/mutants) ± 1.0 mg/mL liposomes ±

1 mM NTP ± 100 nM of *NBS*-harboring/empty plasmid was incubated at room temperature for 10 min. Afterwards, 1 μL of Benzonase (250 units/ μL) was added, and the mixture was incubated for a further 10 min at 22°C before ultracentrifugation.

For re-sedimentation experiments (Fig. 5D), the pellet (from the first round of ultracentrifugation) was resuspended either in 500 μL of binding buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, and 1 mM MgCl$_2$] or in a stripping buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, and 10 mM EDTA]. The resuspended pellet was centrifuged for the second time at 50,000 rpm for 20 min at 22°C. After the second centrifugation, the supernatant was transferred to a new 1.5-mL Eppendorf tube. The pellet was resuspended in 500 μL of binding buffer before being transferred to another 1.5-mL Eppendorf tube. SDS-PAGE sample buffer was then added to the supernatant and the pellet fractions, and the samples were analyzed on denaturing polyacrylamide gels.

Liposome flotation assays

A 200 μL mixture of 0.75 μM Noc (WT/mutants) ± 1.0 mg/mL liposomes ± 1 mM NTP ± 20 nM *NBS*-harboring/empty plasmid was assembled in a 30% sucrose buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, 1 mM MgCl$_2$, and 30% sucrose]. The mixture was incubated at 22°C for 5 min before being overlaid with 250 μL of a 25% sucrose buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, 1 mM MgCl$_2$, and 25% sucrose]. Finally, 150 μL of a 0% sucrose buffer [100 mM Tris-HCl pH 7.4, 200 mM NaCl, and 1 mM MgCl$_2$] was added as the top layer. The solution was incubated for 15 min at 22°C before being centrifuged at 70,000 rpm at 22°C for 20 min (TLA120.2 rotor, Optima Max-E Benchtop Ultracentrifuge). After centrifugation, three equal fractions (200 μL each) were gently drawn sequentially from bottom of the ultracentrifugation tube using a Hamilton syringe. SDS-PAGE sample buffer was added to each fraction, and samples were heated at 70°C for 5 min before being loaded onto either 12% WedgeWell Tris-Glycine polyacrylamide gels or 1% agarose gels. Gels were subsequently stained in an InstantBlue Coomassie solution (to detect protein bands) or in a Sybr Green solution (to detect DNA bands). Each assay was triplicated. Protein/DNA band intensity was quantified using Image Studio Lite (LI-COR

Biosciences). The protein/DNA fractions were averaged, and their standard errors were calculated in Excel.

Protein crystallization, structure determination and refinement

Crystallization screens were set up in sitting-drop vapor diffusion format in MRC2 96-well crystallization plates with drops comprised of 0.3 µL precipitant solution and 0.3 µL of protein and incubated at 293 K. After optimization of initial hits, suitable crystals were cryoprotected and mounted in Litholoops (Molecular Dimensions) before flash-cooling by plunging into liquid nitrogen. X-ray data were recorded either on beamline I04 or I04-1 at the Diamond Light Source (Oxfordshire, UK) using either an Eiger2 XE 16M or a Pilatus 6M-F hybrid photon counting detector (Dectris), with crystals maintained at 100 K by a Cryojet cryocooler (Oxford Instruments). Diffraction data were integrated and scaled using DIALS[50] via the XIA2 expert system[51] then merged using AIMLESS[52]. Data collection statistics are summarized in Supplementary Table S3. The majority of the downstream analysis was performed through the CCP4i2 graphical user interface[53].

Geobacillus thermoleovorans NocΔCTD Noc: His-tagged NocΔCTD Noc (~10 mg/mL) was premixed with 1 mM CTP and 1 mM MgCl₂ in buffer [10 mM Tris-HCl pH 8.0 and 250 mM NaCl] before crystallization. Crystals grew in a solution containing 2.0 M ammonium sulphate and 50 mM tri-sodium citrate and were cryoprotected in the crystallisation solution supplemented with 20% (v/v) glycerol, 1 mM CTP and 1 mM MgCl₂. For iodide derivatization, the cryoprotectant comprised the crystallization solution supplemented with 25% (v/v) ethylene glycol, 1 mM CTP, 1 mM MgCl₂, and 500 mM potassium iodide; crystals were soaked in this solution for less than 30 seconds before cryo-cooling. Three 360° passes of X-ray data were taken at a wavelength of 1.8 Å from different parts of a single crystal and merged to give a highly redundant dataset to 3.4 Å resolution in space group $P2_13$ with cell parameters $a = b = c = 136.6$ Å. Solvent content estimation gave a value of 66% for two copies of the 29 kDa subunit per asymmetric unit (ASU). The structure was subsequently solved by single-wavelength anomalous dispersion using the CRANK2

pipeline[54], which located 12 iodide sites, and BUCCANEER[55] was able to build and sequence 339 residues in two chains corresponding to 67% of those expected for two monomers, giving R_{work} and R_{free} values of 0.318 and 0.362, respectively, to 3.4 Å resolution after refinement with REFMAC5[56]. At this point, this preliminary model was used as a starting point for refinement against a native data set processed to 2.5 Å resolution in the same space group, but with a significantly longer cell edge of 146.8 Å corresponding to a solvent content of 72.6%. Thus, it was necessary to resolve the structure by molecular replacement using PHASER[57] before further refinement in REFMAC5. After a complete rebuild in BUCCANEER, several iterations of model building in COOT[58] and REFMAC5 refinement jobs yielded the final model with R_{work} and R_{free} values of 0.210 and 0.240, respectively, to 2.5 Å resolution.

Despite the presence of CTP in the crystallization buffer, no density for CTP was found in the NocΔCTD structure, presumably because *NBS* DNA was not included to facilitate CTP-binding or the high concentrations of sulfate (2M) from the precipitant excluding the ligand.

Geobacillus thermoleovorans NocNΔ26ΔCTD: His-tagged NocNΔ26ΔCTD Noc (~10 mg/mL) was premixed with 1 mM CTPγS and 1 mM MgCl₂ in buffer [10 mM Tris-HCl pH 8.0 and 250 mM NaCl] before crystallization. Crystals grew in a solution containing 0.2 M di-ammonium phosphate and 2.3 M ammonium sulphate and were cryoprotected in this solution supplemented with 20% (v/v) glycerol. X-ray data were processed to a resolution of 2.95 Å in space group $C222_1$ with cell parameters of $a = 105.1$, $b = 106.6$, $c = 42.2$ Å. Analysis of the likely composition of the ASU suggested that it contained a single copy of the 26 kDa NocNΔ26ΔCTD Noc monomer, giving an estimated solvent content of 46%. The structure was solved by molecular replacement with PHASER[57] using chain A from the *G. thermoleovorans* NocΔCTD structure above as the template. The search model was split into three separate ensembles comprising residues 26-100, 101-140, and 141-230, respectively. PHASER successfully placed all three ensembles, although one of these had to be interchanged with a symmetry mate in order to restore the connectivity of the starting template.

Several iterations of model building in COOT and refinement REFMAC5 yielded the final model with R_{work} and R_{free} values of 0.267 and 0.288, respectively, to 2.95 Å resolution.

Fluorescence microscopy

Cells containing fluorescent protein fusions were grown at 30°C. Xylose (0.5% w/v) was included in the media to induce the expression of YFP fusions in *B. subtilis*. Cell membranes were stained by mixing 10 µL of culture with 0.2 µL of membrane dye FM5-95 (200 µg/ml, Invitrogen). Cells were mounted on microscope slides covered with a thin agarose pad (1.2% w/v in dH$_2$O) and images were acquired with a Rolera EM-C2 (Q-imaging) camera attached to a Nikon Ti microscope using METAMORPH version 6 (Molecular Devices), with an exposure time of 400 ms for YFP and 1000 ms for membranes. Images were prepared for publication using ImageJ.

REFERENCES

1. Osorio-Valeriano, M. *et al.* ParB-type DNA Segregation Proteins Are CTP-Dependent Molecular Switches. *Cell* **179**, 1512-1524.e15 (2019).
2. Soh, Y.-M. *et al.* Self-organization of parS centromeres by the ParB CTP hydrolase. *Science* **366**, 1129–1133 (2019).
3. Jalal, A. S. & Le, T. B. Bacterial chromosome segregation by the ParABS system. *Open Biology* **10**, 200097 (2020).
4. Breier, A. M. & Grossman, A. D. Whole-genome analysis of the chromosome partitioning and sporulation protein Spo0J (ParB) reveals spreading and origin-distal sites on the Bacillus subtilis chromosome. *Molecular Microbiology* **64**, 703–718 (2007).
5. Murray, H., Ferreira, H. & Errington, J. The bacterial chromosome segregation protein Spo0J spreads along DNA from parS nucleation sites. *Molecular Microbiology* **61**, 1352–1361 (2006).
6. Graham, T. G. W. *et al.* ParB spreading requires DNA bridging. *Genes Dev.* **28**, 1228–1238 (2014).
7. Sanchez, A. *et al.* Stochastic Self-Assembly of ParB Proteins Builds the Bacterial DNA Segregation Apparatus. *Cell Syst.* **1**, 163–173 (2015).
8. Funnell, B. E. ParB Partition Proteins: Complex Formation and Spreading at Bacterial and Plasmid Centromeres. *Front Mol Biosci* **3**, 44 (2016).
9. Jalal, A. S., Tran, N. T. & Le, T. B. ParB spreading on DNA requires cytidine triphosphate in vitro. *eLife* **9**, e53515 (2020).
10. Lim, H. C. *et al.* Evidence for a DNA-relay mechanism in ParABS-mediated chromosome segregation. *Elife* **3**, e02758 (2014).
11. Hwang, L. C. *et al.* ParA-mediated plasmid partition driven by protein pattern self-organization. *The EMBO Journal* **32**, 1238–1249 (2013).
12. Vecchiarelli, A. G., Neuman, K. C. & Mizuuchi, K. A propagating ATPase gradient drives transport of surface-confined cellular cargo. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 4880–4885 (2014).
13. Vecchiarelli, A. G., Mizuuchi, K. & Funnell, B. E. Surfing biological surfaces: exploiting the nucleoid for partition and transport in bacteria. *Molecular Microbiology* **86**, 513–523 (2012).
14. Vecchiarelli, A. G., Hwang, L. C. & Mizuuchi, K. Cell-free study of F plasmid partition provides evidence for cargo transport by a diffusion-ratchet mechanism. *Proc. Natl. Acad. Sci. U.S.A.* **110**, E1390-1397 (2013).
15. Sievers, J., Raether, B., Perego, M. & Errington, J. Characterization of the parB-Like yyaA Gene of Bacillus subtilis. *J. Bacteriol.* **184**, 1102–1111 (2002).
16. Wu, L. J. & Errington, J. Nucleoid occlusion and bacterial cell division. *Nat. Rev. Microbiol.* **10**, 8–12 (2011).
17. Jalal, A. S. B. *et al.* Diversification of DNA-Binding Specificity by Permissive and Specificity-Switching Mutations in the ParB/Noc Protein Family. *Cell Rep* **32**, 107928 (2020).
18. Wu, L. J. & Errington, J. Coordination of cell division and chromosome segregation by a nucleoid occlusion protein in Bacillus subtilis. *Cell* **117**, 915–925 (2004).
19. Veiga, H., Jorge, A. M. & Pinho, M. G. Absence of nucleoid occlusion effector Noc impairs formation of orthogonal FtsZ rings during Staphylococcus aureus cell division. *Mol. Microbiol.* **80**, 1366–1380 (2011).
20. Pang, T., Wang, X., Lim, H. C., Bernhardt, T. G. & Rudner, D. Z. The nucleoid occlusion factor Noc controls DNA replication initiation in Staphylococcus aureus. *PLOS Genetics* **13**, e1006908 (2017).
21. Adams, D. W., Wu, L. J. & Errington, J. Nucleoid occlusion protein Noc recruits DNA

to the bacterial cell membrane. *EMBO J.* **34**, 491–501 (2015).

22. Rodrigues, C. D. A. & Harry, E. J. The Min System and Nucleoid Occlusion Are Not Required for Identifying the Division Site in Bacillus subtilis but Ensure Its Efficient Utilization. *PLOS Genetics* **8**, e1002561 (2012).

23. Yu, Y., Zhou, J., Gueiros-Filho, F. J., Kearns, D. B. & Jacobson, S. C. Noc Corrals Migration of FtsZ Protofilaments during Cytokinesis in Bacillus subtilis. *mBio* **12**, (2021).

24. Wu, L. J. *et al.* Noc protein binds to specific DNA sequences to coordinate cell division with chromosome segregation. *EMBO J.* **28**, 1940–1952 (2009).

25. Leonard, T. A., Butler, P. J. G. & Löwe, J. Structural analysis of the chromosome segregation protein Spo0J from Thermus thermophilus. *Mol. Microbiol.* **53**, 419–432 (2004).

26. Toro, E., Hong, S.-H., McAdams, H. H. & Shapiro, L. Caulobacter requires a dedicated mechanism to initiate chromosome segregation. *PNAS* **105**, 15435–15440 (2008).

27. Krissinel, E. Stock-based detection of protein oligomeric states in jsPISA. *Nucleic Acids Res.* **43**, W314-319 (2015).

28. Buckstein, M. H., He, J. & Rubin, H. Characterization of Nucleotide Pools as a Function of Physiological State in Escherichia coli. *J Bacteriol* **190**, 718–726 (2008).

29. Nyholm, T. K. M., Özdirekcan, S. & Killian, J. A. How Protein Transmembrane Segments Sense the Lipid Environment. *Biochemistry* **46**, 1457–1465 (2007).

30. Hu, Z. & Lutkenhaus, J. A conserved sequence at the C-terminus of MinD is required for binding to the membrane and targeting MinC to the septum. *Mol. Microbiol.* **47**, 345–355 (2003).

31. Hu, Z., Gogol, E. P. & Lutkenhaus, J. Dynamic assembly of MinD on phospholipid vesicles regulated by ATP and MinE. *PNAS* **99**, 6761–6766 (2002).

32. Wu, W., Park, K.-T., Holyoak, T. & Lutkenhaus, J. Determination of the structure of the MinD-ATP complex reveals the orientation of MinD on the membrane and the relative location of the binding sites for MinE and MinC. *Mol Microbiol* **79**, 1515–1528 (2011).

33. Ramm, B., Heermann, T. & Schwille, P. The E. coli MinCDE system in the regulation of protein patterns and gradients. *Cell. Mol. Life Sci.* **76**, 4245–4273 (2019).

34. Lutkenhaus, J. The ParA/MinD family puts things in their place. *Trends Microbiol* **20**, 411–418 (2012).

35. Park, K.-T., Wu, W., Lovell, S. & Lutkenhaus, J. Mechanism of the asymmetric activation of the MinD ATPase by MinE. *Mol Microbiol* **85**, 271–281 (2012).

36. Zhukovsky, M. A., Filograna, A., Luini, A., Corda, D. & Valente, C. Protein Amphipathic Helix Insertion: A Mechanism to Induce Membrane Fission. *Front. Cell Dev. Biol.* **7**, (2019).

37. Lee, M. C. S. *et al.* Sar1p N-Terminal Helix Initiates Membrane Curvature and Completes the Fission of a COPII Vesicle. *Cell* **122**, 605–617 (2005).

38. Krauss, M. *et al.* Arf1-GTP-induced tubule formation suggests a function of Arf family proteins in curvature acquisition at sites of vesicle budding. *J Biol Chem* **283**, 27717–27723 (2008).

39. Hanna, M. G. *et al.* Sar1 GTPase Activity Is Regulated by Membrane Curvature. *J Biol Chem* **291**, 1014–1027 (2016).

40. Dodonova, S. O. *et al.* 9Å structure of the COPI coat reveals that the Arf1 GTPase occupies two contrasting molecular environments. *Elife* **6**, (2017).

41. Beck, R. *et al.* Membrane curvature induced by Arf1-GTP is essential for vesicle formation. *PNAS* **105**, 11731–11736 (2008).

42. Bielli, A. *et al.* Regulation of Sar1 NH2 terminus by GTP binding and hydrolysis promotes membrane deformation to control COPII vesicle fission. *J Cell Biol* **171**, 919–924 (2005).

43. Bos, J. L., Rehmann, H. & Wittinghofer, A. GEFs and GAPs: Critical Elements in the Control of Small G Proteins. *Cell* **129**, 865–877 (2007).

44. Goldberg, J. Structural Basis for Activation of ARF GTPase: Mechanisms of Guanine Nucleotide Exchange and GTP–Myristoyl Switching. *Cell* **95**, 237–248 (1998).

45. Basu, M. K. & Koonin, E. V. Evolution of eukaryotic cysteine sulfinic acid reductase, sulfiredoxin (Srx), from bacterial chromosome partitioning protein ParB. *Cell Cycle* **4**, 947–952 (2005).

46. Partridge, S. R. & Errington, J. The importance of morphological events and

intercellular interactions in the regulation of prespore-specific gene expression during sporulation in Bacillus subtilis. *Mol Microbiol* **8**, 945–955 (1993).

47. Thanbichler, M., Iniesta, A. A. & Shapiro, L. A comprehensive set of plasmids for vanillate- and xylose-inducible gene expression in Caulobacter crescentus. *Nucl. Acids Res.* **35**, e137–e137 (2007).

48. Jenkinson, H. F. Altered arrangement of proteins in the spore coat of a germination mutant of Bacillus subtilis. *J Gen Microbiol* **129**, 1945–1958 (1983).

49. Rejzek, M. & Le, T. B. K. Chemical synthesis and purification of a non-hydrolyzable CTP analog CTPγS. *bioRxiv* 2021.01.13.426546 (2021) doi:10.1101/2021.01.13.426546.

50. Winter, G. *et al.* DIALS: implementation and evaluation of a new integration package. *Acta Crystallogr D Struct Biol* **74**, 85–97 (2018).

51. Winter, G. xia2: an expert system for macromolecular crystallography data reduction. *J Appl Cryst* **43**, 186–190 (2010).

52. Evans, P. R. & Murshudov, G. N. How good are my data and what is the resolution? *Acta Cryst D* **69**, 1204–1214 (2013).

53. Potterton, L. *et al.* CCP4i2: the new graphical user interface to the CCP4 program suite. *Acta Crystallogr D Struct Biol* **74**, 68–84 (2018).

54. Skubák, P. & Pannu, N. S. Automatic protein structure solution from weak X-ray data. *Nat Commun* **4**, 2777 (2013).

55. Cowtan, K. The Buccaneer software for automated model building. 1. Tracing protein chains. *Acta Crystallogr. D Biol. Crystallogr.* **62**, 1002–1011 (2006).

56. Murshudov, G. N., Vagin, A. A. & Dodson, E. J. Refinement of Macromolecular Structures by the Maximum-Likelihood Method. *Acta Cryst D* **53**, 240–255 (1997).

57. McCoy, A. J. *et al.* Phaser crystallographic software. *J Appl Crystallogr* **40**, 658–674 (2007).

58. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Crystallogr. D Biol. Crystallogr.* **60**, 2126–2132 (2004).

59. Williams, C. J. *et al.* MolProbity: More and better reference data for improved all-atom structure validation. *Protein Sci* **27**, 293–315 (2018).

SUPPLEMENTARY FILE S1. DNA, plasmids, and oligos

Plasmids	Description	Source
pET21b::*Bacillus subtilis noc-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc, carbenicillin^R	This study
pET21b::*Bacillus subtilis nocNΔ10-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc that lacks the first 10 amino acids, carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (R89A)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (R89A), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (N121S)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (N121S), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (E29C)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (E29C), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (S253C)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (S253C), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (E29C R89A)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (E29C R89A), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (E29C N121S)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (E29C N121S), carbenicillin^R	This study
pET21b::*Bacillus subtilis noc (E29C S253C)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (E29C S253C), carbenicillin^R	This study
pET21b::*Bacillus subtilis nocNΔ10 (E29C)-his₆*	Overexpression of a C-terminally His$_6$-tagged *B. subtilis* Noc (E29C) variant that also lacks the first 10 amino acids, carbenicillin^R	This study
pET21b::*Geobacillus thermoleovorans NocΔCTD*	Overexpression of a C-terminally His$_6$-tagged *G. thermoleovorans* Noc that lacks the last 42 amino acids, carbenicillin^R	This study
pET21b::*Geobacillus thermoleovorans NocNΔ26ΔCTD*	Overexpression of a C-terminally His$_6$-tagged *G. thermoleovorans* Noc that lacks the first 26 amino acids and the last 42 amino acids, carbenicillin^R	This study
pMCS5-4x*NBS*	pMCS5 plasmid that harbors four *NBS* sites, tetracycline^R	This study
pMCS5::empty	pMCS5 plasmid with an intact multiple cloning site, tetracycline^R	[47]
pUC19-4x*NBS*	pUC19 plasmid that harbors four *NBS* sites, carbenicillin^R	This study
pUC19-scrambled *NBS*	pUC19 plasmid that harbors four scrambled *NBS* sites, carbenicillin^R	This study
pET21b::*Caulobacter crescentus parB-his₆*	Overexpression of a C-terminally His$_6$-tagged *C. crescentus* ParB, carbenicillin^R	Gift from Christine Jacobs-Wagner [10]
pSG4926	*bla amyE' spc Pxyl-noc-yfpmut1 'amyE*; shuttle plasmid harboring a xylose-inducible *noc-yfp* fusion for integration at the *B. subtilis amyE* locus	[24]
pSG4926N121S	*bla amyE' spc Pxyl-noc (N121S)-yfpmut1 'amyE*; *B. subtilis* plasmid harboring a xylose-inducible *noc (N121S)-yfp* fusion for integration at the *B. subtilis amyE* locus	This study
Oligos/gBlocks		
170bp-*parS*	CGCCAGGGTTTTCCCAGTCACGACGTTGTAAAACGACGGCCAGAATTCGCAACGTG<u>TGTTTCACGTGAAAC</u>AGCCTTGAACTGATAACGACTCTATCATTGATAGAGTGTTCTCTCCACG**GGATCC**CCAGGCATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCT	[9]
170bp-*NBS*	CGCCAGGGTTTTCCCAGTCACGACGTTGTAAAACGACGGCCAGAATTCGCAACGTG<u>TATTTCCCGGGAAATA</u>GCCTTGAACTGATAACGACTCTATCATTGATAGAGTGTTCTCTCCACG**GGATCC**CCAGGC	This study

	ATGCAAGCTTGGCGTAATCATGGTCATAGCTGTTTCCT		The
22bp-*parS*	GGAT̲GTTTCACGTGAAACATCC	[17]	
22bp-*NBS*	GGAT̲ATTTCCCGGGAAATATCC	This study	
260bp-4x*NBS* gBlocks	GCCCAGGCCCTGGAGCGCATCTCCGGCT̲ATTTCCCGGGAAATA̲ACGTTCTTGGACGGGTCTTTTCGACCTTCTGTATCGGCAAGTGAT̲ATTTCCCGGGAAATA̲TCCCCAATATTGTCCACAGGCCGCTCACAGCTGCGGTGGGGGT̲ATTTCCCGGGAAATA̲CCGATGAGTCACATCGACCCGCTCGCCTGATTGGCGTATAGATCT̲ATTTCCCGGGAAATA̲CTCGAACCCAGGGGCGTTCGCATTGAGGCGAGCGTCTTG	This study	
4x*NBS*-F	GCCCAGGCCCTGGAGCGCATCTC	This study	
4x*NBS*-R	CAAGACGCTCGCCTCAATGCGAAC		
AJ65	TAACTTTAAGAAGGAGATATACATATGTTGGGTGAAAAGGAGCAAGAACCG	This study	
AJ66	GGTGGTGCTCGAGTGCGGCCGCAAGCTTTTTTGGAATACGGATTGTAAGCTG	This study	
AJ73	CTGAACCGCGCGCCACGCGCGTTCTCCCGCAAT	This study	
AJ74	ATTGCGGGAGAACGCGCGTGGCGCGCGGTTCAG	This study	
AJ76	GGTGGTGCTCGAGTGCGGCCGCAAGCTTGTCGCGGGAAAAGGCTTTACGTTTAGG	This study	
AJ81	TAACTTTAAGAAGGAGATATACATATGGAAGAGGTCCGTCACATCCCCGTCAAA	This study	
AJ84	GATACCAATAAGGAATGCATTTTAGAAATTCCA	This study	
AJ85	TGGAATTTCTAAAATGCATTCCTTATTGGTATC	This study	
AJ86	GTGGCCTTAATTGAGTCTTTGCAACGCGAGGAG	This study	
AJ87	CTCCTCGCGTTGCAAAGACTCAATTAAGGCCAC	This study	
AJ88	AATACTATTCGTCAGTGCTTATCAATGGTGGAA	This study	
AJ89	TTCCACCATTGATAAGCACTGACGAATAGTATT	This study	
M13-F	CGCCAGGGTTTTCCCAGTCACGAC	Lab stock	
M13-R	ATGGTCATAGCTGTTTCCT	Lab stock	
P3296	TTTAACTTTAAGAAGGAGATATACAT	This study	
P3297	TGGTGCTCGAGTGCGGCCGCAAGCTT	This study	
Noc(N121S)-F	GTGGCGTTAATCGAAAGCTTGCAGCGGGAAGAA	This study	
Noc(N121S)-R	TTCTTCCCGCTGCAAGCTTTCGATTAACGCCAC	This study	

sequences of *parS/NBS* are underlined. The sequence of BamHI recognition site is bold.

SUPPLEMENTARY FILE S2. Bacterial strains

Strains	Description	Source
DWA117	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet*	[21]
DWA206	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΩamyE::[spc Pxyl-noc (WT)-myfp)*	[21]
DWA382	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΩamyE::[spc Pxyl-nocNΔ10-myfp)*	[21]
DWA546	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΩamyE::[spc Pxyl-nco (R89A)-myfp)*	[21]
4746	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΩamyE::[spc Pxyl-nco (N121S)-yfpmut1)*	This study
DWA564	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΔminCD::kan ΩamyE::[spc Pxyl-noc (WT)-myfp)*	[21]
DWA566	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΔminCD::kan ΩamyE::[spc Pxyl-nocNΔ10-myfp)*	[21]
DWA600	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΔminCD::kan ΩamyE::[spc Pxyl-nco (R89A)-myfp)*	[21]
4747	*B. subtilis* 168CA (*trpC2*) *Δnoc::tet ΔminCD::kan ΩamyE::[spc Pxyl-noc (N121S)-yfpmut1)*	This study

SUPPLEMENTARY FILE S3. X-ray data collection and processing statistics

Structure	*G. thermoleovorans* NocΔCTD - iodide	*G. thermoleovorans* NocΔCTD - native	*G. thermoleovorans* NocNΔ26ΔCTD
Data collection			
Diamond Light Source beamline	I04	I04-1	I04
Wavelength (Å)	1.800	0.912	0.980
Detector	Eiger2 XE 16M	Pilatus 6M-F	Eiger2 XE 16M
Resolution range (Å)	96.61 – 3.40 (3.67 – 3.40)	84.89 – 2.50 (2.60 – 2.50)	37.44 – 2.95 (3.13 – 2.95)
Space Group	$P2_1 3$	$P2_1 3$	$C222_1$
Cell parameters (Å/°)	$a = b = c = 136.6$	$a = b = c = 146.8$	$a = 105.1$, $b = 106.6$, $c = 42.2$
Total no. of measured intensities	2147484 (426036)	1458851 (163435)	66966 (10641)
Unique reflections	11990 (2445)	36704 (4119)	5285 (835)
Multiplicity	179.1 (174.2)	39.7 (39.7)	12.6 (12.0)
Mean $I/\sigma(I)$	15.9 (3.3)	28.5 (1.8)	5.5 (1.5)
Completeness (%)	100.0 (100.0)	100.0 (100.0)	100.0 (100.0)
R_{merge}[a]	0.500 (3.392)	0.091 (2.654)	0.281 (1.343)
R_{meas}[b]	0.501 (3.402)	0.093 (2.688)	0.293 (1.399)
$CC_{½}$[c]	0.999 (0.894)	1.000 (0.670)	0.997 (0.885)
Wilson B value (Å^2)	88.7	68.0	38.2
Refinement			
Resolution range (Å)	-	84.89 – 2.50 (2.57 – 2.50)	37.44 – 2.95 (3.03 – 2.95)
Reflections: working/free[d]	-	34843/1794	4752/522
R_{work}[e]	-	0.210 (0.328)	0.267 (0.438)
R_{free}[e]	-	0.240 (0.386)	0.288 (0.443)
Ramachandran plot: favored/allowed/disallowed[f] (%)	-	98.1/1.9/0.0	98.0/2.0/0.0
R.m.s. bond distance deviation (Å)	-	0.010	0.007
R.m.s. bond angle deviation (°)	-	1.55	1.33
Mean B factors: protein/sulfate/water/overall (Å^2)	-	85/118/71/86	74/70/0/74
PDB accession code		**7NFU**	**7NG0**

Values in parentheses are for the outer resolution shell.

[a] $R_{merge} = \sum_{hkl} \sum_i |I_i(hkl) - \langle I(hkl)\rangle|/ \sum_{hkl} \sum_i I_i(hkl)$.

[b] $R_{meas} = \sum_{hkl} [N/(N-1)]^{1/2} \times \sum_i |I_i(hkl) - \langle I(hkl)\rangle|/ \sum_{hkl} \sum_i I_i(hkl)$, where $I_i(hkl)$ is the *i*th observation of reflection *hkl*, $\langle I(hkl)\rangle$ is the weighted average intensity for all observations *i* of reflection *hkl* and *N* is the number of observations of reflection *hkl*.

[c] $CC_{½}$ is the correlation coefficient between symmetry equivalent intensities from random halves of the dataset.

[d] The dataset was split into "working" and "free" sets consisting of 95 and 5% of the data respectively. The free set was not used for refinement.

[e] The R-factors R_{work} and R_{free} are calculated as follows: $R = \sum(|F_{obs} - F_{calc}|)/\sum|F_{obs}|$, where F_{obs} and F_{calc} are the observed and calculated structure factor amplitudes, respectively.

[f] As calculated using MolProbity[59]

FIGURE 1

Figure 1. Noc binds and hydrolyzes CTP in the presence of *NBS* DNA. (A) The domain architecture of *B. subtilis* Noc: a membrane-targeting sequence (MTS) that contains an amphipathic helix, an N-terminal domain (NTD), a central DNA-binding domain (DBD), and a C-terminal domain (CTD). **(B-D)** CTP binding as monitored by DRaCALA assay using radiolabeled CTP α-P^{32}. The bulls-eye staining indicates CTP binding due to a more rapid immobilization of protein-ligand complexes compared to free ligands. The starting concentration of Noc used in all panels was 30 μM. The concentrations of CTP α-P^{32}, unlabeled CTP, and a 22-bp *parS*/*NBS* DNA used in all panels were 5 nM, 30 μM, and 1 μM, respectively. **(E)** CTP hydrolysis rate of Noc (WT) and variants, measured by a continuous detection of released inorganic phosphates (see Materials and Methods). CTPase rates were measured at increasing concentration of CTP. All reactions contained 1 μM Noc (WT/variants) ± 1 μM 22-bp *NBS* or *parS* DNA and an increasing concentration of CTP (0, 10, 50, 100, 500, and 1000 μM). Experiments were triplicated and standard deviation of the CTPase rates were presented.

FIGURE 2

Figure 2. CTP and *NBS* DNA promote the self-engagement of the N-terminal domain of Noc.
(A-B) SDS-PAGE analysis of BMOE crosslinking products of 10 µM *B. subtilis* Noc (E29C) ± 1 µM 22-bp *parS*/*NBS* DNA ± 1.0 mM NTP. All crosslinking reactions were performed at 22°C unless indicated otherwise. X indicates a crosslinked form of Noc (E29C). Quantification of the crosslinked fraction is shown below each representative image. Error bars represent SEM from three replicates. **(C)** Same as panel A, but Noc (E29C R89A) was used instead. **(D)** Same as panel A, but Noc (E29C N121S) was used instead. **(E)** SDS-PAGE analysis of BMOE crosslinking of 10 µM dual-cysteine Noc (E29C S253C) variant ± DNA ± 1.0 mM NTP. Different DNAs were employed in crosslinking reactions: a linear 22-bp *NBS* DNA (22bp *NBS* lin, 1.0 µM), a circular 3-kb *NBS* plasmid (3kb *NBS* cir, 100 nM), or a circular 3-kb scrambled *NBS* plasmid (3kb *nonS* cir, 100 nM). The high-molecular weight (HMW) smear near the well of the polyacrylamide gel is marked with a solid box and an asterisk (see also panel F and Fig. S2G). When the crosslinking reaction was subsequently treated with a non-specific DNA nuclease, Benzonase, the HMW smear was no longer observed (dashed box, lane 8). X indicates a singly crosslinked form of Noc (E29C S253C). XX indicates a doubly crosslinked form of Noc (E29C S253C). **(F)** Same as panel E but fivefold more volume of each sample was loaded in each well to enhance the visualization of the HMW smear, only the top of the polyacrylamide gel was shown. The gel was also stained with a DNA-dye, Sybr Green (SYBR), and only the top section of the gel was shown. A smaller 22-bp *NBS* DNA migrated out of the gel, thus was not observed. **(G)** Agarose gel analysis of BMOE crosslinking products. A subset of crosslinking reactions (lanes 6,7,9, and 10) were loaded and resolved on a 1% agarose gel. The gel was subsequently stained with Sybr Green for DNA.

FIGURE 3

Figure 3. CTP facilitates the association of Noc with a closed *NBS* DNA substrate beyond nucleation. (A) Bio-layer interferometry (BLI) analysis of the interaction between a premix of 1.0 μM *B. subtilis* Noc ± 1.0 mM NTP and a 170-bp dual biotin-labeled DNA that contains either an *NBS* or a non-cognate *parS* site. Interactions between a dual biotinylated DNA and a streptavidin (SA)-coated probe created a closed DNA molecule where both ends were blocked (see also the schematic diagram of the BLI probes). Other Noc variants, Noc (R89A) and Noc (N121S), were also analyzed in the same assay. **(B)** BLI analysis of the interaction between a premix of 1.0 μM *B. subtilis* Noc ± 1.0 mM CTP and a BamHI-restricted dual biotinylated *NBS* DNA. The 170-bp *NBS* DNA substrate was also designed with a BamHI recognition site (see the schematic diagram of the probes). The intact dual biotinylated DNA was first immobilized onto the probe surface, then an open end was generated by digestion with BamHI (See Materials and Methods). Afterwards, the probe was used in a BLI analysis with a premix of Noc ± CTP. Each experiment was triplicated and a representative sensorgram was shown.

FIGURE 4

Figure 4. Noc binds liposomes in the presence of CTP and *NBS* DNA, and the phenotypic effects of the Noc variants. (A) Analysis of *B. subtilis* Noc binding to membranes by a liposome co-sedimentation assay. A premix of 0.75 µM *B. subtilis* Noc protein ± 1.0 µM 22-bp linear *parS*/*NBS* DNA ± 1.0 mM CTP ± 1.0 mg/mL liposomes was incubated at 22°C for 5 min before ultracentrifugation. The resulting supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE. Samples were also loaded onto a 20% TBE PAGE, and the gel was subsequently stained with Sybr Green for DNA. Quantification of Noc in each fraction is shown below each representative image. Error bars represent SEM from three replicates. **(B)** Same as panel A but 1.0 mM CTPɣS was used instead. **(C)** Other Noc variants, Noc (R89A), Noc (N121S), and NocNΔ10, were also analyzed in a liposome co-sedimentation assay. **(D)** Complementation of *noc* in a *ΔnocΔminCD* background. Strains *ΔnocΔminCD amyE*::P*xyl*-*noc* (WT or mutant)-*myfp* were streaked on nutrient agar plates supplemented with 0.5% xylose and incubated at 30°C or 39°C. **(E)** Cellular localization of mYFP-labeled Noc (WT or mutants). Representative images of *Δnoc amyE*::P*xyl*-*noc* (WT or mutant)-*myfp* cells grown in the presence of 0.5% xylose. Cell membranes were stained with FM5-95. Scale bar, 3 µm. Inset shows magnification of a section of cells.

FIGURE 5

Figure 5. Noc recruits *NBS* plasmid to liposomes in the presence of CTP. (A) Analysis of *B. subtilis* Noc binding to membranes and the recruitment of plasmid DNA to the membranes by a liposome co-sedimentation assay. A premix of 0.75 µM *B. subtilis* Noc protein ± 100 nM 5-kb plasmid DNA ± 1.0 mM CTP ± 1.0 mg/mL liposomes was incubated at 22°C for 5 min before ultracentrifugation. Either an empty plasmid or an *NBS*-harboring plasmid was employed in this assay. The resulting supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE. Samples were also loaded onto a 1% agarose gel, and was subsequently stained with Sybr Green for DNA. Quantification of Noc or DNA in each fraction is shown below each representative image. Error bars represent SEM from three replicates. **(B)** Similar to panel A, a premix of 100 nM *NBS* plasmid ± 0.75 µM Noc ± 1.0 mM CTP ± 1.0 mg/mL liposomes was first assembled and incubated at 22°C for 5 min. However, before ultracentrifugation, a non-specific DNA nuclease (Benzonase) was either added or omitted from the samples, as indicated. **(C)** Other Noc variants, Noc (R89A), Noc (N121S), and NocNΔ10, were also analyzed in a liposome co-sedimentation assay. Benzonase was either added or omitted, as indicated, before ultracentrifugation. **(D)** The association of Noc-*NBS* DNA with liposomes is reversible. A premix of 0.75 µM *B. subtilis* Noc protein + 100 nM *NBS* plasmid + 1.0 mM CTP + 1.0 mg/mL liposomes was ultracentrifuges, and the resulting fractions were analyzed for protein and DNA contents (lanes 1-2 and 3-4). The resulting pellet (lanes 2 and 4) were subsequently resuspended in either a binding buffer (- EDTA + 1 mM MgCl$_2$) or in a stripping buffer (+ 10 mM EDTA - MgCl$_2$). The resuspensions were ultracentrifuged for the second time, and the resulting fractions were analyzed for protein and DNA contents (lanes 5-6 and 7-8).

FIGURE 6

Figure 6. Crystal structure of *Geobacillus thermoleovorans* NocΔCTD shows the membrane-targeting amphipathic helix in an autoinhibitory conformation. (A) Crystal structure of a *G. thermoleovorans* NocΔCTD monomer (grey) with an N-terminal amphipathic helix α1 (magenta) and helix α2 (pink). Helix α2 is connected to the main N-terminal domain (NTD) via an eight amino-acid loop. The dashed line demarcates the NTD from the DNA-binding domain (DBD). Helix α8 at the DBD is the recognition helix that contributes to the specific recognition of the *NBS* site[17,24]. **(B)** The membrane-targeting amphipathic helix α1. A helical wheel representation of the 10-amino acid at the N-terminus of *G. thermoleovorans* Noc. While the first five amino acids were unresolved in the NocΔCTD crystal structure, the next five amino acids adopt a 3_{10} helical conformation with distinct polar and hydrophobic sides. **(C-D)** Helices α1 (magenta) and α2 (pink) pack themselves into the core N-terminal domain (grey). Hydrogen bonds are shown as dashed lines, water molecules are also shown.

FIGURE 7

Figure 7. The conformation of Noc in the crystal structure of *G. thermoleovorans* NocNΔ26ΔCTD is incompatible with an autoinhibitory state of the amphipathic helix. (A) Crystal structure of *G. thermoleovorans* NocΔCTD (same as Fig. 6A) with helices α5 and α6 highlighted in blue. The amphipathic helix α1 and α2 are shown in magenta and pink, respectively. **(B)** Crystal structure of a *G. thermoleovorans* NocNΔ26ΔCTD variant that lacks both the C-terminal domain (CTD) and helices α1 and α2 (dashed boxes). Helices α5 and α6 are shown in blue. **(C)** A dimer of *G. thermoleovorans* NocNΔ26ΔCTD that self-dimerizes at the N-terminal domain (NTD). Helices α5 and α6 in one of the subunits are shown in blue. **(D)** A superimposition at the NTDs of NocΔCTD monomer (grey) and NocNΔ26ΔCTD dimer (green and light blue) shows a severe clash (arrows) between α1 (magenta), α2 (pink) and the opposite subunit of NocNΔ26ΔCTD (light blue).

FIGURE 8

Figure 8. A model for a CTP-dependent regulation of membrane-binding activity of Noc. (A) Noc binds specifically to *NBS* site to nucleate on DNA. In the apo- or *NBS*-associated form, the amphipathic helix (magenta) adopts an autoinhibitory conformation, thus cannot bind to the membrane. *NBS*-binding stimulates Noc to bind CTP (orange). Concomitantly, CTP induces a sliding clamp conformation in Noc that can run off open ends of a linear 22-bp *NBS* DNA. In this state, the amphipathic helix is likely liberated from the autoinhibitory conformation, thus enables Noc-CTP to bind to the membrane. **(B)** When a circular *NBS* plasmid with no open end was employed, a sliding clamp of Noc entraps plasmid DNA and recruit DNA to the membrane. In the presence of CTD, a tripartite membrane-protein-DNA linkage is formed. CTP hydrolysis is not required for membrane binding or DNA recruitment but might have a role in releasing Noc from DNA and the membrane.

FIGURE S1

A

```
B. subtilis Noc   24  DTNKEEILEIPVNAIVPNRFQPRTIFSDEKIKELAMTIHTHGIIQPIVVRHTEEEGQYEL  83
                      D ++E + EI +  + PN +QPR  F DE + EL  ++  HGI+QP++VR + +   Y++
B. subtilis ParB  17  DLSEETVEEIKIADLRPNPYQPRKHFDDEALAELKESVLQHGILQPLIVRKSLK--GYDI  74

                           89                                     121
B. subtilis Noc   84  IAGERRWRAVQSLEWEKIPAIIKDFSDTETASVALIENLQREELSSIEEAHAYARLLELH  143
                      +AGERR+RA +      + +PAI+++ S+      +AL+ENLQRE+LS +EEA AY  LL+
B. subtilis ParB  75  VAGERRFRAAKLAGLDTVPAIVRELSEALMREIALLENLQREDLSPLEEAQAYDSLLKHL  134
                      arigine-rich patch

B. subtilis Noc  144  DLTQEALAQRLGKGQSTIANKLRLLKLPQPVQEAIMEKKITERHARALIPLKQPELQVTL  203
                      DLTQE LA+RLGK +   IAN LRLL LP+ +Q+ I E  ++  H R L+ LK       L
B. subtilis ParB 135  DLTQEQLAKRLGKSRPHIANHLRLLTLPENIQQLIAEGTLSMGHGRTLLGLKNKNKLEPL  194

B. subtilis Noc  204  LTEIIEKSLNVKQTEDRVVKMLEQ  227
                      + ++I + LNV+Q E ++++ L Q            Identities: 43%
B. subtilis ParB 195  VQKVIAEQLNVRQLE-QLIQQLNQ  217
```

B

Figure S1. Interactions between Noc (WT and mutants) with a non-hydrolyzable analog of CTP, CTPγS. **(A)** A sequence alignment between *B. subtilis* Noc and its paralog ParB. The conserved arginine-rich patch that mediates CTP binding in *B. subtilis* ParB is shown in a solid box. Residues R89 (red) and N121 (red) in *B. subtilis* Noc, whose equivalent substitutions in *B. subtilis* ParB have been showed to impair spreading and CTP binding[2], were substituted by alanine and serine, respectively. **(B)** Analysis of the interaction of *B. subtilis* Noc (WT and mutants) with CTPγS by isothermal calorimetry (ITC). Each experiment was duplicated. Regression curves were fitted, and binding affinities (K_d) are shown.

FIGURE S2

Figure S2. CTP and CTPɣS, but not other nucleotides, promote the engagement of the N-terminal domain of Noc. (A) SDS-PAGE analysis of BMOE crosslinking products of 10 µM *B. subtilis* Noc (WT) protein ± 1.0 µM 22-bp *parS*/*NBS* DNA ± 1.0 mM NTP. Wild-type Noc naturally lacks cysteine, hence does not crosslink in the presence of BMOE. All crosslinking reactions were performed at 22°C unless indicated otherwise. **(B)** Same as panel A but Noc (E29C) was used instead. X indicates a crosslinked form of Noc (E29C). Quantification of the crosslinked fraction is shown below each representative image. Error bars represent SEM from three replicates. **(C)** Same as panel A, but NocNΔ10 (E29C) was used instead. **(D)** Time-course of Noc (E29C) crosslinking with CTP in the presence or absence of 22-bp *NBS* DNA. Purified Noc (E29C) was preincubated with 1.0 mM CTP ± 1 µM 22-bp *NBS* DNA for 1, 5, 10, 15, or 30 min at 4°C (instead of the usual 22°C) before BMOE was added. A lower incubation temperature was needed to slow down the reaction. Quantification of the crosslinked fraction is shown below each representative image. Error bars represent SE from three replicates. **(E)** Same as panel D, but 1.0 mM CTPɣS was used instead. **(F)** Same as panel A, but Noc (S253C) was used instead. **(G)** Magnification of the Coomassie-stained top gel shown in Fig. 2F.

FIGURE S3

Figure S3. CTP and *NBS* DNA enables Noc binding to liposomes. (A) Analysis of *B. subtilis* Noc binding to membranes by a liposome co-sedimentation assay. A premix of 1.0 µM 22-bp linear *NBS* DNA ± 0.75 µM *B. subtilis* Noc protein ± 1.0 mM CTP ± 1.0 mg/mL liposomes was incubated at 22ºC for 5 min before ultracentrifugation. The resulting supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE. Without Noc, 22-bp *NBS* DNA did not co-sediment on its own (lanes 1-2) or with liposomes (lanes 3-4). Samples were also loaded onto a 20% TBE PAGE, and gel was subsequently stained with Sybr Green for DNA. Quantification of Noc in each fraction is shown below each representative image. Error bars represent SEM from three replicates. **(B)** CTP but no other nucleotide enables Noc to co-sediment with liposomes. A premix of 0.75 µM *B. subtilis* Noc protein + 1.0 µM 22-bp *NBS* DNA ± 1.0 mM NTP + 1.0 mg/mL liposomes was incubated at 22ºC for 5 min before ultracentrifugation. The resulting supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE. **(C)** *Caulobacter crescentus* ParB does not co-sediment with liposomes in any tested conditions. A premix of 0.75 µM *C. crescentus* ParB protein ± 1.0 µM 22-bp linear *parS/NBS* DNA ± 1.0 mM CTP ± 1.0 mg/mL liposomes was incubated at 22ºC for 5 min before ultracentrifugation. The resulting supernatant (S) and pellet (P) fractions were analyzed by SDS-PAGE.

FIGURE S4

Figure S4. Liposome flotation assays shows Noc can recruit *NBS* plasmid to membrane in the presence of CTP. (A) The principle of a liposome flotation assay. Liposomes ± purified Noc ± CTP ± *NBS* plasmid were incubated in a 30% sucrose binding buffer. Buffer only with 25% sucrose and 0% sucrose were subsequently layered on top sequentially. After ultracentrifugation, liposomes and associated protein/DNA migrate along the sucrose gradient i.e. floating to the upper most fractions. Three equal fractions (bottom, middle, and top) were drawn out sequentially using a Hamilton syringe, and their protein and DNA contents were analyzed. **(B)** Control experiments: liposome flotation assays in which one component, either Noc protein, a 5-kb plasmid DNA, CTP, or liposomes, was omitted. **(C)** Analysis of *B. subtilis* Noc binding to membranes and the recruitment of plasmid DNA to membranes by a liposome flotation assay. A premix of 0.75 µM *B. subtilis* Noc ± 100 nM 5-kb plasmid DNA ± 1.0 mM CTP ± 1.0 mg/mL liposomes was incubated at 22°C for 5 min before ultracentrifugation. Either an empty plasmid or an *NBS*-harboring plasmid were employed in this assay. The resulting fractions (Bottom B, Middle M, and Top T) were analyzed by SDS-PAGE. Samples were also loaded onto a 1% agarose gel and was subsequently stained with Sybr Green for DNA. Quantification of Noc or DNA in each fraction is shown below each representative image. Error bars represent SEM from three replicates. **(D)** Other Noc variants, Noc (R89A), Noc (N121S), and NocNΔ10, were also analyzed in a liposome flotation assay. A premix of 0.75 µM *B. subtilis* Noc protein (WT or mutants) + 100 nM *NBS* plasmid + 1.0 mM CTP + 1.0 mg/mL liposomes was ultracentrifuges, and the resulting fractions were analyzed for protein and DNA contents.

FIGURE S5

A

```
                                    α1            α2        α2-β1 linker
G. thermoleovorans Noc   1   MKHPFSRLFSFGEKEQEE--MEEKQEREEVRHIPVKSIIPNRFQPRTMFDEEKIDELALT    58
                             MKH FSR F  GEKEQE    E   +EE+   IPV +I+PNRFQPRT+F +EKI ELA+T
        B. subtilis Noc   1  MKHSFSRFFGLGEKEQEPEIAEHDTNKEEILEIPVNAIVPNRFQPRTIFSDEKIKELAMT    60
                                                                  arigine-rich patch
G. thermoleovorans Noc  59   IRTHGIIQPIVVRECGN-GRFEIIAGERRWRAVQKLGWTEIPAIIKNLNDKETASVALIE   117
                             I THGIIQPIVVR     G++E+IAGERRWRAVQ L W +IPAIIK+ +D ETASVALIE
        B. subtilis Noc  61  IHTHGIIQPIVVRHTEEEGQYELIAGERRWRAVQSLEWEKIPAIIKDFSDTETASVALIE   120

G. thermoleovorans Noc 118   NLQREELTPIEEAMAYAKLIELHDLTQEALAQRLGKGQSTIANKLRLLKLPQEVQEALLQ   177
                             NLQREEL+ IEEA AYA+L+ELHDLTQEALAQRLGKGQSTIANKLRLLKLPQ VQEA+++
        B. subtilis Noc 121  NLQREELSSIEEAHAYARLLELHDLTQEALAQRLGKGQSTIANKLRLLKLPQPVQEAIME   180

G. thermoleovorans Noc 178   RAITERHARALIALKDKEKQLKLLQEIIDKQLNVKQTEDRVLKLLEAGERKPKPKRKAFS   237
                             + ITERHARALI LK  E Q+ LL EII+K LNVKQTEDRV+K+LE G+RKPKP+RKAFS
        B. subtilis Noc 181  KKITERHARALIPLKQPELQVTLLTEIIEKSLNVKQTEDRVVKMLEQGQRKPKPRRKAFS   240

G. thermoleovorans Noc 238   RDMRIAVNTIRQSLSMVESSGVSVQSEEEEFDDYYQITIRIAK   280      ┌──────────────────┐
                             RD RIA+NTIRQSLSMVE SGV + +EEEF++Y Q+TIRI K              │ Identities: 72%  │
        B. subtilis Noc 241  RDTRIAMNTIRQSLSMVEDSGVKLNTEEEEFEEYIQLTIRIPK   283      └──────────────────┘
```

B

RMSD = 0.96 Å

— NocΔCTD chain A
— NocΔCTD chain B

Figure S5. The composition of the asymmetric unit of *G. thermoleovorans* NocΔCTD crystal structure. (A) A sequence alignment between *B. subtilis* Noc and its homolog *G. thermoleovorans* Noc. The conserved arginine-rich patch that mediates CTP binding in *B. subtilis* ParB is shown in the solid box. The sequences of amphipathic helix α1, helix α2, and the α2-β1 connecting loop are also indicated on the sequence alignment. **(B)** The asymmetric unit of *G. thermoleovorans* NocΔCTD contains two copies of NocΔCTD monomers (left panel). Chains A and B are structurally similar, RMSD = 0.96 Å (right panel).

FIGURE S6

Figure S6. The conformation of *G. thermoleovorans* NocNΔ26ΔCTD is similar to that of a nucleotide-bound *B. subtilis* ParBΔCTD. (A) Superimposition between a *G. thermoleovorans* NocNΔ26ΔCTD dimer (green) and a *B. subtilis* ParBΔCTD-CDP dimer (yellow). CDP molecules are shown in magenta. **(B)** Magnification of the nucleotide-binding pocket of *B. subtilis* ParBΔCTD and *G. thermoleovorans* NocNΔ26ΔCTD. CDP and Ca²⁺ cations that belong to *B. subtilis* ParBΔCTD-CDP co-crystal structure are highlighted in magenta and yellow, respectively. In the *G. thermoleovorans* NocNΔ26ΔCTD structure, a sulfate ion (dark green) occupies a similar position to the β-phosphate group of CDP. **(C)** A superimposition at the N-terminal domains of a NocΔCTD monomer (grey) and a NocNΔ26ΔCTD monomer (green). The amphipathic helix α1 and helix α2 are shown in magenta and pink, respectively. **(D)** Same as panel B, but only the N-terminal domain (NTD) is shown for clarity.