1    Moving together, foraging apart: effects of the social environment on movement-integrated

2                                      habitat selection

3    Quinn M.R. Webber[1,*], Christina M. Prokopenko[2], Katrien A. Kingdon[2], & Eric Vander Wal[1,2]

4        [1]Cognitive and Behavioural Ecology Interdisciplinary Program, Memorial University of

5                              Newfoundland, St. John's, NL

6        [2]Department of Biology, Memorial University of Newfoundland, St. John's, NL

7                    * Corresponding author: webber.quinn@gmail.com

# Abstract

Movement links the distribution of habitats with the social environment of animals using those habitats; yet integrating movement, habitat selection, and socioecology remains an opportunity. Here, our objective was to disentangle the roles of habitat selection and social association as drivers of collective movement in a gregarious ungulate. To accomplish this objective, we (1) assessed whether socially familiar individuals form discrete social communities and whether social communities have high spatial, but not necessarily temporal, overlap and (2) modelled the relationship between collective movement and selection of foraging habitats using socially informed integrated step selection analysis. Based on assignment of individuals to social communities and home range overlap analyses, individuals assorted into discrete social communities and these communities had high spatial overlap. By unifying social network analysis with movement ecology, we identified state-dependent social association, where individuals were less cohesive when foraging, but were cohesive when moving collectively between foraging patches. Our study demonstrates that social behaviour and space use are inter-related based on spatial overlap of social communities and state-dependent habitat selection. Movement, habitat selection, and social behaviour are linked in theory. Here, we put these concepts into practice to demonstrate that movement is the glue connecting individual habitat selection to the social environment.

**Keywords:** caribou, integrated step selection analysis, movement ecology, social preference, social network analysis

# 1. Introduction

Movement is defined by a change in spatial location and is the behavioural link between the physical space an animal occupies and the resources available to them [1]. In the context of the social environment, movement represents the connection between the distribution of foraging resources and the social structure of animals that consume those resources [2]. Disentangling the social and spatial drivers of movement is a formidable challenge within behavioural ecology. In many cases, research omits the social contexts within which animals move to, from, and within the areas that contain foraging resources [but see 3,4]. Spatially-explicit models of sociality highlight that some gregarious species aggregate at areas associated with profitable foraging resources [5], while territorial species typically only interact at territory edges [6]. Sharing space, either at foraging sites, territory edges, or elsewhere within an animal's range is required to form the social environment. For example, animals are predicted to select habitat as a function of the profitability and availability of the habitat [7]. A logical extension can be made to conspecifics; individuals form groups based on their familiarity with conspecifics. We aim to quantify the relative importance of habitat and conspecifics by developing a novel socially informed integrated step selection analysis, a movement-based method that accounts for the relative intensity of selection for habitats and neighbours.

For social animals, individual movement shapes social encounters and subsequent interactions with conspecifics that can affect collective movement [8]. Further complicating our understanding of collective movement is the idea that the type, quality, and distribution of habitats on the landscape can constrain or promote collective movement [9]. For example, dense vegetation impedes visibility, which could reduce the probability a group remains together. In addition, individual movement and habitat selection are affected by the distribution of resources.

For example, patchily distributed foraging resources could facilitate large aggregations, whereas

homogenously distributed foraging resources could result in a reduction in social associations

[10]. The physical space an individual, or group, occupies and the distribution and availability of

foraging resources within that space are important drivers of animal movement and the social

environment an individual experiences [2].

Animals typically select for habitats that maximize foraging and minimize risk of

predation; an important trade-off because most habitats do not accommodate both high quality

foraging *and* low predation risk. When animals aggregate in large groups, the per capita risk of

predation is lower and individuals in larger groups tend to select more risky habitats, including

foraging in open areas [11]. For example, animals in larger groups reduce time spent vigilant

[12]. However, not all social groups are equal; some groups contain unfamiliar individuals (i.e.,

anonymous groups) [13], while others contain familiar individuals [14]. For anonymous and

familiar groups, social foraging occurs when the costs and benefits of an individual's foraging

behaviour are linked with the foraging behaviour of conspecifics [15]. Social foraging can be

most beneficial when social information about foraging resources comes from familiar

individuals [16]. For example, when foraging resources are unpredictable, familiar individuals

obtain reliable information from conspecifics to increase foraging efficiency [17,18], such that

time searching for forage is reduced in favour of more time spent foraging. In the context of

movement and habitat selection, theory on social foraging and the benefits of social familiarity

provides a framework through which the costs and benefits of collective movement can be

explored [15,19].

Apparent social familiarity or preference is the long-term repeated social association due

to shared space at the same time. Although individuals often interact with many conspecifics,

non-random repeated social interactions or associations with certain individuals form the basis

for social preference [20]. Proximately, long-term social relationships can influence collective

movement via the reliability of information transfer about foraging resources or predator risk

[21,22], while ultimately they can enhance fitness [23]. The social environment can be

influenced by the availability of foraging resources, but social communities can also be

composed of individuals with similar physiological or nutritional requirements that occupy the

same locations. Apparent social preference may therefore arise as a function of spatial

constraints [24], including physical barriers, such as rivers or mountains. Disentangling social

preference from spatial constraint could inform our understanding of collective movement and

habitat selection [25,26].

Here, we develop a unified framework to bridge the gap between social network analysis

and movement ecology. We disentangle the roles of social preference and collective movement

on habitat selection behaviour by parameterizing socially informed integrated step selection

models (Figure 1). Animal social networks often comprise distinct sub-networks, or social

communities, defined by the existence of social preference among discrete clusters of individuals

[20]. Using a social ungulate as a model system, our objective was to disentangle the roles of

habitat selection and social association as drivers of collective movement in a gregarious

ungulate (*Rangifer tarandus*) when the availability and distribution of foraging resources are

variable. We calculated three distinct measures of social preference. First, we assigned

individuals to social communities based on a community detection algorithm. Second, we

assessed the temporal stability of social association among individuals. Third, we estimated

spatial overlap of social communities using home range analyses. Due to variance in the

distribution of foraging resources on the landscape, we expected that access to social information

via close proximity to conspecifics should influence patterns of selection for foraging resources.

Specifically, individuals with stronger social preference should select foraging habitat

collectively. The corollary is that individuals should also take short steps in the presence of

conspecifics, given that from a movement ecology perspective, shorter steps typically represent

foraging behaviour and longer steps represent searching behaviour [27].

## 2. Materials and Methods

*(a) Caribou as a model system*

We investigated patterns of movement, space use, and social behaviour for caribou

(*Rangifer tarandus*) on Fogo Island, Newfoundland, Canada. Fogo Island is a small (~237km$^2$)

island off the northeastern coast of Newfoundland with a humid continental climate (see

Supplementary Materials for details). Between 1964-1967, 26 caribou were introduced to Fogo

Island from the Island of Newfoundland [28]. Currently, Fogo Island has a population of

approximately 300 caribou (Newfoundland and Labrador Wildlife Division, unpublished data).

Caribou live in fission-fusion societies [29] and throughout much of their range caribou forage

primarily on lichen, grasses, sedges, and other deciduous browse with access to these resources

changing between the seasons [30]. During winter (January to March) the landscape is covered

by snow and caribou forage primarily on lichen. Lichen is heterogeneously distributed and

access is impeded by snow and ice cover. Caribou dig holes in the snow, termed craters, to

access lichen in the winter, often where snow depth is relatively shallow (~30–60 cm deep).

Consequently, caribou have limited access to subnivean forage and tend to re-use established

craters. To cope with this limitation, caribou use conspecific attraction and social information

transfer to gain access to foraging opportunities [31]. In winter, caribou activity budgets suggest

that caribou spend ~50% of their time foraging, while ~40% of their time is spent lying down or

ruminating, 7% of their time is spent walking or trotting, and 3% of their time is spent standing

[32,33]. In addition, caribou typically avoid forested habitats due to deep snow in forests and

lack of access to forage opportunities [34], whereas most open habitats on Fogo Island are

windswept in the winter, therefore facilitating foraging and movement [30].

We used GPS location data collected from Fogo Island caribou (2017–2019) to assess the

relationship between social behaviour, habitat selection, and movement (see supplementary

information for details on collaring procedures). Each relocation was assigned to a given habitat

classification that was extracted from Landsat images with 30x30m pixels [35]. Locations were

categorized as one of open foraging (lichen barrens), open moving (wetland, rocky outcrops, and

water/ice), or closed (conifer scrub, mixed wood, and conifer forest). Adult female caribou (n =

26 individual caribou, n = 72 caribou-years) were immobilized and fitted with global positioning

system (GPS) collars (Lotek Wireless Inc., Newmarket, ON, Canada, GPS4400M collars, 1,250

g). Prior to analyses, we removed all erroneous and outlier GPS locations following Bjørneraas

et al. [36]. We did not collar all female caribou in the herds, however, and collared individuals

were randomly selected from the population. We therefore assume that our sample of collared

animals was randomly distributed. Although associations between collared and uncollared

animals were unrecorded, we assumed that our networks (see below) were unbiased

representations of the relative degree of social association among all caribou.

*(b) Formulating integrated step selection models*

Integrated step selection analysis (iSSA) simultaneously incorporates movement and

habitat selection within a conditional logistic regression framework [see Figure 1; 33]. As in

other resource and step selection analyses [38], iSSA models habitat use as a binomial response

variable where 'use' represents the location an animal was observed and 'availability' represents

the geographical area an animal could potentially use but was not necessarily observed (Figure

S1). iSSA defines availability based on empirically fitted distributions of step lengths and turn

angles [37], where a step is the linear connection between consecutive relocations, and turn angle

is the angular deviation between the headings of two consecutive steps [39]. We generated

available steps and turn angles based on the distributions informed by observed population-level

movement behaviour using the *amt* package in R [40]. First, we sampled step lengths from a

gamma distribution of observed step lengths for the study population; values were log-

transformed for analysis. The statistical coefficient of log-transformed step length is a modifier

of the shape parameter from the gamma distribution originally used to generate available steps

[37]. Second, we sampled turn angles (measured in radians) for available steps from observed

values between $-\pi$ and $\pi$ following a Von Mises distribution; values were cosine transformed

for analysis. The statistical coefficient of cosine transformed turn angle is an unbiased estimator

of the concentration parameter of the Von Mises distribution [37,41]. Each observed relocation

was paired through a shared start point with 10 available steps generated from step-length and

turn-angle distributions and compared in a conditional logistic regression framework (see section

*g*). In addition to generating available movement parameters, we also generated an available

social environment (see below).

*(c) Social network analysis*

We used the R [42] package *spatsoc* [43] to generate proximity-based social association

networks from GPS location data. Nodes in the networks were represented by individual caribou

and edges were represented by the frequency of association based on proximity between

individuals. We generated social networks at two scales based on proximity of locations between

individual caribou: (1) seasonal winter networks to assign individuals to social communities and

assess long-term social preference and (2) weekly networks to assess the role of short-term social

preference on patterns of habitat selection (section *b*). Social communities represent a subset of

individuals within a network that are more closely connected with each other than with the rest

of the network. For networks at both seasonal and weekly scales, we assumed association

between two individuals when simultaneous locations (i.e. GPS relocations that occurred within

5 minutes of each other) were within 50 m of one another [29,31]. We selected the 50 m

threshold based on the standard distance applied to assign individuals to groups in studies of

ungulate group size and social behaviour [44]. We applied the 'chain rule', where each discrete

GPS fix was buffered by 50 m and we considered individuals in the same group if 50 m buffers

for two or more individuals were contiguous, even if some individuals were beyond 50 m of one

another. We weighted edges of social networks by the strength of association between dyads of

caribou using the simple ratio index [45], SRI:

$$SRI = \frac{x}{x + y_{AB} + y_A + y_B}$$

where *x* is the number of times individuals A and B were within 50 m of each other, $y_A$ is the

number of fixes from individual A when individual B did not have a simultaneous fix (i.e. within

5 minutes), $y_B$ is the number of fixes from individual B when individual A did not have a

simultaneous fix, and $y_{AB}$ is the number of simultaneous fixes from individuals A and B that

were separated by >50 m [46]. We constructed social networks with the *igraph* package in R

[47]. Nodes in our networks therefore represented individuals and edges represented associations

between individuals.

*(d) Detecting social communities: long-term social preference*

For seasonal winter social networks, we used a community detection algorithm to define

social communities [48]. We assessed social community structure for each winter to determine

the broadest extent of social structure. Modularity is a commonly used measure that defines how

well-connected social communities are to one another. It is calculated from the weighted

proportion of edges that occur within a community, minus the expected proportion of edges, if

edges were distributed randomly in the network [48]. A modularity value close to 1 indicates a

network with a strong clustered structure in which interactions of individuals belonging to

different clusters do not occur. We quantified modularity ($Q$) for observed annual winter

networks. To ensure observed social structure did not occur at random, we compared these

values to null models [24]. Specifically, we generated null models based on GPS fixes to reduce

potential for type II error typically associated with node-based permutations [49]. Following

Spiegel et al. [24], we re-ordered daily GPS movement trajectories for each individual while

maintaining the temporal path sequence within each time block (e.g., day 1 and day 2 may be

swapped). This technique is a robust network randomization procedure for GPS data because: 1)

it maintains the spatial aspects of an individual's movement; 2) by randomizing movement

trajectories of individuals independent of one another, temporal dependencies of movement are

decoupled [24]. We repeated this procedure 100 times for annual winter networks and re-

calculated modularity at each iteration. We then compared observed modularity ($Q$) values to the

null distribution and determined whether the observed $Q$ value fell within the 95% confidence

interval of the distribution of $Q$ values [20].

In addition to comparing observed $Q$ values from annual winter networks to a null

distribution, we also calculated a community assortativity coefficient ($R_{com}$) to assess confidence

in the assignment of an individual to a given community [50]. Specifically, $R_{com} = 0$ indicates no

confidence in the assignment of an individual to a community, while $R_{com} = 1$ indicates certainty

in the assignment of an individual to its community.

*(e) Weekly networks and lagged association rates: short-term social preference*

We iteratively generated weekly social networks using a moving window approach and

calculated the observed SRI to be included as a covariate in iSSA models (see section *b*). The

first network was calculated for 1 January to 7 January, the second was 2 January to 8 January,

and so on. Weekly networks contained 84 relocations per individual (12 relocations per day). For

each of these networks we used dyadic values of SRI as a proxy for short-term social preference.

We used a three-step process. First, to incorporate SRI within the iSSA framework, we

determined the identity and distance (m) of each individual's nearest neighbour at each

relocation [43]. Second, for each focal individual and their nearest neighbour at each relocation,

we matched the dyadic SRI value for the prior week. For example, for individual A at 12:00 on 8

January, we determined the nearest neighbour was individual B and we extracted the dyadic SRI

value for these individuals for the previous week. Third, we repeated steps one and two for all

'available' relocations defined by random steps generated in the iSSA (section *b*). Therefore,

each individual at each relocation had an observed weekly dyadic SRI value and a series of

available weekly dyadic SRI values (see section *b*).

In addition to incorporating social preference directly within iSSA models, we also

assessed social preference by estimating within-season temporal patterns in associations between

individuals by calculating the lagged association rate (LAR). We calculated the LAR within each

seasonal proximity-based social network using the *asnipe* package in R [51]. We calculated

LARs for each winter by generating social networks following the procedure described above

(section *e*) using GPS relocations. LARs measure the probability that pairs of individuals

associating at a given relocation would still associate at subsequent relocations [52]. We

compared within-season LARs for individuals in the same annual winter social community to

LARs for individuals in different annual winter social communities to assess potential for within-

season patterns of association among individuals.

*(f) Home range overlap between social communities*

To determine spatial overlap of social communities we estimated home ranges for winter

social communities using the area of the 95% isopleths from fixed kernel density estimates [53]

for each social community in each year with the *href* smoothing parameter in the *adehabitatHR*

package in R. Data from all individuals in a given social community were pooled to estimate the

community home range. We estimated home range overlap between social communities with the

utilization distribution overlap index (UDOI), where higher values of UDOI represent a greater

proportion of overlap and lower values represent lower proportion of overlap [54].

*(g) Modelling collective movement and habitat selection*

We fitted three separate iSSA models using the *glmmTMB* package in R to fit conditional

logistic regressions [55]. We fit iSSA models with a series of fixed and random effects following

Muff et al. [55], through which we take advantage of the fact that the conditional logistic

regression model is a likelihood-equivalent to a Poisson model with stratum☐specific fixed

intercepts. Specifically, our first model was a 'core model' [39] which is an informed null model

that includes covariates expected to influence patterns of habitat selection regardless of collective

movement (Table S1). The core model contained three fixed effects: habitat type, the natural log-

transformed step length, cosine transformed turn angle, and interactions between habitat and log-

transformed step length and cosine-transformed turn angle. We also modelled individual-specific

slopes for log-transformed step length and habitat within the random effect structure of the

model. We included three distinct habitat types in our models: forest, open-foraging, and open-

movement. The forest category included coniferous forest, conifer scrub, broadleaf forest, and

mixed-wood forest habitat types; open-foraging included lichen habitat; and open-movement

included wetland, rocky barrens, frozen lakes, and anthropogenic habitat types. The core model

was an informed null model that we used as the foundation on to add hypothesis-testing

covariates.

Our second model tested covariates included in the core model and nearest neighbour

distance as a fixed effect. Nearest neighbour distance (m) was measured as the distance between

a focal individual and the nearest collared conspecific and was calculated for all used and

available steps in *spatsoc* [43]. Nearest neighbour distance was transformed using the natural

logarithm (ln) and included in interactions with cosine transformed turn angle, ln step length, and

habitat type within the fixed effect structure of the model (Table S1). In addition, we also

included individual-specific slopes for nearest neighbour distance, the interaction between log-

transformed step length and nearest neighbour distance, and the interaction between habitat type

and nearest neighbour distance within the random effect structure of the model.

Our third model also tested covariates included in the core model, nearest neighbour

distance, and weekly dyadic simple ratio index (section *c*). Specifically, we incorporated weekly

dyadic SRI values for each observed and available step. Dyadic SRI was included in interactions

with turn angle, step length, and habitat type (Table S1).

*(h) Calculating effect sizes*

We calculated population-level relative selection strength (RSS) to demonstrate how

habitat features influenced selection [56]. Specifically, we calculated the strength for selecting

one step over another that differ in the habitat value where those steps end. RSS was calculated

for each habitat type (i.e., forest, lichen, or open habitats) as a function of nearest neighbour

distance and the shared dyadic simple ratio index between nearest neighbours.

# 3. Results

We found that individuals associated with members of multiple communities and associations were stronger among members a given community. Depending on the year, social networks comprised 2–6 social communities and although community assortativity ($R_{com}$) was similar across years, there was high certainty (range = 0.95–1.00) of an individual's assignment to a given community in a given year (Table 1). In addition, lagged association rates (LAR) within each winter confirmed temporal stability of community assortment, where association rates for members of the same winter community remained higher than association rates for members of different communities in each year (Figure 2). Seasonal winter values of modularity ($Q$) were significantly lower than the distribution of $Q$ generated from null models (Figure S2), suggesting that although social networks were structured into communities, inter-community social associations are common (Table 1). In support of our expectation, we observed relatively high spatial overlap between different winter social communities (average UDOI = 0.37, SD = 0.34, range = 0–0.98; Figure S3; Table S2), thus facilitating the potential for association between social communities.

We found that models including nearest neighbour distance and the simple ratio index covariates were higher ranked compared to the core and nearest neighbour distance models (Table S3). We found mixed support for our second set of predictions. In contrast to our expectation, individuals avoided conspecifics while selecting foraging lichen habitat ($\beta = 0.021$, 95% CI: 0.0033, 0.038; Figure 3; Table S3). Although individuals tended to move away from conspecifics during foraging ($\beta = 0.239$, 95% CI: 0.007, 0.041), on average, individuals selected nearest neighbours (regardless of distance) that had higher dyadic values of the simple ratio index ($\beta = 2.91$, 95% CI: 2.77, 3.06; Figure 3; Table S3). These findings suggest that although

individuals selected foraging habitat further from conspecifics, they had stronger shared values

of the SRI for their nearest neighbours. We also found support for our expectation of collective

movement, although there was no effect of social proximity on turning angles. Specifically,

individuals took shorter steps when they were in close proximity to conspecifics ($\beta$ = –0.02, 95%

CI: –0.0254, –0.0148), even though individuals took longer steps when they shared a higher

dyadic SRI value with their nearest neighbours ($\beta$ = 0.151, 95% CI: 0.131, 0.172; Figure 3;

Table S3). These findings suggest individuals tend to move shorter distances in close proximity

to any conspecifics, but further distances when they had stronger shared values of the SRI for

their nearest neighbours.

## 4. Discussion

Our study examined apparent social preference in the context of shared space use using socially

informed integrated step selection analysis. We present a framework that unifies social networks

within a traditional movement ecology and habitat selection framework. Although individual

social associations were well mixed at the population level, we found that social networks were

structured into discrete communities. Despite spatial overlap between different social

communities, which suggests an opportunity for individuals to interact with members of other

communities, we highlight two forms of within-community social preference, including long-

term temporal stability of associations among individuals, as well as an effect of short-term

social preference on habitat selection. Further, we found that individuals tended to select

foraging habitat spaced away from conspecifics, but moved between foraging habitats with

conspecifics, suggesting collective movement is state-dependent. The processes underlying

community structure appear to be social, and not spatial. Based on our unification of social

network analysis with integrated step selection analyses, we highlight the influence of collective

movement and preferred associations on habitat selection and foraging.

Testing social preference as a driver of movement and habitat selection required

establishing the existence of discrete communities and long-term social associations within the

population-level network. Indeed, the formation of discrete social communities, in combination

with our lagged association analysis, confirmed the existence of temporal stability in social

associations for members of the same social community. The formation of non-random social

communities was driven in part by social preference, but aspects of space use, including shared

space, could also influence the stability of social communities [57]. We found high spatial

overlap between social communities, suggesting that physical barriers on the landscape do not

explain the formation of discrete social communities. For social communities to emerge from a

well-mixed population, individuals in different communities must have high spatial, but low

temporal overlap in shared geographical space, thus revealing the importance of space and time

in the formation of social communities [58]. Disentangling space and time within the social

environment reveals distinct social communities and groups of individuals that are more likely to

associate than simply by chance [24]. On resource limited landscapes, individuals are expected

to aggregate in close proximity to those resources, for example, elephants (*Loxodonta africana*)

aggregate near water-holes, which are a limiting resource [5]. At the population-level, social

networks were highly connected, thus providing the impetus to quantify socially informed

patterns in movement and habitat selection.

Our findings reveal state-dependent social associations, where selection for nearest

neighbours was reduced when foraging, but strongest when moving between foraging patches.

Individuals are more likely to trust social information from familiar individuals, but the potential

costs are an increase in competition at foraging patches. Individuals may balance the trade-off

between competition and access to information by sharing information about the general location

of foraging patches in transit but spacing apart at patches to reduce competition. Specifically,

when individuals were in closer proximity to one another they selected foraging resources, which

is typically open, allowing them to remain in visual and vocal contact, thereby facilitating social

cohesion during foraging despite physically spacing apart [59]. Individuals also took longer steps

when they had high social preference for their nearest neighbour, indicating increased movement

when nearest neighbours are familiar with each other. Under the assumption that shorter steps

represent foraging behaviour and longer steps represent searching behaviour [27], our results

suggest familiar individuals move between foraging patches together. Our results reveal potential

behavioural mechanisms (i.e., foraging or moving) that influence the frequency and magnitude

of social associations.

The emergent geometry of collective movement and spatial arrangement of individuals in

a group appears to change as individuals adjust their behaviour based on the availability of

resources and the presence of familiar conspecifics [60]. Assame macaques (*Macaca*

*assamensis*) distance from one another during foraging, but move collectively between foraging

sites [61], while individual giraffes (*Giraffa camelopardalis*) show social preference for

conspecifics during foraging, but not during movement [22]. Interestingly, macaques foraged in

closer proximity to individuals of similar dominance rank, while for giraffes it was unclear

whether observed social preference was the result of passive or active assortment. For caribou,

dominance hierarchies are linear and typically driven by body size [62,63], suggesting that social

preference in caribou could also be related to dominance. Our ability to delineate aspects of the

social environment between collective movement and habitat selection within a unified

framework is useful for disentangling passive or active assortment, for example dominance rank,

conspecific attraction, or the transfer of information about foraging resources.

We assumed that dispersed foraging is a function of competition among individual

caribou for craters in the winter [62] and movement in groups could reflect the use of social

information about foraging sites [29] or predation [64]. Craters can vary in size and distribution

[30]; however, craters may only be large enough for a single individual to forage at a time [65].

Solitary foraging is therefore most beneficial but foraging in close proximity to preferred

conspecifics may be an exception. We propose that while caribou generally have larger group

sizes in winter [31,66] groups may be sparsely distributed in space to reduce fine-scale

competition at individual cratering sites. Groups may spread out such that competition at

foraging sites is limited, but group members can retain visual contact. Furthermore, female

caribou have antlers, which unlike males, persist into winter. Females are hypothesized to use

their antlers to defend craters and exert dominance over both males and females without antlers

[62,67]. This interpretation is corroborated by theory used to explain fission-fusion dynamics,

where individuals are expected to split and merge through space and time to reduce conflict and

competition during foraging.

We demonstrate assortment of individuals into distinct social communities, despite high

range overlap with individuals in other communities. Integrating space and time revealed fine-

scale processes that form social communities and the socially mediated nature of movement

ecology and habitat selection. Within a unified socially informed integrated step selection

framework, we bridge the theoretical and methodological gap between social network analysis

[68], movement ecology [69], and habitat selection [70]. We also demonstrate how social

association is context-dependent, where individuals spaced out during foraging, but moved

collectively between foraging patches. Our synthesis of integrated step selection analysis with

social networks to test hypotheses is an important step towards identifying the roles of physical

space and animal space use as factors influencing the social environment [9]. Moreover,

individual variation in phenotypes attributable to movement or habitat selection may affect how

individuals experience the social environment [71]. Movement, habitat selection, and social

behaviour are clearly linked; as van Moorter et al. [1] described movement as the 'glue'

connecting habitat selection to the physical location of a given set of habitats, we posit that

movement is the glue connecting collective habitat selection to the social environment.

**Ethics.** All animal captures and handling procedures were consistent with the American Society

of Mammologist guidelines and were approved by Memorial University Animal Use Protocol

No. 20152067.

**Data accessibility.** All code and data used for statistical analysis and figures will become

available upon acceptance.

**Authors contributions.** QMRW and EVW conceived the ideas and designed method; QMRW

conducted fieldwork QMRW and generated social networks; QMRW, CMP, and KAK

conducted movement and spatial analysis; QMRW led the writing of the manuscript; EVW

developed the research program. All authors contributed critically to drafts and gave final

approval for publication.

**Competing interests.** We have no competing interests.

**Funding.** Funding for this study was provided by Natural Sciences and Engineering Research

Council (NSERC) Vanier Canada Graduate Scholarships to QMRW and CMP, a NSERC

Canada Graduate Scholarship (Masters and Doctoral) to KAK, and a NSERC Discovery Grant to

EVW.

**Acknowledgements.** We respectfully acknowledge the territory in which data were collected

and analyzed as the ancestral homelands of the Beothuk and the Island of Newfoundland as the

ancestral homelands of the Mi'kmaq and Beothuk. We thank M. Laforge, M. Bonar, J. Hendrix,

and R. Huang for help in the field, J. Turner for help with statistical analysis, and D. Wilson, K.

Lewis, I. Fleming, G. Albery, and members of the Wildlife Evolutionary Ecology Lab for helpful

comments on previous versions of this manuscript. We also thank members of the Newfoundland

and Labrador Wildlife Division, including S. Moores, B. Adams, W. Barney, and J. Neville for

facilitating animal captures and for logistical support in the field.

**Table 1.** Summary of winter community assignment for three caribou social networks (2017–

2019), including the number of individuals (N), number of communities, community size,

modularity ($Q$), and the community assortativity coefficient ($R_{com} = 0$ indicates no confidence in

community assignment; $R_{com} = 1$ indicates certainty in community assignment).

| Year | N | Community | Community size | Home range area (km$^2$) | $Q$ | $R_{com}$ |
|---|---|---|---|---|---|---|
| 2017 | 10 | 1 | 3 | 84.71 | 0.13 | 0.95 |
| | | 2 | 7 | 43.43 | | |
| 2018 | 16 | 1 | 5 | 113.09 | 0.17 | 1.00 |
| | | 2 | 7 | 77.31 | | |
| | | 3 | 1 | 13.84 | | |
| | | 4 | 1 | 23.11 | | |
| | | 5 | 1 | 4.94 | | |
| | | 6 | 1 | 19.06 | | |
| 2019 | 12 | 1 | 2 | 32.39 | 0.13 | 1.00 |
| | | 2 | 9 | 33.87 | | |
| | | 3 | 1 | 1.26 | | |

**Figure 1.** Summary of the data pipeline used to generate integrated step selection analysis

(iSSA) models. Primary data sources were landcover data and caribou GPS relocation data,

which were combined to determine the physical locations of animals on the landscape. The

pairing of animal locations and landcover data was used to generate the comparison of used to

available points (panel a), which is the response variable in iSSA models, as well as the habitat

type in which a given relocation occurred: lichen (defined in text as open-forage), open (defined

in text as open-movement), and forest (panel b). Caribou relocation data were also used to

generate two movement parameters (panel c) and aspects of the social environment (panels d and

f). Movement parameters included turn angle, which is the angular deviation between the

headings of two consecutive steps, and step length, which is the linear distance between

consecutive relocations. The social environment included nearest neighbour distance (panel d)

and weekly social networks and the dyadic simple ratio index generated based on a moving-

window as a proxy for short-term social preference (panel e). The bottom row represents a

graphical formulation of our iSSA models, where habitat use (1:10 ratio of used to available

relocations) was regressed against habitat type (lichen, open, and forest), movement parameters

(step length and turn angle), nearest neighbour distance, and weekly dyadic simple ratio index.

**Figure 2.** Observed lagged association rate (LAR) for caribou in the same (blue lines) and

different (orange) annual winter social communities, calculated as the probability that any pair of

individuals associated on a given day, are still associated on subsequent days. Note, the time

period for LAR analysis and social community assignment was 1 January to 16 March. Error

bars represent the standard error of all pairwise association rates calculated on each day.

Individuals in the same social communities (blue lines) generally had higher lagged association

rates, suggesting they were more likely to associate together over time.

**Figure 3.** Habitat selection coefficients and 95% confidence intervals for the top integrated step

selection model. Coefficients that do not overlap zero are deemed significant (see Table S3).

Colours represent the model in which each variable or interaction was introduced and once

introduced into a model a given variable was included in subsequent models. Purple coefficients

represent variables initially included in the core model, orange coefficients represent variables

initially introduced in the nearest neighbour model, and green coefficients represent variables

462     initially introduced in the simple ratio index. Note, step length, nearest neighbour distances (both

463     start and end), and the simple ratio index were log-transformed and turn angle was cosine

464     transformed.

# References

1.  Van Moorter B, Rolandsen CM, Basille M, Gaillard J-M. 2016 Movement is the glue connecting home ranges and habitat selection. *J. Anim. Ecol.* **85**, 21–31. (doi:10.1111/1365-2656.12394)

2.  He P, Maldonado-Chaparro AA, Farine DR. 2019 The role of habitat configuration in shaping social structure: a gap in studies of animal social complexity. *Behav. Ecol. Sociobiol.* **73**, 9.

3.  Spiegel O, Leu ST, Bull CM, Sih A. 2017 What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecol. Lett.* **20**, 3–18. (doi:10.1111/ele.12708)

4.  Strandburg-Peshkin A, Papageorgiou D, Crofoot MC, Farine DR. 2018 Inferring influence and leadership in moving animal groups. *Philos. Trans. R. Soc. B* **373**, 20170006. (doi:10.1098/not)

5.  Chamaillé-Jammes S, Fritz H, Valeix M, Murindagomo F, Clobert J. 2008 Resource variability, aggregation and direct density dependence in an open context: The local regulation of an African elephant population. *J. Anim. Ecol.* **77**, 135–144. (doi:10.1111/j.1365-2656.2007.01307.x)

6.  Spiegel O, Sih A, Leu ST, Bull CM. 2018 Where should we meet? Mapping social network interactions of sleepy lizards shows sex-dependent social network structure. *Anim. Behav.* **136**, 207–215. (doi:10.1016/j.anbehav.2017.11.001)

7.  van Beest FM, Uzal A, Vander Wal E, Laforge MP, Contasti AL, Colville D, Mcloughlin PD. 2014 Increasing density leads to generalization in both coarse-grained habitat selection and fine-grained resource selection in a large mammal. *J. Anim. Ecol.* **83**, 147–156. (doi:10.1111/1365-2656.12115)

8.  Jolles JW, King AJ, Killen SS. 2019 The role of individual heterogeneity in collective animal behaviour. *Trends Ecol. Evol.* **35**, 278–291. (doi:10.1016/j.tree.2019.11.001)

9.  Strandburg-Peshkin A, Farine DR, Crofoot MC, Couzin ID. 2017 Habitat structure shapes individual decisions and emergent group structure in collectively moving wild baboons. *Elife* **6**, e19505. (doi:10.7554/eLife.19505)

10. Spiegel O, Leu ST, Bull CM, Sih A. 2017 What's your move? Movement as a link between personality and spatial dynamics in animal populations. *Ecol. Lett.* **20**, 3–18. (doi:10.1111/ele.12708)

11. Lima SL. 1995 Back to the basics of anti-predatory vigilance: the group-size effect. *Anim. Behav.* **49**, 11–20.

12. Creel S, Schuette P, Christianson D. 2014 Effects of predation risk on group size, vigilance, and foraging behavior in an African ungulate community. *Behav. Ecol.* **25**, 773–784. (doi:10.1093/beheco/aru050)

13. Harel R, Spiegel O, Getz WM, Nathan R. 2017 Social foraging and individual consistency in following behaviour: testing the information centre hypothesis in free-ranging vultures. *Proc. R. Soc. B* **284**, 20162654.

14. Lachlan RF, Crooks L, Laland KN. 1998 Who follows whom? Shoaling preferences and social learning of foraging information in guppies. *Anim. Behav.* **56**, 181–190. (doi:10.1006/anbe.1998.0760)

15. Giraldeau LA, Dubois F. 2008 Social foraging and the study of exploitative behavior. *Adv. Study Behav.* **38**, 59–104. (doi:10.1016/S0065-3454(08)00002-8)

16. Patin R, Fortin D, Sueur C, Chamaillé-Jammes S. 2019 Space use and leadership modify
dilution effects on optimal vigilance under food-safety trade-offs. *Am. Nat.* **193**, E15–E28.
(doi:10.1086/700566)

17. Spiegel O, Crofoot MC. 2016 The feedback between where we go and what we know—
information shapes movement, but movement also impacts information acquisition. *Curr.*
*Opin. Behav. Sci.* **12**, 90–96. (doi:10.1016/j.cobeha.2016.09.009)

18. Jones TB, Patrick SC, Evans JC, Wells MR. 2020 Consistent sociality but flexible social
associations across temporal and spatial foraging contexts in a colonial breeder. *Ecol. Lett.*
**23**, 1085–1096. (doi:10.1111/ele.13507)

19. Giraldeau L-A, Caraco T. 2018 *Social foraging theory*. Princeton University Press.

20. Mourier J, Vercelloni J, Planes S. 2012 Evidence of social communities in a spatially
structured network of a free-ranging shark species. *Anim. Behav.* **83**, 389–401.
(doi:10.1016/j.anbehav.2011.11.008)

21. Best EC, Seddon JM, Dwyer RG, Goldizen AW. 2013 Social preference influences female
community structure in a population of wild eastern grey kangaroos. *Anim. Behav.* **86**,
1031–1040. (doi:10.1016/j.anbehav.2013.09.008)

22. Muller Z, Cantor M, Cuthill IC, Harris S. 2018 Giraffe social preferences are context
dependent. *Anim. Behav.* **146**, 37–49. (doi:10.1016/j.anbehav.2018.10.006)

23. Silk JB. 2007 Social components of fitness in primate groups. *Science (80-. ).* **317**, 1347–
1351.

24. Spiegel O, Leu ST, Sih A, Bull CM. 2016 Socially interacting or indifferent neighbours?
Randomization of movement paths to tease apart social preference and spatial constraints.
*Methods Ecol. Evol.* **7**, 971–979. (doi:10.1111/2041-210X.12553)

25. Pinter-Wollman N *et al.* 2013 The dynamics of animal social networks: Analytical,
conceptual, and theoretical advances. *Behav. Ecol.* **25**, 242–255.
(doi:10.1093/beheco/art047)

26. Croft DP, Darden SK, Wey TW. 2016 Current directions in animal social networks. *Curr.*
*Opin. Behav. Sci.* **12**, 52–58. (doi:10.1016/j.cobeha.2016.09.001)

27. Owen-Smith N, Fryxell JM, Merrill EH. 2010 Foraging theory upscaled: the behavioural
ecology of herbivore movement. *Philos. Trans. R. Soc. B* **365**, 2267–2278.
(doi:10.1098/rstb.2010.0095)

28. Bergerud AT, Mercer WE. 1989 Caribou introductions in eastern North America. *Wildl.*
*Soc. Bull.* **17**, 111–120.

29. Lesmerises F, Johnson CJ, St-Laurent M-H. 2018 Landscape knowledge is an important
driver of the fission dynamics of an alpine ungulate. *Anim. Behav.* **140**, 39–47.
(doi:10.1016/j.anbehav.2018.03.014)

30. Bergerud AT. 1974 Relative abundance of food in winter for Newfoundland caribou.
*Oikos* **25**, 379–387.

31. Peignier M, Webber QMR, Koen EL, Laforge MP, Robitaille AL, Vander Wal E. 2019
Space use and social association in a gregarious ungulate: Testing the conspecific
attraction and resource dispersion hypotheses. *Ecol. Evol.* **9**, 5133–5145.
(doi:10.1002/ece3.5071)

32. Boertje RD. 1985 An Energy Model for Adult Female Caribou of the Denali Herd,
Alaska. *J. Range Manag.* **38**, 468–473. (doi:10.2307/3899725)

33. Duquette LS, Klein DR. 1987 Activity budgets and group size of caribou during spring
migration. *Can. Jounal Zool.* **65**.

34. Fortin D, Courtois R, Etcheverry P, Dussault C, Gingras A. 2008 Winter selection of
landscapes by woodland caribou: Behavioural response to geographical gradients in
habitat attributes. *J. Appl. Ecol.* **45**, 1392–1400. (doi:10.1111/j.1365-2664.2008.01542.x)
35. Lewis KP. 2014 Sustainable Development and Strategic Science Branch Land Cover
Classification. **5**.
36. Bjørneraas K, Van Moorter B, Rolandsen CM, Herfindal I. 2010 Screening Global
Positioning System Location Data for Errors Using Animal Movement Characteristics. *J.*
*Wildl. Manage.* **74**, 1361–1366. (doi:10.2193/2009-405)
37. Avgar T, Potts JR, Lewis MA, Boyce MS. 2016 Integrated step selection analysis:
Bridging the gap between resource selection and animal movement. *Methods Ecol. Evol.*
**7**, 619–630. (doi:10.1111/2041-210X.12528)
38. Fortin D, Beyer HL, Boyce MS, Smith DW, Duchesne T, Mao JS. 2005 Wolves influence
elk movements: Behavior shapes a trophic cascade in Yellowstone National Park. *Ecology*
**86**, 1320–1330.
39. Prokopenko CM, Boyce MS, Avgar T. 2017 Characterizing wildlife behavioural
responses to roads using integrated step selection analysis. *J. Appl. Ecol.* **54**, 470–479.
(doi:10.1111/1365-2664.12768)
40. Signer J, Fieberg J, Avgar T. 2019 Animal movement tools (amt): R package for
managing tracking data and conducting habitat selection analyses. *Ecol. Evol.* **9**, 880–890.
(doi:10.1002/ece3.4823)
41. Duchesne T, Fortin D, Rivest LP. 2015 Equivalence between step selection functions and
biased correlated random walks for statistical inference on animal movement. *PLoS One*
**10**, 1–12. (doi:10.1371/journal.pone.0122947)
42. R Core Team. 2019 R: A language and environment for statistical computing. ,
https://www.r-project.org/.
43. Robitaille AL, Webber QMR, Vander Wal E. 2019 Conducting social network analysis
with animal telemetry data: applications and methods using spatsoc. *Methods Ecol. Evol.*
**10**, 1203–1211. (doi:10.1111/2041-210X.13215)
44. Kasozi H, Montgomery RA. 2020 Variability in the estimation of ungulate group sizes
complicates ecological inference. *Ecol. Evol.* , 1–9. (doi:10.1002/ece3.6463)
45. Cairns SJ, Schwager SJ. 1987 A comparison of association indices. *Anim. Behav.* **35**,
1454–1469.
46. Farine DR, Whitehead H. 2015 Constructing, conducting and interpreting animal social
network analysis. *J. Anim. Ecol.* **84**, 1144–1163. (doi:10.1111/1365-2656.12418)
47. Csárdi G, Nepusz T. 2006 The igraph software package for complex network research.
*InterJournal Complex Syst.* **1695**, 1–9.
48. Newman MEJ. 2006 Modularity and community structure in networks. *Proc. Natl. Acad.*
*Sci.* **103**, 8577–8582.
49. Farine DR. 2014 Measuring phenotypic assortment in animal social networks: Weighted
associations are more robust than binary edges. *Anim. Behav.* **89**, 141–153.
(doi:10.1016/j.anbehav.2014.01.001)
50. Shizuka D, Farine DR. 2016 Measuring the robustness of network community structure
using assortativity. *Anim. Behav.* **112**, 237–246.
51. Farine DR. 2013 Animal social network inference and permutations for ecologists in R
using asnipe. *Methods Ecol. Evol.* **4**, 1187–1194. (doi:10.1111/2041-210X.12121)
52. Whitehead H. 2008 *Analyzing animal societies: quantitative methods for vertebrate social*

*analysis.* University of Chicago Press.

53.　Worton BJ. 1989 Kernel methods for estimating the utilization distribution in home-range
studies. *Ecology* **70**, 164–168.

54.　Fieberg J, Kochanny CO. 2005 Quanitfying home-range overlap: the importance of the
utilization distribution. *J. Wildl. Manage.* **69**, 1346–1359. (doi:10.2193/0022-
541X(2005)69)

55.　Muff S, Signer J, Fieberg J. 2020 Accounting for individual-specific variation in habitat-
selection studies: Efficient estimation of mixed-effects models using Bayesian or
frequentist computation. *J. Anim. Ecol.* **89**, 80–92. (doi:10.1111/1365-2656.13087)

56.　Avgar T, Lele SR, Keim JL, Boyce MS. 2017 Relative Selection Strength: Quantifying
effect size in selection inference. *Ecol. Evol.* **7**, 5322–5330. (doi:10.1002/ece3.3122)

57.　Shizuka D, Chaine AS, Anderson J, Johnson O, Laursen IM, Lyon BE. 2014 Across-year
social stability shapes network structure in wintering migrant sparrows. *Ecol. Lett.* **17**,
998–1007. (doi:10.1111/ele.12304)

58.　Cantor M, Wedekin LL, Guimarães PR, Daura-Jorge FG, Rossi-Santos MR, Simões-
Lopes PC. 2012 Disentangling social networks from spatiotemporal dynamics: The
temporal structure of a dolphin society. *Anim. Behav.* **84**, 641–651.
(doi:10.1016/j.anbehav.2012.06.019)

59.　Jacobs A. 2010 Group cohesiveness during collective movements: Travelling apart
together. *Behav. Processes* **84**, 678–680. (doi:10.1016/j.beproc.2010.03.004)

60.　Morrell LJ, Ruxton GD, James R. 2011 Spatial positioning in the selfish herd. *Behav.*
*Ecol.* **22**, 16–22. (doi:10.1093/beheco/arq157)

61.　Heesen M, Macdonald S, Ostner J, Schülke O. 2015 Ecological and Social determinants
of group cohesiveness and within-group spatial position in wild assamese macaques.
*Ethology* **121**, 270–283. (doi:10.1111/eth.12336)

62.　Barrette C, Vandal D. 1986 Social rank, dominance, antler size, and access to food in
snow-bound wild woodland caribou. *Behaviour* **97**, 118–145.

63.　Hirotani A. 1990 Social organization of reindeer (*Rangifer tarandus*), with a special
reference to relationships among females. *Can. J. Zool.* **68**, 743–749.

64.　Hamilton WD. 1971 Geometry for the selfish herd. *J. Theor. Biol.* **31**, 295–311.
(doi:10.1016/0022-5193(71)90189-5)

65.　Mayor SJ, Schaefer JA, Schneider DC, Mahoney SP. 2009 The spatial structure of habitat
selection: A caribou's-eye-view. *Acta Oecologica* **35**, 253–260.
(doi:10.1016/j.actao.2008.11.004)

66.　Jung TS, Larter NC, Allaire DG. 2019 Social organization of boreal woodland caribou
(*Rangifer tarandus caribou*) in response to decreasing annual snow depth. *Mammal*
*ReLsearch* **64**, 377–385.

67.　Schaefer JA, Mahoney SP. 2001 Antlers on female caribou: Biogeography of the bones of
contention. *Ecology* **82**, 3556–3560.

68.　Croft DP, Madden JR, Franks DW, James R. 2011 Hypothesis testing in animal social
networks. *Trends Ecol. Evol.* **26**, 502–507. (doi:10.1016/j.tree.2011.05.012)

69.　Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE. 2008 A
movement ecology paradigm for unifying organismal movement research. *Proc. Natl.*
*Acad. Sci.* **105**, 19052–19059. (doi:10.1073/pnas.0800375105)

70.　McLoughlin PD, Morris DW, Fortin D, Vander Wal E, Contasti AL. 2010 Considering
ecological dynamics in resource selection functions. *J. Anim. Ecol.* **79**, 4–12.

(doi:10.1111/j.1365-2656.2009.01613.x)

71.    Webber QMR, Vander Wal E. 2018 An evolutionary framework outlining the integration

of individual social and spatial ecology. *J. Anim. Ecol.* **87**, 113–127. (doi:10.1111/1365-

2656.12773)