A benchmarking of human mitochondrial DNA haplogroup classifiers from whole-genome and whole-exome sequence data

Víctor García-Olivares[1], Adrián Muñoz-Barrera[1], José Miguel Lorenzo-Salazar[1], Carlos Zaragoza-Trello[2], Luis A. Rubio-Rodríguez[1], Ana Díaz-de Usera[1], David Jáspez-Toledo[1], Antonio Iñigo Campos[1], Rafaela González-Montelongo[1,3], Carlos Flores[1,3,4,5]

1. Genomics Division, Instituto Tecnológico y de Energías Renovables (ITER), Santa Cruz de Tenerife, Spain.
2. Estación Biológica de Doñana (EBD-CSIC), Integrative Ecology, E-41092 Sevilla, Spain.
3. Instituto de Tecnologías Biomédicas (ITB), Universidad de La Laguna, Santa Cruz de Tenerife, Spain.
4. Research Unit, Hospital Universitario Nuestra Señora de Candelaria, Universidad de La Laguna, Santa Cruz de Tenerife, Spain.
5. CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain.

* Corresponding author: cflores@ull.edu.es

Abstract

The mitochondrial genome (mtDNA) is of interest for a range of fields including evolutionary, forensic, and medical genetics. Human mitogenomes can be classified into evolutionary related haplogroups that provide ancestral information and pedigree relationships. Because of this and the advent of high-throughput sequencing (HTS) technology, there is a diversity of bioinformatic tools for haplogroup classification. We present a benchmarking of the 11 most salient tools for human mtDNA classification using empirical whole-genome (WGS) and whole-exome (WES) short-read sequencing data from 36 unrelated donors. Besides, because of its relevance, we also assess the best performing tool in third-generation long noisy read WGS data obtained with nanopore technology for a subset of the donors. We found that, for short-read WGS, most of the tools exhibit high accuracy for haplogroup classification irrespective of the input file used for the analysis. However, for short-read WES, Haplocheck and MixEmt were the most accurate tools. Based on the performance shown for WGS and WES, and the accompanying qualitative assessment, Haplocheck stands out as the most complete tool. For third-generation HTS data, we also showed that Haplocheck was able to accurately retrieve mtDNA haplogroups for all samples assessed, although only after following assembly-based approaches (either based on a referenced-based assembly or a hybrid *de novo* assembly). Taken together, our results provide guidance for researchers to select the most suitable tool to conduct the mtDNA analyses from HTS data.

Keywords: Mitogenome, lineage, third-generation sequencing, next-generation sequencing.

1. Introduction

The human mtDNA is a circular double-stranded genome of 16,569 base pairs (bp) in the hg38 reference sequence, encoding 37 genes for 13 proteins, 22 tRNAs, and 2 rRNAs. Besides its key role in diverse human diseases (DeBalsi et al., 2017; Pyle et al., 2016; West & Shadel, 2017), distinctive features such as the matrilineal inheritance, the lack of recombination, and a higher mutation rate than the nuclear genome, makes mtDNA analysis a powerful tool also for population genetics (Brotherton et al., 2013; Llamas et al., 2016; Posth et al., 2016) and forensic studies (Børsting & Morling, 2015; Just et al., 2009).

Worldwide human mtDNA diversity has been reconstructed through a genealogy of distinctive lineages, representing mtDNA sequences clustered into evolutionarily related haplotypes (a.k.a. haplogroups). They are linked to human evolutionary history and allow tracing origins and differentiation patterns across populations and over time periods (Balter, 2011; Fu et al., 2013; Gonder et al., 2007; Hajdinjak et al., 2018). As such, many haplogroups are associated with specific biogeographical ancestries (Chan et al., 2019; De Angelis et al., 2018; Maca-Meyer et al., 2001). MtDNA haplogroup classification has become an essential step to recovery ancestry and genealogical information from analyzed samples (Weissensteiner et al., 2016). One of the most popular and continuously updated repositories of mtDNA lineage relationships and nomenclature is Phylotree (http://www.phylotree.org) (van Oven & Kayser, 2009). The latest version of Phylotree (Build 17) contains more than 5,400 haplogroups and it constitutes the central reference for many bioinformatic tools to classify human mtDNA sequences.

The advent of high-throughput sequencing (HTS) technology has allowed the development of a wide range of applications, including whole-genome sequencing (WGS) and whole-exome sequencing (WES). MtDNA studies have also leveraged the power offered by HTS (Churchill et al., 2018; Schönberg et al., 2011; Vasta et al., 2009). In fact, mtDNA can be fully captured by commercial WES solutions, possibly due to the high copy number of mtDNA (Picardi & Pesole, 2012). Hence, mtDNA information can be recovered effectively from WES studies at no extra cost. Furthermore, the sequencing depth associated with WES usually allows reconstructing the entire mitogenome with high quality and detecting heteroplasmic sites (Sosa et al., 2012). Third-generation HTS, such as those based on nanopore technology (ONT, Oxford Nanopore Technologies, Oxford, UK), allows sequencing with long reads, providing an opportunity to generate full-length mtDNA sequences in single reads.

Because of the importance of recovering ancestral information and pedigree relationships of study samples, the number of bioinformatics tools developed for mtDNA haplogroup classification has increased notably in the last decade to adapt to the HTS technologies (Calabrese et al., 2014; Fan & Yao, 2011; Ishiya & Ueda, 2017; Kim et al., 2020; Navarro-Gomez et al., 2015; Röck et al., 2013; Smieszek et al., 2018; Vohr et al., 2017; Weissensteiner et al., 2016, 2020). An unmet need is a benchmarking study of their capabilities, requirements, and usability to correctly classify the mtDNA haplogroups from different source files. Based on short-read WES and WGS data from the same donors, as well as on WGS obtained from long noisy reads from a subset of them, here we present an empirical evaluation of the 11 most salient bioinformatic tools available for human mtDNA classification from HTS data.

2. Materials and methods

Samples, library preparation and sequencing

The study was approved by the Research Ethics Committee of the Hospital Universitario Nuestra Señora de Candelaria and performed according to The Code of Ethics of the World Medical Association (Declaration of Helsinki).

Thirty-six DNA samples from unrelated donors of European descent were used for the study after informed consent (see supplementary **Table S1**). DNA was extracted from peripheral blood using a commercial column-based solution (GE Healthcare, Chicago, IL). DNA quantifications were performed in a Qubit dsDNA HS Assay (Thermo Fisher, Waltham, MA). All samples were sequenced in parallel using short-read WGS and WES. Library constructions were performed with Illumina preparation kits following the manufacturer's recommendations (Illumina Inc., San Diego, CA). The Nextera DNA Prep kit was used for WGS, except for six samples that were processed by Illumina DNA Prep. The same samples were processed with Nextera DNA Exome with a 350 bp insert size as described elsewhere (Díaz-de Usera et al., 2020). The library quality control was carried out in a TapeStation 4200 (Agilent Technologies, Santa Clara, CA). Sequencing was conducted on a HiSeq 4000 Sequencing System (Illumina Inc.) at the Instituto Tecnológico y de Energías Renovables (Santa Cruz de Tenerife, Spain).

Eight of these samples had WGS based on noisy long-read data obtained at KeyGene (Wageningen, The Netherlands). Briefly, sequencing was performed on a PromethION system (ONT) for 64 h using one FLO_PR002 (R9.4.1 pore) flow cell per sample following the manufacturer's recommendations. Basecalling was performed on the PromethION computing module using MinKNOW v1.14.2 with Guppy v2.2.2 and data preprocessing was carried out with PycoQC (Leger & Leonardi, 2019).

Sequence processing and variant calling

Processing of short-read WGS and WES data was carried out using an in-house pipeline (see supplementary **Figure S1**) based on GATK v4.1 for WGS and GATK v3.8 for WES (McKenna et al., 2010). Reads were aligned to the GRCh37/hg19 reference genome and the mtDNA reads realigned to the revised Cambridge Reference Sequence (rCRS), GenBank NC_012920 (Anderson et al., 1981; Andrews et al., 1999), following GATK best practices for this circular genome (see supplementary **Figure S1**). This required two alignment steps: one against the canonical mitogenome reference and another against the same reference but shifted by 8,000 nucleotide positions. This generates a displacement of the mtDNA D-loop to the opposite side of the contig, allowing to better capture variants of this highly variable region. The alignments were then processed for duplicate marking and base quality score recalibration (DePristo et al., 2011). The variant calls were obtained with the "mitochondria mode" of Mutect2 GATK v4.1. This specific mode provides a better sensitivity for this genome as it shows a robust detection of very low fractions of variants once the nuclear mtDNA segments (NuMT) are filtered out. The mtDNA variants were then filtered by using FilterMutectCalls and VariantFiltration tools to keep the variants classified as PASS.

For ONT data, we first extracted all reads aligning to the mtDNA genome and then used four alternative strategies to obtain sequence variation for mtDNA classification: a) one based on the alignment of reads to the rCRS sequence with Minimap2 (Li, 2018) followed by a variant-calling step with Medaka (https://github.com/nanoporetech/medaka); b) another relying on the reference-based assembly performed by Rebaler (https://github.com/rrwick/Rebaler); c) a third strategy based on *de novo* assembly with Miniasm (Li, 2016) followed by nine rounds of polishing with Racon (Vaser et al., 2017) and a final step with RagTag (Alonge et al., 2019) for scaffolding; and d) the last strategy combining ONT and Illumina reads in a hybrid *de novo* assembly built with Unicycler (Wick et al., 2017) and a final step with RagTag for scaffolding (see supplementary **Figure S2**). Quality of assemblies were assessed with QUAST (Gurevich et al., 2013). For all the alternatives, a VCF file per sample was generated for the haplogroup classification step.

For each sample, three different files were generated for short-read data: BAM and VCF files created through the pipeline previously described, and FASTA files obtained by the VCF-consensus script included in vcftools (Danecek et al., 2011) generated from the VCF files. Of note, an additional filter was applied to the VCF files discarding variants that showed an allele fraction below 0.9, a threshold applied by default in Haplogrep during the haplogroup classification. This enabled the harmonization of the genetic variation considered in FASTA and VCF files.

mtDNA haplogroup classification

Among the tools available from the literature, we selected 11 that were published in the last three years or that were cited at least 30 times since its description (**Table 1**). Haplogrep, Haplocheck and Phy-Mer have the option of using alternative input files, fostering an evaluation with the alternative supported format files. Some of the tools provide quality scores supporting the mtDNA haplogroup classification. In these cases, only the classification with the best scoring was considered for the analyses. For the haplogroup classification process, the tools that were designed to be run locally, either through an application or via command line, were executed on a 4-core Intel Core i7 CPU at 2.6 GHz and 16 GB of RAM.

Table 1. List of software assessed for human mtDNA haplogroup classification. All the tools assessed classify mtDNA sequences according to the PhyloTree nomenclature by using the latest version Build 17. Two of them, MitoTool and Phy-Mer, remain outdated in regards of the PhyloTree built when this study was conducted.

Tool	User interface	Release year	Version	Latest release	Database version	Input options	Execution time
James Lick's[a]	Web	2010	0.19a	2013.04.08	Phylotree 17	FASTA	< 1 min
Haplogrep[b]	Web / CLI*	2011	2.2.4	2016.07.08	Phylotree 17	VCF/FASTA	< 1 min
MitoTool[c]	App	2011	1.1.2	2013.11.02	Phylotree 16	FASTA	< 1 min
Haplofind[d]	Web	2013	-	2013.01.12	Phylotree 17	FASTA	< 1 min
EMMA[e]	Web	2013	13	2019.11.28	Phylotree 17	FASTA	< 1 min
Phy-Mer[f]	Web / CLI	2015	1.0.0	2016.01.14	Phylotree 16	BAM/FASTA	1 − 5 min / < 1 min
MitoSuite[g]	App	2017	1.0.9	2017.06.06	Phylotree 17	BAM	1 − 5 min
MixEmt[h]	CLI	2017	0.1	2017.05.09	Phylotree 17	BAM	WGS > 1 hour | WES < 1 hour
Haplocheck[i]	Web / CLI	2019	1.1.0	2019.11.17	Phylotree 17	BAM/VCF	< 1 min
Haplotracker[j]	Web	2020	-	2020.04.23	Phylotree 17	FASTA	< 1 min
HaploGrouper[k]	CLI	2020	-	2020.08.17	Phylotree 17	VCF	< 1 min

a: https://dna.jameslick.com/mthap/; b: (Weissensteiner et al., 2016); c: (Fan & Yao, 2011, 2013); d: (Vianello et al., 2013); e: (Röck et al., 2013); f: (Navarro-Gomez et al., 2015) ; g: (Ishiya & Ueda, 2017); h: (Vohr et al., 2017); i: (Weissensteiner et al., 2020); j: (Kim et al., 2020); k: (Jagadeesan et al., 2020).

Statistical analyses

All statistical analyses were performed in R v.3.3.3 (R Core Team 2017). Model fitting was evaluated visually using the 'DHARMa' package (Hartig, 2017) (v.0.1.5). Prior to analysis, all predictors were standardized by subtracting the mean and dividing by the standard deviation with the 'scale' base function in R.

In order to evaluate the performance of the 11 tools in the classification, we ran generalized linear mixed models (GLMM) with the tools as fixed factor and the classification results (concordance/discordance) as the response variable. This was done separately for WGS and WES data. In the models, we included the sample as a random factor to account for differences introduced by the sequencing runs and the diverse enrichment kits involved. Classification results were transformed to a binary response so that, for each sample, discordance between the consensus haplogroup and the classification result was coded as one, and the concordance as zero. In such binary outcome data, the models may suffer from complete separation when one of the levels of an explanatory variable explains completely the binomial response variable, precluding an optimal fitting of the algorithm (Heinze & Schemper, 2002). This prevents the algorithm from properly fitting the coefficient for this level. To overcome this issue, we fitted the models using the bglmer function of the "blme" v.1.0.4 R package (Chung et al., 2013), which incorporates a similar algorithm to "lmer4" v.1.1.15 R package (Bates et al., 2015) to facilitate the estimation with the incorporation of a prior. Before the analysis, a Pearson's correlation test was conducted to examine the cross-correlation among the different sequencing parameters measured (i.e., mapped reads, duplicate reads, depth, mean mapping quality, and base quality). Pearson's correlation test revealed that the proportion of duplicate reads and the depth of coverage were highly correlated ($r < 0.8$). Therefore, only the mapped reads, mapping quality, and base quality were incorporated as covariates in the model (see supplementary **Table S2**). Finally, to evidence the pairwise differences between the tools, we run a *post-hoc* analysis with the Tukey contrast by using the "multcomp" v.1.4.8 package (Hothorn et al., 2008). Model outputs were visualized using 'effects' v.0.9.4 (Fox, 2003) and 'ggplot2' v.2.2.1 (Wickham, 2016).

3. Results

Short-read sequencing summary of mitogenomes

The mean (± SD) number of mtDNA reads recovered per sample (n=36) for short-read WGS and WES data were 197,717 ± 98,719 and 9,905 ± 6,808, respectively. For WGS, 100% of the mitogenome was covered at least at 1X. For WES, this percentage decreased to a mean (± SD) of 86.79% ± 27.01 of the recovered mitogenome. The mean (± SD) mapping quality for mapped reads had a value of 59.9 ± 0.01 and the Phred base-quality scores estimated were high for both applications (28.38 ± 0.91 for WGS, and 29.74 ± 0.02 for WES). The average (± SD) depth recovered for WGS was 1,119X ± 433 (range: 554-2,577X), decreasing to 37X ± 20 for WES (range: 11-92X). Besides, while WGS provided a homogeneous depth of coverage profile across the mitogenomes, those

recovered by WES showed a highly heterogeneous profile across samples. Interestingly, the region between nucleotide positions 2,000-3,000 was highly enriched in reads from Illumina WES data (**Figure 1**). As for the number of detected variants after filtering, the mean (± SD) depth of coverage per variant call had a value of 958 ± 362 in WGS, decreasing to 40 ± 21 in WES. Out of the 36 samples sequenced, ten showed the equivalent number of variants by WGS and WES. In the remaining samples, the number of variants detected by WES was lower than by WGS: 17 samples missed between 1 and 3 variants, and nine missed more than 30% of the variants (**Table S1**).

Figure 1. Circos plot of the depth of coverage for short-read and long-read sequencing in the mtDNA of an exemplar sample. Short-read WGS and WES data are colored in green and red, respectively. Long-read WGS data is shown in blue.

Haplogroup classification based on short-read data

Twenty-eight samples showed >90% concordance in the haplogroup classification among the 11 tools. The remaining showed a mean concordance rate of 69.23% (see supplementary **Table S3)**. In addition, a higher classification accuracy was provided for WGS, reaching an average of 90.08%, while for WES it decreased to an average of 76.98% (see supplementary **Table S4**). On average (± SD), there were more discordances on the WES classifications (2.97 ± 3.92) than for WGS (1.39 ± 1.48). Based on this evidence, we set the WGS-derived haplogroup as the ground truth.

By the classification accuracy, Haplogrep, Haplocheck, EMMA, and James Lick's mtHap classified all samples correctly based on WGS. For the WES data, only Haplocheck classified all haplogroups correctly, and only when the BAM file was used as input. Concerning the less accurate classifiers, MixEmt showed the lowest accuracy for WGS, only classifying precisely 30.55% of the analyzed samples. MitoTool was the least accurate tool for WES data, yielding an incorrect haplogroup classification in 41.67% of the samples. The execution time required for haplogroup classification was also very different between the tools (**Table 1**). As expected,

classifications that relied on BAM inputs had a processing time higher than those running with FASTA and VCF files. Among the tools that support BAM file inputs, MixEmt was the most time-consuming, in which the heterogeneity of processing time among samples was likely due to the different number of mtDNA reads between datasets. In case of Haplocheck, which was designed along with MixEmt to estimate the potential cross-contamination of samples, it required less time per sample for the haplogroup classification.

Modelling of the haplogroup classification accuracy

Based on these results, we modelled the performance of the 11 tools for short-read mtDNA data classification. The total variance explained by a fixed effects model was 57.69% and 46.19% for WGS and WES data, respectively. The predicted probabilities were then estimated in order to assess the haplogroup classification accuracy of each tool both on WES and WGS data (**Figure 2**). Among all tools, MixEmt (estimate ± SE; 3.55 ± 0.66; $p < 0.001$) was the one with the lowest accuracy in correctly classifying the haplogroup from WGS (80.91% predicted probability of incorrect classification). The rest of the tools showed a negligible probability of misclassifying the haplogroup from WGS (**Figure 2** and **Table S5**). *Post-hoc* tests did not show statistically significant differences among them. For WES, Haplocheck (-4.38, ± 1.99, $p = 0.03$) and MixEmt (-1.13, ± 0.50, $p = 0.02$), both using BAM as the input file, were the tools showing the highest accuracy for haplogroup classification (**Figure 2** and **Table S5**). Haplocheck correctly classified the haplogroup for all samples, while MixEmt incorrectly classified four of the 36 samples analyzed. Despite that, *post-hoc* tests did not show significant differences in the classification performance between Haplocheck and MixEmt. On the contrary, MitoTool (0.97, ± 0.49, $p = 0.05$) showed the lowest accuracy in the haplogroup classification based on WES, for which the predicted probability of incorrectly classifying an haplogroup was ≥25%. Regarding the effect of the covariates, their effect was negligible for WGS data. However, for the WES data, a relatively high effect size was found for the mapped reads (-5.96 ± 3.55; $p = 0.09$) and the mapping quality (1.16 ± 0.39; $p < 0.001$) (**Table S5**). Taken together, these results suggest that the number of reads available and their mapping quality have stronger effects on the probability of inaccurate haplogroup classification based on WES data than based on WGS data.

Figure 2. Predicted probability values (and 95% confidence intervals) of the GLMM model estimated for each tool for a) WGS and b) WES datasets. In light grey, the raw data from the haplogroup classification results.

JML: James Lick's, **HPG**: Haplogrep, **MTO**: MitoTool, **HPF**: Haplofind, **EMA**: EMMA, **PHY**: Phy-Mer, **MTS**: MitoSuite, **MIX**: MixEmt, **HPC**: Haplocheck, **HPT**: Haplotracker, **HPR**: HaploGrouper.

Qualitative assessment of haplogroup classification tools

Due to the large number of haplogroup classification tools, a secondary aim of this study was to provide a guidance for researchers to select the most suitable tool for their analyses, not only based on the mtDNA haplogroup classification accuracy but also based on different software features and the usability of the evaluated tools. In order to facilitate the comparison among different tools, a qualitative assessment table is provided with the advantages and limitations of each tool. For this, the following characteristics were considered: haplogroup classification accuracy, computation time for classification, whether or not the latest Phylotree database is used, ability to process cohorts, versatility in the input files supported, user-friendly interface, frequency of tool maintenance, and the presence of others major functions (**Table 2**). Overall, Haplocheck proved to be the most complete tool, achieving the best performance in over 90% of the evaluated features. At the opposite end, Phy-Mer, with more than 50% of the features resulting poorly classified was the tool with the worst performance among all the assessed tools in this study.

Table 2. Qualitative assessment of mtDNA haplogroup classification tools. Performance of each tool is evaluated across different features and represented on a color scale based on the level of performance: green for good, orange for fair, and red for low performance.

Haplogroup classification accuracy of each tool was categorized into three ranges based on the predicted probabilities by each application: tools with a predicted probability of incorrectly classifying a haplogroup below 1% were represented as good performance, for a fair performance was established a range between 1.01% and 10%, and as low performance tools with a predicted probability higher than 10.01%. For computation time, three intervals were established: tools that classified samples in less than a minute were defined as good performance, from 1 to 5 minutes were categorized as fair performance, and those tools that required more than an hour were represented as low performance. Regarding the PhyloTree database used, it was represented as low performance those tools which are not updated to the latest version of Phylotree, Build 17, and those that allow the latest version as high performance. Multi-sample function was evaluated based on the possibility of cohort analysis. Tools that allow to process these functions were defined as good performance, tools that allow processing several samples by using a loop through a command-line were categorized as fair performance, and tools without this ability were represented as low performance. Based on the file format supported two categories were established: tools that support various input format files categorized as high performance and those that only support one format file that are represented as low performance. The user interface was divided into two categories, tools based on web or desktop applications defined as good performance, and tools developed exclusively for CLI as low performance. The tool maintenance was classified into two classes, tools updated continually or have been recently released, both identified as good performance, and tools that have not been updated during the last years. The last feature is the presence or not of additional functions; tools that have other functions implemented are categorized as good performance. Those tools without more functions were determined as low performance.

Assessment of ONT data for mtDNA haplogroup classification

The mean (± SD) number of mtDNA reads per sample (n=8) recovered by ONT was 2,812 ± 1,003. The average mtDNA coverage depth was 576X (range: 358-910X). The ONT-recovered mitogenome profile was uniform. However, a decrease in the coverage depth was observed for the D-loop region (positions between 16,024-576) (**Figure 1**). Regarding the quality of the recovered assemblies, the hybrid *de novo* and reference-based assembly strategies reached a similar value of N50, with 16,571 bp and 16,569 bp, respectively. However, for the long-read only *de novo* assembly, this length decreased to 16,407 bp. The consensus overall identity with the rCRS sequence reached a value of 100% for hybrid *de novo* assembly, 99.97% for reference-based strategy, and 88.99% for the long-read only *de novo* assembly strategy. Taking the called variants by short-read WGS as the ground truth, the variant calling strategy for ONT data shared a consensus average (± SD) of 76.41% (± 5.43), the reference-based assembly shared an average of 93.95% (± 5.76), *de novo* assembly shared an average of 89.13% (± 5.56), and the hybrid *de novo* assembly shared an average of 97.35% (± 3.34). In addition, the strategies that only use ONT reads for mtDNA assembly called a larger number of variants that were undetected by short-read WGS. The reference-based assembly strategy called a total of 61 (± 31) variants and *de novo* assembly 158 (± 173). For the variant calling and hybrid *de novo* assembly strategies, the number of novel variants called was lower, decreasing to an average of 3 (± 2) and 1 (± 2), respectively.

Given that Haplocheck was the best performing classifier based on short-read data, we then used it for ONT mtDNA haplogroup classification. The reference-based assembly and the hybrid *de novo* assembly strategies classified all samples correctly. However, both the variant calling and the *de novo* assembly approaches incorrectly classified one out of the eight samples analyzed. Despite that, all samples were classified correctly at the macro-haplogroup level with all strategies (**Table 3**).

Table 3. Long-read (ONT) sequencing summary and haplogroup classification results for the samples. The results of the short-read whole-genome sequencing (WGS) mtDNA classification are also shown.

Samples	ONT sequencing		Variant-calling		Reference-based assembly		De novo assembly		Hybrid de novo assembly		Short-read WGS	
	Mapped reads	Depth of coverage	Variants	Haplogroup	Variants	Haplogroup	Variants	Haplogroup	Variants	Haplogroup	Variants	Haplogroup
CAN03	3,065	581	14	H1ao1	69	H1ao1	119	H1ao1	16	H1ao1	14	H1ao1
CAN06	2,981	566	36	J2a2d	141	J2a2d	154	J2a2d	46	J2a2d	44	J2a2d
CAN07	2,343	444	15	H6a1b2	132	H6a1b2	175	H6a1b	17	H6a1b2	14	H6a1b2
CAN09	2,126	358	33	U4c1	66	U4c1	602	U4c1	37	U4c1	37	U4c1
CAN17	4,209	821	10	H2a2	59	H+16189	41	H+16189	15	H+16189	12	H+16189
CAN23	1,737	532	29	K1a1b1	69	K1a1b1	78	K1a1b1	37	K1a1b1	35	K1a1b1
CAN31	1,776	398	20	U6b1a1	67	U6b1a1	119	U6b1a1	28	U6b1a1	26	U6b1a1
CAN34	4,256	910	34	L3f1b1a	90	L3f1b1a	174	L3f1b1a	42	L3f1b1a	43	L3f1b1a

4. Discussion

Over the last decade, the number of mtDNA haplogroup classification tools has increased considerably. However, to our best understanding, there was no benchmarking study that assessed the accuracy of the haplogroup classification provided by them until now. This study presents a comparative of the 11 tools that are most widely used. The evaluation was done using empirical HTS data from two of the most widely used applications in human genetics, WES and WGS, in diverse empirical data. Besides, we also evaluated the best performing tool in long noisy ONT reads in a subset of the samples. Our results support that WES offers a suitable solution to allow accurate reconstruction of the mtDNA sequence, although at a lower depth of coverage than WGS. Our results also demonstrated that the most accurate tools for human mtDNA haplogroup classification from short-read WGS data are Haplogrep, Haplocheck, James Lick's, EMMA, HaploGrouper, and Haplotracker. On the contrary, for short-read WES, Haplocheck and MixEmt were the most accurate. Considering both the accuracy of classification for both approaches and the qualitative assessment made, Haplocheck demonstrated the best performance overall. In fact, using Haplocheck on long noisy ONT reads, we were able to accurately retrieve the mtDNA haplogroup from all assessed samples.

Previous studies have used WES data for human mitogenome reconstruction (Griffin et al., 2014; Patowary et al., 2017; Picardi & Pesole, 2012; Wortmann et al., 2015). Based on our comparisons between WGS and WES datasets, the most striking difference between their classification results was mainly related to the breadth and depth of coverage, both being better for WGS than for WES under our standards. Differences in mtDNA breadth of coverage were less pronounced. However, the mean depth of mtDNA for WGS was around 30 times higher than that obtained by WES. For WGS, the depth reported in diverse studies ranged between 1,200-4,000X (Puttick et al., 2019; Raymond et al., 2018; Watson et al., 2020), fitting with the expected proportion of mtDNA copy number compared to the nuclear DNA, which theoretically differs by 10 to 100 times (Al-Nakeeb et al., 2017; Robin & Wong, 1988). With respect to WES, the observed depth fits in the range described in the literature (Abicht et al., 2018; Diroma et al., 2020; Griffin et al., 2014; Patowary et al., 2017). Attending to the number of variants compared to those detected by WGS, a loss of variants was observed for WES in several samples. This may be explained by the low number of mapped reads of the mtDNA recovered in these samples, causing a shallower mtDNA depth. Despite that, based on the BAM-deduced haplogroup results, we and others (Diroma et al., 2020) have confirmed that WES can be an efficient approach to recover complete human mitogenomes, which allows retrieving the haplogroup with comparable accuracy to that based on WGS.

In general, a loss of accuracy in mtDNA classification was found for WES data in most of the evaluated tools. This could be explained by the lower depth of mitogenomes in WES compared to WGS, as has been described elsewhere (Ishiya & Ueda, 2019; Yin et al., 2019), given that a low depth of coverage will have negative consequences on the performance of the variant calling algorithms (Jennings et al., 2017; Petrackova et al., 2019). mtDNA haplogroup classification relies on a hierarchical algorithm based on the presence or absence of specific diagnostic variants defining the genealogy (Lee et al., 2008). Therefore, missing key variants due to the low depth may have a strong impact on the correct assignment in those tools that only support VCF and/or FASTA files as input. In contrast, the tools that can rely on BAM as an input file, given that this format contains all mapped reads -- including information about the alignment conditions and the complete sequence together

with the quality for each base -- may facilitate the proper haplogroup classification in sample datasets where certain variant positions are supported by a low number of reads. Hence, the BAM format might be optimal in samples where a low number of mtDNA reads is expected. The use of the BAM file as a possible input file for haplogroup classification has become popular in the last few years (Ishiya & Ueda, 2017; Navarro-Gomez et al., 2015; Vohr et al., 2017; Weissensteiner et al., 2020). With respect to the WES haplogroup results, the tools that support BAM as input files obtained the highest classification accuracy, except Phy-Mer, which was one of the tools performing worst in the classification. Haplocheck and MixEmt reached the highest accuracy among all the tools evaluated. Haplocheck excels in the correct classification of all samples, even with low depth of mtDNA, unlike MixEmt, which resulted in several misclassifications. On the other hand, our results from WES data also showed that the total number of mapped reads, which closely relates to the depth of coverage, has a strong effect on the haplogroup classification, affecting the classification of the samples with a low number of mtDNA reads. However, the effect of this variable was negligible in WGS datasets. Consequently, the total number of mtDNA mapped reads can be considered another key factor to take into account in the mtDNA classification from WES datasets.

Haplogrep, Haplocheck, James Lick's, EMMA, HaploGrouper, and Haplotracker tools yielded the highest accuracy scores based on WGS data, and their accuracy was independent of the input file format. This finding may be related to the high and uniform depth of WGS throughout the mitogenome, translating in strong support of variants by a high number of reads and, therefore, making more equivalent the information contained in the BAM, VCF, and FASTA files. Haplogrep and Haplocheck, both sharing the underlying algorithm developed first for Haplogrep and now integrated as a module directly in Haplocheck (Weissensteiner et al., 2020), were the ones providing the best ratings for all evaluated features. Despite that, taking into account all the evaluated features, Haplocheck stands out as the most complete haplogroup classification tool for WGS data as it also allows detecting potential sample contaminations based on BAM files. For those users who prefer working with mtDNA alignments in FASTA format and with the sole objective of classifying a limited number of samples by web-based user-friendly tools, EMMA, James Lick's, HaploGrouper, and Haplotracker are good choices as they all showed similar accuracies as Haplogrep and Haplocheck. EMMA is integrated into the EMPOP platform and stands out as one of the most complete and up-to-date databases of human mtDNA information, containing high quality representations of haplogroups from all over the world based on logical and phylogenetic measures suitable for forensic purposes (Amorim et al., 2019; W. Parson et al., 2014). James Lick's is one of the first tools released for mitochondrial haplogroup classification that is continually updated to new versions of the PhyloTree database. This tool is widely used among genetic genealogists because it is user friendly and based on a web application. HaploGrouper, the most recently released tool, allows classifying the haplogroups both for mtDNA and the non-recombining portion of the Y-chromosome, being unique in this dual function. However, this tool requires basic bioinformatic skills since it runs from a command-line interface. Finally, Haplotracker has been designed for fragmented DNA samples, such as degraded ones, allowing datasets to be classified using both short reads and complete mtDNA sequences. It runs as a web application with a user-friendly interface that makes it appealing for users without bioinformatics skills.

Long-read sequencing technology allows new genome discoveries leveraging the improvements in genome assemblies and detecting structural variants, among others. Long noisy ONT reads have been applied for studies of the nuclear genome (Beyter et al., 2019; Olson et al., 2020). However, there are few examples of the use of this sequencing technology for mtDNA genome analysis (Franco-Sierra & Díaz-Nieto, 2020; Lindberg et al., 2016). The Achilles' heel of this emerging sequencing technology, the high error rates, is continuously improving mostly based on pore modifications and the development of basecalling methods. Here we showed that reference-based assembly and hybrid *de novo* assembly strategies provide precise results for haplogroup classification. However, despite the high number of artefactual variants detected using only the long reads for assemblies (reference-based and *de novo*), these results could be improved using new methods for basecalling and/or genome assembly. Irrespective of that, our results demonstrate that the ONT reads are appropriate for recovering accurate mtDNA haplogroups from WGS data.

5. Conclusions

With the advent of the HTS technologies, the number of human mtDNA haplogroup classification tools has increased notably in the last decade. Each new tool released incorporates novel features and different analysis functions, but this has not been always linked to an improvement in the haplogroup classification accuracy. In this study, an evidence-based benchmarking effort was proposed to compare the classification accuracy provided by the most salient tools. We conclude that Haplocheck is the most suitable mtDNA haplogroup estimator for WGS and WES datasets, not only because of its classification accuracy but also because of all the included features and its user-friendly web interface. Regarding third-generation HTS, despite the lower *per* base accuracy currently offered by ONT, we found that it does not hinder a precise human mtDNA classification.

6. Acknowledgements

We would like to thank the support from our colleagues from the Teide-HPC Supercomputing facility (http://teidehpc.iter.es/en), which was funded by INP-2011-0063-PCT-430000-ACT (INNPLANTA program) from the Spanish Ministry of Economy and Competitiveness.

7. Author Contributions

Conceptualization, VGO, AMB, JMLS, CF; Data curation, VGO, AMB, JMLS, LARR, DJT, RGM; Methodology, VGO, AMB, JMLS, CZT, LARR, ADU, DJT, AIC, RGM; Supervision, CF; Writing—original draft preparation, VGO, AMB, CF; Writing—review and editing, VGO, AMB, JMLS, RGM, CF. All authors have read and agreed to the published version of the manuscript.

8. Conflicts of Interest

The authors declare no potential conflicts of interest with respect to the authorship and/or publication of this article. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

9. Funding

This research was funded by Ministerio de Ciencia e Innovación (RTC-2017-6471-1; AEI/FEDER, UE), co-financed by the European Regional Development Funds 'A way of making Europe' from the European Union; Cabildo Insular de Tenerife (CGIEU0000219140); and by the agreement OA17/008 with Instituto Tecnológico y de Energías Renovables (ITER) to strengthen scientific and technological education, training, research, development and innovation in Genomics, Personalized Medicine and Biotechnology. ADU was supported by a fellowship from the Spanish Ministry of Education and Vocational Training (grant number FPU16/01435).

10. References

Abicht, A., Scharf, F., Kleinle, S., Schön, U., Holinski-Feder, E., Horvath, R., Benet-Pagès, A., & Diebold, I. (2018). Mitochondrial and nuclear disease panel (Mito-aND-Panel): Combined sequencing of mitochondrial and nuclear DNA by a cost-effective and sensitive NGS-based method. *Molecular Genetics & Genomic Medicine*, *6*(6), 1188–1198.

Al-Nakeeb, K., Petersen, T. N., & Sicheritz-Pontén, T. (2017). Norgal: extraction and *de novo* assembly of mitochondrial DNA from whole-genome sequencing data. *BMC Bioinformatics*, *18*(1), 510.

Alonge, M., Soyk, S., Ramakrishnan, S., Wang, X., Goodwin, S., Sedlazeck, F. J., Lippman, Z. B., & Schatz, M. C. (2019). RaGOO: fast and accurate reference-guided scaffolding of draft genomes. *Genome Biology*, *20*(1), 224.

Amorim, A., Fernandes, T., & Taveira, N. (2019). Mitochondrial DNA in human identification: a review. *PeerJ*, *7*, e7314.

Anderson, S., Bankier, A. T., Barrell, B. G., de Bruijn, M. H., Coulson, A. R., Drouin, J., Eperon, I. C., Nierlich, D. P., Roe, B. A., Sanger, F., Schreier, P. H., Smith, A. J., Staden, R., & Young, I. G. (1981). Sequence and organization of the human mitochondrial genome. *Nature*, *290*(5806), 457–465.

Andrews, R. M., Kubacka, I., Chinnery, P. F., Lightowlers, R. N., Turnbull, D. M., & Howell, N. (1999). Reanalysis and revision of the Cambridge reference sequence for human mitochondrial DNA. *Nature Genetics*, *23*(2), 147.

Balter, M. (2011). Was North Africa the launch pad for modern human migrations? *Science*, *331*(6013), 20–23.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software, Articles*, *67*(1), 1–48.

Beyter, D., Ingimundardottir, H., & Eggertsson, H. P. (2019). Long read sequencing of 1,817 Icelanders provides insight into the role of structural variants in human disease. *Biorxiv*. https://www.biorxiv.org/content/10.1101/848366v1.abstract

Børsting, C., & Morling, N. (2015). Next generation sequencing and its applications in forensic genetics. *Forensic Science International. Genetics*, *18*, 78–89.

Brotherton, P., Haak, W., Templeton, J., Brandt, G., Soubrier, J., Jane Adler, C., Richards, S. M., Der Sarkissian, C., Ganslmeier, R., Friederich, S., Dresely, V., van Oven, M., Kenyon, R., Van der Hoek, M. B., Korlach, J., Luong, K., Ho, S. Y. W., Quintana-Murci, L., Behar, D. M., … Genographic Consortium. (2013). Neolithic

mitochondrial haplogroup H genomes and the genetic origins of Europeans. *Nature Communications*, *4*, 1764.

Calabrese, C., Simone, D., Diroma, M. A., Santorsola, M., Guttà, C., Gasparre, G., Picardi, E., Pesole, G., & Attimonelli, M. (2014). MToolBox: a highly automated pipeline for heteroplasmy annotation and prioritization analysis of human mitochondrial variants in high-throughput sequencing. *Bioinformatics* , *30*(21), 3115–3117.

Chan, E. K. F., Timmermann, A., Baldi, B. F., Moore, A. E., Lyons, R. J., Lee, S.-S., Kalsbeek, A. M. F., Petersen, D. C., Rautenbach, H., Förtsch, H. E. A., Bornman, M. S. R., & Hayes, V. M. (2019). Human origins in a southern African palaeo-wetland and first migrations. *Nature*, *575*(7781), 185–189.

Chung, Y., Rabe-Hesketh, S., Dorie, V., Gelman, A., & Liu, J. (2013). A nondegenerate penalized likelihood estimator for variance parameters in multilevel models. *Psychometrika*, *78*(4), 685–709.

Churchill, J. D., Stoljarova, M., King, J. L., & Budowle, B. (2018). Massively parallel sequencing-enabled mixture analysis of mitochondrial DNA samples. *International Journal of Legal Medicine*, *132*(5), 1263–1272.

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics* , *27*(15), 2156–2158.

De Angelis, F., Scorrano, G., Martínez-Labarga, C., Scano, G., Macciardi, F., & Rickards, O. (2018). Mitochondrial variability in the Mediterranean area: a complex stage for human migrations. *Annals of Human Biology*, *45*(1), 5–19.

DeBalsi, K. L., Hoff, K. E., & Copeland, W. C. (2017). Role of the mitochondrial DNA replication machinery in mitochondrial DNA mutagenesis, aging and age-related diseases. *Ageing Research Reviews*, *33*, 89–104.

DePristo, M. A., Banks, E., Poplin, R., Garimella, K. V., Maguire, J. R., Hartl, C., Philippakis, A. A., del Angel, G., Rivas, M. A., Hanna, M., McKenna, A., Fennell, T. J., Kernytsky, A. M., Sivachenko, A. Y., Cibulskis, K., Gabriel, S. B., Altshuler, D., & Daly, M. J. (2011). A framework for variation discovery and genotyping using next-generation DNA sequencing data. *Nature Genetics*, *43*(5), 491–498.

Díaz-de Usera, A., Lorenzo-Salazar, J. M., Rubio-Rodríguez, L. A., Muñoz-Barrera, A., Guillen-Guio, B., Marcelino-Rodríguez, I., García-Olivares, V., Mendoza-Alvarez, A., Corrales, A., Íñigo-Campos, A., González-Montelongo, R., & Flores, C. (2020). Evaluation of Whole-Exome Enrichment Solutions: Lessons from the High-End of the Short-Read Sequencing Scale. *Journal of Clinical Medicine Research*, *9*(11),

3656.

Diroma, M. A., Varvara, A. S., Attimonelli, M., Pesole, G., & Picardi, E. (2020). Investigating Human

Mitochondrial Genomes in Single Cells. *Genes*, *11*(5). https://doi.org/10.3390/genes11050534

Fan, L., & Yao, Y.-G. (2011). MitoTool: a web server for the analysis and retrieval of human mitochondrial DNA

sequence variations. *Mitochondrion*, *11*(2), 351–356.

Fan, L., & Yao, Y.-G. (2013). An update to MitoTool: using a new scoring system for faster mtDNA haplogroup

determination. *Mitochondrion*, *13*(4), 360–363.

Fox, J. (2003). Effect Displays in R for Generalised Linear Models. *Journal of Statistical Software*, *008*(i15).

https://ideas.repec.org/a/jss/jstsof/v008i15.html

Franco-Sierra, N. D., & Díaz-Nieto, J. F. (2020). Rapid mitochondrial genome sequencing based on Oxford

Nanopore Sequencing and a proxy for vertebrate species identification. *Ecology and Evolution*, *10*(7),

3544–3560.

Fu, Q., Mittnik, A., Johnson, P. L. F., Bos, K., Lari, M., Bollongino, R., Sun, C., Giemsch, L., Schmitz, R., Burger, J.,

Ronchitelli, A. M., Martini, F., Cremonesi, R. G., Svoboda, J., Bauer, P., Caramelli, D., Castellano, S., Reich,

D., Pääbo, S., & Krause, J. (2013). A revised timescale for human evolution based on ancient

mitochondrial genomes. *Current Biology: CB*, *23*(7), 553–559.

Gonder, M. K., Mortensen, H. M., Reed, F. A., de Sousa, A., & Tishkoff, S. A. (2007). Whole-mtDNA genome

sequence analysis of ancient African lineages. *Molecular Biology and Evolution*, *24*(3), 757–768.

Griffin, H. R., Pyle, A., Blakely, E. L., Alston, C. L., Duff, J., Hudson, G., Horvath, R., Wilson, I. J., Santibanez-Koref,

M., Taylor, R. W., & Chinnery, P. F. (2014). Accurate mitochondrial DNA sequencing using off-target reads

provides a single test to identify pathogenic point mutations. *Genetics in Medicine: Official Journal of the*

American College of Medical Genetics, *16*(12), 962–971.

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: quality assessment tool for genome

assemblies. *Bioinformatics* , *29*(8), 1072–1075.

Hajdinjak, M., Fu, Q., Hübner, A., Petr, M., Mafessoni, F., Grote, S., Skoglund, P., Narasimham, V., Rougier, H.,

Crevecoeur, I., Semal, P., Soressi, M., Talamo, S., Hublin, J.-J., Gušić, I., Kućan, Ž., Rudan, P., Golovanova,

L. V., Doronichev, V. B., … Kelso, J. (2018). Reconstructing the genetic history of late Neanderthals.

Nature, *555*(7698), 652–656.

Hartig, F. (2017). DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models. *R*

Package Version 0.1.

https://scholar.google.ca/scholar?cluster=4360617861234309959,11567242814940452891,2963838797
944146284,13693552109982088139,10858470353899442396,13508535809485820766,8264605584777
647235,11449340219754401872,4568553816172706093,7031840379694243239,805624554589352998
6,5011821598121647820&hl=en&as_sdt=0,5&sciodt=0,5

Heinze, G., & Schemper, M. (2002). A solution to the problem of separation in logistic regression. *Statistics in Medicine*, *21*(16), 2409–2419.

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal. Biometrische Zeitschrift*, *50*(3), 346–363.

Ishiya, K., & Ueda, S. (2017). MitoSuite: a graphical tool for human mitochondrial genome profiling in massive parallel sequencing. *PeerJ*, *5*, e3406.

Ishiya, K., & Ueda, S. (2019). *Novel approach for accurate detection of contaminating human mitochondrial DNA in next-generation sequencing data - IOPscience*. iopscience.iop.org.

https://iopscience.iop.org/article/10.1088/1742-6596/1391/1/012045/meta

Jagadeesan, A., Ebenesersdóttir, S. S., Guðmundsdóttir, V. B., Thordardottir, E. L., Moore, K. H. S., & Helgason, A. (2020). HaploGrouper: A generalized approach to haplogroup classification. *Bioinformatics* .

https://doi.org/10.1093/bioinformatics/btaa729

Jennings, L. J., Arcila, M. E., Corless, C., Kamel-Reid, S., Lubin, I. M., Pfeifer, J., Temple-Smolkin, R. L., Voelkerding, K. V., & Nikiforova, M. N. (2017). Guidelines for Validation of Next-Generation Sequencing-Based Oncology Panels: A Joint Consensus Recommendation of the Association for Molecular Pathology and College of American Pathologists. *The Journal of Molecular Diagnostics: JMD*, *19*(3), 341–365.

Just, R. S., Leney, M. D., Barritt, S. M., Los, C. W., Smith, B. C., Holland, T. D., & Parsons, T. J. (2009). The use of mitochondrial DNA single nucleotide polymorphisms to assist in the resolution of three challenging forensic cases. *Journal of Forensic Sciences*, *54*(4), 887–891.

Kim, K., Kim, Y., Kim, D.-H., Kwon, C., & Kim, K.-Y. (2020). Haplotracker: a web application for simple and accurate mitochondrial haplogrouping using short DNA fragments. In *bioRxiv* (p. 2020.04.23.057646).

https://doi.org/10.1101/2020.04.23.057646

Lee, H. Y., Song, I., Ha, E., Cho, S.-B., Yang, W. I., & Shin, K.-J. (2008). mtDNAmanager: a Web-based tool for the management and quality analysis of mitochondrial DNA control-region sequences. *BMC Bioinformatics*,

9, 483.

Leger, A., & Leonardi, T. (2019). pycoQC, interactive quality control for Oxford Nanopore Sequencing. *Journal of Open Source Software*, *4*(34), 1236.

Li, H. (2016). Minimap and miniasm: fast mapping and *de novo* assembly for noisy long sequences. *Bioinformatics* , *32*(14), 2103–2110.

Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* , *34*(18), 3094–3100.

Lindberg, M. R., Schmedes, S. E., Hewitt, F. C., Haas, J. L., Ternus, K. L., Kadavy, D. R., & Budowle, B. (2016). A Comparison and Integration of MiSeq and MinION Platforms for Sequencing Single Source and Mixed Mitochondrial Genomes. *PloS One*, *11*(12), e0167600.

Llamas, B., Fehren-Schmitz, L., Valverde, G., Soubrier, J., Mallick, S., Rohland, N., Nordenfelt, S., Valdiosera, C., Richards, S. M., Rohrlach, A., Romero, M. I. B., Espinoza, I. F., Cagigao, E. T., Jiménez, L. W., Makowski, K., Reyna, I. S. L., Lory, J. M., Torrez, J. A. B., Rivera, M. A., … Haak, W. (2016). Ancient mitochondrial DNA provides high-resolution time scale of the peopling of the Americas. *Science Advances*, *2*(4), e1501385.

Maca-Meyer, N., González, A. M., Larruga, J. M., Flores, C., & Cabrera, V. M. (2001). Major genomic mitochondrial lineages delineate early human expansions. *BMC Genetics*, *2*, 13.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303.

Navarro-Gomez, D., Leipzig, J., Shen, L., Lott, M., Stassen, A. P. M., Wallace, D. C., Wiggs, J. L., Falk, M. J., van Oven, M., & Gai, X. (2015). Phy-Mer: a novel alignment-free and reference-independent mitochondrial haplogroup classifier. *Bioinformatics* , *31*(8), 1310–1312.

Olson, N. D., Wagner, J., McDaniel, J., Stephens, S. H., Westreich, S. T., Prasanna, A. G., Johanson, E., Boja, E., Maier, E. J., Serang, O., Jáspez, D., Lorenzo-Salazar, J. M., Muñoz-Barrera, A., Rubio-Rodríguez, L. A., Flores, C., Kyriakidis, K., Malousi, A., Shafin, K., Pesout, T., … Zook, J. M. (2020). precisionFDA Truth Challenge V2: Calling variants from short- and long-reads in difficult-to-map regions. *Biorxiv*. https://doi.org/10.1101/2020.11.13.380741

Parson, W., & Dür, A. (2007). EMPOP—A forensic mtDNA database. *Forensic Science International. Genetics*, *1*(2), 88–92.

Parson, W., Gusmão, L., Hares, D. R., Irwin, J. A., Mayr, W. R., Morling, N., Pokorak, E., Prinz, M., Salas, A.,

Schneider, P. M., Parsons, T. J., & DNA Commission of the International Society for Forensic Genetics. (2014). DNA Commission of the International Society for Forensic Genetics: revised and extended guidelines for mitochondrial DNA typing. *Forensic Science International. Genetics*, *13*, 134–142.

Patowary, A., Nesbitt, R., Archer, M., Bernier, R., & Brkanac, Z. (2017). Next Generation Sequencing Mitochondrial DNA Analysis in Autism Spectrum Disorder. *Autism Research: Official Journal of the International Society for Autism Research*, *10*(8), 1338–1343.

Petrackova, A., Vasinek, M., Sedlarikova, L., Dyskova, T., Schneiderova, P., Novosad, T., Papajik, T., & Kriegova, E. (2019). Standardization of Sequencing Coverage Depth in NGS: Recommendation for Detection of Clonal and Subclonal Mutations in Cancer Diagnostics. *Frontiers in Oncology*, *9*, 851.

Picardi, E., & Pesole, G. (2012). Mitochondrial genomes gleaned from human whole-exome sequencing. *Nature Methods*, *9*(6), 523–524.

Posth, C., Renaud, G., Mittnik, A., Drucker, D. G., Rougier, H., Cupillard, C., Valentin, F., Thevenet, C., Furtwängler, A., Wißing, C., Francken, M., Malina, M., Bolus, M., Lari, M., Gigli, E., Capecchi, G., Crevecoeur, I., Beauval, C., Flas, D., … Krause, J. (2016). Pleistocene Mitochondrial Genomes Suggest a Single Major Dispersal of Non-Africans and a Late Glacial Population Turnover in Europe. *Current Biology: CB*, *26*(6), 827–833.

Puttick, C., Kumar, K. R., Davis, R. L., Pinese, M., Thomas, D. M., Dinger, M. E., Sue, C. M., & Cowley, M. J. (2019). mity: A highly sensitive mitochondrial variant analysis pipeline for whole genome sequencing data. In *Bioinformatics* (No. biorxiv;852210v1). bioRxiv. https://doi.org/10.1101/852210

Pyle, A., Anugrha, H., Kurzawa-Akanbi, M., Yarnall, A., Burn, D., & Hudson, G. (2016). Reduced mitochondrial DNA copy number is a biomarker of Parkinson's disease. *Neurobiology of Aging*, *38*, 216.e7–e216.e10.

Raymond, F. L., Horvath, R., & Chinnery, P. F. (2018). First-line genomic diagnosis of mitochondrial disorders [Review of *First-line genomic diagnosis of mitochondrial disorders*]. *Nature Reviews. Genetics*, *19*(7), 399–400. nature.com.

Robin, E. D., & Wong, R. (1988). Mitochondrial DNA molecules and virtual number of mitochondria per cell in mammalian cells. *Journal of Cellular Physiology*, *136*(3), 507–513.

Röck, A. W., Dür, A., van Oven, M., & Parson, W. (2013). Concept for estimating mitochondrial DNA haplogroups using a maximum likelihood approach (EMMA). *Forensic Science International. Genetics*, *7*(6), 601–609.

Schönberg, A., Theunert, C., Li, M., Stoneking, M., & Nasidze, I. (2011). High-throughput sequencing of complete human mtDNA genomes from the Caucasus and West Asia: high diversity and demographic inferences. *European Journal of Human Genetics: EJHG*, *19*(9), 988–994.

Smieszek, S., Mitchell, S. L., Farber-Eger, E. H., Veatch, O. J., Wheeler, N. R., Goodloe, R. J., Wells, Q. S., Murdock, D. G., & Crawford, D. C. (2018). Hi-MC: a novel method for high-throughput mitochondrial haplogroup classification. *PeerJ*, *6*, e5149.

Sosa, M. X., Sivakumar, I. K. A., Maragh, S., Veeramachaneni, V., Hariharan, R., Parulekar, M., Fredrikson, K. M., Harkins, T. T., Lin, J., Feldman, A. B., Tata, P., Ehret, G. B., & Chakravarti, A. (2012). Next-generation sequencing of human mitochondrial reference genomes uncovers high heteroplasmy frequency. *PLoS Computational Biology*, *8*(10), e1002737.

van Oven, M., & Kayser, M. (2009). Updated comprehensive phylogenetic tree of global human mitochondrial DNA variation. *Human Mutation*, *30*(2), E386–E394.

Vaser, R., Sović, I., Nagarajan, N., & Šikić, M. (2017). Fast and accurate *de novo* genome assembly from long uncorrected reads. *Genome Research*, *27*(5), 737–746.

Vasta, V., Ng, S. B., Turner, E. H., Shendure, J., & Hahn, S. H. (2009). Next generation sequence analysis for mitochondrial disorders. *Genome Medicine*, *1*(10), 100.

Vianello, D., Sevini, F., Castellani, G., Lomartire, L., Capri, M., & Franceschi, C. (2013). HAPLOFIND: a new method for high-throughput mtDNA haplogroup assignment. *Human Mutation*, *34*(9), 1189–1194.

Vohr, S. H., Gordon, R., Eizenga, J. M., Erlich, H. A., Calloway, C. D., & Green, R. E. (2017). A phylogenetic approach for haplotype analysis of sequence data from complex mitochondrial mixtures. *Forensic Science International. Genetics*, *30*, 93–105.

Watson, E., Davis, R., & Sue, C. M. (2020). New diagnostic pathways for mitochondrial disease. *Journal of Translational Genetics and Genomics*. https://doi.org/10.20517/jtgg.2020.31

Weissensteiner, H., Forer, L., Fendt, L., Kheirkhah, A., Salas, A., Kronenberg, F., & Schoenherr, S. (2020). Haplocheck: Phylogeny-based Contamination Detection in Mitochondrial and Whole-Genome Sequencing Studies. In *bioRxiv* (p. 2020.05.06.080952). https://doi.org/10.1101/2020.05.06.080952

Weissensteiner, H., Pacher, D., Kloss-Brandstätter, A., Forer, L., Specht, G., Bandelt, H.-J., Kronenberg, F., Salas, A., & Schönherr, S. (2016). HaploGrep 2: mitochondrial haplogroup classification in the era of high-throughput sequencing. *Nucleic Acids Research*, *44*(W1), W58–W63.

West, A. P., & Shadel, G. S. (2017). Mitochondrial DNA in innate immune responses and inflammatory pathology. *Nature Reviews. Immunology*, *17*(6), 363–375.

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wick, R. R., Judd, L. M., Gorrie, C. L., & Holt, K. E. (2017). Unicycler: Resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Computational Biology*, *13*(6), e1005595.

Wortmann, S. B., Koolen, D. A., Smeitink, J. A., van den Heuvel, L., & Rodenburg, R. J. (2015). Whole exome sequencing of suspected mitochondrial patients in clinical practice. *Journal of Inherited Metabolic Disease*, *38*(3), 437–443.

Yin, C., Liu, Y., Guo, X., Li, D., Fang, W., Yang, J., Zhou, F., Niu, W., Jia, Y., Yang, H., & Xing, J. (2019). An Effective Strategy to Eliminate Inherent Cross-Contamination in mtDNA Next-Generation Sequencing of Multiple Samples. *The Journal of Molecular Diagnostics: JMD*, *21*(4), 593–601.

Supplementary Materials

Figure S1. Bioinformatic pipeline for short-read WES and WGS data processing and variant calling of human mitogenomes.

Figure S2. Bioinformatic pipeline for processing the long-read ONT data for human mitogenome reconstruction, including the four strategies used in the study: *de novo* assembly using long reads (in green), the hybrid *de novo* assembly using short and long reads (in red), the reference-based assembly using long reads (in blue), and the variant-calling strategy using long reads (in purple).

Table S1. Short-read sequencing summaries of the samples.

Samples	Whole-Genome Sequencing							Whole-Exome Sequencing						
	Protocol	Mapped reads	Duplicate reads	Mapping Quality	Base Quality	Depth	Variants	Protocol	Mapped reads	Duplicate reads	Mapping Quality	Base Quality	Depth	Variants
CAN01	Nextera-DNA-Prep	269.488	77.240	59.98	26.00	1227	42	Nextera-DNA-Exome	15.273	2.438	59.96	29.10	58	42
CAN02	Nextera-DNA-Prep	239,511	65,285	59.99	29.20	1466	11	Nextera-DNA-Exome	13,499	1,809	59.97	29.20	53	10
CAN03	Nextera-DNA-Prep	225,038	63,872	59.96	28.90	1293	14	Nextera-DNA-Exome	3,299	950	60.00	28.70	18	6
CAN04	Nextera-DNA-Prep	239,742	56,631	59.95	27.90	1589	10	Nextera-DNA-Exome	16,786	2,296	59.98	29.30	66	9
CAN05	Nextera-DNA-Prep	140,980	27,795	59.98	28.80	974	28	Nextera-DNA-Exome	9,925	1,274	59.98	29.30	39	25
CAN06	Nextera-DNA-Prep	209,422	86,736	59.98	28.60	977	44	Nextera-DNA-Exome	15,609	2,394	59.98	29.40	60	44
CAN07	Nextera-DNA-Prep	202,279	53,246	59.98	29.20	676	14	Nextera-DNA-Exome	17,727	3,247	59.98	29.20	66	14
CAN08	Nextera-DNA-Prep	102,834	22,765	59.97	28.70	661	41	Nextera-DNA-Exome	2,456	702	60.00	29.20	14	8
CAN09	Nextera-DNA-Prep	288,083	92,888	59.99	29.40	880	37	Nextera-DNA-Exome	36,117	15,507	59.99	29.50	93	34
CAN10	Nextera-DNA-Prep	107,424	17,424	59.98	28.80	760	26	Nextera-DNA-Exome	6,501	1,082	59.97	28.70	25	26
CAN11	Nextera-DNA-Prep	486,352	184,988	59.97	28.60	2577	25	Nextera-DNA-Exome	5,452	1,783	60.00	29.10	29	17
CAN12	Illumina-DNA-Prep	96,563	15,042	59.95	28.10	724	25	Nextera-DNA-Exome	5,297	850	59.95	28.70	20	22
CAN13	Nextera-DNA-Prep	298,707	84,386	59.98	29.00	1822	25	Nextera-DNA-Exome	23,503	3,853	59.99	29.30	89	25
CAN14	Nextera-DNA-Prep	160,800	31,404	59.98	28.60	1130	12	Nextera-DNA-Exome	8,746	3,222	60.00	29.40	25	12
CAN15	Illumina-DNA-Prep	86,288	12,342	59.94	27.90	656	34	Nextera-DNA-Exome	6,208	1,168	59.98	29.50	23	31
CAN16	Nextera-DNA-Prep	145,985	52,881	59.95	27.00	786	33	Nextera-DNA-Exome	5,211	1,964	60.00	29.10	23	20
CAN17	Nextera-DNA-Prep	287,742	80,049	59.96	28.40	1691	12	Nextera-DNA-Exome	5,532	1,705	60.00	28.80	30	9
CAN18	Nextera-DNA-Prep	186,339	78,863	59.99	28.90	874	34	Nextera-DNA-Exome	5,056	2,026	60.00	29.30	22	11
CAN19	Nextera-DNA-Prep	158,760	34,517	59.97	28.60	1021	13	Nextera-DNA-Exome	2,345	698	59.98	28.10	13	4
CAN20	Nextera-DNA-Prep	198,085	48,750	59.99	29.20	1284	13	Nextera-DNA-Exome	9,765	1,655	59.98	36.80	37	12
CAN21	Nextera-DNA-Prep	183,878	37,268	59.96	28.30	1282	17	Nextera-DNA-Exome	6,736	965	59.99	36.60	26	17
CAN22	Nextera-DNA-Prep	466,404	196,793	59.98	29.10	1823	7	Nextera-DNA-Exome	3,911	1,212	59.99	29.20	22	1
CAN23	Nextera-DNA-Prep	149,371	32,434	59.99	28.60	1002	35	Nextera-DNA-Exome	9,810	1,422	59.99	29.20	38	35
CAN24	Nextera-DNA-Prep	75,746	10,746	59.97	28.70	554	34	Nextera-DNA-Exome	6,809	1,010	59.95	29.20	26	32
CAN25	Nextera-DNA-Prep	130,389	23,037	59.97	28.40	935	39	Nextera-DNA-Exome	6,707	1,184	59.99	29.30	25	37
CAN26	Illumina-DNA-Prep	145,401	30,564	59.96	28.40	1024	35	Nextera-DNA-Exome	10,105	1,900	59.99	29.50	37	32
CAN27	Illumina-DNA-Prep	106,440	16,750	59.94	28.00	796	39	Nextera-DNA-Exome	6,087	996	60.00	29.40	23	38
CAN28	Nextera-DNA-Prep	341,321	102,609	59.97	25.80	1594	10	Nextera-DNA-Exome	13,280	2,376	59.98	29.20	50	9
CAN29	Nextera-DNA-Prep	351,160	108,741	59.97	25.80	1618	15	Nextera-DNA-Exome	17,020	5,915	59.99	29.60	50	15
CAN30	Nextera-DNA-Prep	189,623	78,268	59.97	27.90	973	38	Nextera-DNA-Exome	4,347	1,745	60.00	29.20	19	11
CAN31	Nextera-DNA-Prep	143,579	32,325	59.98	29.20	950	26	Nextera-DNA-Exome	10,019	2,159	59.93	28.70	36	25
CAN32	Nextera-DNA-Prep	116,947	21,154	59.98	28.60	824	26	Nextera-DNA-Exome	7,947	1,362	59.98	29.10	30	25
CAN33	Nextera-DNA-Prep	138,396	38,670	59.99	29.30	649	31	Nextera-DNA-Exome	4,038	1,204	60.00	29.10	23	17
CAN34	Nextera-DNA-Prep	130,484	23,846	59.98	28.80	913	43	Nextera-DNA-Exome	14,569	2,380	59.98	29.00	55	42
CAN35	Illumina-DNA-Prep	119,570	22,308	59.96	28.50	867	40	Nextera-DNA-Exome	5,709	732	59.97	35.10	23	38
CAN36	Illumina-DNA-Prep	198,697	40,465	59.95	28.30	1402	26	Nextera-DNA-Exome	15,194	2,908	59.99	29.50	56	26

Table S2. Multicollinearity test results (Pearson correlation coefficients) based on cross-correlation among the selected parameters estimated from the BAM files for short-read A) WGS and B) WES datasets. The statistical significance is indicated by asterisks: * $p \leq 0.05$ | ** $p \leq 0.01$ | *** $p \leq 0.001$

a)

R square WGS	Mapped reads	Duplicate reads	Mapping Quality	Base quality	Coverage
Mapped reads	1.000	0.959***	0.190	-0.197	0.877***
Duplicates reads	0.959***	1.000	0.224	-0.135	0.772***
Mapping quality	0.190	0.224	1.000	0.382*	0.018
Base quality	-0.197	-0.135	0.382*	1.000	-0.172
Depth	0.877***	0.772***	0.018	-0.172	1.000

b)

R square WES	Mapped reads	Duplicate reads	Mapping Quality	Base quality	Coverage
Mapped reads	1.000	0.837***	-0.107	-0.040	0.960***
Duplicates reads	0.837***	1.000	0.095	-0.076	0.677***
Mapping quality	-0.107	0.095	1.000	-0.006	-0.113
Base quality	-0.040	-0.076	-0.006	1.000	-0.064
Depth	0.960***	0.677***	-0.113	-0.064	1.000

Table S3. Mitochondrial DNA haplogroup consensus and concordance rate among the 11 tools based on short-read WGS for the samples assessed.

Samples	Haplogroup	Concordance rate
CAN01	J2a2d	100.00
CAN02	H7	100.00
CAN03	H1ao1	100.00
CAN04	H1t	92.31
CAN05	U6b1a	100.00
CAN06	J2a2d	92.31
CAN07	H6a1b2	100.00
CAN08	L3d1b3a	92.31
CAN09	U4c1	69.23
CAN10	U6b1a	92.31
CAN11	U6b1a	92.31
CAN12	U6b1a	100.00
CAN13	U6b1a	100.00
CAN14	H1e1a	92.31
CAN15	U4c1	69.23
CAN16	T1a	92.31
CAN17	H1+16189	69.23
CAN18	K1a4a1	100.00
CAN19	H1e1a	100.00
CAN20	H1au	92.31
CAN21	HV+16311	69.23
CAN22	H	69.23
CAN23	K1a1b1	92.31
CAN24	K1a14	92.31
CAN25	J1c2c1	92.31
CAN26	J1c2	92.31
CAN27	T2c1d+152	69.23
CAN28	H1cf	69.23
CAN29	H6a1a	92.31
CAN30	U5b2b3a	92.31
CAN31	U6b1a1	100.00
CAN32	U6b1a	92.31
CAN33	X3a	92.31
CAN34	L3f1b1a	92.31
CAN35	L3b1a+@16124	69.23
CAN36	U6b1a1	100.00

Table S4. Per sample mtDNA haplogroup classifications for the 11 tools based on short-read A) WGS and B) WES datasets.

a)

Samples	Haplogrep VCF	Haplogrep FASTA	Haplocheck BAM	Haplocheck VCF	Haplotracker FASTA	HaploGrouper VCF	Phy-Mer FASTA	Phy-Mer BAM	EMMA FASTA	MitoSuite BAM	Haplofind FASTA	MixEmt BAM	MitoTool FASTA	James Lick's FASTA
CAN01	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2a	J2a2d	J2a2d
CAN02	H7	H7	H7	H7	H7	H7	H7	H7	H7	H27e	H7	H22	H7	H7
CAN03	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1	H1ao1
CAN04	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t2	H1t	H1t
CAN05	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN06	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2a	J2a2d	J2a2d
CAN07	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2b	H6a1b2	H6a1b2
CAN08	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d1b3a	L3d4a	L3d1b3a	L3d1b3a
CAN09	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1a	U4c1a	U4c1	U4c1	U4c1	U4a2g	U4c1a	U4c1
CAN10	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a1	U6b1a	U6b1a
CAN11	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a1	U6b1a	U6b1a
CAN12	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN13	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN14	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1a4	H1e1a	H1e1a
CAN15	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1a	U4c1a	U4c1	U4c1	U4c1	U4c1a	U4c1a	U4c1
CAN16	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a1b1	T1a	T1a
CAN17	H1+16189	H1+16189	H1+16189	H1+16189	H1+16189	H1+16189	H1-16189	H1-16189	H1+16189	B	H1	B4a1a1ab	H1	H1+16189
CAN18	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1	K1a4a1
CAN19	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a1	H1e1a	H1e1a
CAN20	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H28	H1au	H1au
CAN21	HV+16311	HV+16311	H+16311	HV+16311	HV+16311	HV+16311	HV-16311	HV-16311	HV+16311	B	HV	R0a'b	HV	HV+16311
CAN22	H	H	H	H	H	H77	H	H-16291	H	H27e	H	H3b	H	H
CAN23	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1b	K1a1b1	K1a1b1
CAN24	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a5a	K1a14	K1a14
CAN25	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2h	J1c2c1	J1c2c1
CAN26	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c4c	J1c2	J1c2
CAN27	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d-152	T2c1d-152	T2c1d+152	T2c1d2	T2c1d2	T2c1d[1]	T2c1d	T2c1d+152
CAN28	H1cf	H1cf	H1cf	H1cf	H1cf	H1cf	H1	H1	H1cf	H1cf	H1cf	H3ac	H1	H1cf
CAN29	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a2b1	H6a1a	H6a1a
CAN30	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b3a	U5b2b	U5b2b3a	U5b2b3a
CAN31	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a	U6b1a1	U6b1a1
CAN32	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a1	U6b1a	U6b1a
CAN33	X3a	X3a	X3a	X3a	X3a	X3a	X3a	X3a	X3a	X3a	X3a	U4b1a1a1	X3a	X3a
CAN34	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a
CAN35	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a-@16124	L3b1a-@16124	L3b1a+@16124	L3b1a11	L3b1a	L3b1a3	L3b1a	L3b1a-@16124
CAN36	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1

b)

Samples	Haplogrep VCF	Haplogrep FASTA	Haplocheck BAM	Haplocheck VCF	Haplotracker FASTA	HaploGrouper VCF	Phy-Mer FASTA	Phy-Mer BAM	EMMA FASTA	MitoSuite BAM	Haplofind FASTA	MixEmt BAM	MitoTool FASTA	James Lick's FASTA
CAN01	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d
CAN02	H7	H7	H7	H7	H7	H7	H7	H7	H7	H27e	H7	H7	H7	H7
CAN03	H1ao	H1ao	H1ao1	H1ao	H2a2a1	H2a2a1	H2a2a1	H1x	H2a2a1	H1ao1	H2a2a1	H1ao1	H2a2a1	H2a2a1
CAN04	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t	H1t
CAN05	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN06	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d	J2a2d
CAN07	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2	H6a1b2
CAN08	L3d	L3d	L3d1b3a	L3d	H2a2a1	H2a2a1	H2a2a1	L3d1b1a	H2a2a1	L3d1b3a	H2a2a1	L3d1b3a	H2a2a1	H2a2a1
CAN09	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1a	U4c1a	U4c1	U4c1	U4c1	U4c1	U4c1a	U4c1
CAN10	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN11	U6b1a	U6b1a	U6b1a	U6b1a	U6+16311	U6a'b'd T16311C	U6a'b'd-16311	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6a'b'd	U6a'b'd
CAN12	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN13	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN14	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a	H1e1a
CAN15	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1	U4c1a	U4a2c	U4c1	U4c1	U4c1	U4c1	U4c1a	U4c1
CAN16	T1a	T1a	T1a	T1a	T1a	T1a	T1a	T1a6	T1a	T1a	T1a	T1a	T1a	T1
CAN17	H2	H2	H1+16189	H2	H1a	H1a	H1a	H1a2	H2	B	H1a	H1a	H1a	H2
CAN18	K1a4	K1a4	K1a4a1	K1a4	H2a2	H2a2	H2a2a	K1a4a1	H2a2a	K1a4a1	H2a2a	K1a4a1	H2a2	H2a2a
CAN19	H2a	H2a	H1e1a	H2a	H2a	H2a	H2a	H1e1b1a	H2a	H1e1a	H2a	H1e1a	H2a	H2a
CAN20	H1au	H1au	H1au	H1au	H1au	H1au	H1au	H1-16311	H1au	H1au	H1au	H1au	H1au	H1au
CAN21	HV+16311	HV+16311	H+16311	HV+16311	HV+16311	HV+16311	HV-16311	HV-16311	HV+16311	B	HV	HV	HV	HV+16311
CAN22	H2	H2	H	H2	H2a2a1	H2a2a1	H2a2a1	H30a	H2a2a1	H27e	H2a2a1	H	H2a2a1	H2a2a1
CAN23	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	k1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1	K1a1b1
CAN24	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14	K1a12	K1a14	K1a14	K1a14	K1a14	K1a14	K1a14
CAN25	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1	J1c2c1
CAN26	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2	J1c2
CAN27	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d+152	T2c1d-152	T2c1d-152	T2c1d+152	T2c1d2	T2c1d2	T2c1d2	T2c1d	T2c1d+152
CAN28	H1cf	H1cf	H1cf	H1cf	H1cf	H1cf	H1	H1	H1cf	H1cf	H1cf	H1cf	H1	H1cf
CAN29	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a	H6a1a
CAN30	U5b2b3	U5b2b3	U5b2b3a	U5b2b3	H2a2a1	H2a2a1	H2a2a1	U5b2b3a	H2a2a1	U5b2b3a	H2a2a1	U5b2b3a	H2a2a1	H2a2a1
CAN31	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1
CAN32	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a	U6b1a
CAN33	X	X	X3a	X	X	X	X	X3a	X	X3a	X	X3a	X	X
CAN34	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a	L3f1b1a
CAN35	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a+@16124	L3b1a-@16124	L3b1a-@16124	L3b1a+@16124	L3b1a11	L3b1a	L3b1a	L3b1a	L3b1a+@16124
CAN36	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1	U6b1a1

Table S5. BGLMM model results for WGS and WES datasets. Summary of values obtained with the haplogroup classification tool as fixed factor. Mapped reads, mapping quality, and base quality were introduced as covariates. Discordance was used as the response variable and the sample was used as a random factor. The statistical significance is marked with asterisks: * $p \leq$ 0.05 | ** $p \leq$ 0.01 | *** $p \leq$ 0.001

Application	Tool	Estimate	Std.Error	zvalue	Pr(>|z|)	
WGS	Haplotracker FASTA	-3.16	0.66	-4.81	0.00	***
-	Phy-Mer FASTA	0.31	0.59	0.52	0.60	
-	EMMA FASTA	-2.20	2.41	-0.91	0.36	
-	Haplofind FASTA	0.56	0.57	0.98	0.33	
-	MitoTool FASTA	1.13	0.55	2.05	0.04	*
-	Haplogrep FASTA	-2.18	2.35	-0.93	0.35	
-	JamesLick's FASTA	-2.16	2.32	-0.93	0.35	
-	HaploCheck BAM	-2.20	2.39	-0.92	0.36	
-	MixEmt BAM	3.55	0.66	5.35	0.00	***
-	MitoSuite BAM	0.76	0.56	1.36	0.17	
-	Phy.mer BAM	0.55	0.57	0.96	0.34	
-	HaploCheck VCF	-2.14	2.27	-0.94	0.35	
-	Haplogrep VCF	-2.13	2.25	-0.95	0.34	
-	HaploGrouper VCF	-0.43	0.71	-0.61	0.54	
-	Mapped Reads	0.37	0.30	1.23	0.22	
-	Mapping Quality	0.06	0.32	-1.86	0.06	.
-	Base Quality	-0.09	0.51	0.17	0.87	
WES	Haplotracker FASTA	-6.84	2.82	-2.42	0.02	*
-	Phy-Mer FASTA	0.47	0.48	0.98	0.33	
-	EMMA FASTA	-0.36	0.51	-0.70	0.48	
-	Haplofind FASTA	0.22	0.49	0.45	0.66	
-	MitoTool FASTA	0.97	0.49	1.98	0.05	*
-	Haplogrep FASTA	-0.55	0.52	-1.07	0.29	
-	JamesLick's FASTA	0.14	0.49	0.28	0.78	
-	HaploCheck BAM	-4.38	1.99	-2.20	0.03	*
-	MixEmt BAM	-1.13	0.50	-2.25	0.02	*
-	MitoSuite BAM	-0.63	0.48	-1.32	0.19	
-	Phy.mer BAM	0.27	0.45	0.60	0.55	
-	HaploCheck VCF	-0.59	0.52	-1.13	0.26	
-	Haplogrep VCF	-0.57	0.52	-1.09	0.27	
-	HaploGoruper VCF	-0.13	0.50	-0.26	0.80	
-	Mapped Reads	-5.96	3.55	-1.68	0.09	.
-	Mapping Quality	1.16	0.39	2.97	0.00	**
-	Base Quality	0.15	0.24	0.61	0.54	