Alternative female and male developmental trajectories in the dynamic balance of human visual perception

Gergő Ziman[*1,2,3], Stepan Aleshin[4], Zsolt Unoka[5], Jochen Braun[6], and Ilona Kovács[1,2,7]

[1]*Laboratory for Psychological Research, Pázmány Péter Catholic University, 1088 Budapest, Hungary*
[2]*Adolescent Development Research Group, Hungarian Academy of Sciences - Pázmány Péter Catholic University, 1088 Budapest, Hungary*
[3]*Doctoral School of Mental Health Sciences, Semmelweis University, 1083 Budapest, Hungary*
[4]*Center for Behavioral Brain Sciences, Otto-von-Guericke-Universität, 39120 Magdeburg, Germany*
[5]*Department of Psychiatry and Psychotherapy, Semmelweis University, 1085 Budapest, Hungary*
[6]*Cognitive Biology Group, Institute of Biology, Otto-von-Guericke Universität, 39120 Magdeburg, Germany*
[7]*Institute of Cognitive Neuroscience and Psychology, Research Centre for Natural Sciences, 1117 Budapest, Hungary*
[*]*e-mail: ziman.gergo@btk.ppke.hu*

June 2021

Abstract

The numerous multistable phenomena in vision, hearing and touch attest that the inner workings of perception are prone to instability. We investigated a visual example – binocular rivalry – with an accurate no-report paradigm, and uncovered developmental and maturational lifespan trajectories that were specific for age and sex. To interpret these trajectories, we hypothesized that conflicting objectives of visual perception – such as *stability* of appearance, *sensitivity* to visual detail, and *exploration* of fundamental alternatives – change in relative importance over the lifespan. Computational modelling of our empirical results allowed us to estimate this putative development of stability, sensitivity, and exploration over the lifespan. Our results confirmed prior findings of developmental psychology and appear to quantify important aspects of neurocognitive phenotype. Additionally, we report atypical function of binocular rivalry in autism spectrum disorder and borderline personality disorder. Our computational approach offers new ways of quantifying neurocognitive phenotypes both in development and in dysfunction.

1 Introduction

An 80 year old grandmother and her 8 year old grandson will react very differently to the same stimulus, e.g. a loud truck passing the street outside the house. The child will be engaged by the sound, turn his body toward the sound source, perhaps even drop his toys and rush to the window for a better look. Grandma, on the other hand, may not even lift her eyes from the page she was reading. While the child cannot help being over-sensitive to novel stimuli, even when it is occupied elsewhere, the grandmother - given her extensive past sensory experience - likely remains indifferent. This scenario illustrates the main hypothesis of this paper: adaptive human functioning balances conflicting objectives, and this balance changes throughout life.

Human neurocognitive development is a multidimensional and ever-changing process determined both by biological mechanisms and the environment over the lifespan. Details on the protracted maturation of human structural brain connectivity are emerging with the advance of brain imaging technology [1–5]. It is becoming increasingly clear from such findings that lifespan trajectories of brain organization improve our current, fragmented view of human development. Nonetheless, little is known about the trajectories of active neurocognitive adaptation to the environment, in other words, about the lifetime development of our own behavioural phenotype.

Because of the aforementioned complexity of human growth, the description of the neurotypical behavioural phenotype ought to be complex as well. It does not seem meaningful to ask what the typical brain structure of an adult human is. One needs to specify at least age and sex, since the brain is continually reorganizing, with its development extending into adulthood [6, 7] and as its structure shows differences between the sexes throughout development [8–10]. Adolescence, in particular, is a period of substantial changes in brain structure – notably, in the association cortex, connectivity is remodelled [11], while cortical shrinkage and myelination are accelerated [12,13]. These organizational changes are coupled with a high vulnerability to mental health disorders [14,15]. Parallel to the changes in the brain's structural organization that take place across the lifespan, we should expect changes in the way it adapts to the environment. Further differences should be expected in the case of atypical development underlying mental health disorders, whether resulting from developmental conditions, environmental factors, or a combination of both.

The aim of the present work is to characterize lifespan trajectories for visual perception. Sensory perception is generally a promising target for such an investigation, because there is considerable flexibility in how sensory perception balances its several conflicting objectives. Accordingly, over the course of development and maturation, sensory dynamics may well have to change in order to accommodate different behavioural strategies or increasing sensory experience.

That successful interaction with volatile and unpredictable environments necessitates trades between conflicting objectives is a central insight from reinforcement learning [16, 17]. For example, relying on past experience in order to make choices in perception or action typically creates an "exploration-exploitation-dilemma" [17–19]. Here, the benefit of safe choices capitalizing on past experience (exploitation) must be weighed against the potential benefit of risky choices flouting precedent (exploration) in order to extend or update this experience. Further tensions arise when a "cost of time" creates an urgency

for reaching timely decisions [20–22]. For example, visual perception generates transiently stable interpretations of continuous sensory input and updates these interpretations at short intervals. Here, the benefit of remaining committed to an interpretation until analysis is complete must be weighed against the benefit of promptly overturning this interpretation in response to ongoing changes in the input. This tension reflects the competing objectives of perceptual stability and sensitivity [22, 23].

To study the inherent dynamics of sensory perception – as well as the trade-offs between conflicting objectives that this may entail – a suitable starting point is "multistable perception", where perceived appearance perpetually changes at irregular intervals, shifting abruptly between distinct alternatives [24]. Presumably, this intrinsic instability reflects a spontaneous reassessment of perceptual decisions, even when the sensory scene remains unchanged [25, 26]. Numerous properties of multistable perception support this notion; notably, the ecological plausibility of alternative appearances and their dependence on behavioural context and prior experience [27]. Interestingly, the dynamics of multistable perception differs substantially between individuals of different age [28–33], and also in individuals with psychiatric disorders, such as autism spectrum disorder [34–38] among many others [39–45]).

To quantify the dynamics of multistable perception in cohorts of different age and psychiatric health, we tracked spontaneous reversals of binocular rivalry that were induced by presenting different images to each eye (Fig. 1). Importantly, we reliably and accurately inferred spontaneous reversals of subjective (covert) eye dominance from objective (overt) eye movements, thereby avoiding the confounds and biases of volitional reports [46]. Briefly, participants dichoptically viewed two horizontally moving gratings through a mirror stereoscope, while we recorded optokinetic nystagmus (OKN) in one eye. The slow phases (smooth pursuit) of the OKN pattern revealed which direction of movement was consciously perceived at the time [47–49].

With this approach, we quantified and characterized binocular rivalry dynamics in a large group of neurotypical participants (N=107) of different ages, from 12-year-old children to senior adults. This broad range of ages was intended to capture both developmental changes during adolescence and maturational changes through adulthood, into senior age. Additionally, we studied two psychiatric groups associated with developmental, genetic and environmental risk factors: Autism spectrum disorder (ASD; all males), which is diagnosed more often in males [50], and where patients are known to present an atypical dynamics of multistable perception [34–38]; and borderline personality disorder (BPD; all females), which is diagnosed predominantly in female adolescents and adults [50, 51]. While multistable perception has not been studied in this latter group, patients are prone to emotion-related misattributions in the perception of emotionally ambiguous faces (i.e., for faces showing no, or neutral emotion [52, 53]).

After establishing three observable variables of binocular rivalry statistics, we were surprised to find that females and males follow quite different developmental trajectories, with particularly pronounced differences during adolescence and menopause. The maturational peak was observed at 19 years for females and 24 years for males, consistent with the earlier puberty onset time seen in girls compared to boys [54]. The intriguing divergence between female and male trajectories during adolescence and menopause raises the possibility

Figure 1: **Binocular rivalry paradigm.** Participants sat in a mirror stereoscope and viewed two separate displays with their left and right eyes. Displays showed gratings of different colour (red and green) and opposite motion, inducing occasional reversals of phenomenal appearance (binocular rivalry). Perceived appearance and its reversals were monitored objectively by recording optokinetic nystagmus of one eye (OKN), using an automated analysis [46].

that the fine-tuning of perceptual function is correlated with sex hormone levels throughout the lifetime. The binocular rivalry statistics of both psychiatric patient groups fell well outside these typical maturational trajectories.

From the observed rivalry statistics, we inferred the changing balance between conflicting objectives, by fitting a computational model of rivalry dynamics [55]. Modelling the rivalry dynamics of each age or patient group allowed us to conduct extensive simulations and to predict perceptual performance in volatile and unpredictable environments. This extrapolated perceptual performance could then be quantified in terms of the conflicting objectives of "stability", "sensitivity", and disposition for "exploration".

Inferred trajectories of performance objectives again differed substantially between females and males. During adolescence, females seemed to gain on all three objectives, reaching a "sweet spot" of high stability, sensitivity, and exploration in their early twenties, where they remained until menopause. In contrast, males attained their highest levels of sensitivity and exploration in early adolescence, subsequently retreating slightly from this peak. Concomitantly, their perceptual stability increased until the mid-twenties and gradually declined thereafter. Psychiatric patient groups differed substantially from age-matched controls, raising the possibility that atypical development due to developmental, genetic, or environmental risk factors alters the functional optimization of perceptual decisions.

2 Results

2.1 Females and males show different developmental trajectories of binocular rivalry dominance statistics

During the experiment, visual stimulation was dichoptic and consisted of two horizontally moving gratings, differing in direction of motion and in colour contrast (red-and-black or green-and-black). This stimulation induced optokinetic nystagmus (OKN) in the direction the grating that currently dominates perception, so that (covert) reversals of subjective dominance (as well as transitional periods) could be inferred accurately from (overt) eye movements [46,48,56]. As this method does not require volitional reports from observers, it is well suited for developmental and patient groups. Participants viewed the dichoptic stimulus for ten trials of 95 seconds duration. The neurotypical participants included 21 developing children (12 twelve-year-olds, 19 sixteen-year-olds) and 52 adults (aged 18-69) (see Methods for details).

We established the statistics of perceptual dominance periods from the recorded eye movements with an automated OKN analysis [46]. Fig. 2 a, b shows the median duration, interquartile range (IQR), and medcouple (MC, a robust measure of skewness) of dominance period distributions, separately for female and male observers, computed with a log-normal weighted sliding average. The resulting trajectories show considerable difference between sexes, e.g. in terms of the starting and turning points, end points, or in terms of the curved segments of the trajectories. It is noticeable that the trajectory of males changes during adolescence, whereas that of females changes between the ages of 40 and 50 years (i.e., during menopause). In participants of the other sex, no such directional change is apparent. Remarkably, the trajectories appear nearly closed, i.e. the values observed at senior age are similar to those observed in children. This may in part be owed to the fact that both younger and older observers exhibited more diverse dominance statistics than middle age observers.

2.2 Females peak earlier in maturation index derived from dominance statistics

To summarize the development of distribution parameters, we defined a maturation index (MI) by performing a principal component analysis on the z-scores of all distribution parameters (from both sexes). As shown in Fig.2 d, the first principal component accounted 79% of the variance. Thus, the projection of distribution parameter triples onto this component summarizes the overall development and was used as a maturational index. In terms of this MI, females peak at an earlier age (age 18.9) than males (age 23.7). The projection of parameter triplets on the second and third PCs accounted for 16% and 3% of the variance, respectively (Fig. 2 e, f).

2.3 Reinterpretation in terms of a computational model of reversal dynamics

To better interpret the results described above, we reproduced the particular dominance statistics of each group of participants with a computational model of binocular rivalry simulating a dynamic interaction of competition, adaptation,

Figure 2: **Developmental and maturational trajectory of distribution parameters.** a, b: Parameters of distribution of dominance durations – median duration (median), interquartile range (IQR), and medcouple (MC) – during development and maturation of neurotypical female and male observers (blue to yellow colour scale). Red stars represent peak age of maturation index (d). The grey mesh represents mean values ± 50% SEM, computed with sliding log-normal weighting and repeated sampling with replacement. The developmental trajectories of the two sexes differ notably within this space. c: Principal component (PC) axes of mean distribution parameters observed over all ages and sexes (computed after conversion to z-score values). First (red), second (blue) and third (green) components account for 79%, 16%, and 3% of the variance, respectively. A maturational index (MI) is obtained by projecting mean parameter triplets onto the first principal component axis. d: Maturation index (MI) of neurotypical female and male observers (magenta and green, respectively), as a function of age. At age 18.9 and 23.7, peak values of 2.11 and 2.42 are reached by female and male observers, respectively (stars). Sliding window average (solid curves) and confidence intervals (±100% SEM, dotted curves) in harmonized units (z-score values). Individual observers are also indicated (circles). e, f: Development and maturation along second (e) and third (f) principal component axes, in harmonized units (z-score values). Note far smaller variance with age compared to (d).

and noise [23, 55] (Fig 3 a, b). The four most consequential parameters of this model were strength of competition (β), adaptation strength (ϕ_a), noise (σ_n) (all dimensionless) and the time-constant of adaptation (τ_a). Simulated rivalry reversals were generated with four levels of competition strength ($\beta = 1$, $\beta = 2$, $\beta = 3$, and $\beta = 4$). All values of β produced qualitatively similar results and illustrations show the results for $\beta = 3$. Input strength was fixed at $I_{1,2} = 1$. Other, less consequential parameters were fixed as well (time-constants of activity τ_n and noise τ_r, inflection point of activation function k; see Methods). We systematically varied ϕ_a, τ_a, and σ_n in order to determine the volume in the ϕ_a-τ_a-σ_n subspace in which the particular dominance distribution parameters of a given group of observers were reproduced within a tolerance of $\pm 5\%$ tolerance (see Methods for details).

The resulting volumes of model parameters in ϕ_a-τ_a-σ_n subspace are shown in Fig. 3 c, d for neurotypical female and male participants. Both females and males show a similar trajectory in terms of adaptation time constant: an increase in development (from childhood to young adulthood), and a decrease during maturation (from young adulthood to senior age). Distinctly, females develop toward a maturational peak with larger values of adaptation strength, and smaller values of noise, while males develop towards a maturational peak with smaller values of adaptation strength, and larger values of noise. This reflects a tight trade-off between adaptation and noise in the dynamics of multistable perception - strong adaptation necessitates weak noise, whereas weak adaptation necessitates strong noise. After the maturational peak, there is a decrease in adaptation time constant for both females and males. The female trajectory from young adulthood to senior age also exhibits a second change of direction, with an increase in noise. There is no similar change of direction seen in the male trajectory. However, the male trajectory exhibits an early change of direction towards more noise in early adolescence. While male parameter trajectories appear less closed (i.e., the centroid values at childhood and old age are more distant, Fig. 3 d), the ambiguity created by the trade-off between adaptation and noise is quite large at low values of τ_a (i.e., the confidence range represented by the grey mesh is rather large).

2.4 Trajectories of perceptual objectives predicted for volatile environments

A second goal of this study was to characterize the perceptual performance of female and male observers of different ages in volatile and unpredictable sensory situations, in order to understand how these groups balance conflicting perceptual objectives such as stability, sensitivity or exploration. Unfortunately, it is not feasible to empirically establish a correlation between stochastic sensory stimuli and stochastic perceptual choice responses, as there would be far too many joint possibilities to consider. However, given a dynamic model for the perceptual choice response, responses to stochastic sensory stimuli can be predicted from exhaustive simulations (approximately 30 hours of simulated viewing time). Of course, the accuracy of such predictions is predicated on the validity of the model.

To predict perceptual performance of observers in a volatile environment, we simulated approximately 10^6 perceptual reversals for any given dynamic model (i.e., any particular set of model parameters), in response to stochastically time-

Dynamic model

Model parameters

Figure 3: **Development and maturation of fitted model parameters.** a: Dynamic model of binocular rivalry, with competition, adaptation, and noise. Two representations (r_1, r_2) are driven by associated visual inputs (I_1, I_2) and independent noise (n_1, n_2). Each representation is inhibited by the other, as well as by the associated adaptive state (a_1, a_2). Free model parameters are competition strength (β), adaptation strength (ϕ_a), adaptation time-scale (τ_a), and noise amplitude (σ_n). Input strength is fixed at $I_1 = I_2 = 1$. b: Representative example of model dynamics with abrupt dominance reversals of activities r_1 and r_2, gradual build-up and recovery of adaptive states a_1 and a_2 (middle), and noise n_1 and n_2 added to visual input $I = 1$ ($\beta = 2$, $\phi_a = 0.7$, $\tau_a = 0.3$, $\sigma_n = 0.2$). Reversals may be triggered by differential adaptive state $a_1 - a_2$, differential noise $n_1 - n_2$, or both. c, d: Parameter triplets ϕ_a-τ_a-σ_n fitted to reproduce (within a $\pm 5\%$) experimentally observed mean distribution parameters. Competition strength was fixed at $\beta = 3$. Centroid values (coloured dots) and confidence range ($\pm$ standard deviation, grey mesh). Red stars represent peak age of maturation index (Fig. 2 d).

varying inputs I_1 and I_2. Specifically, while mean input $(I_1 + I_2)/2$ remained constant, input bias $\Delta I = I_1 - I_2$ varied as a continuously stochastic process (see Methods for details). This allowed us to assess the relative influence on perceptual reversals of the sensory environment (external state I_1, I_2), on the one hand, and of the internal perceptual dynamic (internal state variables r_1, r_2 and a_1, a_2), on the other hand. At any given moment, the model responded to these external and internal factors either with 'perceived stability' (continued dominance) or 'perceived novelty' (reversal of dominance).

Further details of these *in silico* experiments are illustrated in Fig. 4 a. Sensory input bias $\Delta I = I_1 - I_2$ varied as a continuously stochastic process with constant mean, variance, and autocorrelation (see Methods). The superposition of sensory input I_1, I_2 and intrinsic noise n_1, n_2 are also shown. The internal state variables of levels of activity (r_1, r_2) and levels of adaptation (a_1, a_2) respond dynamically and stochastically. Activity levels r_1 and r_2 assume mostly categorical values (near zero or near unity), with intermediate values occurring only transiently during reversals (defined as crossover points with $r_1 = r_2$). In contrast, adaptation levels a_1 and a_2 change gradually over time and their combined state may be summarized conveniently in terms of adaptation bias $\Delta a = a_1 - a_2$. Perceptual performance in this stochastic environment is encapsulated in the joint statistics of input bias ΔI, adaptation bias Δa, and model response (reversal or no reversal).

In order to focus the analysis on the period of initiation or non-initiation of reversals, we disregarded the largely stereotypical time course of reversals themselves. Specifically, we excluded transition periods (20 ms before and after each crossover point) and analyzed both initiation periods (defined as 40 ms to 21 ms before a crossover) and non-initiation periods (all other times) at sampling intervals of 1 ms. The analysis provided a large ensemble of value pairs of ΔI and Δa from initiation periods and an even larger ensemble of value pairs from non-initiation periods. To characterize the stochastic dependence of reversal initiation on external state ΔI and internal state Δa, we computed the conditional probability $P_i nit(\Delta I, \Delta a)$ of reversals as a function of ΔI and Δa. The accuracy of this probability surface was highest in the vicinity of the median state ΔI_m, Δa_m (Fig. 4 b; see Methods for details).

The results showed that the logarithm of initiation probability $\log P_{init}$ varied almost linearly with state variables ΔI and Δa (linear regression $r^2 > 0.99$) in the vicinity of the median state (ΔI_m, Δa_m). As a planar surface, the function $\log P_{init}(\Delta I, \Delta a)$ had three degrees of freedom, which corresponded to three different perceptual objectives: sensitivity, exploration, and stability. (Fig 4 b, c). Specifically, sensitivity represented the maximal gradient of $\log P_{init}$ with respect to state variables ΔI and Δa. Mathematically, it was defined as the length of the gradient vector $\partial \log P_{init}(\Delta I, \Delta a)$ at the median state (or, equivalently, as slope angle β). Similarly, exploration represented the relative influence of internal state Δa, compared to external state ΔI, and was defined mathematically as the direction of the gradient vector $\partial \log P_{init}(\Delta I, \Delta a)$ at the median state (or, equivalently, as angle α). Finally, stability represented the stabilizing or destabilizing effect of the median state and was defined mathematically as the difference between $\log P_{init}$ levels at the median state (ΔI_m, Δa_m) and at the neutral state ($\Delta I = 0$, $\Delta a = 0$), at which the system experiences neither external nor internal biases. For positive values of stability, the median stabilizes the percept and lowers reversal probability. For negative

values of stability, the median state destabilizes the percept and raises reversal probability (see Methods).

In this way, every set of model parameters β-ϕ_a-τ_a-σ_n that reproduced the reversal statistics of a particular group of participants could be mapped onto a corresponding value triplet stability-sensitivity-exploration of perceptual objectives. The resulting trajectories of stability, sensitivity, and exploration associated with different age groups are shown in Fig. 4 d, e. Once again, there are notable differences in the developmental and maturational trajectories of the sexes.

Females gained higher values of stability, sensitivity, and exploration during development, reaching their peak values at young adulthood, with values subsequently decreasing during maturation. Males, on the other hand, exhibited peak values of sensitivity and exploration during adolescence (age 13.5). They subsequently gained stability, peaking at young adulthood, but at the same time lost some sensitivity and exploration. These results hint at different developmental strategies between sexes, in that females gained steadily on all three objectives – stability, sensitivity, and exploration – during adolescence until young adulthood, while adolescent males initially develop sensitivity and exploration at the expense of stability, only to subsequently gain stability at the expense of sensitivity and exploration until young adulthood.

2.5 Clinical populations fall outside of typical developmental and maturational trajectories

In addition to neurotypical observers of various ages, we performed the same set of procedures - binocular rivalry experiment, model fitting, and volatile-environment simulations - with two adult clinical groups: 12 females with borderline personality disorder (BPD; mean age 27.1), and 12 males with autism spectrum disorder (ASD; mean age 28.5). The observed binocular rivalry dominance distribution parameters, and predicted perceptual parameters are shown in Fig 5, contrasted with the corresponding sex's typical developmental and maturational trajectories (results of the model fitting are shown in Supplementary Fig. 1).

The results show that both atypical groups fall outside the typical developmental and maturational trajectories. Despite the different character of their respective disorders, they differ from the typical trajectories in similar ways. In terms of observable dominance distribution parameters (Fig. 5 a, b), both groups show lower median and IQR, and higher MC values. In the predicted perceptual performance in a volatile environment (Fig. 5 c, d), both groups have higher values of sensitivity and exploration, compared to neurotypicals.

3 Discussion

The main purpose of our study was to establish lifespan trajectories of visual perceptual behaviour. With respect to neurotypical brain development and maturation, our results indicate that visual decision making, at least insofar as this is reflected in multistable perception, changes markedly in adolescence, and then more gradually across the lifespan. The fact that the lifetime trajectories we found were sex-specific, suggests that biological maturation plays a major

Simulation

Perceptual predictions

Figure 4: **Predicted perceptual behaviour in a volatile environment.** a: Representative example of simulated dynamics of fitted model: activities r_1, r_2 and adaptive states a_1, a_2 driven by intrinsic noise n_1, n_2 and variable sensory inputs I_1 and I_2. In a volatile environment, reversals (defined by $r_1 = r_2$) may be triggered externally (differential input $\Delta I = I_1 - I_2$) or internally (differential adaptive state $\Delta a = a1 - a2$) or both. To assess reversal initiation, we distinguished between transition periods (yellow stripes, 20ms before and after $r_1 = r_2$), immediately preceding periods (purple stripes, 40 to 21 ms before $r_1 = r_2$), and all other times. b: Based on this classification, we computed the conditional likelihood of reversals (coloured contours) as a function of ΔI and Δa values during initiation periods, in the vicinity of the median ΔI and Δa values (over all periods, red dot). In this vicinity, the reversal probability grows exponentially in a particular direction (red dashed line). The length of the gradient vector $\partial \log P(\Delta I, \Delta a)$ represents the 'sensitivity' of reversal probability to ΔI and/or Δa values, and the 'exploration' angle α represents the relative influence of internal state Δa, compared to external state ΔI (external state). A larger value implies that the system responds less consistently to external state, behaving in a more explorative manner. c: Enlarged cut through the $\log P(\Delta I, \Delta a)$ surface in the direction of the the gradient vector (red arrow), showing the neutral level (defined by $\Delta I = \Delta a = 0$, black dashed line) and median level (red dashed line). The distance between levels (blue arrow), represents 'stability' and measures the stabilizing or destabilizing effect of external and internal median states. For positive values (within green area), median states lower reversal probability, stabilizing the percept; vice versa for negative values (within grey area). d, e,: Developmental and maturational trajectories of predicted perceptual parameters (stability, sensitivity, exploration) for females (d) and males (e). Mean values (coloured dots), peak age of maturation index (red stars), and confidence intervals (standard deviations, grey volumes).

Figure 5: **Observed distribution parameters and predicted perceptual behaviour of atypical groups.** Observed distribution parameters (a, b), and predicted perceptual behaviour in volatile environments (c, d) for atypical groups, compared to neurotypical trajectories. Observers with borderline personality disorder (BPD, all female) are compared to neurotypical females of different ages. Observers with autism spectrum disorder (ASD, all male) are compared to neurotypical males of different ages. (a, b) Ellipsoids represent mean SEM values for BPD (purple) and for ASD (in orange). (c, d) Coloured volumes represent confidence intervals (standard deviation) for BPD (purple) and for ASD (in orange). The dominance period statistics and perceptual predictions of both groups fall outside the typical developmental and maturational trajectory.

role in visual decision making. Neuroendocrine changes, especially levels of sex steroid hormones, are known to pace bodily growth, emotional development, and cortical pruning in adolescence [5,57,58]. Neuroendocrine changes also seem to influence cognitive function during [59,60] and after the teenage years [61,62]. The fact that the maturation index in our study peaks earlier in females (around 19 years of age) than in males (around 24 years of age) is consistent with the relative onset times of puberty in girls and boys [54]. The comparative lateness of these maturational peaks may also be significant. For example, it may be that visual decision making is a neotenous function in humans, which involves the latest maturing brain areas, such as prefrontal cortex and precuneus [63,64]. Looking further into maturation, after the apex in young adulthood, males exhibit an uninterrupted linear change, perhaps reflecting the progressive decline of sex hormones in males [65]. In contrast, females exhibit a second abrupt turn in the direction of change around the time of menopause. This second turn might be the consequence of the sharp gonadal hormone falloff later in life in females [66].

By themselves, these results are not easily interpreted and the implications for visual decision making, if any, are not immediately evident. To generate an interpretive hypothesis, we sought to predict the perceptual behaviour of observers in volatile and unpredictable sensory situations by means of extensive simulations (approximately 30 hours of simulated viewing time). Such extensive data would have been difficult to obtain with human observers. As a first step, we fitted a stochastic dynamic model to reproduce the reversal statistics (i.e., median, IQR and medcouple of dominance time distributions) of a particular group of human observers (e.g., females aged 25 years). In nutshell, this model allowed tension (or mismatch) between external input and internal state to build up, but also occasionally relieved this tension by executing a perceptual reversal. As a second step, we characterized the perceptual behaviour of this model when image contrast varied unpredictably over time. This simulated behaviour was then summarized quantitatively by three measures, as described below.

The first measure, *stability*, captured the influence of the median model state. Note that this influence may be stabilizing or destabilizing, depending on how 'loosely' or 'tightly' the model interacts with the fluctuating input. The second measure, *sensitivity*, reflects how much reversal probability changes when model state is perturbed in the vicinity of the median state. The third measure, *exploration*, describes the relative influence of internal tension (differential adaptation) compared to external tension (differential input).

Although defined in the context of a stochastic dynamic model, these three measures are broadly analogous to competing objectives of perception prescribed by the framework of reinforcement learning. When objects can appear or disappear at any time, the thoroughness of object classifications must be balanced against the frequency of such classifications [20, 21]. In our context, this corresponds to the tension between *stability* (temporal persistence of current choice) and *sensitivity* (susceptibility to changed circumstances, internal or external) [22, 23]. Additionally, in an unstable world, the value of past experience diminishes with time, so that the benefit of past experience must be weighed against the benefit of discovering something new (exploitation-exploration dilemma [17–19]). In our context, this corresponds to the influence of external states (input bias), relative to internal states (adaptation bias) that prompt a fundamental reassessment of such evidence. Hence the relative weight

of adaptation bias represents a tendency for *exploration*.

In terms of behavioural measures predicted by our model, females are characterised by increasing values of stability, sensitivity, and exploration during adolescent development. Peak values of all three measures are reached by twenty, defining a sweet-spot of optimal function for perceptual decisions. After the peak, a moderate decline can be observed until menopause, which is similar in all three parameters. Signs of a stability-sensitivity trade-off can be observed after menopause when stability declines sharply, with slight increases in sensitivity and exploration, following the male pattern in this age-group. Males, on the other hand, have peak values of sensitivity and exploration by the age of sixteen, exceeding the highest female levels. The heightened sensitivity is coupled with low levels of stability, demonstrating a clear trade-off between these measures, at least at the extreme end of the range. Until the mid-twenties, stability rises at the expense of sensitivity and exploration. The male sweet-spot of optimal function seems to be characterised by lower sensitivity and exploration levels, but a higher level of stability, when compared to females. After the peak, males decline progressively in stability. This intriguing and complex pattern in the lifespan evolution of the two sexes suggests, on one hand, that the rules of growth are not uniform even in the developmental period, and on the other hand, that while parameter values in senior age are similar to those in childhood, ageing is not simply the "reverse" of development, as it has been suggested with respect to other cognitive functions as well [67].

Given the biological determination reflected in the age- and sex-specific differences, we interpret this pattern of findings as a result of alternative developmental strategies. It seems that females approach the peak in all three parameters faster, and although this provides them with lower peaks as compared to males, it also seems to provide for a greater stability throughout the childbearing years, with a major decline only after menopause. The trade-off between stability and sensitivity is more obvious in males who start with maximum levels of sensitivity and exploration mid-adolescence, and in parallel to stability building up by the mid-twenties, a great deal of sensitivity is lost. The greater exploration range in the early years, the higher apex, and the continuous decline after the apex might indicate a developmental strategy to fine-tune the individual for the age of highest fertility in males [68]. These alternative developmental strategies suggested by our simulated experiments - one focusing on stability throughout the childbearing period, and the other, focusing on highest performance by the peak fertility age - are particularly interesting in the context of perceptual decisions. Decisions made between alternatives, even if not volitional or highly conscious, make up our everyday life, and the particular style with which we are dealing with those decisions may have crucial effects on our lives. It makes a difference, for example, whether our brain is tuned to be relatively insensitive to environmental changes, rendering decisions stable and stereotypical and preventing extensive exploration or, in another scenario, whether higher sensitivity is combined with a heightened inner drive for exploration.

After obtaining developmental and maturational lifetime trajectories of neurotypical subjects, it is of particular interest how these would be drawn when biological conditions or environmental factors are not adequate for typical development. To this end, we have tested two psychiatric groups: adult participants with autism spectrum disorder (ASD, all males) and with borderline personality disorder (BPD, all females). Both the experimentally observed distribution pa-

rameters and the predicted behavioural measures fell completely outside of the typical developmental trajectories for their respective sex – i.e., adults with these disorders differed not only from neurotypical adults of a similar age, but from the measures of any examined ages, probably missing the sweet-spot of optimal function at an earlier age. In terms of the trade-off in predicted behavioural measures, both patient groups demonstrate sensitivity and exploration levels beyond the range of neurotypicals, while stability is reduced markedly only in males, resembling the pattern seen in neurotypicals. As all psychiatric patients in the study were adults, we cannot pinpoint the when-and-how of the deviations from typical trajectories in ASD and BPD from these results. It is likely that the onset of deviations is present from early childhood in ASD as sensory symptoms and differences in visual perception are already present in childhood [69], and multistable perception already differs from that of typically developing children before adolescence [37]. Since BPD is not characterised as a neurodevelopmental disorder, deviations probably have a later onset. By adolescence, the disorder can be reliably diagnosed [51], so the deviations from typical trajectories likely occur during, or somewhat before adolescence.

In interpreting the clinical findings, it should be mentioned that BPD is a disorder that predominantly affects females [50] (although one study found equal prevalence of BPD among both sexes [70]). Distinctive patterns in hormone levels, especially the relative changes in ovarian hormones may induce the expression of BPD features [71]. Our BPD participants seem to be in a higher sensitivity and exploration range than neurotypical women of the same age, however, this is obtained at the expense of stability, demonstrating the force of the trade-off, especially outside of the sweet-spot. With respect to ASD which is more often diagnosed in males [50], a similar tendency can be observed: increased levels of sensitivity and exploration and reduced stability as compared to neurotypicals of similar age. Although female-underdiagnosis [72] should not be overlooked, a popular theory of autism claims that the autistic brain is a hypermasculinized version of the male brain due to increased fetal testosterone levels [73]. In terms of excessive levels of sensitivity and exploration, our result from simulated experiments support this picture, although alternative interpretations, such as a general drawback of sex differentiation [74,75] cannot be ruled out. Including male BPD, female ASD participants, larger samples of patient groups, and hormonal assessments would be essential in further studies.

Our results regarding ASD and BPD may inform the field of computational psychiatry, which aims at the large-scale phenotyping of human behaviour using computational models, with the hope that it may structure the search for genetic and neural contributions of healthy and diseased cognition [76]. Within this field, the framework of developmental computational psychiatry aims to establish normative developmental trajectories of computations, relate them to brain maturation, and determine when and how they deviate in mental disorders, in order to help uncover the relationship between the changes of brain organization in childhood and adolescence, and the heightened vulnerability to psychiatric disorders in these periods [15]. Our results fit into this framework by establishing typical developmental trajectories of visual decision making, and relating results from subjects with mental disorders to these trajectories. They also may serve as a small step of the large-scale phenotyping efforts to better understand the nature of mental disorders in terms of aberrant computations.

To conclude, in our detailed study on the lifespan trajectories of perceptual

performance employing a no-response binocular rivalry paradigm combined with dynamic computational modelling, we have found characteristic age- and sex-specific developmental and maturational trajectories, with marked differences between neurotypical and psychiatric populations. These trajectories should serve to better describe our own neurocognitive phenotype and reveal relevant factors behind atypical development underlying mental health disorders.

4 Methods

4.1 Participants

A total of 107 participants took part in the binocular rivalry experiment: 28 twelve-year-old (19 female), and 19 sixteen-year-old (10 female) children; 52 neurotypical adults (average age 35.9, range 18 to 69, 32 female); 12 adults with autism spectrum disorder (ASD, average age 29, range 19 to 44, n=12, all male), and 12 adults with borderline personality disorder (BPD, average age 27, range 20-37, n=12, all female). Participants were considered typically developing, or neurotypical, if they reported no history of mental illness or disorder.

Nine of the twelve participants with ASD were recruited from the Department of Psychiatry and Psychotherapy, Semmelweis University. These participants were diagnosed by a trained psychiatrist. They went through a general psychiatric examination, and their parents were interviewed about early autism-specific developmental parameters. All participants fulfilled the diagnostic criteria of ASD, including autism-specific signs between the ages of 4-5 years. The other three participants in this group were recruited from Aura Organization, a nonprofit organization assisting people with ASD. We did not collect further diagnostic information from these participants. The participants with BPD were all recruited from the Department of Psychiatry and Psychotherapy, Semmelweis University. Their diagnostic status was assessed by the Hungarian version of the Structured Clinical Interview for the Diagnostic and Statistical Manual of Mental Disorders, fourth edition, Axis I and II disorders.

All participants had normal or corrected to normal vision, and reported no colour blindness. Before the experiment, all participants passed a stereoacuity test (Super Stereoacuity Timed Tester, by Stereo Optical Co., U.S. Patent No. 5,235,361, 1993). All adult participants, and all the children's caregivers have provided informed written consent. The study was approved by the Ethical Review Committee of the Institute of Psychology, Pázmány Péter Catholic University for neurotypical participants, and by the Semmelweis University Regional and Institutional Committee of Science and Research Ethics for participants with a psychiatric condition. Participants were given a book voucher for their participation.

4.2 Binocular rivalry experiment

Participants were fixated at a headrest during the experiments (SR Research Head Support, https://www.sr-research.com). The setup for dichoptic stimulation consisted of two LCD displays (subtending 26.6° horizontally and 21.5° vertically, with an approximate resolution of 48 pixels/° of visual angle, and a refresh rate of 120 Hz), which participants viewed through two 45° mirrors,

attached to the headrest. The mirrors were coated, such that they reflected the visible light spectrum, but transmitted infrared light, allowing the use of an infrared camera for optical eye tracking.

The participants viewed green-and-black gratings with one eye, and red-and-black gratings with the other eye. The gratings moved horizontally, either consistently (in the same direction), facilitating perceptual fusion, or inconsistently (in opposing directions), facilitating perceptual rivalry. Each grating subtended a rectangular area of 15.2° width and 8.4° height. The spatial frequency was 0.26 cycles/°, and the temporal frequency 8.7 cycles/s. The motion's speed was 33.5°/s or 1600 pix/s. Gratings were framed in a rectangular box with a random texture pattern, in order to facilitate binocular fusion. Stimuli were generated with Psychophysics Toolbox 3 [77–79] running under MATLAB R2015a. The display's spatial resolution was 48 pix/°, and its temporal refresh rate was 120 Hz.

Before the experiment, participants were asked to view the display as attentively as possible, and to follow the horizontally moving gratings with their gaze. This was introduced with analogies such as "follow them like you would follow passing trees on a moving train". We did not tell them that they will see rivalling stimuli, only that if the direction of the gratings they see will seem to change, let their gaze change direction too. We asked them to refrain from blinking as much as convenient. Participants did not have to report which stimuli they were perceiving at any time. Instead, perceptual states and transitions were calculated from eye-movement recordings.

The experiment consisted of ten trials, each 95 s long. The initial trial (introductory trial) served to familiarize participants with the display, and was not included in the analysis. It started with 22 s of consistent grating motion in alternating directions, followed by 72 s of inconsistent motion, and finishing with 1 s of consistent motion. During the introductory trial, we provided feedback for the participants on the behaviour of their eye movements. The following nine trials (experimental trials) began with 2 s of consistent motion, followed by 92 s of inconsistent motion, and ended with 1 s of consistent motion. The consistent episodes served to reduce eye strain and to test the ocular response to physical motion reversals. On experimental trials, participants received no feedback on their behaviour. Across trials, the colour (either red or green) and direction (either leftward or rightward) shown to each eye was altered. The experiment consisted of three blocks. After the third and sixth experimental trials, participants had a 5-minute break.

4.3 Establishing reversal sequences and dominance statistics from OKN

When a rivalrous display induces horizontal OKN, the direction of smooth pursuit phases provides a reflex-like indication of perceived direction [47–49]. During the experiment, we recorded horizontal eye position of subjects with a sample rate of 1000 Hz, and inferred reversals of perceived direction with the cumulative smooth pursuit (CSP) method, described elsewhere [46]. Briefly, the method removes off-scale values (blinks and other artefacts), and defines slow pursuit segments by a compound criterion (slow velocity $|v| \leq 1.5$ pix/ms, low acceleration $|a| \leq 0.12$ pix/ms^2, duration > 50 ms), aligns slow pursuit segments, interpolates gaps, then subsamples and fits the resulting sequence multiple times

("bagging" [80]). The result of this robust splining procedure is estimated eye velocity (median and 95% CI) at every time-step. Dominance periods were defined as contiguous intervals in which the entire 95% CI is either above or below a gaze velocity threshold of ± 0.1 pix/ms. All other intervals were designated as perceptual transition periods.

As the rate of perceptual reversals often accelerates while viewing a binocular rivalry display [81, 82], the initial 30 seconds of each trial were discarded from analysis. We pooled the remaining reversal sequences obtained for each observer across different trials, and calculated the median (M), interquartile range (IQR), and medcouple (MC) of dominance durations. These robust statistical measures were used to reduce the effect of outliers. Median and interquartile range (IQR) provide robust alternatives for first and second moments (mean and variance), while medcouple (MC) offers a robust alternative for the third moment (skewness), which is particularly sensitive to outliers.

4.4 Establishing developmental and maturational trajectories

For each observer i, three distribution parameters were established as described above: M_i, IQR_i, MC_i, and age a_i. From the individual observer values, average values were computed separately for female and male observers. As development slows down with age, sliding averages were computed with log-normal weighting, so that window size increased proportionally with age. For median M_i, $\overline{M_i}$ was computed as follows:

$$\overline{M_i} = \frac{\sum_j M_j LN(a_j|a_i)}{\sum_j LN(a_j|a_i)}, \qquad LN(a_j|a_i) = \frac{1}{a_j - a_0} \exp[-\ln(a_j - a_0) - a_i + a_0 / 2\sigma^2],$$

where $a_0 = 8$ and $\sigma = 0.35$.

$\overline{IQR_i}$ and $\overline{MC_i}$ were obtained analogously. Confidence intervals were computed by repeatedly sampling observers with replacement, and recomputing the average parameters.

4.5 Computing maturation index

As all parameters tended to change concomitantly, we sought to summarize the development of all three parameters in terms of a single maturation index. As a first step, averaged distribution parameters $\overline{M_i}$, $\overline{IQR_i}$, $\overline{MC_i}$ were normalized (z-scored), to obtain values $\widehat{M_i}$, $\widehat{IQR_i}$, $\widehat{MC_i}$ with zero mean and unit variance, over all typically neurotypical observers. Next, we computed the principal component direction over all typical observers (both male and female), which captured most of the variance ($\sim 80\%$). We defined the maturation index (MI) as the projection of normalized average parameters ($\widehat{M_i}$, $\widehat{IQR_i}$, $\widehat{MC_i}$) onto this direction, or equivalently, a linear combination of these parameter triplets:

$$MI = PC1 = 0.542 \ \widehat{M_i} - 0.567 \ \widehat{IQR_i} - 0.620 \ \widehat{MC_i} \qquad \text{(79\% variance)}$$

The maturation index provided a convenient summary of binocular rivalry statistics over different ages and sexes. The other components were:

$$PC2 = 0.765\ \widehat{M_i} + 0.639\ \widehat{IQR_i} + 0.084\ \widehat{MC_i} \qquad (16\%\ \text{variance})$$
$$PC3 = 0.349\ \widehat{M_i} - 0.520\ \widehat{IQR_i} + 0.780\ \widehat{MC_i} \qquad (3\%\ \text{variance})$$

Confidence intervals were computed by repeatedly sampling observers with replacement, and recomputing the average parameters.

4.6 Computational model implementation

Bistable perception was modeled in terms of a dynamic system with competition, adaptation, and noise. The specific formulation we used was introduced by Laing and Chow [55, 83, 84] and has been analyzed and extended by several other groups [23, 85–88].

The dynamic response $r_{1,2}$ of each neural representations is given by

$$\tau_r \dot{r}_{1,2} = -r_{1,2} + F\left(-\beta r_{2,1} - \phi_a a_{1,2} + I_{1,2} + n_{1,2}\right)$$

with sensory input $I_{1,2}$, intrinsic noise $n_{1,2}$, adaptive state $a_{1,2}$ and activation function

$$F(x) = \left[1 + \exp\left(-\frac{x}{k}\right)\right]^{-1}$$

The dynamics of adaptive states $a_{1,2}$ is given by

$$\tau_a \dot{a}_{1,2} = -a_{1,2} + r_{1,2}$$

and intrinsic noise $n_{1,2}$ is generated from two independent Ornstein–Uhlenbeck processes:

$$\dot{n}_{1,2} = -\frac{n_{1,2}}{\tau_n} + \sqrt{\frac{2\sigma_n^2}{\tau_n}}\xi$$

where ξ is normally distributed white noise. Different dynamical regimes may be obtained by varying competition (β), adaptation (ϕ_a, τ_a), and noise (σ_n), while keeping time constants τ_r and τ_n, and activation parameter k fixed. Inputs were set as equal ($I_1 = I_2 = 1$).

4.7 Fitting computational model parameters

We performed grid simulations for competition strengths $\beta = 1$, $\beta = 2$, $\beta = 3$, and $\beta = 4$, because competition strength is not well constrained by our observations. Each simulation lasted 104 s and covered 100-by-100-by-100 value triplets of the other critical model parameters, adaptation strength (ϕ_a), time constant of adaptation (τ_a), and noise (σ_n). The explored range of parameter values was $\phi_a \in [0.1, 0.5]$, $\sigma_n \in [0, 0.1]$ for $\beta = 1$; $\phi_a \in [0.3, 1.2]$, $\sigma_n \in [0, 0.4]$ for $\beta = 2$; $\phi_a \in [0.5, 2.0]$, $\sigma_n \in [0, 0.4]$ for $\beta = 3$; and $\phi_a \in [1, 4]$, $\sigma_n \in [0, 0.5]$ for $\beta = 4$. For all values of β, τ_a was in a range of $[0.1, 1.3]$ s. The remaining, non-critical parameters were $I_{1,2} = 1$, $\tau_n = 0.1$ s, $\tau_r = 0.02$ s, $k = 0.1$, and $dt = 0.002$ s in all four cases.

For each of the 3-by-106 value quadruplets of β, ϕ_a, σ_n, and τ_a, we parsed the resulting reversal sequence of 104 s duration into dominance periods by taking $sign(r_1 - r_2)$, and calculated three summary statistics of dominance duration: median, interquartile range (IQR), and medcouple (MC).

To compare simulations to the reversal statistics of human observers, we identified combinations of model parameters that reproduce the observed binocular rivalry statistics (median, IQR, MC). Specifically, those parameter combinations were selected for which the simulated median, IQR and MC falls within 5% of the respective mean values obtained for any given participant.

4.8   Simulating experiments with modulated inputs

After fitting model parameter combinations to observers, we performed simulations on these parameter combinations with a time-varying input bias, ΔI_t, generated as an Ornstein-Uhlenbeck process with standard deviation $\sigma_I = 0.3$ and autocorrelation time $\tau_I = 0.2$ s. To keep total input constant $I_1(t) + I_2(t) = 2$, the bias was applied anti-symmetrically $I_1(t) = 1 + \Delta I(t)$, $I_2(t) = 1 - \Delta I(t)$.

The simulated dynamics were obtained in time-steps of 1 ms, and included response $\Delta r = r_1 - r_2$, perceptual dominance $sign(\Delta r)$, differential adaptation $\Delta a = (a_1 - a_2)\ sign(r)$, and differential input $\Delta I = (I_1 - I_2)\ sign(\Delta r)$. Note that positive Δa favours the suppressed percept, whereas positive ΔI favours the dominant percept. Intrinsic noise $n_{1,2}$ was treated as unobservable, and was subsumed in the probabilistic analysis described in the next section. In other words, probabilities and expectation values were obtained by averaging over intrinsic noise.

4.9   Obtaining perceptual parameters

We calculated perceptual parameters from the simulations described above. In the simulated time series, reversals were defined as time-points where responses were equal, $r_1 = r_2$. To calculate perceptual parameters, we disregarded transition periods, defined as 20 ms before and after a reversal. We classified all other time points as either initiation periods preceding a reversal (40-21 ms before a reversal), or as periods not closely preceding a reversal. Based on this classification, we established the following probabilities: the joint probability of $P(\Delta I, \Delta a)$, the conditional joint probability given a subsequent reversal (being in initiation) $P(\Delta I, \Delta a \mid init)$, and the conditional joint probability of Δa and ΔI given no subsequent reversal $P(\Delta I, \Delta a \mid no\ rev)$. From this, the we calculated conditional probability of a subsequent reversal (i.e., initiation) given input bias and model response, $P(init \mid \Delta I, \Delta a)$, for each time-point.

In the vicinity of the median state $(\Delta I_M, \Delta a_M)$, the log reversal probability varies almost linearly with ΔI and Δa (Fig. 4 b, quality of linear regression $r^2 > 0.99$). The three perceptual parameters - sensitivity, stability, and exploration - were defined to parametrize this dependence. Stability was defined as the gradient vector, or, equivalently, as the tangent of the slope, at the median state $tan(\beta) = \sqrt{\gamma_I^2 + \gamma_A^2}$. Stability was defined as the difference between the median and neutral levels of $\log P_{init}$, $-\gamma_I \Delta I_M - \gamma_A \Delta A_M$. Exploration was defined as the direction of the gradient vector at the median state $sin(\alpha) = \frac{\gamma_A}{\sqrt{\gamma_I^2 + \gamma_A^2}}$. These definitions are presented alongside visualizations in Supplementary figure 2.

5 Data availability

The eye tracking datasets are available from the authors on request.

6 Computer code

The code used to extract reversal sequences is available via `https://github.com/cognitive-biology/Cumulative-smooth-pursuit-analysis-of-BR-OKN`. The code used for computational modelling and simulated experiments is available from the authors on request.

References

[1] Sowell, E. R. et al. Mapping cortical change across the human life span. Nat. Neurosci. 6, 309–315 (2003).

[2] Paus, T. et al. Maturation of white matter in the human brain: A review of magnetic resonance studies. Brain Res. Bull. 54, 255–266 (2001).

[3] Geidd, J. N. Structural magnetic resonance imaging of the adult brain. Ann. N. Y. Acad. Sci. 1021, 77–85 (2004).

[4] Gomez-Robles, A., Hopkins, W. D., Schapiro, S. J. & Sherwood, C. C. Relaxed genetic control of cortical organization in human brains compared with chimpanzees. Proc. Natl. Acad. Sci. U. S. A. 112, 14799–14804 (2015).

[5] Foulkes, L. & Blakemore, S. J. Studying individual differences in human adolescent brain development. Nat. Neurosci. 21, 315–323 (2018).

[6] Stiles, J., Brown, T. T., Haist, F. & Jernigan, T. L. Brain and Cognitive Development. in Handbook of Child Psychology and Developmental Science 1–54 (John Wiley & Sons, Inc., 2015). doi:10.1002/9781118963418.childpsy202.

[7] Lebel, C., Treit, S. & Beaulieu, C. A review of diffusion MRI of typical white matter development from early childhood to young adulthood. NMR Biomed. 32, 1–23 (2019).

[8] Giedd, J. N., Raznahan, A., Mills, K. L. & Lenroot, R. K. Review: Magnetic resonance imaging of male/female differences in human adolescent brain anatomy. Biol. Sex Differ. 3, 1–9 (2012).

[9] Ruigrok, A. N. V. et al. A meta-analysis of sex differences in human brain structure. Neurosci. Biobehav. Rev. 39, 34–50 (2014).

[10] Xin, J., Zhang, Y., Tang, Y. & Yang, Y. Brain Differences Between Men and Women: Evidence From Deep Learning. Front. Neurosci. 13, (2019).

[11] Váša, F. et al. Conservative and disruptive modes of adolescent change in human brain functional connectivity. Proc. Natl. Acad. Sci. U. S. A. 117, 3248–3253 (2020).

[12] Khundrakpam, B. S., Lewis, J. D., Zhao, L., Chouinard-Decorte, F. & Evans, A. C. Brain connectivity in normally developing children and adolescents. Neuroimage 134, 192–203 (2016).

[13] Whitaker, K. J. et al. Adolescence is associated with genomically patterned consolidation of the hubs of the human brain connectome. Proc. Natl. Acad. Sci. U. S. A. 113, 9105–9110 (2016).

[14] Paus, T., Keshavan, M. & Giedd, J. N. Why do many psychiatric disorders emerge during adolescence? Nat. Rev. Neurosci. 9, 947–957 (2008).

[15] Hauser, T. U., Will, G. J., Dubois, M. & Dolan, R. J. Annual Research Review: Developmental computational psychiatry. J. Child Psychol. Psychiatry Allied Discip. 60, 412–426 (2019).

[16] Bellman, R. Dynamic Programming. (1957).

[17] Sutton, R. S. & Barto, A. G. Reinforcement Learning: An Introduction. (MIT Press, 2018).

[18] Yang, S. C. H., Wolpert, D. M. & Lengyel, M. Theoretical perspectives on active sensing. Current Opinion in Behavioral Sciences vol. 11 100–108 (2016).

[19] Noel, J.-P. et al. Supporting generalization in non-human primate behavior by tapping into structural knowledge: Examples from sensorimotor mappings, inference, and decision-making. Prog. Neurobiol. 101996 (2021) doi:10.1016/j.pneurobio.2021.101996.

[20] Cisek, P., Puskas, G. A. & El-Murr, S. Decisions in changing conditions: The urgency-gating model. J. Neurosci. 29, 11560–11571 (2009).

[21] Carland, M. A., Thura, D. & Cisek, P. The Urge to Decide and Act: Implications for Brain Function and Dysfunction. Neuroscientist vol. 25 491–511 (2019).

[22] Tajima, S., Drugowitsch, J., Patel, N. & Pouget, A. Optimal policy for multi-alternative decisions. Nat. Neurosci. 22, 1503–1511 (2019).

[23] Pastukhov, A. et al. Multi-stable perception balances stability and sensitivity. Front. Comput. Neurosci. 7, 1–18 (2013).

[24] Leopold, D. A., & Logothetis, N. K. Multistable phenomena: changing views in perception. Trends in cognitive sciences, 3(7), 254-264. (1999)

[25] Braun, J. & Mattia, M. Attractors and noise: Twin drivers of decisions and multistability. NeuroImage vol. 52 740–751 (2010).

[26] Cao, R., Pastukhov, A., Aleshin, S., Mattia, M. & Braun, J. Instability with a purpose: how the visual brain makes decisions in a volatile world. bioRxiv 2020.06.09.142497 (2020) doi:10.1101/2020.06.09.142497.

[27] Brascamp, J., Sterzer, P., Blake, R. & Knapen, T. Multistable Perception and the Role of the Frontoparietal Cortex in Perceptual Inference. Annu. Rev. Psychol. 69, 77–103 (2018).

[28] Ukai, K., Ando, H. & Kuze, J. Binocular rivalry alternation rate declines with age. Percept. Mot. Skills 97, 393–397 (2003).

[29] Kovács, I. & Eisenberg, M. Human Development of Binocular Rivalry. in Binocular Rivalry 101–116 (2004). doi:10.7551/mitpress/1605.003.0008.

[30] Hudak, M. et al. Increased readiness for adaptation and faster alternation rates under binocular rivalry in children. Front. Hum. Neurosci. 5, 1–7 (2011).

[31] Beers, A. M. Characterizing Binocular Rivalry Across the Lifespan. (2016).

[32] Pitchaimuthu, K. et al. Occipital GABA levels in older adults and their relationship to visual perceptual suppression. Sci. Rep. 7, 1–11 (2017).

[33] Abuleil, D., Mcculloch, D. L. & Thompson, B. Older adults exhibit greater visual cortex inhibition and reducedvisual cortex plasticity compared to younger adults. Front. Neurosci. 13, 1–7 (2019).

[34] Robertson, C. E., Kravitz, D. J., Freyberg, J., Baron-Cohen, S. & Baker, C. I. Slower Rate of Binocular Rivalry in Autism. J. Neurosci. 33, 16983–16991 (2013).

[35] Freyberg, J., Robertson, C. E. & Baron-Cohen, S. Reduced perceptual exclusivity during object and grating rivalry in autism. J. Vis. 15, 1–12 (2015).

[36] Robertson, C. E., Ratai, E. M. & Kanwisher, N. Reduced GABAergic Action in the Autistic Brain. Curr. Biol. 26, 80–85 (2016).

[37] Karaminis, T., Lunghi, C., Neil, L., Burr, D. & Pellicano, E. Binocular rivalry in children on the autism spectrum. Autism Res. 10, 1096–1106 (2017).

[38] Spiegel, A., Mentch, J., Haskins, A. J. & Robertson, C. E. Slower Binocular Rivalry in the Autistic Brain. Curr. Biol. 29, 2948-2953.e3 (2019).

[39] Miller, S. M. et al. Slow binocular rivalry in bipolar disorder. Psychol. Med. 33, 683–692 (2003).

[40] Antonio Aznar Casanova, J., Amador Campos, J. A., Moreno Sánchez, M. & Supèr, H. Onset time of binocular rivalry and duration of inter-dominance periods as psychophysical markers of ADHD. Perception 42, 16–27 (2013).

[41] Vierck, E. et al. Further evidence for slow binocular rivalry rate as a trait marker for bipolar disorder. Aust. N. Z. J. Psychiatry 47, 371–379 (2013).

[42] Jia, T. et al. Difference in the binocular rivalry rate between depressive episodes and remission. Physiol. Behav. 151, 272–278 (2015).

[43] Xiao, G. et al. Slow Binocular Rivalry as a Potential Endophenotype of Schizophrenia. Front. Neurosci. 12, 1–9 (2018).

[44] Ye, X., Zhu, R. L., Zhou, X. Q., He, S. & Wang, K. Slower and less variable binocular rivalry rates in patients with bipolar disorder, OCD, major depression, and schizophrenia. Front. Neurosci. 13, 1–11 (2019).

[45] Jia, T. et al. Difference in binocular rivalry rate between major depressive disorder and generalized anxiety disorder. Behav. Brain Res. 391, 112704 (2020).

[46] Aleshin, S., Ziman, G., Kovács, I. & Braun, J. Perceptual reversals in binocular rivalry: Improved detection from OKN. J. Vis. 19, 5–5 (2019).

[47] Naber, M., Frässle, S. & Einhäuser, W. Perceptual Rivalry: Reflexes Reveal the Gradual Nature of Visual Awareness. PLoS One 6, e20910 (2011).

[48] Frässle, S., Sommer, J., Jansen, A., Naber, M. & Einhäuser, W. Binocular rivalry: Frontal activity relates to introspection and action but not to perception. J. Neurosci. 34, 1738–1747 (2014).

[49] Tsuchiya, N., Wilke, M., Frässle, S. & Lamme, V. A. F. No-Report Paradigms: Extracting the True Neural Correlates of Consciousness. Trends Cogn. Sci. 19, 757–770 (2015).

[50] American Psychological Association. Diagnostic and statistical manual of mental disorders. (2013).

[51] Guilé, J. M., Boissel, L., Alaux-Cantin, S. & Garny de La Rivière, S. Borderline personality disorder in adolescents: prevalence, diagnosis, and treatment strategies. Adolesc. Health. Med. Ther. Volume 9, 199–210 (2018).

[52] Daros, A. R., Zakzanis, K. K. & Ruocco, A. C. Facial emotion recognition in borderline personality disorder. Psychol. Med. 43, 1953–1963 (2013).

[53] Daros, A. R., Uliaszek, A. A. & Ruocco, A. C. Perceptual biases in facial emotion recognition in borderline personality disorder. Personal. Disord. Theory, Res. Treat. 5, 79–87 (2014).

[54] Martos-Moreno, G. Á., Chowen, J. A. & Argente, J. Metabolic signals in human puberty: Effects of over and undernutrition. Mol. Cell. Endocrinol. 324, 70–81 (2010).

[55] Laing, C. R. & Chow, C. C. A Spiking Model for Binocular Rivalry. J. Comput. Neurosci. 12, 1–18 (2002).

[56] Fujiwara, M. et al. Optokinetic nystagmus reflects perceptual directions in the onset binocular rivalry in Parkinson's disease. PLoS One 12, 1–22 (2017).

[57] Heidi, L. The Shifting Boundaries Of Adolescence. Nature 554, 429–431 (2018).

[58] Johnson, M. B. & Stevens, B. Pruning hypothesis comes of age. Nature 554, 438–439 (2018).

[59] Fung, M. H. et al. Pubertal Testosterone Tracks the Developmental Trajectory of Neural Oscillatory Activity Serving Visuospatial Processing. Cereb. Cortex 30, 5960–5971 (2020).

[60] Killanin, A. D. et al. Development and sex modulate visuospatial oscillatory dynamics in typically-developing children and adolescents. Neuroimage 221, 117192 (2020).

[61] Hamson, D. K., Roes, M. M. & Galea, L. A. M. Sex Hormones and Cognition: Neuroendocrine Influences on Memory and Learning. Compr. Physiol. 6, 1295–1337 (2016).

[62] Gurvich, C., Hoy, K., Thomas, N. & Kulkarni, J. Sex differences and the influence of sex hormones on cognition through adulthood and the aging process. Brain Sci. 8, (2018).

[63] Shaw, P. et al. Neurodevelopmental trajectories of the human cerebral cortex. J. Neurosci. 28, 3586–3594 (2008).

[64] Li, R. et al. Developmental maturation of the precuneus as a functional core of the default mode network. J. Cogn. Neurosci. 31, 1506–1519 (2019).

[65] Araujo, A. B. & Wittert, G. A. Endocrinology of the aging male. Best Pract. Res. Clin. Endocrinol. Metab. 25, 303–319 (2011).

[66] Rannevik, G. et al. A longitudinal study of the perimenopausal transition: altered profiles of steroid and pituitary hormones, SHBG and bone mineral density. Maturitas 21, 103–113 (1995).

[67] Craik, F. I. M. & Bialystok, E. Cognition through the lifespan: Mechanisms of change. Trends Cogn. Sci. 10, 131–138 (2006).

[68] Harris, I. D., Fronczak, C., Roth, L. & Meacham, R. B. Fertility and the aging male. Rev. Urol. 13, e184-90 (2011).

[69] Simmons, D. R. et al. Vision in autism spectrum disorders. Vision Res. 49, 2705–2739 (2009).

[70] Grant, B. F. et al. Prevalence, correlates, disability, and comorbidity of DSM-IV borderline personality disorder: Results from the Wave 2 National Epidemiologic Survey on Alcohol and Related Conditions. J. Clin. Psychiatry 69, 533–545 (2008).

[71] Eisenlohr-Moul, T. A., DeWall, C. N., Girdler, S. S. & Segerstrom, S. C. Ovarian hormones and borderline personality disorder features: Preliminary evidence for interactive effects of estradiol and progesterone. Biol. Psychol. 109, 37–52 (2015).

[72] Kreiser, N. L. & White, S. W. ASD in Females: Are We Overstating the Gender Difference in Diagnosis? Clin. Child Fam. Psychol. Rev. 17, 67–84 (2014).

[73] Baron-Cohen, S. et al. Why are Autism Spectrum conditions more prevalent in Males? PLoS Biol. 9, (2011).

[74] Alaerts, K., Swinnen, S. P. & Wenderoth, N. Sex differences in autism: A resting-state fMRI investigation of functional brain connectivity in males and females. Soc. Cogn. Affect. Neurosci. 11, 1002–1016 (2016).

[75] Ferri, S. L., Abel, T. & Brodkin, E. S. Sex Differences in Autism Spectrum Disorder: a Review. Curr. Psychiatry Rep. 20, (2018).

[76] Montague, P. R., Dolan, R. J., Friston, K. J. & Dayan, P. Computational psychiatry. Trends in Cognitive Sciences vol. 16 72–80 (2012).

[77] Brainard, D. H. The Psychophysics Toolbox. Spat. Vis. 10, 433–436 (1997).

[78] Pelli, D. G. The VideoToolbox software for visual psychophysics: Transforming numbers into movies. Spat. Vis. 10, 437–442 (1997).

[79] Kleiner, M. et al. What's new in psychtoolbox-3. Perception 36, 1–16 (2007).

[80] Bühlmann, P. & Yu, B. Analyzing bagging. Ann. Stat. 30, 927–961 (2002).

[81] Lehky, S. R. Binocular rivalry is not chaotic. Proc. R. Soc. B Biol. Sci. 259, 71–76 (1995).

[82] Mamassian, P. & Goutcher, R. Temporal dynamics in bistable perception. J. Vis. 5, 361–375 (2005).

[83] Vattikuti, S. et al. Canonical Cortical Circuit Model Explains Rivalry, Intermittent Rivalry, and Rivalry Memory. PLOS Comput. Biol. 12, e1004903 (2016).

[84] Cohen, B. P., Chow, C. C. & Vattikuti, S. Dynamical modeling of multi-scale variability in neuronal competition. Commun. Biol. 2, 1–11 (2019).

[85] Moreno-Bote, R., Rinzel, J. & Rubin, N. Noise-induced alternations in an attractor network model of perceptual bistability. J. Neurophysiol. 98, 1125–1139 (2007).

[86] Shpiro, A., Curtu, R., Rinzel, J. & Rubin, N. Dynamical Characteristics Common to Neuronal Competition Models. J. Neurophysiol. 97, 462–473 (2007).

[87] Curtu, R., Shpiro, A., Rubin, N. & Rinzel, J. Mechanisms for frequency control in neuronal competition models. SIAM J. Appl. Dyn. Syst. 7, 609–649 (2008).

[88] Shpiro, A., Moreno-Bote, R., Rubin, N. & Rinzel, J. Balance between noise and adaptation in competition models of perceptual bistability. J. Comput. Neurosci. 27, 37–54 (2009).

7   Acknowledgements

We acknowledge support from NKFI (Nemzeti Kutatási, Fejlesztési és Innovációs Hivatal) 110466 and 134370 to IK, and from DFG (Deutsche Forschungsgemeinschaft) BR987/3 and BR987/4 to JB, as well as helpful discussions with Alexander Pastukhov. We thank Péter Soltész, Doctoral School of Psychology, Eötvös Loránd University, Budapest, Hungary for carrying out an extensive pilot study before the current project, and Kinga Farkas, MD, PhD, Department of Psychiatry and Psychotherapy, Semmelweis University, for recruiting and diagnosing patients. We are indebted for the work of Tímea Jáger, a graphic artist who helped us to generate the final versions of the figures.

8 Author contributions

GZ designed, developed and performed behavioral experiments; SA analyzed eye movements and conducted computational modelling; ZU recruited and diagnosed BPD patients, all authors discussed the results and contributed to the final manuscript.

9 Competing interest

The authors declare no competing interests.