Distinct function of Mediator subunits in fungal development, stress response and

secondary metabolism in fungal pathogen *Fusarium verticillioides*

Zehua Zhou[1,2], Huijuan Yan[1,3], Man S. Kim[1,4] and Won Bo Shim[1]*

[1] Department of Plant Pathology and Microbiology, Texas A&M University, College Station,

TX, 77843, USA

[2] College of Plant Protection, Nanjing Agricultural University, Nanjing, 210095, China

[3] Current Address: Department of Microbiology and Immunology, University of California at

San Francisco, San Francisco, CA 94143, USA

[4] Current Address: Department of Biomedical and Health Informatics, The Children's Hospital

of Philadelphia, Philadelphia, PA 19104, USA

Running title: Mediator subunits in *Fusarium verticillioides*

*Corresponding author: Won Bo Shim

E-mail: wbshim@tamu.edu

ABSTRACT

Mediator is a conserved, nucleus-localized, multi-subunit complex highly conserved across eukaryotes. It interacts with RNA polymerase II transcription machinery as well as various transcription factors to regulate gene expression. However, systematic characterization of Mediator complex has not be thoroughly performed in filamentous fungi. In current study, our aim was to investigate key biological functions of Mediator subunits in a mycotoxigentic plant pathogen *Fusarium verticillioides*. While we recognized some level of divergence in subunits constituent, overall structure remained conserved between *Saccharomyces cerevisiae* and *F. verticillioides*. We were able to generate 11 Mediator subunit deletion mutants and characterize serveal important phenotypes associated with fungal development, environmental stress, nutrient utilization, virulence as well as secondary metabolism. While each Mediator subunit deletion mutant showed deficiencies in at least three of the phenotypes tested, it is important to note that different modules or subunits showed unique regulatory role in different cellular processes. Significantly, the deletion of *FvMed1* led to increased FB_1 production, and we were able to confirm that FvMed1 is transported from the nucleus to the cytoplasm under fumonisin-producing conditions. Taken together, our study characterized the different biological functions of Mediator subunits in plant pathogen *F. verticillioides* and the possibility of select subunits unique cytoplasmic function independent of the core Mediator.

IMPORTANCE (150 words)

Mediator is a highly conserved component of RNA polymerase II pre-initiation complex (PIC) in eukaryotes. Studies in yeast have demonstrated the important roles of Mediator subunits, but we lack deeper understanding in filamentous fungi. Here, we selected *Fusarium verticillioides*,

a major fungal pathogen of maize worldwide, to study transcriptional profile followed by

mutational analyses of all Mediator genes. Each Mediator subunit deletion mutant showed

some shared deficiencies but it is also important that different modules or subunits showed

unique regulatory role in different cellular processes. Significantly, *FvMed1* deletion mutant

showed elevated levels of FB_1 production, and we were able to confirm that FvMed1

translocates from the nucleus to cytoplasmic organelles under fumonisin-producing conditions.

This is a first study demonstrating that a Mediator subunit regulate cellular functions while not

directly associating with PIC in filamentous fungi.

KEYWORDS *Fusarium verticillioides*, Mediator, fungal development, fumonisin B1,

cytoplasmic translocation

INTRODUCTION

Cellular growth, differentiation and development is a complex and highly regulated process that relies on differentially and methodically coordinated expression of numerous genes in fungi. The coordinated expression, including protein-coding genes and various non-coding RNAs, is accomplished by the RNA polymerase II (Pol II) complex. Notably, Pol II holoenzyme does not carry out transcription alone but performs function as a macromolecular structure known as the pre-initiation complex (PIC), which consist of Pol II, Mediator, TFIIA, TFIIB, TFIID, TFIIE, TFIIF and TFIIH (1, 2). Among these components, Mediator serves as the central scaffold of PIC and is known as the main regulator of Pol II activity. However, owing to its structural complexity and expansive size, our understanding of the biochemical and genetic functions of Mediator in filamentous fungi is rudimentary.

Mediator complex was first discovered in *Saccharomyces cerevisiae* (3), and their organization and structure seem to be evolutionarily conserved among eukaryotes. Mediator complex contains 30 (human) or 25 (yeast) subunits which can be classified into four functional modules: Head, Middle, Tail and Kinase. Significantly, the subunit 14 (Med14) serves as a scaffold and holds the Head, Middle and Tail modules together as a complex, which is defined as the 'core Mediator' (4-6). Published studies indicated that the 'core Mediator' serves as a bridge between DNA-binding transcription factors (TFs), Pol II as well as other components in the transcription machinery, which collectively plays a crucial role in transcriptional activation (7-10). The Kinase module, which consists of four subunits Med12, Med13, Cdk8 (cyclin-dependent kinase 8) and Ssn (cyclin C), could associate reversibly with the 'core Mediator', thereby regulating Pol II activity and transcription (11-13).

While the functions of Mediator appear highly conserved among eukaryotes, analyses of Mediator mutants in various cells and tissues have shown that each subunit of the Mediator

complex regulates distinct subsets of genes during various developmental processes. For

example, Ssn3 and Ssn8 are required for glucose metabolism and biofilm formation in *C.*

albicans (14). In *S. cerevisiae*, Med5 is involved in expression of genes associated with nuclear

and mitochondrial oxidative phosphorylation (OXPHOS) (15). It is interesting to note that

while Med7 is essential for yeast viability, its ortholog in *C. albicans* is not essential and is

involved crucial developmental control (16). Med31 is a maternal-associated gene in

Drosophila (17), whereas no defects have been detected during early embryonic growth in

mouse mutants (18). These reports support gene- and species-specific functions within

Mediator even though it has many conserved roles across eukaryotes. Although the biological

and physiological functions have been extensively studied in a select number of model

organisms, little is known about Mediator functions filamentous fungi.

Fusarium verticillioides is the fungal pathogen responsible for devastating stalk rot, ear

rot and seedling blight diseases in maize, which can lead to severe yield losses worldwide (19,

20). Importantly, *F. verticillioides* produces a variety of mycotoxins during maize infestation

in the field or in storage. Among these mycotoxins, fumonisins B_1 (FB_1) poses a particular

concern, with links to birth defects and cancer in humans as well as acute and chronic toxicosis

in animals (21-24). Therefore, the concentration of fumonisins in maize and food product is

regulated by governmental agencies worldwide (25). One of the key challenges in developing

effective mitigation strategies for fumonisins in food and feeds is our limited understanding of

how fumonisin biosynthesis is regulated in *F. verticillioides*. Published studies showed that

Kinase module subunits, namely cyclin C and cyclin C-dependent kinase, play a crucial role in

mycotoxin biosynthesis in *Fusarium* species (26-29), but the role of other Mediator subunits is

uncharacterized to date.

In order to investigate the functions of Mediator subunits, we implemented a systematic

study of all Mediator subunit genes in *F. verticillioides* by transcriptional analyses and gene

knockout strategy. In our study, 23 Mediator subunit genes were found in *F. verticillioides*, and

the functional roles of 11 Mediator subunits were investigated in this study. Our results provide

further evidence that different modules and subunits play a unique role in transcriptionl

regulation of specific subsets of genes in *F. verticillioides*. Significantly, the translocation of a

Mediator subunit from the nucleus to the cytoplasm was observed in fumonisin-producing

conditions and non-producing conditions, suggesting that Mediator subunits may perform yet-

to-be determined functions in the cytoplasm independent of Pol II.

116   **RESULTS**

Mediator subunit composition in *S. cerevisiae* and F. *verticillioides*. In all eukaryotes,

including *S. cerevisiae*, Mediator subunits have been categorized into four functional modules:

Head, Middle, Tail and Kinase (4, 15, 30, 31). We used *S. cerevisiae* subunits as queries and

screened *F. verticillioides* predicted proteins for putative orthologs. Table 1 describes how

Mediator subunits in two fungi can be comparatively organized. Although the Mediator subunit

organization is conserved between *F. verticillioides* and *S. cerevisiae*, we did learn that *S.*

cerevisiae subunits Med20, Med10, Med19 and Med2 were not identifiable in *F. verticillioides*.

Also, we found that *F. verticillioides* harbors 15 copies of *FvMED15* gene while the gene

encoding *MED15* resides as a single copy locus in *S. cerevisiae*. These genes are named as the

telomere-associated genes (*TLOs*) and may suggest the evolutional divergence of Med15

between *F. verticillioides* and *S. cerevisiae*.

Transcription profile of Mediator genes. To determine whether certain Mediator

subunits are transcriptionally associated with FB_1 biosynthesis regulation, we analyzed their

expression levels in RNA-Seq libraries generated from *F. verticillioides* cultured in maize

inbred line B73 kernels (moderate FB_1 production) versus hybrid 33K44 kernels (high FB_1

production). We followed the relative transcription levels of Mediator subunits in 2, 4, 6, 8 days

post inoculation (dpi) samples (32). Overall, the expression of each Mediator gene increased

through the time course and highest level at 8 dpi, especially noticeable for *FvMed14*

(*FVEG_01411*), *FvMed12* (*FVEG_02030*), *FvMed31* (*FVEG_02237*), *FvMed7*

(*FVEG_04284*), *FvMed21* (*FVEG_06141*) and *FvMed3* (*FVEG_07603*) (Fig. S1). However, it

is important to note that the expression of all Mediator genes showed no statistical difference

between two RNA-Seq libraries as shown in Fig. 1 where we highlighted key Head

(*FVEG_03198*), Middle (*FVEG_02872*, *FVEG_02237*), Tail (*FVEG_07603*) and Kinase

(*FVEG_11159*, *FVEG_11306*) subunits. Notably, transcription profile of cyclin C *FvFcc1*

(*FVEG_11306*) and Cyclin-dependent kinase *FvFck1* (*FVEG_11159*) were in agreement with

earlier published studies (26, 27). These outcomes led us to predict Mediator subunits do not

regulate FB_1 biosynthesis transcriptionally but rather by forming complex with core Mediator

or PIC in *F. verticillioides*.

Construction of Mediator complex deletion mutants. In an effort to test the biological

functions of Mediator subunits in *F. verticillioides*, we generated knockout mutation of each

gene via split-marker approach (Fig. S2). All transformants were analyzed by PCR and qPCR

assays with corresponding primers to verify null mutation (Table S1). For 11 Mediator subunits,

at least two knockout mutants with similar phenotypes were obtained (Table S2). As for the

other 10 putative Mediator subunits (Table 1), we were not able to retrieve viable gene-deletion

mutants after screening more than 200 transformants from at least four independent

transformation experiments. This outcome led us to conclude that the null mutation of these

subunits is deemed lethal in *F. verticillioides*. Significantly, these subunits are also recognized

as essential genes for *S. cerevisiae* viability (33).

Mediator complex play different roles in fungal development processes. When we

tested vegetative growth in mutants, strains with Head module gene deletion (ΔFvMed18) and

Kinase module gene deletion (ΔFvMed12, ΔFvMed13, ΔFvFcc1 and ΔFvFck1) exhibited

significantly reduced growth rate compared to the wild-type progenitor on V8 agar (Fig. 2A

and Table 2). We also compared colony morphology and growth rate of 11 Mediator mutants

on 0.2×PDA and Myro agar plates (Table 2 and Fig. S3). Among these 11 mutants, 6 showed

reduced conidiation by > 30% compared to the wild-type progenitor. Surprisingly, ΔFvMed9

and ΔFvMed5 exhibited significantly increased conidiation (Fig. 2B and Table 2). We also

measured conidia germination rate in all Mediator subunit mutants and found that 9 mutants

exhibited dramatic defect in conidia germination (Fig. 2C and Table 2). Interestingly,

ΔFvMed18 conidia fail to germinate even after a 6.5-hour incubation on water agar (WA) plate.

In addition, we observed that mycelial surface of some Mediator mutants exhibited different

level of hydrophobicity compared with the wild-type progenitor. As showed in Fig. 2D, a 5-μL

water droplet could persist for more than 1 h on WT and the majority of Mediator mutants.

However, ΔFvMed18, ΔFvMed5, ΔFvMed16 and ΔFvMed12 exhibited different degree of

reduced hydrophobicity (Fig. 2D). Taken together, these results suggested that Mediator

complex is critical for fungal development processes in *F. verticillioides*, but each subunit

likely have different principal functions.

Involvement of Mediator complex in plant virulence. To test the role Mediator subunits

play in *F. verticillioides* virulence, we used maize seedling infection assay as previously

described (34). After a 10-day incubation at room temperature, all mutants exhibited reduced

virulence when compared to the wild-type progenitor. In particular, seedling rot symptoms

caused by ΔFvMed18, ΔFvMed12, ΔFvMed13, ΔFvFcc1 and ΔFvFck1 were restricted around

the inoculation sites (Fig. 3), suggesting that Head and Kinase modules in the Mediator

complex play more crucial roles compared to Middle and Tail modules in fungal virulence.

Mediators show different levels of stress and fungicide sensitivity in *F. verticillioides*.

We asked whether Mediator complex subunits play a role in regulating responses to

environmental stress agents and subsequently tested all mutants against cell wall-damaging

agent, oxidative and osmotic stresses, and fungicides. Our results showed that only 5 mutants

(ΔFvMed18, ΔFvMed12, ΔFvMed13, ΔFvFcc1 and ΔFvFck1) exhibited slightly decreased

sensitivity to cell wall-damaging agent Congo Red (Fig. 4 and Fig. S4). Overall, no sensitivity

differences were observed among 11 Mediator mutants to osmotic stress in comparison with

the wild-type progenitor in our experiments (data not shown).

In human pathogens, Mediator has been shown to play an important role in azole fungicide

resistance (35-39). Intriguingly, when we assayed the sensitivity of *F. verticillioides* Mediator

mutants, deletion of Head and Kinase module subunits (ΔFvMed18, ΔFvMed12, ΔFvMed13,

ΔFvFcc1 and ΔFvFck1) caused significant increase in ketoconazole and fluconazole resistance

compared to the wild-type progenitor. However, no sensitivity difference was detected among

other mutants (Table S3). These results demonstrated that Head and Kinase modules are

negatively associated with azole resistance in *F. verticillioides*.

Mediators are involved in carbon and fatty acid utilization. In earlier published studies,

two homologs of Kinase module, CycC and Cdk8, were suggested as a link between ambient

environment sensing and transcriptional regulation in *F. verticillioides* (26, 27). To further

understand the roles of the Mediators in carbon source utilization, each mutant was cultured on

minimal medium (MM) amended with glucose, fructose, lactose, starch or acetate as a sole

carbon nutrient. As results showed, the vegetative growth of wild-type strain and all mutants

were similar on MM amended with fructose, lactose and starch when compared to glucose (Fig.

S5A). Interestingly, most of the mutants in 'core Mediator' exhibited reduced growth rate with

acetate (except for ΔFvMed31 and ΔFvMed5). In contrast, mutants in Kinase module showed

increased relative growth rate on MM amended with acetate as compared with glucose (Fig. 5).

These results indicated that core Mediator plays contrasting role when compared to Kinase

module in acetate utilization in *F. verticillioides*.

We further tested carbon nutrient utilization using fatty acids. We observed that ΔFvMed18

exhibit reduced relative growth rate with butyrate sodium but increased relative growth rate

with palmitic acid and corn oil compared with glucose (Fig. S5B). However, mutants in Kinase

module showed increased relative growth rate with all fatty acids (excluding Med13 with

butyrate sodium) tested in our study (Fig. S5B). These results suggested that subunits in Head

and Kinase module are involved in fatty acids utilization, but their functions are discrepant.

Mediator complexes play distinct roles in secondary metabolism. Fumonisin B_1 (FB_1)

is the most harmful secondary metabolite produced by *F. verticillioides* (40). We assayed the

FB_1 production in Mediator mutants when they were incubated for 7 days on autoclaved corn

kernels (26). After normalization with fungal ergosterol level in each sample set, we were able

to confirm that FB_1 production was significantly reduced in 10 Mediator mutants in comparison

with the wild-type progenitor. One unexpected outcome was that ΔFvMed1 exhibited

drastically higher FB_1 biosynthesis than the wild type (Fig. 6A and Table S4). Consistently, the

expression level of *PKS11* (*FUM1*) was significantly up-regulated in ΔFvMed1 (Table S5).

These results led us to hypothesize that FvMed1 functions as a negative regulator of FB_1

biosynthesis in *F. verticillioides* while majority of other Mediator subunits serve as the positive

regulator.

To test the scope of Mediator complex subunit in regulating secondary metabolism in *F.*

verticillioides, all mutants were cultured in YEPD (to promote vegetative growth) and Myro

(to promote secondary metabolite production) liquid media for 3 days. As results showed,

ΔFvMed3, ΔFvMed12, ΔFvMed13, ΔFvFcc1 and ΔFvFck1 exhibited higher levels of dark

pigmentation in comparisons with the wild-type progenitor (Fig. 6B). Furthermore, qRT-PCR

assays showed that the expression of gene associated with fusarubins biosynthesis, *PKS3*, is

significant up-regulated in ΔFvMed18 and ΔFvFcc1, while deletion of ΔFvMed1 significant

decrease *PKS3* expression (Table S5). Taken together, our results indicated that different

Mediator modules and subunits regulate specific subset of genes involved in secondary

metabolism in *F. verticillioides*.

Subcellular localization analysis indicates functional variety of FvMed1. It is

recognized that Mediator subunits are localized in cell nucleus and regulates gene transcription

processes as an indispensable component of PIC (29, 41, 42). However, our FB_1 assay results

indicated that FvMed1 plays a unique role unlike other subunits under FB_1 producing

conditions. When we analyzed *in silico* putative Med1-interacting protein in *S. cerevisiae*,

majority of the published and predicted interacting proteins were nuclear proteins, *e.g.*,

subunits of Mediator complexes and histone proteins. However, we learned that certain non-

nuclear proteins, e.g. cytoplasmic proteins, mitochondrial proteins and endoplasmic reticulum

proteins, were identified as Med1-interacting proteins in *S. cerevisiae*. This raised a question

whether Med1 can translocate outside the nucleus in response to certain ambient signals or

conditions. In order to test this idea, we generated a FvMed1-GFP fusion construct with its

native promoter and transformed it into the wild-type protoplast. All transformants were

verified by PCR assay and observed under fluorescence microscope. Under FB_1 non-producing

condition, we observed weak GFP signals likely in the nucleus but clearly not in vacuoles in

each septate mycelial cell (Fig. 7A). Under FB_1-producing condition, although Med1-GFP

could be found in the nucleus, we also observed multiple punctate and rounded structures in

the cytoplasm (Fig. 7B - D). Significantly, we did not observe GFP signals localizing to

vacuoles as well as a large number of lipid bodies, which are commonly observed when F.

verticillioides is grown in secondary metabolite-inducing media. Also, that fact that the GFP

signal was not diffused throughout the cytoplasm suggests that FvMed1 translocates to a

defined cellular organelle when it moves out of the nucleus. When assayed for *FvMed1*

expression in YEPD and Myro cultures, the transcript level of *FvMed1* showed no difference

between two media (data not shown), suggesting that *FvMed1* may be constitutively expressed

in different growth conditions. These results showed that FvMed1 proteins were transported to

the cytoplasm in response to FB_1 promoting signals.

DISCUSSION

Mediator is a conserved protein complex that performs indispensable cellular functions as

a regulator of gene transcription. Here, we identified 21 genes encoding putative *F.*

verticillioides Mediator homologs by using 25 *S. cerevisiae* Mediator subunits as queries.

Although the modular organization of Mediator complexes are predicted to be similar between

S. cerevisiae and *F. verticillioides*, the biological functions of certain subunits show significant

differences in *F. verticillioides*. For instance, the gene encoding *Med15* resides as a single copy

locus in *S. cerevisiae*, but the homolog in *F. verticillioides* seems to have expanded to 11 copies

of telomere-associated genes (*TLO*s). We are making this prediction base on previous report

that showed *S. cerevisiae Med2* being replaced by a subset of *TLO*s in *C. albicans* (42).

Furthermore, when we studied 11 viable gene knockout mutants, we observed some distinct

phenotypic outcomes in different mutants.

It is perceived that Mediator is important for fungal development processes, secondary

metabolism and plant infection. Intriguingly, we observed phenotypic variations when

comparing between modules and subunits, indicating that certain modules or subunits of Mediator complex preferentially control the expression of different subset of target genes, thus regulating specific developmental processes. Disruption of a Kinase module exhibited defects in growth and conidia production in *F. graminearum* (28, 29). Another study showed that *CaMed7* is required for hyphal development (16). In our study, 5 Mediator mutants displayed significant defect in vegetative growth, including mutants in Head and Kinase modules. In addition, while the conidia production in ΔFvMed18 was similar to the wild-type strain, the conidial germination was conspicuously suppressed. Furthermore, we observed that conidia produced in ΔFvMed9 and ΔFvMed5 were significantly increased compared to the wild-type progenitor. These results indicated that each module or subunit of Mediator have unique functions during fungal development processes in *F. verticillioides*.

In *Cryptococcus neoformans*, *Cdk8* and *Ssn8* deletion mutants showed increased sensitivity to cell wall and oxidative stress (44). The authors provided further evidence that this outcome was due to mitochondria morphology defect and the resulting transcriptional regulation defects were associated with oxidation-reduction mechanism. In our study, we observed that mutants in Head and Kinase modules exhibited reduced tolerance to cell wall damage agent. However, no significant difference was found in wild-type strain and all Mediator mutants in regard to oxidative stress agents (data not shown). These results suggested that Mediator is involved in regulating environment stress responses, but there may be species-specific functions against various stress agents. Elucidating the mechanism in which these processes are disrupted in *F. verticillioides* may require further study.

Glucose is the most suitable carbon source for eukaryotes, especially fungi. However, fungi can utilize a wide array of carbon sources, even the most complex carbohydrates, to survive and thrive in the most challenging environmental conditions. Previous research showed that certain subunits of Mediator complex are involved in glucose metabolism and carbon

source utilization (14, 44-46). Our study showed how each Mediator mutant can differ in its

ability to utilize different carbon sources. While most of the mutants in the core Mediator

showed defects in using acetate as sole carbon source, mutants in Kinase module exhibited

increased growth rate in MM amended with acetate when compared to glucose. Surprisingly,

these results do not agree with those observed in *C. neoformans* ΔCdk8 and ΔSsn8 mutants

(44), indicating that Mediator has species-specific functions in regulating genes involved in

glucose metabolism.

A number of studies have demonstrated the importance role of Mediator in virulence. In

Candida glabrata, the Med2 subunit was shown to control the Epa adhesins biosynthesis, with

the mutant displaying elevated adherence to epithelial cells (37). In *C. albicans*, Med13 and

Med31 subunits can activate genes associated with adhesion thereby stimulating fungal

infection (47). Here, we observed that all Mediator mutants in *F. verticillioides* exhibited

attenuated disease symptoms on maize seedling, despite each Mediator mutant showing

varying reduced virulence levels. Additionally, our results are not consistent with those

observed in *C. albicans*, further supporting the divergence of function within Mediator despite

their conserved roles across eukaryotes.

Our effort to effectively control fungal diseases in crop production faces persistent

difficulty due to the limitation in antifungal agent availability. More seriously, this difficulty is

exacerbated due to the emergence of resistant pathogen population in the field. Many studies

showed that Mediator plays an important role in azoles resistance by regulating transcriptional

level of efflux pump genes. In *C. glabrata*, the expression of efflux pump genes *CgCDR1* and

CgCDR2 were significantly decreased in *CgMed15A* mutant, which resulted in increased

sensibility to azole fungicides (36, 48). In addition, deletion of *CgMed2* also resulted in the

reduction of *CgCDR1* expression and sensitivity to azoles (37). However, *CgMed1* deletion

mutant did not exhibit any changes to azole sensitivity and *CDR2* expression (48). In our study,

we tested the sensitivity of wild-type and all Mediator mutants against two azole fungicides

ketoconazole and fluconazole. We observed that mutants in Head and Kinase module exhibited

increased drug resistance compared to the wild-type and other mutant strains. However, qPCR

assays showed that the expression of efflux pump genes in these mutants showed no statistical

difference in comparison to the wild-type progenitor and other mutants. Intriguingly, recent

studies showed that the deletion of *Ssn3* or *Ssn8* could increase azole resistance, which is

associated with chromatin remodeling (49, 50). Thus, we hypothesize that the increased drug

resistance in *F. verticillioides* Mediator mutants share similar mechanism to that in *C. albicans*.

In our study, FB_1 biosynthesis was significantly reduced in majority of the Mediator mutant

except ΔFvMed1 and ΔFvMed9. Similar results were observed in *F. graminearum* (28, 29).

Surprisingly, we found that FB_1 biosynthesis in ΔFvMed1 was about 2-fold higher than wild-

type progenitor. We questioned whether FvMed1 plays different roles in other secondary

metabolism compared to other Mediator subunits, and analyzed the expression of all 15 *PKS*

genes in ΔFvMed1 under toxin-inducing conditions. We learned that the expression of *PKS11*

(*FUM1*) was significantly higher in ΔFvMed1. Additionally, we observed that the expression

of other *PKS* genes showed significantly difference among 4 Mediator mutants tested (Table

S5). These results implied that Mediator complex is critical for secondary metabolism in

phytopathogen, but each subunit played unique roles and regulated specific subsets of genes.

Although many reports focused on the functions of Mediator subunits in the nucleus and

as a component to RNA Pol II, an earlier study revealed their possible roles in the cytoplasm

(51). In lung cancer cells, Med12 was found in both the nucleus and the cytoplasm. When

localized in the cytoplasm, Med12 participated in TGF-beta signaling regulation via binding

with TGF-beta receptor 2, thereby affecting its glycosylation and cell membrane expression

(51). We hypothesize that FvMed1 can translocate to the cytoplasm under FB_1-producing

conditions and perhaps perform its unique function as a cytoplasmic protein. Our result raised

a possibility that FvMed1 can control yet-to-be determined cytoplasmic signaling pathways

associated with FB_1 biosynthesis. Although the potential cytoplasmic FvMed1-interacting

proteins remains unknown, this outcome provides us with motivation to expand our efforts to

gain novel insight into the cytoplasmic function of Mediator subunits in filamentous fungi.

<div align="center">**MATERIALS AND METHODS**</div>

Strains and culture assays. *F. verticillioides* wild-type strain M3125 was used for the

construction of the derived mutants used in this study (34). All strains used in this study were

grown and evaluated at 28˚C on V8, 0.2×PDA and Myro agar plates for mycelial growth (34).

For conidia gemination assay, conidia (10^6) grown on V8 plates for 6 days were harvested and

subsequently cultivated in water agar plates for 6.5 h at 28˚C in dark. For stress response assays,

5-mm diameter mycelial plugs taken from a 6-day-old colony of wild-type and all mutant

strains grown on 0.2×PDA were inoculated on 0.2×PDA plate containing 0.05% Congo red

(w/v) or NaCl (1.2 M). Carbon source utilization assays were performed on minimal medium

(MM) supplemented with different carbon sources at concentrations of 1% (w/v). All

experiments were repeated three times independently.

RNA-Seq data analysis. RNA-seq data were obtained through sequencing platform,

Illumina HiSeq 2500, from Texas A&M AgriLife Genomics and Bioinformatics Services

(College Station, Texas). The platform processed fourteen independent samples composed of

seven biological replicates and seven technical replicates at four distinct time points (2 dpi, 4

dpi, 6 dpi, and 8 dpi). The sequencing data were generated as paired-end reads with 125-bp for

two different maize kernels (hybrid 33k44 vs. inbred B73), where *F. verticillioides* inoculated.

We first aligned and quantified the RNA-seq by TopHat2 (51) along with HTSeq (53).

Subsequently, we filtered scarcely expressed genes out and normalized the remaining quantity

by two steps, TPM (54) and TMM (55). Our dataset is publicly available at NCBI Gene

Expression Omnibus (Accession GSE146208).

For those selected genes, we analyzed whether their expression levels were

directly/indirectly associated with the virulence by comparing expression levels of 33K44 and

those of B73 at all the time points. *P value* for comparison was computed by the non-parametric

Van der Waerden (VDW) test while not having any assumptions for normality or equality of

variance between 33K44 and B73. The bar plots of their mean expression levels with *P values*

for the comparison are shown in Fig. 1 and Fig. S1.

Strains construction. All gene deletion mutants were constructed using our standard split-

marker gene knock out protocol as previously described (34). Hygromycin B

phosphotransferase gene (*HPH*) was split into two fragments *PH* (929 bp) and *HP* (765 bp)

and fused with left and right flanking sequence fragments of each targeted gene, respectively

(Fig. S2). As the result of this homologous recombination, the open reading frame (ORF) of

each gene was replaced with *HPH* gene. We confirmed targeted gene replacement by PCR,

and further tested the absence of gene transcription by qPCR. FvMed1-GFP fusion cassette

was constructed and transformed into *F. verticillioides* following our standard protocol

described previously (34). All transformants were verified by PCR and observed under an

Olympus BX51 microscope (Olympus America, Melville, NY, USA). A detailed description

of features used for imaging fluorescence with this microscope and the image processing has

been described previously (34).

Sensitivity assay against ketoconazole and fluconazole. To determine the sensitivity to

ketoconazole and fluconazole, a mycelial plug of all strains from a 6-day-old 0.2×PDA plate

was transferred to the center of 0.2×PDA plates amended with ketoconazole and fluconazole

at different concentrations. Three replicates for all concentrations were tested. After incubation

at 28°C for 6 days, the colony diameter in all plates were measured. For each plate, the average

colony diameter was used to calculate the fungicide concentration that resulted in 50% mycelial

growth inhibition (EC_{50}). The EC_{50} values were calculated with the Data Processing System

(DPS) software. Ketoconazole was added to 0.2×PDA plates at 0, 0.01, 0.04, 0.2, 1.0 and 5.0

μg/mL. Fluconazole was added to 0.2×PDA plates at 0, 0.2, 1.0, 5.0, 10.0 and 50.0 μg/mL. The

assays were repeated three time.

RNA preparation and qPCR assays for Mediator subunit genes. To analyze relative

gene expression of Mediator subunits, fungal mycelia were harvested from 3-days-old YEPD

or 3-days-old Myro cultures and used for RNA isolation with the GeneJet RNA Purification

Kit (Thermo Scientific). First-strand cDNA was synthesis by Verso cDNA synthesis kit

(Thermo Scientific) and subsequently used for qPCR assays with DyNAmo ColorFlash SYBR

Green qPCR Kit (Thermo Scientific). The β-tubulin gene of *F. verticillioides* was used as the

internal control. The relative quantification of all genes was calculated with the $2^{-\Delta\Delta Ct}$ method

(56). All qPCR assays were conducted with a StepOne plus realtime PCR system. All results

were calculated with the data from three biological replicates.

Fumonisin B_1 assay and Maize seedling infection assay. To determine FB_1 and

ergosterol production, we followed the standard method described previously (34, 57). FB_1

levels were normalized with ergosterol levels. All results were calculated with the data from

three biological replicates. Stalk rot virulence assays was conducted using silver queen hybrid

seeds as previously described (34). After growth on V8 plates for 6 days, conidia of each mutant

were collected and re-suspended in sterile distilled water to a concentration of 1×10^7 mL^{-1}.

Seedlings were inoculated with 10 µL conidia suspension. After 10 days inoculation in the dark

room, the infected seedlings were recorded and imaged. For all mutants, 15 seedlings were

inoculated.

ACKNOWLEDGMENTS

The research was supported by Texas A&M T3: Triads for Transformation Program and the

China Scholarship Council State Scholarship Fund (201906850058). The authors declare no

conflicts of interest.

REFERENCES

1. Hahn S. 2004. Structure and mechanism of the RNA polymerase II transcription machinery. Nat Struct Mol Biol 11:394-403.

2. Thomas MC, Chiang CM. 2006. The general transcription machinery and general cofactors. Crit Rev Biochem Mol Biol 41:105-178.

3. Myers LC, Kornberg RD. 2006. Mediator of transcriptional regulation. Annu Rev Biochem 69:729-749.

4. Tsai KL, Tomomori-Sato C, Sato S, Conaway RC, Conaway JW, Asturias FJ. 2014. Subunit architecture and functional modular rearrangements of the transcriptional Mediator complex. Cell 157:1430-1444.

5. Plaschka C, Larivière L, Wenzeck L, Seizl M, Hemann M, Tegunov D, Petrotchenko EV, Borchers CH, Baumeister W, Herzog F, Villa E, Cramer P. 2015. Architecture of the RNA polymerase II-Mediator core initiation complex. Nature 518:376-380.

6. Robinson PJ, Trnka MJ, Bushnell DA, Davis RE, Mattei PJ, Burlingame AL, Kornberg RD. 2016. Structure of a complete Mediator-RNA polymerase II pre-initiation complex. Cell 166:1411-1422 .

7. Kelleher RJ 3rd, Flanagan PM, Kornberg RD. 1990. A novel mediator between activator proteins and the RNA polymerase II transcription apparatus. Cell 61:1209-1215.

8. Flanagan PM, Kelleher RJ 3rd, Sayre MH, Tschochner H, Kornberg RD. 1991. A Mediator required for activation of RNA polymerase II transcription in vitro. Nature 350:436-438.

9. Koleske AJ, Young RA. 1994. An RNA polymerase II holoenzyme responsive to activators. Nature 368:466-469.

10. Balamotis MA, Pennella MA, Stevens JL, Wasylyk B, Belmont AS, Berk AJ. 2009. Complexity in transcription control at the activation domain-mediator interface. Sci Signal, 2:ra20.

11. Holstege FC, Jennings EG, Wyrick JJ, Lee TI, Hengartner CJ, Green MR, Golub TR, Lander ES, Young RA. 1998. Dissecting the regulatory circuitry of a eukaryotic genome. Cell 95:717-728.

12. Andrau J, van de Pasch L, Lijnzaad P, Bijma T, Koerkamp MG, van de Peppel J, Werner M, Holstege FCP. 2006. Genome-wide location of the coactivator Mediator: binding without activation and transient Cdk8 interaction on DNA. Mol Cell 22:179-192.

13. Tsai KL, Sato S, Tomomori-Sato C, Conaway RC, Conaway JW, Asturias FJ. 2013. A conserved Mediator CDK8 kinase module association regulates Mediator-RNA polymerase II interaction. Nat Struct Mol Biol 20:611-619.

14. Lindsay AK, Morales DK, Liu Z, Grahl N, Zhang A, Willger SD, Myers LC, Hogan DA. 2014. Analysis of *Candida albicans* mutants defective in the Cdk8 module of Mediator reveal links between metabolism and biofilm formation. PLoS Genet 10:e1004567.

15. Beve J, Hu G, Myers LC, Balciunas D, Werngren O, Hultenby K, Wibom R, Ronne H, Gustafsson CM. 2005. The structural and functional role of Med5 in the yeast Mediator tail module. J Biol Chem 280:41366-41379.

16. Tebbji F, Chen Y, Albert RJ, Gunsalus KTW, Kumamoto CA, Nantel A, Sellam A,

Whiteway M. 2014. A functional portrait of Med7 and the Mediator complex in *Candida albicans*. PLoS Genet 10:e1004770.

17. Bosveld F, van Hoek S, Sibon OC. 2008. Establishment of cell fate during early Drosophila embryogenesis requires transcriptional Mediator subunit dMED31. Dev Biol 313:802-813.

18. Risley MD, Clowes C, Yu M, Mitchell K, Hentges KE. 2010. The Mediator complex protein Med31 is required for embryonic growth and cell proliferation during mammalian development. Dev Biol 342:146-156.

19. Munkvold GP. 2003. Epidemiology of Fusarium diseases and their mycotoxins in maize ears. Eur J Plant Pathol 109:705-713.

20. Blacutt AA, Gold SE, Voss KA, Gao M, Glenn AE. 2018. *Fusarium verticillioides*: Advancements in understanding the toxicity, virulence, and niche adaptations of a model mycotoxigenic pathogen of maize. Phytopathology 108:312-326.

21. Hendricks K. 1999. Fumonisins and neural tube defects in south Texas. Epidemiology 10:198-200.

22. Howard PC, Eppley RM, Stack ME, Warbritton A, Voss KA, Lorentzen RJ, Kovach RM, Bucci TJ. 2001. Fumonisin B1 carcinogenicity in a two-year feeding study using F344 rats and B6C3F1 mice. Environ Health Perspect 109 Suppl 2:277-282.

23. Marasas WFO. 2001. Discovery and occurrence of the fumonisins: A historical perspective. Environ Health Perspect 109 Suppl 2:239-243.

24. Desai K, Sullards MC, Allegood J, Wang E, Schmelz EM, Hartl M, Humpf HU, Liotta DC, Peng Q, Merrill AH Jr. 2002. Fumonisins and fumonisin analogs as inhibitors of ceramide synthase and inducers of apoptosis. Biochim Biophys Acta 1585:188-192.

25. Woloshuk CP, Shim WB. 2013. Aflatoxins, fumonisins, and trichothecenes: a convergence of knowledge. FEMS Microbiol Rev 37:94-109.

26. Shim WB, Woloshuk CP. 2001. Regulation of fumonisin B1 biosynthesis and conidiation in *Fusarium verticillioides* by a cyclin-like (C-type) gene, *FCC1*. Appl Environ Microbiol 67:1607-1612.

27. Bluhm BH, Woloshuk CP. 2006. Fck1, a C-type cyclin-dependent kinase, interacts with Fcc1 to regulate development and secondary metabolism in *Fusarium verticillioides*. Fungal Genet Biol 43:146-154.

28. Zhou XY, Heyer C, Choi YE, Mehrabi R, Xu JR. 2010. The CID1 cyclin C-like gene is important for plant infection in *Fusarium graminearum*. Fungal Genet Biol 47:143-151.

29. Cao, SL, Zhang SJ, Hao CF, Liu HQ, Xu JR, Jin QJ. 2016. FgSsn3 kinase, a component of the mediator complex, is important for sexual reproduction and pathogenesis in *Fusarium graminearum*. Sci Rep 6:22333.

30. Kang JS, Kim SH, Hwang MS, Han SJ, Lee YC, Kim YJ. 2001. The structural and functional organization of the yeast mediator complex. J Biol Chem 276:42003-42010.

31. Harper TM, Taatjes DJ. 2018. The complex structure and function of Mediator. J Biol Chem 293:13778-13785.

32. Zhang H, Kim MS, Hung J, Yan H, Yang T, Song L, Yu W, Shim WB. 2019. FvLcp1, a type-D fungal LysM protein with Chitin-binding domains, is a secreted protein involved in host recognition and fumonisin production in *Fusarium verticillioides* - maize kernel interaction. BioRxiv (**doi:** https://doi.org/10.1101/789925)

33. Jeronimo C, Robert F. 2017. The Mediator complex: At the nexus of RNA polymerase II
transcription. Trends Cell Biol 27:765-783.
34. Yan HJ, Shim WB. 2020. Characterization of non-canonical G beta-like protein FvGbb2
and its relationship with heterotrimeric G proteins in *Fusarium verticillioides*. Environ
Microbiol 22:615-628.
35. Shahi P, Gulshan K, Näär AM, Moye-Rowley WS. 2010. Differential roles of
transcriptional mediator subunits in regulation of multidrug resistance gene expression in
Saccharomyces cerevisiae. Mol Biol Cell 21:2469-2482.
36. Paul S, Schmidt JA, Moye-Rowley WS. 2011. Regulation of the CgPdr1 transcription
factor from the pathogen *Candida glabrata*. Eukaryot Cell 10:187-197.
37. Borah S, Shivarathri R, Srivastava VK, Ferrari S, Sanglard D, Kaur R. 2014. Pivotal role
for a tail subunit of the RNA polymerase II Mediator complex CgMed2 in azole tolerance
and adherence in *Candida glabrata*. Antimicrob Agents Chemother 58:5976-5986.
38. Cowen LE, Sanglard D, Howard SJ, Rogers PD, Perlin DS. 2014. Mechanisms of
antifungal drug resistance. Cold Spring Harb Perspect Med 5:a019752.
39. Liu Z, Myers LC. 2017. Mediator tail module is required for Tac1-activated CDR1
expression and azole resistance in *Candida albicans*. Antimicrob Agents Chemother
61:e01342-17.
40. Rheeder JP, Marasas WFO, Vismer HF. 2002. Production of fumonisin analogs by
Fusarium species. Appl Environ Microbiol 68:2101-2105.
41. Zhu X, Zhang Y, Bjornsdottir G, Liu Z, Quan A, Costanzo M, Dávila López M, Westholm
JO, Ronne H, Boone C, Gustafsson CM, Myers LC. 2011. Histone modifications
influence mediator interac tions with chromatin. Nucleic Acids Res 39:83428354.
42. Liu Z, Myers LC. 2012. Med5 (Nut1) and Med17 (Srb4) are direct targets of mediator
histone H4 tail interactions. PLoS One 7:e38416.
43. Zhang A, Petrov KO, Hyun ER, Liu Z, Gerber SA, Myers LC. 2012. The Tlo proteins are
stoichiometric components of *Candida albicans* Mediator anchored via the Med3 subunit.
Eukaryot Cell 11:874-884.
44. Chang AL, Kang YM, Doering TL. 2019. Cdk8 and Ssn801 regulate oxidative stress
resistance and virulence in *Cryptococcus neoformans*. mBio 10:e02818-18.
45. Chen W, Zhang XT, Birsoy K, Roeder RG. 2010. A muscle-specific knockout
implicates nuclear receptor coactivator MED1 in the regulation of glucose and
energy metabolism. Proc Natl Acad Sci U S A 107:10196-10201.
46. Galbraith MD, Donner AJ, Espinosa JM. 2010. CDK8: a positive regulator of
transcription. Transcription 1:4-12.
47. Uwamahoro N, Qu Y, Jelicic B, Lo TL, Beaurepaire C, Bantun F. Quenault T, Boag
PR, Ramm G, Callaghan J, Beilharz TH, Nantel A, Peleg AY, Traven A. 2012. The
functions of Mediator in *Candida albicans* support a role in shaping species-specific gene
expression. PLoS Genet 8:e1002613.
48. Thakur JK, Arthanari H, Yang F, Pan SJ, Fan X, Breger J, Frueh DP, Gulshan K, Li
DK, Mylonakis E, Struhl K, Moye-Rowley WS, Cormack BP, Wagner G, Näär AM. 2008.
A nuclear receptorlike pathway regulating multidrug resistance in fungi. Nature 452: 604-
609.
49. Liu Z, Myers LC. 2017. *Candida albicans* Swi/Snf and mediator complexes differentially

regulate Mrr1-induced MDR1 expression and fluconazole resistance. Antimicrob Agents
Chemother 61: e01344-17.

50. Long NB, Zeng LP, Qiao SL, Li L, Zhong GW. 2018. *Aspergillus fumigatus* Afssn3-
Afssn8 pair reverse regulates azole resistance by conferring extracellular polysaccharide,
sphingolipid pathway intermediates, and efflux pumps to biofilm. Antimicrob Agents
Chemother 62:e01978-17.

51. Huang SD, Hölzel M, Knijnenburg T, Schlicker A, Roepman P, McDermott U, Garnett
M, Grernrum W, Sun C, Prahallad A, Groenendijk FH, Mittempergher L, Nijkamp
W, Neefjes J, Salazar R, Dijke PT, Uramoto H, Tanaka F, Beijersbergen RL, Wessels
LFA, Bernards R. 2012. MED12 controls the response to multiple cancer drugs through
regulation of TGF-β receptor signaling. Cell 151:937-950.

52. Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions and gene
fusions. Genome Biol 14:R36.

53. Anders S, Pyl PT, Huber W. 2015. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics 31:166-169.

54. Wagner GP, Kin K, Lynch VJ. 2012. Measurement of mRNA abundance using RNA-seq
data: RPKM measure is inconsistent among samples. Theory Biosci 131:281-285.

55. Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics 26:139-40.

56. Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the $2^{-\Delta\Delta Ct}$ method. Methods 25:402-408.

57. Christensen S, Borrego E, Shim WB, Isakeit T, Kolomiets M. 2012. Quantification of
fungal colonization, sporogenesis, and production of mycotoxins using kernel bioassays. J
Vis Exp 62:e3727.

603   **TABLE 1** Comparisons of Mediator modular organization and subunit composition between *Fusarium*

604   *verticillioides* and *Saccharomyces cerevisiae*.

		S. cerevisiae		*F. verticillioides*	
Module	Subunit	Gene ID	Essential	Gene ID	Essential
Scaffold	Med14	Rrg1	Yes	FVEG_01411	Yes
Head	Med6	Med6	Yes	FVEG_04734	Yes
	Med8	Med8	Yes	FVEG_09579	Yes
	Med11	Med11	Yes	FVEG_07465	Yes
	Med17	Srb4	Yes	FVEG_02224	Yes
	Med18	Srb5	No	FVEG_03198	No
	Med20	Srb2	No	-	-
	Med22	Srb6	Yes	FVEG_05568	Yes
Middle	Med1	Med1	No	FVEG_02872	No
	Med4	Med4	Yes	FVEG_06276	Yes
	Med7	Med7	Yes	FVEG_04284	Yes
	Med9	Cse2	No	FVEG_04688	No
	Med10	Nut2	Yes	-	-
	Med19	Rox3	Yes	-	-
	Med21	Srb7	Yes	FVEG_06141	Yes
	Med31	Soh1	No	FVEG_02237	No
Tail	Med2	Med2	No	-	-
	Med3	Pgd1	No	FVEG_07603	No
	Med5	Nut1	No	FVEG_07865	No
	Med15	Gal11	No	11 copies	N/A
	Med16	Sin4	No	FVEG_01083	No
Kinase	Med12	Srb8	No	FVEG_02030	No
	Med13	Ssn2	No	FVEG_02107	No
	Cdk8	Ssn3	No	FVEG_11159	No
	CycC	Ssn8	No	FVEG_11306	No

609
610
611

TABLE 2 Hyphal growth, conidiation and conidial germination of the Mediator deletion mutants.

Strain	Colony diameter (mm)[a]			Conidiation $(10^5/cm^2)$[b]	Percentage of germinated conidia (%)[c]
	V8	PDA	Myro		
WT	80.0±1.4a	63.0±0.8c	66.8±0.9b	104.1±3.9c	89.0±2.6a
ΔFvMed18	44.7±1.2g	38.2±0.8h	29.5±1.4i	100.2±5.0c	-
ΔFvMed1	78.0±1.4a	59.8±1.1e	54.5±0.8e	87.8±2.9d	55.8±9.1c
ΔFvMed9	66.3±0.9d	59.0±1.2e	61.3±1.4d	182.2±6.2a	62.5±9.3bc
ΔFvMed31	70.5±1.3c	61.0±1.0de	62.8±0.9cd	73.7±3.7e	74.8±8.4bc
ΔFvMed3	78.4±1.3a	70.1±1.1a	71.2±0.9a	66.6±3.9e	88.6±1.8a
ΔFvMed5	74.0±1.0b	62.8±1.2cd	68.0±1.4b	115.4±3.8b	92.2±2.8a
ΔFvMed16	75.0±1.0b	65.2±0.9b	64.3±1.3c	85.2±3.8d	70.2±4.7b
ΔFvMed12	64.1±1.2e	54.8±1.2f	40.5±1.2g	26.4±1.4g	31.8±1.2d
ΔFvMed13	69.0±1.1c	60.0±0.9e	43.6±1.4f	47.6±2.4f	38.2±7.9d
ΔFvFcc1	62.5±1.1ef	53.4±1.2f	35.1±0.8h	42.5±2.1f	17.6±2.1e
ΔFvFck1	61.5±0.9f	50.7±1.4g	41.7±0.8fg	40.0±5.7f	20.0±1.8e

[*]Means and standard deviations (SD) were calculated from three independent experiments. Values in table followed by the same letter are not significantly different (P < 0.05).
[a]Colonies were cultured on V8, 0.2×PDA or Myro plates for 6 days at 28°C.
[b]Conidiation inV8 plate was examined after incubation for 6 days at 28°C.
[c]Conidial germination was assayed on Water Agar (WA) after 6.5 h of incubation at 28°C.

Figure legends

Fig. 1 Transcription profile of 6 key Mediator genes in moderately resistant (B73) versus highly

susceptible (33K44) maize seeds. *F. verticillioides* was inoculated on surface sterilized B73

and 33K44 seeds, and RNA-Seq was performed on 2, 4, 6, 8 days-post-inoculation (dpi)

samples. We analyzed whether transcription of Mediator genes show difference in two maize

seeds. *P value* for comparison was computed by the non-parametric Van der Waerden (VDW)

test while not having any assumptions for normality or equality of variance between 33K44

and B73. A complete set of transcriptional comparisons for all *F. verticillioides* Mediator genes

is shown in Fig. S1.

Fig. 2 Phenotypes of the Mediator deletion mutants in vegetative growth, conidiation, conidial

germination and hydrophobicity. (A) Colony morphology of wild-type and Mediator mutants

grown on V8 plate at 28°C for 6 days. (B) Deletion of Mediator subunits on conidia production.

Conidia of all strains were quantified with a hemacytometer after incubation on V8 plate for 6

days. (C) Conidia of each strain were cultivated in WA medium. After incubation at 28°C for

6.5 h, conidial germination of 200 conidia was examined. (D) Hydrophobicity assays of wild-

type and all mutants. Bars denote standard errors from three repeated experiments. Values on

the bars followed by the same letter are not significantly different at P = 0.05, according to

Fisher's LSD test.

Fig. 3 Impact of Mediator gene mutation in virulence when tested on maize seedlings. (A)

Maize seedlings were grown in the dark and inoculated with conidia suspension from all strains.

The images were taken after 10 days of incubation. Sterile water was used as negative control

(NC). (B) The relative lesion size in maize seedling rot assays was measured by image J. (C)

The relative lesion size was normalized with strain growth rate on V8 plate. Bars denote

standard errors from three repeated experiments. Values on the bars followed by the same letter

are not significantly different at P = 0.05, according to Fisher's LSD test.

Fig. 4 Gene disruption of Head and Kinase modules of Mediator increases the sensitivity

towards cell-wall damage agent. (A) Growth phenotype of the wild type (WT) and mutants

with Congo Red amended in 0.2xPDA plates after 6 days of incubation at 28°C. (B) Statistical

analysis of the growth inhibition rate of each strain with Congo Red. Bars denote standard

errors from three repeated experiments. Values on the bars followed by the same letter are not

significantly different at P = 0.05, according to Fisher's LSD test.

Fig. 5 Mediator complex is involved in acetate utilization. (A) All strains were cultivated on

MM with glucose or acetate, and incubated at 28°C for 6 days. (B) The growth inhibition rate

of each strain was analyzed by comparing the growth with acetate against glucose. Bars denote

standard errors from three repeated experiments. Values on the bars followed by the same letter

are not significantly different at P = 0.05, according to Fisher's LSD test.

Fig. 6 Roles of Mediator complex on FB_1 and secondary metabolite biosynthesis. (A) FB_1

production in the wild type (WT) and Mediator mutants grown on autoclaved kernels for 7

days at room temperature. FB_1 production levels were normalized with ergosterol production

in each strain. (B) Comparisons of secondary metabolite biosynthesis among the WT and

Mediator mutants. Images were taken after all strains were grown on liquid Myro for 7 days.

Bars denote standard errors from three repeated experiments. Values on the bars followed by

the same letter are not significantly different at P = 0.05, according to Fisher's LSD test.

Zhou et al. 2021

Fig. 7 Subcellular localization of Med1-GFP in *F. verticillioides* mycelial cells grown in (A)

FB$_1$-non-inducing medium YEP and (B~D) FB$_1$-inducing medium Myro. We inoculated *F.*

verticillioides Med1-GFP conidia in two cultures and took images after 2-day incubation. GFP

expression was observedunder an Olympus BX51 microscope and the images were processed

by ImageJ. Bar = 10 μm.

Zhou et al. Fig 1.

685   Zhou et al. Fig 2.

690   Zhou et al. Fig 3.

695   Zhou et al. Fig 4.

700 Zhou et al. Fig 5.

705 Zhou et al. Fig 6.

710 Zhou et al. Fig 7.

Table S1. Primers used in this study.

Primer	Sequence (5'-3')	Application
Med1_LF/FN	TCCCTAAGGATTTAGACTTC	Amplify *FvMed1* 5' flank sequence, for gene knock out
Med1_LF/R1	TAGATGCCGACCGGGAACGAGCTCCAGGTTGAAAGCAA	
Med1_RF/F1	CCACTAGCTCCAGCCAAGGGCGGAGAAACTGAACGAC	Amplify *FvMed1* 3' flank sequence, for gene knock out
Med1_RF/RN	GCTTCCTCGCTGAGCACTT	
Med4_LF/FN	TTCCTCAAGTCCACCAGAT	Amplify *FvMed4* 5' flank sequence, for gene knock out
Med4_LF/R1	TAGATGCCGACCGGGAACGGTCAATGTGGGTCTTTAGT	
Med4_RF/F1	CCACTAGCTCCAGCCAAGGGTAAACGCGAATTCCAGAT	Amplify *FvMed4* 3' flank sequence, for gene knock out
Med4_RF/RN	ATACTGTCCTCGGCTTCGT	
Med5_LF/FN	GAACGACCACGAACCTATG	Amplify *FvMed5* 5' flank sequence, for gene knock out
Med5_LF/R1	TAGATGCCGACCGGGAACTGGGCTTGGTGTTGACTTT	
Med5_RF/F1	CCACTAGCTCCAGCCAAGGTCAGTCCGTCTATTTTCTTG	Amplify *FvMed5* 3' flank sequence, for gene knock out
Med5_RF/RN	GTCGGTGACACTAATGGTTTG	
Med7_LF/FN	CGTCTTCATCGTATGTGGC	Amplify *FvMed5* 5' flank sequence, for gene knock out
Med7_LF/R1	TAGATGCCGACCGGGAACCGTACCATCCAACGTCATA	
Med7_RF/F1	CCACTAGCTCCAGCCAAGTCAGTAATCTACGCAAGCATG	Amplify *FvMed5* 3' flank sequence, for gene knock out
Med7_RF/RN	ATGCGAGATAGAGGACGGT	
Med9_LF/FN	CAGTGTAGCCCTTGGAGTC	Amplify *FvMed9* 5' flank sequence, for gene knock out
Med9_LF/R1	TAGATGCCGACCGGGAACGGTCTGTCGTTCTATGATATG	
Med9_RF/F1	CCACTAGCTCCAGCCAAGTATCAGCAGGGTGTTTGGG	Amplify *FvMed9* 3' flank sequence, for gene knock out
Med9_RF/RN	CTCTTTGACTCGGTGCTTC	
Med21_LF/FN	GGAGGGCTCGCTTGACTAA	

Table S2. Wild-type and mutant strains of *F. verticillioides* used in this study.

Strain	Genotype description	Reference
M3125	Wild-type strain of *F. verticillioides*	Lab preservation
ΔFvMed18	*FvMed18* deletion mutant of M3125	This study
ΔFvMed1	*FvMed1* deletion mutant of M3125	This study
ΔFvMed9	*FvMed9* deletion mutant of M3125	This study
ΔFvMed31	*FvMed31* deletion mutant of M3125	This study
ΔFvMed3	*FvMed3* deletion mutant of M3125	This study
ΔFvMed5	*FvMed5* deletion mutant of M3125	This study
ΔFvMed16	*FvMed16* deletion mutant of M3125	This study
ΔFvMed12	*FvMed12* deletion mutant of M3125	This study
ΔFvMed13	*FvMed13* deletion mutant of M3125	This study
ΔFvFcc1	*FvFcc1* deletion mutant of M3125	Shim and Woloshuk, 2001 and this study
ΔFvFck1	*FvFck1* deletion mutant of M3125	Bluhm and Woloshuk, 2005 and this study

Table S3. Sensitivity of *F. verticillioides* to ketoconazole and fluconazole.

Strain	EC$_{50}$ (μg/mL)	
	Ketoconazole	Fluconazole
WT	1.41±0.11c	1.32±0.10c
ΔFvMed18	2.76±0.30ab	4.09±0.86a
ΔFvMed1	1.48±0.11c	1.38±0.05c
ΔFvMed9	1.43±0.11c	1.34±0.03c
ΔFvMed31	1.32±0.16c	1.29±0.16c
ΔFvMed3	1.45±0.15c	1.36±0.10c
ΔFvMed5	1.55±0.09c	1.45±0.05c
ΔFvMed16	1.36±0.32c	1.28±0.16c
ΔFvMed12	3.08±0.24a	2.17±0.21b
ΔFvMed13	2.65±0.08b	2.06±0.06b
ΔFvFcc1	2.43±0.19b	2.10±0.07b
ΔFvFck1	2.84±0.22ab	2.17±0.18b

[*]Means and standard deviations (SD) were calculated from three independent experiments. Values in table followed by the same letter are not significantly different (P < 0.05).

EC$_{50}$: the fungicide concentration resulting in 50% inhibition of mycelial growth.

Table S4. Fumonisin B$_1$ (FB$_1$) and ergosterolproduction of the wild-type and Mediator deletion mutants.

	FB$_1$ (μg/L)[a]	Ergosterol (mg/L)[b]	FB$_1$/Ergosterol (μg/mg)[c]
WT	73915.6±12127.6a	375.0±49.8a	197.1±32.3b
ΔFvMed18	4797.6±712.0e	213.1±29.4b	22.5±3.4d
ΔFvMed1	84499.1±4949.8a	210.6±9.9b	401.3±23.5a
ΔFvMed9	6719.5±799.1d	248.3±27.5b	27.1±7.1d
ΔFvMed31	9551.1±1822.6c	234.3±41.0b	39.3±7.5d
ΔFvMed3	13488.1±1790.0b	370.2±53.6a	36.4±4.8d
ΔFvMed5	10845.0±1175.8bc	254.9±24.7b	42.5±6.5d
ΔFvMed16	3270.1±353.0f	264.5±47.0b	12.4±1.3d
ΔFvMed12	2475.8±196.8f	68.7±14.3cd	36.0±2.9d
ΔFvMed13	6349.3±407.3d	73.5±16.7cd	86.4±5.5c
ΔFvFcc1	1636.1±327.9g	62.0±8.8d	26.4±5.3d
ΔFvFck1	2859.8±583.3f	83.5±6.0c	34.2±7.0d

[*]Means and standard deviations (SD) were calculated from three independent experiments. Values in table followed by the same letter are not significantly different (P < 0.05).

[a]FB$_1$ production in wild-type and mutants grown on autoclaved kernels for 7 days at room temperature.

[b]Ergosterol production in wild-type and mutants grown on autoclaved kernels for 7 days at room temperature.

[c]FB$_1$ levels were normalized with fungal ergosterol levels.

Table S5. Relative expression level of 15 *PKS* genes in wild-type and Mediator mutants.

Gene	Relative expression level				
	WT	ΔFvMed1	ΔFvMed3	ΔFvMed18	ΔFvFcc1
PKS1	1	2.55±0.33	ND	ND	8.78±2.52
PKS2	1	0.97±0.10	0.98±0.22	0.64±0.06	3.74±0.37
PKS3	1	0.36±0.03	1.93±0.77	8.66±0.84	31.17±3.65
PKS4	1	8.41±1.12	0.24±0.11	0.29±0.11	9.92±1.52
PKS5	1	5.18±0.78	1.00±0.15	ND	1.14±0.17
PKS6	1	2.30±0.21	1.85±0.31	0.02±0.01	3.70±0.26
PKS7	1	0.47±0.08	2.44±0.36	3.18±0.45	0.63±0.16
PKS8	1	3.70±0.35	11.03±1.96	0.03±0.003	0.05±0.01
PKS9	1	0.31±0.05	0.62±0.04	0.56±0.06	0.27±0.02
PKS10	1	0.14±0.04	2.34±0.34	0.07±0.04	0.58±0.04
PKS11	1	2.49±0.27	0.62±0.2	0.04±0.01	0.06±0.01
PKS12	ND	ND	ND	ND	ND
PKS13	1	1.13±0.07	2.25±0.13	1.21±0.07	1.80±0.11
PKS14	1	0.06±0.01	ND	0.24±0.06	0.08±0.01
PKS15	1	0.66±0.2	0.23±0.07	65.78±21.22	1.42±0.33

Mycelia of each mutants were incubated in myro liquid medium for 3 days at 28℃. Each gene expression were normalized with β-tubulin expression level. Gene expressions were calculated using $2^{-\Delta\Delta Ct}$. Gene expression level of wild-type strain was standardized to 1.0. Three replicates were performed for each test. ND: not detected.

725 **FIG S1.** Assessment of the RNA-Seq Data Set of *F. verticillioides* Mediator genes under
726 different conditions. The Mediator genes expression were normalized with β-tubulin
727 expression. Error bars indicate SD of 7 biological replicates.

730 **FIG S2** Schematic description of split marker approach for constructing Mediator deletion
731 mutants in *F. verticillioides.*
732

735
736
737 **FIG S3** Colony morphology of wild-type and Mediator mutants grown on 0.2×PDA and
738 Myro agar plate at 25°C for 6 days
739

742 **FIG S4** Deletion of Mediator subunits on cell-wall damage agent response in *F.*
743 *verticillioides*
744

747 **FIG S5** Relative growth rates of wild-type and Mediator mutants on various carbon sources.
748 Values on the bars followed by the same letter are not significantly different at P = 0.05,
749 according to Fisher's LSD test