The genetic basis of adaptation through the evolution of self-fertilization

Kuangyi Xu

Department of Biology, University of North Carolina, CB#3280, Coker Hall, University of North Carolina,
Chapel Hill, NC, USA.

kyxu@email.unc.edu

Abstract

Although adaptation can be realized through the fixation of beneficial alleles that increase

viability, many plant populations may adapt through the evolution of self-fertilization, especially

when pollination becomes inefficient. However, the genetic basis of adaptation through the

evolution of selfing remains unclear. Using population genetic models, I study adaptation

through the fixation of alleles that increase the selfing rate (selfing modifiers) from new

mutations or/and standing variation. For adaptive alleles unrelated to selfing, it is known that

selfing promotes adaptation from a new mutation only when the beneficial alleles are recessive,

and the probability of adaptation from standing variation is nearly independent of dominance,

and always decreases with the selfing rate. In contrast, for adaptation through the evolution of

selfing, when it occurs by fixation of a newly arisen mutation, a population that already has a

high selfing rate may be more (less) likely to adapt than outcrossers even when the modifier is

dominant (recessive) if the modifier is weakly (strongly) selected. Also, adaptation from standing

variation is more likely through recessive modifier alleles, with the highest fixation probability

found in partially selfing populations, but fixation is fastest when dominance is intermediate.

When there are multiple modifiers, adaptation through new mutations is more likely when selfing

is controlled by few large-effect rather than many slight-effect modifiers. This study suggests

that to understand the genetic basis of adaptation, it is necessary to determine the ecological

and genetic advantages of adaptive alleles.

Significance statement

This study, by deriving the selective coefficient and effective population size, investigated the

genetic basis of adaptation through fixation of modifier alleles that increase the selfing rate,

which is shown to differ in several aspects from that through evolution of mating-unrelated

alleles. Specifically, when adaptation is from new mutations, the dominance of a selfing modifier

allele below which selfing increases the fixation probability depends on the strength of pollen

limitation and pollen discounting. Adaptation from standing variation is more likely through

recessive modifier alleles and in populations with an intermediate selfing rate. This work

suggests it is important to have a mechanistic understanding of how adaptive alleles increase

individual fitness in environment that is unfavorable to the population.

Introduction

Adaptation often involves fixation of alleles that increase individual fitness, either through new beneficial mutations, or through alleles that are previously present in the population before an environmental change, often called standing genetic variation (Hermisson and Pennings 2005, Barret and Schluter 2008, Orr 2010, Lai 2019). In many plant populations, adaptation may occur through the evolution of their mating systems, especially when environmental change reduces mating success (Barret et al. 2008, Eckert et al. 2010, Pannell 2015, Cheptou 2019). Although our understanding of adaptation through mating-unrelated beneficial alleles has been greatly promoted by previous studies (Orr 2001, Hermisson and Penings 2005, Barret and Schluter 2008, Orr 2010, Ronfort and Glemin 2013, Glemin and Ronfort 2013), how the genetic properties of alleles which modify the selfing rate (hereafter referred as selfing modifiers or modifiers) influence the probability of successful adaptation remains unclear.

Adaptation through the evolution of mating systems may be particularly common for today's plant populations, since an increased level of human disturbance and widespread declines in pollinator populations have intensified limitation of pollen (Larson and Barrett 2000, Knight et al. 2005) and reduced plant reproductive success (Eckert et al. 2010, Thomann et al. 2013). Compared to outcrossing, selfing depends less on extrinsic factors, and thus may offer assurance for seed reproduction when pollination service is scarce (Holsinger 2000, Eckert et al. 2006). Indeed, an increased level of self-fertilization has been widely documented in response to various environmental conditions, including human disturbances (Eckert et al. 2010), habitat fragmentation (Aguilar et al. 2006), range expansion (Levin 2012), and colonization (Barrett et al. 2008, Pannell 2015). Also, the evolution of selfing is often associated with morphological changes in several floral characters (Goodwillie et al. 2010, Sicard and Lenhard 2011), which are analyzed to be controlled by multiple loci with both large and small effects (Fishman et al 2002, 2015, Goodwillie et al. 2006).

Adaptation through the fixation of selfing modifiers differs from that of mating-unrelated alleles in several aspects. First, the selective advantage of a modifier allele depends on both genetic and ecological factors. Regarding genetic factors, selfing enjoys a transmission advantage over outcrossing (Fisher 1941), but suffers the cost of inbreeding depression (Charlesworth and Willis 2009). In the ecological aspect, selfing imposes a cost as it may reduce opportunities for pollen export, often called pollen discounting (Harder and Wilson 1998, Porcher and Lande 2005), but is more advantageous when pollen limitation becomes severe. Second, in a finite population, a selfing modifier should have a different effective population size from a mating-unrelated locus, since the selfing rate is polymorphic at the modifier locus, and during the fixation process, the level of inbreeding is no longer constant but changes with the frequency of the modifier. Third, although previous studies show that the fixation probability of mating-unrelated alleles will often be lower in populations with higher selfing rates (Caballero and Hills 1992, Glemin and Ronfort 2013), this may not hold for selfing modifier alleles, as the selective advantage and effective population size of a modifier may depend on the selfing rate that is already in the population The answer to this question may also help us understand the speed of the evolution of selfing rates during the transition from outcrossing to selfing, which is commonly seen in plant populations (Igic and Busch 2013).

To examine the genetic basis of adaptation through the evolution of selfing, I first derive the effective selection coefficient and the effective population size for a selfing modifier. I then use the diffusion approximation to obtain the fixation probability of such a modifier from a new mutation and from standing variation. I consider adaptation through fixation at a single modifier locus in three demographic situations: a constant population, a declining population, and a population after a bottleneck. For adaptive alleles unrelated to selfing, selfing promotes adaptation from a new mutation only when it is recessive, and the probability of adaptation from

standing variation is independent of dominance. In contrast, for adaptation through the evolution

of selfing, when adaptation is from a new mutation, selfing may inhibit adaptation even when the

modifier is recessive if it is strongly favored, and can facilitate adaptation for dominant, weakly-

favored modifiers, while adaptation from standing variation is more likely through recessive

modifiers. Generally, partial-selfing populations tend to have the highest fixation probability of

selfing modifiers, suggesting a nonlinear evolutionary rate from outcrossing to selfing. An

extension to multiple loci suggests that adaptation through the evolution of selfing is more likely

when selfing is controlled by few large-effect modifiers rather than many small-effect modifiers.

Models and results

I consider a selfing rate modifier locus with two alleles A and a in a hermaphroditic, diploid

population with non-overlapping generations. Asexual reproduction, that is, apomixis (Asker and

Jerling 1992), which only happens in less than 1% of the species (Mogie 1992), is not

considered and thus seeds will be produced from either selfing or outcrossing. The population

already has a background selfing rate r_0, and the selfing rate of the three genotypes AA, Aa and

aa are $r_0 + r$, $r_0 + hr$ and r_0, where h is the dominance coefficient. It is assumed that the

genetic properties of the selfing modifier locus are independent of the environment, so r and h

are the same before and after pollen limitation. The inbreeding depression in the population is d

and the level of pollen discounting is c, which means for an individual with a selfing rate r_i, a

proportion $r_i c$ of its pollen will not be disseminated compared to a completely outcrossing

individual. The population suffers from pollen limitation with a strength m, i.e., a proportion m of

its ovules which should have been outcrossed will not be fertilized. Since the number of viable

offspring from selfing relative to outcrossing is $(1 - d)/(1 - m)$, I can define an effective

inbreeding depression, which incorporates the effect of pollen limitation as

$$d_e = 1 - \frac{1-d}{1-m}.\tag{1}$$

Evolution towards a higher selfing rate will increase the average fitness only when $1 - m < 1 -$

d, so that $d_e < 0$, but $d_e > 0$ will also be considered for a thorough analysis.

To model drift at a modifier locus, the key is to obtain the effective selective coefficient and the

effective population size, based on which the fixation probability of alleles starting from any

frequency can be calculated. This is done by first considering the frequency change and its

variance after one discrete generation (see section 1 in SI), and then using the diffusion

approximation by considering an infinitesimal time interval $dt \to 0$. The overall allele frequency

change is obtained by a weighted average of changes among selfed and outcrossed seeds. For

outcrossed seeds, the allele frequency is the average of those in the pollen and ovule pool. The

overall variance is the sum of two components: variance in selfed and outcrossed offspring

given that the proportion of selfed seeds is fixed, and variation of the actual proportion of selfed

offspring. The first component is calculated by assuming the number of selfed and outcrossed

offspring per individual is subject to a Poisson distribution. After that, the fixation probability of a

modifier allele starting from any frequency can be obtained. The expected fixation probability

from standing variation is integrated over the initial frequency distribution under the selection-

drift-mutation equilibrium before pollen limitation. For declining populations, the fixation

probability is modified to account for the reduction of effective population size caused by

demographic decline, and the population goes to extinction when all pre-existing alleles and all

mutations that appears before the population size declines to 0 will not go fixation. I also extend

the model to investigate adaptation after a bottleneck, the effect of genetic architecture and

coevolution of inbreeding depression. A full presentation can be found in the Methods section.

In the section 2 of SI, I show that the effective selection coefficient R the effective population

size of the selfing enhancer allele A are

$$140 \qquad R = \frac{1 - 2d_e - c(1 - 2d_e r_0)}{2(1 - d_e r_0)(1 - c r_0)} r, \qquad (2a)$$

$$141 \qquad N_e = \frac{2(1 - d_e r_0)}{2 + [1 - 3d_e + F(1 - d_e)]r_0} N. \qquad (2b)$$

F is the inbreeding coefficient, which measures the deviation of the frequency of homozygotes

from the Hardy-Weinberg equilibrium. Although F will change with time, it approaches the

equilibrium much more quickly than the allele frequency changes (Hartfield and Glemin 2016).

Therefore, F is first approximated with its equilibrium value, which depends on the allele

frequency. However, it is shown that a secondary approximation of F as a frequency-

independent constant works well (Table S1), given by

$$148 \qquad F \approx \frac{(1 - d_e)r_0}{2 - (1 + d_e)r_0} + \left(2 - \frac{1 - c}{1 - c r_0}\right) \frac{(1 - d_e)r_0 + h(1 - r_0)}{(2 - (1 + d_e)r_0)^2} r, \qquad (3)$$

Equation (2a) shows the selective advantage depends on the background selfing rate, and

unlike mating-unrelated loci, equations (2b) and (3) show that the effective population size

depends on the effect of the modifier r. N_e also decreases with d_e and c (see section 5 in SI),

and thus is regulated by ecological factors. In contrast, the effective population size of mating-

unrelated alleles should increase with d_e since d_e lowers the level of inbreeding (Caballero and

Hill 1992).

156   Adaptation in a constant population

157   When fixation is from a single new modifier mutant, equation (7) in the Methods section shows

158   that fixation is more likely through dominant modifiers. However, whether a higher background

159   selfing rate will promote or inhibit fixation depends on the dominance coefficient. For a selfing

160   modifier, the critical dominance coefficient h_c below which selfing increases the fixation

161   probability (obtained from equation (8) in Methods) is

$$162 \qquad h_c \approx \frac{1 - 2d_e}{2(1 - d_e)(1 - c - 2d_e)}. \qquad (4)$$

As shown in Fig. 1, h_c is higher than 0.5 under weak pollen limitation and strong pollen

discounting (large d_e and c). The critical dominance above which adaptation through evolution

of selfing is less likely in selfers than outcrossers becomes more restricted compared to

adaptation through mating-unrelated alleles. On the contrary, when the modifier is strongly

favored (d_e and c small), h_c is often smaller than 0.5. Therefore, when the evolution of selfing

will increase the average fitness (i.e., $d_e < 0$), outcrossers may be much more likely to adapt

than selfers. In contrast, h_c is always 0.5 for mating-unrelated alleles. This difference is mainly

because the selective advantage of a selfing modifier depends on the background selfing rate

(see equations (2) and (8) Methods).

When fixation is through standing variation, for mating-unrelated alleles, the fixation probability

is nearly independent of the dominance (Orr and Betancourt 2001), which holds for both

outcrossing and selfing populations (Glemin and Ronfort 2013). However, for selfing modifier

alleles, fixation from standing variation is more likely through recessive alleles, as shown in Fig.

2. This pattern is more prominent when the background selfing rate r_0 is intermediate, when the

modifier is more strongly selected against before pollen limitation, and when the population size

is larger (see Fig. S1). Also, for mating-unrelated alleles, the fixation probability from standing

variation decreases with the selfing rate (Glemin and Ronfort 2013). For adaptation from the

evolution of selfing, this result usually holds when the modifier is not very recessive ($h = 0.5, 0.9$

in Fig. 2), but when the modifier allele is recessive enough, the fixation probability will be highest

when the background selfing rate is at an intermediate level ($h = 0.1$ in Fig. 2). However, when

pollen limitation is weak and pollen discounting is strong (Fig. 2(c)), in which case selfing is

more likely to enhance the fixation probability after pollen limitation (see Fig.1), fixation

probabilities increase almost monotonically with the background selfing rate. The above findings

are robust to the effect of the selfing modifier r, as the fixation probability is nearly independent

of r (Fig. S2).

Despite a higher fixation probability for recessive modifier alleles when adaptation is from

standing variation, generally, fixation is fastest when the allele has intermediate dominance,

while it is slowest for dominant alleles (Fig. S3). However, when the allele is under weak

selection fixation is quickest for recessive alleles (Fig. S3(c)). The above findings are robust to r

(Fig. S4). As a contrast, for mating-unrelated alleles, more dominant alleles will go to fixation

more quickly (Glemin and Ronfort 2013).

The above differences mainly result from the fact that the evolution at a selfing modifier locus

will change the level of inbreeding at the locus itself. At the selection-mutation-drift equilibrium

before pollen limitation, because selfing enhancer alleles are exposed more in homozygotes,

they will be kept at an even lower frequency than mating-unrelated alleles of the same selective

disadvantage. Moreover, a higher dominance will strengthen the selection in two ways. First, it

directly increases the selfing rate of the heterozygote. Second, an enhanced selfing rate of the

heterozygotes further increases the homozygosity, which indirectly increases the strength of

selection. Therefore, compared to dominant mating-unrelated alleles, which increases strength

of selection only through heterozygotes, a dominant selfing modifier will be more strongly

selected against before pollen limitation, and although it will also be more strongly favored after

pollen limitation, this advantage usually cannot compensate for the disadvantage before pollen

limitation.

Adaptation in a declining population

The previous section considers populations that are still able to persist and maintain a constant

size after pollen limitation. However, some populations may not produce sufficient offspring and

thus will suffer a demographic decline and go extinct. To investigate adaptation in declining

populations, I consider a population which has an expected offspring number $1 - \lambda$ after pollen

limitation, and will be successfully rescued when the selfing enhancer allele A goes to fixation.

For the case of evolutionary rescue occuring through new mutations, equation (16) in the

Methods shows that the critical dominance at which selfers and outcrossers have the same

rescue probability is higher than that in a constant population. Therefore, demographic decline

favors selfers to be more likely to adapt than outcrossers.

For rescue from standing variation, demographic decline favors adaptation through dominant

alleles compared to a case with the same conditions in a constant population. This can be seen,

for example, by comparing $r_0 = 0$ in Figs. 3(a) and 2(a). This difference is more prominent

when r is small (Fig. S5). Moreover, for all levels of dominance (Fig. 3(b)), the proportion of

rescue from standing variation is lower for dominant alleles, and decreases with the background

selfing rate, suggesting an increased relative contribution from new mutations in selfing

populations. However, how dominance influences the overall rescue probability from both

sources should depend on the relative contribution from standing variation versus new

mutations. Rescue will be more likely through recessive alleles when the contribution from

standing variation is larger, which happens under a smaller effect r (Fig. S5), a larger initial

population size and a faster decline rate (Glemin and Ronfort 2013).

Adaptation after a bottleneck

In the previous sections, I consider that the pollen limitation happens in the original habitat.

However, the evolution of self-fertilization is often associated with bottleneck effects to some

degree (Eckert et al. 2010, Aguilar et al. 2006, Levin 2012, Barrett et al. 2008, Pannell 2015). I

consider a population of an ancestral size N_0 that suffers from a bottleneck effect of N_c

individuals in a new environment. Equation (7) indicates that a bottleneck will have little effect

on the fixation probability of a new mutant. For adaptation from standing variation, compared to

a constant population, a bottleneck lowers the fixation probability (compare the y axis in Figs.

3(c) and 2(a)). Fixation is even more biased for recessive alleles, especially when the

background selfing is low, which can be easily seen by comparing Fig. 3(c) with Fig. 2(a) at $r_0 =$

0. In contrast, the fixation probability is usually insensitive to dominance for mating-unrelated

alleles (Fig. S6). Moreover, for all levels of dominance, the fixation probability always decreases

with the background selfing rate. It is often considered that self-compatible populations are more

likely to establish after long-distance dispersal, known as Baker's law (Baker 1951, Cheptou

2012). However, it is often misinterpreted as an association between higher selfing rates and

successful colonization. Our result indicates that establishment of a colony through the genetic

evolution of selfing is more probable for outcrossing populations, which supports the view of

Pannell et al. (2015) that Baker's law should be interpreted in terms of an enhanced capacity of

selfing associated with colonization.

254   Effects of the genetic architecture of selfing on adaptation

As selfing is often controlled by multiple modifier loci, it is critical to ask whether adaptation is

more likely when the selfing rate is controlled by many slight-effect modifier loci or few large

effect loci. For adaptation from standing variation, as the fixation probability at a single locus is

insensitive to its effect r (Fig. S2), adaptation is more likely when there is a larger number of

loci. However, the answer is not clear for adaptation from new mutations. Although genetic

architecture of many slight-effect loci has a higher overall mutation rate, the fixation probability

of a single modifier is lower. Here I assume modifier loci act additively on the selfing rate, as

selfing rates sometimes have an approximately linear relationship with phenotypic values of

several quantitative characters (Karron et al. 1997, Strand et al. 2017, Fishman et al. 2002,

2015). Equation (19) in Methods and Fig. S7 show that in constant populations, adaptation from

new mutations is always faster under few large-effect modifier loci, consistent with the empirical

findings (Bodbyl Roels and Kelly 2011), and rescue of declining populations is also more likely

through the genetic architecture of few large-effect loci (see discussion below equation (19)).

Incorporating the genetic basis of inbreeding depression

Up to now, inbreeding depression is treated as a constant parameter. However, inbreeding

depression is caused by deleterious mutations and will coevolve with the mating system.

Previous models and empirical evidence have shown that a higher selfing rate will lower the

inbreeding depression at the equilibrium (Lande and Schemske 1985, Roze 2015). Moreover,

selection against deleterious mutations may also inhibit fixation of a linked beneficial mutation,

known as background selection (Charlesworth et al. 1993, Charlesworth 2012), which is more

severe in selfing populations due to a reduced effective recombination rate.

When inbreeding depression is fixed, adaptation probability tends to either decreases or

increases monotonically with the selfing rate. In contrast, under the coevolution of inbreeding

depression, Fig. 4 shows that for both adaptation from new mutations and standing variation,

the fixation probability is highest at an intermediate selfing rate, suggesting that a trade-off

between the benefit of lower inbreeding depression and the cost of higher background selection.

More specifically, the selfing rate corresponding to the highest fixation probability is higher for

recessive alleles when fixation is from a new mutation (Fig. 4(a)), but lower when it is from

standing variation (Fig. 4(b)). Although the rate of the transition from outcrossing to selfing has

been estimated at the phylogenetic scale (Igic and Busch 2013), the above result implies that at

a smaller scale during this transition, the evolution of self-fertilization accelerates when the

population initially starts from being predominantly outcrossing, while slows down later when it

approaches predominantly selfing.

Conclusion

While the genetic basis of adaptation through the fixation of mating-unrelated adaptive alleles has been well studied (Haldane 193, Orr 2001, Hermisson and Pennings 2005, Glemin and Ronfort 2013), it is not clear whether these results can apply to the case when adaptation is through the evolution of sefl-fertilization. In this study, I investigate fixation of selfing modifier alleles from new mutations and standing variation in populations under different demographic situations. Although for mating-unrelated alleles, selfing increases adaptation from new mutations as long as they are recessive, selfing can lower the fixation probability of a recessive modifier if pollen limitation is strong, but makes dominant modifier more likely to fix under weak pollen limitation (Fig. 1). On the other hand, adaptation from standing variation is more likely through recessive modifier alleles (Fig. 1), which seems to be supported by previous genetic analysis of the rapid evolution of selfing rates in *Mimulus guttatus* (Bodbyl Roels and Kelly 2011). However, fixation is fastest for intermediate dominant modifier alleles (Fig. 2). Generally, adaptation is more likely through dominant alleles under a demographic decline, consistent with adaptation while a bottleneck effect favors recessive alleles.

These characteristics of adaptation through the evolution of mating systems result from two factors. First, a selfing modifier allele has a different inbreeding coefficient from mating-unrelated alleles, as it directly influences the level of inbreeding at the locus itself. Second, dominance has two-fold effects, as it directly increases the selfing rate of the heterozygote, which further increases the homozygosity. Although the dominance coefficient distribution of mating-unrelated alleles is generally unknown (Orr 2010), for selfing modifier alleles, it may be possible to estimate dominance by combining QTL analyses of selfing-related characters (Fishman et al. 2002, 2014) with studies on the association between selfing rate and phenotypic values. This work also provides insights on how the selfing rate of the population affects adaptation through evolution towards a higher selfing rate. Generally, a higher selfing rate limits

the further increase of self-fertilization, but this may be counterbalanced by the coevolution of

deleterious mutations in the genome (Fig.5).

Although I contrast adaptation through the evolution of mating system with that through increase

of viability or fertility, adaptation may involve both paths through fixation of alleles that

simultaneously influence several aspects of individual fitness. For example, an allele that

reduces the flower display may not only allow more resources to be allocated for fertility

(Sargent et al. 2007), but also increase the selfing rate (Sicard and Lenhard 2011). On the other

hand, alleles that increase selfing may impose a fitness cost. For example, reduction of flower

size may be caused by deleterious mutations that inhibit development (Kelly and Willis 2001).

This pleiotropy may make the genetic basis underlying adaptation more complicated. All in all,

this work suggests the importance of a mechanistic understanding of the source of the selective

advantage of adaptive alleles.

Methods

Adaptation in a constant population size

The fixation probability of a selfing modifier allele with an initial frequency x is given by (Kimura

1962)

$$\pi(x) = \frac{\int_0^x G(p)\,dp}{\int_0^1 G(p)\,dp},$$

335   (5)

336 where

$$G(p) = \exp\left[-\int \frac{2M(p)}{V(p)}\,dp\right],$$

337   (6a)

$$M = Rp(1-p)[h + F - hF + (1-F)(1-2h)\,p],$$

338   (6b)

$$V = \frac{p(1-p)}{2N_e}.$$

339   (6c)

340   M and V are the expected change of the allele frequency and its variance under the limit of an

341   infinitesimal time interval. When $N_e(h + F - hF)R \gg 1$, the fixation probability can be

342   approximated as

$$P_{fix}^{new} \approx \frac{(2h(1 - r_0) + (1 - d_e)r_0)\big(1 - 2d_e - c(1 - 2d_e r_0)\big)}{2(1 - cr_0)(1 - d_e r_0)^2} r + o(r). \tag{7}$$

344   The fixation probability of a selfing modifier mutant in a completely outcrossing ($r_0 = 0$) and a

345   nearly completely selfing ($r_0 = 1 - r$) population can be approximated as

$$P_{fix}^{new}(r_0 = 0) \approx (1 - c - 2d_e)hr + o(r), \tag{8a}$$

$$P_{fix}^{new}(r_0 = 1 - r) \approx \frac{1 - 2d_e}{2(1 - d_e)} r + o(r). \tag{8b}$$

When the above two equations are equal, we can solve for the critical dominance h_c at which

the fixation probability is the same for outcrossers and selfers, given in equation (4).

For adaptation from standing variation, denote the mutation rate from allele a to A as v_1, and the

backward mutation rate as v_2. Before pollen limitation happens, allele A is selected against, and

in a finite population under selection-drift-mutation balance, the stationary distribution of

frequency of allele A is (Ewens 2012)

$$\Psi(x) = \frac{C_1}{V(x)} \exp\left[2 \int_0^x \frac{M_d(p)}{V(p)} dp\right], \tag{9}$$

356   where C_1 is a constant which normalizes the integral as $\int_0^1 \Psi(x)dx = 1$, and $M_d(p) = M(p) +$

357     $(1 - p)v_1 - pv_2$, with d_e replaced by d. Therefore, the expected fixation probability from

358   standing genetic variation is (Orr 2001)

$$P_{fix}^{sv} = \int_0^1 \pi(x)\Psi(x)dx. \tag{10}$$

361   **Adaptation in a declining population**

362　Suppose the maximum number of offspring an individual can produce is W_0, the absolute fitness

363　of aa individuals after pollen limitation happens is

$$W = W_0\big((1-d)r_0 + (1-r_0)(1-m)\big) = W_0\big((1-m) + (m-d)r_0\big). \tag{11}$$

365　The absolute fitness of AA individuals is

$$W' = W_0\big((1-d)(r_0+r) + (1-r_0-r)(1-m)\big) = W\left(1 - \frac{d_e}{1-d_e r_0}r\right). \tag{12}$$

367　Therefore, for the evolution of selfing to rescue the population ($W' > 1$), it requires $r > (1 -$

368　$d_e r_0)(1 - W^{-1})/d_e$. Denote the rate of demographic decline as $\lambda = 1 - W$, the fixation

369　probability from a single mutant is reduced to (Glemin and Ronfort 2013)

$$P_{fix}^{new*} \approx 2\frac{N_e}{N}\big((h + F - hF)R - \lambda\big). \tag{13}$$

371　We assume the strength of pollen limitation is independent of the population size, so the

372　probability for a population starting from an initial size N_0 to be rescued by new mutations is

373　approximately (Orr and Unckless 2008)

$$P_{res}^{new} = 1 - \exp\left[-\frac{2N_0 v_1 P_{fix}^{new*}}{\lambda}\right]. \tag{14}$$

375　To obtain the critical dominance at which the fixation probability is the same for outcrossers and

376　selfers, note that the rescue probability from new mutations in a completely outcrossing ($r_0 = 0$)

377　and a nearly completely selfing ($r_0 = 1 - r$) population are

$$P_{res}^{new}(r_0 = 0) \approx 1 - \exp\left[2N_0 v_1\left(2 - \frac{(1 - c - 2d_e)hr}{\lambda}\right)\right], \tag{15a}$$

$$P_{res}^{new}(r_0 = 1 - r) \approx 1 - \exp\left[2N_0 v_1\left(1 - \frac{(2(1 - 2d_e) - 3\lambda)r}{4(1 - d_e)\lambda}\right)\right]. \tag{15b}$$

380　Therefore, the critical dominance is

$$h_{c2} = h_c + \frac{\lambda}{1 - c - 2d_e}\left(\frac{1}{r} - \frac{3}{4(1 - d_e)}\right). \tag{16}$$

382　Since r is assumed to be small and $d_e < 0.5$, the second term is usually positive, and thus $h_{c2} >$

383　h_c. The probability for the population to be rescued by standing variation is approximately

$$384 \qquad P_{res}^{sv} = \int_0^1 \left(1 - P_{fix}^{new*}\right)^{2N_0 x} \Psi(x) dx. \qquad (17)$$

385 The population will go to extinction when alleles from both standing variation and new mutations

386 cannot fix, so the overall rescue probability is $P_{res} = 1 - (1 - P_{res}^{sv})(1 - P_{res}^{new})$, and the relative

387 contribution to rescue from standing variation is $\Pi_{sv} = P_{res}^{sv}/P_{res}$.

Adaptation after a bottleneck

390 Here we consider a population with an initial size N_0 that suffers a bottleneck, so that the

391 population size is suddenly reduced to N_c. Suppose the allele frequency in the ancestral

392 population is x, given that $N_0 \gg N_c$, the probability that the allele frequency after the bottleneck

393 x' is approximately a binomial distribution. Therefore, the expected fixation probability from

394 standing variation is approximately

$$395 \qquad P_{fix}^{sv*} = \int_0^{2N_c} \pi(k/2N_c)\binom{2N_c}{k} dk \int_0^1 x^2 (1-x)^2 \psi(x) dx. \qquad (18)$$

Adaptation from multiple loci

398 For a certain overall increase of the selfing rate Δr, I suppose there are k identical modifier loci,

399 so the effect of each modifier is $r = \Delta r/n$ when modifiers act additively. For adaptation from

400 new mutations, if the waiting time for a mutation to happen is longer than the time for a modifier

401 to sweep to high frequency, i.e., $\frac{1}{2Nv_1} > 4N$. In this case, modifiers fix successively, and the time

402 for all the modifiers to fix is

$$403 \qquad T_{fix(multi)}^{new} \approx \sum_{k=n}^{k=1} \frac{1}{2Nv_1 P_{fix}^{new}\big((r_0 + (n-k)r)\big)}. \qquad (19)$$

Fig. S7 shows that $T_{fix(multi)}^{new}$ increases with n. For rescue of a declining population, if fixation of

a single modifier is enough to stop demographic decline, the probability of rescue from multiple

loci is still given by equation (14), with v_1 replaced by nv_1. Therefore, the effect of genetic

architecture on rescue can be assessed from $nP_{fix}^{new*}(\Delta r/n)$, which decreases with n.

Incorporating the genetic basis of inbreeding depression

It has been firmly established that inbreeding depression is mainly contributed to by two

categories of deleterious mutations (Charlesworth and Willis 2009): slightly deleterious, partially

dominant mutations, and large-effect, highly recessive lethal and sublethal mutations. Slightly

deleterious, partially dominant mutations also cause background selection (Charlesworth 2012).

I denote the inbreeding depression contributed by highly recessive large-effect mutations as d_0,

and previous models (Lande et al. 1994) show that d_0 decreases slowly as the selfing rate

increases. Therefore, here I assume d_0 is the same for populations with different selfing rates,

and I only consider the influence of the selfing rate on inbreeding depression contributed by

slightly deleterious mutations. I denote the haploid genomic mutation rate as U, and assume all

deleterious mutations are identical with selection coefficient s and dominance h_d. For a

population at the equilibrium state, the inbreeding depression can be approximated by (Roze

2015)

$$d = 1 - (1 - d_0)\left(1 + \frac{I_2}{4}\right)\exp\left[-U\frac{1 - 2h_d}{2h_d + r_0(1 - 2h_d)}(1 + I_1) + \frac{r_0}{2 - r_0}I_2\right], \tag{20}$$

423   where

$$I_1 = 2U(1 - h)(1 - 2h)\frac{2 + r_0}{2 - r_0}T, \tag{21a}$$

$$I_2 = U^2(1 - 2h_d)^2 T, \tag{21b}$$

$$T = \frac{2r_0(1 - r_0)}{(4 - r_0)[2h_d + r_0(1 - 2h_d)]^2}. \tag{21c}$$

427   The effect of the background selection on reducing the effective population size (Nordborg et al.

428   1996) is

429
$$\alpha \approx \exp\left[-\frac{2U}{(h_d + F_d - h_d F_d)s + L(1 - F_d)}\right],$$
(22)

430 where L is the number of crossovers along the chromosomes and $F_d = r_0/(2 - r_0)$. The fixation

431 probability can be calculated by replacing N_e with αN_e in equations (5) and (10).

Simulations

Simulations are written in R 4.0.2 and the code is archived on Dryad [filed upon acceptance].

Every generation, reproduction occurs first based on the individual selfing rate, followed by

selection. The next generation is drawn from a multinomial distribution based on the expected

genotype frequencies after selection. For adaptation from standing variation, I run $6N_0$

generations to reach the selection-drift-mutation equilibrium (Hermisson and Pennings 2005)

with forward and backward mutations, and after pollen limitation happens, the input of mutations

stops. For adaptation in a declining population, the population size in the next generation is

drawn from a Poisson distribution with the mean being the product of the current population size

and the average fitness. The population is considered as having gone to extinction when $N = 0$,

and is considered as rescued when the modifier allele goes to fixation or the population size

exceeds $10N_0$.

References

1. Aguilar, R., Ashworth, L., Galetto, L., & Aizen, M. A. (2006). Plant reproductive susceptibility
to habitat fragmentation: review and synthesis through a meta-analysis. Ecology letters, 9(8),
968-980.

2. Asker, S., & Jerling, L. (1992). Apomixis in plants. CRC press.

3. Baker, H. G. (1955). Self-compatibility and establishment after 'long-distance' dispersal.
Evolution, 9(3), 347-349.

4. Barrett, S. C., Colautti, R. I., & Eckert, C. G. (2008). Plant reproductive systems and evolution
during biological invasion. Molecular ecology, 17(1), 373-383.

5. Barrett, R. D., & Schluter, D. (2008). Adaptation from standing genetic variation. Trends in
ecology & evolution, 23(1), 38-44.

6. Bell, G. (2017). Evolutionary rescue. Annual Review of Ecology, Evolution, and Systematics,
48, 605-627.

7. Bodbyl Roels, S. A., & Kelly, J. K. (2011). Rapid evolution caused by pollinator loss in
Mimulus guttatus. Evolution: International Journal of Organic Evolution, 65(9), 2541-2552.

8. Busch, J. W., & Schoen, D. J. (2008). The evolution of self-incompatibility when mates are
limiting. Trends in plant science, 13(3), 128-136.

9. Caballero, A., & Hill, W. G. (1992). Effective size of nonrandom mating populations. Genetics,
130(4), 909-916.

10. Charlesworth, B., M. T. Morgan, and D. Charlesworth. 1993a. The effect of deleterious
mutations on neutral molecular variation. Genetics.134:1289–1303.

11. Charlesworth, D., & Willis, J. H. (2009). The genetics of inbreeding depression. Nature
reviews genetics, 10(11), 783-796.

12. Cheptou, P. O. (2012). Clarifying Baker's law. Annals of botany, 109(3), 633-641.

13. Cheptou, P. O. (2019). Does the evolution of self-fertilization rescue populations or increase
the risk of extinction?. Annals of Botany, 123(2), 337-345.

14. Eckert, C. G., Kalisz, S., Geber, M. A., Sargent, R., Elle, E., Cheptou, P. O., ... & Porcher, E.
(2010). Plant mating systems in a changing world. Trends in Ecology & Evolution, 25(1), 35-43.

15. Eckert, C. G., Samis, K. E., & Dart, S. (2006). Reproductive assurance and the evolution of
uniparental reproduction in flowering plants. Ecology and evolution of flowers, 183, 203.

Ewens, W. J. (2012). Mathematical population genetics 1: theoretical introduction (Vol. 27).
Springer Science & Business Media.

Fisher, R. A. (1941). Average excess and average effect of a gene substitution. Annals of
Eugenics, 11(1), 53-63.

Fishman, L., Beardsley, P. M., Stathos, A., Williams, C. F., & Hill, J. P. (2015). The genetic
architecture of traits associated with the evolution of self-pollination in Mimulus. New
Phytologist, 205(2), 907-917.

Fishman, L., Kelly, A. J., & Willis, J. H. (2002). Minor quantitative trait loci underlie floral traits
associated with mating system divergence in Mimulus. Evolution, 56(11), 2138-2155.

Glémin, S., & Ronfort, J. (2013). Adaptation and maladaptation in selfing and outcrossing
species: new mutations versus standing variation. Evolution: International Journal of Organic
Evolution, 67(1), 225-240.

Goodwillie, C., Ritland, C., & Ritland, K. (2006). The genetic basis of floral traits associated with
mating system evolution in Leptosiphon (Polemoniaceae): an analysis of quantitative trait loci.
Evolution, 60(3), 491-504.

Goodwillie, C., Sargent, R. D., Eckert, C. G., Elle, E., Geber, M. A., Johnston, M. O., ... & Winn,
A. A. (2010). Correlated evolution of mating system and floral display traits in flowering plants
and its implications for the distribution of mating system variation. New Phytologist, 185(1), 311-
321.

Haldane, J. B. S. (1927). A mathematical theory of natural and artificial selection, part V:
selection and mutation. In Mathematical Proceedings of the Cambridge Philosophical Society
(Vol. 23, No. 7, pp. 838-844). Cambridge University Press.

Harder, L. D., & Wilson, W. G. (1998). A clarification of pollen discounting and its joint effects
with inbreeding depression on mating system evolution. The American Naturalist, 152(5), 684-
695.

Hermisson, J., & Pennings, P. S. (2005). Soft sweeps: molecular population genetics of
adaptation from standing genetic variation. Genetics, 169(4), 2335-2352.

Holsinger, K. E. (2000). Reproductive systems and evolution in vascular plants. Proceedings of
the National Academy of Sciences, 97(13), 7037-7042.

Igic, B., Lande, R., & Kohn, J. R. (2008). Loss of self-incompatibility and its evolutionary
consequences. International Journal of Plant Sciences, 169(1), 93-104.

Igic, B., & Busch, J. W. (2013). Is self-fertilization an evolutionary dead end?. New Phytologist,
198(2), 386-397.

Karron, J. D., Jackson, R. T., Thumser, N. N., & Schlicht, S. L. (1997). Outcrossing rates of
individual Mimulus ringens genets are correlated with anther–stigma separation. Heredity, 79(4),
365-370.

Kelly, J. K., & Willis, J. H. (2001). Deleterious mutations and genetic variation for flower size in
Mimulus guttatus. Evolution, 55(5), 937-942.

Kimura, M. (1962). On the probability of fixation of mutant genes in a population. Genetics,
47(6), 713.

Knight, T. M., Steets, J. A., Vamosi, J. C., Mazer, S. J., Burd, M., Campbell, D. R., ... & Ashman,
T. L. (2005). Pollen limitation of plant reproduction: pattern and process. Annu. Rev. Ecol. Evol.
Syst., 36, 467-497.

Lai, Y. T., Yeung, C. K., Omland, K. E., Pang, E. L., Hao, Y., Liao, B. Y., ... & Hung, H. Y.
(2019). Standing genetic variation as the predominant source for adaptation of a
songbird. *Proceedings of the National Academy of Sciences*, *116*(6), 2152-2157.

Lande, R., & Schemske, D. W. (1985). The evolution of self-fertilization and inbreeding
depression in plants. I. Genetic models. Evolution, 39(1), 24-40.

Lande, R., Schemske, D. W., & Schultz, S. T. (1994). High inbreeding depression, selective
interference among loci, and the threshold selfing rate for purging recessive lethal mutations.
Evolution, 48(4), 965-978.

Larson, B. M., & Barrett, S. C. (2000). A comparative analysis of pollen limitation in flowering
plants. Biological journal of the Linnean Society, 69(4), 503-520.

Levin, D. A. (2012). Mating system shifts on the trailing edge. Annals of Botany, 109(3), 613-
620.

Miller, J. S., Blank, C. M., & Levin, R. A. (2019). Colonization, Baker's law, and the evolution of
gynodioecy in Hawaii: implications from a study of Lycium carolinianum. American journal of
botany, 106(5), 733-743.

Mogie, M. 1992. The Evolution of Asexual Reproduction in Plants. Chapman & Hall, London.

Nordborg, M., Charlesworth, B., & Charlesworth, D. (1996). The effect of recombination on
background selection. Genetics Research, 67(2), 159-174.

Orr, H. A. (2005). The genetic theory of adaptation: a brief history. Nature Reviews Genetics,
6(2), 119-127.

Orr, H. A. (2010). The population genetics of beneficial mutations. Philosophical Transactions of
the Royal Society B: Biological Sciences, 365(1544), 1195-1201.

Orr, H. A., & Betancourt, A. J. (2001). Haldane's sieve and adaptation from the standing genetic
variation. Genetics, 157(2), 875-884.

Orr, H. A., & Unckless, R. L. (2008). Population extinction and the genetics of adaptation. The
American Naturalist, 172(2), 160-169.

Pannell, J. R. (2015). Evolution of the mating system in colonizing plants. Molecular Ecology,
24(9), 2018-2037.

Pannell, J. R., Auld, J. R., Brandvain, Y., Burd, M., Busch, J. W., Cheptou, P. O., ... & Hovick, S.
M. (2015). The scope of Baker's law. New Phytologist, 208(3), 656-667.

Porcher, E., & Lande, R. (2005). The evolution of self-fertilization and inbreeding depression
under pollen discounting and pollen limitation. Journal of evolutionary biology, 18(3), 497-508.

Ronfort, J., & Glemin, S. (2013). Mating system, Haldane's sieve, and the domestication
process. Evolution, 67(5), 1518-1526.

Roze, D. (2015). Effects of interference between selected loci on the mutation load, inbreeding
depression, and heterosis. Genetics, 201(2), 745-757.

Sargent, R. D., Goodwillie, C., Kalisz, S., & Ree, R. H. (2007). Phylogenetic evidence for a
flower size and number trade-off. American Journal of Botany, 94(12), 2059-2062.

Sicard, A., & Lenhard, M. (2011). The selfing syndrome: a model for studying the genetic and
evolutionary basis of morphological adaptation in plants. Annals of botany, 107(9), 1433-1443.

Spigler, R. B., & Ashman, T. L. (2012). Gynodioecy to dioecy: are we there yet?. Annals of
Botany, 109(3), 531-543.

Strandh, M., Jönsson, J., Madjidian, J. A., Hansson, B., & Lankinen, Å. (2017). Natural selection
acts on floral traits associated with selfing rate among populations of mixed-mating Collinsia
heterophylla (Plantaginaceae). International Journal of Plant Sciences, 178(8), 594-606.

Thomann, M., Imbert, E., Devaux, C., & Cheptou, P. O. (2013). Flowering plants under global
pollinator decline. Trends in plant science, 18(7), 353-359.

Table

	Mating-unrelated allele	Selfing modifier allele
Adaptation from new mutations in a constant population	1. Fixation probability is higher for dominant alleles. 2. $h_c = 0.5$	1. Fixation probability is higher for dominant alleles. 2. h_c can vary from 0 to 1, depending on the strength of pollen limitation and pollen discounting.
Adaptation from standing variation in a constant population	1. Fixation probability is nearly independent of dominance. 2. Fixation probability decreases with the selfing rate. 3. Fixation is quickest for dominant alleles.	1. Fixation probability is higher for recessive alleles. 2. Fixation probability is highest at an intermediate selfing rate. 3. Fixation is quickest when alleles are intermediate dominant, while slowest for dominant alleles;
Adaptation in declining populations	1. h_c increases. 2. Rescue is more likely through dominant alleles. 3. Rescue probability from standing variation is lower in selfers	1. h_c increases. 2. Rescue is more likely through dominant alleles. 3. Rescue probability from standing variation is lower in selfers.
Adaptation in populations after bottleneck	1. Fixation probability is insensitive to dominance. 2. Adaptation is less likely in selfers.	1. Recessive alleles have a much higher fixation probability. 2. Adaptation is less likely in selfers.
Genetic architecture of adaptation	N/A	Adaptation from standing variation more likely when selfing is controlled by many small-effect loci, while adaptation from new mutations should be faster and more likely through few large-effect loci.
Under the coevolution of inbreeding depression	N/A	The fixation probability from both new mutations and standing variation is highest at an intermediate selfing rate.

Table 1. Comparison of the genetics of adaptation for mating-unrelated alleles with selfing
modifier alleles. h_c is the critical dominance above which fixation probability is lower in selfers
than outcrossers.

Figures

Figure 1. Effects of pollen discounting c and effective inbreeding depression d_e on the value of
critical dominance h_c. Results are calculated based on equation (4), but the value is set to be 1
if $h_c > 1$. The grey area is where for all levels of dominance, the selfing modifier is selected
against in a completely outcrossing population and thus cannot fix, but favored in a highly
selfing population (see equation (8)).

Figure 2. Effects of background selfing rate on fixation probability from standing variation in a constant population. Lines are model predictions from equation (10), and dots are simulation results. For all panels, $r = 0.01, d = 0.6, N = 20000, v_1 = v_2 = 10^{-6}$.

Figure 3. Effects of the background selfing rate on the rescue probability in declining

populations (panels (a), (b)) and populations suffering a bottleneck (panel (c)). Lines in panels

(a), (b) and (c) are model predictions from equation (14), (17) and (18), respectively, and dots

are simulation results. For panels (a) and (b), $r = 0.05, d = 0.6, d_e = -0.3, c = 0, N_0 = 20000, \lambda =$

$0.002, v_1 = v_2 = 10^{-6}$. For panel (c), $r = 0.01, d = 0.6, d_e = -0.3, N_0 = 20000, N_c = 500, v_1 =$

$v_2 = 10^{-6}$.

Figure 4. Effects of the background selfing rate on the population survival probability from a new mutation (panel (a)) and standing variations (panel (b)) when incorporating the genetic basis of inbreeding depression and background selection. Parameters related to deleterious mutations are $U = 0.4, h_d = 0.2, s = 0.1, d_0 = 0.4, L = 10$; and other parameters are $r = 0.01, c = 0, m = 0.5, N = 20000, v_1 = v_2 = 10^{-6}$.