1

The relationship between affective visual mismatch negativity and interpersonal difficulties

across autism and schizotypal traits

Talitha C. Ford*[1,2], Laila E. Hugrass[2,3], & Bradley N. Jack[4]

[1] Cognitive Neuroscience Unit, School of Psychology, Deakin University, Geelong, Victoria,

Australia

[2] Centre for Human Psychopharmacology, Faculty of Heath, Arts and Design, Swinburne

University of Technology, Melbourne, Victoria, Australia

[3] Department of Psychology and Counselling, School of Psychology and Public Health, La

Trobe University, Melbourne, VIC, Australia

[4] Research School of Psychology, The Australian National University, Canberra, Australia

* Corresponding Author

Dr Talitha C. Ford

Cognitive Neuroscience Unit, School of Psychology, Deakin University

221 Burwood Hwy, Burwood, Victoria 3125, Australia

Email: talitha.ford@deakin.edu.au

Phone: +61 3 9246 8730

25
26
27

Highlights

- The visual Mismatch Negativity (vMMN) is larger in response to happy compared to sad or neutral facial expressions
- The vMMN to happy facial expressions is specifically associated with more social communication difficulties
- vMMN to positive affect may be a neural mechanism associated with social communication difficulties in autism and schizophrenia

Abstract

Objective: Sensory deficits are a feature of autism and schizophrenia, as well as the upper end of their non-clinical spectra. The mismatch negativity (MMN), an index of pre-attentive auditory processing, is particularly sensitive in detecting such deficits; however, little is known about the relationship between the visual MMN (vMMN) to emotional faces and autism and schizophrenia spectrum symptom domains.

Methods: We probed the vMMN to happy, sad, and neutral faces in 77 young adults (18-40 years, 41 female), and evaluated their degree of autism and schizophrenia spectrum traits using the Autism Spectrum Quotient (AQ) and Schizotypal Personality Questionnaire (SPQ).

Results: The vMMN to happy faces was significantly larger than the vMMNs to sad and neutral faces ($p > 0.05$). The vMMN to happy faces was associated with interpersonal difficulties as indexed by AQ Communication and Attention to Detail subscales, and SPQ Interpersonal Features ($p < 0.05$, *uncorrected*), such that a larger vMMN was associated with more interpersonal difficulties.

Conclusions: Pre-attentive processing of positive affect might be more specific to the interpersonal features associated with autism and schizophrenia.

Significance: These findings add valuable insights into the growing body of literature investigating symptom-specific neurobiological markers of autism and schizophrenia spectrum conditions.

Keywords: visual mismatch negativity, electroencephalography, autism, schizotypy, facial emotion processing.

63 **The relationship between affective visual mismatch negativity to interpersonal difficulties**
64 **across autism and schizotypal traits.**
65
Several lines of evidence point to poor sensory processing in autism and schizophrenia
spectrum disorders, particularly pre-attentive processing of visual and auditory information
(Jahshan et al., 2012; Lanillos et al., 2020; Randeniya et al., 2018; Thye et al., 2018). The
mismatch negativity (MMN) is a well-established marker of pre-attentive processing of
auditory information. To elicit a MMN, typically the repeated presentation of a predictable
"standard" stimulus is interrupted at random intervals by an unpredictable "deviant"
stimulus, and the MMN represents the difference in the neural response to the standard
and deviant stimuli (Naatanen et al., 2007, 2012). Several theories attempt to explain the
MMN, such as neural fatigue, neural population sharpening, facilitation of stimulus
processing, and cortical predictive coding (see Kremláček et al., 2016; Naatanen et al., 2007;
Stefanics et al., 2014). The most popular explanation for the MMN is the predictive coding
theory, which proposes that the brain generates a probabilistic model of the sensory input
(i.e., the standard stimuli), which explains the attenuation of the neural response to the
standard over time. The deviant stimulus is a violation of the probabilistic model, and hence
results in a larger neural response (Friston, 2005).
81
Deficits in the auditory MMN have been consistently replicated across the schizophrenia
spectrum (Naatanen et al., 2011; Umbricht & Krljes, 2005) – from chronic illness to high-risk
groups (Fulham et al., 2014; Jahshan et al., 2012; Lavoie et al., 2018). Deficits in the auditory
MMN in autism are less well established, however, with inconsistencies between studies
attributed to the heterogeneity of samples investigated (Schwartz et al., 2018).
Nevertheless, recent meta-analyses reported generally reduced auditory MMN in children
with ASD, but not adults (Chen et al., 2020; Schwartz et al., 2018). Importantly, the auditory
MMN has been associated with poorer social/interpersonal functioning in both autism (Fan
& Cheng, 2014) and schizophrenia (Featherstone et al., 2017; Lee et al., 2014), as well as in
non-clinical groups (Ford et al., 2017).
92
Far fewer studies have investigated whether the visual counterpart of the auditory MMN,
the vMMN, is similarly affected in autism and schizophrenia. In schizophrenia studies, a
smaller vMMN has been reported in response to changes on motion direction (Urban et al.,
2008), letters (Neuhaus et al., 2013), happy and fearful facial expressions (Csukly et al.,
2013; Vogel et al., 2018), and sequence of stimulus presentation (Vogel et al., 2018), but not
to changes in Gabor patch orientation (Farkas et al., 2015). The vMMN amplitude to motion
direction has also been associated with negative symptom severity (Urban et al., 2008), and
the vMMN to happy deviants was associated with better facial emotion recognition (Csukly
et al., 2013).
102

Studies investigating the vMMN in autism have reported reduced vMMN amplitude in
adults (Cléry, Roux, et al., 2013) and longer vMMN latency to object shape deformation in
children with ASD (Cléry, Bonnet-Brilhault, et al., 2013). No difference in vMMN amplitude
or latency was reported for windmill pattern deviants in a small sample of adults (n = 11)
with ASD (Maekawa et al., 2011). One study investigating the relationship between facial
emotion vMMN and autistic traits reported a reduced vMMN to happy but not sad deviants
that was associated with higher overall levels of autistic traits, as quantified using the
autism spectrum quotient (AQ) (Gayle et al., 2012). These findings, along with those
demonstrating that the vMMN to happy deviants is associated with better facial emotion
recognition (Csukly et al., 2013), suggest that the vMMN elicited by changes in emotional
expression might index affective reactivity and social competency (Gayle et al., 2012).
114
Although there is some evidence to suggest that pre-attentive facial emotion processing is
associated with autism and schizophrenia spectrum symptomatology, the specificity of the
vMMN differences to autism and schizophrenia spectrum symptom dimensions remains
unknown. This study, therefore, extends that of Gayle et al. (2012) by investigating the
extent to which pre-attentive processing of changes in facial emotions are associated with
specific autism and schizophrenia spectrum traits. It was predicted that a smaller vMMN to
happy face deviants would be associated with higher autism and schizophrenia trait
severity, particularly in the social domain. The extent to which vMMN to happy and sad face
deviants predict the positive and disorganised schizotypy dimensions and attention and
imagination dimensions of autism were explored.
125
126   **Methods**
127   Ethical approval for this study was granted by the Swinburne University Human Research
128   Ethics Committee (2016/092) in accordance with the Declaration of Helsinki. All participants
129   provided written informed consent prior to commencing the study.
130
Participants
A total of 77 (41 female, 36 male) health adults aged 18-40 years participated in this study
(age mean (*SE*): total = 24.96 (0.58), male = 25.39 (0.86), female = 25.59 (0.78)). All
participants had normal or corrected to normal vision, and were free from psychotropic
medications and psychiatric illness, except for one male participant taking an SSRI for mild
depression who was excluded from analyses. One male participant was removed due to
incomplete psychometric data, and an additional 15 were excluded due to poor EEG data (9
female, 6 male). See Table 1 for final sample participant characteristics, including Mann-
Whitney *U* non-parametric tests to identify sex differences in continuous variable, and χ^2 to
identify sex differences in handedness. Males scored slightly higher than females on AQ
Attention Switching and SPQ Disorganisation dimensions ($p < 0.05$).
142
143

Measures

The full 50-item autism spectrum quotient (AQ) was used to measure the five autism trait dimensions of Social Skill, Communication, Attention Switching, Attention to Detail and Imagination (Baron-Cohen et al., 2001). The 74-item schizotypal personality questionnaire (SPQ) was used to measure schizophrenia domains of Cognitive-Perceptual Features, Interpersonal Features and Disorganised Features (Raine, 1991). AQ and SPQ items were pseudo-randomised and presented on a 4-point Likert scale from 1 (*strongly disagree*) to 4 (*strongly agree*) in order to improve reliability (Ford & Crewther, 2014; Wuthrich & Bates, 2005). Items were scaled such that *strongly agree* and *agree* = 1, and *strongly disagree* and *disagree* = 0. All statistical analyses were replicated using the full scale scoring system, which retained 1-4 scores for each item, and is presented in the supplementary material. Participants trait depression, anxiety and stress scores were collected using the DASS-21 (Lovibond & Lovibond, 1995), in order to control for these effects on the relationships between AQ and SPQ domains and the vMMN.

Stimuli

Face stimuli were selected from the Nimstim database of standardised facial expressions (Tottenham et al., 2009). Five grey-scale female faces were selected; we used female faces only to remove gender bias toward faces. All faces were of caucasian appearance, and the happy, sad and neutral emotions for each face was included. We used the SHINE toolbox for Matlab to match faces for mean luminance (Willenbockel et al., 2010).

EEG procedure

Participants were seated 57 cm from the stimulus presentation computer screen (inside an electrically shielded room). Participants were instructed to focus on a fixation cross (i.e., "+") in the centre of the screen that randomly alternated in colour between red and green, and to respond as quickly as possible via keypress when the cross changed colour. The face stimuli for the vMMN task were presented directly behind the fixation cross during the colour-change distractor task.

The vMMN task consisted of 6 blocks containing a total of 750 trials (600 standards, 75 deviants, 75 oddballs), with each facial emotion serving as standard, deviant and oddball across the blocks (e.g., neutral standard - happy deviant - sad deviant, neutral standard - sad deviant - happy deviant, happy standard - neutral deviant - sad deviant, etc.). Each face identity was presented an equal number of times in each block (i.e. 120 presentations of each face as standard emotion, and 15 presentations of each face as deviant emotion). Block order was counterbalanced across participants using a randomised balanced Latin-square. Face stimulus presentation was pseudo-randomised, ensuring that the same identity was not shown on consecutive trials, and that standard emotions were presented for the first three and final two trials of each block, and at least two standard emotions were presented between each deviant emotion. Face stimuli were presented for 400 ms,

with an inter-stimulus interval of 150 ms and a jitter of ± 50 ms. The participant's task was to attend to a fixation cross in the centre of the face stimuli, and to press one of two keys when the colour of the fixation cross changed from red to green, or vice versa (see Figure 1). This task ensured that participants were looking at the face stimuli, but were not attending to it. Analyses of the key press data confirmed that the distractor task was performed to a satisfactory level throughout the vMMN recording, with all participants responding within two seconds on at least 80% of trials (M = 97.93%, SD = 2.75%, M_{RT}= 650ms SD_{RT} = 99ms).

ERP pre-processing

The EEG data were pre-processed in BrainVision Analyzer, Version 2.1 (Brain Products GmBH, Germany). We re-referenced the data offline to the average of all scalp electrodes, and filtered the data using a half-amplitude 0.1 to 30 Hz phase-shift free Butterworth filter (24 dB/Oct slope). EEG data were epoched from -100 to 500 ms in relation to face stimulus onset, and epochs were baseline-corrected to their mean voltage from -100 to 0 ms. Epochs with signals exceeding peak-to-peak amplitudes of 100 microvolts at EEG or EOG channel were excluded. This resulted in the rejection of 14% of all trials. All standard trials immediately following a deviant were excluded from analyses.

ERPs for each emotion (happy, neutral, sad) and for each context (standard, deviant) were computed from occipital electrodes (PO7, O1, Oz, O2, PO8). The time-window for the vMMN was identified using a collapsed localiser approach (Luck & Gaspelin, 2017). All standards and deviants were collapsed across all electrodes and facial emotions, and a difference wave (vMMN) was calculated by subtracting the collapsed standards from the collapsed deviants. The vMMN was then identified as the negative peak between 150 ms and 300 ms, which occurred at 237 ms. The vMMN time-window was then determined as the 100 ms around the vMMN peak; i.e., 187-287 ms. This time-window was used to analyse the uncollapsed vMMNs at each electrode, which were calculated by substracting the mean standard ampitude from the mean deviant amplitude for each facial emotion (i.e., happy vMMN = happy deviant – happy standard).

Statistical analysis

An *a priori* power analysis conducted using the pwr package in R (Champely, 2018) indicated that 63 participants were required to detect a moderate effect (*r* =.343, based on Gayle et al. [2012]) with 80% power using correlations with alpha = .05.

A 5 (electrode: PO7, O1, Oz, PO8, O2) x 3 (emotion: happy, sad, neutral) linear mixed effects analysis was conducted to investigate vMMN differences across electrodes and emotions, with subject entered as a random factor. To investigate the extent to which the happy, sad, and neutral vMMN at each electrode were related to autism and schizophrenia trait domains, non-parametric Spearman rank order correlations were calculated. We calculated non-parametric correlation coefficients due to significantly skewed distributions of most

226 trait domains and vMMN values. Trait depression, anxiety and stress were not related to the
227 happy or sad vMMN at any electrode site, thus were not added as covariates in the
228 analyses. Family-wise error was corrected for by applying false discovery rate (FDR;
229 Benjamini & Hochberg, 1995). All statistical analyses were conducted using Jamovi version
230 1.2 (*Jamovi*, 2020).
231
232 **Results**
233 The grand mean average vMMN ERP for each facial emotion at each visual electrode are
234 shown in Figure 2, and the electrode x emotion mixed effects plot of mean vMMN is shown
235 in Figure 3. There was a clear negativity between 150 ms and 300 ms for both happy and
236 sad facial expressions, with the happy vMMN appearing to peak earlier than the sad vMMN.
237
The linear mixed effects analysis (electrode x emotion) revealed a significant main effect for
emotion ($F(2, 824.11) = 8.10$, $p < 0.001$), with the happy vMMN significantly larger than the
neutral vMMN ($t(824.08) = -3.99$, $p < 0.001$) and sad vMMN ($t(824.08) = -2.48$, $p = 0.013$).
There was no difference between sad and neutral vMMN ($p = 0.132$). No significant
hemisphere main effect or emotion x hemisphere interactions was found (ps < 0.1; Figure
3).
244
Spearman rank order correlations investigating the relationship between the vMMN to
happy, sad, and neutral facial expressions found significant moderate negative correlations
between the happy vMMN at O1, Oz and O2 and social autism and schizophrenia spectrum
domains ($p < 0.05$; see Table 2), indicating that those with more social and communication
difficulties and more attention to detail exhibit a larger (more negative) vMMN at O1 and
O2 (Figure 4); however, these correlations did not survive correction for FDR. There were no
relationships between sad and neutral vMMN amplitude at any electrode site and any of
the AQ or SPQ domains ($p > 0.05$).
253
In order to compare the current data with previous results (Gayle et al., 2012), we re-ran
the AQ and SPQ correlations with the vMMN calculated for happy and sad deviants relative
to the neutral standards (Table 3). Similar to the original analyses (i.e., when the vMMN was
calculated for the same emotion when it was deviant relative to when it was the standard;
see Table 2), the communication domain of AQ and the total SPQ score were negatively
correlated with the vMMN at O2 ($p < .05$). There were no other significant correlations.
260
Discussion
This study is the first to investigate the relationship between emotion-related vMMN
amplitudes and specific autism and schizophrenia spectrum traits within the non-clinical
population. This work extends from Gayle et al. (2012) who reported an association
between the emotion-related vMMN and total AQ scores only. As expected, there were
strong bilateral occipitotemporal vMMN responses to the happy and sad expressions.

Although vMMN amplitudes were not significantly correlated with the total AQ score, we
found that people with higher levels of interpersonal difficulty specifically (as indexed by the
AQ Communication and Attention to Detail, and SPQ Interpersonal Features subscales)
tended to exhibit stronger vMMN responses to happy expressions. Despite the large sample
size, however, none of these correlations survived correction for family-wise error. Given
the theoretical importance of an association between autistic traits and vMMN amplitudes,
it is necessary to consider why the correlations between autistic personality traits and the
vMMN are in the opposite direction to those previously reported.
275
These data reveal a weak-moderate relationship between interpersonal difficulties and the
vMMN to happy faces in the opposite direction to Gayle et al. (2012). A notable
methodological difference between the two studies is that the previous study calculated the
vMMN responses to happy and sad deviants relative to neutral standards. Hence,
differences in the morphology of the neural response to the standard and deviant might
reflect differences in response to the specific stimulus being presented (i.e., a sad face
versus a neutral face), as opposed to a neural response to prediction violation. It has been
shown that within the vMMN time-window, evoked responses to emotional faces tend to
be lower amplitude for people with ASD or high levels of autistic traits than for people with
low levels of autistic traits (Csukly et al., 2013; Gayle et al., 2012). Hence, Gayle et al.'s
results may have reflected a correlation between AQ and the evoked responses to facial
emotional, rather than a correlation between AQ and responses to emotion prediction
error. However, our subsequent analyses showed that regardless of whether we compared
deviants and standards for the same emotion (i.e., happy deviant vs happy standard), or
compared emotional deviants with neutral standards, correlations between the happy
vMMN amplitude and autistic and schizotypal personality traits tended to be negative. Thus,
our results indicate that predictive error responses to unexpected happy faces tend to be
stronger for people with higher levels of interpersonal difficulties, as indexed by the
subscales of the AQ and SPQ.
295
Although the increase in vMMN to happy deviants (compared to happy standards) with
more social and interpersonal difficulties was unexpected, previous studies of the auditory
vMMN in Asperger's Syndrome (AS) or ASD have also reported conflicting findings. For
instance, Lepisto et al. (2007) found that adults with AS tended to exhibit enhanced MMNs
to speech sounds, but not for non-speech sounds, suggesting an auditory hypersensitivity or
filtering difficulty in AS (Lepistö et al., 2007). Furthermore, Schwartz and colleagues' (2018)
meta-analysis found that while the auditory MMN was generally reduced in children with
ASD, adults with ASD elicit a larger MMN to speech deviants compared to controls.
However, the meta-analysis highlighted that many studies of the auditory MMN had small
sample sizes and did not properly counterbalance their stimuli.
306

We observed moderate relationships between autistic and schizotypal traits and emotion-
related vMMN amplitudes for happy facial expressions (uncorrected), but not for sad or
neutral expressions. This finding is consistent with Gayle et al., who reported that the total
AQ score was selectively related to vMMN amplitude in response to happy expressions but
are surprising in relation to evidence that people with ASD exhibit more general deficits in
affective processing (Eack et al., 2015; Fan & Cheng, 2014; Wright et al., 2008). It could be
that different aspects of affective processing have different relationships with autistic traits;
for instance, Wilbarger et al. (2009) found that people with ASD exhibit an atypical startle
response to positively valanced faces, despite showing typical implicit responses with
automatic facial mimicry and overt valence ratings. This indicates a disruption in an early
stage of affective processing, such that there is a dissociation between higher-order
assignment of valence and basic automatic defensive responses. The authors suggest that
this ambiguity may mean that positive stimuli are experienced as unreliable or potentially
harmful, which could in turn lead to interpersonal difficulties. The current results are
consistent with the idea that people with higher self-report levels of interpersonal difficulty
tend to exhibit a heightened automatic alerting response to unexpected emotions.
323
The vMMN to happy faces over O1 and O2 was also shown to increase with more AQ
Attention to Detail (uncorrected), which is somewhat intuitive given better preattentive
discrimination between emotions could lend itself to more superior attention to detail in
the social environment. Again, better discrimination of happy deviants among neutral and
sad standard faces may not necessarily translate to correct interpretation of facial
expressions. Finally, SPQ scores were associated with larger vMMN amplitudes for the
happy facial expressions (uncorrected). This effect was also specific to occipital sites (Oz and
O2) and was driven by weak-moderate relationships across the three schizotypy dimensions
(Cognitive-Perceptual, Interpersonal and Disorganised Features). In contrast, the
relationship between AQ subscales Imagination and Attention Switching and the happy
vMMN were weak across electrode sites, driving down the relationship between AQ total
and the vMMN.
336
Our design allowed the vMMN to be computed as the difference between deviant and
standard of each emotion and removed a particular potential bias toward happy and sad
emotion compared to the neutral emotion. A further strength was the use of grey-scale
faces of four different female actors. The use of female-only stimuli and equal number of
male and female participants recruited in this study removed any potential sex vMMN
(Kecskes-Kovacs et al., 2013), and allowed us to be more confident that the vMMN was due
to emotion prediction error, rather than a sex prediction error. Furthermore, through
investigating the relationship between the vMMN to facial emotions and varying degrees of
specific autism and schizophrenia spectrum symptom traits in a non-clinical sample, these
findings highlight the utility of this population to better understand the neurobiology of
symptoms at a clinical level. Indeed, the relationship between sub-clinical and clinical

auditory MMN profiles has been well established, with high-risk individuals and first-degree
relatives showing a similar MMN deficit, albeit less pronounced, to clinical groups
(Brockhaus-Dumke et al., 2005; Lavoie et al., 2018).
351
Despite the robust research methods and large sample size employed herein, there are
some limitations that should be taken into consideration when interpreting these data.
First, we did not include a non-emotional deviant, such as stimulus tint. Instead, we
calculated the vMMN by subtracting the deviant of each emotion from the standard of the
same emotion (i.e., happy deviant minus happy standard), which ensured the detected
vMMN was not simply a response to the physical characteristics of faces or to the change in
emotional expression. Furthermore, although participants were excluded if they were taking
psychoactive medications, they were not asked to abstain from caffeine, nicotine, alcohol,
cannabis, or other illicit substances prior to testing. Only moderate alcohol consumption
(.04-.07% BAC) has been shown to affect the vMMN (Kenemans et al., 2010), but acute
doses of nicotine, alcohol and cannabis affect the auditory MMN, an thus may also effect
the visual counterpart. Given testing for this study occurred on weekdays, it is likely that
participants were only acutely affected by caffeine, which has been shown not to affect
auditory MMN (Rosburg et al., 2004). Nevertheless, these data should be interpreted with
the above limitations in mind. Finally, due to the large number of statistical tests performed,
none of the correlations survived correction for FDR. Future research should be conducted
on a larger sample, or with more targeted hypotheses, to replicate these findings.
369
This study sought to further explore the findings of Gayle et al. (2012) by investigating the
relationship between the vMMN to happy and sad facial expressions and specific autism and
schizophrenia spectrum traits in a large sample of healthy adults. We addressed the
research design limitations outlined in previous studies and found that communication and
interpersonal difficulties, as well as attention to detail, were associated with a greater
vMMN amplitude in response to happy faces, however, the statistical significance of these
effects diminished following correction for family-wise error. The large early visual response
to happy faces might suggest a hyper-responsivity to the change in emotional expression,
with this hyper-responsivity possibly leading to poorer interpersonal outcomes. In utilising a
non-clinical population, these findings add valuable insights into the growing body of
literature investigating symptom-specific neurobiological markers of autism and
schizophrenia spectrum conditions.
382

Acknowledgements
We thank Swinburne Neuroimaging for use of the EEG facilities, and our third year
Psychophysics students, Tomas Cox, Edenn Baczyk, Jodeci Cowell and Alexandra Suvorova,
for their help with recruitment and data collection. Dr Ford is supported by a Deakin
University Dean's Postdoctoral Research Fellowship.

Declaration of interests: None
390
391   **References**
392

393   Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-
394   Spectrum Quotient (AQ): Evidence from Asperger Syndrome/high-functioning
395   autism, males and females, scientists and mathematicians. *Journal of Autism and*
396   *Developmental Disorders*, *31*(1), 5–17. https://doi.org/10.1023/A:1005653411471
397   Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: A practical and
398   powerful approach to multiple testing. *Journal of the Royal Statistical Society. Series*
399   *B (Methodological)*, 289–300.
400   Brockhaus-Dumke, A., Tendolkar, I., Pukrop, R., Schultze-Lutter, F., Klosterkotter, J., &
401   Ruhrmann, S. (2005). Impaired mismatch negativity generation in prodromal
402   subjects and patients with schizophrenia. *Schizophrenia Research*, *73*(2–3), 297–310.
403   https://doi.org/10.1016/j.schres.2004.05.016
404   Champely, S. (2018). *pwr: Basic Functions for Power Analysis* (R package version 1.2-2)
405   [Computer software]. https://CRAN.R-project.org/package=pwr
406   Chen, T.-C., Hsieh, M. H., Lin, Y.-T., Chan, P.-Y. S., & Cheng, C.-H. (2020). Mismatch negativity
407   to different deviant changes in autism spectrum disorders: A meta-analysis. *Clinical*
408   *Neurophysiology*, *131*(3), 766–777. https://doi.org/10.1016/j.clinph.2019.10.031
409   Cléry, H., Bonnet-Brilhault, F., Lenoir, P., Barthelemy, C., Bruneau, N., & Gomot, M. (2013).
410   Atypical visual change processing in children with autism: An electrophysiological
411   Study: Visual mismatch response in children with autism. *Psychophysiology*, *50*(3),
412   240–252. https://doi.org/10.1111/psyp.12006
413   Cléry, H., Roux, S., Houy-Durand, E., Bonnet-Brilhault, F., Bruneau, N., & Gomot, M. (2013).
414   Electrophysiological evidence of atypical visual change detection in adults with
415   autism. *Frontiers in Human Neuroscience*, *7*.
416   https://doi.org/10.3389/fnhum.2013.00062
417   Csukly, G., Stefanics, G., Komlósi, S., Czigler, I., & Czobor, P. (2013). Emotion-Related Visual
418   Mismatch Responses in Schizophrenia: Impairments and Correlations with Emotion
419   Recognition. *PLoS ONE*, *8*(10), e75444.
420   https://doi.org/10.1371/journal.pone.0075444
421   Eack, S. M., Mazefsky, C. A., & Minshew, N. J. (2015). Misinterpretation of facial expressions
422   of emotion in verbal adults with autism spectrum disorder. *Autism*, *19*(3), 308–315.
423   https://doi.org/10.1177/1362361314520755
424   Fan, Y. T., & Cheng, Y. (2014). Atypical mismatch negativity in response to emotional voices
425   in people with autism spectrum conditions. *PLoS One*, *9*(7), e102471.
426   https://doi.org/10.1371/journal.pone.0102471
427   Farkas, K., Stefanics, G., Marosi, C., & Csukly, G. (2015). Elementary sensory deficits in
428   schizophrenia indexed by impaired visual mismatch negativity. *Schizophrenia*
429   *Research*, *166*(1–3), 164–170. https://doi.org/10.1016/j.schres.2015.05.011
430   Featherstone, R. E., Melnychenko, O., & Siegel, S. J. (2017). Mismatch negativity in
431   preclinical models of schizophrenia. *Schizophr Res*.
432   https://doi.org/10.1016/j.schres.2017.07.039
433   Ford, T. C., & Crewther, D. P. (2014). Factor Analysis Demonstrates a Common Schizoidal
434   Phenotype within Autistic and Schizotypal Tendency: Implications for Neuroscientific

Studies. *Frontiers in Psychiatry*, *5*(117), 1–11.
https://doi.org/10.3389/fpsyt.2014.00117
Ford, T. C., Woods, W., & Crewther, D. P. (2017). Mismatch field latency, but not power,
may mark a shared autistic and schizotypal trait phenotype. *International Journal of*
Psychophysiology, *116*, 60–67. https://doi.org/10.1016/j.ijpsycho.2017.02.008
Friston, K. (2005). A theory of cortical responses. *Philos Trans R Soc Lond B Biol Sci*,
360(1456), 815–836. https://doi.org/10.1098/rstb.2005.1622
Fulham, W. R., Michie, P. T., Ward, P. B., Rasser, P. E., Todd, J., Johnston, P. J., Thompson, P.
M., & Schall, U. (2014). Mismatch negativity in recent-onset and chronic
schizophrenia: A current source density analysis. *PLoS One*, *9*(6), e100221.
https://doi.org/10.1371/journal.pone.0100221
Gayle, L. C., Gal, D. E., & Kieffaber, P. D. (2012). Measuring affective reactivity in individuals
with autism spectrum personality traits using the visual mismatch negativity event-
related brain potential. *Frontiers in Human Neuroscience*, *6*.
https://doi.org/10.3389/fnhum.2012.00334
Jahshan, C., Cadenhead, K. S., Rissling, A. J., Kirihara, K., Braff, D. L., & Light, G. A. (2012).
Automatic sensory information processing abnormalities across the illness course of
schizophrenia. *Psychol Med*, *42*(1), 85–97.
https://doi.org/10.1017/S0033291711001061
Jamovi (1.2). (2020). [Computer software]. https://www.jamovi.org
Kecskes-Kovacs, K., Sulykos, I., & Czigler, I. (2013). Is it a face of a woman or a man? Visual
mismatch negativity is sensitive to gender category. *Frontiers in Human*
Neuroscience, *7*, 532. https://doi.org/10.3389/fnhum.2013.00532
Kenemans, J. L., Hebly, W., Heuvel, E. van den, & Grent-'T-Jong, T. (2010). Moderate alcohol
disrupts a mechanism for detection of rare events in human visual cortex. *Journal of*
Psychopharmacology, *24*(6), 839–845. https://doi.org/10.1177/0269881108098868
Kremláček, J., Kreegipuu, K., Tales, A., Astikainen, P., Põldver, N., Näätänen, R., & Stefanics,
G. (2016). Visual mismatch negativity (vMMN): A review and meta-analysis of studies
in psychiatric and neurological disorders. *Cortex*, *80*, 76–112.
https://doi.org/10.1016/j.cortex.2016.03.017
Lanillos, P., Oliva, D., Philippsen, A., Yamashita, Y., Nagai, Y., & Cheng, G. (2020). A review on
neural network models of schizophrenia and autism spectrum disorder. *Neural*
Networks, *122*, 338–363. https://doi.org/10.1016/j.neunet.2019.10.014
Lavoie, S., Jack, B. N., Griffiths, O., Ando, A., Amminger, P., Couroupis, A., Jago, A., Markulev,
C., McGorry, P. D., Nelson, B., Polari, A., Yuen, H. P., & Whitford, T. J. (2018).
Impaired mismatch negativity to frequency deviants in individuals at ultra-high risk
for psychosis, and preliminary evidence for further impairment with transition to
psychosis. *Schizophrenia Research*, *191*, 95–100.
https://doi.org/10.1016/j.schres.2017.11.005
Lee, S. H., Sung, K., Lee, K. S., Moon, E., & Kim, C. G. (2014). Mismatch negativity is a
stronger indicator of functional outcomes than neurocognition or theory of mind in
patients with schizophrenia. *Prog Neuropsychopharmacol Biol Psychiatry*, *48*, 213–
219. https://doi.org/10.1016/j.pnpbp.2013.10.010
Lepistö, T., Nieminen-von Wendt, T., von Wendt, L., Näätänen, R., & Kujala, T. (2007).
Auditory cortical change detection in adults with Asperger syndrome. *Neuroscience*
Letters, *414*(2), 136–140. https://doi.org/10.1016/j.neulet.2006.12.009

Lovibond, P. F., & Lovibond, S. H. (1995). The structure of negative emotional states: Comparison of the Depression Anxiety Stress Scales (DASS) with the Beck Depression and Anxiety Inventories. *Behav Res Ther*, *33*(3), 335–343. https://doi.org/10.1016/0005-7967(94)00075-U

Luck, S. J., & Gaspelin, N. (2017). How to get statistically significant effects in any ERP experiment (and why you shouldn't): How to get significant effects. *Psychophysiology*, *54*(1), 146–157. https://doi.org/10.1111/psyp.12639

Maekawa, T., Tobimatsu, S., Inada, N., Oribe, N., Onitsuka, T., Kanba, S., & Kamio, Y. (2011). Top-down and bottom-up visual information processing of non-social stimuli in high-functioning autism spectrum disorder. *Research in Autism Spectrum Disorders*, *5*(1), 201–209. https://doi.org/10.1016/j.rasd.2010.03.012

Naatanen, R., Kujala, T., Escera, C., Baldeweg, T., Kreegipuu, K., Carlson, S., & Ponton, C. (2012). The mismatch negativity (MMN)—A unique window to disturbed central auditory processing in ageing and different clinical conditions. *Clin Neurophysiol*, *123*(3), 424–458. https://doi.org/10.1016/j.clinph.2011.09.020

Naatanen, R., Kujala, T., Kreegipuu, K., Carlson, S., Escera, C., Baldeweg, T., & Ponton, C. (2011). The mismatch negativity: An index of cognitive decline in neuropsychiatric and neurological diseases and in ageing. *Brain*, *134*(Pt 12), 3435–3453. https://doi.org/10.1093/brain/awr064

Naatanen, R., Paavilainen, P., Rinne, T., & Alho, K. (2007). The mismatch negativity (MMN) in basic research of central auditory processing: A review. *Clin Neurophysiol*, *118*(12), 2544–2590. https://doi.org/10.1016/j.clinph.2007.04.026

Neuhaus, A. H., Brandt, E. S. L., Goldberg, T. E., Bates, J. A., & Malhotra, A. K. (2013). Evidence for impaired visual prediction error in schizophrenia. *Schizophrenia Research*, *147*(2–3), 326–330. https://doi.org/10.1016/j.schres.2013.04.004

Raine, A. (1991). The SPQ- A Scale for the Assessment of Schizotypal Personality Based on DSM-III-R Criteria. *Schizophrenia Bulletin*, *17*(4), 555–564.

Randeniya, R., Oestreich, L. K. L., & Garrido, M. I. (2018). Sensory prediction errors in the continuum of psychosis. *Schizophr Res*, *191*, 109–122. https://doi.org/10.1016/j.schres.2017.04.019

Rosburg, T., Marinou, V., Haueisen, J., Smesny, S., & Sauer, H. (2004). Effects of lorazepam on the neuromagnetic mismatch negativity (MMNm) and auditory evoked field component N100m. *Neuropsychopharmacology*, *29*(9), 1723–1733. https://doi.org/10.1038/sj.npp.1300477

Schwartz, S., Shinn-Cunningham, B., & Tager-Flusberg, H. (2018). Meta-analysis and systematic review of the literature characterizing auditory mismatch negativity in individuals with autism. *Neuroscience & Biobehavioral Reviews*, *87*, 106–117. https://doi.org/10.1016/j.neubiorev.2018.01.008

Stefanics, G., Kremlacek, J., & Czigler, I. (2014). Visual mismatch negativity: A predictive coding view. *Front Hum Neurosci*, *8*, 666. https://doi.org/10.3389/fnhum.2014.00666

Thye, M. D., Bednarz, H. M., Herringshaw, A. J., Sartin, E. B., & Kana, R. K. (2018). The impact of atypical sensory processing on social impairments in autism spectrum disorder. *Developmental Cognitive Neuroscience*, *29*, 151–167. https://doi.org/10.1016/j.dcn.2017.04.010

Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., Marcus, D. J., Westerlund, A., Casey, B., & Nelson, C. (2009). The NimStim set of facial expressions:

Judgments from untrained research participants. *Psychiatry Research*, *168*(3), 242–
249. https://doi.org/10.1016/j.psychres.2008.05.006
Umbricht, D., & Krljes, S. (2005). Mismatch negativity in schizophrenia: A meta-analysis.
Schizophr Res, *76*(1), 1–23. https://doi.org/10.1016/j.schres.2004.12.002
Urban, A., Kremlacek, J., Masopust, J., & Libiger, J. (2008). Visual mismatch negativity among
patients with schizophrenia. *Schizophrenia Research*, *102*(1–3), 320–328.
https://doi.org/10.1016/j.schres.2008.03.014
Vogel, B. O., Stasch, J., Walter, H., & Neuhaus, A. H. (2018). Emotional context restores
cortical prediction error responses in schizophrenia. *Schizophrenia Research*, *197*,
434–440. https://doi.org/10.1016/j.schres.2018.02.030
Wilbarger, J. L., McIntosh, D. N., & Winkielman, P. (2009). Startle modulation in autism:
Positive affective stimuli enhance startle response. *Neuropsychologia*, *47*(5), 1323–
1331. https://doi.org/10.1016/j.neuropsychologia.2009.01.025
Willenbockel, V., Sadr, J., Fiset, D., Horne, G. O., Gosselin, F., & Tanaka, J. W. (2010).
Controlling low-level image properties: The SHINE toolbox. *Behavior Research*
Methods, *42*(3), 671–684. https://doi.org/10.3758/BRM.42.3.671
Wright, B., Clarke, N., Jordan, J., Young, A. W., Clarke, P., Miles, J., Nation, K., Clarke, L., &
Williams, C. (2008). Emotion recognition in faces and the use of visual context Vo in
young people with high-functioning autism spectrum disorders. *Autism*, *12*(6), 607–
626. https://doi.org/10.1177/1362361308097118
Wuthrich, V., & Bates, T. C. (2005). Reliability and validity of two Likert versions of the
Schizotypal Personality Questionnaire (SPQ). *Personality and Individual Differences*,
38(7), 1543–1548. https://doi.org/10.1016/j.paid.2004.09.017

551
552   **Legends**
553 Figure 1: Example stimulus presentation in the neutral standard - happy deviant - sad
554 deviant condition. Stimuli were presented for 400 ms with a 150 ms ± 5 ms jitter. At least 2
555 standard faces were presented between each deviant. Participants attended to the cross in
556 the centre of the face and pressed the relevant key when the cross changed colour.
557
558 Figure 2: The line graph shows the grand-averaged deviant-minus-standard difference
559 waves at occipital electrodes (PO7, O1, Oz, O2, PO8), showing time (ms) on the x-axis, with 0
560 indicating stimulus onset of the emotional face, and voltage (µV) on the y-axis, with positive
561 voltages plotted upwards. The voltage maps show the distribution of voltages over the back
562 of the scalp during the vMMN time-window (187 to 287 ms).
563
564 Figure 3: Mean and standard error bars for visual mismatch negativity (vMMN) amplitude
565 (µV) by visual electrode and facial emotion.
566
567 Figure 4: Scatter plots of uncorrected correlations between visual mismatch negativity
568 (vMMN) amplitude (µV) to happy faces and Communication, Attention to Detail and
569 Interpersonal subscales at O1 and O2 electrodes.
570

                                          Tables and Figures

573   Table 1. Final sample characteristics for full-scale AQ and SPQ total and dimension scores.

	Total Mean(*SE*)	Female Mean(*SE*)	Male Mean(*SE*)	Mann-Whitney *U*	*p*-value
N	60	32	28		
Age	25.02(0.6)	24.91(0.84)	25.14(0.86)	435.0	0.853
AQ total	105.28(2.93)	99.03(3.84)	112.43(4.16)	306.5	0.037*
Social Skills	20.2(0.81)	19.16(1.04)	21.39(1.25)	350.0	0.148
Communication	19.13(0.77)	17.69(0.99)	20.79(1.15)	321.5	0.061
Attention Switching	23.28(0.67)	21.34(0.91)	25.5(0.81)	240.0	0.002*
Attention to Detail	23.2(0.72)	22.41(1.06)	24.11(0.96)	358.5	0.186
Imagination	19.47(0.68)	18.44(0.75)	20.64(1.14)	375.0	0.281
SPQ total	145.85(4.94)	136.75(6.29)	156.25(7.4)	323.5	0.066
Cognitive-Perceptual	56.82(2.4)	52.88(2.99)	61.32(3.72)	340.0	0.111
Interpersonal	54.63(1.92)	52.34(2.39)	57.25(3.05)	380.5	0.320
Disorganisation	34.4(1.4)	31.53(1.68)	37.68(2.18)	315.0	0.049*
Handedness (R:L)	53:7	26:6	27:1	$\chi^2 = 3.34$	0.068

574   Notes: *SE* = standard error, AQ = autism spectrum quotient, SPQ = schizotypal personality

575   questionnaire, χ^2 = chi square test.

578 Table 2. Spearman rank order correlations between AQ and SPQ dimensions and average vMMN amplitudes across visual electrode sites for
579 each emotional expression.

	Happy vMMN					Sad vMMN					Neutral vMMN				
	PO7	O1	Oz	O2	PO8	PO7	O1	Oz	O2	PO8	PO7	O1	Oz	O2	PO8
Social skills	-0.11	-0.19	-0.18	-0.22	-0.04	0.17	0.08	0.02	0.04	0.05	-0.12	-0.12	-0.12	-0.12	0.03
Communication	-0.21	-0.29*	-0.23	-0.28*	-0.07	0.21	0.14	0.12	0.07	0.11	-0.09	-0.07	-0.11	-0.08	0.05
Attention Switching	-0.10	-0.07	-0.21	-0.19	-0.11	0.15	0.14	0.10	0.21	0.20	-0.04	-0.07	-0.14	-0.08	-0.08
Attention to Detail	-0.19	-0.30*	-0.13	-0.27*	-0.09	0.0	-0.02	0.02	0.01	-0.04	0.02	0.06	0.08	0.03	0.15
Imagination	0.02	-0.08	-0.02	-0.03	0.17	0.03	0.03	0.03	0.01	0.08	-0.21	-0.21	-0.15	-0.21	-0.17
AQ total	-0.15	-0.24	-0.19	-0.25	-0.05	0.13	0.10	0.08	0.06	0.07	-0.13	-0.13	-0.12	-0.13	0.0
Cognitive-Perceptual	-0.14	-0.20	-0.20	-0.14	0.10	0.12	0.06	0.04	0.04	0.11	-0.12	-0.04	-0.06	-0.13	0.04
Interpersonal	-0.16	-0.26*	-0.25	-0.28*	0.0	0.16	0.10	0.04	0.01	0.02	-0.02	-0.07	-0.04	-0.12	0.10
Disorganised	-0.10	-0.10	-0.18	-0.24	0.05	0.20	0.10	0.05	-0.02	0.06	-0.01	0.01	-0.05	-0.09	0.0
SPQ total	-0.18	-0.23	-0.27*	-0.27*	0.03	0.16	0.07	0.03	0.01	0.07	-0.05	-0.01	-0.04	-0.11	0.06

580 Notes: AQ = Autism Spectrum Quotient, SPQ = Schizotypal Personality Questionnaire. *p < .05 uncorrected. No correlations survived correction
581 for false discovery rate.

584 Table 3. Spearman rank order correlations between AQ and SPQ dimensions and average
585 vMMN amplitudes when the mismatch response was calculated for happy and sad deviants
586 relative to neutral standards

	Happy vMMN					Sad vMMN				
	PO7	O1	Oz	O2	PO8	PO7	O1	Oz	O2	PO8
Social skills	-0.11	-0.16	-0.10	-0.25	-0.13	0.06	0.05	-0.04	-0.04	-0.02
Communication	-0.11	-0.16	-0.16	-0.27*	-0.09	0.10	0.03	0.05	-0.02	-0.01
Attention Switching	0.03	0.05	0.06	0.01	0.04	0.02	0.10	0.06	0.11	0.10
Attention to Detail	-0.09	-0.13	-0.14	-0.15	-0.02	-0.04	0.01	0.09	-0.02	-0.11
Imagination	0.05	0.02	0.16	0.02	0.22	-0.07	0.01	-0.03	-0.04	-0.03
AQ total	-0.07	-0.10	-0.05	-0.18	-0.01	0.01	0.05	0.03	-0.02	-0.04
Cognitive-Perceptual	-0.09	-0.22	-0.21	-0.23	0.04	-0.02	-0.05	-0.07	-0.12	-0.10
Interpersonal	-0.04	-0.18	-0.13	-0.24	-0.08	0.01	-0.03	0.02	-0.12	-0.14
Disorganised	0.10	-0.01	-0.09	-0.24	0.07	0.03	-0.04	-0.03	-0.12	-0.16
SPQ total	-0.07	-0.21	-0.21	-0.29*	-0.02	-0.12	-0.05	-0.02	-0.14	-0.17

587 Notes: AQ = Autism Spectrum Quotient, SPQ = Schizotypal Personality Questionnaire. *$p <$
588 .05 uncorrected. No correlations survived correction for false discovery rate.

589
590 **Figure 1**

Stimulus = 400ms

ISI = 150ms ± 5ms

OR

≤ 2 standards

591
592

593   **Figure 2**

Happy vMMN

Neutral vMMN

Sad vMMN

594
595

596   **Figure 3**

597
598

599   **Figure 4**