Emerin oligomerization and nucleoskeletal coupling at the nuclear envelope regulate nuclear mechanics against stress

Anthony Fernandez[1], Markville Bautista[2] and Fabien Pinaud[1,2,3*]

[1]Department of Biological Sciences, University of Southern California, 1050 Childs Way, Los Angeles, CA 90089, USA.

[2]Department of Chemistry, University of Southern California, 1050 Childs Way, Los Angeles, CA 90089, USA.

[3]Department of Physics and Astronomy, University of Southern California, 1050 Childs Way, Los Angeles, CA 90089, USA.

*Correspondence should be addressed to F.P. (pinaud@usc.edu)

ABSTRACT

Emerin is an integral membrane protein of the nuclear envelope that binds to various nucleoskeletal partners and participates in the maintenance of nuclear shape. When mutated or absent, emerin causes X-linked Emery-Dreifuss muscular dystrophy (EDMD). To define how emerin takes parts in molecular scaffolding at the nuclear envelope and helps protect the nucleus against mechanical stress, we established the nanoscale spatial organizations of wild-type emerin and EDMD-inducing emerin mutants in human cells using single molecule tracking and super-resolution microscopy. We show that emerin is distributed as monomers and oligomeric clusters at the inner nuclear membrane and that its interactions with lamin A/C, nuclear actin and BAF differentially modulate its mobility and its ability to oligomerize. In nuclei under stress, the mechanotransducing functions of emerin are coupled to its oligomeric state and the oligomerization of wild-type emerin or that of emerin mutants determine normal or aberrant nuclear deformations against increasing forces. These findings place emerin at the center of the molecular processes that regulate nuclear shape remodeling and maintain the structural integrity of the nuclear envelope in response to mechanical stress.

INTRODUCTION

Emerin is a conserved inner nuclear membrane (INM) protein that participates in the maintenance of the nuclear architecture by interacting with the nucleoskeletal lamina and elements of the linker

of the nucleoskeleton and cytoskeleton (LINC) complex[1,2]. Emerin also contributes to tethering

chromatin at the nuclear envelope by binding the DNA-bridging barrier-to-autointegration factor

(BAF)[3] and it regulates the activity of chromatin compaction modulators[4]. Both the nucleoskeleton

and the compaction state of chromatin contribute to the mechanical responses of nuclei to forces[5]

and emerin is deemed a pivotal actor of mechanotransducing processes at the nuclear envelope[6].

When mutated or absent, emerin causes X-linked Emery-Dreifuss muscular dystrophy (EDMD)[7,8],

a disease part of a larger of group of laminopathies[9] associated with structural perturbations of the

nuclear envelope and its underlying lamina. While emerin is ubiquitously expressed, mutations in

the *EMD* gene primarily affect cells exposed to extensive mechanical stress, such as skeletal and

cardiac muscle cells. Emerin-null muscle tissues display deformed and disorganized nuclei,

impaired myogenesis and improper muscle fiber formation, which participate in the muscle

wasting and cardiac disease phenotypes of EDMD[10-12]. In non-muscle cells, loss of emerin also

leads to altered nuclear envelope elasticity and increased nuclear fragility[13], impaired expression

of mechanosensitive genes[14] and enhanced apoptosis after continuous mechanical strain[14]. EDMD

phenotypes might stem from both an altered structural integrity of the nuclear envelope and

modified gene expression profiles.

The structure of emerin and its binding to various partners have been extensively characterized *in*

vitro[1]. Emerin is 254 amino acids (aa) long, with a N-terminal globular LAP2-emerin-MAN1

(LEM, aa: 1-45) domain, followed by a long intrinsically disordered region (IDR, aa: 46-222)[15]

and a transmembrane domain close to the C-terminus (aa: 223-234) that allows anchoring to the

nuclear envelope. Prominent binding partners of emerin include lamin A/C and lamin B[16-18],

actin[19], BAF[20], nesprins and SUN proteins from the LINC complex[21,22], chromatin modification

enzymes and other factors[1]. While binding of emerin to BAF takes place via its structured LEM

domain, interactions with other binding partners and emerin self-association[18,23] are mediated by

the IDR. Emerin can indeed oligomerize via regions of the IDR that serve as LEM binding and

self-assembly sites between emerin monomers[18], and, *in vitro*, emerin self-association into

filamentous structures[24] impact binding to lamin A/C and BAF[18]. Post-translational modifications

in the IDR and the LEM domain also influence the oligomerization of emerin and its interactions

with binding partners[1,25,26]. Phosphorylation of emerin residues Y74 and Y95 are required for

lamin A/C recruitment to the LINC complex and nuclear envelope stiffening in response to

mechanical stress[27]. Additional post-translational modifications regulate the affinity of the LEM

domain to BAF[26] and binding to actin[28]. How emerin organizes at the nuclear envelope and

participates in protecting the nucleus against mechanical strains remains unclear.

EDMD often involves frame-shift deletions and nonsense mutations of emerin that effectively

ablates expression in cells[29]. Yet, a few sets of point mutations and small deletions, including Δ95-

99, Q133H and P183H/T, also cause EDMD despite an apparently correct localization of these

mutated emerins at the nuclear envelope and normal cellular expression levels[19,20,30,31]. These

emerin mutants, however, display altered self-association and binding to nucleoskeletal proteins

in vitro[1,23] and possible changes in molecular organizations at the nuclear envelope might explain

why they act as null emerin in cells.

Here, we used super-resolution imaging by stochastic optical reconstruction microscopy

(STORM) and single particle tracking by photoactivated localization microscopy (sptPALM) to

investigate the nanoscale organization of wild-type emerin and emerin mutants at the nuclear

envelope in cells. By modulating the mechanical landscape of nuclei using cell micropatterning

techniques[32], we also studied how emerin participates in nuclear envelope mechanotransduction

processes during stress. We show that emerin oligomerization and its differential interactions with

key structural partners at the nuclear envelope is central to nuclear shape adaptation against

mechanical challenges.

RESULTS

Emerin displays distinct diffusive behaviors at the nuclear envelope

To study the nuclear envelope dynamics of wild-type and dystrophy causing mutations of emerin,

we made PA-TagRFP-emerin fusions that we expressed in emerin-null human dermal fibroblasts

(HDF) from an EDMD patient (*EMD*[-/y])[33]. PA-TagRFP-emerin expression led to the expected

localization of emerin at the nuclear envelope, as observed for endogenous emerin in HDF from a

healthy individual (*EMD*[+/y]) (Figure 1a). For sptPALM, PA-TagRFP-emerin was excited at 405

nm and 561 nm using highly inclined and laminated optical sheet (HILO)[34] illumination to

photoactivate single emerin molecules and image their diffusion at the bottom nuclear envelope.

Diffusion tracks were generated from localized emerin appearances (Figure 1b and Supplementary

Movie 1) and diffusion analyses were performed using the probability distribution of square

displacement (PDSD)[35] to separate unique diffusive behaviors (Figure 1c).

From the PDSD analysis of thousands PA-TagRFP-emerin trajectories, we observed that wild-

type emerin exhibits four distinct diffusive behaviors with diffusion values D_1: $2.21 \times 10^{-1} \pm 4.9 \times 10^{-2}$

$\mu m^2 s^{-1}$, a value similar to that previously reported for diffusion at the ER membrane[36], D_2:

$1.48 \times 10^{-2} \pm 1.5 \times 10^{-3}$ $\mu m^2 s^{-1}$, D_3: $1.73 \times 10^{-3} \pm 1.1 \times 10^{-4}$ $\mu m^2 s^{-1}$, and D_4: $2.6 \times 10^{-4} \pm 1 \times 10^{-5}$ $\mu m^2 s^{-1}$

(Figure 1d and Supplementary Table 1). The putative ER population D_1 makes up 1% of the

detected behaviors, while the next slowest population D_2 represents 9%, a small fraction of emerin

similar to that expected at the outer nuclear membrane (ONM)[2,6]. The two slowest populations D_3

and D_4 collectively represent 90% of emerin diffusive behaviors. To determine if these four

diffusive behaviors are associated with specific positions of emerin at the nuclear envelope, we

examined the diffusion coefficient of single emerins from individual mean square displacements.

Based on their diffusion coefficients, we grouped emerin trajectories into four diffusion ranges

spanning the D values measured from PDSD analyses and plotted them as maps (Figure 1e). In

trajectory maps, the fastest emerin population D_1 is primarily distributed in the immediate

proximity of nuclear envelope (Figure 1e), a location consistent with diffusion in the ER

membrane. The three slower populations D_2, D_3 and D_4 display enrichment at the nuclear

membrane (Figure 1e). Compared to diffusion analyses by fluorescence recovery after

photobleaching, where we could detect only two diffusive behaviors of wild-type emerin

(Supplementary Figure 1), these measurements show that the mobility of emerin involves four

very different types of diffusions at and around the nuclear envelope.

Emerin organizes as slowly diffusing monomers or oligomers at the inner nuclear membrane

Emerin primarily localizes at the INM but a small fraction is often detected at the ONM[2,6], where

it cannot interact with the nuclear lamina. We therefore determined if interactions with lamin A/C

would affect specific diffusing populations of wild-type emerin. We found that downregulation of

lamin A/C expression by siRNA (Supplementary Figure 2) increases the mobility of the two

slowest populations D_3 and D_4 ($p<0.01$, Figure 2a and Supplementary Table 1), but does not affect

the faster populations D_1 and D_2 (p not significant, Figure 2a). This shows that D_3 (40%) and D_4

(50%) are two distinct emerin populations diffusing at the INM and interacting with lamin A/C,

while population D_2 (9%) corresponds to emerin diffusing at the ONM. A similar increase in the

mobility of emerin populations D_3 and D_4 is also observed when nuclear actin is depleted after

downregulating the expression of the actin nuclear import factor, importin-9[37] (IPO9,

Supplementary Figure 2), although decreased nuclear actin levels additionally induce slightly

faster emerin diffusion at the ONM (Figure 2a and Supplementary Table 1). These results indicate

that a large majority of emerin is organized as two distinct INM pools, both interacting with the

nucleoskeleton.

Emerin can also self-assemble *in vitro*[23,24], and its oligomerization at the INM could result in the

slow mobility we observed, in particular if emerin oligomers are tethered by the nucleoskeleton.

To determine if diffusing populations D_3 or D_4 correspond to emerin monomers or oligomers, we

tracked individual emerins in their dimeric and oligomeric forms. We co-expressed wild-type

emerin fusions to self-complementary split-GFP fragments[38] (sGFP-emerin and M3-emerin,

Figure 2b) to irreversibly induce the formation of complemented emerin-GFP-emerin species and

we studied their lateral mobility by complementation activated light microscopy[39] (CALM). We

found that emerin-GFP-emerin species localize almost exclusively at the nuclear envelope (Figure

2c and Supplementary Movie 2), where they diffuse as two separate populations: a dominant 90%

population with a mobility similar to D_4 (*p* not significant, Figure 2d), and a minor 10% population

with a mobility similar to the ONM population D_2 (*p* not significant, Figure 2d). No fluorescent

species with diffusion coefficients matching those of the ER population D_1 or the INM population

D_3 are detected. Emerin-GFP-emerin trajectories often spatially overlap, indicating that emerin

forms multimers rather than strictly dimers at the INM (Figure 2c). These results show that

population D_3 detected by sptPALM but not by CALM represents emerin monomers, while the

slowest population D_4 represents dimers or, more likely, larger emerin oligomers at the INM, as

we subsequently confirmed by super-resolution imaging. The nearly 10-fold difference in

translational diffusion coefficient between populations D_3 and D_4 is close to the 10-40 fold

reduction in mobility reported when monomeric membrane proteins transition to oligomers[40]. The

detection of a few emerin-GFP-emerin species at the ONM is likely due to split-GFP-induced

emerin dimers that cannot translocate through peripheral nuclear pores channels[9] and do not access

the INM.

BAF binding modulates the mobility of both emerin monomers and oligomers distributed

across the inner nuclear membrane.

BAF binds the LEM domain of emerin with high affinity[18] and, because it also binds to lamin

A/C[41] and chromatin[42], it likely participates in the slow mobility of emerin at the INM[43]. To study

how binding of emerin monomers and oligomers to BAF might influence their respective

diffusion, endogenous BAF was knockdown by siRNA (Supplementary Figure 2) and replaced by

the expression of BAF L58R, a mutant that does not bind LEM domains[44]. When unable to bind

BAF L58R, emerin monomers diffuse significantly faster with a lateral mobility higher than for

lamin A/C or IPO9 knock down ($p<0.01$, Figure 2a and Supplementary Table 1). Emerin oligomers

also diffuse faster ($p<0.01$, Figure 2a), indicating that BAF modulates the INM dynamics of emerin

when it oligomerizes. While emerin diffusion at the ER is undisturbed, its mobility at the ONM

increases ($p<0.01$, Figure 2a), which implies that cytoplasmic BAF additionally interacts with

emerin at the ONM. BAF is highly mobile in the nucleus[43] and the comparatively slow mobility

of BAF-bound emerin at the INM likely stems from the formation of ternary complexes with lamin

A/C or chromatin. The stronger influence of BAF L58R on emerin dynamics compared to lamin

A/C depletion indeed indicates that BAF binding modulates the mobility of emerin not only via

lamin A/C, but also via interactions with other nuclear components, potentially chromatin.

We additionally built diffusion maps of wild-type PA-tagRFP-emerin at the nuclear envelope by

spatial rendering of diffusion coefficients at the position of each tracked molecule (Figure 2e).

Those maps are dominated by the two slowest INM emerin populations, which represent 90% of

the total pool detected. They show that domains with very slow diffusion and attributed to emerin

oligomers are distributed throughout the nuclear envelope and are surrounded by diffusion areas

where the mobility of emerin is faster and matches that of monomers (Figure 2e). This indicates a

membrane-wide, yet locally structured distribution of emerin at the INM, with monomers

surrounding emerin oligomer domains in an organization compatible with emerin

monomer/oligomer exchanges in distinct INM areas.

Together, these diffusion analyses show that newly synthesized emerin diffuse rapidly at the ER

membrane on their way to the juxtaposed ONM, where emerin diffusion is slower, in part by

interaction with cytoplasmic BAF. Once at the INM, the mobility of emerin is further reduced,

with emerin monomers and oligomers interacting with nuclear BAF and the nucleoskeleton. This

distribution is consistent with a combined BAF-induced and lamina-induced retention and

accumulation of emerin at the INM.

Emerin forms discrete oligomeric nanodomains surrounded by monomers

We then probed the structural organization of emerin in more detail using super-resolution microscopy. Wild-type emerin fused to a SNAP-tag (SNAP-emerin) was expressed in *EMD*[-/y] HDF and, after chemical fixation, cells were stained with benzylguanine-FluorAlexa647 to image individual emerins by dSTORM (Figure 3a). An emerin neighborhood density function (NDF) built from localized emerin positions across multiple nuclei was compared to MonteCarlo simulations of completely random distributions and fitted with exponential decay models[45] to measure molecular densities and length scales of significant clustering. We found that wild-type emerin is not distributed randomly at the nuclear envelope, as it displays density distributions significantly higher than expected for a completely random organization (Figure 3b). The NDF curve is best fitted with a two exponential decay model, indicating that emerin organizes into two distinct clustering states across the nuclear envelope. The first type of cluster is small with a typical cluster length of 22 ± 11 nm and a molecular density of emerin 8.2 ± 0.2 fold higher than expected for a random distribution (Figure 3c and Supplementary Table 2). The second type of cluster is significantly larger with a typical cluster length of 236 ± 30 nm and an emerin density slightly above the expected value of 1 for complete randomness (1.3 ± 0.1 fold, Figure 3c). In cluster maps built from local cluster analyses[46], wild-type emerin forms small high density clusters that are scattered throughout the nuclear envelope and are surrounded by larger emerin areas with densities close to random (Figure 3d). These cluster maps are strikingly similar to our sptPALM diffusion maps in live cells (Figure 2e) and indicate that small, high emerin density clusters are oligomers while the surrounding larger areas are populated by emerin monomers. While it is intriguing that emerin monomers appear as slightly clustered, a fully random distribution is not to be expected because they do not populate membrane-less areas of the nuclear envelope, such as nuclear pores. These observations confirm that, across the INM, emerin organizes into small and discrete oligomeric units surrounded by dispersed monomers.

Emerin oligomers are primarily stabilized by lamin A/C and partially maintained by nuclear actin and BAF

We then determined the importance of lamin A/C, nuclear actin and BAF in maintaining the nanoscale spatial organization of emerin at the INM. Wild-type emerin was imaged by super-resolution microscopy after siRNA knock down of lamin A/C, IPO9, or replacement of

endogenous BAF with BAF L58R. After lamin A/C depletion, the molecular density of emerin is
dramatically reduced at all length scales (Figure 3b). The NDF distribution is comprised of small,
35±17 nm emerin clusters with molecular density close to random (1.2±0.1 fold, Figure 3c and
Supplementary Table 2) and of large 950±173 nm emerin monomer domains with density just
above random (1.1±0.1 fold, Figure 3c). Compared to emerin distributions in cells having normal
lamin A/C levels, emerin monomers are dispersed over larger areas and, importantly, emerin
oligomers nearly vanish. Such changes in organization are seen in emerin cluster maps of lamin
A/C depleted nuclei, where most oligomeric nanodomains have dissipated, and where monomers
cover a significantly larger part of the nuclear envelope (Figure 3d). This shows that lamin A/C
plays an important structural role for the stabilization of emerin oligomers and the general spatial
distribution of emerin monomers at the INM.
When nuclear actin is depleted after knocking down of IPO9, wild-type emerin forms oligomers
having a reduced molecular density of 3.6±0.1 fold above random and a larger size of 65±13 nm
compared to untreated cells (Figure 3c and Supplementary Table 2). Emerin monomers are
dispersed over membrane domains having a typical length scale of 382±81 nm, which is larger
than in untreated cells (236±30 nm, Figure 3c), but not as large as for lamin A/C knock down
(950±173 nm, Figure 3c). Thus, despite reduced nuclear actin levels, emerin retains the ability to
oligomerize, although it disperses over large INM areas. These results show that, together with
lamin A/C, nuclear actin contributes to maintaining the overall spatial distribution of emerin at the
INM and additionally participates in the structural maintenance of emerin oligomers, albeit to a
lesser degree than the lamina itself.
When endogenous BAF is replaced by BAF L58R, wild-type emerin oligomers are still formed
but they have a reduced molecular density of 2.0±0.1 fold above random and a larger size of 85±19
nm compared to cells expressing wild-type BAF (Figure 3c and Supplementary Table 2). The
inability to bind BAF also induces a re-distribution of emerin monomers over large INM domains
with typical sizes of 718±250 nm (Figure 3c). Consistent with our diffusion results, emerin binding
to BAF modulates the organization of both oligomeric and monomeric emerin and strongly
influences the spatial distribution of emerin monomers at the INM.
Together, these results show that wild-type emerin monomers are distributed throughout the INM
where their dispersion is maintained by direct binding to BAF and additional spatial constrains
imposed by lamin A/C and nuclear actin. Emerin additionally organizes into discrete and small

oligomeric nanodomains that are surrounded by monomers and are structurally stabilized by lamin

A/C and, to a lesser or more indirect extent, by BAF and nuclear actin.

252 Increased diffusion of emerin monomers and oligomerization upon nuclear adaptation to

253 mechanical stress

We then sought to define the mechanotransducing functions of emerin by studying how its

diffusion and its nanoscale organization change when nuclei are subjected to increasing

mechanical stress. SptPALM and super-resolution imaging of wild-type emerin were done in

EMD[-/y] HDF grown on increasingly narrow rectangular micropatterns with widths of 15 µm, 10

µm or 5 µm, to impose steady-state mechanical stress to the nucleus[6,32,47](Figure 4a and 4b). First,

we verified that changes in nuclear shape index (NSI) on these micropatterns reflect the mechanical

adaptation of nuclei against forces by depleting lamin A/C or nuclear actin, two key nucleoskeletal

proteins involved in maintaining nuclear shape during stress. In non-patterned cells, nuclei are

slightly more deformed after lamin A/C knock down or nuclear actin depletion compared to

controls ($p<0.01$, Figure 4c). These effects are further amplified on micropatterns where nuclei

become increasingly deformed as patterns get narrower ($p<0.01$, Figure 4c). This demonstrates

that cell micropatterning elicits increased nuclear stress and effectively induces mechanical

responses of the nucleus that implicate the nucleoskeleton.

When we tracked wild-type PA-tag-RFP-emerin on deformed nuclei in 15 µm and 10 µm patterns

(Supplementary Movie 3), we found that its ER mobility is unchanged compared to non-patterned

cells (p not significant, Figure 4d and Supplementary Table 1), while its ONM diffusion is faster

($p<0.01$, Figure 4d). At the INM, emerin monomers also diffuse significantly faster ($p<0.01$,

Figure 4d) and the mobility of oligomers increases compared to non-patterned cells, albeit short

of statistical significance for 15 µm patterns (p not significant and $p<0.01$, Figure 4d).

Interestingly, there is no significant difference in the mobility of all four ER, ONM and INM

emerin populations in both types of micropatterns compared to nuclear actin depletion (p not

significant, Supplementary Figure 3). This suggests that nuclear shape adaptation to mechanical

cues entails modified interactions of emerin with nuclear actin, notably for monomers.

Super-resolution imaging and spatial distribution analyses of wild-type SNAP-emerin further

indicate that, as the NSI decreases, monomers disperse over increasingly large INM domains with

sizes of 382±62 nm and 460±136 nm, compared to 236±30 nm for non-deformed nuclei (Figure

4e). Concurrently, oligomer densities drop from 8.2±0.2 fold to 3.6±0.1 fold above random in 15

µm patterns, before increasing slightly to 4.6±0.1 fold above random in 10 µm patterns (Figure

4e). Remarkably, oligomeric nanodomains become larger during nuclear stress and their size

expands from 22±11 nm to 60±13 nm in both micropatterns (Figure 4e). When considering this

wider spatial distribution of oligomers, the relative oligomerization of emerin compared to non-

deformed nuclei increases by 3.4 fold and by 4.1 fold as the nucleus adapts to incremental

mechanical stress (Figure 4f). This shows that nuclear shape adaptation to forces is associated with

a gradual change in the oligomerization potential of emerin at the INM. This enhanced

oligomerization of emerin over larger nanodomains is accompanied by a faster lateral mobility of

emerin and is likely triggered by nucleoskeletal re-arrangements. Indeed, the stress-induced spatial

reorganizations of emerin in micropatterns, including oligomer densities and emerin domain sizes,

is strikingly similar to those observed when nuclear actin is depleted in non-patterned cells

(Supplementary Figure 3 and Supplementary Table 2). This indicates that nuclear shape

deformation involves a disengagement of nuclear actin from the nucleoskeleton that leads to a

faster diffusion of emerin monomers and to their increased oligomerization. An increased lateral

mobility of emerin at the INM could indeed facilitate molecular collisions between monomers and

the formation of oligomers stabilized by lamin A/C.

Together, these results show that the mechanotransducing functions of emerin are coupled to its

oligomeric state along the INM and are modulated by emerin interactions with nucleoskeletal

partners, including nuclear actin, in order to ensure appropriate nuclear deformation and response

to mechanical challenges.

Emerin mutants induce abnormal nuclear deformation against mechanical stress

To establish the biological significance of emerin oligomerization at the INM and its importance

for nuclear adaptation to mechanical stress, we then compared the organization of wild-type

emerin with that of mutated forms of emerin known to induce EDMD. We studied how emerin

mutation Q133H, deletion Δ95-99, and mutation P183H (Figure 5a) impact the lateral diffusion

and the nanoscale distribution of emerin at the nuclear envelope. First, we verified that mutated

emerins effectively induce defective mechanical responses of nuclei when expressed in $EMD^{-/y}$

HDF and we compared changes in NSI after random cell plating or plating on increasingly narrow

rectangular micropatterns (Figure 5b, c). In randomly plated cells expressing mutated emerins,

nuclei are slightly less circular compared to cells expressing wild-type emerin (Figure 5b). With

increasing mechanical stress in micropatterns, cells expressing emerin mutants display

significantly higher NSI values than wild-type (Figure 5b), indicative of a failure to correctly

modify the shape of the nucleus in response to forces. These deficient changes in nuclear shape

are accompanied by a mispositioning of the nucleus relative to the cell major axis, nucleus

crumpling, abnormal organization of the actin cytoskeleton and failure of cells to properly fit

within micropatterns, specifically in cell areas adjacent to the misshaped nucleus (Figure 5c). It

indicates that expression of mutated emerin in $EMD^{-/y}$ HDF impedes nuclear adaptation to

mechanical stress.

Emerin mutants display defective oligomerization at the inner nuclear membrane

Q133H mutation

Using sptPALM and super-resolution imaging, we then characterized the nuclear envelope

dynamics of each mutated emerin and their respective nanoscale organization, starting with Q133H

emerin. We found that the lateral mobility of Q133H PA-TagRFP-emerin is similar to that of wild-

type emerin at the ER and the ONM (p not significant, Figure 6a and Supplementary Table 1).

However, both Q133H monomers and oligomers diffuse significantly faster at the INM ($p<0.01$

and $p<0.05$, Figure 6a). The Q133H mutation disrupts emerin binding to actin[48] but does not

impede interactions with lamin A/C[31], BAF or other partners[49]. The increased lateral diffusion of

Q133H, only at the INM, therefore indicates that it does not bind nuclear actin. This faster INM

diffusion of Q133H also resembles the increased mobility of wild-type emerin when nuclear actin

is depleted (Figure 2a), which further underlines that nucleoskeletal actin modulates the diffusion

of emerin monomers and oligomers. This is consistent with previous observations that emerin

expression influences the mobility of nuclear actin[50] and indicative of a reciprocal effect of

emerin/nuclear actin interactions on their respective mobility. The diffusion of Q133H at the INM

is also similar to that of wild-type emerin under mechanical stress, in both 15 µm and 10 µm wide

micropatterns (p not significant, Supplementary Figure 4 and Supplementary Table 1). This

signifies that nuclear deformations against stress involve a dissociation of emerin from nuclear

actin that leads to faster emerin diffusion at the INM.

Although, it was reported that Q133H has a reduced capacity to self-assemble in $vitro$[23], our

analyses of its spatial distribution and cluster maps show that, like wild-type emerin, it organizes

into monomers and oligomers across the INM (Figures 6b and 6c). Q133H monomers are

distributed over length scales of 213±62 nm, similar to those of wild-type emerin (Figure 6b),

indicating that the inability to bind the nuclear actin induces a faster diffusion of Q133H

monomers, but does not affect their overall spatial distribution at the INM. Q133H also retains the

ability to self-assemble at the INM where it forms oligomeric clusters (Figure 6c). These oligomers

have sizes of 19±12 nm, similar to the 22±11 nm size of wild-type emerin oligomers, but they have

molecular densities 12.2±0.2 fold above random, a 50% increase in oligomerization compared to

wild-type emerin (Figure 6b and Supplementary Table 2). This indicates that the deficient binding

of Q133H to nucleoskeletal actin leads to a disproportionate self-association of emerin into

oligomeric nanodomains. It also implies that direct binding to nuclear actin normally reduces the

oligomerization potential of wild-type emerin.

Δ95-99 mutation

When we performed similar studies with Δ95-99 emerin, we found that its diffusion at the ER

membrane and the ONM is similar to that of wild-type emerin (*p* not significant, Figure 6d and

Supplementary Table 1), but that its lateral mobility at the INM is significantly reduced ($p<0.05$

and $p<0.01$, Figure 6d). This suggests that Δ95-99 interacts more strongly or more frequently than

wild-type emerin with some of its binding partners on the nucleoplasmic side of the nuclear

envelope. Previous biochemical studies have shown that the small Δ95-99 deletion eliminates

emerin interactions with most of its binding partners, including lamin A/C and actin, but not BAF[1].

Considering that binding BAF strongly influences the mobility of wild-type emerin (Figure 2a),

the reduced diffusion of Δ95-99 at the INM could stem from its increased interaction with BAF,

as recently proposed[24]. To destabilize these interactions, we attempted to track Δ95-99 in *EMD*[-/y]

HDF knocked down for endogenous BAF and expressing BAF L58R. Co-expression of both Δ95-

99 and BAF mutants, however, proved toxic to cells, indicating that interactions between Δ95-99

emerin and BAF are important to maintain cell viability.

Beside a slower emerin mobility at the INM, the Δ95-99 deletion also induces defective

oligomerization. Δ95-99 is distributed randomly over large, 420±51 nm nuclear envelope domains

and in smaller, 48±14 nm nanodomains where the molecular density of 1.3±0.1 fold above random

is dramatically reduced compared to wild-type emerin oligomers (Figure 6e and Supplementary

Table 2). In cluster maps, Δ95-99 also displays less dense and fewer oligomerization nanodomains

(Figure 6f). Thus, Δ95-99 does not efficiently oligomerize at the INM, consistent with its impaired

self-assembly *in vitro* and observations that the deletion lowers emerin/emerin proximity[23].

Interestingly, despite an intact lamina, the molecular densities and cluster maps of Δ95-99 are

similar to those of wild-type emerin when the expression of lamin A/C is reduced (Figure 3c, d).

The decreased oligomerization of Δ95-99 to levels seen after lamin A/C depletion implies that

binding to lamin A/C is required to stabilize emerin oligomers at the INM. Such a stabilizing role

of lamin A/C is consistent with prior proximity ligation assays where Δ95-99 was found less close

to lamin A/C than wild-type emerin[23] and *in vitro* studies showing that the deletion abolishes lamin

A/C binding to emerin[20].

Together, these results show that Δ95-99 fails at oligomerizing and primarily distributes at random

throughout the INM due to its reduced self-association, its inability to directly bind lamin A/C and

its slow mobility.

P183H mutation

SptPALM tracking of P183H emerin reveals that its lateral diffusion at the ER membrane and the

ONM is unchanged compared to wild-type emerin (*p* not significant, Figure 6g and Supplementary

Table 1). However, its mobility at the INM is reduced for the populations attributed to monomers

and oligomers ($p<0.05$ and $p<0.01$, Figure 6g). This slow diffusion is similar to that observed for

Δ95-99 and again implies that P183H interacts more frequently than wild-type emerin with some

of its nucleoplasmic binding partners. *In vitro*, P183H maintains its binding to many emerin

partners, including BAF and actin[1] and displays enhanced binding to lamin A/C compared to wild-

type emerin[1,20]. The observed reduced INM diffusion of P183H might therefore be linked to an

increased binding frequency to BAF, like for Δ95-99, or to its enhanced binding to lamin A/C.

This slow INM mobility could also be due to the formation of dimers as recently suggested based

on the strong propensity of P183H to self-assemble *in vitro*[23] and observations that residue P183

is positioned in the 168-186 emerin region required to limit emerin-emerin association[18]. To

determine if P183H tends to form dimers at the nuclear envelope, we performed single particle

tracking by CALM after co-expression of sGFP-P183H and M3-P183H emerin. Three different

populations of fluorescently activated P183H-GFP-P183H emerin species were detected at the

nuclear envelope: a 2% population with a mobility slightly slower than P183H at the ER membrane

($p<0.05$, Figure 6g), a larger 39% population with a mobility comparable to the ONM behavior of

P183H (*p* not significant, Figure 6g), and a dominant 59% population with a lateral diffusion

similar to the INM population initially attributed to P183H monomers by sptPALM (*p* not

significant, Figure 6g and Supplementary Table 1). Surprisingly, no fluorescent species with

diffusion coefficient matching that of the slowest P183H oligomers are detected (Figure 6g). The

high frequency detection of complemented split-GFP for P183H compared to wild-type emerin at

both the ER membrane and the ONM (Figure 2d) indicates that P183H is more prone to self-

assemble and to form dimers than wild-type emerin before reaching the INM. This apparent biased

monomer:dimer equilibrium towards dimers is maintained at the INM where it precludes an

efficient association of P183H into oligomer domains, in particular when dimers are further

stabilized by irreversible assembly of the split-GFP fragments. This shows that P183H has a

propensity to form dimers that could impact oligomerization at the INM.

Consistent with these observations, spatial analyses reveal that the dimerization of P183H leads to

a significantly reduced oligomerization at the INM. P183H monomers/dimers are distributed in

domains with typical sizes of 321±29 nm and molecular densities of 1.3±0.1 fold above random

(Figure 6h and Supplementary Table 2). Smaller nanodomains with sizes of 35±12 nm are also

observed, but their molecular density is reduced to 2.2±0.1 compared to wild-type emerin

oligomers (Figure 6h). Indeed, while P183H can still bind lamin A/C, it forms oligomeric domains

having lower molecular density than wild-type emerin in cluster maps (Figure 6j). This indicates

that the dimerization of P183H hinders further self-association into dense oligomers at the INM.

Abnormal reorganization of Δ95-99 emerin in response to mechanical stress

To further characterize the importance of emerin oligomerization for nuclear deformation and

responses to mechanical stress, we studied the nanoscale organization of Δ95-99 in micropatterned

cells. For deformed nuclei in 15 µm patterns, Δ95-99 monomers are dispersed over large, 810±215

nm INM domains, almost double the size of monomer domains in non-stressed nuclei (420±51

nm, Figure 7a). However, as abnormal nuclear deformations become more pronounced in 10 µm

patterns, the distribution of Δ95-99 monomers return to non-stress levels, with dispersions in

domains 499±250 nm in size (Figure 7a). Contrary to wild-type emerin, a progressive dispersion

of Δ95-99 monomers over increasingly large INM areas is not observed with growing stress. At

the same time, the formation of oligomers remains very limited, with Δ95-99 molecular densities

increasing slightly from 1.3±0.1 to 1.7±0.1 and 2.0±0.1 fold above random for 15 µm and 10 µm

micropatterns (Figure 7a and Supplementary Table 2). With the concurrent size enlargement of

nanodomains from 48±14 nm to 81±16 nm and 75±20 nm, the relative oligomerization of Δ95-99

compared to non-mechanically stressed nuclei increases by 3.9 fold initially, but does not rise

further as nuclear stress intensifies (Figure 7b). Δ95-99 oligomeric nanodomains remains sparser

and significantly less dense than for wild-type emerin in cluster maps of nuclei under stress (Figure

7c). Thus, compared to wild-type emerin, the failure of Δ95-99 to gradually self-associate at

sufficiently high molecular densities leads to defective nuclear responses to force. It underlines the

importance of modulating the oligomerization of emerin as a function of mechanical stress for

adaptive nuclear deformations.

DISCUSSION

Using a combination of single molecule imaging and quantitative measurements, we showed that

emerin is distributed as monomers and oligomers at the INM and that clinically relevant EDMD

mutations induce modified emerin mobility and nanoscale organizations. The oligomerization of

emerin at specific sites across the nuclear envelope is modulated by its ability to engage or

disengage interactions with different structural elements juxtaposed to the INM, including lamin

A/C, nuclear actin and BAF. Lamin A/C is essential for a stabilization of emerin oligomers and

balanced interactions of emerin with BAF and nuclear actin further modulate its diffusion and its

oligomerization potential. We also showed that mechanotransducing functions of emerin are

coupled to its the oligomeric state, with the formation, stabilization and maintenance of emerin

oligomers being central to nuclear shape adaptation to forces. Indeed, EDMD-inducing mutations

that affect the oligomerization of emerin and its binding to structural elements at the INM lead to

abnormal nuclear deformation and defective nuclear positioning in response to mechanical stress.

While the altered *in vitro* binding properties of Q133H, Δ95-99 and P183H emerin mutants result

in expected differences in organization *in vivo*, they do not necessarily induce fully predicable

changes in diffusion and distribution at the nuclear envelope. In effect, via its flexible IDR, emerin

appears to mediate complex interactions with itself, lamin A/C, nuclear actin and BAF, where

binding to one partner impacts oligomerization and, interdependently, affects interactions with

other partners. As shown from the organization of wild-type emerin after nuclear actin depletion

and our study of the Q133H emerin mutant, binding to nuclear actin significantly influences the

mobility and the oligomerization potential of emerin. Compared to emerin, actin is relatively large

and its binding to the IDR of emerin could mask interaction domains that normally serve as LEM

binding and self-association sites between emerin monomers[18]. IDR masking, combined with the

reduced lateral mobility of actin-bound emerin, might therefore modulate the formation of

oligomers by limiting molecular collisions between emerin monomers. Consistent with the need

to precisely regulate emerin oligomerization for appropriate nuclear shape deformation, increased

emerin diffusion and controlled formation of emerin oligomers appear to be coupled molecular

events during nuclear adaptation against stress, as our measurements in micropatterns indicate.

The deficient binding of Q133H emerin to nuclear actin and its over-oligomerization at the INM

might thus be linked to the abnormal nuclear deformation observed in cells expressing this mutant.

The influence of emerin differential interactions with nucleoplasmic components on its mobility

and its oligomerization is also underlined by the surprisingly slow lateral diffusion of the Δ95-99

emerin mutant, which does not bind lamin A/C nor actin *in vitro*, but retains binding to BAF. The

48-118 region of emerin, where Δ95-99 is located, was suggested to act as a potential binding site

for the LEM domain[18] and through altered region flexibility, the deletion might reduce the efficacy

of such interactions, potentially causing the LEM domain to bind BAF more frequently. Repeated

interactions of Δ95-99 with BAF and with the lamina[41] or chromatin[42] via ternary BAF complexes,

could slow its diffusion, limit molecular collisions, and impede its oligomerization by sequestering

the LEM domain and reducing bridging interactions with LEM binding sites on other emerins[18].

The Δ95-99 deletion, within the 55-132 lamin tail-binding region of emerin[18], could also prevent

a stabilization of already sparse Δ95-99 oligomers by lamin A/C. As such, the aberrant nuclear

deformation against stress observed for cells expressing Δ95-99 might stem from out of balance

interactions of the deletion mutant with BAF and its inability to form lamin A/C-stabilized

oligomers.

A similar impediment of *in trans* interactions between the LEM domain and self-association sites

along the joined IDR of P183H dimers likely result in the slow mobility and the reduced ability of

this emerin mutant to oligomerize. The reported enhanced binding of lamin A/C to P183H

emerin[1,20], in close proximity to these same self-association sites could also interfere with inter-

emerin bridging interactions. In both cases, the inability of the LEM domain to access binding sites

along the IDR would elicit its repeated interactions with BAF, leading to the observed slow

diffusion of P183H at the INM. The defective nuclear shape adaptation to mechanical challenges

of cells expressing P183H might therefore stem from emerin dimerization and excessive

interactions with BAF or lamin A/C that prevent an efficient formation of emerin oligomers and a

remodeling of the nucleoskeleton.

As we have shown, Δ95-99 emerin does not self-assemble in dense oligomers at the INM and

induces aberrant nuclear shape remodeling on micropatterns. The phosphorylation of emerin

residues Tyr 74 and Tyr 95 by Scr kinase[51] was recently shown to mediate the recruitment of lamin

A/C to the LINC complex during nuclear stiffening in response to force[27]. Our observation that

Δ95-99 fails at promoting normal nuclear deformation against mechanical stress by its reduced

ability to form lamin A/C-stabilized emerin oligomers is consistent with the major role played by

Tyr 95 phosphorylation for emerin-mediated mechanotransduction at the nuclear envelope. The

abnormal organization of cytoskeletal actin in mechanically challenged cells expressing Δ95-99,

akin to disorganizations observed with a non-phosphorylable tyrosine 74-95FF emerin mutant[27],

also suggests that emerin oligomers are required to strengthen the connection between lamin A/C,

the LINC complex and cytoplasmic actin filaments in order to promote correct nucleus positioning

and deformation. This points towards a link between Tyr 95 phosphorylation, emerin

oligomerization and recruitment of lamin A/C for stiffening the nuclear envelope at LINC

complexes, that are likely driven by phosphorylation-induced changes in emerin conformation and

binding to nuclear partners[26,27,51]. It is thus possible that oligomeric nanodomains at the INM are

enriched in Tyr 95 phosphorylated emerin, lamin A/C and LINC complex components, while

emerin monomers populate the rest of the INM, consistent with emerin enrichments in distinct

nucleoskeletal "niches" at the nuclear envelope[26].

We propose that, during stress responses, transient unbinding of emerin monomers from BAF and

nuclear actin favors oligomerization by increasing lateral diffusion at the INM and exposing both

the LEM domain and self-association sites along the IDR for intermolecular binding between

emerins. Within emerin oligomers, stabilization by direct interaction with the lamin A/C and

additional modulation of emerin self-association by nuclear actin and BAF likely allow for a

precise regulation of the oligomerization potential and the size of oligomeric domains, as required

for adaptive nuclear deformation in response to forces. The localized INM distribution of emerin

oligomers and their reliance on a part of the emerin IDR that requires phosphorylation for the

recruitment of lamin A/C to the LINC complex[27], also suggest that they are sites where interactions

between emerin, lamin A/C and LINC components are strengthened. Emerin oligomerization

might therefore contribute to the increased connectivity between the nucleoskeleton, the nuclear

envelope and the cytoskeleton for anchoring the nucleus and for providing force absorption contact points during nuclear deformation. In response to surging mechanical stress, the incremental oligomerization of emerin in larger nanodomains we observed could be part of a mechanism that redistributes increasing forces over wider areas to maintain a basal membrane pressure at local anchoring points between the nucleoskeleton, the nuclear envelope and the cytoskeleton. Outside these anchoring oligomeric domains, disengagement of emerin monomers from nuclear actin likely provides additional modulations of nucleoskeletal contacts with the nuclear envelope, for instance by reducing the connections of the INM with the nucleoskeleton, while maintaining its connectivity with nuclear chromatin, via BAF. Indeed, both tethering of chromatin to the nuclear envelope and the nucleoskeleton participate in nuclear mechanics[5,52]. Together, strengthening the connections between the nuclear envelope and the lamina at specific cytoskeletal anchor points, but relaxing them in the rest of the membrane could provide a means to couple controlled nuclear deformation with nucleus positioning in cells. Such coupling is defective with emerin mutants, which display abnormal oligomerization and modified interactions with nucleoplasmic partners. The structural interdependency between emerin monomers, emerin oligomers, BAF, and key elements of the nucleoskeleton likely provides an integrated mechanism where INM/lamina contacts and INM/chromatin contacts are spatially regulated and where reinforced connectivity between the nuclear envelope and the cytoskeleton is provided where needed. At the center of such processes, emerin diffusion and monomer/oligomer exchanges ensure that mechanical cues are transduced throughout the nuclear envelope, for coordinated changes in local nuclear stiffness and appropriate remodeling of the nuclear shape.

METHODS

Cell culture, emerin expression and cell staining

Emerin-null human dermal fibroblast (*EMD*[-/y] HDF) and normal dermal fibroblasts (*EMD*[+/y] HDF) were kindly provided by Dr. Howard Worman, Columbia University, New York, USA. *EMD*[-/y] HDF are derived from a male EDMD patient (G-9054) and carry a 59 nucleotide deletion within the *EMD* gene (*EMD* g.329del59) that ablates emerin expression[22]. HDF were grown in DMEM (Lonza) with 10% fetal bovine serum (Gibco-Life Technologies), 50 units ml^{-1} penicillin and 50 µg ml^{-1} streptomycin and maintained at 37°C in a humidified atmosphere with 5% CO_2.

Human wild-type emerin cDNA was kindly provided by Dr. Juliet Ellis, University College
London, UK. For the expression of PA-TagRFP-emerin, a pEGFP-N1 plasmid backbone encoding
emerin fused to the C-terminus of PA-TagRFP was produced by XbaI and KpnI insertion and PCR
fusion of the human emerin cDNA. Cells plated on fibronectin-coated glass coverslips were
transfected with PA-TagRFP-emerin using X-tremeGENE HP (Roche). 48-72 hours post-
transfection, live cells were imaged by sptPALM in HBSS buffer at 37°C. For micropatterning
experiments, cells grown on 6-well plates were trypsinized after 48-72 hours of transfection and
plated on fibronectin-micropatterned coverslips.

To express SNAP-emerin, human emerin was first fused to the C-terminus of a SNAP tag by AscI
and XhoI insertion in a pSNAP-tag(m) plasmid (NEB). SNAP-emerin was then subcloned into a
modified pFUW lentiviral vector by NheI and AgeI insertion. Lentiviral particles for the
expression SNAP-emerin were produced by the UCLA Vector Core. Transduction of HDF grown
at 70% confluence on 6-well plates was done for 48 hours, using 25 ng ml^{-1} of lentiviral particles
in complete growth medium containing 8 μg ml^{-1} of polybrene, after which the medium was
replaced. Following another 24 hours incubation, cells were trypsinized, and plated on fibronectin-
coated or fibronectin-micropatterned coverslips. For imaging, cells were fixed with 4%
paraformaldehyde in PBS for 15 min, permeabilized with 0.1% Triton X-100 (Sigma-Aldrich) for
15 min and blocked with 4% bovine serum albumin (BSA, Sigma-Aldrich) + 0.1% Tween-20
(Sigma-Aldrich) for 30 min, at room temperature. Cells were then stained with 1 μM of SNAP-
Surface-AlexaFluor 647 (BG-A647, NEB) in 4% BSA + 0.1% Tween-20 for 1 hour at 37°C, then
thoroughly washed before super-resolution imaging.

For the expression of sGFP-emerin, humanized cDNA for split-GFP 1-10[39] was inserted by NheI
and XbaI digestion in the pEGFP-N1 plasmid backbone encoding emerin, and expressed as an N-
terminal fusion to emerin. The shorter 11[th] β-sheet M3 fragment[39] was also fused to the N-terminus
of emerin by PCR cloning using primers encoding the M3 fragment sequence and subcloning into
the pEGFP-N1 backbone via NheI and XhoI digestions. Both plasmids were co-transfected in HDF
using X-tremeGENE HP (Roche) as described for PA-TagRFP-emerin, and cells were imaged by
CALM 48-72 hours post-transfection. All constructs were verified by sequencing. Upon re-
expression of wild-type emerin fusions in *EMD*$^{-/y}$ HDF, nuclear deformations against mechanical
stress recover to levels similar to those of normal *EMD*$^{+/y}$ HDF[47].

For immunostaining of emerin, cells were grown on coverslips, fixed and permeabilized as

described for SNAP-emerin staining. Cells were labeled with a rabbit anti-emerin antibody (1:500,

Santa Cruz Biotechnology, sc-15378) for 45 min, washed in PBS and further labeled with a goat

anti-rabbit Alexa fluor 488 antibody (1:400, Invitrogen) for 45 min. After washing, cells were

mounted in a DAPI-Fluoromount G (Electron Microscopy Sciences) and imaged by confocal

microscopy.

To label cytoplasmic actin and measure nuclear shape indices, cells were fixed with 4%

paraformaldehyde in PBS for 15 min, permeabilized with 0.1% Triton X-100 for 10 min, and

blocked with 4% bovine serum albumin + 0.1% Tween-20 for 1 hour. Cells were stained with

phalloidin-iFluor 488 (1:1000, Abcam) for 1 hour, washed with PBS, mounted in DAPI-

Fluoromount G and imaged by confocal or wide field microscopy.

Mutations, siRNA and BAF L58R expression

Emerin mutations Q133H and P183H were introduced in PA-TagRFP-emerin, SNAP-emerin,

sGFP-emerin and M3-emerin by site-directed mutagenesis using QuickChange Lightning Site

Directed Mutagenesis (Agilent) and mutagenic primer pairs for:

Q133H: 5'-CGCTTTCCATCACCATGTGCATGATGA-3' and

606   5'-GATCGTCATCATGCACATGGTGATGGA-3'

607   P183H: 5'-CCTGTCCTATTATCATACTTCCTCCTC-3' and

608   5'-GTGGAGGAGGAAGTATGATAATAGGA-3'.

609   The Δ95-99 emerin deletion was produced using partially phosphorothioated PCR primers and T7

610   exonuclease digestion, as previously described[53]. Primers pairs for the Δ95-99 deletion were:

611   5'-GACTACTTCACCA*C*C*A*GGACTTAT-3' and

612   5'-GGTGAAGTAGTCG*T*C*A*TTGTAGCC-3',

where * denotes phosphorothioate modifications. All primers were obtained from Integrated DNA

Technologies (IDT) and all mutations were verified by sequencing.

siRNA duplex for BAF and Dicer siRNA for lamin A/C were obtained from IDT. The sequences

of sense nucleotides were as follows: BAF siRNA: 5'-AGAUUGCUAUUGUCGUACUUU-3',

lamin A/C DsiRNA: 5'-AGCUGAAAGCGCGCAAUACCAAGaa-3'. siRNA duplex for IPO9

was obtained from Ambion (id # S31299). All siRNA were transfected or co-transfected with

619 emerin plasmids at 25 nM using X-tremeGENE HP (Roche). When associated with lentiviral

620 expression of emerin, siRNA transfection was done 2 hours before viral titer application.

621 BAF L58R was expressed from an EGFP-BAF L58R lentiviral plasmid[44] (Addgene #101776) and

622 lentiviral particles were produced by the UCLA Vector Core. HDF cells were transduced with 25

623 ng ml^{-1} of lentiviral particles as described for SNAP-emerin.

Cell micropatterning and nuclear shape index measurements

626 HDF were micropatterned as described previously[32,47]. Briefly, hexamethyldisilazane-activated

627 glass coverslips (Marienfeld, #1.5, Ø25 mm) were stamped with rectangular and fibronectin-

628 coated polydimethylsiloxane stamps having lengths of 210 µm and widths of 15 µm, 10 µm, or 5

629 µm respectively. Cell attachment outside the fibronectin strips was blocked with a 1% solution of

630 Pluronic F-127. After attachment for 1 hour and removal of unattached cells, HDF were allowed

631 to spread out on the micropatterns for 6 hours at 37°C before being prepared for microscopy

632 imaging.

633 Using ImageJ[54], the nuclear shape index (NSI)[55] was determined by measuring the nuclear cross-

634 sectional area and the nuclear perimeter of DAPI-stained nuclei imaged by wide-field microscopy,

635 and by calculating the ratio:

636
$$\text{NSI} = \frac{4 \times \pi \times \text{area}}{\text{perimeter}^2} \quad (1)$$

638 The NSI measures the roundness of the nucleus with an NSI of 1 corresponding to a circular

639 nuclear shape. Mean NSI values ± standard deviation of the mean are reported for multiple cell

640 nuclei per condition.

Microscopy imaging

643 Confocal imaging of immunostained emerin in normal HDF, emerin-null HDF and emerin null

644 HDF after expression of wild-type PA-TagRFP-emerin were performed on an Olympus Fluoview

645 FV1000 confocal microscope equipped with a 60x/1.40 NA objective, 405 nm and 488 nm lasers

646 and appropriate emission filters for imaging DAPI (450/20 nm) and Alexa-488 labeled antibodies

647 against emerin (510/10 nm).

648 Confocal imaging of nuclear deformation and actin organization in micropatterned *EMD*$^{+/y}$ HDF

649 was performed on a Zeiss LSM 700 microscope equipped with a C-Apochromat 63×/1.15 W Korr

objective, excitation lasers at 405 nm and 488 nm and appropriate beamsplitter and emission
channel settings for dual detection of DAPI and phalloidin-iFluor 488.

Wide-field imaging of labeled actin and labeled nuclei for NSI measurements were performed on
an inverted Nikon Eclipse Ti-E microscope equipped with a 40x objective (Nikon), an iXon Ultra
EMCCD camera (Andor), 405 nm and 488 nm lasers, a T495lpxr dichroic mirror and a 525/50
emission filter (Chroma) for phalloidin-iFluor 488 or a 458Di02 dichroic mirror and a 483/32
emission filter (Semrock) for DAPI.

SptPALM, dSTORM and CALM imaging were performed on an inverted Nikon Eclipse Ti-E
microscope equipped with a 100x 1.49 NA objective (Nikon), an iXon EMCCD camera (Andor),
perfect focus drift compensation optics, an astigmatic lens for 3D super-resolution imaging, a piezo
z-scanner for calibration of 3D super-resolution images (Mad City Labs), laser lines at 405, 488,
561 and 647 nm (Agilent), a multiband pass ZET405/488/561/647x excitation filter (Chroma), a
quad-band ZT405/488/561/647 dichroic mirror (Chroma) and appropriate emission filters for
sptPALM imaging of PA-tagRFP (600/50 nm, Chroma), 3D-dSTORM imaging of Alexa-647
(700/75 nm, Chroma) and CALM imaging of complemented split-GFP (525/50 nm, Semrock).

sptPALM of PA-TagRFP-emerin was performed in 37ºC HBSS buffer (Corning) by HILO
excitation of the bottom nuclear membrane of cells with a continuous and low power
photoactivation at 405 nm and an excitation at 561 nm. The HILO illumination angle was $\theta = 51.6º$.
Images were acquired continuously at a frame rate of 40 ms per frame for no longer than 3 minutes
per cell to limit UV damage. CALM imaging of complemented emerin-GFP-emerin species was
done as described for sptPALM but with a single HILO excitation at 488 nm.

dSTORM of SNAP-emerin labeled with BG-A647 was performed at room temperature in a
photoswitching buffer composed of 10% glucose, 0.5 mg ml^{-1} glucose oxidase (Sigma), 40 μg ml^{-1}
catalase (Sigma), and 1% β-mercaptoethanol (Sigma). Continuous photoswitching was achieved
with a low power 488 nm laser and imaging was done with a 647 nm laser excitation at a frame
rate of 80 ms per frame. Z-calibration and sample drift corrections were done using a few 40 nm
TransFluoSphere beads (488/685 nm, Life Technologies) as fiducial markers spread on the cell
samples.

Analyses of diffusion coefficients

Localization and tracking analyses were performed using the software package SLIMfast, which

uses multiple-target tracing algorithms[56] and was kindly provided by Christian Ritcher and Jacob

Piehler. Localizations were performed by 2D-gaussian fitting of the point-spread-function of each

activated PA-TagRFP-emerin or activated emerin-GFP-emerin species in each frame. Localization

precision was determined as previously described[57], and individual PA-TagRFP-emerin were

localized with a precision of 13±5 nm. Diffusion trajectories were built by linking localizations

frame to frame and accounting for blinking statistics and local particle densities. Trajectories with

fewer than three steps were discarded. Diffusion coefficients were estimated using a probability

density of square displacement (PDSD) analysis[35]. For each time lag t, the PDSD curve was fitted

with the following model:

691
$$P(\vec{r}^2, t) = 1 - \sum_{i=1}^{n} a_i(t)e^{-r^2/r_i^2(t)} \qquad (2)$$

692
$$\sum_{i=1}^{n} a_i(t) = 1$$

where $r_i^2(t)$ is the square displacement and $a_i(t)$ is the population density of i numbers of diffusive

behaviors at each time lag t. To limit the risks of overfitting or underfitting the PDSD curves and

select an appropriate model for i numbers of diffusive behaviors in each data set, we used both an

Akaike information criterion (AIC) and a Bayesian information criterion (BIC) after fitting PDSD

with models where $1 \leq i \geq 5$. Square displacement curves ($r_i^2(t)$) were extracted from PDSD

analyses and reported with error bars determined using $\frac{r_i^2}{\sqrt{N}}$, where N is the number of analyzed

trajectories per time lag, as previously described[35]. Diffusion coefficients (D) representative of

each behaviors were determined by fitting each $r_i^2(t)$ curves over the first four time lags (t_1-t_4)

using OriginPro 2020 software (OriginLab) and a 2D Brownian diffusion model with position

error:

704
$$r^2 = 4Dt + 4\sigma^2 \qquad (3)$$

All diffusion coefficients D are reported in $\mu m^2\ s^{-1}$ ± standard error of fit value (± s.e.m.). Statistical

comparisons between D values were done using F-tests. Population percentages are derived from

the averaged $a_i(t)$ values over the considered time lags.

Individual diffusion coefficients (D_i) were obtained by fitting the individual mean square

displacement (MSD) for each detected emerin over the first three time lags (t_1-t_3), using again a

2D Brownian diffusion model. Based on their individual D_i value, emerin trajectories were

grouped into four diffusion ranges ($D_{i1} > 0.1 \ \mu m^2 \ s^{-1}$, $0.1 < D_{i2} > 0.01 \ \mu m^2 \ s^{-1}$, $0.1 < D_{i3} > 0.001 \ \mu m^2$

s^{-1}, and $D_{i4} < 0.001 \ \mu m^2 \ s^{-1}$) and plotted as maps.

Spatial distribution and cluster analyses from super-resolution images

After 3D-dSTORM super-resolution imaging, the localization of individual emerin molecules and

z-position assignments were performed by Gaussian fitting using rapidSTORM (version 3.3.1)[58].

Sample drift and overcounting corrections for molecules appearing in consecutive frames were

done using PALMsiever[59] and renderings of super-resolved images were done using ImageJ[54].

Localization precisions (σ) in the x, y, and z dimensions were evaluated as previously described[32]

and were σ_x:8.3 nm, σ_y:13.0 nm, and σ_z:28.4 nm. 2D spatial pattern analyses of emerin distributions

were performed on 2 µm x 2 µm regions of interest (ROI) typically chosen in nuclear envelope

areas having homogenous z ranges and away from the edges of nuclei, in order to limit 3D effects.

Emerin clustering was determined using an edge-corrected neighborhood density function (NDF)

as previously described[32]. Briefly, the NDF is a pairwise-correlation function similar to O-ring

statistics that tallies the density of detected emerin within a ring of outer radius r and width Δr

located at a distance r from an emerin position in the ROI and for all $r + \Delta r$ in the ROI. The density

of emerin as a function of distance from an average emerin was obtained with:

729
$$D_r = \frac{\sum N_r}{\sum A_r} \qquad (4)$$

where N_r is the number of neighbors and A_r is the area summed over all detected emerin. NDF

analyses were done over a 1µm distance on selected ROIs and with a fixed ring width of 10 nm

and a ring radius increasing by 10 nm steps. To average NDF statistics from multiple ROIs across

different nuclei and make them sample-size independent, D_r was further standardized by dividing

it by the mean density of detected emerin across the entire ROI. As such, an NDF value at a given

radius indicates the relative clustering of emerin as compared to the average density across the

entire sample. This relative NDF gives a value of 1 for a completely random spatial distribution as

determined by Monte Carlo simulations of random emerin distributions with area and number of

randomly seeded emerin equal to that of each experimental ROIs.

Relative NDF curves averaged across multiple ROIs and multiple nuclei were fitted with a

previously described model[32], which accounts for a distribution of cluster lengths that includes two

populations of emerin (monomer and oligomers) and for a probability density of emerin in 2D

clusters that decays approximately as an exponential function[45]:

$$743 \quad \text{Relative NDF} = \{A_1 \times \exp\left(\frac{-r}{\varepsilon_1}\right) + A_2 \times \exp\left(\frac{-r}{\varepsilon_2}\right) + 1\} * g(r)^{PSF} \quad (5)$$

where, A is the clustering density, ε is the typical half-maximum cluster length, $*$ denotes a 2D

convolution, and $g(r)^{PSF}$ is the correlation function of the effective point spread function of

uncertainty in position determination for the dSTORM experiments. As described previously[32],

$g(r)^{PSF}$ corrects the NDF for contribution of multiple single molecule appearances (blinking) to

the overall spatial distribution. After fitting relative NDF curves, the molecular density above

random for emerin clusters are reported as $A \pm$ standard error of the fit ($\pm$s.e.m) and their typical

length scales as $2 \times \varepsilon \pm$ standard error of the fit and localization precision ($\pm$s.e.m). Relative

increases in emerin oligomer formation during nuclear stress were determined by considering a

circular shape of oligomer nanodomains and multiplying the area of oligomerization by the

measured molecular density.

Cluster maps

Cluster maps were generated from drift- and overcounting-corrected super-resolved emerin

positions by determining local cluster values around each emerin using the Getis and Franklin L

function[46] in spPack[60] and for a distance of 25 nm. Spatial positions in x and y, and cluster values

were plotted as maps in MATLAB (MathWorks) using the meshgrid and griddata functions, a 1

nm x 1 nm pixel size and the 'v4' option when calculating pixel density values. The contour maps

were generated using the countourf function with 200 levels. In contour maps, values $L(r_{25})=25$

represent areas where emerin is randomly distributed and $L(r_{25})=70$ values represent areas with

emerin local density $(70/25)^2 = \sim$8-fold higher than expected for a random distribution.

ACKNOWLEDGMENTS

This work was supported by the National Institute of Arthritis and Musculoskeletal and Skin
Diseases of the National Institutes of Health under award number R21AR076514. We are grateful
to H. Worman for providing emerin null and normal human dermal fibroblasts and to J. Ellis for
providing human emerin cDNA.

772
AUTHOR CONTRIBUTIONS

A.F. and M.B. designed and performed experiments, contributed reagents and analyses, and participated in writing of the manuscript. F.P. conceived and supervised the research, designed and performed experiments, contributed analyses and wrote the manuscript.

COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests

    Figure 1

Figure 1. Emerin displays four distinct diffusive behaviors at endoplasmic reticulum membrane
and the nuclear envelope. a) Immunostaining and confocal imaging of emerin in emerin-null HDF
($EMD^{-/y}$), HDF with endogenous emerin expression ($EMD^{+/y}$) and $EMD^{-/y}$ HDF after expression
of wild-type PA-TagRFP-emerin. Scale: 10 µm. b) Map of emerin trajectories built from
localization and tracking of individual PA-TagRFP-emerin at the bottom nuclear envelope. Scale:
5 µm. c) Probability distribution of square displacement (PDSD) for wild-type emerin (71004
trajectories in 14 cells) at time lag $\Delta t = 160$ ms. The distribution is composed of four diffusive
behaviors. d) Square displacement curves (r^2_i) for each emerin diffusive behavior generated from
PDSD analyses on the first ten time lags Δt 40-400 ms. Curves are fitted over the first four values
with a Brownian diffusion model (red line) to estimate the respective diffusion coefficients of each
emerin behaviors. e) Separated emerin trajectory maps as a function of individual diffusion
coefficient (D_i). Scale: 5 µm.
799

Figure 2

Figure 2. Emerin diffuses as inner nuclear membrane monomers and oligomers that interact with lamin A/C, nuclear actin and BAF. a) Diffusion coefficients (± s.e.m.) and population percentages of wild-type emerin (untreated, *n*=71004 trajectories in 14 nuclei) after depletion of lamin A/C (lamin A/C KD, *n*=60569 trajectories in 11 nuclei), depletion of nuclear actin (IPO9 KD, *n*=74501 trajectories in 17 nuclei) or replacement of endogenous BAF with BAF L58R mutant (BAF KD + BAF L68R, *n*=62714 trajectories in 8 nuclei) (F-test, ns: non-significant, **: *p*<0.01). b) Schematic of emerin fused to complementary split-GFP fragments and co-expressed to track the mobility of emerin oligomers by single molecule imaging of complemented GFP. c) Map of individual trajectories for complemented emerin-GFP-emerin species at the nuclear envelope (left) and examples of oligomer domains (squares) where trajectories often overlap (left). Scales: 2 µm (left) and 50 nm (right). d) Comparison of diffusion coefficients (± s.e.m.) and population percentages for individual wild-type emerin assessed by sptPALM and complemented emerin-GFP-emerin species (n=4833 trajectories in 13 nuclei) assessed by CALM (F-test, ns: non-significant). e) Diffusion map of wild-type PA-TagRFP-emerin at the nuclear envelope where emerin oligomers form slow mobility domains (blue, O) surrounded by larger areas where emerin monomers diffuse faster (red, M). Scale: 500 nm.

 Figure 3

Figure 3. Emerin forms oligomeric nanodomains stabilized by lamin A/C and surrounded by
larger emerin monomer areas across the nuclear membrane. a) Super-resolved 3D-dSTORM
positions (top half) and non-super-resolved imaging (bottom half) of wild-type SNAP-emerin at
the bottom nuclear envelope of an *EMD*$^{-/y}$ HDF. Scale: 5 µm. b) Neighborhood densities (± s.d.)
and cluster analyses of wild-type emerin at the nuclear membrane of untreated HDF (n=189,331
localizations in 10 nuclei) and HDF after lamin A/C knock down (n=178,206 localizations in 6
nuclei). Neighborhood densities at various length scales are compared to MonteCarlo simulations
of complete spatial randomness and fitted to measure molecular densities above random and length
scale of significant clustering (green). c) Relative nuclear envelope molecular densities above
random (± s.e.m.) for wild-type emerin oligomers (O) and monomers (M) in untreated *EMD*$^{-/y}$

HDF, HDF depleted for lamin A/C, HDF depleted for nuclear actin after IPO9 knock down
(n=225,394 localizations in 9 nuclei) and HDF where endogenous BAF is replaced by BAF L58R
(n=90,241 localizations in 6 nuclei). Values in parenthesis represent the typical length scale ($\pm$
s.e.m.) of each domain in nanometers. d) Local cluster maps of wild-type emerin in nuclear
envelope regions of untreated (left) and lamin A/C knocked down HDF (right) for a search radius
of 25 nm ($L(r_{25})$). Cluster values in maps are assigned to all localized emerin, where values
$L(r_{25})$=25 represent areas where emerin is randomly distributed and $L(r_{25})$=70 values represent
areas with emerin local densities $(70/25)^2 = \sim$8-fold higher than expected for a random distribution.
M: monomer areas, O: oligomer nanodomains. Scale: 250 nm.
839

Figure 4

Figure 4. Mechanical stress on the nucleus increases emerin mobility and the formation of emerin oligomers. a) Schematic for increasing mechanical stress on the nucleus by cell micropatterning on rectangular fibronectin strips with decreasing widths of 15, 10 and 5 µm. Arrows represent forces. b) Fluorescence confocal imaging of actin (green) and the nucleus (blue) in micropatterned $EMD^{+/y}$ HDF. Scale: 50 µm. c) Nuclear shape index as a function of micropattern width for untreated $EMD^{+/y}$ HDF (n=54, 70, 61 and 62 nuclei for non-patterned, 15, 10 and 5 µm wide patterns, respectively), HDF depleted for lamin A/C (n=75, 60, 70 and 66 nuclei) and HDF depleted for nuclear actin (n=62, 71, 66 and 46 nuclei). Box length: index interquartile range; central square: mean; central bar: median; error bars: ± s.d. (T-test, ** p < 0.01). d) Diffusion coefficients (± s.e.m.) and population percentages of wild-type emerin in non-patterned cells or after nuclear deformation on 15 µm wide (n=27266 trajectories in 10 nuclei) and 10 µm wide micropatterns (n=12915 trajectories 8 nuclei) (F-test, ns: non-significant, **: p<0.01. e) Relative nuclear envelope molecular densities above random (± s.e.m.) for wild-type emerin oligomers (O) and monomers (M) in non-patterned $EMD^{-/y}$ HDF or after nuclear deformation on 15 µm wide (n=151,647 localizations in 10 nuclei) or 10 µm wide micropatterns (n=56,563 localizations in 6 nuclei). Values in parenthesis represent the typical length scale (± s.e.m.) of each domain in nanometers. f) Increase in emerin oligomers as a function of nuclear stress on 15 and 10 µm wide micropatterns compared to non-deformed nuclei in non-patterned cells.

  Figure 5

Figure 5. Emerin mutations induce defective nuclear shape adaptation against mechanical stress. a) Diagram of wild-type emerin (WT) showing its transmembrane (TM) and LEM domains, its disordered central region with relevant binding domains and self-association domain (self-ass.) and the position of Δ95-99, Q133H and P183H EDMD-inducing mutations. b) Nuclear shape index as a function of micropattern width for *EMD*$^{-/y}$ HDF expressing wild-type emerin (n=38, 33, 26 and 57 nuclei for non-patterned, 15, 10 and 5 µm wide patterns, respectively), Q133H emerin (n= 74, 58 and 37 nuclei), Δ95-99 emerin (n= 64, 89, 45 and 46 nuclei) and P183H emerin (n=82, 78 and 28 nuclei). Box length: index interquartile range; central square: mean; central bar: median; error bars: ± s.d. (T-test, ns: non-significant, *: *p*<0.05, **: *p*<0.01. c) Fluorescence wide-field imaging of actin (green) and the nucleus (blue) in micropatterned *EMD*$^{-/y}$ HDF expressing Q133H, Δ95-99 or P183H emerin mutants. Scales: 50 µm.

Figure 6

Figure 6. Emerin mutants exhibit modified lateral mobilities and defective oligomerization at the inner nuclear membrane. a) Diffusion coefficients (± s.e.m.) and population percentages of Q133H emerin (*n*=105050 trajectories in 13 nuclei) compared to wild-type emerin (n=71004 trajectories in 14 nuclei) (F-test, ns: non-significant, *: *p*<0.05, **: *p*<0.01). b) Relative nuclear envelope molecular densities above random (± s.e.m.) for Q133H emerin oligomers (O) and monomers (M) (n=149,340 localizations in 6 nuclei) compared to wild-type emerin (n=189,331 localizations in 10 nuclei). c) Local cluster map of Q133H emerin at the nuclear envelope for a search radius of 25 nm (L(r$_{25}$)). M: monomer areas, O: oligomer nanodomains. Scale: 250 nm. d) Diffusion coefficients (± s.e.m.) and population percentages of Δ95-99 emerin (*n*=76944 trajectories in 14 nuclei) compared to wild-type emerin (F-test, ns: non-significant, *: *p*<0.05, **: *p*<0.01. e) Relative nuclear envelope molecular densities above random (± s.e.m.) for Δ95-99 emerin oligomers (O) and monomers (M) (n=208,092 localizations in 8 nuclei) compared to wild-type emerin. f) Local cluster map of Δ95-99 emerin. M: monomer areas, O: oligomer nanodomains. Scale: 250 nm. g) Diffusion coefficients (± s.e.m.) and population percentages of P183H emerin

(n=86529 trajectories in 21 nuclei) compared to wild-type emerin and compared to complemented
P183H emerin-GFP-emerin species (n=10519 trajectories in 21 nuclei) assessed by CALM (F-test,
ns: non-significant, *: $p<0.05$, **: $p<0.01$). h) Relative nuclear envelope molecular densities above
random (± s.e.m.) for P183H emerin oligomers (O) and monomers/dimers (M/D) (n=138,075
localizations in 6 nuclei) compared to wild-type emerin. i) Local cluster map of P183H emerin.
M/D: monomer/dimer areas, O: oligomer nanodomains. Scale: 250 nm. Values in parenthesis in
b, e and h represent the typical length scale (± s.e.m.) of each domain in nanometers.
903

Figure 7

Figure 7. Insufficient oligomerization of EDMD-inducing Δ95-99 emerin mutant against
mechanical stress. a) Relative nuclear envelope molecular densities above random (± s.e.m.) for
Δ95-99 emerin oligomers (O) and monomers (M) in non-patterned *EMD*$^{-/y}$ HDF or after nuclear
deformation on 15 μm wide (n=138,119 localizations in 5 nuclei) or 10 μm wide micropatterns
(n=135,143 localizations in 6 nuclei). Values in parenthesis represent the typical length scale (±
s.e.m.) of each domain in nanometers. b) Increase in Δ95-99 emerin oligomers as a function of
nuclear stress on 15 and 10 μm wide micropatterns compared to non-deformed nuclei in non-
patterned cells. c) Local cluster maps of wild-type and Δ95-99 emerin after nuclear deformation
on 15 μm wide micropatterns. M: monomer areas, O: oligomer nanodomains. Scale: 250 nm.

REFERENCES

1. Berk, J.M., Tifft, K.E. & Wilson, K.L. The nuclear envelope LEM-domain protein emerin. *Nucleus* **4**, 298-314 (2013).

2. Kirby, T.J. & Lammerding, J. Emerging views of the nucleus as a cellular mechanosensor. *Nature Cell Biology* **20**, 373-381 (2018).

3. Cui, Y. et al. Cyclic stretching of soft substrates induces spreading and growth. *Nature Communications* **6**, 6333 (2015).

4. Demmerle, J., Koch, A.J. & Holaska, J.M. The nuclear envelope protein emerin binds directly to histone deacetylase 3 (HDAC3) and activates HDAC3 activity. *The Journal of Biological Chemistry* **287**, 22080-8 (2012).

5. Stephens, A.D., Banigan, E.J., Adam, S.A., Goldman, R.D. & Marko, J.F. Chromatin and lamin A determine two different mechanical response regimes of the cell nucleus. *Molecular Biology of the Cell* **28**, 1984-1996 (2017).

6. Maurer, M. & Lammerding, J. The Driving Force: Nuclear Mechanotransduction in Cellular Function, Fate, and Disease. *Annual Review of Biomedical Engineering* **21**, 443-468 (2019).

7. Emery, A.E. Emery-Dreifuss muscular dystrophy - a 40 year retrospective. *Neuromuscular Disorders* **10**, 228-32 (2000).

8. Emery, A.E. & Dreifuss, F.E. Unusual type of benign x-linked muscular dystrophy. *J Neurol Neurosurg Psychiatry* **29**, 338-42 (1966).

9. Maggi, L., Carboni, N. & Bernasconi, P. Skeletal Muscle Laminopathies: A Review of Clinical and Molecular Features. *Cells* **5**(2016).

10. Frock, R.L. et al. Lamin A/C and emerin are critical for skeletal muscle satellite cell differentiation. *Genes & Development* **20**, 486-500 (2006).

11. Fidzianska, A. & Hausmanowa-Petrusewicz, I. Architectural abnormalities in muscle nuclei. Ultrastructural differences between X-linked and autosomal dominant forms of EDMD. *Journal of the Neurological Sciences* **210**, 47-51 (2003).

12. Park, Y.-E. et al. Nuclear changes in skeletal muscle extend to satellite cells in autosomal dominant Emery-Dreifuss muscular dystrophy/limb-girdle muscular dystrophy 1B. *Neuromuscular Disorders* **19**, 29-36 (2009).

13. Rowat, A.C., Lammerding, J. & Ipsen, J.H. Mechanical properties of the cell nucleus and the effect of emerin deficiency. *Biophysical Journal* **91**, 4649-64 (2006).

14. Lammerding, J. et al. Abnormal nuclear shape and impaired mechanotransduction in emerin-deficient cells. *Journal of Cell Biology* **170**, 781-91 (2005).

15. Yuan, J. & Xue, B. Role of structural flexibility in the evolution of emerin. *Journal of Theoretical Biology* **385**, 102-11 (2015).

16. Clements, L., Manilal, S., Love, D.R. & Morris, G.E. Direct interaction between emerin and lamin A. *Biochemical and Biophysical Research Communications* **267**, 709-14 (2000).

17. Sakaki, M. et al. Interaction between emerin and nuclear lamins. *Journal of Biochemistry* **129**, 321-7 (2001).

18. Berk, J.M. et al. The molecular basis of emerin-emerin and emerin-BAF interactions. *Journal of Cell Science* **127**, 3956-69 (2014).

19. Fairley, E.A., Kendrick-Jones, J. & Ellis, J.A. The Emery-Dreifuss muscular dystrophy
phenotype arises from aberrant targeting and binding of emerin at the inner nuclear membrane.
Journal of Cell Science **112 (Pt 15)**, 2571-82 (1999).

20. Lee, K.K. et al. Distinct functional domains in emerin bind lamin A and DNA-bridging protein
BAF. *Journal of Cell Science* **114**, 4567-73 (2001).

21. Mislow, J.M.K. et al. Nesprin-1α self-associates and binds directly to emerin and lamin A in
vitro. *FEBS Letters* **525**, 135-140 (2002).

22. Haque, F. et al. Mammalian SUN protein interaction networks at the inner nuclear membrane
and their role in laminopathy disease processes. *Journal of Biological Chemistry* **285**, 3487-98
(2010).

23. Herrada, I. et al. Muscular Dystrophy Mutations Impair the Nuclear Envelope Emerin Self-
assembly Properties. *ACS Chemical Biology* **10**, 2733-42 (2015).

24. Samson, C. et al. Emerin self-assembly mechanism: role of the LEM domain. *FEBS Journal*
284, 338-352 (2017).

25. Roberts, R.C. et al. The Emery-Dreifuss muscular dystrophy associated-protein emerin is
phosphorylated on serine 49 by protein kinase A. *FEBS Journal* **273**, 4562-75 (2006).

26. Berk, J.M. et al. O-Linked beta-N-acetylglucosamine (O-GlcNAc) regulates emerin binding to
barrier to autointegration factor (BAF) in a chromatin- and lamin B-enriched "niche". *Journal*
of Biological Chemistry **288**, 30192-209 (2013).

27. Guilluy, C. et al. Isolated nuclei adapt to force and reveal a mechanotransduction pathway in
the nucleus. *Nature Cell Biology* **16**, 376-81 (2014).

28. Hirano, Y. et al. Dissociation of Emerin from Barrier-to-autointegration Factor Is Regulated
through Mitotic Phosphorylation of Emerin in a Xenopus Egg Cell-free System*. *Journal of*
Biological Chemistry **280**, 39925-39933 (2005).

29. Yates, J.R. & Wehnert, M. The Emery-Dreifuss Muscular Dystrophy Mutation Database.
Neuromuscular Disorders **9**, 199 (1999).

30. Ellis, J.A., Craxton, M., Yates, J.R. & Kendrick-Jones, J. Aberrant intracellular targeting and
cell cycle-dependent phosphorylation of emerin contribute to the Emery-Dreifuss muscular
dystrophy phenotype. *Journal of Cell Science* **111 (Pt 6)**, 781-92 (1998).

31. Holt, I., Clements, L., Manilal, S. & Morris, G.E. How does a g993t mutation in the emerin
gene cause Emery-Dreifuss muscular dystrophy? *Biochemical and Biophysical Research*
Communications **287**, 1129-33 (2001).

32. Fernandez, A., Bautista, M., Stanciauskas, R., Chung, T. & Pinaud, F. Cell-Shaping
Micropatterns for Quantitative Super-Resolution Microscopy Imaging of Membrane
Mechanosensing Proteins. *ACS Applied Materials & Interfaces* **9**, 27575-27586 (2017).

33. Talkop, U.A. et al. Early onset of cardiomyopathy in two brothers with X-linked Emery-
Dreifuss muscular dystrophy. *Neuromuscul Disord* **12**, 878-81 (2002).

34. Tokunaga, M., Imamoto, N. & Sakata-Sogawa, K. Highly inclined thin illumination enables
clear single-molecule imaging in cells. *Nature Methods* **5**, 159-61 (2008).

35. Schutz, G.J., Schindler, H. & Schmidt, T. Single-molecule microscopy on model membranes
reveals anomalous diffusion. *Biophysical Journal* **73**, 1073-80 (1997).

36. Ostlund, C., Ellenberg, J., Hallberg, E., Lippincott-Schwartz, J. & Worman, H.J. Intracellular trafficking of emerin, the Emery-Dreifuss muscular dystrophy protein. *Journal of Cell Science* **112 (Pt 11)**, 1709-19 (1999).

37. Dopie, J., Skarp, K.P., Rajakyla, E.K., Tanhuanpaa, K. & Vartiainen, M.K. Active maintenance of nuclear actin by importin 9 supports transcription. *Proceedings of the National Academy of Sciences of the United States of America* **109**, E544-52 (2012).

38. Cabantous, S., Terwilliger, T.C. & Waldo, G.S. Protein tagging and detection with engineered self-assembling fragments of green fluorescent protein. *Nature Biotechnology* **23**, 102-107 (2005).

39. Pinaud, F. & Dahan, M. Targeting and imaging single biomolecules in living cells by complementation-activated light microscopy with split-fluorescent proteins. *Proceedings of the National Academy of Sciences* **108**, E201-E210 (2011).

40. Iino, R., Koyama, I. & Kusumi, A. Single Molecule Imaging of Green Fluorescent Proteins in Living Cells: E-Cadherin Forms Oligomers on the Free Cell Surface. *Biophysical Journal* **80**, 2667-2677 (2001).

41. Samson, C. et al. Structural analysis of the ternary complex between lamin A/C, BAF and emerin identifies an interface disrupted in autosomal recessive progeroid diseases. *Nucleic Acids Research* **46**, 10460-10473 (2018).

42. Burger, M., Schmitt-Koopmann, C. & Leroux, J.-C. DNA unchained: two assays to discover and study inhibitors of the DNA clustering function of barrier-to-autointegration factor. *Scientific Reports* **10**, 12301 (2020).

43. Shimi, T. et al. Dynamic interaction between BAF and emerin revealed by FRAP, FLIP, and FRET analyses in living HeLa cells. *Journal of Structural Biology* **147**, 31-41 (2004).

44. Samwer, M. et al. DNA Cross-Bridging Shapes a Single Nucleus from a Set of Mitotic Chromosomes. *Cell* **170**, 956-972 e23 (2017).

45. Sengupta, P. et al. Probing protein heterogeneity in the plasma membrane using PALM and pair correlation analysis. *Nature Methods* **8**, 969-75 (2011).

46. Getis, A. & Franklin, J. Second-Order Neighborhood Analysis of Mapped Point Patterns. *Ecology* **68**, 473-477 (1987).

47. Bautista, M., Fernandez, A. & Pinaud, F. A Micropatterning Strategy to Study Nuclear Mechanotransduction in Cells. *Micromachines (Basel)* **10**(2019).

48. Holaska, J.M., Kowalski, A.K. & Wilson, K.L. Emerin caps the pointed end of actin filaments: evidence for an actin cortical network at the nuclear inner membrane. *PLoS Biology* **2**, E231 (2004).

49. Bengtsson, L. & Wilson, K.L. Multiple and surprising new functions for emerin, a nuclear membrane protein. *Current Opinion in Cell Biology* **16**, 73-9 (2004).

50. Ho, C.Y., Jaalouk, D.E., Vartiainen, M.K. & Lammerding, J. Lamin A/C and emerin regulate MKL1–SRF activity by modulating actin dynamics. *Nature* **497**, 507-511 (2013).

51. Tifft, K.E., Bradbury, K.A. & Wilson, K.L. Tyrosine phosphorylation of nuclear-membrane protein emerin by Src, Abl and other kinases. *Journal of Cell Science* **122**, 3780-90 (2009).

52. Schreiner, S.M., Koo, P.K., Zhao, Y., Mochrie, S.G.J. & King, M.C. The tethering of chromatin to the nuclear envelope supports nuclear mechanics. *Nature Communications* **6**, 7159 (2015).

53. Stoynova, L., Solórzano, R. & Collins, E.D. Generation of large deletion mutants from plasmid
DNA. *Biotechniques* **36**, 402-4, 406 (2004).

54. Schneider, C.A., Rasband, W.S. & Eliceiri, K.W. NIH Image to ImageJ: 25 years of image
analysis. *Nature Methods* **9**, 671-5 (2012).

55. Versaevel, M., Grevesse, T. & Gabriele, S. Spatial coordination between cell and nuclear shape
within micropatterned endothelial cells. *Nature Communications* **3**, 671 (2012).

56. Serge, A., Bertaux, N., Rigneault, H. & Marguet, D. Dynamic multiple-target tracing to probe
spatiotemporal cartography of cell membranes. *Nature Methods* **5**, 687-94 (2008).

57. Thompson, R.E., Larson, D.R. & Webb, W.W. Precise nanometer localization analysis for
individual fluorescent probes. *Biophysical Journal* **82**, 2775-83 (2002).

58. Wolter, S. et al. rapidSTORM: accurate, fast open-source software for localization microscopy.
Nature Methods **9**, 1040-1 (2012).

59. Pengo, T., Holden, S.J. & Manley, S. PALMsiever: a tool to turn raw data into results for
single-molecule localization microscopy. *Bioinformatics* **31**, 797-8 (2015).

60. Perry, G.L.W. SpPack: spatial point pattern analysis in Excel using Visual Basic for
Applications (VBA). *Environmental Modelling & Software* **19**, 559-569 (2004).