1 Impact of an oral amino acid provision on Achilles peritendinous amino acid concentrations in
2 young and older adults
3
4
5

Chad C. Carroll, PhD[1]
Samantha Couture, MS[1]
Dominick O. Farino[1]
Shivam H. Patel, PhD[1]
Nathan W.C. Campbell[1]
Julianne Stout, MD[2]
Arman Sabbaghi, PhD[3]
18 [1]Department of Health and Kinesiology, Purdue University, West Lafayette, IN
19 [2]College of Veterinary Medicine, Purdue University, West Lafayette, IN
20 [3]Department of Statistics, Purdue University, West Lafayette, IN

21
22
23
24 Running Title: Achilles peritendinous amino acid concentrations in humans
25
26
27

28 Address for correspondence:
29
Chad C. Carroll, PhD
Assistant Professor
Purdue University
Department of Health and Kinesiology
800 W. Stadium Ave
West Lafayette, IN 47907
Phone: (765) 496-6002
carrol71@purdue.edu

38
39

40   **ABSTRACT**
41
Recent studies have indicated that consumption of amino acid-rich compounds can increase

tendon collagen content and enhance biomechanical function. Still, it is not clear as to what

extent oral consumption of amino acids alters peritendinous amino acid concentrations.

Whether aging alters the delivery of amino acids to tendon tissue after oral consumption is also

not known. Using microdialysis, we determined the impact of a single oral essential amino acid

bolus on Achilles peritendinous amino acid concentrations in younger (n=7; 27±1 yr.) and older

adults (n=6; 68±2 yrs.) over four hours. The peritendinous concentration of all amino acids in the

beverage except methionine (p=0.136) and glycine (p=0.087) increased with time (p<0.05).

Additionally, the concentrations of glycine and arginine were greater in older adults (p≤0.05).

We also accessed the impact of amino acid consumption on peritendinous concentrations of

pro-collagen Iα1, a marker of collagen synthesis. Pro-collagen Iα1 tended to change with time

(p=0.071) but was not altered age (p=0.226). We demonstrate that an oral amino acid bolus

leads to modest increases in Achilles peritendinous amino acid concentrations in young and

older adults. The concentration of some amino acids was also greater in older adults. However,

the amino acid bolus did not significantly impact peritendinous pro-collagen concentrations.

Keywords: tendon, amino acids, exercise, pro-collagen, microdialysis
59

INTRODUCTION

Recent investigations have implied that tendons are responsive to amino acids. Oral consumption of amino acids may improve clinical outcomes in patients with tendinopathy (15, 30) and optimize tendon adaptations to exercise training (13). Work in rodents has implied that dietary amino acid supplementation can provide an effective means to promote recovery from tendon injury recovery and enhance collagen synthesis (28, 32). In contrast to skeletal muscle, the impact of amino acid consumption on the health of tendon tissue has not been extensively evaluated, especially in humans.

Tendon collagen content is greater in rats given a leucine-rich diet during recovery from malnourishment, suggesting that a diet rich in leucine could increase collagen synthesis. Additionally, a glycine-enriched diet improved tendon collagen content and biomechanical properties in rats after the induction of inflammation via collagenase injection (32). In support of these preclinical findings, patients treated with an arginine-rich oral supplement had modest improvements in pain (15, 25) and rotator cuff repair integrity (15) over those not consuming the supplement. Further, oral supplementation with glycine-rich collagen-peptides improved clinical outcomes in Achilles tendinopathy patients completing a calf-strengthening program (27).

Healthy tendons also appear to be responsive to dietary amino acid/protein-based interventions. Consumption of a vitamin C-enriched gelatin by healthy young men increased serum glycine, proline, and hydroxyproline concentrations, as well as collagen synthesis (30). The serum collected after subjects consumed the vitamin C-enriched gelatin was also used to treat engineered ligaments *ex vivo*. Treated ligaments had greater collagen content and superior biomechanical properties than ligaments treated with serum take before gelatin consumption (30). Consistent with the anabolic effect of leucine on skeletal muscle (10), tendon cross-sectional area (CSA) increased to a greater extent in young adults consuming a leucine-rich whey isolate during a resistance training program compared to a placebo group (13).

However, Achilles tendon stiffness was lower in mice given branched-chain amino acids during

an exercise intervention than mice on a standard diet (3).

Surprisingly, studies examining the impact of amino acid consumption on tendon properties

are limited to young and middle-aged adults. Age-related declines in tendon function (9) and

morphological properties (5, 8) and a higher incidence of tendinopathies with age (2, 20) make

older adults excellent candidates for amino acid interventions that are proposed to improve

tendon properties, yet critical knowledge gaps remain. The extent to which oral consumption of

amino acids increases the local delivery of amino acids to tendons is unknown. Knowing this

information could guide the development of future oral amino acid beverages to optimize amino

acid delivery to tendon tissues. Additionally, whether peritendinous amino acid content is

influenced by aging has not been determined. It is well established that oral amino acid

provisions increase serum and skeletal muscle amino acid concentrations (17, 22), but such

work in human tendon *in vivo* has not been completed.

In this investigation, we utilized microdialysis to determine the impact of oral amino acid

consumption on Achilles peritendinous amino acid concentrations in young and older adults (1,

16). We also assessed peritendinous concentrations of pro-collagen Iα1 as a marker of local

collagen synthesis in response to the amino acid provision (29). Based on the work described

above and work in skeletal muscle (10, 12), we utilized an amino acid bolus rich in leucine and

glycine. To our knowledge, no studies have assessed the effects of oral amino acid

supplementation on peritendinous amino acid concentrations in young or older adults *in vivo*. To

set the framework for future clinical investigations and the optimization of amino acid beverages

for therapeutic purposes, establishing the extent to which peritendinous amino acid

concentrations change after oral consumption is critical.

METHODS

Subjects. Seven young (4 men, 3 women; 27±1 yr.; BMI: 24±1) and six elderly (1 man, 5

women; 68±2 yrs.; BMI: 24±2) adults participated in this research study. Exclusion criteria

included body mass index (BMI) greater than 35 kg/m^2, chronic use of medications known to

affect protein or collagen metabolism, a previous history of tendinopathies or diabetes, and

sensitivity to study beverage ingredients. All subjects were sedentary (one day or less per week

of aerobic or resistance exercise for at least one year) or recreationally active (not training for

competitive events). All participants provided voluntary informed written consent. The

Institutional Review Board of Purdue University, West Lafayette, IN (IRB#1904022075),

approved this research study. This study was registered at ClinicalTrials.gov (NCT04064528).

Microdialysis. To assess amino acid concentrations in the peritendinous space of the Achilles

tendon, we utilized microdialysis (1, 16, 18). For each microdialysis experiment, subjects fasted

for 12-hours before arrival at Purdue University. After preparation of the skin with an antiseptic

(povidone-iodine) and local anesthetic (lidocaine 1%), an ethylene oxide sterilized microdialysis

fiber was inserted in the peritendinous space anterior to the Achilles tendon (1, 16). One hour

after fiber insertion, subjects consumed a mixed amino acid beverage prepared from individual

amino acid stocks (Ajinomoto Health & Nutrition North America, Inc, Raleigh, NC). Microdialysis

samples were collected every 30-minutes for four hours after amino acid consumption.

Amino Acid Supplement. Subjects received an oral bolus of essential amino acids containing

3.5 grams of leucine (11, 12, 14), 3 grams of proline, 2 g glycine, 1.1 g histidine, 1.0 g

isoleucine, 1.55 g lysine, 0.30 g methionine, 1.55 g phenylalanine, 1.45 g threonine, and 1.2 g

valine (11). Amino acids (Ajinomoto Health & Nutrition North America, Inc) were mixed in a

noncaloric, non-caffeinated carbonated beverage (Crystal Light, Kraft Foods, Inc). The leucine

dose was chosen based on work demonstrating its effectiveness at stimulating skeletal muscle

protein synthesis (10) and for comparison to previous studies (10, 12). We decided to include

greater glycine because of the recent preclinical evidence implying that glycine can improve

tendon properties (32). The proline and glycine were meant to provide additional material for

collagen synthesis without providing unhealthy levels of these amino acids.

Sample Analysis. Amino acid concentrations were determined with high-performance liquid

chromatography (Agilent Technologies 1100 HPLC System, Santa Clara, CA). Microdialysis

samples (15 μl) were deproteinized with 15 μl of 10%TCA and further diluted with 30 μl 0.1N

HCl. Diluted samples were immediately centrifuged for 10 minutes at 4°C (10,000 g). The

supernatant was removed and transferred to an HPLC vial. Amino acids were eluted using

gradient elution with mobile phase A (10 mM Na_2HPO_4, 10 mM $Na_2B_4O_7$, pH 8.2, and 5 mM

NaN_2, pH 8.2) and mobile phase B (45:45:10 of HPLC-grade acetonitrile, methanol, and water

(21). Separation of amino acids was achieved using an Eclipse Plus C18 4.6x100 mm, 3.5μm

column (Agilent) with a Restek Ultra C18 Guard Column (Restek Corporation, Bellefonte, PA).

Peaks were monitored at 230 nm excitation/450 nm emission (G1321A, Agilent). The

concentration of individual amino acid concentrations was determined by comparison with a

standard curve (AAS18, MilliporeSigma, St. Louis, MO).

The concentration of pro-collagen 1α1 concentration in the peritendinous space was

determined at select time points after amino acid consumption using a DuoSet® ELISA from

R&D Systems (DY6220-05, Minneapolis, MN). Due to the large dialysate volume needed for the

pro-collagen assay, we could not include every sampling point. Microdialysis samples were

diluted 1:9.5 with Reagent Dilutant and assayed in duplicate per the manufacture instructions.

Statistics

Several participants required a restroom break during the microdialysis experiment resulting

in a small number of missed collection points after amino acid consumption. Thus, individual

amino acid concentrations were evaluated with a mixed-effects model. For amino acids and pro-

collagen, noted residuals were not normally distributed and assumptions of excepted non-

constant variance were not correct, thus raw data was log-transformed before analysis. Multiple

comparisons were performed using a Fisher's LSD test. The area under the curve values were

compared using an unpaired, two-tailed t-test. All statistical analysis was completed in

GraphPad Prism 9.0.1.

RESULTS
168
Mean values for all measured amino acids are provided in Table 1. When comparing

individual amino acids, peritendinous glycine concentrations (p=0.015, Figure 1) were greater in

older adults than young adults after beverage consumption through 150 minutes post-beverage

consumption. The age-related difference in glycine concentrations was no longer significant

beginning at 180 minutes post-beverage consumption (Figure 1). In contract, glycine did not

significantly change with time (p=0.087, Figure 1). Threonine, valine, isoleucine, lysine, leucine,

histidine, and phenylalanine concentrations increased with time (p≤0.05, Figure 1) with all of

these amino acids returning to baseline by 180 minutes post-beverage consumption. For

threonine, valine, isoleucine, lysine, leucine, histidine, and phenylalanine, no differences

between young and older adults were detected (p>0.05, Figure 1). The dose of methionine in

the study bolus was not sufficient to increase peritendinous concentrations (p=0.136, Figure 1)

and no significant difference was noted between young and older adults (p=0.077, Figure 1). No

differences across time were noted for the amino acids that were not included in the essential

amino acid bolus (p>0.05, Table 1). Additionally, except for arginine (p=0.05), no differences

between young and older adults were noted in the peritendinous concentrations of amino acids

not included in the essential amino acid bolus.

In contrast to the mixed model analysis, AUC values for glycine, histidine, methionine, and

phenylalanine were greater in older adults compared to young (Table 2). AUC for lysine and

threonine were also higher in older adults but did not reach statistical significance (p=0.06 and

p=0.07, respectively). Peritendinous pro-collagen Iα1 concentrations tended to change with time

(p=0.071) but were not altered by age (mixed-effect analysis, p=0.226 and AUC, p=0.109,

Figure 2).

DISCUSSION

It is well-established that an elevation of serum amino acids can have a significant anabolic effect on skeletal muscle (6, 11, 31). A limited number of investigations in humans and rodents have indicated that tendon tissue is also sensitive to elevations in amino acids. Amino acid or protein beverages have increased tendon CSA in adults completing resistance training (13) and improved outcomes during tendinopathy rehabilitation (15, 30). In rodents, amino acids solutions, especially those rich in leucine or glycine, improve tendon properties in various disease models (28, 32). Consumption of amino acid or protein beverages increases serum and skeletal muscle amino acids. However, to our knowledge, the extent to which oral amino acid ingestion alters local (peritendinous) concentrations of amino acids in humans has not been described. The potential impact of amino acids on tendon properties in older adults is also relevant, given the changes in tendon properties with aging (5). Thus, we also considered the impact of aging on the ability of oral amino acids to increase peritendinous amino acid concentrations. Additionally, we determined if a provision of amino acids would increase the peritendinous concentration of a common collagen synthesis marker.

In this study, we demonstrate that amino acids can be detected in the Achilles peritendinous space of humans. The concentrations of the amino acids not included with the beverage were stable across time, suggesting a minimal impact of fiber insertion (Table 1). Of the amino acids in the drink, all increased with time (p<0.05), except methionine and glycine. The change in Achilles peritendinous amino acid concentrations across time followed a similar pattern to those previously reported in serum and skeletal muscle (1, 10) with levels peaking at 1-2 hours post-ingestion then returning to baseline levels at approximately three hours (Figure 2).

While direct statistical comparisons are not possible, the observed basal peritendinous amino acid concentrations (Table 1) were substantially lower than previously reported in serum and skeletal muscle (7, 10). Additionally, the peak concentrations obtained in the peritendinous space after consuming the amino acid bolus were generally lower than skeletal muscle or serum

(10). For example, utilizing a similar amino acid bolus, Dickinson et al. (10) reported peak serum

concentrations of approximately 1000 μM and 250 μM for leucine and phenylalanine,

respectively. In comparison, the highest concentration obtained in the Achilles peritendinous

space for leucine was 115 μM for young and 143 μM for older adults. Phenylalanine Achilles

peritendinous concentrations peaked at 20 μM for young and 38 μM for older adults. We did not

account for probe recovery due to the large number of analytes examined. However, we have

previously reported a probe recovery range of 50-65% for the amino acid sarcosine (16). Even

accounting for estimation of recovery, the values obtained in the Achilles peritendinous space

are lower than those obtained in serum or skeletal muscle. Further, the increase in

peritendinous amino acid concentrations after the oral bolus was modest when compared to

serum or skeletal muscle. Even with the larger dose of glycine, we did not observe a significant

increase in peritendinous glycine concentrations with time. This suggests that delivery of amino

acids to the Achilles tendon is limited when compared to skeletal muscle and serum. Larger oral

doses of amino acids may be needed to achieve greater benefits for tendons.

A lower concentration of amino acids after oral consumption is consistent with our previous

work with acetaminophen (16). Achilles peritendinous levels of acetaminophen achieved

maximum values that were approximately 50% lower than those seen in serum or skeletal

muscle after oral consumption of the drug (24). It is not yet clear why the peritendinous

concentration of compounds is lower than serum or skeletal muscle compared to the Achilles

peritendinous space. While the tendon is poorly vascularized, blood flow in the peritendinous

space during non-exercise conditions is similar to the surrounding calf muscle (4) implying that

delivery of amino acids to the tendon would not be impaired.

Interestingly, when accounting for the area under the curve, the concentration of several of

the amino acid included in the beverage achieved higher peritendinous concentrations in older

adults than in young (Figure 1). Arginine, which was not included in the amino acid bolus, was

also higher in the older participants compared to young. A greater baseline peritendinous amino

acid concentration is surprising as serum amino acid concentrations are typically lower in older

adults than young (26).

With the apparent anabolic effect of leucine and glycine on tendon collagen (32) and mass

(13), we evaluated peritendinous pro-collagen Iα1 as a marker of collagen synthesis. Even with

the large proportion of leucine and glycine in the amino acid beverage, we did not observe a

significant increase in peritendinous levels of pro-collagen Iα1 (p=0.071). One limitation of the

peritendinous microdialysis approach is that we assume that any pro-collagen detected is a

reflection of intratendinous events. Extensive work by Langberg and colleagues (19) has

validated the peritendinous microdialysis technique providing evidence that peritendinous

measures correlate with intratendinous measures. However, using stable isotope methods to

directly assess collagen synthesis (23) would provide more detailed results but would require

invasive tissue sampling.

In future work, it would be interesting to determine if a different source of protein (e.g., whey

or soy) or variations in beverage amino acid content would alter peritendinous amino acid

concentrations, as reported for serum (31). The limited rodent and human work suggest that

tendon tissue is indeed sensitive to serum amino acids changes. However, given the lower

concentrations obtained in the peritendinous space, we should carefully explore the dose-

response impact of amino acids on tendon tissue using *in vitro* or *ex vivo* models. Such

information is critical for optimizing oral doses of amino acids for human consumption to

maximize the stimulation of collagen synthesis. Also, understanding the anabolic potential of

each amino acid would aid in optimizing beverage content. It would be exciting to determine if a

larger bolus or repeated smaller doses of amino acids would increase peritendinous amino acid

concentrations to a greater extent than seen in the current investigation. Understanding the

impact of these beverages on peritendinous amino acid concentrations will optimize beverage

content to maximize the clinical benefit of such compounds.

While much work has attempted to define an optimal protein dose to stimulate skeletal

muscle protein synthesis, such information is not yet available for the tendon. Developing a

nutritional cocktail to optimize tendon and skeletal muscle health is appropriate given the

importance of both tissues to overall musculoskeletal function. The small number of human

studies suggest the exciting possibility that amino acid beverages could be optimized to

increase peritendinous amino acids level for optimization of tendon health and recovery while

providing the established benefits for skeletal muscle.

<u>Acknowledgments</u>

We thank the subjects for their participation and commitment to the investigation. This project

was funded by Purdue University Research Initiative Funds to CCC. The study was designed by

CCC; data were collected and analyzed by all authors; data interpretation and manuscript

preparation were undertaken by CCC and AS. All authors approved the final version of the

paper.

Figure Legend

Figure 1: Line graphs of Achilles peritendinous amino acid concentrations over time. Data

expressed as mean±standard error. No interactions were noted. Main effect p values for time

and age are presented above each figure insert. Multiple comparisons were performed on

significant main effects. *p<0.05 versus Young vs. Old; [#]p<0.05 versus 30 minutes. [$]p=0.06

versus 30 minutes.

Figure 2: Line graph of Achilles peritendinous pro-collagen concentrations over time. Data

expressed as mean±standard error.

300
301

REFERENCES

1.   **Astill BD, Katsma MS, Cauthon DJ, Greenlee J, Murphy M, Curtis D, and Carroll CC**. Sex-based difference in Achilles peritendinous levels of matrix metalloproteinases and growth factors after acute resistance exercise. *J Appl Physiol (1985)* 122: 361-367, 2017.

2.   **Birch HL, Peffers MJ, and Clegg PD**. Influence of Ageing on Tendon Homeostasis. *Adv Exp Med Biol* 920: 247-260, 2016.

3.   **Boivin GP, Platt KM, Corbett J, Reeves J, Hardy AL, Elenes EY, Charnigo RJ, Hunter SA, and Pearson KJ**. The effects of high-fat diet, branched-chain amino acids and exercise on female C57BL/6 mouse Achilles tendon biomechanical properties. *Bone Joint Res* 2: 186-192, 2013.

4.   **Boushel R, Langberg H, Olesen J, Nowak M, Simonsen L, Bulow J, and Kjaer M**. Regional blood flow during exercise in humans measured by near-infrared spectroscopy and indocyanine green. *J Appl Physiol (1985)* 89: 1868-1878, 2000.

5.   **Carroll CC, Dickinson JM, Haus JM, Lee GA, Hollon CJ, Aagaard P, Magnusson SP, and Trappe TA**. Influence of aging on the in vivo properties of human patellar tendon. *J Appl Physiol (1985)* 105: 1907-1915, 2008.

6.   **Carroll CC, Fluckey JD, Williams RH, Sullivan DH, and Trappe TA**. Human soleus and vastus lateralis muscle protein metabolism with an amino acid infusion. *Am J Physiol Endocrinol Metab* 288: E479-485, 2005.

7.   **Corso G, Cristofano A, Sapere N, la Marca G, Angiolillo A, Vitale M, Fratangelo R, Lombardi T, Porcile C, Intrieri M, and Di Costanzo A**. Serum Amino Acid Profiles in Normal Subjects and in Patients with or at Risk of Alzheimer Dementia. *Dement Geriatr Cogn Dis Extra* 7: 143-159, 2017.

8.   **Couppe C, Hansen P, Kongsgaard M, Kovanen V, Suetta C, Aagaard P, Kjaer M, and Magnusson SP**. Mechanical properties and collagen cross-linking of the patellar tendon in old and young men. *J Appl Physiol (1985)* 107: 880-886, 2009.

9.   **Csapo R, Malis V, Hodgson J, and Sinha S**. Age-related greater Achilles tendon compliance is not associated with larger plantar flexor muscle fascicle strains in senior women. *J Appl Physiol (1985)* 116: 961-969, 2014.

10.   **Dickinson JM, Gundermann DM, Walker DK, Reidy PT, Borack MS, Drummond MJ, Arora M, Volpi E, and Rasmussen BB**. Leucine-Enriched Amino Acid Ingestion after Resistance Exercise Prolongs Myofibrillar Protein Synthesis and Amino Acid Transporter Expression in Older Men. *J Nutr* 144: 1694-1702, 2014.

11.   **Dickinson JM, Gundermann DM, Walker DK, Reidy PT, Borack MS, Drummond MJ, Arora M, Volpi E, and Rasmussen BB**. Leucine-enriched amino acid ingestion after resistance exercise prolongs myofibrillar protein synthesis and amino acid transporter expression in older men. *J Nutr* 144: 1694-1702, 2014.

12.   **Dickinson JM, Reidy PT, Gundermann DM, Borack MS, Walker DK, D'Lugos AC, Volpi E, and Rasmussen BB**. The impact of postexercise essential amino acid ingestion on the ubiquitin proteasome and autophagosomal-lysosomal systems in skeletal muscle of older men. *J Appl Physiol (1985)* 122: 620-630, 2017.

13.   **Farup J, Rahbek SK, Vendelbo MH, Matzon A, Hindhede J, Bejder A, Ringgard S, and Vissing K**. Whey protein hydrolysate augments tendon and muscle hypertrophy independent of resistance exercise contraction mode. *Scand J Med Sci Spor* 24: 788-798, 2014.

14.   **Glynn EL, Fry CS, Drummond MJ, Timmerman KL, Dhanani S, Volpi E, and Rasmussen BB**. Excess leucine intake enhances muscle anabolic signaling but not net protein anabolism in young men and women. *J Nutr* 140: 1970-1976, 2010.

15.   **Gumina S, Passaretti D, Gurzi MD, and Candela V**. Arginine L-alpha-ketoglutarate, methylsulfonylmethane, hydrolyzed type I collagen and bromelain in rotator cuff tear repair: a prospective randomized study. *Curr Med Res Opin* 28: 1767-1774, 2012.

16.   **Gump BS, McMullan DR, Cauthon DJ, Whitt JA, Del Mundo JD, Letham T, Kim PJ,**
Friedlander GN, Pingel J, Langberg H, and Carroll CC. Short-term acetaminophen
consumption enhances the exercise-induced increase in Achilles peritendinous IL-6 in humans.
J Appl Physiol (1985) 115: 929-936, 2013.
17.   **Gutierrez A, Anderstam B, and Alvestrand A**. Amino acid concentration in the
interstitium of human skeletal muscle: a microdialysis study. *Eur J Clin Invest* 29: 947-952,
1999.
18.   **Langberg H, Boushel R, Skovgaard D, Risum N, and Kjaer M**. Cyclo-oxygenase-2
mediated prostaglandin release regulates blood flow in connective tissue during mechanical
loading in humans. *J Physiol* 551: 683-689, 2003.
19.   **Langberg H, Olesen JL, Bulow J, and Kjaer M**. Intra- and peri-tendinous microdialysis
determination of glucose and lactate in pigs. *Acta Physiol Scand* 174: 377-380, 2002.
20.   **Lewis JS**. Rotator cuff tendinopathy. *Br J Sports Med* 43: 236-241, 2009.
21.   **Long W**. Automated Amino Acid Analysis
Using an Agilent Poroshell HPH-C18
Column. Agilent Technologies, 2017.
22.   **McCormack WG, Cooke JP, O'Connor WT, and Jakeman PM**. Dynamic measures of
skeletal muscle dialysate and plasma amino acid concentration in response to exercise and
nutrient ingestion in healthy adult males. *Amino Acids* 49: 151-159, 2017.
23.   **Miller BF, Olesen JL, Hansen M, Dossing S, Crameri RM, Welling RJ, Langberg H,**
Flyvbjerg A, Kjaer M, Babraj JA, Smith K, and Rennie MJ. Coordinated collagen and muscle
protein synthesis in human patella tendon and quadriceps muscle after exercise. *J Physiol-*
London 567: 1021-1033, 2005.
24.   **Muller M, Schmid R, Georgopoulos A, Buxbaum A, Wasicek C, and Eichler HG**.
Application of microdialysis to clinical pharmacokinetics in humans. *Clin Pharmacol Ther* 57:
371-380, 1995.
25.   **Notarnicola A, Pesce V, Vicenti G, Tafuri S, Forcignano M, and Moretti B**. SWAAT
Study: Extracorporeal Shock Wave Therapy and Arginine Supplementation and Other
Nutraceuticals for Insertional Achilles Tendinopathy (vol 29, pg 799, 2012). *Adv Ther* 29: 992-
992, 2012.
26.   **Pitkanen HT, Oja SS, Kemppainen K, Seppa JM, and Mero AA**. Serum amino acid
concentrations in aging men and women. *Amino Acids* 24: 413-421, 2003.
27.   **Praet SFE, Purdam CR, Welvaert M, Vlahovich N, Lovell G, Burke LM, Gaida JE,**
Manzanero S, Hughes D, and Waddington G. Oral Supplementation of Specific Collagen
Peptides Combined with Calf-Strengthening Exercises Enhances Function and Reduces Pain in
Achilles Tendinopathy Patients. *Nutrients* 11: 2019.
28.   **Scott A, and Nordin C**. Do Dietary Factors Influence Tendon Metabolism? *Adv Exp*
Med Biol 920: 283-289, 2016.
29.   **Seo WY, Kim JH, Baek DS, Kim SJ, Kang S, Yang WS, Song JA, Lee MS, Kim S,**
and Kim YS. Production of recombinant human procollagen type I C-terminal propeptide and
establishment of a sandwich ELISA for quantification. *Sci Rep* 7: 15946, 2017.
30.   **Shaw G, Lee-Barthel A, Ross MLR, Wang B, and Baar K**. Vitamin C-enriched gelatin
supplementation before intermittent activity augments collagen synthesis. *Am J Clin Nutr* 105:
136-143, 2017.
31.   **Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, and Phillips SM**. Ingestion of
whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at
rest and following resistance exercise in young men. *J Appl Physiol (1985)* 107: 987-992, 2009.
32.   **Vieira CP, De Oliveira LP, Da Re Guerra F, Dos Santos De Almeida M, Marcondes**
MC, and Pimentel ER. Glycine improves biochemical and biomechanical properties following
inflammation of the achilles tendon. *Anat Rec (Hoboken)* 298: 538-545, 2015.
402

	30		60		90		120		150		180		210		240	
	Y	O	Y	O	Y	O	Y	O	Y	O	Y	O	Y	O	Y	O
Glu	12±2	31±15	11±2	31±11	9±2	22±5	11±1	21±3	8±2	27±9	8±2	17±6	10±2	16±5	12±4	15±4
Asn	11±2	16±2	11±2	17±2	9±2	14±2	9±1	15±1.5	8.5±1	16±4	9±2	9±3	10±2	10±2	10±1	10±1
Ser	28±4	47±9	30±5	56±10	27±6	52±10	28±5	54±10	27±4	67±22	29±7	33±8	31±6	34±7	30±5	30±3
Gln	273±47	555±119	292±50	612±110	241±50	468±59	253±44	557±94	245±40	614±215	272±68	302±75	298±62	322±65	279±42	320±36
His$^{\$}$	18±3	36±11	22±3	44±13	21±4	45±9	20±1	38±3	18±3	46±11	18±3	20±6	20±2	20±4	18±2	18±3
Gly	66±9	133±24*	85±11	182±30*	87±16	197±38*	98±18	199±32*	84±13	217±47*	86±22	109±28	87±17	107±18	79±11	102±7
Thr	32±9	42±8	43±10	58±8	42±9.5$^{\#}$	78±17$^{\#}$	41±8$^{\#}$	78±13$^{\#}$	40±8$^{\#}$	85±22$^{\#}$	42±10	41±12	42±8	42±10	89±51	38±7
Arg*	17±3	30±6	16±3	33±7	11±2	27±5	14±3	30±5	11±2	31±9	13±4	15±4	15.5±4	18±4	15±3	17±3
Ala	62±13	117±26	67±14	125±13	51±14	116±7	63±13	125±10	53±11	128±22	57±17	84±23	65±17	84±18	58±10	78±10
Tyr	13±2	19±2	14±2	23±2	12±3	21±2	11±2	22±2	11±2	23±4	11±2	15±4	12±2	15±3	12±2	14±1
Cys	70±10	127±31	92±13	166±42	88±20	159±23	105±20	177±32	82±15	202±63	85±20	116±36	94±19	118±32	86±11	105±24
Val	61±17	65±14	79±19$^{\#}$	113±20$^{\#}$	71±15$^{\#}$	104±14$^{\#}$	65±7	125±27	65±12$^{\#}$	136±49$^{\#}$	64±13	56±15	65±8	56±10	62±8	52±5
Met	10±2	14±2	12±2	19±3	10±2	18±2	10±1	17±2	9±2	18±3	9±2	12±0.6	9±1	12±0.9	10±2	10±0.4
Trp	5±1	7±1	5±1	7±1	6±1	5±2	4±1	5±1	4±1	6±2	4±0.9	3±0.9	6±2	3±0.5	4±0.9	2±0.3
Phe	14±3	18±3	19±3$^{\#}$	30±4$^{\#}$	20±4$^{\#}$	33±4$^{\#}$	18±2$^{\#}$	34±4$^{\#}$	18±3$^{\#}$	38±10$^{\#}$	18±4	19±5	18±3	18±4	18±3	16±2
Iso	16±3	18±4	26±4$^{\#}$	43±8$^{\#}$	24±5$^{\#}$	42±6$^{\#}$	23±3$^{\#}$	46±10$^{\#}$	19±4$^{\#}$	43±16$^{\#}$	35±17	16±5	20±3	15±3	18±4	13±2
Leu	35±6	37±7	115±55$^{\#}$	119±26$^{\#}$	74±15$^{\#}$	134±18$^{\#}$	73±10$^{\#}$	143±29$^{\#}$	63±10$^{\#}$	137±46$^{\#}$	57±12	54±17	55±7$^{\#}$	49±11$^{\#}$	51±7$^{\#}$	44±6$^{\#}$
Lys$^{\$}$	32±6	45±8	41±8	67±15	36±7	59±13	34±6	64±12	33±5	61±14	33±8	31±9	33±6	31±8	30±4	29±4

Table 1: Bolded amino acids were included in the study beverage. Values are expressed in μM with mean±SE. Y = young adults; O = older adults. Numbers in first row represent minutes post-amino acid consumption. *p≤0.05 young versus old, $^{\#}$p≤0.05 different from 30 minutes. $^{\$}$p<0.05, main effect for time.

Table 2

Amino Acid	Young Adults	Older Adults
Glycine	17974±2249	33843±3655*
Histidine	4079±414	7188±970*
Isoleucine	4948±1086	6629±970
Leucine	14412±3158	20294±2889
Lysine	7234±938	10504±1327[#]
Methionine	2069±253	3244±239*
Phenylalanine	3821±450	5687±603*
Threonine	8574±1267	12645±1587[&]
Valine	14092±1831	19471±2826

Date presented as mean±standard error. *p<0.05, [#]p=0.0642, [&]p=0.0673

Time: p=0.087, Age: p=0.015

Time: p=0.05, Age: p=0.197

Time: p=0.042, Age: p=0.268

Time: p=0.004, Age: p=0.349

Time: p=0.136, Age: p=0.077

Time: p=0.046, Age: p=0.198

Time: p=0.0002, Age: p=0.213

Time: p=0.047, Age: p=0.089

Time: p=0.027, Age: p=0.167