Salmonella Typhimurium outer membrane protein A (OmpA) renders protection against nitrosative stress by promoting SCV stability in murine macrophages

Atish Roy Chowdhury[1, 2], Shivjee Sah[1, 2], Umesh Varshney[1, 2] and Dipshikha Chakravortty[1, 2, 3*]

[1]Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, Karnataka, India-560012

[2]Division of Biological Sciences, Indian Institute of Science, Bangalore, Karnataka, India-560012

[3]Centre for Biosystems Science and Engineering, Indian Institute of Science, Bangalore, Karnataka, India-560012.

Current address: Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore, Karnataka, India-560012

* *Corresponding author*

Dipshikha Chakravortty

Email: dipa@ iisc.ac.in

Tel: 0091 80 2293 2842

Fax: 0091 80 2360 2697

Abstract

Porins are highly conserved bacterial outer membrane proteins involved in the selective transport of charged molecules across the membrane. Despite their significant contributions to the pathogenesis of Gram-negative bacteria, their precise role in salmonellosis remains elusive. In this study, we investigated the role of porins (OmpA, OmpC, OmpD, and OmpF) in *Salmonella* Typhimurium (STM) pathogenesis. OmpA played a multifaceted role in STM pathogenesis, and a strain deleted for *ompA* (STM *ΔompA*) showed enhanced proneness to phagocytosis and compromised proliferation in macrophages. However, in the epithelial cells, despite being invasion deficient, it was hyper-proliferative. The poor colocalization of STM *ΔompA* with LAMP-1 confirmed impaired stability of SCV membrane around the intracellular bacteria, resulting in its (STM *ΔompA*) release into the cytosol of macrophages where it is assaulted with reactive nitrogen intermediates (RNI). The cytosolic localization of STM *ΔompA* was responsible for the downregulation of SPI-2 encoded virulence factor SpiC, which is required to suppress the activity of iNOS. The reduced recruitment of nitrotyrosine on STM in the macrophage cytosol upon ectopically expressing Listeriolysin O (LLO) explicitly supported the pro-bacterial role of OmpA against the host nitrosative stress. Further, we show that the generation of time-dependent redox burst could be responsible for the enhanced sensitivity of STM *ΔompA* towards nitrosative stress. The absence of OmpA in STM *ΔompA* resulted in the loss of integrity and enhanced porosity of the bacterial outer membrane, which was attributed to the upregulated expression of *ompC*, *ompD,* and *ompF*. We showed the involvement of OmpF in the entry of excess nitrite in STM *ΔompA,* thus increasing the susceptibility of the bacteria towards *in vitro* and *in vivo* nitrosative stress. In conclusion, we illustrated a mechanism of strategic utilization of OmpA compared to other porins by wildtype *Salmonella* for combating the nitrosative stress in macrophages.

Keywords: *Salmonella* Typhimurium, Outer membrane protein A, *Salmonella* containing vacuole (SCV), LAMP-1, Nitrosative stress, Reactive Nitrogen Intermediates (RNI), Nitrotyrosine, Outer membrane protein F.

Introduction

The pathogenicity of many Gram-negative bacteria of the family *Enterobacteriaceae* is regulated by porins, commonly known as outer membrane proteins (or Omp). Porins are outer membrane-bound β barrel proteins with 8 to 24 antiparallel β strands connected by loops and are well known for their role in the selective diffusion of ions and solutes across the outer membrane of bacteria [1]. Despite their roles in maintaining outer membrane stability, their pathogenic functions have also been well documented. OmpA, one of the most abundant porins of the bacterial outer membrane, is extensively utilized by *Klebsiella pneumoniae* to prevent the IL-8 dependent pro-inflammatory response in the airway epithelial A549 cells [2]. The deletion of *ompC* and *ompF* from pathogenic *E. coli* not only impaired its invasion in bEnd.3 cells but also reduced its virulence in a mouse model [3]. OprF, an OmpA ortholog in the outer membrane of *Pseudomonas sp.,* has been reported to function as a sensor of quorum signaling and to induce virulence [4]. *E. coli* OmpW has been reported to play a significant role against phagocytosis and complement activation [5, 6].

The outer membrane of *Salmonella* Typhimurium, a member of the *Enterobacteriaceae* family that causes typhoid fever-like symptoms in mice and self-limiting gastroenteritis in humans, is densely populated with many porins such as OmpA, OmpC, OmpD, and OmpF. Unlike OmpA, which has a significant role in the tight attachment of the outer membrane to the underlying peptidoglycan layer with its periplasmic tail [7], the other porins facilitate transportation of charged ions [8]. The connection between the outer membrane porins of *Salmonella* Typhimurium and its pathogenesis remains elusive due to the lack of detailed studies. Earlier, Heijden *et al*. proposed a mechanism of OmpA and OmpC dependent regulation of outer membrane permeability in *Salmonella* in response to H_2O_2 and antibiotic stresses [9].

In the current study, we have investigated the individual roles of OmpA, OmpC, OmpD, and

OmpF in the pathogenesis of *Salmonella* Typhimurium with a profound focus on OmpA. We

found a strong dependence of wild-type *Salmonella* on OmpA for its survival in macrophages

and the mouse model. Our study illustrates a mechanism of strategic utilization of OmpA by

intracellular *Salmonella* in combatting the nitrosative stress of macrophages by enhancing the

outer membrane stability of the bacteria. To the best of our knowledge, this is the first study to

demonstrate the impact of outer membrane porins in maintaining the stability of *Salmonella*-

containing vacuole (SCV) in macrophages and epithelial cells.

Results

OmpA promotes the evasion of phagocytosis and intracellular survival of *Salmonella* in

macrophages.

The transcriptomic analyses of intracellular *Salmonella* Typhimurium infecting J774-A.1 and

HeLa cells have demonstrated almost 2 to 2.5 fold induction in the expression level of *ompA*

during early (4h), middle (8h), and late (12h) stages of infection in comparison with other major

membrane porins namely *ompC*, *ompD*, and *ompF* [10, 11]. To validate this observation, we

isolated total RNA from *Salmonella* Typhimurium strain 14028S grown in nutritionally

enriched Luria-Bertani broth, low magnesium minimal acidic F media (pH= 5.4) mimicking

the nutrient-deprived acidic environment of *Salmonella* containing vacuole (SCV) [12], and

RAW264.7 cells at different time points (3, 6, 9, 12 h post-inoculation/ infection) and

investigated the transcript levels of *ompA* (**Figure S1A**), *ompC* (**Figure S1B**), *ompD* (**Figure**

S1C), and *ompF* (**Figure S1D**) by real-time PCR. Unlike the steady decrease in the expression

levels of *ompC*, *ompD,* and *ompF*, whose expression steadily decreased in wild type bacteria

growing in LB broth, F media, and macrophages at all the time points except for *ompF* in

macrophages at 9^{th} h post-infection, there was a significant increase in the transcript level of

ompA during the late phase of infection (9h and 12 h post-infection) in macrophages (**Figure S1A, S1D**). These observations suggest a preference of *ompA* over the other membrane-bound larger porins to enable *Salmonella* to thrive inside macrophages. The observation is also consistent with the microarray data published by Eriksson *et al*. and Hautefort *et al.* [10, 11].

To investigate the importance of *ompA* in the intracellular survival of *Salmonella*, we knocked out *ompA* from the genome of *Salmonella* Typhimurium using a protocol demonstrated earlier by Datsenko and Wanner (Data not shown) [13]. The complementation of *ompA* (1.1 kb) in the knockout bacteria was done using the pQE60 plasmid. A ~1.42-fold upregulation of *ompA* was found by real-time PCR analysis compared to the wild type *S.* Typhimurium (Data not shown). Unpublished data from our lab suggested that knocking out *ompA* from *Salmonella* did not alter the *in vitro* growth of the bacteria in LB broth. To determine the role of *S.* Typhimurium *ompA* in phagocytosis, we checked phagocytosis of the wild-type and knockout strains in macrophages (**Figure 1A**). We found that the OmpA of STM (WT) helps evade phagocytosis by RAW264.7 and activated U937 cells (**Figure 1A**). The percent phagocytosis of STM *ΔompA* in RAW 264.7 and U937 cells is more than STM (WT).

On the contrary, the complemented strain was less prone to phagocytosis by both the macrophages. To mimic the physiological condition, RAW264.7 cells were further infected with STM (WT), and *ΔompA* coated with 10% mouse complement sera (**Figure 1B**). A marked increase in the phagocytosis of complement coated STM *ΔompA* in comparison with complement uncoated STM *ΔompA* and coated STM (WT) confirms the role of *Salmonella* OmpA against complement recruitment and phagocytosis by macrophages (**Figure 1B**). Adhesion of bacteria onto the host cell surface occurs before it enters the host cell, either by phagocytosis or by invasion [14]. Since we found enhanced phagocytosis of STM *ΔompA* in macrophages compared to wild-type bacteria, we decided to evaluate the bacterial attachment on the macrophage surface by *in vitro* adhesion assay (**Figure 1C**). The adhesion of STM

ΔompA on phagocytic RAW 264.7 cells is more than STM (WT) (**Figure 1C**). The enhanced

adhesion of STM *ΔompA* was abrogated upon complementation with *ompA*. We then performed

the intracellular survival assay of STM (WT) and STM *ΔompA* strains in RAW264.7, activated

U937 cells, respectively (**Figure 1D**). We observed that the intracellular growth of STM

ΔompA (fold proliferation in RAW264.7 cells- 11.45± 3.349, U937 cells- 5.075± 1.157) is

significantly attenuated in RAW 264.7 (by 2.75 folds) and activated U937 cells (by 3.08 folds)

compared to the wild type parent (fold proliferation in RAW264.7 cells- 31.5± 5.347, U937

cells- 15.65± 3.981) (**Figure 1D**). When the cell lines were infected with the complemented

strain, there was a recovery of the intracellular proliferation of bacteria (**Figure 1D**). Hence,

we conclude that *Salmonella* utilizes OmpA as a double-edged sword to protect the bacteria

from phagocytosis and then helps it to survive within macrophages.

Once ingested, *Salmonella* starts invading M cells of the Peyer's patches in the small intestine

with the help of the SPI-1 encoded type 3 secretion system. Hence, we checked the role of

Salmonella OmpA in bacterial invasion of non-phagocytic epithelial cells (**Figure 2A**).

Compared to the wild-type bacteria, STM *ΔompA* exhibited significant attenuation in the

invasion of the human colorectal adenocarcinoma cell line- Caco-2 and human cervical cancer

cell line- HeLa (**Figure 2A**), and the invasiveness was rescued upon complementation (**Figure

2A**). To further validate this observation, we carried out an *in vitro* adhesion assay using HeLa

cells as hosts (**Figure 2B**). STM *ΔompA* showed reduced attachment on the surface of HeLa

cells compared to the wild-type bacteria (**Figure 2B**). The inefficiency of the knockout strain

to attach to the epithelial cell surface was rescued in the complemented strain (**Figure 2B**).

This observation is consistent with the result obtained from the invasion assay and showed

utilization of OmpA by *Salmonella* as an important adhesion and invasion tool. We further

verified the role of OmpA in maintaining the intracellular life of bacteria in epithelial cells

(**Figure 2C**). Surprisingly, compared to wild-type bacteria, STM *ΔompA* exhibited

hyperproliferation in Caco-2 and HeLa cells (**Figure 2C**). The intracellular proliferation of

complement strain was comparable to the wild-type bacteria in both the cell lines. Since we

found opposite outcomes in the intracellular survival of the *ompA* knockout strain of

Salmonella in two different cell types (attenuation in macrophages and hyper-proliferation in

the epithelial cells), we decided to monitor the intracellular niche of the bacteria in both cell

types. After entering the host cells, *Salmonella* resides inside a modified phagosomal

compartment called SCV, which is acidic. The intracellular life and proliferation of *Salmonella*

depend upon the stability and integrity of SCV. *Salmonella* recruits a plethora of host proteins

to maintain the sustainability of SCV. During the early stage of infection in macrophages and

epithelial cells, SCV is characterized by the presence of the markers of early endocytic pathway

such as EEA1, Rab5, Rab4, Rab11 and transferrin receptors, etc. These proteins are replaced

by late endosome markers such as LAMP-1, Rab7, vATPase, etc., within 15 to 45 min post-

infection [15, 16]. Since we found an attenuation in the intracellular proliferation of STM

ΔompA in macrophages and hyperproliferation in epithelial cells, we decided to check the

intracellular niche of the bacteria using LAMP-1 as a marker of SCV in RAW264.7 cells

(**Figure 1E**) and Caco-2 cells (**Figure 2D**) 16 h post-infection. It was observed that the percent

colocalization of LAMP-1 with STM *ΔompA* is less compared to STM (WT) in RAW264.7

cells (**Figure 1E**). The colocalization with LAMP-1 increased when the macrophage cells were

infected with complement strain. These observations corroborated not only the gradual loss of

the SCV membrane from its surrounding but also its enhanced cytosolic localization in

macrophages (**Figure 1E**). The PFA-fixed dead bacteria lacking the ability to quit vacuole was

used as a positive control in this study (**Figure 1E**). The environment within the SCV is acidic

(pH= 5.4) in comparison with the cytosol (pH= 7.4) of the macrophages [17], [18]. This acidic

environment of the SCV is sensed by the wild-type *Salmonella* Typhimurium with the help of

envZ/ ompR and *phoP/ phoQ* two-component systems, which activates the expression of SPI-

2 genes [17, 19, 20]. The SPI-2 codes for a multiprotein needle-like complex called type III

secretion system (T3SS) and several effector proteins (called translocon), which are primarily

accumulated on the surface of the matured SCV and secreted into the cytosol of the host cells

during the late stage of infection to enhance the severity of the infection (**Figure S2A**). The

assembly of the SPI-2 encoded effector proteins, namely SseB, SseC, SseD, on the surface of

the SCV facilitates the formation of a functionally active T3SS needle complex, which in turn

helps in the systemic colonization of the bacteria (**Figure S2A)** [21, 22]. Since the cytosolic

population of STM *ΔompA* lacks the vacuolar membrane, we anticipated an interruption in the

accumulation of these virulent proteins on the bacterial surface and their secretion into the host

cytosol. In line with our previous observations, we found a marked reduction in the

accumulation and secretion (the area of the infected macrophage demarcated with a dotted line

for the wild-type *Salmonella*) of SseC (**Figure S2B and S2C**) and SseD (**Figure S2B and**

S2D) on or from the surface of STM *ΔompA* into the host cytosol. In continuation of this

observation, we found a dampened expression of *sseC* (**Figure S2E**) and *sseD* (**Figure S2F**)

genes in intracellularly growing STM *ΔompA* compared to the wild-type bacteria. We further

wanted to test the impact of the cytosolic localization of STM *ΔompA* on the expression profile

of several other important SPI-2 virulent genes such as *ssaV* (**Figure S2G**) and *sifA* (**Figure**

S2H). The attenuated expression of *ssaV* and *sifA* suggested that unlike the acidic pH of SCV,

the cytosolic pH of the macrophage does not favor the expression of SPI-2 virulent genes,

which could be a reason behind the compromised growth of *ompA* deficient bacteria in

macrophage. To verify whether this is a cell type-specific phenomenon or not, we further

investigated the intracellular niche of wild type, knockout, and complement strains of

Salmonella in epithelial Caco-2 cells. The quantitative percent colocalization between LAMP-

1 and all three bacterial strains (**Figure 2D**) demonstrated that STM *ΔompA* comes into the

cytoplasm of Caco-2 cells after being released from the SCV during the late phase of infection.

In contrast, wild-type and the complemented strains abstained themselves from doing so
(**Figure 2D**). These findings were further supported by chloroquine resistance assay of STM-
(WT), *ΔompA*, *ΔompA*: pQE60-*ΔompA* in RAW264.7 (**Figure 1F**) and Caco-2 cells (**Figure
2E**). After being protonated inside SCV because of the acidic pH, chloroquine cannot quit the
vacuole and kills the vacuolar population of *Salmonella*, allowing the growth of the cytosolic
population. During the late phase of infection (16 h post-infection) in RAW264.7 cells (**Figure
1F**) and Caco-2 cells (**Figure 2E**), the cytosolic abundance of STM *ΔompA* was found more
comparable to the wild type and complement strain. Taken together, these results authenticate
that, irrespective of the cell type, the intravacuolar life of the intracellular *Salmonella* strongly
depends upon OmpA.

OmpA dependent protection of *Salmonella* against nitrosative stress inside RAW264.7
cells.

The intracellular population of *Salmonella* is heavily encountered by NADPH phagocytic
oxidase-dependent oxidative burst and iNOS dependent nitrosative burst during early and late
stages of infection in macrophages, respectively [23, 24]. The SCV membrane protects the
vacuolar niche of wild-type *Salmonella* from the potential threats present in the cytoplasm of
macrophages in the form of Reactive Oxygen Species (ROS), Reactive Nitrogen Intermediates
(RNI), antimicrobial peptides (AMPs), etc. [25, 26]. Our study confirmed the proneness of
STM *ΔompA* release from SCV into the cytoplasm of macrophages and epithelial cells during
the late stage of infection. Hence, we assumed the increasing possibility of the mutant bacteria
being targeted with ROS and RNI in the cytoplasm of the macrophages, which could be a
probable reason for attenuation of intracellular proliferation. Our hypothesis was verified by
measuring nitrosative (**Figure 3A and 3B**) and oxidative (**Figure S3A – S3C**) stress response
of macrophages infected with wild type, knockout, and complemented strains. We quantified

the extracellular (**Figure 3A**) [NO] response produced by infected macrophages by Griess

assay. It was found that during the late stage of infection (16 h post-infection), the accumulation

of nitrite in the culture supernatant of RAW264.7 cells infected with STM *ΔompA* was

significantly higher compared to the wild type. This heightened [NO] response was revoked

when the RAW264.7 cells were infected with complement strain (**Figure 3A**). This result was

further validated by quantifying intracellular [NO] response using DAF2-DA (**Figure 3B**).

Only 2.22 % of wild-type bacteria-infected macrophages produced [NO] (Figure 3.6B), which

increased to 6.17 % when the cells were infected with STM *ΔompA* (**Figure 3B**). The

percentage of RAW264.7 cells producing [NO] after being infected with complement strains

3.94 % was comparable to the wild-type STM infected cells (Figure 3B). During the late stage

of infection (16 h post-infection) in macrophages, the intracellular ROS and extracellular H_2O_2

levels were estimated using H_2DCFDA (**Figure S3A and S3B**) and phenol red assay (**Figure**

S3C), respectively.

We did not find any considerable change in the population of cells producing ROS, infected

with wild-type (3.06%; **Figure S3A**) (2.157 ± 0.1611) % (**Figure S3B**), knockout (4.09%;

Figure S3A) (2.192 ± 0.2955) % (**Figure S3B**), and the complement strains (3.81%; **Figure**

S3A) (2.61 ± 0.2244) % (**Figure S3B**) of *Salmonella* (**Figure S3A and S3B**). We further

checked the accumulation of H2O2 in the culture supernatants of infected RAW264.7 cells to

validate this observation. But there was hardly any difference in H_2O_2 production (**Figure**

S3C). The [NO] produced from the cellular pool of L- arginine by the catalytic activity of

inducible nitric oxide synthase (iNOS) is further oxidized into [NO]-adducts (NONOates,

peroxynitrite, nitrite, etc.) with the help of the superoxide ions which have higher oxidation

potential and bactericidal activity [27]. The damage caused by peroxynitrite (ONOO⁻) can be

monitored by checking the recruitment of nitrotyrosine on the surface of intracellular bacteria

using specific anti-nitrotyrosine antibodies by confocal microscopy (**Figure 3C**). During the

late stage of infection in RAW264.7 cells, the STM *ΔompA* strain showed higher surface

recruitment of nitrotyrosine residues and greater colocalization than the wild-type bacteria

(**Figure 3C**), indicating massive damage caused by the bactericidal function of peroxynitrite.

Listeria monocytogenes produce listeriolysin O (LLO), a pore-forming toxin to degrade the

vacuolar membrane to escape lysosomal fusion [28]. Despite having intact OmpA on its

surface, wild-type *Salmonella* residing in the cytoplasm due to the ectopic expression of

listeriolysin O (LLO) showed poor colocalization with nitrotyrosine (**Figure 3C**). These results

suggest the ability of OmpA to protect the cytosolic population of wild-type *Salmonella* against

the RNI of murine macrophages. The greater recruitment of nitrotyrosine on STM *ΔompA*:

LLO, which is comparable to STM *ΔompA,* suggests that LLO does not play any role in

modulating the activity of iNOS. Consistent with this, the intracellular survival of STM (WT):

LLO in macrophages was better than STM *ΔompA* and comparable to that of the wild-type

bacteria (**Figure 3D**). Taken together, we show that OmpA protects intracellular wild-type

Salmonella Typhimurium against the nitrosative stress of macrophages. While proliferating in

macrophages, it utilizes SPI-2 encoded virulent factor SpiC to downregulate inducible nitric

oxide synthase activity in SOCS-3 dependent manner [29, 30]. To verify whether the OmpA

dependent downregulation of the action of iNOS happens in SpiC dependent manner, the

transcript level of *spiC* was measured from wild-type and the mutant bacteria growing in

macrophages (**Figure 3E**). Unlike the wild-type *Salmonella*, STM *ΔompA* was unable to

produce *spiC* within macrophages. The reduced expression of *spiC* by STM (WT): *LLO* and

bafilomycin A treated cells infected with wild-type bacteria not only shows the requirement of

the acidic pH of SCV for the expression of *spiC* but also indirectly authenticates the cytosolic

localization of STM *ΔompA*. To validate this observation, the promoter activity of *spiC* was

measured in STM (WT) and *ΔompA* growing in acidic F media (**Figure 3F**), LB media (**Figure**

3F), and macrophages (**Figure 3G**) by beta-galactosidase assay. No significant change was

observed in the activity of the *spiC* promoter between the wild-type and mutant bacteria growing in acidic F media and LB broth for 12 hours (**Figure 3F**). However, the enhanced activity of the *spiC* promoter in STM (WT) and *ΔompA* growing in F media (pH= 5.4) compared to the bacterial culture obtained from LB broth (which is less acidified compared to the F media) suggests that the expression of the *spiC* gene is regulated by the acidification of the environment around the bacteria. Inside the macrophages, where there exists a notable difference between the localization of wild-type and *ompA* deficient bacteria, a significant drop in the activity of the *spiC* promoter was observed in STM *ΔompA* (**Figure 3G**). The localization of STM *ΔompA* in the cytosol of macrophages where the pH is relatively higher than SCV can be held accountable for the abrogated expression of *spiC*. To determine the degree of acidification of the cytosol of STM (WT) and *ΔompA* upon alteration in the pH of the surrounding media, we have used a pH-sensitive dye BCECF-AM. We observed a higher 488 nm/ 405 nm ratio of STM *ΔompA* labeled with 20 µM concentration of BCECF-AM resuspended in phosphate buffer of acidic pH (range- 5.5, 6, and 6.5) in comparison with STM (WT) and the complemented strain (*ΔompA*: pQE60-*ompA*) (**Figure 3H**). This result suggests reduced acidification of the cytosol of STM *ΔompA* compared to STM (WT) and STM *ΔompA*: pQE60-*ompA* even when they are present in the same (*in vitro*) acidic environment (**Figure 3H**). Surprisingly when all these strains were incubated separately in a phosphate buffer of pH= 7, we found a comparable 488 nm/ 405 nm ratio of BCECF-AM, which unveils an uncharacterized novel but controversial role of OmpA in the acidification of the cytosol of *S.* Typhimurium in response to extracellular acidic stress.

Inhibition of the activity of iNOS improves the survival of STM *ΔompA* in the *in vitro* and *in vivo* infection models.

To investigate further the pro-bacterial role of *Salmonella* OmpA against the nitrosative stress

of macrophages, we treated the cells with 1400W dihydrochloride (10 µM), an irreversible

inhibitor of inducible nitric oxide synthase (iNOS), the key enzyme in the production of [NO]

(**Figure 4A, 4C, and 4D**), or mouse IFN-ɣ that upregulates the expression of iNOS (at 100 U/

mL concentration) (**Figure 4B, 4C, and 4E**). Inhibition of iNOS using 1400W completely

restored the intracellular proliferation of STM *ΔompA* (43.61 ± 6.837) compared to the

untreated reference (19.32 ± 3.913) (**Figure 4A**). Consistent with this finding, STM *ΔompA*

showed poor colocalization with nitrotyrosine under 1400W treatment (**Figure 4C and 4D**),

which can be attributed to the poor biogenesis of RNI due to the inhibition of iNOS.

Augmenting the iNOS activity of macrophages using mouse IFN-ɣ hindered the intracellular

proliferation of STM *ΔompA* (12.52 ± 1.334) compared to the IFN-ɣ untreated cells (19.64 ±

2.11) (**Figure 4B**). As demonstrated by the confocal image, the enhanced biogenesis and

colocalization of nitrotyrosine with STM ΔompA under IFN-ɣ treatment can be considered for

its higher attenuation in intracellular proliferation (**Figure 4C and 4E**). The reduced CFU

burden of *ompA* deficient *Salmonella* in the liver, spleen, and MLN of C57BL/6 mice

(compared to the wild-type bacteria) strongly supports the role of OmpA in bacterial

pathogenesis (**Figure 4G**). In our *ex vivo* studies, we established the role of OmpA in protecting

the wild-type *Salmonella* against the nitrosative stress of murine macrophages. Further, we

used *iNOS*$^{-/-}$ C57BL/6 mice and treated the wild type C57BL/6 mice with a pharmacological

inhibitor of iNOS called aminoguanidine hydrochloride at a dose of 10 mg/ kg of body weight

from 0th day (On the day of infecting the mice) to 5thday (the day of sacrificing and dissecting

the mice) (**Figure 4G**) [31]. We found a comparable CFU burden of wild type and *ompA*

deficient bacteria in the liver, spleen, and MLN of *iNOS*$^{-/-}$ and wild type C57BL/6 mice

administered with the inhibitor. The higher bacterial burden of STM *ΔompA* from the *iNOS*$^{-/-}$

and aminoguanidine-treated mice compared to PBS treated mice (**Figure 4G**) concomitantly

authenticates our results obtained from *ex vivo* experiments. Likewise, the use of the *gp91phox⁻*

ᐟ⁻ mice unable to produce ROS upon bacterial infection showed comparable CFU burden for

both the wild type and *ΔompA* STM in the liver, spleen, and MLN. A significantly higher

bacterial load of *ompA* deficient STM in the liver of *gp91phox⁻ᐟ⁻* vs. wild type C57BL/6 mice

indicates the inability of *gp91phox⁻ᐟ⁻* mice to generate peroxynitrite due to dampened

production of superoxide ions (ROS) (Figure 3.8G). However, a comparable burden of STM

ΔompA and the wild-type strain in the spleen and MLN of *gp91phox⁻ᐟ⁻* mice and the wild-type

C57BL/6 mice suggests that ROS alone cannot clear the *in vivo* infection by STM *ΔompA*.

Earlier, we found that the infection of RAW264.7 cells with STM *ΔompA* does not induce the

level of intracellular ROS. To show the role of *ompA* in the *in vivo* infection of *Salmonella*

Typhimurium, we challenged 4- 6 weeks old adult BALB/c and C57BL/6 mice (**Figure 4H**)

with a lethal dose (10^8 CFU of bacteria/ animal) of wild type and knockout strains by oral

gavaging. Almost 80% of BALB/c mice infected with STM *ΔompA* survived compared to the

group of mice infected with wild-type bacteria (**Figure 4H**). On the other side, the C57BL/6

mice infected with STM *ΔompA* showed better survival and retarded death than those infected

with the wild-type strain, suggesting a critical role of ompA in the infection caused by

Salmonella.

OmpA dependent regulation of outer membrane permeability in *Salmonella* controls

cytoplasmic redox homeostasis in response to *in vitro* nitrosative stress.

In a mildly acidic environment (pH= 5 – 5.5), $NaNO_2$ dissociates to form nitrous acid (HNO_2),

which undergoes dismutation reaction upon oxidation and eventually generates a wide range

of reactive nitrogen intermediates (RNI) such as nitrogen dioxide (NO_2), dinitrogen trioxide

(N_2O_3), nitric oxide (NO), etc. [32]. After entering the bacterial cells, RNIs cause irreparable

deleterious effects targeting multiple subcellular components such as nucleic acids (cause

deamination of nucleotides), proteins (disruption of Fe-S clusters, heme group; oxidation of

thiols and tyrosine residues), lipids (cause peroxidation), etc., and finally kill the pathogen [33].

To investigate the role of *Salmonella* OmpA against *in vitro* nitrosative stress, we decided to

check *in vitro* sensitivity of the wild type and *ompA* knockout strains in the presence of a

varying concentration (0-5 mM) of H_2O_2 (**Figure S3D**), $NaNO_2$ (**Figure S3E**), and a

combination of H_2O_2 and $NaNO_2$ for 12 h (**Figure S3F**) by CFU and resazurin test. Compared

with wild-type bacteria, STM *ΔompA* did not show any significant difference in viability when

exposed to peroxide (**Figure S3D**). The knockout strain displayed a substantial reduction in

viability compared to the wild-type bacteria at 800 µM concentration when incubated with

$NaNO_2$ alone (**Figure S3E**). The sensitivity of the knockout strain towards acidified nitrite

further increased when H_2O_2 was added to $NaNO_2$ (where the growth inhibition started at 600

µM concentration) (**Figure S3F**). To ensure OmpA dependent protection of wild-type

Salmonella against the damage caused by *in vitro* nitrosative stress, we performed a death

kinetics experiment with wild-type and knockout bacteria in the presence of 800 µM acidified

nitrite (**Figure 5A**). Consistent with our previous observations, the knockout strain showed a

notable hindered growth compared to the wild type at 12 h post-inoculation. Considering the

impeded growth of the mutant strain in the presence of in vitro nitrosative stress, we

hypothesized enhanced entry of nitrite into the *ΔompA* strain compared to the wild type. To

verify this, we performed nitrite uptake assay with the wild type, knockout, complement, and

empty vector strain in MOPS-NaOH buffer with 200 µM initial nitrite concentration. We

noticed a greater time-dependent nitrite uptake by the *ΔompA* and empty vector strain than

wild-type and complemented strains (**Figure 5B**).

To verify OmpA dependent redox homeostasis of *Salmonella* in response to *in vitro* nitrosative

stress, we exposed both the wild type and knockout strains harboring pQE60-Grx1-roGFP2

plasmid to 800 µM, 1 mM, and 5 mM (**Figure 5C**) concentration of acidified nitrite for 15, 30,

45 and 60 minutes. Glutathione (GSH), a low molecular weight thiol of Gram-negative

bacteria, maintains the reduced state of the cytoplasm. In the presence of external ROS/ RNI

stress cytosolic GSH pool is oxidized to form glutathione disulfide (GSSG) [34]. The redox-

sensitive GFP2 (roGFP2) having two redox-sensitive cysteine residues at 147th and 204th

positions (which form disulfide bond upon oxidation) can absorb at two wavelengths, 405 nm

and 488 nm depending upon its oxidized and reduced state, respectively. It has a fixed emission

at 510 nm [35]. The glutaredoxin protein (Grx1) fused to the redox-sensitive GFP2 can

reversibly transfer electrons between the cellular (GSH/GSSG) pool and the thiol groups of

roGFP2 at a much faster rate. The ratio of fluorescence intensity of Grx-roGFP2 measured at

405 nm and 488 nm demonstrates the redox status of the cytoplasm of bacteria [34]. In all the

three concentrations of acidified nitrite {800 µM, 1 mM, 5 mM (**Figure 5C**)}, we found a time-

dependent increase in the 405/ 488 ratio of Grx1-roGFP2 in STM *ΔompA* strain in comparison

with STM (WT), suggesting a heightened redox burst in the cytoplasm of ompA knockout

strain in response to RNI in vitro. Taken together, our data indicate the importance of OmpA

in maintaining the cytosolic redox homeostasis of wild-type *Salmonella*. Earlier, we have

noticed a significant downregulation in the transcript levels of outer membrane-bound larger

porins (*ompC*, *ompD*, and *ompF*) compared to *ompA* in wild-type *Salmonella* growing in acidic

F media and RAW264.7 cells. Our previous study depicted enhanced nitrite uptake by STM

ΔompA strain compared to STM (WT), which further indicates increased permeability of the

bacterial outer membrane. Hence, we decided to check the expression of larger porins in STM

ΔompA strain growing in *in vitro* and *ex vivo* conditions. Surprisingly, in contrast to the wild-

type bacteria, we have found elevated expression of *ompC*, *ompD*, and *ompF* in STM *ΔompA*

strain growing in nutritionally enriched LB media, nutrient-depleted acidic F media (pH= 5.4),

and RAW264.7 macrophage cells (**Figure 5D**). This increased expression of larger porins was

revoked in the complemented strain. In this regard, the enhanced outer membrane

depolarization of STM *ΔompA* growing in acidic F media was further tested (during stationary

phase) using a membrane-permeant negatively charged dye called DiBAC$_4$ (**Figure 5E**) [36].

Because of the enhanced outer membrane permeability, when the negative charge of the bacterial cytosol is diminished by the inflow of cations (depolarization), the cell allows the entry of DiBAC$_4$, which binds to the cell membrane proteins and starts fluorescing.

Contrary to the wild-type and the complemented strain, the higher DiBAC$_4$ positive population and greater median fluorescence intensity of DiBAC$_4$ corresponding to STM *ΔompA* and the empty vector strain (**Figure 5E**) ensures enhanced outer membrane permeability of *Salmonella* in the absence of OmpA. We used another porin-specific DNA binding fluorescent dye called bisbenzimide (Figure 5F) [41] to strengthen this observation. In line with our previous observation, we found that the fluorescence intensity of bisbenzimide taken up by STM *ΔompA* growing in acidic F media is more than the wild-type and complemented bacterial strain (**Figure 5F**). Compared to the wild-type bacteria, the greater fluorescence intensity of bisbenzimide corresponding to STM *ΔompA* grown intracellularly in murine macrophages for 12h (**Figure 5G**) firmly endorsed the result obtained from the in vitro experiment. To show explicitly that the increased expression of larger porins such as *ompC*, *ompD*, and *ompF* on bacterial outer membrane enhances the outer membrane porosity, we have expressed *ompC,* *ompD,* and *ompF* in wild-type *Salmonella* with pQE60 plasmid. We have observed that the increased expression of *ompD* and *ompF* enhanced the permeability of the outer membrane of wild-type *Salmonella* growing overnight in acidic F media (**Figure S4A and S4B**). We have further elevated the expression of *ompC, ompD,* and *ompF* in wild-type *Salmonella* by adding IPTG to the LB broth. Our data suggested that the over-expression of *ompF* in the wild-type *Salmonella* causes massive depolarization (61.62%) of the outer membrane compared to the other porins, namely *ompC* (2.73%) and *ompD* (8.06%) (**Figure S4C and S4D**). Hence, we can conclude that in the absence of *ompA*, the expression of larger porins such as *ompC, ompD,*

and *ompF* increases on the outer membrane of *Salmonella*. However, the enhanced outer

membrane porosity of the bacteria majorly depends upon the elevated expression of *ompF*.

The maintenance of the integrity of the SCV membrane inside RAW264.7 macrophages

solely depends upon OmpA, not on other larger porins such as OmpC, OmpD, and

OmpF.

To strengthen our previous observation, we decided to knockout *ompC*, *ompD*, and *ompF*

individually in the kanamycin-resistant *ΔompA* background of *Salmonella*. We have generated

STM *ΔompA ΔompC*, STM *ΔompA ΔompD,* and STM *ΔompA ΔompF* using chloramphenicol

resistant gene cassette (data not shown). We further investigated the intracellular niche of these

double knockout bacterial strains in RAW264.7 cells during the late phase of infection (16 h

post-infection). In line with our previous finding, compared to the wild-type bacteria, STM

ΔompA showed poor colocalization with SCV marker LAMP-1 (**Figure 6A and 6B**). The

drastic loss of SCV membrane from the surroundings of STM *ΔompA ΔompC*, *ΔompA ΔompD*,

and *ΔompA ΔompF* as demonstrated by their poor colocalization with LAMP-1 (**Figure 6B**),

indicating the cytosolic localization of these double knockout strains in macrophages during

the late phase of infection. The reduced colocalization of STM (WT) expressing LLO (which

usually stays in the cytoplasm) with LAMP-1 (**Figure 6A and 6B**) authenticates the cytosolic

phenotype of the double knockout strains. To rule out the possibility that the lack of OmpC,

OmpD and OmpF also contributes to the cytosolic localization of STM *ΔompA ΔompC*, *ΔompA*

ΔompD, and *ΔompA ΔompF* (**Figure 6A and 6B**), we generated single knockout strains of

Salmonella lacking *ompC*, *ompD*, and *ompF* using the method demonstrated earlier. We

observed that the STM *ΔompC*, *ΔompD*, and *ΔompF* strains colocalized with LAMP-1

similarly to the wild-type strain (**Figure 6C and 6D**). Hence, it can be concluded that the

maintenance of the vacuolar life of wild-type *Salmonella* depends on OmpA and not on OmpC,

OmpD, and OmpF. A decreased recruitment of nitrotyrosine on STM *ΔompC*, *ΔompD*, and

ΔompF in comparison with STM *ΔompA* while growing intracellularly in RAW264.7 cells

further supports the presence of intact SCV membrane around them (**Figure S5A**). The *in vitro*

sensitivity of STM *ΔompC*, *ΔompD*, and *ΔompF* against the acidified nitrite was also checked

(**Figure S5B**). It was observed that, unlike the *ompA* deficient strain of *Salmonella*, the ability

of STM *ΔompC*, *ΔompD*, and *ΔompF* to withstand the bactericidal effect of acidified nitrite is

comparable with the wild type bacteria (**Figure S5B**). To further support this observation, we

checked nitrite consumption by STM *ΔompC*, *ΔompD*, and *ΔompF* (**Figure S5C**). It was found

that in comparison to STM *ΔompA* strain, *ΔompC*, *ΔompD*, and *ΔompF* are more efficient in

restricting the entry nitrite (**Figure S5C**), which can be attributed to their better survival in the

presence of *in vitro* acidified nitrite (**Figure S5B**).

In the absence of OmpA, porins OmpC and OmpF enhance the susceptibility of

***Salmonella* against nitrosative stress of RAW264.7 cells.**

To dissect the role of each larger porin, namely OmpC, OmpD, and OmpF in the entry of nitrite

into bacteria during the absence of OmpA, we performed nitrite uptake assay using the double

knockout bacterial strains STM *ΔompA ΔompC*, *ΔompA ΔompD*, and *ΔompA ΔompF* (**Figure**

7A). In comparison with the wild-type bacteria, the rapid disappearance of nitrite, which

corroborates substantial nitrite uptake from the media by STM *ΔompA* and *ΔompA ΔompD,*

demonstrates the involvement of OmpC and OmpF (present in both STM *ΔompA* and *ΔompA*

ΔompD strains) in this entry process (**Figure 7A**). This result was further validated by the

measurement of the reduced *in vitro* percent viability of STM *ΔompA* and *ΔompA ΔompD* in

comparison with wild type, *ΔompA ΔompC* and *ΔompA ΔompF* strains of *Salmonella* in the

presence of acidified nitrite (800 μM) (**Figure 7B**). Using the *ex vivo* system of infection in

murine macrophages, we have found enhanced recruitment of nitrotyrosine on STM *ΔompA*

and *ΔompA ΔompD* as depicted by their higher colocalization in comparison with wild type,

ΔompA ΔompC, and *ΔompA ΔompF* (**Figure 7C**) *Salmonella*. The intracellular nitrosative burst of macrophages infected with STM (WT), *ΔompA, ΔompA: ompA, ΔompA ΔompC, ΔompA ΔompD*, and *ΔompA ΔompF* was tested using DAF2DA (**Figure 7E**). Only 1.94% of macrophages infected with the wild-type bacteria produced [NO], which is comparable to STM *ΔompA: ompA* (1.66%), *ΔompA ΔompC* (1.15%) and *ΔompA ΔompF* (1.62%) infected macrophages population producing [NO]. The greater percent population of DAF2DA positive infected macrophages corresponding to STM *ΔompA* (3.78%) and *ΔompA ΔompD* (3.77%) (**Figure 7E**) strengthens the conclusion obtained from confocal data (**Figure 7C**). The macrophages infected with STM (WT): *LLO* showed a very low intracellular nitrosative burst (0.9%), suggesting that the cytosolic population of wild-type *Salmonella* expressing LLO was protected by the presence of OmpA (**Figure 7E**). This data was further verified by the attenuated intracellular proliferation of STM *ΔompA* and *ΔompA ΔompD* inside RAW264.7 cells (**Figure 7D**). It was found that unlike STM (WT), *ΔompA ΔompC*, and *ΔompA ΔompF*, which are showing very poor colocalization with intracellular nitrotyrosine, the intracellular proliferation of STM *ΔompA* and *ΔompA ΔompD* was severely compromised in murine macrophages (**Figure 7C and 7D**). Taken together, it can be concluded that in the absence of OmpA, the elevated expression of two major larger porins, namely OmpC and OmpF, on the bacterial outer membrane may help in the entry of nitrite into the bacterial cytoplasm and makes the bacteria highly susceptible to the intracellular nitrosative burst. To validate this observation, we have overexpressed *ompC, ompD,* and *ompF* in wild-type *Salmonella* and checked the susceptibility of these bacterial strains against *in vitro* nitrosative stress. We have incubated the wild-type *Salmonella,* overexpressing *ompA, ompC, ompD,* and *ompF* in the presence of acidified nitrite for 12 hours and checked the viability of the bacteria by flow cytometry (propidium iodide staining) (**Figure S6A and S6B**) and resazurin assay (**Figure S6C, S6D, and S6E**). Earlier, we have shown that over-expression of both *ompD* and *ompF* enhances the

permeability of the bacterial outer membrane. In line with our previous observation, the

flowcytometric data has shown that compared to *ompC,* the over-expression of *ompD* (14.97%)

and *ompF* (16.17%) in wild-type *Salmonella* can induce bacterial death in the presence of

acidified PBS (**Figure S6A**). However, over-expressing *ompF* in the wild-type *Salmonella*

alone is responsible for increasing the susceptibility of the bacteria (22.8%) towards acidified

nitrite (**Figure S6A- S6E**). This suggests that out of all three major larger porins, OmpF plays

a pivotal role in increasing the susceptibility of *ompA* deficient *Salmonella* against *in vitro* and

in vivo nitrosative stress.

Discussion

Bacterial pathogens (such as *Acinetobacter baumannii*, *E. coli*, *K. pneumoniae,* etc.) can

restrict the entry of toxic molecules such as antibiotics and cationic antimicrobial peptides

either by changing the outer membrane permeability [37, 38] or by augmenting options/

proteases that degrade the antimicrobial peptides [39]. The alteration in the outer membrane

permeability of Gram-negative pathogen is strictly regulated by the core oligosaccharide

composition of outer membrane lipopolysaccharide and differential expression of outer

membrane porins [40-42]. OmpA, OmpC, OmpD, OmpF, and PhoE, the predominantly found

porins on the outer membrane of *Salmonella* Typhimurium, are involved in a wide range of

functions. OmpA tightly holds the outer membrane of the bacteria to the peptidoglycan layer of the cell

wall, and hence, deletion of OmpA aggravates the biogenesis of outer membrane vesicles (OMV) [7, 43].

OmpD, the most abundant porin on *Salmonella* outer membrane, is involved in the uptake of

H_2O_2 [44]. OmpC in conjunction with OmpF contributes to the acquisition of cations, whereas

PhoE majorly helps in the uptake of negatively charged phosphate groups [8]. The correlation

between outer membrane porins and *Salmonella* pathogenesis has been poorly understood due

to the lack of extensive studies. Earlier, Heijden *et al.* showed that HpxF⁻ *Salmonella* altered

outer membrane permeability by reciprocally regulating the expression of OmpA and OmpC

when exposed to H_2O_2 [9]. OmpA deficient *Salmonella* was found to be incapable of reaching

the mouse brain [45]. Alternative sigma factor of *Salmonella* Typhimurium regulates the

expression of OmpA within macrophages to trhive in the presence of oxidative stress [46].

Deletion of *ompD* from the genome of *Salmonella* made it hyper-proliferative in RAW264.7

macrophages and BALB/c mouse models [8]. This study delineated the individual

contributions of OmpA, OmpC, OmpD, and OmpF in *Salmonella* pathogenesis. Our study

revealed the role of porins in the maintenance of the intravacuolar life of wild-type *Salmonella*

Typhimurium and deciphered their role in the regulation of outer membrane permeability and

bacterial resistance to the nitrosative stress of macrophages.

We have found a time-dependent steady decrease in the transcript levels of *ompC*, *ompD*, and

ompF (**Figure S1B, S1C, and S1D**) in *Salmonella* growing in RAW264.7 cells. On the

contrary, the consistently elevated expression of *ompA* during the late phase of infection in

RAW264.7 cells (9 h and 12 h post-infection) (**Figure S1A**) suggested the importance of this

porin in bacterial survival inside the macrophages. Our observations corroborate the findings

of Eriksson *et al.* and Hautefort *et al.* [10, 11]. Unlike other significant porins, OmpA has a

small pore size and a unique periplasmic domain (**Figure S1A, S1B, S1C, and S1D**), which

can act as a gate to restrict the entry of many toxic molecules [9]. Hence it can be concluded

that the wild-type bacteria growing in a nutrient-depleted stressed environment of SCV within

the macrophages, where the bacterial growth is severely challenged by acidic pH, will prefer

OmpA, a porin with a smaller pore size and periplasmic gating mechanism to be expressed on

its outer membrane for the survival. OmpA, one of the most abundant porins of the outer

membrane of *E. coli* K1, the causative agent of neonatal meningitis, is highly conserved in the

family of *Enterobacteriaceae* [4]. Besides its structural role in the bacterial outer membrane,

porins interact with host immune cells. OmpA of *E. coli* K1 and *Enterobacter sakazakii* has

been reported to be involved in the invasion of hBMEC cells and INT407 cells by multiple studies [47-49]. Earlier, multiple groups reported that an eight-stranded β barrel outer membrane porin (*ompW*) of *Escherichia coli* helps the bacteria evade phagocytosis and confers resistance against alternative complement activation pathway mediated killing by the host [5, 6]. Another study carried out by March C *et al*. suggested that the *ompA* mutant strain of *Klebsiella pneumoniae* is severely attenuated in the pneumonia mouse model [2]. OmpA of *E. coli* K1 not only augments complement resistance by recruiting C4 binding protein (C4BP) on the surface of the bacteria [50] but also aggravates their intracellular survival in murine and human macrophages [51]. In our study, the OmpA deficient strain of *Salmonella* has shown an inclination towards greater phagocytosis and severe attenuation in intracellular proliferation in macrophages. The increased macrophage-mediated intake of complement coated STM *ΔompA* compared to complement coated wild-type bacteria proved that wild-type *Salmonella* Typhimurium impairs the complement activation in OmpA dependent manner. However, the detailed mechanism will be addressed in the future. Surprisingly, the deletion of *ompA* makes *Salmonella* invasion deficient and hyper-proliferative in the epithelial cells. The successful systemic infection of *Salmonella* in macrophages and epithelial cells depends upon its intravacuolar inhabitation. The acidification inside the SCV is a prerequisite for the synthesis and secretion of SPI-2 encoded virulent proteins required for *Salmonella*'s successful proliferation [17]. Mild tampering with the integrity of SCV may create unusual outcomes in the bacterial burden of the cells. Earlier it has been reported that the intracellular proliferation of *sifA* mutant of *Salmonella,* which comes into the cytosol after quitting the SCV, was severely abrogated in macrophages. On the contrary, the bacteria become hyperproliferative in the cytosol of epithelial cells [26]. The introduction of point mutations in the Rab5 or Rab7 proteins (markers of SCV) can also trigger the release of wild-type bacteria from SCV to the cytosol of epithelial cells [52]. Our immunofluorescence microscopy data and the result from chloroquine

resistance assay showed the cytosolic localization of ompA deficient strain of *Salmonella* in

macrophages and epithelial cells. The wild-type intracellular *Salmonella* recruits SPI-2

encoded translocon proteins SseC and SseD on the surface of SCV to form a functionally active

T3SS needle complex. Eventually, wild-type bacteria secret these translocon proteins into the

cytosol of the host cells to establish an actively proliferating niche by manipulating the host

signaling cascade. The synthesis, surface accumulation, and secretion of SseC and SseD into

the host cell cytosol depend upon the acidic pH and integrity of SCV [21]. In line with our

expectation, STM *ΔompA* staying in the cytoplasm of macrophages (neutral pH) has exhibited

poor colocalization with SseC and SseD. We further checked the expression of *sseC* and *sseD*

from intracellularly growing bacteria. We have found that STM *ΔompA* is unable to produce

sseC and *sseD* like wild-type bacteria, which is the reason behind the poor secretion of SseC

and SseD into the cytosol of the macrophages. The cytosolic stay of STM *ΔompA* hampers the

acidification of the cytosol of bacteria within macrophages, which further reduces the

expression of several SPI-2 encoded virulent genes such as *ssaV* and *sifA*. Earlier, it has been

proved that *sifA⁻ Salmonella* comes into the cytosol of the host cells after quitting the SCV and

exhibits defective proliferation in the macrophages [53]. Taken together, the result obtained

from the intracellular proliferation assay of STM *ΔompA* in macrophages and epithelial cells

and the immunofluorescence microscopy data on the vacuolar/ cytosolic status of the bacteria

are consistent with the available pieces of literature. Hence, to the best of our knowledge, this

is the first report commenting on the role of *Salmonella* Typhimurium outer membrane protein

A (OmpA) to maintain the stability of the SCV within macrophages. However, the effect

exerted by OmpA in maintaining the integrity of SCV membrane is indirect and dependent on

the reduced expression of *sifA*.

The SCV membrane functions as a protective barrier around wild-type bacteria. Once the

intactness of the SCV membrane is lost, the bacteria will eventually be exposed to the threats

present in the cytosol in the form of ROS and RNI [25]. Earlier, Bonocompain. G *et al.* reported that *Chlamydia trachomatis* infection in HeLa cells transiently induces ROS for the initial few hours of infection [54]. The epithelial cells (HeLa) cannot challenge wild-type *Salmonella* with ROS during infection as efficiently as the macrophages [25]. The generation of RNI indirectly depends upon the ROS burden of a cell. The epithelial cells, a poor producer of ROS, are assumed to be generating a lower level of RNI., which is insufficient to kill the cytosolic population STM *ΔompA* in epithelial cells. On the contrary, the RAW264.7 macrophages can produce both ROS and RNI upon bacterial infection, which might explain the attenuated proliferation of STM *ΔompA* in macrophages. Hence, the oxidative and nitrosative burst of macrophages infected with wild type and the *ompA* knockout strain of *Salmonella* was checked. Surprisingly, we found a remarkable rise in the level of intracellular and extracellular [NO] of macrophages infected with STM *ΔompA*, indicating the protective role of OmpA against the nitrosative stress of macrophages. In continuation with the previous observation, STM *ΔompA,* which has already quit the SCV and stayed in the cytosol of macrophages, showed greater colocalization with nitrotyrosine when compared with the wild-type bacteria protected inside the SCV. To further establish the role of *Salmonella* OmpA against the cytosolic nitrosative stress of macrophages, we have ectopically expressed listeriolysin O (LLO) in wild-type *Salmonella* Typhimurium. The intracellular population of *Listeria monocytogenes*, a causative agent of listeriosis, utilizes LLO to degrade the phagosomal membrane for escaping lysosomal fusion [28, 55]. Expressing LLO in wild-type *Salmonella* will force the bacteria to quit the SCV and be released into the cytosol with intact OmpA on their outer membrane. A decreased recruitment of nitrotyrosine on STM (WT): *LLO* staying in the cytosol of macrophages and their better survival compared to STM *ΔompA* and *ΔompA*: *LLO* proved the role of OmpA in defending the cytosolic population of wild-type *Salmonella* from the harmful effect of RNIs. On the contrary, the higher recruitment of the nitrotyrosine on STM *ΔompA* and STM *ΔompA*:

LLO can be attributed to their attenuated intracellular proliferation compared to STM (WT) and STM (WT): *LLO.* The alteration in the proliferation of STM *ΔompA* in macrophages and the recruitment profile of nitrotyrosine upon manipulating iNOS activity using specific, irreversible inhibitor 1400W dihydrochloride and activator IFN-γ fell in line with the previous results. Wild-type *Salmonella*, with the help of its pathogenicity island (SPI)- 2 encoded virulent factor SpiC activates the suppressor of cytokine signaling 3 (SOCS-3) which inhibit IFN-γ signaling and thus eventually represses the activity of iNOS [29, 30]. As discussed earlier, the acidification of the SCV compartment acidifies the cytoplasm of wild-type *Salmonella*. This process of acidification is essential to synthesize and secret SPI-2 effector proteins into the host cell's cytoplasm [17]. We have also found that STM *ΔompA,* which stays in the neutral pH of cytosol in macrophages cannot produce SpiC. The decreased expression of *spiC* activates the iNOS in the macrophages infected with STM *ΔompA*. We further checked the promoter activity of *spiC* in wild-type and mutant bacteria growing extracellularly and intracellularly by β galactosidase assay. Compared to the intracellularly growing STM *ΔompA*, the higher activity of *spiC* promoter in the wild-type *Salmonella* simultaneously supports the cytosolic inhabitation of STM *ΔompA* and answers the reason behind the nitrosative burst of macrophages. The downregulation of SPI-2 encoded gene *spiC* in STM *ΔompA* can also be attributed to its incompetence to acidify its cytosol in response to extracellular acidic response, which was determined by a higher 488 nm/ 405 nm ratio of BCECF-AM. But strikingly, STM *ΔompA* was unable to induce intracellular and extracellular ROS production while infecting macrophages. NADPH phagocytic oxidase is the key enzyme of macrophages involved in the synthesis of superoxide ions. It is a multimeric protein consisting of two membrane-bound subunits such as $gp91^{phox}$, $p22^{phox}$, and four cytosolic subunits such as $p47^{phox}$, $p40^{phox}$, $p67^{phox,}$ and RacGTP [56]. Continuing with our previous observations, we found poor colocalization of $gp91^{phox}$ with STM (WT) and *ΔompA* in macrophages during the late phase of infection (data

not shown). Wild-type *Salmonella* can inhibit the recruitment of NADPH oxidase on the surface of the SCV membrane with the help of the SPI-2 encoded type 3 secretion system [57]. The restriction in the recruitment of NADPH oxidase on the surface of the *ompA* knockout bacteria lacking SCV membrane and staying in the cytosol is the probable reason behind the dampened oxidative stress inside the infected macrophages. On the contrary, the ability of iNOS to maintain its uninterrupted catalytic activity, despite being recruited on the cortical actin of macrophages [58], is the most probable reason behind elevated production of RNI in RAW264.7 cells infected with STM *ΔompA*. To investigate the role of OmpA in the establishment of *in vivo* infection by wild-type *Salmonella* Typhimurium we have used 4 to 6 weeks old *Nramp*[-/-] C57BL/6 and BALB/c mice. The better survival of the mice infected with *ompA* deficient strains of *Salmonella*, which was attributed to the reduced bacterial burden in the liver, spleen, and MLN, finally endorsed the essential role of OmpA in the *in vivo* infection of *Salmonella*. Administration of iNOS inhibitor aminoguanidine hydrochloride by intraperitoneal injection or direct oral infection of *iNOS*[-/-] mice diminished the *in vivo* attenuation of STM *ΔompA,* suggesting OmpA dependent protection of wild type bacteria against nitrosative stress *in vivo*. The comparable CFU burden of STM (WT) and *ΔompA* in *gp91phox*[-/-] C57BL/6 mice, which cannot produce ROS, can be accounted for the abrogated peroxynitrite response [59, 60]. Acidified nitrite generating a wide range of reactive nitrogen intermediates (RNI) such as nitrogen dioxide (NO_2), dinitrogen trioxide (N_2O_3), nitric oxide (NO), etc., are extensively used for assessing *in vitro* viability of bacteria and fungi [32, 61]. After entering the bacterial cells, RNIs cause enormous irreparable damages to multiple subcellular components such as nucleic acids (cause deamination of nucleotides), proteins (disruption of Fe-S clusters, heme group; oxidation of thiols and tyrosine residues), lipids (cause peroxidation), etc., and eventually destroy the pathogen. The enhanced sensitivity of STM *ΔompA* against acidified nitrite and combination of acidified nitrite with peroxide, which

generates peroxynitrite [62], suggested increased entry of nitrite in the knockout strain comparison with wild type bacteria. The faster depletion of nitrite from the media having STM *ΔompA* strongly supported our hypothesis. The time-dependent gradual increase in the 405/ 488 ratio of pQE60-Grx1-roGFP2 harbored by STM *ΔompA* under acidified nitrite's tested concentrations proved the loss of redox homeostasis in STM *ΔompA*.

As discussed earlier, the outer membrane of Gram-negative bacteria acts as an impenetrable barrier to many toxic compounds (including antibiotics, bile salts, cationic antimicrobial peptides, reactive oxygen species, abnormal pH, osmotic stress, etc.) and protect the bacteria from environmental threats during its extracellular and intracellular life-cycle. The enhanced uptake of nitrite by *Salmonella* in the absence of OmpA and a time-dependent increase in the sensitivity of STM *ΔompA* towards acidified nitrite proves the occurrence of a permanent damage to the outer membrane of the bacteria. The increased uptake of $DiBAC_4$ (measure the cell membrane depolarization) and bisbenzimide (binds to the DNA) by the *ompA* deficient strain of *Salmonella* provided further supports to our conclusion. To rationalize the enhanced uptake of fluorescent probes by STM *ΔompA,* we checked the expression of larger porins such as *ompC*, *ompD*, and *ompF* in wild-type, mutant, and complemented strains of *Salmonella* growing in LB broth, F media, and RAW264.7 macrophages. We found a remarkable increase in the expression of the larger porins in the knockout strain (lacking OmpA) compared to wild-type and complemented strains of *Salmonella*. Hence, we have concluded that OmpA regulates the outer membrane stability of *Salmonella* Typhimurium. When OmpA is deleted, the increased expression of larger porins reduces the stability of the outer membrane and makes it permeable to nitrites, which eventually kills the bacteria. To best of our knowledge, this is the first report where we have experimentally dissected the role of OmpA in maintaining outer membrane stability of *Salmonella* Typhimurium. To strengthen our conclusion, we have further overexpressed *ompC*, *ompD*, and *ompF* in the wild-type *Salmonella* with the help of a low copy

number plasmid pQE60 that has a site for 'lac operator.' We over-expressed these larger porins

in wild-type bacteria after incubating them with appropriate antibiotics and 500 µM of IPTG.

Compared to other larger porins, the remarkably improved uptake of DiBAC$_4$ by wild-type

Salmonella expressing *ompF* pinpoints the contribution of OmpF in escalating the reduction of

outer membrane stability in the absence of OmpA. This conclusion can be further extrapolated

to understand the reason behind the enhanced recruitment of nitrotyrosine on STM *ΔompA*

compared to STM (WT): *LLO* in macrophages during the late phase of infection. Despite

having cytosolic niche STM (WT): *LLO* has intact OmpA in its outer membrane, unlike STM

ΔompA, which maintains the integrity and stability of the outer membrane and prohibits the

entry of peroxynitrite. This further proves that the outer membrane defect of STM *ΔompA*

makes the bacteria accessible to RNI produced by the macrophages.

To ascertain the individual contribution of the larger porins such as *ompC*, *ompD*, and *ompF* in

the entry of nitrite in *ompA* deficient bacteria, we constructed *ompC*, *ompD*, and *ompF* single

and double knockout strains in wild type and *ompA*⁻ background of *Salmonella*, respectively.

The cytosolic localization of double knockout strains, namely STM *ΔompA ΔompC*, *ΔompA*

ΔompD, and *ΔompA ΔompF* and vacuolar imprisonment of the single knockout strains such as

STM *ΔompC*, *ΔompD*, and *ΔompF* provided vital support to our previous conclusion on the

dependence of SCV integrity and stability on OmpA. Under the *in vitro* challenge of acidified

nitrite, the better survival of STM *ΔompC*, *ΔompD*, and *ΔompF* in comparison with *ΔompA* not

only suggested the abrogated consumption of nitrite but also indisputably established the

paramount importance of OmpA in the maintenance of the outer membrane permeability of

wild-type *Salmonella*. STM *ΔompA ΔompD,* which possesses intact OmpC and OmpF on their

outer membrane, showed enhanced nitrite consumption and higher sensitivity towards *in vitro*

nitrosative stress (800 µM of acidified NaNO$_2$). In comparison with STM *ΔompA ΔompC* and

ΔompA ΔompF, greater recruitment of nitrotyrosine on the cytosolic population of STM *ΔompA*

ΔompD (having OmpC and OmpF) due to the significant loss of outer membrane stability is

considered as the sole reason behind their poor proliferation in murine macrophages. These

results were further supported by the inability of STM *ΔompA ΔompC* and *ΔompA ΔompF* to

induce heightened [NO] response while staying inside the macrophages, unlike STM *ΔompA*

ΔompD. To strongly endorse this result, we have decided to verify the viability of wild-type

Salmonella against the *in vitro* nitrosative stress upon expressing *ompC*, *ompD*, and *ompF*. In

line with our previous observations, we have found that compared to *ompD* and *ompF,* the over-

expression of *ompF* in wild-type *Salmonella* drastically increases the bacteria's susceptibility

towards acidified nitrites. These results collectively suggest the pivotal role of OmpF in the

entry of nitrite in *ompA* deficient *Salmonella* Typhimurium by increasing the permeability and

worsening the integrity of the bacterial outer membrane. In this context, we must mention that

STM *ΔompA ΔompC* is also expected to express OmpD and OmpF on its outer membrane.

Earlier, we have shown that compared to OmpC, OmpD and OmpF contribute more in the

depolarization of the bacterial outer membrane. However, OmpF is solely responsible for the

death of the bacteria in the presence of *in vitro* nitrosative stress. Despite having OmpF on their

outer membrane, the better survival of STM *ΔompA ΔompC* compared to STM *ΔompA ΔompD*

in response to *in vitro* and e*x vivo* nitrosative stress might be questioned by the conclusions of

our study, which we will answer in the future.

To summarize, our study claims OmpA of *Salmonella* Typhimurium to be a versatile protein

with a multitude of activities. The deletion of *ompA* from *Salmonella* interrupts the stability of

SCV and imposes significant paradoxical consequences on the intracellular proliferation of

bacteria in the macrophages and epithelial cells, respectively. We have experimentally proved

that OmpA maintains the stability of the bacterial outer membrane. In the absence of OmpA,

the porosity of the outer membrane increases, which makes the bacteria vulnerable to *in vitro*

and *in vivo* nitrosative stress. We proposed an OmpA dependent mechanism that regulated the

stability of the bacterial outer membrane and was employed cleverly by *Salmonella* to fight

against the nitrosative stress of murine macrophages.

Abbreviations

STM: *Salmonella* Typhimurium

OmpA: Outer membrane protein A

OmpC: Outer membrane protein C

OmpD: Outer membrane protein D

OmpF: Outer membrane protein F

LLO: Listeriolysin O

SCV: *Salmonella* containing vacuole

LAMP-1: Lysosome associated membrane protein-1

iNOS: Inducible nitric oxide synthase

RNI: Reactive nitrogen intermediates

ROS: Reactive oxygen species

RFP: Red fluorescent protein

roGFP2: Redox sensitive green fluorescent protein

Materials and Methods

Bacterial strains, media, and culture conditions

The wild type (WT) bacteria *Salmonella enterica* serovar Typhimurium [STM- (WT)] strain

14028S used in this study was a generous gift from Professor Michael Hensel, Max Von

Pettenkofer-Institute for Hygiene und Medizinische Mikrobiologie, Germany. The bacterial

strains were revived from glycerol stock (stored in -80^{0}C) and plated either on LB agar

(purchased from HiMedia) or LB agar along with appropriate antibiotics like- kanamycin (50

μg/mL), ampicillin (50 μg/mL), Chloramphenicol (25 μg/mL), kanamycin and ampicillin (both

50 μg/mL), Kanamycin and Chloramphenicol (Kanamycin= 50 μg/mL, Chloramphenicol= 25

μg/mL) for wild type, knockout (single and double), complement and mCherry expressing

strains respectively. *Salmonella- Shigella* agar was used to plating cell lysates/ cell suspensions

to calculate the bacterial burden in infected cell lines and several organs of infected mice.

Bacterial LB broth cultures were grown in a shaker incubator at 180 rpm, either 37^{0}C for typical

wild type and knockout strains or 30^{0}C for the strains having temperature-sensitive pKD46

plasmid or the strains harboring pQE60-Grx1-roGFP2 plasmid and undergoing IPTG

(concentration= 500 μM) treatment. For growth curve experiments and *in vitro* RNA extraction

studies, a single colony was inoculated in 5mL of LB broth and grown overnight with or

without antibiotics at 37^{0}C. Overnight-grown stationary phase bacteria were sub-cultured at a

ratio of 1: 100 in freshly prepared LB or minimal F media (acidic) and kept in a 37^{0}C shaker

incubator. At different time intervals, aliquots were taken for RNA isolation, serial dilution,

plating, and [OD]600nm measurement by TECAN 96 well microplate reader. The complete

list of strains and plasmids has been listed below. (Description in Table- 1) Dead bacteria used

in several experiments were produced from viable wild-type bacteria either heating at 650C for

20 minutes or treating the bacteria with 3.5% paraformaldehyde for 30 minutes.

Eukaryotic cell lines and growth conditions

The murine macrophage-like cell line RAW 264.7, human cervical adenocarcinoma cell line

HeLa, human colorectal adenocarcinoma cell line Caco-2 were maintained in Dulbecco's

Modified Eagle's Media (Sigma-Aldrich) supplemented with 10% FCS (Fetal calf serum,

Gibco) at 37^{0}C temperature in the presence of 5% CO_2. Human monocyte cell line U937 cells

were maintained in Roswell Park Memorial Institute 1640 media (Sigma-Aldrich)

supplemented with 10% FCS (Fetal calf serum, Gibco). For polarizing the Caco-2 cells,

DMEM media was further supplemented with 1% non-essential amino acid solution (Sigma-

Aldrich). Phorbol Myristate Acetate (Sigma-Aldrich) (concentration- 20 ng/ mL) was used for

the activation of U937 cells for 24 hours at 37^0C temperature in the presence of 5% CO_2,

followed by the replacement of the media carrying PMA with normal RPMI supplemented with

10% FCS and further incubating the cells for 24 hours before starting the experiments.

Construction of *ompA, ompC, ompD, & ompF* knockout strains of *Salmonella*

The knockout strains of *Salmonella enterica* serovar Typhimurium (strain 14028S) were made

using one step chromosomal gene inactivation method demonstrated by Datsenko and Wanner

[13]. Briefly, STM (WT) was transformed with pKD46 plasmid, which has a 'lambda red

recombinase system' under arabinose inducible promoter. The transformed cells were grown

in LB broth with ampicillin (50 μg/mL) and 50 mM arabinose at 30^0C until the $[OD]_{600nm}$

reached 0.35 to 0.4. Electrocompetent STM pKD46 cells were prepared after washing the

bacterial cell pellet thrice with double autoclaved chilled Milli Q water and 10% (v/v) glycerol.

Finally, the electrocompetent STM pKD46 cells were resuspended in 50 μL of 10% glycerol.

Kanamycin resistant gene cassette (Kan^R, 1.6kb- for knocking out *ompA, ompD*) and

chloramphenicol resistant gene cassette (Chl^R, 1.1 kb- for knocking out *ompC, ompD & ompF*)

were amplified from pKD4 and pKD3 plasmids, respectively using knockout primers (Table-

3.2). The amplified Kan^R and Chl^R gene cassettes were subjected to phenol-chloroform

extraction and electroporated into STM (WT) pKD46. The transformed cells were plated on

LB agar with kanamycin (50 μg/mL) for selection of *ΔompA, ΔompD* strains and LB agar with

chloramphenicol (25 μg/mL) for selection of *ΔompC, ΔompD,* and *ΔompF* strains. The plates

were further incubated overnight at 37^0C. The knockout colonies were confirmed by

confirmatory and kanamycin/ chloramphenicol internal primers (Table- 2).

Construction of *ompA, ompC, ompD, ompF* complemented strain of *Salmonella*

The *ompA, ompC, ompD,* and *ompF* genes were amplified with their respective cloning primers (Description in Table- 3.2) by colony PCR. The amplified PCR products, purified by phenol-chloroform extraction, was subjected to restriction digestion by specific restriction enzymes such as NcoI (NEB) and HindIII (NEB) for *ompA*, BamHI (NEB), and HindIII (NEB) for *ompC, ompD,* and *ompF* in the CutSmart buffer (NEB) at 37^0C for 2-3 h along with the empty pQE60 vector backbone. Double digested insert and vector were subjected to ligation by a T_4 DNA ligase in 10X ligation buffer (NEB) overnight at 16^0C. The ligated products and the empty vector were transformed into respective bacterial strains separately to generate complemented, over-expression, and empty vector strains. Complementation and over-expression were initially confirmed by colony PCR with cloning and *ompA, ompC, ompD,* and *ompF* internal primers (data not shown) and finally by restriction digestion of recombinant plasmid. The expression level of *ompA* in the knockout, complemented, and empty vector strains were further confirmed by RT PCR using *ompA* specific RT primers (Description in Table- 2).

Construction of *ΔompA ΔompC*, *ΔompA ΔompD*, & *ΔompA ΔompF* double knockout strains of *Salmonella*

The double knockout strains of *Salmonella enterica* serovar Typhimurium (strain 14028S) were made by slightly modifying the one-step chromosomal gene inactivation strategy demonstrated by Datsenko and Wanner [13]. Briefly, STM *ΔompA*, where the *ompA* gene has already been replaced with a kanamycin-resistant gene cassette, was transformed with a pKD46 plasmid. The transformed cells were grown in LB broth with ampicillin (50 μg/mL) and 50 mM arabinose at 30^0C until $[OD]_{600nm}$ reached 0.35 to 0.4. Electrocompetent STM *ΔompA* pKD46 cells were made using the protocol mentioned above. Chloramphenicol resistant gene

cassette (Chl^R, 1.1 kb) was amplified from pKD3 plasmid using knockout primers having a

stretch of oligos at the 5' end homologous to the flanking region of the target genes- *ompC*,

ompD, *ompF* (Table- 3.2). The amplified Chl^R gene cassette was subjected to phenol-

chloroform extraction and electroporated into STM *ΔompA* pKD46. The transformed cells

were plated on LB agar with kanamycin (50 μg/mL) and chloramphenicol (25 μg/mL) both for

selection of double knockout strains (*ΔompA ΔompC*, *ΔompA ΔompD*, and *ΔompA ΔompF*).

The plates were further incubated overnight at 37^0C. The knockout colonies were confirmed

by confirmatory primers (Table- 2).

RNA isolation and RT PCR

The bacterial cell pellets were lysed with TRIzol reagent (RNAiso Plus, Takara) and stored at

-80^0C overnight. The lysed supernatants were further subjected to chloroform extraction

followed by precipitation of total RNA by adding an equal volume of isopropanol. The pellet

was washed with 70% RNA-grade ethanol, air-dried, and suspended in 20 μL of DEPC treated

water. RNA concentration was measured in nano-drop and analyzed on 1.5% agarose gel to

assess RNA quality. To make cDNA, 3 μg of RNA sample was subjected to DNase treatment

in the presence of DNase buffer (Thermo Fischer Scientific) at 37^0C for 2h. The reaction was

stopped by adding 5mM Na$_2$EDTA (Thermo Fischer Scientific), followed by heating at 65^0C

for 10 min. The samples were incubated with random hexamer at 65^0C for 10 min and then

supplemented with 5X RT buffer, RT enzyme, dNTPs, and DEPC treated water at 42^0C for 1h.

Quantitative real-time PCR was done using SYBR/ TB Green RT PCR kit (Takara Bio) in Bio-

Rad real-time PCR detection system. The expression level of target genes was measured using

specific RT primers (Table- 2). 16S rRNA was used to normalize the expression levels of the

target genes.

Percent phagocytosis calculation/ invasion assay

1.5 to 2 X 10^5 cells (RAW264.7, U937, Caco-2, and HeLa) were seeded into the wells of 24

well plates and incubated for 6-8 h at 37^0C in the presence of 5% CO_2. As demonstrated earlier,

the protocol for calculating percent phagocytosis by macrophage cells has been modified a

little [63]. Briefly, the phagocytic macrophage cells (RAW264.7 and activated U937 cells)

were infected with 10- 12 h grown stationary phase cultures of STM (WT), *ΔompA, ΔompA*:

pQE60-*ompA* and *ΔompA*: pQE60 at MOI 10. For assays with the complemented strains, the

strains were incubated with 10% mouse complement sera for 2h before the experiment, and

MOI 50 was used for the infection. Non-phagocytic epithelial cells (Caco-2 and HeLa cells)

were infected with the cells from 3 to 4 h old log phase culture of all four bacterial strains at

MOI 10. The attachment of bacteria to the cell surface was increased by centrifuging the cells

at 800 rpm for 5 min, followed by incubating the infected cells at 37^0C in the presence of 5%

CO_2 for 25 min. Next, the cells were washed thrice with PBS to remove unattached bacteria

and subjected to 100 μg/ mL and 25 μg/ mL concentration of gentamycin treatment for 1h each.

2 h post-infection, the cells were lysed with 0.1% triton-X-100. The lysate was plated on

Salmonella- Shigella agar, and the corresponding CFUs were enumerated. Percent

phagocytosis (for phagocytic macrophage cells)/ percent invasion (for non-phagocytic

epithelial cells) was determined using the following formula-

Percent phagocytosis/ percent invasion= [CFU at 2 h]/ [CFU of pre-inoculum] X 100

Adhesion assay

The protocol of adhesion assay was as described before [14]. Briefly, 1.5 to 2 X 10^5 cells

(RAW264.7 and HeLa) were seeded on the top of sterile glass coverslips. Phagocytic

macrophage cells (RAW264.7) were infected with 10- 12 h old stationary phase culture of STM

(WT), *ΔompA, ΔompA*: pQE60-*ompA,* and *ΔompA*:pQE60 at MOI 50. Non-phagocytic

epithelial cells (HeLa) were infected with 3 -4 hours old log phase culture of all four bacterial

strains at the same MOI. After centrifuging the cells at 800 rpm for 5 min, the infected cells

were incubated at 37^0C temperature in the presence of 5% CO_2 for 15 minutes (for RAW264.7

cells) and 25 minutes (for HeLa cells), respectively. After the incubation period, the cells were

washed twice with sterile PBS and fixed with 3.5% PFA. To visualize the externally attached

bacteria, the cells were primarily treated with anti-*Salmonella* antibody raised in rabbit

(dilution 1: 100, duration 6 to 8 hours at 4^0C temperature), which was followed by the treatment

of the cells with secondary antibody conjugated to an appropriate fluorophore (dilution 1: 200,

duration 1 hour at room temperature), dissolved in 2.5% BSA solution without saponin. Images

were obtained by confocal laser scanning microscopy (Zeiss LSM 710) using a 63X oil

immersion objective lens. The number of bacteria adhering per cell was calculated by dividing

the total number of bacteria attached by the total number of host cells in a single microscopic

field. The counting and analysis were done with the help of ZEN Black 2009 software provided

by Zeiss.

Intracellular proliferation assay

The protocol of intracellular proliferation assay has been followed, as demonstrated earlier

[45]. Briefly, the seeded RAW264.7, U937, Caco-2, and HeLa cells (1.5 to 2 X 10^5 cells per

well) were infected with STM (WT), *ΔompA, ΔompA*: pQE60-*ompA* and *ΔompA*:pQE60 at

MOI 10, as mentioned earlier in this study. After centrifuging the cells at 800 rpm for 5 minutes,

the infected cells were incubated at 37^0C temperature in the presence of 5% CO_2 for 25 minutes.

Next, the cells were washed thrice with PBS to remove all the unattached extracellular bacteria

and subjected to 100 μg/ mL concentrations of gentamycin treatment for 1 hour. After that, the

cells were washed thrice with sterile PBS and further incubated with25 μg/ mL concentrations

of gentamycin till the lysis. The cells were lysed with 0.1% triton-X-100 at 2 hours and 16

hours post-infection.   The lysates were plated on *Salmonella- Shigella* Agar, and the

corresponding CFU at 2 hours and 16 hours were determined. The intracellular proliferation of bacteria (Fold proliferation) was determined using a simple formula-

Fold proliferation= [CFU at 16 hours]/ [CFU at 2 hours]

In some sets of experiments, the fold proliferation of STM (WT) and *ΔompA* in the macrophages (RAW 264.7) was measured in the presence of iNOS inhibitor 1400W dihydrochloride [10µM] and activator mouse IFN-γ [100U/ mL]. Both inhibitor (1400W) and activator (IFN-γ) were added to the cells infected with STM (WT) and *ΔompA* along with 25 µg/ mL of gentamycin solution. As usual, the cells were lysed with 0.1% triton-X-100 at 2 hours and 16 hours post-infection. The lysates were plated on *Salmonella- Shigella* Agar, and the corresponding CFU at 2 hours and 16 hours were calculated to determine Fold proliferation. In the intracellular survival assay, two consecutive dilutions were made from each technical replicates at 2 hours and 16 hours. After plating each dilution at 2 hours and 16 hours, the obtained CFU was used to calculate the fold proliferation.

Chloroquine resistance assay

To estimate the number of intracellular bacteria localized in the cytoplasm of macrophages and epithelial cells, a chloroquine resistance assay was performed using a modified protocol as described previously [64, 65]. Briefly, the seeded RAW264.7 and Caco-2 cells (density- 1.5 to 2 X 10^5 cells per well) were infected with STM- (WT), *ΔompA,* and *ΔompA*: pQE60-*ompA* at MOI 10, as mentioned earlier in this study. After centrifuging the cells at 800 rpm for 5 minutes, the infected cells were incubated at 37^0C temperature in the presence of 5% CO_2 for 25 minutes. Next, the cells were washed with PBS and subjected to 100 µg/ mL and 25 µg/ mL of gentamycin treatment, respectively. The old DMEM (having 25 µg/ mL gentamycin) was replaced from the wells with freshly prepared DMEM, supplemented with 25 µg/ mL gentamycin and 50 µg/ mL chloroquine two hours before lysis (14 hours post-infection). The

cells were lysed with 0.1% triton-X-100 at 16 hours post-infection. The lysates were plated on

Salmonella- Shigella agar, and the corresponding CFU at 16 hours was determined. Percent

abundance of cytosolic and vacuolar bacteria was obtained after dividing the CFU from

chloroquine treated set with chloroquine untreated set.

Percentage of cytosolic bacteria= [CFU at 16 hours with chloroquine]/ [CFU at 16 hours

without chloroquine] X 100%

Percentage of vacuolar bacteria= (100 - percentage of cytosolic bacteria) %

Confocal microscopy

For immunofluorescence study, RAW 264.7 or Caco-2 cells seeded at a density of 1.5 to 2 X

10^5 cells per sterile glass coverslip were infected with appropriate bacterial strains at MOI 20.

The cells were washed thrice with PBS and fixed with 3.5% paraformaldehyde for 15minutes

at indicated time points post-infection. The cells were first incubated with specific primary

antibody raised against *Salmonella* SseC/ SseD proteins, mouse lysosome-associated

membrane protein-1 (LAMP-1) (rat anti-mouse LAMP-1), mouse nitrotyrosine (mouse anti-

mouse nitrotyrosine), and S. Typhimurium (anti- *Salmonella* O antigen) as per the requirements

of experiments, diluted in 2.5% BSA and 0.01% saponin (dilution 1: 100, duration 6 to 8 hours

at 4^0C temperature). This was followed by incubating the cells with appropriate secondary

antibodies conjugated with fluorophores (dilution 1: 200, duration 1 hours at room

temperature).

The coverslips were mounted with anti-fade reagent and fixed on a glass slide with transparent

nail paint. Samples were imaged by confocal laser scanning microscopy (Zeiss LSM 710) using

a 63X oil immersion objective lens. The images were analyzed with ZEN Black 2009 software

provided by Zeiss.

Griess assay to measure extracellular nitrite concentration

Extracellular nitrite from infected macrophage cells was measured using a protocol described

earlier [66]. 3.13 μM, 6.25 μM, 12.5 μM, 25 μM, 50 μM, 100 μM standard $NaNO_2$ solutions

were prepared from main stock [0.1 (M) $NaNO_2$] by serial dilution in deionized distilled water.

The [OD]$_{545nm}$ of the standard solutions were measured after adding and incubating them with

Griess reagent, and the standard curve was drawn. Culture supernatants were collected from

RAW264.7 cells infected with STM- (WT), *ΔompA, ΔompA*: pQE60-*ompA*, *ΔompA*: pQE60,

and heat-killed bacteria at 16 hours post-infection and subjected to nitrite estimation by adding

Griess reagents. To 50 μL of culture supernatant, 50 μL of 1% sulphanilamide (made in 5%

phosphoric acid), and 50 μL of 0.1% NED (N-1-naphthyl ethylene diamine dihydrochloride)

were added and incubated for 10 minutes in darkness at room temperature. The [OD]$_{545nm}$ was

measured within 30 minutes of the appearance of a purple-colored product.

Measurement of intracellular nitric oxide

The level of intracellular nitric oxide of infected macrophages was measured using cell

membrane-permeable fluorescent nitric oxide probe 4, 5- diaminofluorescein diacetate (DAF2-

DA) [67]. Briefly, RAW264.7 cells were infected with STM- (WT), *ΔompA, ΔompA*: pQE60-

ompA, *ΔompA ΔompC*, *ΔompA ΔompD*, *ΔompA ΔompF* & LLO at MOI 10 as described before.

16 hours post-infection, the culture supernatants were replaced with DMEM media,

supplemented with 5μM concentration of DAF2-DA, followed by further incubation of the

infected cells at 370C temperature in the presence of 5% CO_2 for 30 minutes. The cells were

washed with sterile PBS and acquired immediately for analysis by flow cytometry (BD

FACSVerse by BD Biosciences-US) using a 491 nm excitation channel and 513 nm emission

channel.

Measurement of the activity of the *spiC* promoter

The activity of *spiC* promoter in STM (WT) and *ΔompA* was measured by mild alteration of a protocol described earlier [66]. Briefly, 1.5 mL of overnight grown stationary phase culture of STM (WT) and *ΔompA* carrying pHG86 *spiC-lacZ* construct were centrifuged at 6000 rpm for 10 minutes, and the pellet was resuspended in 500 µL of Z-buffer (Na2HPO4, 60 mM; NaH2PO4, 40 mM; KCl, 10 mM; MgSO4.7H2O, 1mM). The OD of the Z-buffer was measured at 600 nm after resuspension. The cells were permeabilized by adding 5 µL of 0.1% SDS and 20 µL of chloroform and incubated at room temperature for 5 minutes. 100 µL of 4 mg/ mL of o-nitrophenyl *β*-D galactopyranoside was added in the dark and incubated till the color appeared. The reaction was stopped using 250 µL 1 M Na_2CO_3. The reaction mixture was centrifuged at 6000 rpm for 10 minutes, and the OD of the supernatant was measured at 420 nm and 550 nm on flat bottom transparent 96 well plates. STM (WT) and *ΔompA* harboring promoter-less empty pHG86 *LacZ* construct were used as control. The activity of the *spiC* promoter was measured in Miller Unit using the following formula

$$\textbf{Miller Unit (MU)= } 1000[\textbf{ OD}_{420nm} - \textbf{OD}_{550nm} * 1.75]/ \textbf{ T*V*OD}_{600nm}$$

OD_{420nm}= Absorbance by o-nitrophenol and light scattering by cell debris

OD_{550nm}= light scattering by cell debris

OD_{600nm} = bacterial cell density in the washed media

T= Time of reaction in minutes

V= Volume of the culture in mL.

Measurement of the activity of the *spiC* promoter from intracellular bacteria

RAW264.7 cells were infected with STM (WT): pHG86 *spiC-LacZ* and *ΔompA*: pHG86 *spiC-LacZ* at MOI 50. After centrifuging the cells at 800 rpm for 5 minutes, the infected cells were incubated at 37^0C temperature in the presence of 5% CO_2 for 25 minutes. Next, the cells were

washed thrice with PBS to remove all the unattached extracellular bacteria and subjected to

100 µg/ mL and 25 µg/ mL concentrations of gentamycin treatment for 1 hour each. The cells

were lysed with 0.1% triton-X-100 at 12 hours post-infection. The lysates were centrifuged at

14,000 rpm for 30 minutes, and the pellet was resuspended in 500 µL of Z-buffer. The OD of

the Z-buffer was measured at 600 nm after resuspension. The cells were permeabilized by

adding 5 µL of 0.1% SDS and 20 µL of chloroform and incubated at room temperature for 5

minutes. 100 µL of 4 mg/ mL of o-nitrophenyl β-D galactopyranoside was added in the dark

and incubated till the color appeared. The reaction was stopped using 250 µL 1 M Na_2CO_3.

The reaction mixture was centrifuged at 6000 rpm for 10 minutes, and the OD of the

supernatant was measured at 420 nm and 550 nm on flat bottom transparent 96 well plates. The

activity of the *spiC* promoter was measured in Miller Unit using the formula mentioned earlier.

Measurement of the cytosolic acidification of the bacteria using BCECF-AM

The acidification of the cytosol of the bacteria in the presence of *in vitro* acidic stress was

measured using a cell-permeable dual excitation ratiometric dye called 2',7'-Bis-(2-

Carboxyethyl)-5-(and-6)-Carboxyfluorescein, Acetoxymethyl Ester (BCECF-AM). 4.5×10^7

CFU of STM (WT), *ΔompA,* and *ΔompA*: pQE60-*ompA* from 12 hours old overnight grown

stationary phase culture was resuspended in phosphate buffer (of pH 5.5, 6, 6,5, and 7

respectively) separate microcentrifuge tubes and incubated for 2 hours in a shaker incubator at

37^0C temperature. 30 minutes before the flow cytometry analysis, BCECF-AM (1mg/ mL) was

added to each tube to make the final concentration 20 µM. At the end of the incubation period

of 2 hours, the bacterial cells were analyzed in flow cytometry (BD FACSVerse by BD

Biosciences-US) using 405 nm and 488 nm excitation and 535 nm emission channel. The

median fluorescence intensity (MFI) of the bacterial population at 488 nm and 405 nm was

obtained from BD FACSuite software. The 488/ 405 ratio was determined to estimate the level

of acidification of the bacterial cytosol.

Measurement of extracellular H_2O_2 by phenol red assay

H_2O_2 produced by RAW264.7 cells infected with STM- (WT), $\Delta ompA$, $\Delta ompA$: pQE60-*ompA* at MOI 10 was measured by modifying a protocol of phenol red assay as demonstrated before [68]. Briefly, two hours post-infection, infected RAW264.7 cells were supplemented with phenol red solution having potassium phosphate (0.01 M; pH 7.0), glucose (0.0055 M), NaCl (0.14 M), phenol red (0.2 g/ L), and HRPO (8.5 U/ mL; Sigma- Aldrich). 16 hours post-infection, the culture supernatant was collected and subjected to the [OD] measurement at 610 nm wavelength in TECAN 96 well microplate reader. In the presence of H_2O_2 produced by macrophages, horseradish peroxidase (HRPO) converts phenol red into a compound that has enhanced absorbance at 610 nm. The concentration of H_2O_2 produced by macrophages was measured using a standard curve of H_2O_2 in phenol red solution with known concentrations ranging from 0.5 to 5 µM.

Measurement of intracellular ROS

The level of intracellular ROS in infected macrophages was measured using membrane-permeable redox-sensitive probe 2',7'- dichlorodihydrofluorescein diacetate (H_2DCFDA) [68]. Upon its oxidation by intracellular esterases, this non-fluorescent dye is converted into highly fluorescent 2',7'- Dichlorofluorescein (H_2DCF), which has emission at 492- 495 nm and excitation at 517 to 527 nm. Briefly, RAW264.7 cells were infected with STM- (WT), $\Delta ompA$, $\Delta ompA$: pQE60-*ompA* at MOI 10 as described before. 16 hours post-infection, the culture supernatants were replaced with DMEM media, supplemented with 10 µM concentration of H_2DCFDA, followed by further incubation of the infected cells at 370C temperature in the presence of 5% CO_2 for 30 minutes. The cells were washed with sterile PBS and acquired immediately for analysis by flow cytometry (BD FACSVerse by BD Biosciences-US) using a 492 nm excitation channel and 517 nm emission channel.

Sensitivity assay of bacteria against *in vitro* nitrosative and oxidative stress

The sensitivity of STM (WT) and *ΔompA* was tested against *in vitro* nitrosative and oxidative stress. H_2O_2 dissolved in PBS of pH 5.4 was used for creating *in vitro* oxidative stress. Acidified nitrite ($NaNO_2$ in PBS of pH 5.4) alone and a combination of acidified nitrite and H_2O_2 were used to generate *in vitro* nitrosative stress [32]. Sensitivity was checked in both concentration and time-dependent manner.

Concentration-dependent sensitivity

10^8 CFU of overnight grown stationary phase cultures of STM- (WT) and *ΔompA* were inoculated in varying concentrations of acidified nitrite and peroxide ranging from 200 µM to 5 mM and further incubated for 12 hours. At the end of the incubation period, supernatants from each concentration were collected, serially diluted, plated on *Salmonella- Shigella* agar and the log_{10}[CFU/ mL] values were acquired to determine the inhibitory concentrations of nitrite, peroxide, and both together.

Time-dependent sensitivity

10^8 CFU of overnight grown stationary phase cultures of STM- (WT) and *ΔompA* were inoculated in 800µM concentration of acidified nitrite and further incubated for 12 hours. Aliquots were collected, serially diluted, and plated on *Salmonella- Shigella* agar at 0, 3, 6, 9, 12 hours post-inoculation to monitor the CFU.

Bacterial cell viability assay by resazurin

Bacterial cell viability under the treatment of acidified nitrite and peroxide was measured using resazurin assay. Resazurin, a blue-colored non-fluorescent redox indicator, is reduced into resorufin, a pink-colored fluorescent compound (having excitation at 540 nm and emission at 590 nm) by aerobic respiration of metabolically active cells. Briefly, STM- (WT) and *ΔompA*

were treated with varying concentrations of acidified nitrite and peroxide, as mentioned above.

At the end of the incubation period, supernatants were collected and incubated with resazurin

(1 µg/ mL) in a 37^0C shaker incubator at 180 rpm for 2 hours in a 96 well plate. At the end of

the incubation period, the fluorescence intensity was measured using TECAN 96 well

microplate reader, and percent viability was calculated.

Nitrite uptake assay

Nitrite uptake by different bacterial strains [STM- (WT), $\Delta ompA$, $\Delta ompA$: pQE60-$ompA$,

$\Delta ompA$:pQE60, $\Delta ompA\Delta ompC$, $\Delta ompA\Delta ompD$, $\Delta ompA\Delta ompF$, $\Delta ompC$, $\Delta ompD$, $\Delta ompF,$ and

PFA fixed dead bacteria] was determined by measuring the remaining concentration of nitrite

in the uptake mixture using a protocol described earlier [66]. Briefly, 10^8 CFU of overnight

grown stationary phase bacterial cultures were inoculated in an uptake mixture consisting of

40 mM glucose, 80 mM MOPS-NaOH buffer (pH= 8.5), and nitrite (50/ 100/ 200 µM) in a

final volume of 5 mL. The assay mixtures were kept in a 37^0C shaker incubator after the

inoculation was done. At indicated time points, 150 µL of suspension from each assay mixture

was collected and subjected to Griess assay to determine the level of remaining nitrite, as

mentioned earlier.

Examination of *in vitro* redox homeostasis of STM (WT) and *$\Delta ompA$* in response to

acidified nitrite

Stationary phase cultures of STM- (WT) and $\Delta ompA$ harboring pQE60-Grx1-roGFP2 plasmid

were sub-cultured in freshly prepared 5 mL LB broth at 1: 33 ratios in the presence of

appropriate antibiotic in 37^0C shaker incubator at 175 rpm. Once the $[OD]_{600\ nm}$ has reached

0.3 to 0.4, 500 µM of IPTG (Sigma-Aldrich) was added, and the cells were further grown at

30^0C temperature at 175 rpm for 10 to 12 hours. At the end of the incubation period, 4.5 X 10^7

CFU of bacteria were subjected to the treatment of acidified nitrite [varying concentrations (as

mentioned in the figure legend) of $NaNO_2$ in PBS of pH= 5.4] for 15, 30, 45, and 60 minutes.

At the end of every indicated time point, the cells were analyzed in flow cytometry (BD

FACSVerse by BD Biosciences-US) using 405 nm and 488 nm excitation and 510 nm emission

channel. The mean fluorescence intensity at 405 nm and 488 nm was obtained from the FITC

positive (GFP expressing) population, and the 405/ 488 ratio was determined.

Determination of outer membrane porosity of intracellular and extracellular bacteria by

bisbenzimide

The outer membrane porosity of STM- (WT), $\Delta ompA$, $\Delta ompA$: pQE60-*ompA*, and

$\Delta ompA$:pQE60 grown in low magnesium acidic N s medium (pH= 5.4) was measured using

bisbenzimide (Sigma-Aldrich) by modifying a protocol as specified previously [69]. The

bacterial strains were grown in low magnesium acidic F medium for 12 hours. At the end of

the incubation period, the culture supernatants were collected, and the $[OD]_{600\,nm}$ was adjusted

to 0.1 with sterile PBS. 20 µL of bisbenzimide (10 µg/ mL) solution was added to 180 µL of

culture supernatants (whose $[OD]_{600\,nm}$ has already been adjusted to 0.1) in 96 well a microplate

and further incubated for 10 minutes in 37^0C shaker incubator. Because of enhanced outer

membrane porosity, when bisbenzimide is taken up by the bacterial cells, it binds to the

bacterial DNA and starts fluorescing. The fluorescence intensity of DNA bound bisbenzimide

was measured in TECAN 96 well microplate reader using 346 nm excitation and 460 nm

emission filter.

To check the outer membrane porosity of intracellular STM- (WT) and $\Delta ompA$, infected

RAW264.7 macrophage cells were lysed with 0.1% Triton X-100. The lysate was collected

and centrifuged at 300g for 5 minutes to settle down eukaryotic cell debris. The sup was

collected and further centrifuged at 5000 rpm for 20 minutes to settle down the bacteria. This

was followed by decanting the sup and resuspending the bacterial pellets with PBS. Finally,

the suspension was subjected to bisbenzimide treatment to measure the fluorescence intensity

as mentioned above.

Determination of bacterial membrane depolarization using DiBAC$_4$

Outer membrane depolarization of STM (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*,

ΔompA:pQE60, STM (WT): pQE60, STM (WT): pQE60-*ompA*, STM (WT): pQE60-*ompC*,

STM (WT): pQE60-*ompD*, and STM (WT): pQE60-*ompF* grown in low magnesium acidic F

medium (pH= 5.4) for 12 hours was measured using a fluorescent membrane potential sensitive

dye called bis-(1,3-dibutyl barbituric acid)-trimethylene oxonol (Invitrogen). Briefly, 4.5×10^7

CFU of each bacterial strain was incubated with 1 µg/ml of DiBAC$_4$ for 15 minutes in a 37^0C

shaker incubator. The DiBAC$_4$ treated bacterial cells were further analyzed by flow cytometry

(BD FACSVerse by BD Biosciences-US) to evaluate the change in membrane depolarization

upon knocking out *ompA*.

Stationary phase cultures of STM (WT), STM (WT): pQE60, STM (WT): pQE60-*ompA*, STM

(WT): pQE60-*ompC*, STM (WT): pQE60-*ompD*, and STM (WT): pQE60-*ompF* were sub-

cultured in freshly prepared 5 mL LB broth at 1: 33 ratios in the presence of appropriate

antibiotics in 37^0C shaker incubator at 175 rpm. Once the [OD]$_{600 \text{ nm}}$ has reached 0.3 to 0.4,

500 µM of IPTG (Sigma-Aldrich) was added, and the cells were further grown at 30^0C

temperature at 175 rpm for 10 to 12 hours. At the end of the incubation period, 4.5×10^7 CFU

of bacteria were subjected to the treatment of 1 µg/mL of DiBAC$_4$ for 15 minutes in a 37^0C

shaker incubator. At the end of the incubation period, the cells were analyzed in flow cytometry

(BD FACSVerse by BD Biosciences-US) to measure the outer membrane depolarization.

Expression profiling of *ompC*, *ompD*, *ompF* in STM- (WT), *ΔompA* and complement

strains growing in LB broth, acidic F media, and macrophages

Overnight grown stationary phase cultures of STM- (WT), $\Delta ompA$, & $\Delta ompA$: pQE60-$ompA$

were inoculated in freshly prepared LB broth, low magnesium acidic F media (pH= 5.4) at a

ratio of 1: 100. The cells were further grown in a 37^0C shaker incubator at 180 rpm for 12

hours. RAW264.7 cells were infected with above mentioned bacterial strains at MOI 50 and

incubated further for 12 hours, as mentioned earlier. At the end of the specified incubation,

period RNA was isolated, cDNA was synthesized, and the expression of $ompC, ompD, ompF$

were checked, as mentioned earlier.

Live dead assay by propidium iodide

10^8 CFU of overnight grown stationary phase cultures of STM (WT): pQE60, STM (WT):

pQE60-$ompA$, STM (WT): pQE60-$ompC$, STM (WT): pQE60-$ompD$, STM (WT): pQE60-

$ompF$ were inoculated in 1 mM concentration of acidified nitrite (total volume 10mL) and

further incubated for 12 hours. 300 µL of aliquots (corresponding to 10^5 to 10^6 CFU of bacteria)

were collected 12 hours post-inoculation and subjected to the treatment with propidium iodide

(PI) (Sigma-Aldrich) (concentration- 1µg/ mL) for 30 minutes at 37^0C temperature. After the

incubation, the PI-treated bacterial samples were analyzed by flow cytometry (BD FACSVerse

by BD Biosciences-US) to estimate the percent viability.

Animal survival assay

4-6 weeks old BALB/c and C57BL/6 mice housed in the specific-pathogen-free condition of

central animal facility of Indian Institute of Science, Bangalore was used for all the *in vivo*

infection and survival studies. The Institutional Animal Ethics Committee approved all the

animal experiments, and the National Animal Care Guidelines were strictly followed. Two

cohorts of twenty 4-6 weeks old BALB/c and C57BL/6 mice were infected with 10-12 hours

old overnight grown stationary phase cultures of STM (WT) and $\Delta ompA$ by oral gavaging at

lethal dose 10^8 CFU/ animal respectively **(n= 10)**. The survival of infected mice was observed

for the next few days until all the mice infected with STM (WT) died. The survival was

recorded, and the data was represented as percent survival.

Determination of bacterial burden in different organs

Four cohorts of five 4-6 weeks old C57BL/6 mice were infected with STM (WT) and *ΔompA*

by oral gavaging at sub-lethal dose 10^7 CFU/ animal, respectively (**n= 5**). Two of these cohorts

infected with STM- (WT) and *ΔompA* strains respectively were further intraperitoneally

injected with iNOS inhibitor aminoguanidine hydrochloride (AGH- 10mg/ kg of body weight)

regularly for five days post-infection. The other two cohorts were treated with a placebo. Two

cohorts of *iNOS$^{-/-}$* mice were orally infected with STM (WT) and *ΔompA* at 10^7 CFU/ animal

(**n=5**). Two cohorts of five *gp91phox$^{-/-}$* mice unable to generate ROS were orally gavaged with

STM- (WT) and *ΔompA* at 10^7 CFU/ animal (**n=5**). On the 5th day post-infection, all the mice

were sacrificed, followed by isolation, weighing, and homogenization of specific organs like-

liver, spleen, and MLN. The organ lysates were plated on *Salmonella Shigella* agar to

determine the bacterial burden in different organs. The CFU corresponds to an organ was

normalized with organ weight and the $\log_{10}$[CFU/ gm-wt.] value has been plotted.

Statistical analysis

Each assay has been independently repeated 2 to 5 times [as mentioned in the figure legends].

The *in vitro* data and the results obtained from cell line experiments were analyzed by unpaired

student's *t*-test, and *p* values below 0.05 were considered significant. Results received from *in*

vitro sensitivity assays were analyzed by 2way ANOVA. Data obtained from *in vivo* infection

of mice were analyzed by Mann- Whitney *U* test from GraphPad Prism 8.4.3 (686) software.

Flow cytometry data were analyzed and plotted using BD FACSuite (by BD Biosciences-US)

and CytoFLEX (by Beckman Coulter Life Sciences) software. The results are expressed as

mean $\pm$ SD or mean $\pm$ SEM. Differences between experimental groups were considered

significant for $p < 0.05$.

1208

1209

1210 **Table 1. Strains and plasmids used in this study**

Strains/ plasmids	Characteristics	Source/ references
Salmonella enterica serovar Typhimurium ATCC strain14028S	Wild type (WT)	Gifted by Prof. M. Hensel
S. Typhimurium *ΔompA*	Kan^R	This study
ΔompA: pQE60-*ompA*	Kan^R, Amp^R	This study
ΔompA: pQE60	Kan^R, Amp^R	This study
S. Typhimurium *ΔompC*	Chl^R	This study
S. Typhimurium *ΔompD*	Chl^R	This study
S. Typhimurium *ΔompD*	Kan^R	This study
S. Typhimurium *ΔompF*	Chl^R	This study
S. Typhimurium *ΔompA ΔompC*	Kan^R, Chl^R	This study

S. Typhimurium *ΔompA ΔompD*	Kan^R, Chl^R	This study
S. Typhimurium *ΔompA ΔompF*	Kan^R, Chl^R	This study
pKD4	Plasmid with FRT-flanked Kanamycin resistance gene	KA Datsenko & BL Wanner, PNAS, 2000
pKD46	Plasmid expressing λ red recombinase, Amp^R	KA Datsenko & BL Wanner, PNAS, 2000
pQE60 vector	Low copy number plasmid, Amp^R	Laboratory stock
pFV- mCherry (RFP)	Amp^R	Laboratory stock
pFV: GFP	Amp^R	Laboratory stock
STM (WT): *LLO*	Amp^R	Laboratory stock
STM *ΔompA*: *LLO*	Amp^R	This study
S. Typhimurium wild type: pHG86 *spiC-LacZ*	Amp^R	This study
S. Typhimurium *ΔompA*: pHG86 *spiC-LacZ*	Amp^R	This study
S. Typhimurium wild-type: pQE60-*ompA*	Amp^R	This study

S. Typhimurium wild-type: pQE60-*ompC*	Amp^R	This study
S. Typhimurium wild-type: pQE60-*ompD*	Amp^R	This study
S. Typhimurium wild-type: pQE60-*ompF*	Amp^R	This study
S. Typhimurium wild-type: pQE60	Amp^R	This study
pHG86 *spiC-LacZ*	Amp^R	Laboratory stock
pQE60-Grx1-roGFP2	Amp^R	Gifted by Dr. Amit Singh, CIDR, IISc

1211

Table 2. Primer sequences (5' to 3')

ompA knockout forward-

TCGTTGGAGATATTCATGGCGTATTTTGGATGATAACGAGCATATGAATATCCTC

CTTAG

ompA knockout reverse-

AAGAAGTAACGCTGAAAGGCGTTGTCATCCAGACCAGAGCGTGTAGGCTGGAGC

TGCTTC

ompC knockout forward-

ATAACTGTAACATCTTAAAAGTTTTAGTATCATATTCGTGGTGTAGGCTGGAGCT

GCTTC

ompC knockout reverse-

TATCAAAACGTCGTATTTGTACGCCGGAATAAGGCATGATGGGAATTAGCCATG

GTCC

ompD knockout forward-

TTATTAAAATGAAACTTAAGTTAGTGGCAGTGGCAGTGTTTAAATGGCGCGCCTT

ACG

ompD knockout reverse-

CAAAATTAGAACTGGTAGTTCAGACCAACAGCAACGATGTGGAAGATCACTTCG

CAGAA

ompF knockout forward-

ATTGACGGAATTTATTGACGGCAGTGGCAGGTGTCATAGTGTAGGCTGGAGCTGC

TTC

ompF knockout reverse-

TACAAAATGCCAACCGTTAGCGCTAAAAAGCCCGCCTGTTATGGGAATTAGCCA

TGGTCC

ompA cloning forward- CATG**CCATGG**ATGAAAAAGACAGCTATCGC

ompA cloning reverse- CCC**AAGCTT**TTGTCATCCAGACCAGAG

ompC cloning forward- CGC**GGATCC**ATGAAAGTTAAAGTACTGTCC

ompC cloning reverse- CCC**AAGCTT**GCTGATTAGAACTGGTAAACC

ompD cloning forward- CGC**GGATCC**ATGAAACTTAAGTTAGTGGC

ompD cloning reverse- CCC**AAGCTT**CTACAACAAAATTAGAACTGG

ompF cloning forward- CGC**GGATCC**ATGATGAAGCGCAAAATCC

ompF cloning reverse- CCC**AAGCTT**TCAGAACTGGTAAGTAATACC

ompA confirmatory forward- CGGTAGAGTAACTATTGAG

ompA confirmatory reverse- TTACAGGCGTTATTAGGC

ompA expression forward- ATCCAATCACTGACGATCTG

ompA expression reverse- GCATCACCGATGTTGTTAGT

ompC confirmatory forward- GGTAAACAGACATTCAGA

ompC confirmatory reverse- AGTCATTTTCATCGCTGTT

ompD confirmatory forward- GAACTTATGCCACTCCGTCATT

ompD confirmatory reverse- CAGCATTTCGACGTCAACGGTA

ompF confirmatory forward- GTCAGACACATAAAGACACC

ompF confirmatory reverse- CGAGGTTCCATTATAGTTACAG

Kanamycin[R] internal forward- CGGTGCCCTGAATGAACTGC

Kanamycin[R] internal reverse- CGGCCACAGTCGATGAATCC

Chloramphenicol[R] internal forward- ACAAACGGCATGATGAACCT

Chloramphenicol[R] internal reverse- GCTCTGGAGTGAATACCACG

spiC expression forward- ACCTAAGCCTTGTCTTGCCT

spiC expression reverse- CCATCCGCTGTGAGCTGTAT

sseC expression forward- TTTGGCGAGGAAGTGGTTGA

sseC expression reverse- AGCCATTTCACGTTCAAGCG

sseD expression forward- TGTTGTCGGGTGTACTGACG

sseD expression reverse- ACGGCTTGACCCGCTATAAG

sifA expression forward- CCACACGAGAGCGGCTTACA

sifA expression reverse- GCCGTCATTTGTGGATGCGA

ssaV expression forward- CGCCGCAAAAAGTCTGTGGT

ssaV expression reverse- GGGACGCCGGTATCCTCAAA

16srRNA forward- GAGCGCAACCCTTATCCTTTG

16srRNA forward- CACTTTATGAGGTCCGCTTGCT

Acknowledgments

Departmental Confocal Facility, Departmental Real-Time PCR Facility, Central Bioimaging

Facility, Central Flowcytometry Facility, Department of Biochemistry Flowcytometry Facility,

and Central Animal Facility at IISc are duly acknowledged. Ms. Anusha and Ms. Navya are

acknowledged for their help in image acquisition. Mrs. Ranjitha is acknowledged for her help

in flow cytometry data acquisition. Dr. Amit Singh (Centre for Infectious Disease Research

and Department of Microbiology and Cell Biology, Indian Institute of Science, Bangalore) is

acknowledged for providing pQE60-Grx1-roGFP2 construct. Ms. Debapriya Mukherjee and

Ms. Dipasree Hajra are recognized for their timely technical support.

Funding

This work was supported by the DAE SRC fellowship (DAE00195) and DBT-IISc partnership umbrella program for advanced research in biological sciences and Bioengineering to DC Infrastructure support from ICMR (Centre for Advanced Study in Molecular Medicine), DST (FIST), and UGC (special assistance) is acknowledged. DC acknowledges the ASTRA Chair professorship grant from IISc. ARC sincerely acknowledges IISc Fellowship from MHRD, Govt. of India, and the estate of the late Dr. Krishna S. Kaikini for the Shamrao M. Kaikini and Krishna S. Kaikini scholarship.

Availability of data and materials

All data generated and analyzed during this study, including the supplementary information files, have been incorporated in this article. The data is available from the corresponding author on reasonable request.

Author Contributions

ARC and DC conceived the study and designed the experiments. ARC performed all the experiments, analyzed the data, and wrote the original draft of the manuscript. SS constructed pQE60-*ompA* recombinant plasmid under the supervision of UV. SS, UV, and DC reviewed and edited the manuscript. DC supervised the study. All the authors have read and approved the manuscript.

Declarations

Ethics statement

All the animal experiments were approved by the Institutional Animal Ethics Committee, and the Guidelines provided by National Animal Care were strictly followed. (Registration No: 48/1999/CPCSEA).

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no conflict of interest.

References

1.   Vergalli, J., et al., *Porins and small-molecule translocation across the outer membrane of*
Gram-negative bacteria. Nat Rev Microbiol, 2020. **18**(3): p. 164-176.
2.   March, C., et al., *Klebsiella pneumoniae outer membrane protein A is required to prevent the*
activation of airway epithelial cells. J Biol Chem, 2011. **286**(12): p. 9956-67.
3.   Hejair, H.M.A., et al., *Functional role of ompF and ompC porins in pathogenesis of avian*
pathogenic Escherichia coli. Microb Pathog, 2017. **107**: p. 29-37.
4.   Krishnan, S. and N.V. Prasadarao, *Outer membrane protein A and OprF: versatile roles in Gram-*
negative bacterial infections. FEBS J, 2012. **279**(6): p. 919-31.
5.   Wu, X.B., et al., *Outer membrane protein OmpW of Escherichia coli is required for resistance*
to phagocytosis. Res Microbiol, 2013. **164**(8): p. 848-55.
6.   Li, W., et al., *Contribution of the outer membrane protein OmpW in Escherichia coli to*
complement resistance from binding to factor H. Microb Pathog, 2016. **98**: p. 57-62.
7.   Park, J.S., et al., *Mechanism of anchoring of OmpA protein to the cell wall peptidoglycan of the*
gram-negative bacterial outer membrane. FASEB J, 2012. **26**(1): p. 219-28.
8.   Ipinza, F., et al., *Participation of the Salmonella OmpD porin in the infection of RAW264.7*
macrophages and BALB/c mice. PLoS One, 2014. **9**(10): p. e111062.
9.   van der Heijden, J., et al., *Salmonella Rapidly Regulates Membrane Permeability To Survive*
Oxidative Stress. mBio, 2016. **7**(4).
10. Eriksson, S., et al., *Unravelling the biology of macrophage infection by gene expression*
profiling of intracellular Salmonella enterica. Mol Microbiol, 2003. **47**(1): p. 103-18.
11. Hautefort, I., et al., *During infection of epithelial cells Salmonella enterica serovar*
Typhimurium undergoes a time-dependent transcriptional adaptation that results in
simultaneous expression of three type 3 secretion systems. Cell Microbiol, 2008. **10**(4): p. 958-
84.
12. Deiwick, J., et al., *Environmental regulation of Salmonella pathogenicity island 2 gene*
expression. Mol Microbiol, 1999. **31**(6): p. 1759-73.
13. Datsenko, K.A. and B.L. Wanner, *One-step inactivation of chromosomal genes in Escherichia*
coli K-12 using PCR products. Proc Natl Acad Sci U S A, 2000. **97**(12): p. 6640-5.
14. Garai, P., et al., *Peptide utilizing carbon starvation gene yjiY is required for flagella mediated*
infection caused by Salmonella. Microbiology (Reading), 2016. **162**(1): p. 100-116.
15. Garai, P., D.P. Gnanadhas, and D. Chakravortty, *Salmonella enterica serovars Typhimurium*
and Typhi as model organisms: revealing paradigm of host-pathogen interactions. Virulence,
2012. **3**(4): p. 377-88.
16. Knuff, K. and B.B. Finlay, *What the SIF Is Happening-The Role of Intracellular Salmonella-*
Induced Filaments. Front Cell Infect Microbiol, 2017. **7**: p. 335.
17. Chakraborty, S., H. Mizusaki, and L.J. Kenney, *A FRET-based DNA biosensor tracks OmpR-*
dependent acidification of Salmonella during macrophage infection. PLoS Biol, 2015. **13**(4): p.
e1002116.

18. Canton, J., et al., *Contrasting phagosome pH regulation and maturation in human M1 and M2*
macrophages. Mol Biol Cell, 2014. **25**(21): p. 3330-41.
19. Martin-Orozco, N., et al., *Visualization of vacuolar acidification-induced transcription of genes*
of pathogens inside macrophages. Mol Biol Cell, 2006. **17**(1): p. 498-510.
20. Choi, J. and E.A. Groisman, *Acidic pH sensing in the bacterial cytoplasm is required for*
Salmonella virulence. Mol Microbiol, 2016. **101**(6): p. 1024-38.
21. Chakravortty, D., et al., *Formation of a novel surface structure encoded by Salmonella*
Pathogenicity Island 2. EMBO J, 2005. **24**(11): p. 2043-52.
22. Holzer, S.U. and M. Hensel, *Functional dissection of translocon proteins of the Salmonella*
pathogenicity island 2-encoded type III secretion system. BMC Microbiol, 2010. **10**: p. 104.
23. Foster, N., S.D. Hulme, and P.A. Barrow, *Induction of antimicrobial pathways during early-*
phase immune response to Salmonella spp. in murine macrophages: gamma interferon (IFN-
gamma) and upregulation of IFN-gamma receptor alpha expression are required for NADPH
phagocytic oxidase gp91-stimulated oxidative burst and control of virulent Salmonella spp.
Infect Immun, 2003. **71**(8): p. 4733-41.
24. Umezawa, K., et al., *Induction of nitric oxide synthesis and xanthine oxidase and their roles in*
the antimicrobial mechanism against Salmonella typhimurium infection in mice. Infect Immun,
1997. **65**(7): p. 2932-40.
25. van der Heijden, J., et al., *Direct measurement of oxidative and nitrosative stress dynamics in*
Salmonella inside macrophages. Proc Natl Acad Sci U S A, 2015. **112**(2): p. 560-5.
26. Beuzon, C.R., S.P. Salcedo, and D.W. Holden, *Growth and killing of a Salmonella enterica*
serovar Typhimurium sifA mutant strain in the cytosol of different host cell lines. Microbiology
(Reading), 2002. **148**(Pt 9): p. 2705-2715.
27. Chakravortty, D., I. Hansen-Wester, and M. Hensel, *Salmonella pathogenicity island 2*
mediates protection of intracellular Salmonella from reactive nitrogen intermediates. J Exp
Med, 2002. **195**(9): p. 1155-66.
28. Ruan, Y., et al., *Listeriolysin O Membrane Damaging Activity Involves Arc Formation and*
Lineaction -- Implication for Listeria monocytogenes Escape from Phagocytic Vacuole. PLoS
Pathog, 2016. **12**(4): p. e1005597.
29. Uchiya, K.I. and T. Nikai, *Salmonella virulence factor SpiC is involved in expression of flagellin*
protein and mediates activation of the signal transduction pathways in macrophages.
Microbiology (Reading), 2008. **154**(Pt 11): p. 3491-3502.
30. Uchiya, K. and T. Nikai, *Salmonella pathogenicity island 2-dependent expression of suppressor*
of cytokine signaling 3 in macrophages. Infect Immun, 2005. **73**(9): p. 5587-94.
31. Pearse, D.D., et al., *Comparison of iNOS inhibition by antisense and pharmacological inhibitors*
after spinal cord injury. J Neuropathol Exp Neurol, 2003. **62**(11): p. 1096-107.
32. Heaselgrave, W., P.W. Andrew, and S. Kilvington, *Acidified nitrite enhances hydrogen peroxide*
disinfection of Acanthamoeba, bacteria and fungi. J Antimicrob Chemother, 2010. **65**(6): p.
1207-14.
33. Fang, F.C., *Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related*
antimicrobial activity. J Clin Invest, 1997. **99**(12): p. 2818-25.
34. Tung, Q.N., et al., *Application of genetically encoded redox biosensors to measure dynamic*
changes in the glutathione, bacillithiol and mycothiol redox potentials in pathogenic bacteria.
Free Radic Biol Med, 2018. **128**: p. 84-96.
35. Reuter, W.H., et al., *Utilizing redox-sensitive GFP fusions to detect in vivo redox changes in a*
genetically engineered prokaryote. Redox Biol, 2019. **26**: p. 101280.
36. Marathe, S.A., S. Ray, and D. Chakravortty, *Curcumin increases the pathogenicity of Salmonella*
enterica serovar Typhimurium in murine model. PLoS One, 2010. **5**(7): p. e11511.
37. Smani, Y., et al., *Role of OmpA in the multidrug resistance phenotype of Acinetobacter*
baumannii. Antimicrob Agents Chemother, 2014. **58**(3): p. 1806-8.

38.   Ghai, I. and S. Ghai, *Understanding antibiotic resistance via outer membrane permeability.*
1397   Infect Drug Resist, 2018. **11**: p. 523-530.

1398   39.   Kukkonen, M. and T.K. Korhonen, *The omptin family of enterobacterial surface*
1399   *proteases/adhesins: from housekeeping in Escherichia coli to systemic spread of Yersinia*
1400   *pestis.* Int J Med Microbiol, 2004. **294**(1): p. 7-14.

1401   40.   Ebbensgaard, A., et al., *The Role of Outer Membrane Proteins and Lipopolysaccharides for the*
1402   *Sensitivity of Escherichia coli to Antimicrobial Peptides.* Front Microbiol, 2018. **9**: p. 2153.

1403   41.   Iyer, R., et al., *Acinetobacter baumannii OmpA Is a Selective Antibiotic Permeant Porin.* ACS
1404   Infect Dis, 2018. **4**(3): p. 373-381.

1405   42.   Choi, U. and C.R. Lee, *Distinct Roles of Outer Membrane Porins in Antibiotic Resistance and*
1406   *Membrane Integrity in Escherichia coli.* Front Microbiol, 2019. **10**: p. 953.

1407   43.   Nevermann, J., et al., *Identification of Genes Involved in Biogenesis of Outer Membrane*
1408   *Vesicles (OMVs) in Salmonella enterica Serovar Typhi.* Front Microbiol, 2019. **10**: p. 104.

1409   44.   Calderon, I.L., et al., *Response regulator ArcA of Salmonella enterica serovar Typhimurium*
1410   *downregulates expression of OmpD, a porin facilitating uptake of hydrogen peroxide.* Res
1411   Microbiol, 2011. **162**(2): p. 214-22.

1412   45.   Chaudhuri, D., et al., *Salmonella Typhimurium Infection Leads to Colonization of the Mouse*
1413   *Brain and Is Not Completely Cured With Antibiotics.* Front Microbiol, 2018. **9**: p. 1632.

1414   46.   Li, J., et al., *ChIP-Seq Analysis of the sigmaE Regulon of Salmonella enterica Serovar*
1415   *Typhimurium Reveals New Genes Implicated in Heat Shock and Oxidative Stress Response.*
1416   PLoS One, 2015. **10**(9): p. e0138466.

1417   47.   Mohan Nair, M.K. and K. Venkitanarayanan, *Role of bacterial OmpA and host cytoskeleton in*
1418   *the invasion of human intestinal epithelial cells by Enterobacter sakazakii.* Pediatr Res, 2007.
1419   **62**(6): p. 664-9.

1420   48.   Mittal, R., et al., *Deciphering the roles of outer membrane protein A extracellular loops in the*
1421   *pathogenesis of Escherichia coli K1 meningitis.* J Biol Chem, 2011. **286**(3): p. 2183-93.

1422   49.   Singamsetty, V.K., et al., *Outer membrane protein A expression in Enterobacter sakazakii is*
1423   *required to induce microtubule condensation in human brain microvascular endothelial cells*
1424   *for invasion.* Microb Pathog, 2008. **45**(3): p. 181-91.

1425   50.   Abreu, A.G. and A.S. Barbosa, *How Escherichia coli Circumvent Complement-Mediated Killing.*
1426   Front Immunol, 2017. **8**: p. 452.

1427   51.   Sukumaran, S.K., H. Shimada, and N.V. Prasadarao, *Entry and intracellular replication of*
1428   *Escherichia coli K1 in macrophages require expression of outer membrane protein A.* Infect
1429   Immun, 2003. **71**(10): p. 5951-61.

1430   52.   Brumell, J.H., et al., *Disruption of the Salmonella-containing vacuole leads to increased*
1431   *replication of Salmonella enterica serovar typhimurium in the cytosol of epithelial cells.* Infect
1432   Immun, 2002. **70**(6): p. 3264-70.

1433   53.   Beuzon, C.R., et al., *Salmonella maintains the integrity of its intracellular vacuole through the*
1434   *action of SifA.* EMBO J, 2000. **19**(13): p. 3235-49.

1435   54.   Boncompain, G., et al., *Production of reactive oxygen species is turned on and rapidly shut*
1436   *down in epithelial cells infected with Chlamydia trachomatis.* Infect Immun, 2010. **78**(1): p. 80-
1437   7.

1438   55.   Cheng, C., et al., *Listeriolysin O Pore-Forming Activity Is Required for ERK1/2 Phosphorylation*
1439   *During Listeria monocytogenes Infection.* Front Immunol, 2020. **11**: p. 1146.

1440   56.   Panday, A., et al., *NADPH oxidases: an overview from structure to innate immunity-associated*
1441   *pathologies.* Cell Mol Immunol, 2015. **12**(1): p. 5-23.

1442   57.   Gallois, A., et al., *Salmonella pathogenicity island 2-encoded type III secretion system mediates*
1443   *exclusion of NADPH oxidase assembly from the phagosomal membrane.* J Immunol, 2001.
1444   **166**(9): p. 5741-8.

1445   58.   Webb, J.L., et al., *Macrophage nitric oxide synthase associates with cortical actin but is not*
1446   *recruited to phagosomes.* Infect Immun, 2001. **69**(10): p. 6391-400.

59. Zielonka, J., et al., *Mitigation of NADPH Oxidase 2 Activity as a Strategy to Inhibit Peroxynitrite*
Formation. J Biol Chem, 2016. **291**(13): p. 7029-44.
60. Vlahos, R., et al., *Inhibition of Nox2 oxidase activity ameliorates influenza A virus-induced lung*
inflammation. PLoS Pathog, 2011. **7**(2): p. e1001271.
61. Weller, R., et al., *Antimicrobial effect of acidified nitrite on dermatophyte fungi, Candida and*
bacterial skin pathogens. J Appl Microbiol, 2001. **90**(4): p. 648-52.
62. Kono, Y., et al., *Lactate-dependent killing of Escherichia coli by nitrite plus hydrogen peroxide:*
a possible role of nitrogen dioxide. Arch Biochem Biophys, 1994. **311**(1): p. 153-9.
63. Balakrishnan, A., M. Schnare, and D. Chakravortty, *Of Men Not Mice:*
Bactericidal/Permeability-Increasing Protein Expressed in Human Macrophages Acts as a
Phagocytic Receptor and Modulates Entry and Replication of Gram-Negative Bacteria. Front
Immunol, 2016. **7**: p. 455.
64. Wrande, M., et al., *Genetic Determinants of Salmonella enterica Serovar Typhimurium*
Proliferation in the Cytosol of Epithelial Cells. Infect Immun, 2016. **84**(12): p. 3517-3526.
65. Knodler, L.A., V. Nair, and O. Steele-Mortimer, *Quantitative assessment of cytosolic*
Salmonella in epithelial cells. PLoS One, 2014. **9**(1): p. e84681.
66. Das, P., et al., *Novel role of the nitrite transporter NirC in Salmonella pathogenesis: SPI2-*
dependent suppression of inducible nitric oxide synthase in activated macrophages.
Microbiology (Reading), 2009. **155**(Pt 8): p. 2476-2489.
67. Lahiri, A., P. Das, and D. Chakravortty, *Arginase modulates Salmonella induced nitric oxide*
production in RAW264.7 macrophages and is required for Salmonella pathogenesis in mice
model of infection. Microbes Infect, 2008. **10**(10-11): p. 1166-74.
68. Lahiri, A., P. Das, and D. Chakravortty, *The LysR-type transcriptional regulator Hrg counteracts*
phagocyte oxidative burst and imparts survival advantage to Salmonella enterica serovar
Typhimurium. Microbiology (Reading), 2008. **154**(Pt 9): p. 2837-2846.
69. Coldham, N.G., et al., *A 96-well plate fluorescence assay for assessment of cellular*
permeability and active efflux in Salmonella enterica serovar Typhimurium and Escherichia
coli. J Antimicrob Chemother, 2010. **65**(8): p. 1655-63.

1475

Figure Legends

Figure 1.

OmpA promotes the evasion of phagocytosis and intracellular survival of *Salmonella* in macrophages.

(A) RAW 264.7 cells and PMA activated U937 cells were infected with STM wild type (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60 respectively at MOI 10. 2 hours post-infection, the cells were lysed and plated. The CFU at 2 hours was normalized with pre-inoculum to calculate the percent phagocytosis. Data are represented as mean ± SEM (n=3, N=3 for RAW264.7 cells and n=3, N=2 for activated U937 cells). (B) RAW264.7 cells were

infected with either 10% mouse complement sera treated or untreated STM (WT) and *ΔompA*

respectively at MOI of 50. 2 hours post-infection, the cells were lysed and plated. The CFU at

2 hours was normalized with pre-inoculum to calculate the percent phagocytosis. Data are

represented as mean ± SEM (n=3, N=2). (C) RAW 264.7 cells were infected with STM (WT),

ΔompA, ΔompA: pQE60-*ompA*, & *ΔompA*: pQE60 respectively at MOI 50. Cells were fixed at

15 minutes post-infection with PFA. Externally attached bacteria were probed with anti-

Salmonella antibody without saponin. 20 microscopic fields were analyzed to calculate the no.

of adherent bacteria/ total no. of cells per field. Scale bar = 20μm. Data are represented as mean

± SEM (n=20, N=3). (D) RAW264.7 cells and PMA activated U937 cells were infected with

STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60 respectively at MOI 10. 16-

and 2-hours post-infection, the cells were lysed and plated. The CFU at 16 hours was

normalized with the CFU at 2 hours to calculate the fold proliferation of bacteria in

macrophages. Data are represented as mean ± SEM (n=3, N=3 for RAW 264.7 cells and n=3,

N=2 for activated U937 cells). (E) RAW264.7 cells were infected with STM- (WT): RFP,

ΔompA: RFP, *ΔompA*: pQE60-*ompA* at MOI of 20. PFA-fixed wild-type bacteria were used for

infection at MOI 25. Cells were fixed at 16 hours post-infection & LAMP-1 was labeled with

anti-mouse LAMP-1 antibody. To stain the complemented strain and PFA fixed dead bacteria

anti-*Salmonella* antibody was used. The quantification of LAMP-1 recruitment on bacteria in

RAW 264.7 cells has been represented in a graph. Percent colocalization was determined after

analyzing 50 different microscopic stacks from three independent experiments. Scale bar =

5μm. Data are represented as mean ± SEM. (n=50, N=3). (F) Chloroquine resistance assay of

RAW 264.7 cells infected with STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA* strains,

respectively. Data are represented as mean ± SEM (n=3, N=2).

(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant,

(Student's *t*-test- unpaired).

Figure 2.

OmpA dependent invasion and intracellular proliferation of *Salmonella* inside the epithelial cell.

(A) Caco-2 cells and HeLa cells were infected with log phase culture of STM (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60 respectively at MOI 10. 2 hours post-infection, the cells were lysed and plated. The CFU at 2 hours was normalized with pre-inoculum to calculate the percent invasion in epithelial cells. Data are represented as mean ± SEM (n=3, N=3). (B) HeLa cells were infected with the log phase culture of STM (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60, respectively at MOI of 50. Cells were fixed at 25 minutes post-infection with PFA. Externally attached bacteria were probed with anti-*Salmonella* antibody without saponin. 20 microscopic fields were analyzed to calculate the no. of adherent bacteria/ total no. of cells per field. Data are represented as mean ± SEM (n=20, N=3). Scale bar = 20μm. (C) Caco-2 and HeLa cells were infected with log phase culture of STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60 respectively at MOI 10. 16- and 2-hours post-infection, the cells were lysed and plated. The CFU at 16 hours was normalized with the CFU at 2 hours to calculate the fold proliferation of bacteria in epithelial cells. Data are represented as mean ± SEM (n=3, N=3). (D) Caco-2 cells were infected with STM (WT): RFP, *ΔompA*: RFP, and *ΔompA*: pQE60-*ompA* at MOI 20. PFA-fixed wild-type bacteria were used for infection at MOI 25. Cells were fixed at 16 hours post-infection & LAMP-1 was labeled with anti-human LAMP-1 antibody. The complemented strain and PFA fixed dead bacteria were tracked with an anti-*Salmonella* antibody. The quantification of LAMP-1 recruitment on bacteria in Caco-2 cells has been represented in the form of a graph. Percent colocalization was determined after analyzing 50 different microscopic stacks from three independent experiments. Scale bar = 5μm. Data are represented as mean ± SEM (n=50, N=3). (E)

Chloroquine resistance assay of Caco-2 cells infected with STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA* strains. Data are represented as mean ± SEM (n=3, N=2).

(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant, (Student's *t*-test- unpaired).

Figure 3.

OmpA dependent protection of *Salmonella* against nitrosative stress inside RAW264.7 cells.

(A) Estimation of extracellular nitrite from the culture supernatant of RAW264.7 cells infected with STM (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, *ΔompA*: pQE60, & heat-killed bacteria respectively at MOI 10. 16 hours post-infection, the culture supernatants were collected and subjected to Griess assay. Data are represented as mean ± SEM (n=3, N=5). (B) Estimation of intracellular nitric oxide level in RAW 264.7 cells infected with STM (WT), *ΔompA*, and *ΔompA*: pQE60-*ompA* at MOI 10, 16 hours post-infection using DAF-2 DA [5 µM] by flow cytometry. Unstained and uninfected RAW264.7 cells have also been used as controls. Both dot plots (SSC-A vs. DAF-2 DA) and histograms (Count vs. DAF-2 DA) have been represented. The percent population of DAF-2 DA positive RAW264.7 cells has been represented in the form of a graph. Data are represented as mean ± SEM (n≥3, N=4). (C) Immunofluorescence image of RAW264.7 cells infected with STM (WT): RFP, *ΔompA*: RFP, (WT): *LLO*, and *ΔompA*: *LLO* at MOI 20. 16 hours post-infection, the cells were fixed with PFA and labeled with anti-*Salmonella* antibody and anti-mouse nitrotyrosine antibody followed by visualization under confocal microscopy. Percent colocalization of STM (WT), *ΔompA*, (WT): *LLO,* and *ΔompA*: *LLO* with nitrotyrosine was determined by analyzing 50 different microscopic stacks from three independent experiments. Data are represented as mean ± SEM (n=50, N=3). Scale bar = 5µm. (D) RAW264.7 cells were infected with STM (WT), *ΔompA*, (WT): *LLO,* and *ΔompA*: *LLO* at MOI 10. 16- and 2-hours post-infection, the cells

were lysed and plated. The CFU at 16 hours was normalized with the CFU at 2 hours to calculate the fold proliferation of bacteria. Data are represented as mean ± SEM (n≥3, N=2). (E) The transcript-level expression profile of *spiC* from RAW264.7 cells infected with STM (WT), *ΔompA* & (WT): *LLO* at MOI 50. STM (WT) infected RAW264.7 cells treated with bafilomycin A (50 nM) were used as a control. 12 hours post-infection, the cells were lysed, and total RNA was extracted. After the synthesis of cDNA, the expression of *spiC* was measured by RT PCR. Data are represented as mean ± SEM (n=3, N=3). (F) The measurement of the activity of *spiC* promoter in stationary phase culture of STM (WT) and *ΔompA* growing overnight in acidic F media and neutral LB media. Data are represented as mean ± SD (n=6). (G) The measurement of the activity of *spiC* promoter in STM (WT) and *ΔompA* proliferating intracellularly in RAW264.7 cells (MOI= 50) 12 hours post-infection. Macrophages infected with STM (WT) and further treated with bafilomycin A (50 nM) have been used as a negative control. Data are represented as mean ± SEM (n=5, N=2). (H) Estimation of intracellular acidification of STM (WT), *ΔompA*, and *ΔompA*: pQE60-*ompA* in phosphate buffer of pH 5.5, 6, 6.5, and 7, respectively using 20 µM of ratiometric pH indicator BCECF-AM by flow cytometry. The ratio of median fluorescence intensity of BCECF-AM labeled on STM- (WT), *ΔompA*, and *ΔompA*: pQE60-*ompA* at 488 and 405 nm, respectively. Data are represented as mean ± SEM (n=4, N=3).

(P) *< 0.05, *(P)* **< 0.005, *(P)* ***< 0.0005, *(P)* ****< 0.0001, ns= non-significant, (Student's *t* test- unpaired).

Figure 4.

Alteration in the activity of iNOS by using a specific inhibitor or activator changes the fate of STM *ΔompA* in *in vitro* and *in vivo* infection models.

Intracellular survival of STM (WT) and *ΔompA* (MOI= 10) in RAW264.7 cells (16 hours post-infection) in presence and absence of iNOS (A) inhibitor- 1400W dihydrochloride [10 µM]

and (B) activator- mouse IFN-ɣ [100U/ mL]. Fold proliferation of bacteria was calculated by

normalizing the CFU at 16 hours to CFU at 2 hours. Data are represented as mean ± SEM (n=3,

N=3). (C) Immunofluorescence image of RAW264.7 cells infected with STM- (WT) and

ΔompA (MOI=20) in presence and absence of iNOS inhibitor- 1400W dihydrochloride [10 μM]

and activator- mouse IFN-ɣ [100U/ mL]. 16 hours post-infection, the cells were fixed with PFA

and probed with anti-*Salmonella* antibody and anti-mouse nitrotyrosine antibody. The

quantification of the recruitment of nitrotyrosine on STM (WT) and *ΔompA* (MOI-20) in

RAW264.7 cells (16 hours post-infection) in presence and absence of iNOS (D) inhibitor-

1400W dihydrochloride [10 μM] and (E) activator- mouse IFN-ɣ [100U/ mL]. Percent

colocalization of bacteria with nitrotyrosine was determined after analyzing more than 50

different microscopic stacks from two independent experiments. Data are represented as mean

± SEM (n≥ 50, N=2). (F) The schematic representation of the strategy of animal experiments.

(G) Four cohorts of 4 to 6 weeks old five C57BL/6 mice were orally gavaged with STM- (WT)

and *ΔompA* at a sublethal dose of 10^7 CFU/ animal. Two of these four cohorts were

intraperitoneally injected with iNOS inhibitor aminoguanidine hydrochloride (10mg/ kg of

body weight) regularly for 5 days post-infection. Two cohorts of five *iNOS$^{-/-}$* and *gp91phox$^{-/-}$*

mice were orally gavaged with STM- (WT) and *ΔompA* at a sublethal dose of 10^7 CFU/ animal

separately. On the 5^{th}-day post-infection, the mice were sacrificed, followed by isolation,

homogenization of liver, spleen, and MLN. The organ lysates were plated on *Salmonella*

Shigella agar. The colonies were counted 16 hours post-plating. The $\log_{10}$[CFU/ gm-wt.] for

each CFU has been plotted. Data are represented as mean ± SEM (n=5, N=3). (H) Two cohorts-

each consisting of 4 to 6 weeks old 20- BALB/c and C57BL/6 mice infected with a lethal dose

(10^8 CFU/ animal) of STM- (WT) and *ΔompA*. After they were infected by oral gavaging, the

survival of the mice was monitored till the death of all the wild-type infected mice (n=10).

*(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant,*

(Student's *t* test- unpaired, Mann-Whitney *U* test- for animal survival assay).

Figure 5.

OmpA dependent regulation of outer membrane permeability in *Salmonella* controls

cytoplasmic redox homeostasis in response to *in vitro* nitrosative stress.

(A) Time-dependent *in vitro* death kinetics of STM (WT) and *ΔompA* in the presence of

acidified nitrite (Nitrite concentration 800 µM in PBS of pH 5.4). 10^8 CFU of overnight grown

stationary phase culture of both the strains were inoculated in acidified nitrite. The survival of

both the strains was determined by plating the supernatant with appropriate dilution in SS agar

plates at 0, 3, 6, 9, 12 hours post-inoculation. Data are represented as mean ± SEM (N=5). (B)

In vitro nitrite uptake assay of STM (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, *ΔompA*: pQE60, &

PFA fixed dead bacteria. 10^8 CFU of overnight grown stationary phase culture of all the strains

were inoculated in MOPS- NaOH buffer (pH= 8.5) with an initial nitrite concentration of 200

µM. The remaining nitrite concentration of the media was determined by Griess assay at 0, 3,

6, 9, 12 hours post-inoculation. Data are represented as mean ± SEM (n=3, N=4). (C)

Measurement of redox homeostasis of STM (WT) and *ΔompA* harboring pQE60-Grx1-roGFP2

in response to varying concentrations of acidified nitrite in a time-dependent manner. $4.5X\ 10^7$

CFU of overnight grown stationary phase culture of STM (WT) and *ΔompA* expressing pQE60-

Grx1-roGFP2 were subjected to the treatment of acidified nitrite (concentration of 800µM,

1mM, 5mM) for 15, 30, 45, and 60 minutes. Median fluorescence intensities of Grx1-roGFP2

at 405nm and 488nm for the FITC positive population were used to obtain the 405/ 488 ratio

in 800 µM, 1mM, 5 mM concentration of acidified nitrite, respectively. Data are represented

as mean ± SEM (n=3, N=3). (D) The transcript level expression profiling of larger porin genes,

namely- *OmpC*, *OmpD*, *OmpF* by real-time PCR in STM (WT), *ΔompA*, & *ΔompA*: pQE60-

ompA in nutritionally enriched LB media, low magnesium acidic F media (pH= 5.4) mimicking

the internal environment of SCV, and RAW264.7 murine macrophage cells (MOI 50) [12 hours

post inoculation]. Data are represented as mean ± SEM (n=3, N=2). (E) Measurement of

membrane porosity of STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, & *ΔompA*: pQE60 in acidic

F media (12 hours post-inoculation) using a slightly negatively charged dye named DiBAC$_4$

(final concentration- 1 µg/ mL) by flow cytometry. Unstained bacterial cells were used as

control. Both dot plots (SSC-A vs. DiBAC$_4$) and histograms (Count vs. DiBAC$_4$) have been

represented. The median fluorescence intensity of DiBAC$_4$ has been represented here. Data are

represented as mean ± SEM (n=3, N=2). (F) Measurement of outer membrane porosity of

STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA* & *ΔompA*: pQE60 in acidic F media [12 hours

post inoculation] using a nuclear binding fluorescent dye bisbenzimide [excitation- 346 nm and

emission- 460 nm] (final concentration- 1 µg/ mL). (WT) treated with 0.1% saponin for 15

minutes has been used as a positive control. Data are represented as mean ± SEM (n=8, N=3).

(G) Measurement of outer membrane porosity of STM (WT) and *ΔompA* isolated from infected

RAW264.7 cells 12 hours post-infection by bisbenzimide (Sigma) (final concentration- 1 µg/

mL). Data are represented as mean ± SEM (n=6, N=3).

(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant, (2way

ANOVA, Student's *t*-test- unpaired).

Figure 6.

The maintenance of the integrity of the SCV membrane inside RAW264.7 macrophages

solely depends upon OmpA, not on other larger porins such as OmpC, OmpD, and

OmpF.

(A) RAW264.7 cells were infected with STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*,

ΔompAΔompF, & WT: *LLO* at MOI 20. Cells were fixed at 16 hours post-infection, followed

by labeled with anti-*Salmonella* antibody and anti-mouse LAMP-1 antibody. (B)

Quantification of LAMP-1 recruitment on STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*,

ΔompAΔompF, (WT): *LLO*. Percent colocalization between bacteria and LAMP-1 was

determined after analyzing more than 50 different microscopic stacks from two independent

experiments. Data are represented as mean ± SEM (n≥50, N=2). Scale bar = 5μm. (C)

RAW264.7 cells were infected with STM (WT), *ΔompA*, *ΔompC*, *ΔompD*, and *ΔompF* at MOI

20. Cells were fixed at 16 hours post-infection, followed by labeled with anti-*Salmonella*

antibody and anti-mouse LAMP-1 antibody, respectively. (D) Quantification of LAMP-1

recruitment on STM (WT), *ΔompA*, *ΔompC*, *ΔompD*, and *ΔompF*. Percent colocalization of

bacteria with LAMP-1 has been determined after analyzing more than 60 different microscopic

stacks from two independent experiments. Data are represented as mean ± SEM (n≥60, N=2).

Scale bar = 5μm.

*(P) ****< 0.0001, (Student's t test- unpaired).*

Figure 7.

In the absence of OmpA, porins OmpC and OmpF enhance the susceptibility of

***Salmonella* against the nitrosative stress of RAW264.7 cells.**

(A) *In vitro* nitrite uptake assay of STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*,

ΔompAΔompF, & PFA fixed dead bacteria. 10^8 CFU of overnight grown stationary phase

culture of all the strains was inoculated in MOPS- NaOH buffer (pH= 8.5) with 100 μM initial

nitrite concentration. The remaining nitrite in the media was estimated by Griess assay at 0, 2,

4, 6, 8 hours post-inoculation. Data are represented as mean ± SEM (n=3, N=6). (B) *In vitro*

viability assay of STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*, & *ΔompAΔompF* in the

presence of acidified nitrite [800 μM] 12 hours post-inoculation using resazurin solution. Data

are represented as mean ± SEM (n=3, N=3). (C) Immunofluorescence image of RAW264.7

cells infected with STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*, *ΔompAΔompF*, &

(WT): *LLO* at MOI 20. Cells were fixed at 16 hours post-infection, followed by labeled with

anti-*Salmonella* antibody and anti-mouse nitrotyrosine antibody. Quantification of

nitrotyrosine recruitment on STM (WT), *ΔompA*, *ΔompAΔompC*, *ΔompAΔompD*,

ΔompAΔompF, (WT): *LLO* has been represented in the form of a graph. Percent colocalization

between bacteria and nitrotyrosine was determined after analyzing more than 60 different

microscopic fields from two independent experiments. Data are represented as mean ± SEM

(n≥60, N=3). Scale bar = 5μm. (D) Intracellular survival of STM (WT), *ΔompA*,

ΔompAΔompC, *ΔompAΔompD*, & *ΔompAΔompF*, & (WT): *LLO* respectively (MOI 10) in

RAW264.7 cells. 16- and 2-hours post-infection, the cells were lysed and plated. The CFU at

16 hours was normalized with the CFU at 2 hours to calculate the fold proliferation of bacteria

in macrophages. Data are represented as mean ± SEM (n=3, N=3). (E) Estimation of the

intracellular nitrite level in RAW 264.7 cells infected with STM (WT), *ΔompA*,

ΔompA:pQE60-*ompA*, *ΔompAΔompC*, *ΔompAΔompD*, *ΔompAΔompF*, and (WT): *LLO*

respectively at MOI 10, 16 hours post-infection using DAF-2DA [5 μM] by flow cytometry.

Unstained and uninfected macrophages have also been used as a control. Both dot plots (SSC-

A vs. DAF-2 DA) and histograms (Count vs. DAF-2 DA) have been represented. The percent

population of DAF-2DA positive macrophages has been represented in the form of a bar graph.

Data are represented as mean ± SEM (n≥3, N=5).

(P) *< 0.05, (P) **< 0.005, (P) *< 0.0005, (P) ****< 0.0001, ns= non-significant,**

(Student's *t* test- unpaired).

Figure 8.

The hypothetical working model of intracellular survival of STM (WT), STM *ΔompA*,

and STM (WT): *LLO*.

The hypothetical model depicts the fate of (A) STM (WT), (B) STM *ΔompA*, and (C) STM

(WT): *LLO* inside the murine macrophages. (A) STM (WT) staying inside the acidic SCV can

proliferate efficiently by suppressing the activity of iNOS by SPI-2 encoded virulent factor

SpiC. The acidification of the cytosol of wild-type bacteria due to the acidic pH of SCV triggers the expression of SPI-2 genes. (B) STM *ΔompA,* which comes into the cytosol of macrophages after quitting SCV, exhibits an attenuated expression of SpiC because of their stay in a less acidic environment. It is unable to suppress the activity of iNOS and is heavily bombarded with RNI in the cytosol of macrophages. The enhanced outer membrane permeability of the cytosolic population of STM *ΔompA* due to the upregulation of *ompC*, *ompD*, and *ompF* makes them vulnerable towards RNI. (C) STM (WT): *LLO* quits the SCV by expressing LLO. Unlike STM (WT), the cytosolic niche of STM (WT): *LLO* cannot produce SpiC. It can protect itself from RNI by reducing its outer membrane permeability by expressing *ompA* and survive efficiently in the cytosol of macrophages.

Supplementary Figures

Figure S1.

OmpA plays a crucial role in the survival of *Salmonella* Typhimurium in murine macrophages.

The transcript level expression profile of (A) *ompA*, (B) *ompC*, (C) *ompD*, and (D) *ompF* in STM- (WT) at indicated time points (3, 6, 9, 12 hours) in LB broth (n=3, N=3), low magnesium acidic F media (pH=5.4) (n=3, N=3), and RAW264.7 murine macrophage cells (MOI= 50), (n=3, N=3). The time-dependent relative expression of *ompA, ompC, ompD, ompF* have been represented in the log2 scale. The predicted structures of porins (A) OmpA, (B) OmpC, (C) OmpD, and (D) OmpF using SWISS-MODEL protein structure homology-modeling server. *(P) *< 0.05, (P) **< 0.005, (Student's t test- unpaired).*

Figure S2.

STM *ΔompA* comes out into the cytoplasm of the macrophages after quitting the vacuole.

(A) The schematic representation of the SPI-2 encoded T3SS and the secreted effector proteins SseC and SseD around intracellular wild-type *Salmonella*. (B-D) RAW264.7 cells were infected with STM (WT)-: RFP, *ΔompA*: RFP, at MOI 20. Cells were fixed at 16 hours post-infection, *Salmonella* SPI-2 encoded translocon proteins SseC and SseD were labeled with anti- *Salmonella* SseC/ SseD antibody. (C), (D) The quantification of SseC and SseD recruitment on bacteria in RAW 264.7 cells, respectively. Percent colocalization between the bacteria and the effector was determined after analyzing 50 different microscopic stacks from three independent experiments. Scale bar = 5μm, (n=50, N=3). (E-H) The transcript level expression of (E) *sseC*, (F) *sseD*, (G) *ssaV*, and (H) *sifA* in STM (WT) and *ΔompA* growing intracellularly in RAW264.7 cells 12 hours post-infection, (n=3, N=3).

(P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant, (Student's t test-unpaired).

Figure S3.

OmpA does not have a significant role in building up *in vitro* and *in vivo* protection of *Salmonella* against oxidative stress.

(A) Estimation of the level of intracellular reactive oxygen species (ROS) in RAW 264.7 cells infected with STM- (WT), *ΔompA*, and *ΔompA*: pQE60-*ompA* at MOI 10, 16 hours post-infection using DCFDA [10 µM] by flow cytometry. Unstained and uninfected RAW264.7 cells have also been used as controls. Both dot plots (SSC-A vs. DCFDA) and histograms (Count vs. DCFDA) have been represented. (B) The percent population of DACFDA positive RAW264.7 cells, (n=4, N=3). (C) Estimation of extracellular ROS from the culture supernatant of RAW264.7 cells infected with STM- (WT), *ΔompA*, *ΔompA*: pQE60-*ompA*, *ΔompA*: pQE60, & PFA fixed dead bacteria respectively at MOI 10. 2 hours post-infection, the cells were

supplemented with phenol red solution having horseradish peroxidase. 16 hours post-infection,

the culture supernatants were collected, and the OD was measured at 610 nm (n=3, N=2). (D)

Checking the *in vitro* sensitivity of STM- (WT) and *ΔompA* in the presence of H_2O_2 by serial

dilution, plating, and CFU calculation and resazurin test. 10^8 CFU of overnight grown

stationary phase culture of STM (WT) and *ΔompA* were inoculated in PBS with varying

concentrations of H_2O_2. 12 hours post-inoculation, the supernatants were collected for plating

on SS agar to determine the log_{10}[CFU/mL] (N=3) and resazurin test to determine percent

viability. (n=3, N=3). (E) Checking the *in vitro* sensitivity of STM– (WT) and *ΔompA* in the

presence of acidified nitrite by serial dilution, plating, and CFU calculation and resazurin test.

10^8 CFU of overnight grown stationary phase culture of STM (WT) and *ΔompA* were

inoculated in PBS (pH=5.4) with varying concentrations of $NaNO_2$. 12 hours post-inoculation,

the supernatants were collected for plating on SS agar to determine the log_{10}[CFU/mL], (N=3)

and resazurin test to determine percent viability. (n=3, N=3). (F) Checking the *in vitro*

sensitivity of STM– (WT) and *ΔompA* in the presence of $NaNO_2$ and H_2O_2 combined, by serial

dilution, plating, and CFU calculation and resazurin test. 10^8 CFU of overnight grown

stationary phase culture of STM (WT) and *ΔompA* were inoculated in PBS (pH=5.4) with

varying concentrations of $NaNO_2$ and H_2O_2. 12 hours post-inoculation, the supernatants were

collected for plating on SS agar to determine the log_{10}[CFU/mL] (N=3) and resazurin test to

determine percent viability (n=3, N=3).

(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant, (2way

ANOVA), (Student's *t* test- unpaired).

Figure S4.

Over-expression of *ompF* in wild-type *Salmonella* enhances the outer membrane porosity

of the bacteria.

Measurement of outer membrane porosity of STM (WT), STM (WT): pQE60, STM (WT): pQE60-*ompA*, STM (WT): pQE60-*ompC*, STM (WT): pQE60-*ompD*, and STM (WT): pQE60-*ompF* in (A) acidic F media and (C) LB broth with 500 μM of IPTG [12 hours post inoculation] using DiBAC$_4$ (final concentration- 1 μg/ mL) by flow cytometry. Unstained bacterial cells were used as control (A) and (C). Both dot plots (SSC-A vs. DiBAC$_4$) and histograms (Count vs. DiBAC$_4$) have been represented. Percent population of DiBAC$_4$ positive cells in acidic F media (B) and LB broth culture (D) has been represented here in the form of a bar graph (n=6, N=3 for B and n=6 for D).

(P) *< 0.05, (P) **< 0.005, (P) *< 0.0005, (P) ****< 0.0001, ns= non-significant, (Student's *t* test-unpaired)**

Figure S5.

Unlike *ompA*, the deletion of *ompC, ompD*, and *ompF* do not hamper the viability of *Salmonella* against intracellular and extracellular nitrosative stress.

(A) RAW264.7 cells were infected with STM- (WT), *ΔompA*, *ΔompC*, *ΔompD*, and *ΔompF* at MOI 20. Cells were fixed at 16 hours post-infection, followed by labeled with anti-*Salmonella* antibody and anti-mouse nitrotyrosine antibody, respectively. Quantification of nitrotyrosine recruitment on STM- (WT), *ΔompA*, *ΔompC*, *ΔompD*, and *ΔompF* has been represented in the form of a graph. Percent colocalization of bacteria with nitrotyrosine has been determined after analyzing more than 60 different microscopic fields from two independent experiments (n≥60, N=3). Scale bar = 5μm. (B) Checking the *in vitro* sensitivity of STM– (WT), *ΔompA*, *ΔompC*, *ΔompD*, and *ΔompF* in the presence of acidified nitrite by serial dilution, plating, and CFU calculation. 10^8 CFU of overnight grown stationary phase culture of STM (WT), *ΔompA, ΔompC, ΔompD,* and *ΔompF* were inoculated in PBS (pH=5.4) with varying concentrations of NaNO$_2$. 12 hours post-inoculation, the supernatants were collected for plating on SS agar to

determine the $\log_{10}$[CFU/mL], (N=3). (C) *In vitro* nitrite uptake assay of STM- (WT), *ΔompA*,

ΔompC, ΔompD, ΔompF & PFA fixed dead bacteria. 10^8 CFU of overnight grown stationary

phase culture of all the strains were inoculated in MOPS- NaOH buffer (pH= 8.5) with an initial

nitrite concentration of 50 µM. The remaining nitrite concentration of the media was

determined by Griess assay at 0, 1, 2, 3-, 4-, 5-, and 6-hours post-inoculation (n=3, N=3).

(P) *< 0.05, (P) **< 0.005, (P) *< 0.0005, (P) ****< 0.0001, ns= non-significant, (2way**

ANOVA), (Student's *t* test).

Figure S6.

Overexpression of *ompF* enhances the susceptibility of wild-type *Salmonella* towards *in*

***vitro* nitrosative stress.**

Measurement of *in vitro* viability of STM (WT), STM (WT): pQE60, STM (WT): pQE60-

ompA, STM (WT): pQE60-*ompC*, STM (WT): pQE60-*ompD*, and STM (WT): pQE60-*ompF*

in acidified nitrite (PBS of pH= 5.4 and 1 mM NaNO$_2$ at 12 hours post inoculation using

propidium iodide (final concentration- 1 µg/ mL) by flow cytometry (A). Unstained bacterial

cells were used as control. Bacterial cells grown in acidified PBS were stained with propidium

iodide to determine the bacterial death contributed by *in vitro* reactive nitrogen intermediates.

Both dot plots (SSC-A vs. DiBAC$_4$) and histograms (Count vs. DiBAC$_4$) have been

represented. Percent population of propidium iodide positive cells from acidified nitrite have

been represented here in the form of bar graph (n=8, N=2). Measurement of *in vitro* viability

of STM (WT), STM (WT): pQE60, STM (WT): pQE60-*ompA*, STM (WT): pQE60-*ompC*,

STM (WT): pQE60-*ompD*, and STM (WT): pQE60-*ompF* in acidified PBS (C and E) and

acidified nitrite (PBS of pH= 5.4 and 1 mM NaNO$_2$) (D and E) at 12 hours post inoculation

using resazurin (final concentration- 0.002 mg/ mL) (n=6 for C and n=8 for D).

(P) *< 0.05, (P) **< 0.005, (P) ***< 0.0005, (P) ****< 0.0001, ns= non-significant,

(Student's *t* test- unpaired).

1826

1827

1828

(A) RAW264.7   U937

(B) RAW264.7

(C) RAW264.7

STM (WT)   STM *ΔompA*

STM *ΔompA*: pQE60   STM *ΔompA*: pQE60-*ompA*

(D) RAW264.7   U937

(E) [16 hours post infection]

STM (WT)

STM *ΔompA*

STM *ΔompA*: pQE60-*ompA*

PFA fixed STM (WT)

(F)

Mouse complement serum [10%]

🟩 **LAMP-1**   🟥 **STM**   🟨 **Colocalization**

🟥 **Cytosolic**
🟨 **Vacuolar**

(A) Caco-2 / HeLa — Percent Invasion [X 10⁻³]

(B) HeLa — STM (WT), STM ΔompA, STM ΔompA: pQE60-ompA, STM ΔompA: pQE60 — No of Adherent Bacteria/Total No of Cells

(C) Caco-2 (Fold Change) / HeLa (Fold Proliferation)

(D) [16 hours post infection] — STM (WT), STM ΔompA, STM ΔompA: pQE60-ompA, PFA fixed STM (WT)

LAMP-1 (green) | STM (red) | Colocalization (yellow)

(E) Percent abundance of bacteria — STM WT, STM ΔompA, STM ΔompA: pQE60-ompA

Cytosolic (red) | Vacuolar (yellow)

Percent Colocalization between bacteria and LAMP-1 — STM (WT), STM ΔompA, STM ΔompA: pQE60-ompA

(A) RAW264.7

(B) RAW264.7

(C) [16 hours post infection]

Nitrotyrosine ▮ STM ▮ Colocalization

(D) RAW264.7

(E) RAW264.7

(F)

(G) RAW264.7

(H) Phosphate Buffer

(A)

[16 hours post infection]

STM (WT)

STM ΔompA

STM ΔompA ΔompC

STM ΔompA ΔompD

STM ΔompA ΔompF

STM (WT) LLO

LAMP-1
STM
Colocalization

(B)

Percent Colocalization between bacteria and LAMP-1

****
****
****
****
****

STM (WT) STM ΔompA STM ΔompAΔompC STMΔompAΔompD STMΔompAΔompF STM (WT)- LLO

(C)

[16 hours post infection]

STM (WT)

STM ΔompA

STM ΔompC

STM ΔompD

STM ΔompF

(D)

Percent Colocalization between bacteria and LAMP-1

****
****
****

STM (WT) STM ΔompA STM ΔompC STM ΔompD STM ΔompF

(A)

(B)

(C) [16 hours post infection]

Nitrotyrosine
STM
Colocalization

(D)

(E)

DAF2DA

(A)

16 hours post infection

STM (WT)

STM *ΔompA*

STM *ΔompC*

STM *ΔompD*

STM *ΔompF*

■ **Nitrotyrosine** ■ **STM** ■ **Colocalization**

(B)

(C) Concentration of [NO₂⁻]

(A)

STM (WT):
pQE60

STM (WT):
pQE60-*ompA*

STM (WT):
pQE60-*ompC*

STM (WT):
pQE60-*ompD*

STM (WT):
pQE60-*ompF*

0.04% 0.62% 0.64%

0.02% 0.63% 0.7%

0.04% 0.72% 0.89%

0.00% 14.97% 16.17%

0.16% 7.54% 22.8%

Autofluorescence   **PBS (pH= 5.4)**   **PBS (pH= 5.4) +
1mM NaNO₂**

(B)

Propidium Iodide positive population

****   ****

STM (WT): pQE60
STM (WT): pQE60-*ompA*
STM (WT): pQE60-*ompC*
STM (WT): pQE60-*ompD*
STM (WT): pQE60-*ompF*

(C) **Acidified PBS**

Percent viability of bacterial cells
[normalised with untransformed wild type]

STM (WT)
STM (WT); pQE60
STM (WT); pQE60-*ompA*
STM (WT); pQE60-*ompC*
STM (WT); pQE60-*ompD*
STM (WT); pQE60-*ompF*

(D) **Acidified PBS +
1mM NaNO₂**

Percent viability of bacterial cells
[normalised with untransformed wild type]

****

****

STM (WT)
STM (WT); pQE60
STM (WT); pQE60-*ompA*
STM (WT); pQE60-*ompC*
STM (WT); pQE60-*ompD*
STM (WT); pQE60-*ompF*

(E)

Acidified PBS

Acidified PBS +
1mM NaNO₂

STM (WT)
STM (WT): pQE60
STM (WT): pQE60-*ompA*
STM (WT): pQE60-*ompC*
STM (WT): pQE60-*ompD*
STM (WT): pQE60-*ompF*