Quantifying habitat and landscape effects on composition and structure

of plant-pollinator networks in the US Northern Great Plains

Isabela B. Vilella-Arnizaut, Corresponding Author[1]

Department of Biology and Microbiology, South Dakota State University, 1390 College

10 Ave, Brookings, SD 57007

Charles Fenster[2]

Director Oak Lake Field Station, South Dakota State University, Brookings, South

Dakota, USA

Henning Nottebrock[3]

Department of Biology and Microbiology, South Dakota State University, Brookings,

South Dakota, USA

[1]izzyvilella22@yahoo.com

[2]charles.fenster@sdstate.edu

[3]henning.nottebrock@uni-bayreuth.de

[3]Current address: Plant Ecology, Bayreuth Center of Ecology and Environmental Research (BayCEER), University
of Bayreuth, Universitätsstr. 30, Bayreuth, Germany

Abstract

Community structure contributes to ecosystem persistence and stability. To understand the

mechanisms underlying pollination and community stability of natural areas in a human

influenced landscape, a better understanding of the interaction patterns between plants and

pollinators in disturbed landscapes is needed. The Northern Great Plains still retain extensive

tracts of remnant temperate grassland habitat within a matrix of varying land-uses. We used a

network-based approach to quantify how temperate grassland attributes and landscape

heterogeneity influence plant-pollinator community structure in natural habitats. We also

quantified pollinator diversity and floral diversity to assess the functional role of temperate

grassland attributes and the surrounding landscape on the composition of the plant-pollinator

communities in natural habitats. We found that the amount of local nectar sugar and increased

proportions of certain land-uses contribute to pollinator diversity that in turn influences the

structure of interactions between plants and pollinators. Understanding the factors contributing to

plant-pollinator network structure can guide management decisions to support resilient plant-

pollinator communities and conserve the stability of pollination services.

KEYWORDS: Plant-pollinator interactions, Landscape Ecology, Pollination, Pollinator

Diversity, Nectar, Network Analysis, Natural Areas, Conservation

BODY OF ARTICLE:

Introduction

Habitat loss and fragmentation are among the most prominent anthropogenic pressures impacting

biodiversity globally (Foley et al. 2005; Tscharntke et al. 2005; Lundgren and Fausti 2015; Greer

et al. 2016). Increased intensive agriculture has facilitated the conversion of natural vegetation to

crop monoculture (Ramankutty and Foley 1999; Rashford et al. 2011; Wright and Wimberly

2013). Globally, temperate grasslands are considered to be at the greatest risk for biodiversity

loss and ecosystem dysfunction, specifically due to extensive landscape conversion and low rates

of habitat protection (Hoekstra et al. 2005). The disparity between landscape conversion and

protection for grasslands is particularly concerning since grasslands represent essential habitat

and resources for multiple species including insect pollinators. Animal-driven pollination is

essential for the production of approximately 35% of crops worldwide (Klein et al. 2007;

Vanbergen et al. 2013) and contributes to the reproduction of over 70% of flowering plant

species (Potts et al. 2010). Nevertheless, there is an alarming global decline of insect pollinators

largely attributed to anthropogenic pressures such as land-use intensification and widespread use

of pesticides (Kearns et al. 1998; Steffan-Dewenter et al. 2005). Given these documented

declines, there is an increased interest in conserving pollinator communities and their habitats to

stabilize these critical ecosystem services and in turn, ecosystem function (Steffan-Dewenter et

al. 2002; Kremen et al. 2002; Olesen et al. 2007; Peterson et al. 2010; Redhead et al. 2018;

Jauker et al. 2019).

An understudied determinant of pollinator communities in natural areas is the diversity and

configuration of the landscape that exists in the mosaic of a mixed-use agricultural and natural

landscape. Pollinator communities are negatively impacted when their populations become

increasingly isolated from valuable resources (i.e., nectar resources, pollen resources, nesting

resources, etc.) (Olesen et al. 1994; Kearns et al. 1998; Garibaldi et al. 2011). Landscape

composition (e.g., patch size) and configuration (e.g., fragmentation) can influence insect

pollinator behavior and movement (Fahrig et al. 2011; Hadley and Betts 2012; Moreira et al.

2015; Sakai et al. 2016). For example, proximity to semi-natural habitats increases the

abundance of both wild bees and honey bees in crops (Steffan-Dewenter et al. 2002; Kremen et

al. 2004; Heard et al. 2007). Likewise, increased availability and diversity of flowering plants in

natural areas benefits pollinator populations (Potts et al. 2003; Potts et al. 2005; Ponisio et al.

2019; Requier et al. 2020). Recent studies (Nottebrock et al. 2017) demonstrate that the level of

available sugar resources to pollinators can have considerable effects on the outcome of plant-

pollinator interactions. Thus, the consequences of landscape conversion could impact bee

populations on numerous levels (i.e., hive survival and productivity) by decreasing availability

and diversity of nutritional floral resources (Naug 2009; Pettis et al. 2013; Otto et al. 2016).

Within the midwestern United States, the Northern Great Plains has served as a refuge for

approximately 40% of the commercial honey bee colonies from May through October (USDA,

2014). The regional blooms provided by livestock-grazed pastures and grasslands in the

Northern Great Plains have sustained transported honey bee colonies due to the presence and

abundance of floral resources (Otto et al. 2016). However, honey bee colonies in the US and

Europe continue to sustain annual losses which can be attributed to a combination of factors such

as disease, pests, and pesticides (Williams 2002; Cox-Foster et al. 2007; Cox-Foster et al. 2009;

Alaux et al. 2010; Spleen et al. 2013). These detrimental health factors may be due in part to

landscape simplification limiting the abundance and diversity of floral resources available to

pollinators (Tscharntke et al. 2005; Smart et al. 2016).

A second gap in the literature concerns the extent to which landscape use determines the ways in

which pollinators and plants interact with one another. It is important to understand how

interactions between plants and pollinators may be affected across spatial-temporal scales in

order to preserve the stability of pollination services (Burkle and Alarcón 2011). While it is

understood that land-use intensification negatively affects both plant and pollinator diversity

(Vinson et al. 1993; Williams et al. 2002; Kremen et al. 2002; Potts et al. 2003; Knight et al.

2009; Potts et al. 2010; Garibaldi et al. 2011; Spiesman and Inouye 2013; Habel et al. 2019),

relatively few studies have utilized a network-based approach to consider the ecological impacts

of both landscape composition and configuration on plant-pollinator community structure and

stability (Weiner et al. 2014; Moreira et al. 2015; Tylianakis and Morris 2017; Redhead et al.

2018; Jauker et al. 2019; Lazaro et al. 2020). In an ecological context, network theory has been

utilized to examine how the mutualistic interactions within plant-pollinator communities

influences their structure and in a broader sense, interpret the mechanisms behind biodiversity

and community resilience (Memmott et al. 2004; Bascompte et al. 2006; Blüthgen et al. 2008;

Dupont et al. 2009*a*; Hadley and Betts 2012; Spiesman and Inouye 2013; Soares et al. 2017;

Redhead et al. 2018). Mutualistic networks, such as plant-pollinator networks, tend to exhibit

structural patterns such as nestedness, which demonstrates a degree of interaction redundancy in

the community and is associated with overall community stability (Jordano et al. 2003;

Bascompte et al. 2003;2006). Using a network-based approach can help discern the overall

structure of plant-pollinator communities embedded within varying disturbed landscapes and

their response to resource availability across space and season.

We focus our study in the Northern Great Plains within a region in eastern South Dakota known

as the Prairie Coteau. Historically, the Great Plains stretched across approximately 60 million

hectares in North America, though presently approximately 4% of this temperate grassland

biome remains undisturbed from cropland conversion (Bauman et al. 2016). South Dakota is a

part of a greater region known as the Western Corn Belt where the rate of grassland conversion

to corn and soy agriculture fields reached ~1.0-5.4% annually from 2006 to 2011 (Wright and

Wimberly 2013). Even though agriculture remains a primary land-use in much of the Upper

Midwest, the Prairie Coteau region is unique in that approximately 17% of its original temperate

grassland cover remains intact (Bauman et al. 2016). This provides an ideal study region to

investigate plant-pollinator communities in remnant habitats established within an actively

transforming working landscape. Considering the alarming decline of insect pollinators and

grassland communities, there is a need to understand the community-level impact of habitat

attributes and the surrounding landscape within an increasingly disturbed environment.

Agriculture-dominated landscapes such as the Western Corn Belt will need to coexist with the

remaining patches of conserved natural habitat in order to continue utilizing the ecosystem

services these habitats provide. For example, oil and pulse crops (e.g., sunflower, canola,

legumes), which consist of approximately 1.1% of the landcover in South Dakota in 2019 (Han

et al. 2012), benefit from insect pollination through increased crop yield (Tamburini et al. 2017;

Mallinger et al. 2019). Within the Northern Great Plains, insect pollination increases sunflower

crops by up to 45%, which translates to a regional economic value of $40 million (Mallinger et

al. 2019). Even soybean crops, which are primarily autogamous, have been shown to benefit

from insect pollination through increased crop yield (Chiari et al. 2005; Lautenbach et al. 2012;

Milfont et al. 2013; Cunningham-Minnick et al. 2019). Soybean crops occupy 7.1% of the

landcover in South Dakota and are the second largest agricultural crop in the state behind corn

(landcover of 9.01%) (Han et al. 2012). The need to conserve stable plant-pollinator interactions

extends beyond the boundaries of remnant natural habitats and understanding how compounding

local and landscape variables influence their outcome should be prioritized.

Our overall goal is to understand how plant-pollinator community structure is influenced by

factors acting at various spatial-temporal scales. Rather than focus on the contribution of the

surrounding natural landscape on pollinator services in agricultural fields, we take a novel

approach to address how a mixed-use landscape affects pollinators found in natural habitats. Our

study is sequential. We quantify the effects of habitat attributes and the surrounding landscape on

pollinator diversity and then examine the relationship between pollinator diversity and the

overall structure of the plant-pollinator communities by addressing the following questions: 1)

How do North American temperate grassland attributes (i.e., patch size, floral diversity, latitude,

nectar resources) affect pollinator diversity? 2) Is there a relationship between landscape and

pollinator diversity, and if so, at what scale? Furthermore, does the proportion of land-uses

surrounding North American temperate grasslands influence pollinator diversity, and if so, at

what scale? and 3) How does pollinator diversity influence the overall structure of plant-

pollinator communities? Essentially, given the plant community present at a site, how does

pollinator diversity affect the community structure of interactions? These questions become

especially relevant considering landscape conversion is expected to continue to reduce the

natural landscape (Benton et al. 2003; Bauman et al. 2016; Liu et al. 2019). A better

understanding of plant-pollinator networks and how their structure may be impacted by habitat

or landscape attributes can help future management decisions focus on strategies promoting

resilient plant-pollinator communities and conserve the stability of pollination services.

Materials and Methods

Study area

Formed by glacial uplift, the rolling hills of the Prairie Coteau span roughly 810,000 hectares

across southeast North Dakota, eastern South Dakota, and southwest Minnesota (Bauman et al.

2016). Within South Dakota, this region covers approximately 17 counties and contains a matrix

of land uses ranging from undisturbed grasslands to intensive cropland. Approximately 66% of

eastern South Dakota has some type of crop disturbance history (Bauman et al. 2016). Currently,

undisturbed grassland under protection from future conversion represents about 4% of the total

land base in eastern South Dakota (Bauman et al. 2016). Thus, the grasslands that remain are

fragments nestled within an increasingly disturbed landscape. The Prairie Coteau region

represents a valuable resource for remnant temperate grassland habitat considering 17% of

undisturbed grassland in eastern South Dakota still remains intact (Bauman et al. 2016). We

selected fifteen remnant temperate grassland sites within the Prairie Coteau based on size, local

landscape use, and proximity to other semi-disturbed grasslands (Table 1). Sites ranged in size

from 8 to > 400 hectares and are managed by the US Fish & Wildlife Service (2), The Nature

Conservancy (5), South Dakota Game, Fish & Parks (5) and the city of Brookings (1) with varied

management regimes.

Data collection

Pollinator observation surveys

Between May and October of 2019, surveys were conducted for 30 minutes along 30 x 1 m

transects on days warm enough to allow insect flight and in time periods when pollinators are

expected to be active (15-35º C, between 08:00 - 17:00 hours). We divided the sampling into

three seasons: early (May-June), mid (July-August) and late (September-October). Season

intervals were selected based on consistent flowering phenology shifts found in the plant

communities of the Prairie Coteau. For example, species belonging to the genera *Anemone*

(*Ranunculaceae*), *Viola (Violaceae)*, and *Sisyrinchium (Iridaceae)* predominately bloomed in the

early season, while the mid and late seasons were dominated by species in the *Fabaceae* and

Asteraceae (legume and sunflower families, respectively). Though these two families were

predominately found in both seasons, the mid-season is distinct as this period marked a peak in

the number of families in bloom with approximately six times more families present in our

surveys in comparison to other seasons. Late season was characterized by a distinct shift in floral

composition in which Asteraceae became the most prominent family in all sites with nearly all

other families no longer flowering for the year. We sampled for one year and completed 114

transects.

From our roster of fifteen temperate grassland sites, we randomly sampled each site until

flowering ceased at each location. Location and direction of transects were randomized at each

visit using a list of randomly generated numbers to determine number of steps and cardinal

directions before placing transects down. Transects were geospatial referenced using a Trimble

Geo 7x GPS unit with 1-100 cm accuracy. We walked the entire length of the transect and

recorded all plant-pollinator interactions from within one meter of the transect line on both sides.

We defined pollinators as insect floral visitors that made contact with both the male and female

reproductive parts of the flower, a commonly used criterion (Fenster et al. 2004). We

documented each pollinator and the associated biotically-pollinated plant species when an

interaction occurred. Additionally, we documented pollinator return visits to plants.

Pollinators were identified in situ to family and genus, then to morphospecies in order to

quantify insect diversity. The pollinator observations in our study only focus on diurnal

pollinators, however, this does not present a significant bias in our sampling. Our data set

portrays a robust, representative sample of the plant-pollinator networks in this region

considering only one species (*Silene vulgaris*) detected in our floral surveys (described in the

next section) relies on nocturnal pollination, and this one species was only present in 1 transect

of the 114 sampled. Insect voucher specimens were collected in the field with an aspirator and

net, later identified to lowest taxonomic level and then categorized into functional groups.

Specimens were identified using resources available through discoverlife.org, bugguide.net, and

Key to the Genera of Nearctic Syrphidae (Miranda et al. 2013). Voucher specimens were verified

for sampling completeness using the help of experts and the Severin-McDaniel Insect Research

Collection available at South Dakota State University.

Floral Surveys

Floral surveys were conducted directly after insect pollinator observation surveys along the same

transect with a 1 m^2 quadrat. We placed the quadrat at each meter mark from 0 to 30 m and

surveyed only one side of the transect considering each side mirrored the other with regards to

species composition. Within the quadrat, we documented the presence of each biotically-

pollinated plant species, number of individuals per species, number of flowering units defined as

a unit of one (e.g., Ranunculaceae) or a blossom (e.g., Asteraceae) requiring a small pollinator to

fly in order to access another flowering unit per species, and percent cover within quadrat per

species. Plant voucher specimens were collected and identified using Van Bruggen (1985),

verified with the help of experts (see *Acknowledgements*), and are curated at the C. A. Taylor

Herbarium at South Dakota State University. Digitized plant collections for this study may be

accessed on the Consortium of Northern Great Plains Herbaria (https://ngpherbaria.org/portal/).

To quantify the relationship between sugar resources at a site and pollinator diversity, nectar

sugar content was collected using a handheld sucrose refractometer (Bellingham & Stanley

Eclipse model) for each species during the growing season with a minimum of two samples per

species. Nectar was collected in the field by collecting one flowering unit and compressing it into

the refractometer lens to read volume and sucrose concentration based on the Brix scale. For

composite flowers, the capitulum was first removed from the stem and sectioned into thirds.

Then, a third of the ray and disc florets were compressed into the refractometer. Nectar volume

was calibrated and standardized in lab using a micropipette and handheld sucrose refractometer.

Volumes were categorized as low (<3.5 µl), medium (5.5 µl) and high (7.5 µl) based on how

much liquid covered the main body lens of the refractometer during calibration. For example, a

high liquid volume measuring at approximately 7.5 µl would completely cover the lens. These

standardized volumes were applied to all biotically-pollinated plant species sampled in the field.

Although these nectar measurements are a gross proxy of sucrose considering we are crushing

the flowers to extract our measurements, they provided an efficient, consistent, and standardized

method.

Nectar sugar was estimated by calculating an average nectar sugar concentration for each plant

species and average volume from values recorded over the entire sampling season. We then

estimated sugar for one flowering unit of each species by multiplying average volume and sugar

concentration, and then converting to reflect grams of sugar per liter. Then, we calculated sugar

for the entire plant for each species by multiplying estimated flower sugar and the number of

flowers found on each transect. Finally, we summed all plant sugar found on each transect to

obtain total transect sugar. Nectar data was not recorded for the following species: *Anticlea*

elegans, Helenium autumnale, Grindelia squarrosa, Lobelia spicata, Packera plattensis,

Sisyrinchium montanum, Viola nephrophylla as they were regionally rare and uncommon across

sampling sites. When combined, all regionally rare species were found on approximately 9% of

total transects. However, these species were accounted for in all other measurements

implemented in the floral surveys listed in the previous section.

Pollinator and plant diversity

Pollinator and plant Shannon diversity were calculated using the 'vegan' package version 2.5-6

in R version 3.6.3 (R Core Team 2013; Oksanen et a. 2019). The Shannon index is calculated

using the following formula:

$$\mathbf{H'} = -\sum p_i * \ln p_i$$

Where p_i is the proportions of each species found in a community and ln is the natural log. All

diversity indices were natural log transformed. Shannon is the only diversity index presented in

this study as it accounts for abundance, richness, and evenness. Pollinator diversity was

calculated at the functional, family, genus, and species level by transect and season. Values used

in pollinator diversity did not include return visits recorded during observation surveys.

Likewise, plant diversity was calculated at the family, genus, and species level by transect and

season using number of individuals recorded during floral surveys.

Landscape diversity and proportion

To estimate landscape diversity surrounding North American temperate grassland remnants, we

used QGIS version 2.8.3 – with GRASS and 'rgdal', 'maptools', 'rgeos', 'rasterVis'packages in

R (Bivand and Rundel 2020*a*; Bivand et al. 2020*b*; Bivand and Lewin-Koh 2020*c*; Perpinan

and Hijmans 2020). We obtained a 2019 cropland raster layer (CropScape; CDL) from USDA

Ag Data Commons for South Dakota. We then created a vector data layer for the Prairie Coteau

region from Google Earth Pro. The CDL 2019 raster layer was then clipped onto the Prairie

Coteau layer and warped in order to view cropland use in our focus region with the correct

projection (i.e., WGS 84/ UTM zone 14N). Further, we used QGIS to process and create a vector

shapefile with each transect line collected from the Trimble Geo 7x.

We imported the vector shapefile produced in QGIS into R in order to create buffer layers at

three different scales (500 m, 1000 m, and 3000 m) around each transect using the *gBuffer*

function in the 'rgeos' package (Bivand and Rundel 2020*a*). The output of this function creates a

table with pixel counts corresponding to each land-use in the CDL layer found within each

transect buffer. The land-use pixel values were used to create Shannon diversity indices for each

of our transects. Shannon diversity was calculated using the 'vegan' community ecology package

version 2.5-6 in R. Landscape diversity was measured at the transect level due to the variation

we found surrounding the transects in QGIS even at large scales (i.e., 1000 m and 3000 m).

We used the five most prominent land-uses (i.e., grassland, idle cropland, herbaceous wetlands,

corn and soy) that comprise over 75% of the surrounding landscape to investigate the influence

of land-use proportion on pollinator diversity. We used the same pixel values derived for

landscape diversity to calculate land-use proportion for each transect. Land-use pixels were

saved in Excel and proportion was calculated using simple equations for all scales (i.e., 500 m,

1000 m, 3000 m).

Network analysis

To quantify plant-pollinator community structure, we built quantitative visitation networks for

each site by each of the three seasons. We collapsed all transects and created networks for each

site by season considering that transect level analysis for network-level metrics does not provide

enough interactions to include all sampled transects. By collapsing our samples into one season,

we can include all observations and provide context of the entire community at a site in a given

season in order to understand how network-level metrics are influenced by predictor variables

(i.e., pollinator diversity) in our models. We focus on the mid-season in our results as it is the

only season where all sites were able to be sampled more than once due to flooding conditions

barring access to half of the sites in the early season. Sites were sampled only once during the

late season, however, approximately half of the transects in the late season did not have enough

interactions to derive network metrics. Nevertheless, we performed network analyses for the

early and late seasons with halved sample sizes and found similar results to those found in the

mid-season (Supplemental tables 1-3). The Deer Creek site was excluded from all network

analyses as there were too few interactions even when transects were collapsed to generate

network-level metrics.

Networks were constructed using a matrix of interactions between plants and pollinators

including unique and return visits recorded during observation surveys. Documenting return

visits allows us to quantify plant-pollinator communities using weighted network values that also

account for visitation frequency. For each network, we calculated network specialization (H2'),

connectance and nestedness. The H2' specialization metric measures the degree of specialization

at the community level (Blüthgen et al. 2006;2008) and is measured on a scale from 0 to 1,

where a value approaching one indicates a network with increased selective interactions.

Network-level specialization incorporates the interaction frequency between nodes and has been

established as a more robust metric than species-level specialization given that network-level

specialization is independent of network size (Blüthgen et al. 2006). Connectance represents the

proportion of realized links out of all possible interactions within a network and is measured on a

scale from 0 to 1. Connectance is used to describe the degree of generalization and interaction

complexity within a network with a value approaching one indicating a more connected network

with increased generalization (Dunne et al. 2002; Cusser and Goodell 2013). Nestedness

describes a pattern of asymmetry in the interactions of a network (Bascompte et al. 2003).

Within a perfectly nested network, specialist species would interact with a subset of species that

interact with generalists; thus, increasing the redundancy of interactions. Nestedness is measured

on a scale from 0 to 100 with 0 describing perfect nestedness (Rodriguez-Girones & Santamaría

2006). Trends were analyzed using linear regressions. Using these network parameters, we can

determine the influence of certain habitat attributes and the surrounding landscape on the

preservation of specialized interactions and overall resilience of the community. All network

metrics were calculated using the 'bipartite' package version 2.15 in R (Dormann et al. 2009).

Statistical Analyses

Relationship between landscape diversity, North American temperate grassland attributes and

pollinator diversity.

To quantify the relationship between pollinator diversity and attributes found within and outside

of North American temperate grasslands, we used linear mixed effects models with site as the

random effect. We used the packages 'lme4' version 1.1-23 and 'nlme' version 3.1-144 in R for

the linear mixed effects models (Bates et al. 2015; Pinheiro et al. 2020). All variables in the

model are at the transect level. The full model including both fixed and random effects was:

Pollinator Genus Diversity ~ scale(Landscape Diversity at 500 m) + scale(Landscape Diversity

at1000 m) + scale(Landscape Diversity at 3000 m) + Plant Species Diversity + Transect Sugar +

352   (1|Site)

We then used stepwise-backward variable selection to simplify the model. The dropterm

function in the 'MASS' package version 7.3-51.5 was used because it considers each variable

individually and we can specify what test to use to compare the initial model as well as each of

the possible alternative models with one less variable (Venables and Ripley 2020). For model

assessment, we used AIC (Akaike Information Criterion) values and chi-square as criteria when

dropping each variable one at a time from the full model stated above. Size and latitude were not

included in the model to avoid collinearity. Thus, we used simple linear regressions for both size

and latitude separately with pollinator diversity as the response variable.

Relationship between landscape proportions, North American temperate grassland attributes

and pollinator diversity.

To quantify the relationship between landscape proportion, North American temperate grassland

attributes and pollinator diversity, we used linear mixed effects models with site and season as

the random effects. We used the packages 'lme4' and 'nlme' in R for the linear mixed effects

models. All variables in the model are on the transect level. The full model including both fixed

and random effects was:

Pollinator Genus Diversity ~ Corn Proportion at 500 m + Corn Proportion at 1000 m + Corn

Proportion at 3000 m + Soy Proportion at 500 m + Soy Proportion at 1000 m + Soy Proportion at

3000 m + Grassland Proportion at 500 m + Grassland Proportion at 1000 m + Grassland

Proportion at 3000 m + Herbaceous Wetland Proportion at 500 m + Herbaceous Wetland

Proportion at 1000 m + Herbaceous Wetland Proportion at 3000 m + Idle Cropland Proportion at

500 m + Idle Cropland Proportion at 1000 m + Idle Cropland Proportion at 3000 m + Plant

Species Diversity + Transect Sugar + (1|Site) + (1|Season)

We implemented stepwise-backward variable selection to simplify this model. We incorporated

the same model selection process that was applied to the landscape diversity model by using the

dropterm function. For model assessment, we used AIC (Akaike Information Criterion) values

and chi-square as criteria when dropping each variable one at a time from the full model stated

above.

Relationship between pollinator diversity and network structure.

We used generalized linear models with a Gaussian distribution to determine the relationship

between network structure and pollinator diversity due to some of our response variables being

non-normally distributed. All variables were scaled to season level. Univariate regression models

were used for each network parameter with pollinator diversity as the predictor variable and each

network parameter as the response variable.

Results

North American temperate grassland attributes and the surrounding landscape

Landscape Shannon diversity varied from 0.47 to 2.10 across transects and spatial scales (Fig. 1).

The most common land-use pixels found within the 3000 m buffers were grassland, herbaceous

wetlands, fallow/idle cropland, corn, and soybean fields. When averaged across all sites at the

3000 m scale, grassland pixels covered 39%, fallow/idle cropland covered 11.5%, corn fields

covered 9%, soybean fields covered 9%, and herbaceous wetlands covered 8% of 3000 m

buffers. When combined, all five land-uses accounted for 76.5% of the landscape within 3000 m

of our sites.

Pollinator community

Among all 15 sites, 79 genera of pollinators representing 10 functional groups and 45 families

were collected throughout the sampling season (see Appendix A for full list). 77% of total

samples collected and observed were identified to genus. Samples that were not identified to

genus were identified to the next taxonomic level (family 23%) and given a morphospecies

classification which was used for pollinator genus analyses. Though we recorded morphospecies

in the field, we found genus to be the lowest, most robust taxonomic level in the data set that

could be identified with accuracy. Insect pollinators that could not be identified to genus were

placed in a catch-all genus that consisted of the first five letters of their family name. For

example, for a fly pollinator in the family Muscidae, we created a genus named Gen_Musci in

the dataset in order to include these visitors in the analyses. We found pollinator genus diversity

to be correlated with functional group diversity and family diversity (Supplemental Table 4).

Thus, we focus on pollinator genus diversity in our analyses to reduce the number of analyses.

Shannon diversity of pollinator groups ranged from 0 to 2.2 among sites throughout all three

seasons (Fig. 2). The majority of observations in the early season where diversity ranged from 0

– 1.7 included Syrphidae (36%), Muscidae (20%), Chloropidae (14.5%), and Halictidae (14%)

(Supplemental Fig. 1). Mid-season diversity ranged from 0 – 2.2 with Apidae (27%), Syrphidae

(25.7%), Cantharidae (12.5%), and Halictidae (11%) comprising the majority of observations

(Supplemental Fig. 2). Late season diversity ranged from 0 – 1.8 with Syrphidae and Halictidae

encompassing 58 and 28 percent of all observations, respectively (Supplemental Fig. 3).

Plant community

Among all 15 sites, 87 species representing 24 families and 61 genera were collected throughout

the growing season (see Appendix B for full list). Shannon diversity of biotically-pollinated

plants ranged from 0 to 2 among sites throughout all three seasons (Fig. 3). Early season

diversity ranged from 0 – 2 with *Anemone canadensis*, *Gallium boreali* and *Fragaria virginiana*

as the most common species found on the transects (Supplemental Fig. 4). Mid-season diversity

ranged from 0 – 2 with *Melilotus albus, Anemone canadensis,* and *Amorpha canescens* as the

most common species found on the transects (Supplemental Fig. 5). Late season diversity ranged

from 0 – 1.5 with *Symphyotrichum lanceolatum*, *Symphyotrichum ericoides* and *Heliopsis*

helianthoides as the most common species found on the transects (Supplemental Fig. 6). We

found that plant species diversity was correlated with family diversity and genus diversity

(Supplemental Table 4). Thus, we focus on plant species diversity in our analyses to reduce the

number of analyses.

North American temperate grassland attributes and landscape diversity effects on pollinator

diversity

After model assessment, we found the final, simplified model explaining pollinator genus

diversity included transect sugar as the only predictor variable (Table 2). Transect sugar

demonstrates a significant positive relationship with pollinator genus diversity (Table 3). We

found no relationship between pollinator diversity and North American temperate grassland size

or latitude.

North American temperate grassland attributes and land-use proportion effects on pollinator

diversity

We found the final, simplified model explaining pollinator genus diversity included corn

proportion at 500 m, soy proportion at 1000 m, herbaceous wetland proportion at 1000 m, and

transect sugar (Table 4). All predictor variables in the model demonstrated a positive relationship

with pollinator genus diversity (Table 5).

Plant-pollinator network analysis

From May through October, we observed 518 unique plant-pollinator interactions and a total of

9,877 observations of pollinators visiting plants. The most common floral visitors throughout the

entire sampling period were Syrphidae (31%), Apidae (19%), and Halictidae (13%). The plant

species with the most interactions throughout the entire sampling period include *Melilotus albus*

(20.7%), *Amorpha canescens* (10.6 %), *Euphorbia esula* (8.5%), and *Achillea millefolium*

(6.7%).

Network metrics and pollinator genus diversity

Network-level specialization (H2') ranged from 0.35 – 0.81 between temperate grassland sites

during mid-season. Pollinator genus diversity had a significant positive relationship with

network-level specialization ($p < 0.002$, $t = 4.313$, slope coefficient: 0.304 ± 0.072) (Fig. 4).

Connectance ranged from 0.10 – 0.29 between temperate grassland sites at the mid-season.

Pollinator genus diversity demonstrated a negative relationship with connectance ($p < 0.04$, $t = -$

2.39, slope coefficient: -0.077 ± 0.032) (Fig. 4). Nestedness ranged from 9 – 25 between

temperate grassland sites at the mid-season. We found no significant relationship between

pollinator genus diversity and nestedness ($p < 0.50$, $t = -0.70$, slope coefficient: -3.015 ± 4.29).

Discussion

Our study expands on previous research (Aizen et al. 2003; Ashworth et al. 2004; Heard et al.

2007; Carvalheiro et al. 2010; Ferreira et al. 2013; Boyle et al. 2019; Ponisio et al. 2019; Twerd

and Banaszak-Cibicka 2019) documenting the influence of landscape modification and habitat

fragmentation on the persistence of plant-pollinator communities in natural habitats through

changes in pollinator availability and diversity. We further develop these approaches by

investigating a combination of both local habitat attributes (i.e., patch size, latitude, plant species

diversity, nectar resources) and landscape effects on pollinator diversity and linking the effects

of pollinator diversity to the structure of plant-pollinator communities using a network-based

approach. Our results indicate the most important habitat attribute predicting pollinator genus

diversity in the Prairie Coteau region is the amount of floral sugar found at remnant temperate

grassland sites as extrapolated from transect sampling. Additionally, we found landscape

diversity had no effect on pollinator genus diversity. Instead, the proportion of certain land-uses

surrounding remnant temperate grassland sites had a significant effect on pollinator genus

diversity. In turn, pollinator genus diversity demonstrated a positive effect for overall network

specialization and consequentially a negative effect on connectance. Below, we discuss the local

and landscape effects on pollinator diversity followed by the effects of pollinator diversity on

overall community structure and how our findings may influence future planning for the

conservation of temperate grassland remnants and pollinator diversity in the Northern Great

Plains.

The role of local resources and North American temperate grassland attributes on pollinator

diversity.

Nectar resources (i.e., plant sugar) were an important determinant of pollinator diversity at

remnant temperate grassland sites in the agriculturally dominated landscape of the Northern

Great Plains. These findings indicate floral abundance and nectar rewards influence pollinator

behavior, also observed by others (Potts et al. 2009; Fowler et al. 2016), especially in disturbed,

fragmented habitats (Potts et al. 2003; Stang et al. 2006; Ouvrard et al. 2018). The common, non-

native forb *Melilotus albus* recorded in 10 of our 15 sites, had greater average nectar

contribution per transect (based on number of flowers and sugar concentration per flower) and

comprised 20% of total interactions throughout the entire sampling season, making it a core node

in our plant-pollinator networks. Our findings with *M. albus* are also consistent with other

studies demonstrating bees will heavily utilize non-native plants in disturbed habitats, even

though they do not exhibit a preference for non-native plants over native plants (Hinners et al.

2009; Williams et al. 2011). Considering the impacts non-native plant species can have on

pollinators, it is imperative we consider the role that non-native plants play in shaping

community interactions, particularly in heavily disturbed and modified landscapes such as those

dominated by agriculture. The role of non-native pollinator species in plant-pollinator networks

should also be examined considering honey bees could compete with native pollinators for

foraging resources (Sugden et al. 1996; Thomson 2004). In the Northern Great Plains, a study

conducted from 2015-2017 found both honey bees and wild bees visiting *Melilotus* with honey

bees utilizing *Melilotus* more than wild bees (Otto et al. 2020). However, honey bees in our

study were only present in 7 out of 15 temperate grassland sites and made up 12% of total

interactions. Honey bee observations only occurred due to presence of commercial honey bee

hives within 1 km of our sites and overall, account for a low frequency of visitation in networks.

Melilotus is considered an attractive resource to insect pollinators because of their high nectar

production, strong scent, and abundance of flowers produced with up to 350,000 flowers per

plant (Royer and Dickinson 1999; Spellman et al. 2015). As *Melilotus* continues to establish

within the Northern Great Plains, we can expect it to become a prominent, stabilizing anchor in

plant-pollinator networks that could facilitate increased pollinator visitation and diversity

(Tepedino and Griswold 2008; Spellman et al. 2015). Other invasive species in temperate

grasslands such as *Cirsium arvense* have been found to influence the topology of plant-pollinator

networks by decreasing modularity, which is associated with greater network stability; however,

the effects of non-native plant species on native plant reproduction were not investigated (Larson

et al. 2016). Despite the warranted concern over the impacts of invasive plant species on native

plant communities (i.e., reduction of pollen quality and quantity) (Waser 1983; Campbell 1985;

Brown et al. 2002; Morales and Traveset 2008; Kandori et al. 2009), some studies have shown

that *Melilotus* can benefit community interactions by increasing pollinator visitation, diversity,

and abundance (Morales and Traveset 2005; Tepedino and Griswold 2008; Spellman et al.

2015). In the Northern Great Plains, Otto et al. (2020) found both species of *Melilotus* were

among the three most abundant flowering species in their transects, corresponding with our

findings that *Melilotus* is a core node in the plant-pollinator networks in our study region.

Landscape effects on pollinator diversity

Landscape diversity did not exhibit a relationship with pollinator genus diversity at any

scale (i.e., 500 m, 1000 m, and 3000 m). However, we found that increased proportions of

corn fields at 500 m, soybean fields at 1000 m, and herbaceous wetlands at 1000 m demonstrated

a positive effect on pollinator genus diversity. Herbaceous wetlands adjacent to perennial

grasslands, specifically those found in the Upper Midwest and Prairie Coteau regions, are

valuable resources for insect pollinators and help increase native bee diversity and

abundance (Vickruck et al. 2019; Begosh et al. 2020). The positive relationship between

pollinator genus diversity and proportion of herbaceous wetlands in the Prairie Coteau may be

related to the resources provided by the semi-natural habitats within the landscape.

Conversely, intensive agricultural fields (i.e., corn and soy fields) do not provide ample foraging

or nesting resources for pollinators throughout the growing season and have been documented

to negatively affect the hive health of certain pollinators (e.g., bumblebees) by reducing the

quality and diversity of pollen available (Hass et al. 2018). The rise in pollinator genus diversity

as intensive agriculture increases at various scales (i.e., 500 m and 1000 m) in our study could

indicate an oasis effect occurring where insect pollinators are crossing substantial distances to

forage on the ample nectar resources provided by temperate grassland patches in the Northern

Great Plains (Bock et al. 2008; Gotlieb et al. 2011). Approximately 50% of the interactions in

our plant-pollinator networks were dominated by syrphid flies and apid bees, which are highly

mobile and utilize a variety of habitat types, indicating the insect pollinators in our observed

networks may have traveled considerable distances to utilize the semi-natural reservoirs

embedded within this cropland matrix. (Morris 1993; Sommaggio et al. 1999; Beekman and

Ratneiks 2000; Klecka et al. 2018; Jauker et al. 2019). These findings are also consistent

with other studies such as Winfree et al. (2007) and Hoehn et al. (2008) demonstrating that

landscapes dominated by intensive agriculture can still maintain relatively diverse pollinator

communities provided patches of floral resources were present.

How does pollinator diversity influence the overall plant-pollinator community structure?

While other studies have focused on the botanical side of plant-pollinator networks in order to

investigate network structure (Larson et al. 2016; Robinson et al. 2018; Cunningham-Minnick

and Crist 2020; Hernandez-Castellano et al. 2020), we instead use pollinator diversity

as our predictor variable to investigate network structure in perennial grasslands in the Northern

Great Plains. The availability and abundance of nectar resources influence pollinator behavior

which can potentially impact community structure on a larger scale. For example, Nottebrock et

al. (2017) found that pollinator visitation rates strongly depended on phenological variation of

site-scale sugar amounts. Further, the seed sets of their focal plants increased with both site-scale

sugar amounts and nectar sugar amounts of conspecific neighbors. While their study did not

investigate network structure, it demonstrates that the quantity and availability of nectar

resources can have broader community impacts for both plants and pollinators. Our results

indicate pollinator genus diversity influences connectance and network specialization (H2') in

reinforcing ways, with connectance decreasing and specialization increasing with greater

pollinator genus diversity. We did not observe any relationship between nestedness and

pollinator genus diversity, however we did find that networks were nested, as expected of

asymmetric, mutualistic networks which contain hubs (i.e., highly linked species) that form a

central core of interactions linking other species within the network (Bascompote et al. 2003;

Jordano et al. 2003). Networks in our study fell within a range which indicates a nested pattern

(9-25) that has also been observed by others within working grasslands (Vanbergen et al. 2014;

Jauker et al. 2019).

Our findings align with predictions by Montoya and Yvon-Durocher (2007) who suggested that

interspecific competition among pollinators may be driving network specialization more so than

floral diversity (see also Fründ et al. 2010; Valdovinos et al. 2016). Network-level specialization

can be beneficial for plant-pollinator communities as it may allow for increased efficiency of

pollen transfer which can reflect a higher resilience and stability in the community (Arceo-

Gomez 2020). Likewise, the negative relationship between connectance and pollinator genus

diversity suggests plant-pollinator networks in the Prairie Coteau region can be prone to being

less complex and connected (Dunne et al. 2002). Nested networks displaying a higher degree

of connectance are considered more resilient and stable, making them important considerations

for conservation value (Memmott et al. 2004; Okuyama and Holland 2008; Thébault and

Fontaine 2010). The nested pattern found in the networks indicates a degree of interaction

redundancy that likely contributes to community stability (Bascompte et al. 2003; Nielsen

and Bascompte 2007). Nonetheless, nestedness does not seem to be influenced by the diversity

of pollinators in these remnant temperate grassland communities. Overall, our results suggest

that temperate grasslands with increased pollinator diversity can maintain more specialized

plant-pollinator interactions that can potentially promote resilient plant-pollinator

communities in the Prairie Coteau region.

Conservation implications

Our results demonstrate that managing pollinator resources (sucrose availability) within

remaining habitats and the composition of the surrounding landscape should be carefully

considered when seeking to promote pollinator diversity and preserve pollination services. These

results agree with other studies (Potts et al. 2005; Tscharntke et al. 2005; Potts et al. 2009; Heard

et al. 2007; Cusser and Goodell 2013; Moreira et al. 2015; Smart et al. 2016), which identified

how pollinator populations are influenced by a series of interacting factors (i.e., landscape

configuration and floral community composition), rather than a singular attribute such as patch

size. Landscape conversion in agricultural regions is not likely to diminish soon, making it

imperative that we recognize the conservation value of the remaining habitat fragments such as

those found in the Prairie Coteau.

Our study demonstrated plant species with high nectar rewards promoted pollinator diversity

within temperate grasslands of the Northern Great Plains. While we cannot conclude whether

non-native nectar resources provide more benefits to the overall health of pollinators in this

region, future research should address the effects of native vs. non-native floral resources on

pollinator health and network community structure in disturbed, fragmented habitats. Several

studies (Lopezaraiza–Mikel et al. 2007; Aizen et al. 2008; Bartomeus and Santamaría 2008;

Padron et al. 2009) have indicated that non-native plants may alter the structure of plant-

pollinator networks by monopolizing the interactions of the most generalist pollinators. Recently,

Hernandez-Castellano et al. (2020) and Cunningham-Minnick and Crist (2020) found that the

integration of non-native plant species into plant-pollinator networks could dilute pollination

services for native plants, with potential risks for pollinators that may not be able to use invasive

floral resources. Thus, it would be valuable to apply this question to long-term monitoring of

non-native plant species and their pollinators to determine if there will be any consequential

changes to network structure over time. Long-term network studies are not common (Burkle and

Alarcón 2011) and the matter of network stability in agriculture-dominated landscapes should be

further investigated to ascertain how local and landscape changes affect plant-pollinator

networks on a wider temporal scale.

While current evidence indicates network metrics remain stable across years (Alarcón et al.

2008; Petanidou et al. 2008 ; Burkle and Irwin 2009 ; Dupont et al. 2009*b*), consideration should

be made for temporal shifts in plant and pollinator communities that could occur due to climate

change (Roy and Sparks 2000; Fitter and Fitter 2002; Forister and Shapiro 2003; Bartomeus et

al. 2011; Calinger et al. 2013; CaraDonna et al. 2014). Changes in flowering time and pollinator

activity could result in temporal mismatching between actors in a network, potentially shifting

network structure and stability (Albrecht et al. 2010; Morton and Rafferty 2017). Network

interactions are considered fairly dynamic. Overall network structure has shown to remain stable

even when network 'rewiring' occurs as a result of shifts in actors due to changes in abundance

of plants or pollinators between years (Petanidou et al. 2008; Dupont et al 2009*b*); however,

most multi-year network studies only extend to 2-4 years (Burkle and Alarcón 2011). As

ecosystems continue to experience habitat disturbance on various scales along with climate

change, it would be valuable to understand if long-term changes to species compositions would

impact the structure of future plant-pollinator networks. Additionally, further research on

understanding the influence of land management strategies (e.g., prescribed burning and grazing)

within remnant, semi-natural habitats could provide insight on resource changes and availability

at a local scale.

Native pollinators provide valuable ecosystem services for both native plant communities and

agricultural fields by increasing plant reproductive success and crop yield (Fontaine et al. 2005;

Albrecht et al. 2007; Albrecht et al. 2012; Rogers et al. 2014). Increasing pollinator abundance

and diversity while maintaining specialized interactions can promote stability of this essential

ecosystem service by enhancing the overall visitation and pollination of the plant communities

present (Klein et al. 2007; Hoehn et al. 2008; Albrecht et al. 2012). Conserving these pollination

services is important for sustaining future food security of the 35% of agricultural crops that rely

on pollination globally, in addition to local crops such as sunflower and soybean which are

economically valuable for the Northern Great Plains region (Klein et al. 2007; Vanbergen et al.

2013; Mallinger et al. 2019). Additionally, the protection and maintenance of ecosystem services

(Costanza et al. 1998) directly supports ecosystem function (Peterson et al. 2010). Land

managers should bear in mind the influence of nectar rewards on pollinator movement and

behavior, and develop decisions focused on providing ample resources within the

landscape. Furthermore, the conservation of other semi-natural habitats (i.e., herbaceous

wetlands), which provide resources beyond nectar rewards should be prioritized to promote

diverse pollinator communities, at least in the Northern Great Plains. The oasis effect

demonstrated in our study will likely only intensify as time passes and the role of habitat

reservoirs, like the temperate grassland remnants, will become increasingly important as oases

for pollinator communities in agriculture-dominated regions.

ACKNOWLEDGMENTS

South Dakota State University and the Prairie Coteau region is located on the ancestral territory of the Oceti Sakowin [oh-CHEH-tee SHAW-koh-we], meaning Seven Council Fires which is the proper name for the people commonly called Sioux. The tribal alliance made up of individual bands, of the Seven Council Fires is based on kinship, location, and dialects: Santee-Dakota, Yankton-Nakota and Teton-Lakota. The seven tribes now occupy nine reservations in South Dakota: Cheyenne River Sioux Tribe, Crow Creek Sioux Tribe, Flandreau Santee Sioux Tribe, Lower Brule Sioux Tribe, Oglala Sioux Tribe, Rosebud Sioux Tribe, Sisseton-Wahpeton Oyate, Standing Rock Sioux Tribe and Yankton Sioux Tribe.

We thank S. Stiles, J. Gelderman, T. Wallner, N. Petersen, E. Baier, and S. Daniels for all of their assistance during the field season. Thank you to J. Purintun for providing botanical expertise. We would also like to thank Drs. G. Hatfield and H. Moradi for their analytical advice.

Thank you to all the landowners and managers that provided permissions and advice on site selection for the following: SD State Oak Lake Field Station, South Dakota Game Fish and Parks, U.S. Fish and Wildlife Services, The Nature Conservancy, and the City of Brookings. Finally, this project would not have been possible without support from several Hatch grants and the North Central Sun Grant Initiative (USDA/DOE) SA1500640.

Literature Cited

Aizen, M. A., and P. Feinsinger. 2003. Bees not to be? Responses of insect pollinator faunas and flower pollination to habitat fragmentation. *In* G. A. Bradshaw and P. A. Marquet [eds.], How landscapes change, 111–129. Springer-Verlag Berlin, Germany. Doi:10.1007/978-3-662-05238-9_7

Aizen, M. A., C. L. Morales, and J. M. Morales. 2008. Invasive mutualists erode native pollination webs. PloS Biology, 6:396–403. Doi:10.1371/journal.pbio.0060031

Alarcón, R., N. M. Waser, and J. Ollerton. 2008. Year-to-year variation in the topology of a plant – pollinator interaction network. Oikos, 117:1796 – 1807. Doi:10.1111/j.0030-1299.2008.16987.x

Alaux, C., J. L. Brunet, C. Dussaubat, F. Mondet, S. Tchamitchan, M. Cousin, J. Brillard, A. Baldy, L. P. Belzunces, and Y. Le Conte. 2010. Interactions between Nosema microspores and a neonicotinoid weaken honey bees (*Apis mellifera*). Environmental Microbiology, 12:774-782. Doi:10.1111/j.1462-2920.2009.02123.x

Albrecht, M., P. Duelli, C. Müller, D. Kleijn, and B. Schmid. 2007. The Swiss agri-environment scheme enhances pollinator diversity and plant reproductive success in nearby intensively managed farmland. Journal of Applied Ecology, 44:813-822. Doi:10.1111/j.1365-2664.2007.01306.x

Albrecht, M., M. Riesen , and B. Schmid. 2010. Plant – pollinator network assembly along the chronosequence of a glacier foreland. Oikos, 119:1610 – 1624. Doi: 10.1111/j.1600-0706.2010.18376.x

Albrecht, M., B. Schmid, Y. Hautier, and C. B. Müller. 2012. Diverse pollinator communities enhance plant reproductive success. Proceedings of the Royal Society B: Biological Sciences, 279:4845-4852. Doi: 10.1098/rspb.2012.1621

Arceo-Gómez, G., D. Barker, A. Stanley, T. Watson, and J. Daniels. 2020. Plant-pollinator network structural properties differentially affect pollen transfer dynamics and pollination success. Oecologia, 192:1037–1045. Doi:10.1007/s00442-020-04637-5

Ashworth, L., R. Aguilar, L. Galetto, and M. A. Aizen. 2004. Why do pollination generalist and
specialist plant species show similar reproductive susceptibility to habitat fragmentation?
Journal of Ecology, 92:717-719. Doi: 10.1111/j.0022-0477.2004.00910.x

Bartomeus, I., V. Montserrat, and L. Santamaría. 2008. Contrasting effects of invasive plants in
plant–pollinator networks. Oecologia, 155:761-770. Doi:10.1007/s00442-007-0946-1

Bartomeus, I., J. S. Ascher, D. Wagner, B. N. Danforth, S. Colla, S. Kornbluth, and R. Winfree.
2011. Climate-associated phenological advances in bee pollinators and bee-pollinated
plants. Proceedings of the National Academy of Sciences, 108:20645-20649.
Doi:10.1073/pnas.1115559108

Bascompte, J., P. Jordano, C. J. Melián, and J. M. Olesen. 2003. The nested assembly of plant–
animal mutualistic networks. Proceedings of the National Academy of Sciences,
100:9383-9387. Doi:10.1073/pnas.1633576100

Bascompte, J., P. Jordano, and J. M. Olesen. 2006. Asymmetric coevolutionary networks
facilitate biodiversity maintenance. Science, 312:431-433. Doi: 10.1126/science.1123412

Bates, D., M. Maechler, B. Bolker, and S. Walker. 2015. Fitting Linear Mixed-Effects Models
Using lme4. Journal of Statistical Software, 67:1-48. Doi:10.18637/jss.v067.i01.

Bauman, P., J. Blastick, C. Grewing, and A. Smart. 2016. Quantifying undisturbed land on South
Dakota's Prairie Coteau. A report to The Nature Conservancy from South Dakota State
University based on the Prairie Coteau boundary as defined by the April 30, 2010 TNC
National Fish and Wildlife Foundation Business Plan "Conserving and Restoring
Tallgrass Prairie: Prairie Coteau, South Dakota and Minnesota".

Beekman, M., and F. L. W. Ratnieks. 2000. Long-range foraging by the honeybee, *Apis mellifera*
L. Functional Ecology, 14:490–496. Doi:10.1046/j.1365-2435.2000.00443.x

Begosh, A., L. M. Smith, S. T. McMurry, and J. P. Harris. 2020. Influence of the Conservation
Reserve Program (CRP) and playa wetlands on pollinator communities in the Southern
High Plains, USA. Journal of Environmental Management, 256, 109910.
Doi:10.1016/j.jenvman.2019.109910

Benton, T. G., J. A. Vickery, and J. D. Wilson. 2003. Farmland biodiversity: is habitat
heterogeneity the key? Trends in Ecology & Evolution, 18:182-188. Doi:10.1016/s0169-
5347(03)00011-9

Bivand, R., and C. Rundel. 2020*a*. rgeos: Interface to Geometry Engine -Open Source ('GEOS').
R package version 0.5-5.https://CRAN.R-project.org/package=rgeos

Bivand, R., T. Keitt and B. Rowlingson. 2020*b*. Rgdal: Bindings for the 'Geospatial' Data
Abstraction Library. R package version 1.5-18.https://CRAN.R-
project.org/package=rgdal

Bivand, R., and N. Lewin-Koh. 2020*c*. Maptools: Tools for Handling Spatial Objects. R package
version 1.0-2. https://CRAN.R-project.org/package=maptools

Blüthgen, N., F. Menzel, and N. Blüthgen. 2006. Measuring specialization in species interaction
networks. BMC ecology, 6:9. doi:10.1186/1472-6785-6-9

Blüthgen, N., J. Fründ, D. P. Vázquez, and F. Menzel. 2008. What do interaction network
metrics tell us about specialization and biological traits. Ecology, 89:3387-3399.
doi:10.1890/07-2121.1

Bock, C. E., Z. F. Jones, and J. H. Bock. 2008. The oasis effect: response of birds to exurban
development in a southwestern savanna. Ecological Applications, 18:1093-1106.
doi:10.1890/07-1689.1

Boyle, N. K., and T. L. Pitts-Singer. 2019. Assessing blue orchard bee (*Osmia lignaria*)
propagation and pollination services in the presence of honey bees (*Apis mellifera*) in
Utah tart cherries. PeerJ, 7:e7639. doi:10.7717/peerj.7639

Brown, B.J., R. J. Mitchell, and S. A. Graham. 2002. Competition for pollination between an
invasive species (purple loosestrife) and a native congener. Ecology, 83:2328-2336.
doi:10.1890/0012-9658(2002)083[2328:cfpbai]2.0.co;2

Burkle, L. A., and R. Alarcón. 2011. The future of plant–pollinator diversity: understanding
interaction networks across time, space, and global change. American Journal of Botany,
98:528-538. doi:10.3732/ajb.1000391

Burkle, L., and R. Irwin. 2009. The importance of interannual variation and bottom-up nitrogen
770   enrichment for plant – pollinator networks. Oikos, 118:1816 – 1829. doi: 10.1111/j.1600-
771   0706.2009.17740.x,

772   Calinger, K. M., S. Queenborough, and P. S. Curtis. 2013. Herbarium specimens reveal the
773   footprint of climate change on flowering trends across north-central North America.
774   Ecology Letters, 16:1037–1044. doi: 10.1111/ele.12135

775   Campbell, D.R. 1985. Pollinator sharing and seed set of Stellaria pubera: competition for
776   pollination. Ecology, 66:544-553. doi:10.2307/1940403

777   CaraDonna, P. J., A. M. Iler, and D. W. Inouye. 2014. Shifts in flowering phenology reshape a
778   subalpine plant community. Proceedings of the National Academy of Sciences, USA 111:
779   4916-4921. doi:10.1073/pnas.1323073111

780   Carvalheiro, L. G., C. L. Seymour, R. Veldtman, and S. W. Nicolson. 2010. Pollination services
781   decline with distance from natural habitat even in biodiversity-rich areas. Journal of
782   Applied Ecology, 47:810-820. doi:10.1111/j.1365-2664.2010.01829.x

783   Chiari, W.C., V.D.A.A.D. Toledo, M.C.C. Ruvolo-Takasusuki, A.J.B.D. Oliveira, E.S. Sakaguti,
784   V.M. Attencia, F.M. Costa, and M.H. Mitsui. 2005. Pollination of soybean (Glycine max
785   L. Merril) by honeybees (*Apis mellifera L.*). Brazilian archives of biology and
786   technology, 48:31-36. doi:10.1590/s1516-89132005000100005

787   Costanza, R., R. d'Arge, R. De Groot, S. Farber, M. Grasso, B. Hannon, K. Limburg, S. Naeem,
788   R.V. O'Neill, J. Paruelo and R.G. Raskin. 1998. The value of ecosystem services: putting
789   the issues in perspective. Ecological economics, 25:67-72.

790   Cox-Foster, D.L., S. Conlan, E. C. Holmes, … D. M. Geiser, and V. Martinson. 2007. A
791   metagenomic survey of microbes in honeybee colony collapse disorder. Science,
792   318:283-287. doi:10.1126/science.1146498

793   Cox-Foster, D. L., and D. VanEngelsdorp. 2009. Saving the honeybee. Scientific American,
794   300:40-47. doi:10.1038/SCIENTIFICAMERICAN0409-40

Cunningham-Minnick, M.J., Peters, V.E. and Crist, T.O., 2019. Nesting habitat enhancement for
wild bees within soybean fields increases crop production. Apidologie, 50:833-844.
doi:10.1007/s13592-019-00691-y

Cunningham-Minnick, M. J., and T. O. Crist. 2020. Floral resources of an invasive shrub alter
native bee communities at different vertical strata in forest-edge habitat. Biological
Invasions, 22:2283-2298. doi:10.1007/s10530-020-02248-y

Cusser, S., and K. Goodell. 2013. Diversity and distribution of floral resources influence the
restoration of plant–pollinator networks on a reclaimed strip mine. Restoration Ecology,
21:713-721. doi:10.1111/rec.12003

Dormann, C. F., J. Fruend, B. Gruber, and M. C. F. Dormann. 2009. Package bipartite. version
2.15

Dunne, J. A., R. J. Williams, and N. D. Martinez. 2002. Food-web structure and network theory:
the role of connectance and size. Proceedings of the National Academy of Sciences,
99:12917-12922. doi:10.1073/pnas.192407699

Dupont, Y. L., and J. M. Olesen. 2009*a*. Ecological modules and roles of species in heathland
plant-insect flower visitor networks. Journal of Animal Ecology, 78:346-353.
doi:10.11/j.l365-2656.2008.01501.x

Dupont, Y. L., B. Padrón , J. M. Olesen, and T. Petanidou. 2009*b*. Spatio-temporal variation in
the structure of pollination networks. Oikos, 118:1261 – 1269. doi: 10.1111/j.1600-
0706.2009.17594.x

Fahrig, L., J. Baudry, L. Brotons, F. G. Burel, T. O. Crist, R. J. Fuller, C. Sirami, G. M.
Siriwardena, and J. L Martin. 2011. Functional landscape heterogeneity and animal
biodiversity in agricultural landscapes. Ecology Letters, 14:101-112. doi:10.1111/j.1461-
0248.2010.01559.x

Fenster, C. B., W. S. Armbruster, P. Wilson, M. R. Dudash, and J. D. Thomson. 2004.
Pollination syndromes and floral specialization. Annu. Rev. Ecol. Evol. Syst., 35:375-
403. doi:10.1146/annurev.ecolsys.34.011802.132347

Ferreira, P. A., D. Boscolo, and B. F. Viana. 2013. What do we know about the effects of
landscape changes on plant–pollinator interaction networks? Ecological Indicators,
31:35-40. doi:10.1016/j.ecolind.2012.07.025

Fitter, A. H. and R. S. R. Fitter. 2002. Rapid Changes in Flowering Time in British Plants.
Science, 296:1689–1691. doi:10.1126/science.1071617

Foley, J. A., R. DeFries, G. P. Asner, … N. Ramankutty, and P. K. Snyder. 2005. Global
consequences of land use. Science, 309:570-574. doi:10.1126/science.1111772

Fontaine, C., I. Dajoz, J. Meriguet, and M. Loreau. 2005. Functional diversity of plant–pollinator
interaction webs enhances the persistence of plant communities. PLoS Biology, 4:e1.
doi:10.1371/journal.pbio.0040001

Forister, M. L., and A. M. Shapiro. 2003. Climatic trends and advancing spring flight of
butterflies in lowland California. Global Change Biology 9: 1130–1135.
doi:10.1046/j.1365-2486.2003.00643.x

Fowler, R.E., E. L. Rotheray, and D. Goulson. 2016. Floral abundance and resource quality
influence pollinator choice. Insect Conservation and Diversity, 9:481-494.

Fründ, J., K. E. Linsenmair, and N. Blüthgen. 2010. Pollinator diversity and specialization in
relation to flower diversity. Oikos, 119:1581-1590. doi:10.1111/j.1600-
0706.2010.18450.x

Garibaldi, L. A., I. Steffan-Dewenter, C. Kremen, … S. S. Greenleaf, and A. Holzschuh. 2011.
Stability of pollination services decreases with isolation from natural areas despite
honeybee visits. Ecology Letters, 14:1062-1072. doi:10.1111/j.1461-0248.2011.01669.x

Gotlieb, A., Y. Hollender, and Y. Mandelik. 2011. Gardening in the desert changes bee
communities and pollination network characteristics. Basic and Applied
Ecology, 12:310-320. doi:10.1016/j.baae.2010.12.003

Greer, M. J., K. K. Bakker, and C. D. Dieter. 2016. Grassland bird response to recent loss and
degradation of native prairie in central and western South Dakota. The Wilson Journal of
Ornithology, 128:278-289. doi:10.1676/wils-128-02-278-289.1

Habel, J. C., W. Ulrich, N. Biburger, S. Seibold, and T. Schmitt. 2019. Agricultural
intensification drives butterfly decline. The Royal Entomological Society, 12:289-295.
doi:10.1111/icad.12343

Hadley, A. S., and M. G. Betts. 2012. The effects of landscape fragmentation on pollination
dynamics: absence of evidence not evidence of absence. Biological Reviews, 87:526-544.
doi:10.1111/j.1469-185x.2011.00205.x

Han, W., Z. Yang, L. Di, and R. Mueller. 2012. CropScape: A Web service based application for
exploring and disseminating US conterminous geospatial cropland data products for
decision support. Computers and Electronics in Agriculture, 84, 111–123.

Hass, A. L., L. Brachmann, P. Batáry, Y. Clough, H. Behling, and T. Tscharntke. 2019. Maize-
dominated landscapes reduce bumblebee colony growth through pollen diversity loss.
Journal of Applied Ecology, 56:294-304. doi:10.1111/1365-2664.13296

Heard, M. S., C. Carvell, N. L. Carreck, P. Rothery, J. L. Osborne, and A. F. G. Bourke. 2007.
Landscape context not patch size determines bumble-bee density on flower mixtures
sown for agri-environment schemes. Biology Letters, 3:638-641.
doi:10.1098/rsbl.2007.0425

Hernández-Castellano, C., A. Rodrigo, J. M. Gómez, C. Stefanescu, J. A. Calleja, S. Reverté,
and J. Bosch. 2020. A new native plant in the neighborhood: effects on plant–pollinator
networks, pollination, and plant reproductive success. Ecology, 101:e03046.
doi:10.1002/ecy.3046

Hinners, S. J., and M. K. Hjelmroos-Koski. 2009. Receptiveness of foraging wild bees to exotic
landscape elements. The American Midland Naturalist, 162:253-265. doi:10.1674/0003-
0031-162.2.253

Hoehn, P., T. Tscharntke, J. M. Tylianakis, and I. Steffan-Dewenter. 2008. Functional group
diversity of bee pollinators increases crop yield. Proceedings of the Royal Society B:
Biological Sciences, 275:2283-2291. doi:10.1098/rspb.2008.0405

Hoekstra, J. M., T. M. Boucher, T. H. Ricketts, and C. Roberts. 2005. Confronting a biome
crisis: global disparities of habitat loss and protection. Ecology letters, 8:23-29.
doi:10.1111/j.1461-0248.2004.00686.x

Jauker, F., B. Jauker, I. Grass, I. Steffan-Dewenter, and V. Wolters. 2019. Partitioning wild bee
and hoverfly contributions to plant–pollinator network structure in fragmented habitats.
Ecology, 100:e02569. doi:10.1002/ecy.2569

Jordano, P., J. Bascompte, and J. M. Olesen. 2003. Invariant properties in coevolutionary
networks of plant–animal interactions. Ecology Letters, 6:69–81. doi:10.1046/j.1461-
0248.2003.00403.x

Kandori, I., T. Hirao, S. Matsunaga, and T. Kurosaki. 2009. An invasive dandelion unilaterally
reduces the reproduction of a native congener through competition for pollination.
Oecologia, 159:559-569.

Kearns, C. A., D. W. Inouye, and N. M. Waser. 1998. Endangered mutualisms: the conservation
of plant-pollinator interactions. Annu. Rev. Ecol. Evol. Syst, 29:83-112.
doi:10.1146/annurev.ecolsys.29.1.83

Klecka, J., J. Hadrava, P. Biella, and A. Akter. 2018. Flower visitation by hoverflies (Diptera:
Syrphidae) in a temperate plant-pollinator network. PeerJ, 6:e6025.
doi:10.7717/peerj.6025

Klein, A., B. E. Vaissiere, J. H. Cane, I. Steffan-Dewenter, S. A. Cunningham, C. Kremen, and
T. Tscharntke. 2007. Importance of pollinators in changing landscapes for world crops.
Proceedings of the Royal Society, 274:303-313. doi:10.1098/rspb.2006.3721

Knight, M. E., J. L. Osborne, R. A. Sanderson, R. J. Hale, A. P. Martin, and D. Goulson. 2009.
Bumblebee nest density and the scale of available forage in arable landscapes. Insect
Conservation and Diversity, 2:116-124. doi:10.1111/j.1752-4598.2009.00049.x

Kremen, C., N. M. Williams, and R. W. Thorp. 2002. Crop pollination from native bees at risk
from agricultural intensification. Proceedings of the National Academy of Sciences,
99:16812-16816. doi:10.1073/pnas.262413599

Kremen, C., N. M. Williams, R. L. Bugg, J. P. Fay, and R. W. Thorp. 2004. The area
903   requirements of an ecosystem service: crop pollination by native bee communities in
904   California. Ecology Letters, 7:1109-1119. doi:10.1111/j.1461-0248.2004.00662.x

905   Larson, D.L., P.A. Rabie, S. Droege, J. L Larson, and M. Haar. 2016. Exotic plant infestation is
906   associated with decreased modularity and increased numbers of connectors in mixed-
907   grass prairie pollination networks. PloS One, 11:e0155068.
908   doi:10.1371/journal.pone.0155068

909   Lautenbach, S., R. Seppelt, J. Liebscher, and C.F. Dormann. 2012. Spatial and temporal trends of
910   global pollination benefit. PLoS One, 7:e35954. doi:10.1371/journal.pone.0035954

911   Lázaro, A., F. Fuster, D. Alomar, and Ø. Totland. 2020. Disentangling direct and indirect effects
912   of habitat fragmentation on wild plants' pollinator visits and seed production. Ecological
913   Applications, e02099. doi:10.1002/eap.2099

914   Liu, Z., Y. Liu, and M. H. A. Baig. 2019. Biophysical effect of conversion from croplands to
915   grasslands in water-limited temperate regions of China. The Science of the Total
916   Environment, 648:315-324. doi:10.1016/j.scitotenv.2018.08.128

917   Lopezaraiza–Mikel, M. E., R. B. Hayes, M. R. Whalley, and J. Memmott. 2007. The impact of
918   an alien plant on a native plant–pollinator network: an experimental approach. Ecology,
919   Letters 10:539-550. doi:10.1111/j.1461-0248.2007.01055.x

920   Lundgren, J. G., and S. W. Fausti. 2015. Trading biodiversity for pest problems. Science
921   Advances, 1: e1500558. doi:10.1126/sciadv.1500558

922   Mallinger, R.E., J. Bradshaw, A.J. Varenhorst, and J.R. Prasifka. 2019. Native solitary bees
923   provide economically significant pollination services to confection sunflowers
924   (*Helianthus annuus L.*)(Asterales: Asteraceae) grown across the Northern Great Plains.
925   Journal of Economic Entomology, 112:40-48. doi:10.1093/jee/toy322

926   Memmott, J., N. M. Waser, and M. V. Price. 2004. Tolerance of pollination networks to species
927   extinctions. Proceedings of the Royal Society of London. Series B: Biological Sciences,
928   271:2605-2611. doi:10.1098/rspb.2004.2909

Milfont, M.D.O., E.E.M Rocha, A.O.N. Lima, and B.M. Freitas. 2013. Higher soybean
production using honeybee and wild pollinators, a sustainable alternative to pesticides
and autopollination. Environmental Chemistry Letters, 11:335-341. doi:10.1007/s10311-
013-0412-8

Miranda, G. F. G., A. D. Young, M. M. Locke, S. A. Marshall, J. H. Skevington, and F. C.
Thompson. 2013. Key to the genera of Nearctic Syrphidae. Canadian Journal of
Arthropod Identification, 23:351. doi:10.3752/cjai.2013.23

Montoya, J. M., and G. Yvon-Durocher. 2007. Ecological networks: information theory meets
Darwin's entangled bank. Current Biology, 17:128-130. doi:10.1016/j.cub.2007.01.028

Morales, C.L. and A. Traveset. 2008. Interspecific pollen transfer: magnitude, prevalence and
consequences for plant fitness. Critical Reviews in Plant Sciences, 27:221-238.
doi:10.1080/07352680802205631

Moreira, E. F., D. Boscolo, and B. F. Viana. 2015. Spatial heterogeneity regulates plant-
pollinator networks across multiple landscape scales. PloS One, 10:e0123628.
doi:10.1371/journal.pone.0123628

Morris, W. F. 1993. Predicting the Consequence of Plant Spacing and Biased Movement for 754
Pollen Dispersal by Honey bees. Ecology, 74:493–500. doi:10.2307/1939310

Morton, E.M. and N.E. Rafferty. 2017. Plant–pollinator interactions under climate change: The
use of spatial and temporal transplants. Applications in Plant Sciences, 5:1600133.

Naug, D. 2009. Nutritional stress due to habitat loss may explain recent honeybee colony
collapses. Biological Conservation, 142:2369-2372. doi:10.1016/j.biocon.2009.04.007

Nielsen, A., and J. Bascompte. 2007. Ecological networks, nestedness and sampling effort.
Journal of Ecology, 95:1134-1141. doi: 10.11111/j.1365-2745.2007.01271.

Nottebrock, H., B. Schmid, K. Mayer, C. Devaux, K. J. Esler, K. Böhning-Gaese, M.
Schleuning, J. Pagel, and F. M. Schurr. 2017. Sugar landscapes and pollinator-mediated
interactions in plant communities. Ecography, 40:1129-1138. doi:10.1111/ecog.02441

Oksanen, J., F. G. Blanchet, M. Friendly, R. Kindt, P. Legendre, D. McGlinn, P. R. Minchin, R.
B. O'Hara, G. L. Simpson, and H. Wagner. 2019. vegan: Community Ecology Package. R
package version 2.5-6. https://CRAN.R-project.org/package=vegan

Okuyama, T., and J. N. Holland. 2008. Network structural properties mediate the stability of
mutualistic communities. Ecology Letters, 11:208-216. doi:10.1111/j.1461-
0248.2007.01137.x

Olesen, J. M. and S. K. Jain. 1994. D: Fragmented plant populations and their lost interactions.
In V. Loeschcke, S. K. Jain, and J. Tomiuk [eds.], Conservation Genetics, 417-426.
Birkhäuser, Basel. doi:10.1007/978-3-0348-8510-2

Olesen, J. M., J. Bascompte, Y. L. Dupont, and P. Jordano. 2007. The modularity of pollination
networks. Proceedings of the National Academy of Sciences, 104:19891-19896.
doi:10.1073/pnas.0706375104

Otto, C. R., C. L. Roth, B. L. Carlson, and M. D. Smart. 2016. Land-use change reduces habitat
suitability for supporting managed honeybee colonies in the Northern Great Plains.
Proceedings of the National Academy of Sciences, 113:10430-10435.
doi:10.1073/pnas.1603481113

Otto, C.R.V., A. Smart, R. S. Cornman, M. Simanonok, and D.D. Iwanowicz. 2020. Forage and
habitat for pollinators in the northern Great Plains—Implications for U.S. Department of
Agriculture conservation programs: U.S. Geological Survey Open-File Report 2020–
1037, 64 p. doi: 10.3133/ ofr20201037

Ouvrard, P., J. Transon, and A. L. Jacquemart. 2018. Flower-strip agri-environment schemes
provide diverse and valuable summer flower resources for pollinating insects.
Biodiversity and Conservation, 27:2193-2216. doi: 10.1007/s10531-018-1531-0

Padrón, B., A. Traveset, T. Biedenweg, D. Díaz, M. Nogales, and J. M. Olesen. 2009. Impact of
alien plant invaders on pollination networks in two archipelagos. PLoS One, 4:e6275.
doi:10.1371/journal.pone.0006275

Perpinan, O., and R. Hijmans. 2020. rasterVis. R package version 0.49.

Petanidou, T., A. S. Kallimanis, J. Tzanopoulos, S. P. Sgardelis, and J.P. Pantis . 2008. Long-
983   term observation of a pollination network: Fluctuation in species and interactions, relative
984   invariance of network structure and implications for estimates of speciation. Ecology
985   Letters, 11:564 – 575. doi: 10.1111/j.1461-0248.2008.01170.x

986   Peterson, M.J., D.M. Hall, A.M. Feldpausch-Parker and T.R. Peterson. 2010. Obscuring
987   ecosystem function with application of the ecosystem services concept. Conservation
988   Biology, 24:113-119. doi: 10.1111/j.1523-1739.2009.01305.x

989   Pettis, J. S., E. M. Lichtenberg, M. Andree, J. Stitzinger, and R. Rose. 2013. Crop pollination
990   exposes honey bees to pesticides which alters their susceptibility to the gut pathogen
991   *Nosema ceranae*. PloS One, 8:e70182. doi:10.1371/journal.pone.0070182

992   Pinheiro J., D. Bates, S. DebRoy, D. Sarkar, and R Core Team. 2020. nlme: Linear and
993   Nonlinear Mixed Effects Models. R package version 3.1-144, <URL: https://CRAN.R-
994   project.org/package=nlme>.

995   Ponisio, L. C., P. de Valpine, L. K. M'Gonigle, and C. Kremen. 2019. Proximity of restored
996   hedgerows interacts with local floral diversity and species' traits to shape long-term
997   pollinator metacommunity dynamics. Ecology Letters, 22:1048-1060.
998   doi:10.1111/ele.13257

Potts, S. G., B. Vulliamy, A. Dafni, G. Ne'eman, and P. Willmer. 2003. Linking bees and
flowers: how do floral communities structure pollinator communities? Ecology, 84:2628-
2642. doi:10.1890/02-0136

Potts, S. G., J. C. Biesmeijer, C. Kremen, P. Neumann, O. Schweiger, and W. E. Kunin. 2010.
Global pollinator declines: trends, impacts and drivers. Trends in Ecology & Evolution,
25:345-353. doi:10.1016/j.tree.2010.01.007

Potts, S. G., B. Vulliamy, S. Roberts, C. O'Toole, A. Dafni, G. Ne'eman, and P. Willmer. 2005.
Role of nesting resources in organizing diverse bee communities in a Mediterranean
landscape. Ecological Entomology, 30:78-85. doi:10.1111/j.0307-6946.2005.00662.x

Potts, S. G., B. A. Woodcock, S. P. M. Roberts, T. Tscheulin, E. S. Pilgrim, V. K. Brown, and J.
R. Tallowin. 2009. Enhancing pollinator biodiversity in intensive grasslands. Journal of
Applied Ecology, 46:369-379. doi:10.1111/j.1365-2664.2009.01609.x

R Core Team. 2013. R: A language and environment for statistical computing. version 3.6.3

Ramankutty, N., and J. A. Foley. 1999. Estimating historical changes in global land cover:
Croplands from 1700 to 1992. Global Biogeochemical Cycles, 13:997-1027.
doi:10.1029/1999gb900046

Rashford, B. S., J. A. Walker, and C. T. Bastian. 2011. Economics of grassland conversion to
cropland in the Prairie Pothole Region. Conservation Biology, 25:276-284.
doi:10.1111/j.1523-1739.2010.01618

Redhead, J. W., B. A. Woodcock, M. J. Pocock, R. F. Pywell, A. J. Vanbergen, and T.H. Oliver.
2018. Potential landscape-scale pollinator networks across Great Britain: structure,
stability, and influence of agricultural land cover. Ecology Letters, 21:1821-1832.
doi:10.1111/ele.13157

Requier, F., K. K. Jowanowitsch, K. Kallnik, and I. Steffan-Dewenter. 2020. Limitation of
complementary resources affects colony growth, foraging behavior, and reproduction in
bumble bees. Ecology, 101:e02946. doi:10.1002/ecy.2946

Robinson, S. V. J., G. Losapio, and G. H. R. Henry. 2018. Flower-power: Flower diversity is a
stronger predictor of network structure than insect diversity in an Arctic plant–pollinator
network. Ecological complexity, 36:1-6. doi:10.1016/j.ecocom.2018.04.005

Rodríguez-Gironés, M. A., and L. Santamaría. 2006. A new algorithm to calculate the nestedness
temperature of presence–absence matrices. Journal of Biogeography, 33:924-935.
doi:10.1111/j.1365-2699.2006.01444

Rogers, S. R., D. R. Tarpy, and H. J. Burrack. 2014. Bee species diversity enhances productivity
and stability in a perennial crop. PloS One, 9:e97307. doi:10.1371/journal.pone.0097307

Roy, D. B., and T. H. Sparks. 2000. Phenology of British butterflies and climate change. Global
Change Biology, 6:407–416. doi:10.1046/j.1365-2486.2000.00322.x

Royer, F., and R. Dickinson. 1999. Weeds of the Northern US and Canada. The University of
Alberta Press, Edmonton, Canada.

Sakai, S., S. Metelmann, Y. Toquenaga, and A. Telschow. 2016. Geographical variation in the
heterogeneity of mutualistic networks. Royal Society Open Science, 3:150630.
doi:10.1098/rsos.150630

Smart M. D., J. S. Pettis, N. H. Euliss, and M. S. Spivak. 2016. Land use in the Northern Great
Plains region of the U.S. influences the survival and productivity of honeybee colonies.
Agriculture, Ecosystems, and Environment, 230:139–149.
doi:10.1016/j.agee.2016.05.030

Soares, R. G. S., P. A. Ferreira, and L. E. Lopes. 2017. Can plant-pollinator network metrics
indicate environmental quality? Ecological Indicators, 78:361-370.
doi:10.1016/j.ecolind.2017.03.037

Sommaggio, D. 1999. Syrphidae: can they be used as environmental bioindicators? Agriculture,
Ecosystems & Environment, 74:343-356. doi:10.1016/s0167-8809(99)00042-0

Spellman, K.V., L. C. Schneller, C. P. Mulder, C.P. and M. L. Carlson. 2015. Effects of non-
native Melilotus albus on pollination and reproduction in two boreal shrubs. Oecologia,
179:495-507. doi:10.1007/s00442-015-3364-9

Spiesman, B. J., and B. D. Inouye. 2013. Habitat loss alters the architecture of plant–pollinator
interaction networks. Ecology, 94:2688-2696. doi:10.1890/13-0977.1

Spleen, A. M., E. J. Lengerich, K. Rennich, … J. Stitzinger, and K. Lee. 2013. A national survey
of managed honeybee 2011–12 winter colony losses in the United States: results from the
Bee Informed Partnership. Journal of Apicultural Research, 52:44-53.
doi:10.3896/IBRA.1.52.2.07

Stang, M., P. G. L. Klinkhamer, and E. van der Meijden. 2006. Size constraints and flower
abundance determine the number of interactions in a plant–flower visitor web. Oikos,
112:111-121. doi:10.1111/j.0030-1299.2006.14199

Steffan-Dewenter, I., U. Münzenberg, C. Bürger, C. Thies, and T. Tscharntke. 2002. Scale-
dependent effects of landscape context on three pollinator guilds. Ecology, 83:1421-1432.
doi:10.1890/0012-9658(2002)083[1421:sdeolc]2.0.co;2

Steffan-Dewenter, I., S. G. Potts, and L. Packer. 2005. Pollinator diversity and crop pollination
services are at risk. Trends in Ecology & Evolution, 20:651-652.
doi:10.1016/j.tree.2005.09.00

Sugden, E. A., R. W. Thorp, and S. L. Buchmann. 1996. Honeybee–native bee competition:
focal point for environmental change and apicultural response in Australia. Bee World,
77:26–44. doi:10.1080/0005772x.1996.11099280

Tamburini G., F. Lami, L. Marini. 2017. Pollination benefits are maximized at intermediate
nutrient levels. Proc. R. Soc. B 284:20170729. doi:10.1098/rspb.2017.0729

Tepedino, V.J., B.A. Bradley, and T.L. Griswold. 2008. Might flowers of invasive plants
increase native bee carrying capacity? Intimations from Capitol Reef National Park,
Utah. Natural Areas Journal, 28:44–50. doi: 10.3375/0885-
8608(2008)28[44:MFOIPI]2.0.CO;2

Thébault, E., and C. Fontaine. 2010. Stability of ecological communities and the architecture of
mutualistic and trophic networks. Science, 329:853-856. doi: 10.1126/science.1188321

Thomson, D. 2004. Competitive interactions between the invasive European honeybee and
native bumble bees. Ecology, 85:458-470. doi:10.1890/02-0626

Tscharntke, T., A. M. Klein, A. Kruess, I. Steffan-Dewenter, and C. Thies. 2005. Landscape
perspectives on agricultural intensification and biodiversity–ecosystem service
management. Ecology Letters, 8:857-874. doi: 10.1111/j.1461-0248.2005.00782

Twerd, L., and W. Banaszak-Cibicka. 2019. Wastelands: their attractiveness and importance for
preserving the diversity of wild bees in urban areas. Journal of Insect Conservation,
23:573-588. doi:10.1007/s10841-019-00148-8

Tylianakis, J. M., and R. J. Morris. 2017. Ecological networks across environmental gradients.
Annual Review of Ecology, Evolution, and Systematics, 48:25-48. doi:10.1146/annurev-
ecolsys-110316-022821

US Department of Agriculture, National Agricultural Statistics Service. 2014. US Department of
Agriculture, Washington, DC. Available at
http://usda.mannlib.cornell.edu/MannUsda/viewDocumentInfo.do?documentID=1191.

USDA National Agricultural Statistics Service Cropland Data Layer. 2019. Published crop-
specific data layer [Online]. Available at https://nassgeodata.gmu.edu/CropScape/
(accessed in 2019; verified in 2019. USDA-NASS, Washington, DC.

Valdovinos, F. S., B. J. Brosi, H. M. Briggs, P. Moisset de Espanés, R. Ramos-Jiliberto, and N.
D. Martinez. 2016. Niche partitioning due to adaptive foraging reverses effects of
nestedness and connectance on pollination network stability. Ecology Letters, 19:1277–
1286. doi:10.1111/ele.12664

Van Bruggen, T. 1985. The Vascular Plants of South Dakota. Second edition. Iowa State
University Press, Ames, Iowa, USA.

Vanbergen, A. J., and the Insect Pollinators Initiative. 2013. Threats to an ecosystem service:
pressures on pollinators. Frontiers in Ecology and the Environment, 11:251-259.
doi:10.1890/120126

Vanbergen, A.J. 2014. Landscape alteration and habitat modification: impacts on plant–
pollinator systems. Current Opinion in Insect Science, 5:44-49.

Venables, W. N. and B. D. Ripley. 2002. Modern Applied Statistics with S. Fourth Edition.
Springer, New York. ISBN 0-387-95457-0

Vickruck, J. L., L. R. Best, M. P. Gavin, J. H. Devries, and P. Galpern. 2019. Pothole wetlands
provide reservoir habitat for native bees in prairie croplands. Biological Conservation,
232:43-50. doi:10.1016/j.biocon.2019.01.015

Vinson S.B., G. W. Frankie, and J. Barthell. 1993. Threats to the diversity of solitary bees in a
neotropical dry forest in Central America. See Ref. 121a:53–82

Waser, N.M. 1983. Competition for pollination and floral character differences among sympatric
plant species: a review of evidence. *In* C. E. Jones and R. J. Little [eds.], Handbook of
experimental pollination biology, 277-293.

Weiner, C.N., M. Werner, K. E. Linsenmair, and N. Blüthgen. 2014. Land-use impacts on plant–
pollinator networks: interaction strength and specialization predict pollinator declines.
Ecology, 95:466-474. doi:10.1890/13-0436.1

Williams, I. H. 2002. Insect pollination and crop production: a European perspective. *In*
Pollinating Bees-The Conservation Link Between Agriculture and Nature, 59-65.
Ministry of Environment, Brasilia, Brazil.

Williams, N. M., D. Cariveau, R. Winfree, and C. Kremen . 2011. Bees in disturbed habitats use,
but do not prefer, alien plants. Basic and Applied Ecology, 12:332-341.
doi:10.1016/j.baae.2010.11.008

Winfree, R., T. Griswold, and C. Kremen. 2007. Effect of human disturbance on bee
communities in a forested ecosystem. Conservation Biology, 21:213-223.
doi:10.1111/j.1523-1739.2006.00574

Wright, C. K., and M. C. Wimberly. 2013. Recent land use change in the Western Corn Belt
threatens grasslands and wetlands. Proceedings of the National Academy of Sciences,
110:4134-4139. doi:10.1073/pnas.1215404110

1131

Figure Legends

Figure 1. Size of remnant temperate grassland sites and the surrounding landscape diversity within 3000 meters of the sites in the Prairie Coteau Region near Brookings, South Dakota. Remnant temperate grasslands are ordered by size from smallest to largest in hectares. The size of the circle indicates the landscape diversity surrounding that site in 2019.

Figure 2. Shannon diversity of pollinators at the functional, family and genus level from May through October 2019 in the Prairie Coteau region near Brookings, South Dakota. Site abbreviations are as follows: DC = Deer Creek, AG = Altamont GPA, AR = Aurora, BP = Brookings Prairie, OL = Oak Lake Field Station, S7 = Seven-mile Fen, SV = Severson WMA, GG = Gary Gulch, CP = Coteau Prairie WMA, CL = Coteau Lakes GPA, RBH2 = Round Bullhead Lakes Site 2, SP = Sioux Prairie, AP = Altamont Prairie, RBH1 = Round Bullhead Site 1, J = Jacobsen Fen.R5T

Figure 3. Shannon diversity of biotically-pollinated plants at the family, genus, and species level from May through October 2019 in the Prairie Coteau region near Brookings, South Dakota. Site abbreviations are as follows: DC = Deer Creek, AG = Altamont GPA, AR = Aurora, BP = Brookings Prairie, OL = Oak Lake Field Station, S7 = Seven-mile Fen, SV = Severson WMA, GG = Gary Gulch, CP = Coteau Prairie WMA, CL = Coteau Lakes GPA, RBH2 = Round Bullhead Lakes Site 2, SP = Sioux Prairie, AP = Altamont Prairie, RBH1 = Round Bullhead Site 1, J = Jacobsen Fen.

Figure 4. (A) Effect of pollinator genus diversity on network-level specialization (H2') in the plant-pollinator communities found within the Prairie Coteau region near Brookings, South Dakota in 2019. Pollinator genus diversity refers to the Shannon diversity of pollinators in the

sampled remnant temperate grasslands within the Prairie Coteau region. Network specialization

ranges from 0 -1 with values approaching 1 denoting more specialized interactions in a

community. Pollinator genus diversity has a significant positive effect on network-level

specialization at the mid-season ($p < 0.002$, $t = 4.313$, std. error = 0.07). (B) Effect of pollinator

genus diversity on connectance in the plant-pollinator communities found within the Prairie

Coteau region near Brookings, South Dakota in 2019. Pollinator genus diversity refers to the

Shannon diversity of pollinators in the sampled remnant temperate grasslands within the Prairie

Coteau region. Connectance ranges from $0 - 1$ with the value of 1 indicating a maximally

connected community. Pollinator genus diversity has a significant negative effect on network

connectance ($p < 0.04$, $t = -2.39$, std. error = 0.032).

1164

1165   *Figure 1*

1166

1167

1168   *Figure 2*

1169

1170

1171   *Figure 3*

1172

1173

1174   *Figure 4*

1175

1176

Tables

Table 1

Description of temperate grassland sites in eastern South Dakota including site name, county name, latitude & longitude, size (in hectares) and ownership of sites.

Site Name	County	Latitude and Longitude	Size (hectares)	Ownership
Sioux Prairie	Moody	44.031099, -96.783710	81	Nature Conservancy land
Brookings Prairie	Brookings	44.252489, -96.810310	16	City of Brookings
Aurora Prairie	Brookings	44.262012, -96.704581	12	Nature Conservancy land
Oak lake	Deuel	44.509589, -96.532529	231	South Dakota State University
Deer Creek	Brookings	44.469750, -96.502181	91	South Dakota Game, Fish and Parks
Severson WPA	Brookings	44.7145257,-96.5003879	81	U.S. Fish and Wildlife Service
Seven-mile fen	Deuel	44.748073, -96.535572	88	Nature Conservancy land
Jacobson Fen	Deuel	44.793818, -96.627707	65	Nature Conservancy land
Coteau Lakes	Deuel	44.819894, -96.728137	239	South Dakota Game, Fish and Parks
Altamont	Deuel	44.860664, -96.710675	81	South Dakota Game, Fish and Parks
Altamont Prairie	Deuel	44.887370, -96.533403	25	Nature Conservancy land
Coteau Prairie	Deuel	44.904428, -96.719979	169	U.S. Fish and Wildlife Service
Round Bullhead 1	Deuel	44.933864, -96.828411	8	South Dakota Game, Fish and Parks
Round Bullhead 2	Deuel	44.920024, -96.828519	433	South Dakota Game, Fish and Parks
Gary Gulch	Deuel	44.790391, -96.468345	49	South Dakota Game, Fish and Parks

Table 2

North American temperate grassland attributes and landscape diversity effects on pollinator

diversity in the Prairie Coteau region near Brookings, South Dakota in 2019. Pollinator genus

diversity is the response variable in the following models and refers to the Shannon diversity of

pollinators in the sampled remnant temperate grasslands within the Prairie Coteau region. The

landscape diversity variables refer to the scales at which we measured landscape diversity (500

m, 1000 m, 3000 m). Plant species diversity refers to the Shannon diversity of biotically-

pollinated plant species in the sampled remnant temperate grasslands and Transect Sugar refers

to the amount of sugar that was available on each transect. Transect sugar values were derived

from the nectar estimations collected from floral surveys. Random effects in the model include

the remnant temperate grassland site itself. Backwards stepwise selection is applied to the

following models in the table using single term deletion in the dropterm function in 'MASS'

package in R. We list each model in the table and its associated AIC value. The first AIC values

refer to the AIC score of the current model listed. The AIC values next to each variable refer to

the AIC score of the model if the associated variable was removed. The p-values in the table are

outputs from the likelihood ratio test for each variable. * values indicate that removing the

associated variable would significantly change the model. *Indicates significance at the $\alpha = 0.05$

level. $P < 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Model Selection for Pollinator Genus Diversity =	Likelihood ratio test	AIC	p-value
Landscape Diversity at 500 m + Landscape Diversity at 1000 m + Landscape Diversity at 3000 m + Plant Species Diversity + Transect Sugar		177.79	
Landscape Diversity at 500 m	1.32	177.12	0.250
Landscape Diversity at 1000 m	0.28	176.07	0.596
Landscape Diversity at 3000 m	0.02	175.81	0.881
Plant Species Diversity	0.68	176.47	0.409
Transect Sugar	4.43	180.22	0.035*
Landscape Diversity at 500 m + Landscape Diversity at 1000 m + Plant Species Diversity + Transect Sugar		175.81	
Landscape Diversity at 500 m	1.35	175.16	0.246
Landscape Diversity at 1000 m	0.41	174.22	0.522
Plant Species Diversity	0.66	174.48	0.415
Transect Sugar	4.42	178.23	0.036*
Landscape Diversity at 500 m + Plant Species Diversity +Transect Sugar		174.22	
Landscape Diversity at 500 m	1.02	173.24	0.313
Plant Species Diversity	0.72	172.94	0.396
Transect Sugar	4.30	176.53	0.038*
Landscape Diversity at 500 m + Transect Sugar		172.94	
Landscape Diversity at 500 m	1.01	171.95	0.315
Transect Sugar	4.61	175.56	0.032*
Transect Sugar		171.95	
Transect Sugar	6.41	176.37	0.011*

1199

1200

1201　*Table 3*

1202　Linear mixed effect analysis of transect sugar on pollinator genus Shannon diversity in the

1203　Prairie Coteau region near Brookings, South Dakota. P < 0.05; * P < 0.05; ** P < 0.01; *** P <

1204　0.001

Variables	Coefficient	Standard Error	Degrees of Freedom	p-value
Intercept	1.16	0.06	14.78	5.97e-12***
Transect Sugar	0.03	0.01	104.86	0.013*

1205

1206

Table 4

North American temperate grassland attributes and landscape proportion effects on pollinator

diversity in the Prairie Coteau region near Brookings, South Dakota in 2019. Pollinator genus

diversity is the response variable in the following models and refers to the Shannon diversity of

pollinators in the sampled remnant temperate grasslands within the Prairie Coteau region.

Landscape proportion for the five most prominent land-uses were measured for the same scales

used for landscape diversity (500 m, 1000 m, 3000 m). Plant species diversity refers to the

Shannon diversity of biotically-pollinated plant species in the sampled remnant temperate

grasslands and Transect Sugar refers to the amount of sugar that was available on each transect.

Transect sugar values were derived from the nectar estimations collected from floral surveys.

Random effects in the model include season (i.e., early, mid and late) and the remnant temperate

grassland site itself. Backwards stepwise selection was applied to the initial model presented in

the table using single term deletion in the dropterm function in 'MASS' package in R. We

present the initial model in the table and then the final, simplified model after single term

deletion. The AIC values at the top refer to the AIC score of the current model listed. The AIC

values next to each variable refer to the AIC score of the model if the associated variable was

removed. The p-values in the table are outputs from the likelihood ratio test for each variable. *

values indicate that removing the associated variable would significantly change the model.

*Indicates significance at the $\alpha = 0.05$ level. $P < 0.05$; * $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

1226

1227

1228

Model Selection for Pollinator Genus Diversity =	Likelihood ratio test	AIC	p-value
Initial Model: **Corn Proportion at 500 m + Corn Proportion at 1000 m + Corn Proportion at 3000 m + Soy Proportion at 500 m + Soy Proportion at 1000 m + Soy Proportion at 3000 m + Fallow/Idle Cropland Proportion at 500 m + Fallow/Idle Cropland Proportion at 1000 m + Fallow/Idle Cropland Proportion at 3000 m + Grassland Proportion at 500 m + Grassland Proportion at 1000 m + Grassland Proportion at 3000 m + Herbaceous Wetland Proportion at 500 m + Herbaceous Wetland Proportion at 1000 m + Herbaceous Wetland Proportion at 3000 m + Plant Species Diversity + Transect Sugar**		185.90	
	1.71	185.61	0.19
Corn Proportion at 500 m	0.67	184.57	0.41
Corn Proportion at 1000 m	0.30	184.19	0.59
Corn Proportion at 3000 m	0.03	183.93	0.86
Fallow/Idle Cropland Proportion at 500 m	0.17	184.07	0.68
Fallow/Idle Cropland Proportion at 1000 m	3.22	187.12	0.07
Fallow/Idle Cropland Proportion at 3000 m	0.48	184.38	0.49
Grassland Proportion at 500 m	0.32	184.22	0.57
Grassland Proportion at 1000 m	0.58	184.48	0.45
Grassland Proportion at 3000 m	0.76	184.66	0.38
Herbaceous Wetland Proportion at 500 m	2.52	186.42	0.11
Herbaceous Wetland Proportion at 1000 m	0.60	184.50	0.44
Herbaceous Wetland Proportion at 3000 m	0.09	183.99	0.76
Soy Proportion at 500 m	2.61	186.51	0.11
Soy Proportion at 1000 m	0.68	184.58	0.41
Soy Proportion at 3000 m	1.16	185.06	0.28
Plant Species Diversity	6.04	189.94	0.01 *
Transect Sugar			
Final Model: **Corn Proportion at 500 m + Herbaceous Wetland Proportion at 1000 m + Soy Proportion at 1000 m + Transect Sugar**		167.22	

Corn Proportion at 500 m	6.90	172.12	0.009 **
Herbaceous Wetland Proportion at 1000 m	4.92	170.14	0.026 *
Soy Proportion at 1000 m	6.82	172.04	0.009 **
Transect Sugar	5.99	171.20	0.014 *

1230

1231

1232 *Table 5*

1233   Linear mixed effects analysis results of land-use proportions and transect sugar on pollinator

1234 genus Shannon diversity in the Prairie Coteau region near Brookings, South Dakota. $P < 0.05$; *

1235 $P < 0.05$; ** $P < 0.01$; *** $P < 0.001$

Variable	Coefficient	Standard Error	Degrees of Freedom	p-value
Intercept	8.45e-01	1.10e-01	1.09e+02	6.06e-12 ***
Corn Proportion at 500 m	1.80e+00	6.89e-01	1.09e+02	0.0086 *
Herbaceous Wetland Proportion at 1000 m	1.61e+00	7.36e-01	1.09e+02	0.0264 *
Soy Proportion at 1000 m	3.02e+00	1.16e+00	1.09e+02	0.0090 *
Transect Sugar	2.41e-02	9.92e-03	1.09e+02	0.0144 *

1236

Appendices

1237

The table below lists all of the pollinators observed and identified in the Prairie Coteau region

near Brookings, South Dakota in 2019 down to lowest taxonomic level. Insect pollinators that

could not be identified to genus were placed in a catch-all genus that consisted of the first five

letters of their family name. For example, for a fly pollinator in the family Muscidae, we created

a genus named Gen_Musci in the dataset in order to include these visitors in the analyses. Table

also includes number of remnant temperate grassland sites the pollinators were present and the

number of total observed interactions of each pollinator from May through October 2019.

Pollinators	Family	Functional Group	Number of sites present Out of 15 total sties	Number of total interactions from May through October
Agapostemon spp.	Halictidae	Small bee	10	163
Agapostemon sp. 1	Halictidae	Small bee	3	8
Agapostemon sp. 2	Halictidae	Small bee	1	96
Allographta spp.	Syrphidae	Syrphid fly	1	1
Andrena spp.	Andrenidae	Small bee	3	8
Apis mellifera	Apidae	Honey bee	7	1218
Archytas sp. 1	Tachinidae	Non-syrphid fly	6	70
Augochlora spp.	Halictidae	Small bee	1	7
Augochlorella sp. 1	Halictidae	Small bee	1	3
Boloria bellona	Nymphalidae	Lepidopteran	3	4
Bombus bimaculatus	Apidae	Large bee	2	5
Bombus borealis	Apidae	Large bee	2	32
Bombus griseocollis	Apidae	Large bee	9	332
Bombus huntii	Apidae	Large bee	1	2

Bombus impatiens	Apidae	Large bee	6	36
Bombus pensylvanicus	Apidae	Large bee	5	64
Bombus spp.	Apidae	Large bee	2	162
Bracon spp.	Braconidae	Small wasp	1	4
Ceratina spp.	Apidae	Small bee	3	9
Cercyonis pegalla	Nymphalidae	Lepidopteran	4	12
Chauliognathus pensylvanicus	Cantharidae	Beetle	14	850
Cisseps fulvicollis	Erebidae	Lepidopteran	9	173
Coleophora sp. 1	Coleophoridae	Lepidopteran	1	2
Coleophora sp. 2	Coleophoridae	Lepidopteran	4	44
Coleophora spp.	Coleophoridae	Lepidopteran	1	2
Colias philodice	Pieridae	Lepidopteran	5	6
Cupido comyntas	Lycaenidae	Lepidopteran	2	5
Danaus plexippus	Nymphalidae	Lepidopteran	8	11
Diabrotica sp. 1	Chrysomelidae	Beetle	2	11
Enodia anthedon	Nymphalidae	Lepidopteran	3	6
Eristalis transversa	Syrphidae	Syrphid fly	3	11
Eristalis-BFBF sp. 1	Syrphidae	Syrphid fly	3	24
Eristalis-BWFBF sp. 2	Syrphidae	Syrphid fly	3	33
Eristalis-FBF sp. 3	Syrphidae	Syrphid fly	10	402
Eristalis-FBFLA sp. 4	Syrphidae	Syrphid fly	1	5
Eristalis-OFBF sp. 5	Syrphidae	Syrphid fly	7	77
Eristalis-WFBF sp. 6	Syrphidae	Syrphid fly	2	24
Gen_Andren spp.	Andrenidae	Small bee	2	14
Gen_Antho spp.	Anthomyiidae	Non-syrphid fly	4	14
Gen_Bombylii spp.	Bombyliidae	Non-syrphid fly	2	4
Gen_Calliphor spp.	Calliphoridae	Non-syrphid fly	11	259
Gen_Ceramby spp.	Cerambycidae	Beetle	3	5

Gen_Chironom spp.	Chironomidae	Non-syrphid fly	2	3
Gen_Chloro spp.	Chloropidae	Non-syrphid fly	13	401
Gen_Chryso sp. 1	Chrysomelidae	Beetle	8	143
Gen_Chryso spp.	Chrysomelidae	Beetle	3	35
Gen_Chrysopid spp.	Chrysopidae	Non-syrphid fly	1	1
Gen_Crambi spp.	Crambidae	Lepidopteran	2	3
Gen_Culici spp.	Culicidae	Non-syrphid fly	3	7
Gen_Dolicho spp.	Dolichopodidae	Non-syrphid fly	1	27
Gen_Elachis spp.	Elachisitidae	Lepidopteran	1	1
Gen_Fannii sp. 1	Fanniidae	Non-syrphid fly	3	49
Gen_Fannii spp.	Fanniidae	Non-syrphid fly	1	3
Gen_Halic spp.	Halictidae	Small bee	7	46
Gen_Hesp spp.	Hesperidae	Lepidopteran	5	10
Gen_Ichi spp.	Ichneumonidae	Small wasp	4	11
Gen_Lampyr sp. 1	Lampyiridae	Beetle	1	2
Gen_Mega spp.	Megachilidae	Small bee	1	35
Gen_Meloi sp. 1	Meloidae	Beetle	1	2
Gen_Meloi spp.	Meloidae	Beetle	8	24
Gen_Mely spp.	Melyridae	Beetle	1	7
Gen_Miri spp.	Miridae	Hemiptera	3	8
Gen_Musc sp. 1	Muscidae	Non-syrphid fly	1	3
Gen_Musc spp.	Muscidae	Non-syrphid fly	13	755
Gen_Noctu spp.	Noctuidae	Lepidopteran	1	1
Gen_Pent	Pentatomidae	Hemiptera	3	8
Gen_Pompi sp. 1	Pompilidae	Small wasp	1	1
Gen_Pompi spp.	Pompilidae	Large wasp	2	6
Gen_Pyrgot sp. 1	Pyrgotidae	Non-syrphid fly	1	1
Gen_Sarco spp.	Sarcophagidae	Non-syrphid fly	10	139
Gen_Spheci spp.	Sphecidae	Large wasp	1	4

Gen_Sphing spp.	Sphingidae	Lepidopteran	1	6
Gen_Strat sp. 1	Stratiomyidae	Non-syrphid fly	4	22
Gen_Strat spp.	Stratiomyidae	Non-syrphid fly	3	33
Gen_Syr sp. 1	Syrphidae	Syrphid fly	1	1
Gen_Syr sp. 2	Syrphidae	Syrphid fly	2	15
Gen_Syr spp.	Syrphidae	Syrphid fly	1	10
Gen_Tachin spp.	Tachinidae	Non-syrphid fly	1	1
Gen_Tephrit spp.	Tephritidae	Non-syrphid fly	3	27
Gen_Ulidii spp.	Ulidiidae	Non-syrphid fly	7	80
Gen_Weev spp.	Curculionidae	Beetle	9	36
Halictus spp.	Halictidae	Small bee	10	542
Helophilus sp. 1	Syrphidae	Syrphid fly	5	24
Helophilus spp.	Syrphidae	Syrphid fly	2	39
Hemyda spp.	Tachinidae	Non-syrphid fly	1	1
Heriades spp.	Megachilidae	Small bee	1	13
Hylaeus sp. 1	Colletidae	Small bee	2	13
Hylaeus spp.	Colletidae	Small bee	2	2
Junonia coenia	Nymphalidae	Lepidopteran	1	1
Lasioglossum spp.	Halictidae	Small bee	9	457
Lithurgopsis spp.	Megachilidae	Small bee	1	1
Megachile spp.	Megachilidae	Small bee	1	8
Melissodes spp.	Apidae	Small bee	6	23
Merodon equestris	Syrphidae	Syrphid fly	1	2
Nomada spp.	Apidae	Small bee	1	6
Parasyrphus sp. 1	Syrphidae	Syrphid fly	3	88
Parhelophilus spp.	Syrphidae	Syrphid fly	3	35
Phyciodes pulchella	Nymphalidae	Lepidopteran	5	8
Phyllotreta sp. 1	Chrysomelidae	Beetle	3	110
Pseudopanurgus spp.	Andrenidae	Small bee	2	2

Speyeria idalia	Nymphalidae	Lepidopteran	2	2
Sphaerophoria spp.	Syrphidae	Syrphid fly	5	54
Stratiomys spp.	Stratiomyidae	Non-syrphid fly	3	3
Syritta sp. 1	Syrphidae	Syrphid fly	7	82
Toxomerus spp.	Syrphidae	Syrphid fly	15	2125

1246

1247

Appendix B

The table below lists all of the biotically-pollinated flowering plants identified in the Prairie

Coteau region near Brookings, South Dakota in 2019 down to lowest taxonomic level. All

species except *Agrimonia sp.* was identified to species level. Plant code format is derived from

United States Department of Agriculture plant database format. Average sugar contribution

refers to average nectar sugar each plant species contributed per transect based on amount of

sugar per inflorescence and number of inflorescences.

Species	Family	Common name	Number of sites present Out of 15 total sites	Plant code	Average sugar contribution
Achillea millefolium	Asteraceae	Common yarrow	6	ACHMIL	1.39e-01
Agoseris glauca	Asteraceae	False dandelion	2	AGOGLA	4.62e-03
Agrimonia sp.	Rosaceae	Agrimony	2	AGRIMONIA	8.78e-04
Allium stellatum	Amaryllidaceae	Prairie onion	7	ALLSTE	1.57e-02
Amorpha canescens	Fabaceae	Lead plant	6	AMOCAN	2.18e-02
Anemone canadensis	Ranunculaceae	Meadow anemone	9	ANCA	1.64e-03
Anticlea elegans	Melanthiaceae	Mountain deathcamas	3	ANTELE	-
Apocynum cannabinum	Apocynaceae	Prairie dogbane	1	APOCAN	1.10e-02
Artemisia absinthium	Asteraceae	Absinthe wormwood	1	ARTABS	2.77e-02
Asclepias stenophylla	Apocynaceae	Slim leaf milkweed	2	ASCSTE	2.94e-02
Asclepias syriaca	Apocynaceae	Common milkweed	3	ASCSYR	1.03e-01
Astragalus agrestis	Fabaceae	Purple milkvetch	5	ASTAGR	1.68e-03
Astragalus canadensis	Fabaceae	Canadian milkvetch	1	ASTCAN	1.17e-02
Brickellia eupatorioides	Asteraceae	False boneset	1	BRIEUP	5.01e-03
Carduus nutans	Asteraceae	Musk thistle	4	CARNUT	9.80e-04
Cerastium arvense	Caryophyllaceae	Prairie chickweed	1	CERAVE	1.25e-03
Cirsium arvense	Asteraceae	Canada thistle	11	CIRARV	1.04e-03
Cirsium floodmanii	Asteraceae	Floodman's thistle	4	CIRFLO	4.20e-04
Dalea candida	Fabaceae	White prairie clover	5	DALCAN	5.08e-02
Dalea purpurea	Fabaceae	Purple prairie clover	5	DALPUR	9.64e-02

Daucus carota	Apiaceae	Queen Anne's lace	1	DAUCAR	2.89e-01
Desmodium illinoense	Fabaceae	Illinois tick trefoil	1	DEIL2	2.74e-02
Doellingeria umbellata	Asteraceae	Flat-top aster	2	DOEUMB	1.07e-02
Echinacea angustifolia	Asteraceae	Purple coneflower	2	ECHANG	1.01e-03
Erigeron philidelphicus	Asteraceae	Daisy fleabane	3	ERIPHI	1.56e-02
Erigeron strigosus	Asteraceae	Prairie fleabane	2	ERISTR	2.31e-03
Euphorbia esula	Euphorbiaceae	Leafy spurge	4	EUES	1.93e-02
Eupatorium maculatum	Asteraceae	Spotted joe-pye weed	2	EUPMAC	1.48e-02
Fragaria virginiana	Rosaceae	Wild strawberry	2	FRAVIR	5.53e-04
Gaillardia aristata	Asteraceae	Common gaillardia	1	GAIAIR	1.07e-03
Gallium boreale	Rubiaceae	Northern bedstraw	4	GALBOR	4.75e-02
Gentiana andrewsii	Gentianaceae	Bottle gentian	1	GENAND	3.14e-03
Geum triflorum	Rosaceae	Prairie smoke	2	GETR	3.01e-04
Glycyrrhiza lepidota	Fabaceae	American licorice	3	GLYLEP	3.62e-02
Grindelia squarrosa	Asteraceae	Curly-cup gumweed	1	GRISQU	-
Helenium autumnale	Asteraceae	Sneezeweed	3	HELAUT	-
Heliopsis helianthoides	Asteraceae	Smooth oxeye	8	HELHEL	1.63e-03
Helianthus maximiliani	Asteraceae	Maximilian sunflower	3	HELMAX	6.44e-04
Helianthus nuttallii	Asteraceae	Nutall's sunflower	5	HELNUT	1.99e-03
Helianthus pauciflorus	Asteraceae	Stiff sunflower	8	HELPAU	5.85e-03
Hypoxis hirsuta	Amaryllidaceae	Common goldstar	3	HYPHIR	8.21e-04
Liatris ligulistylis	Asteraceae	Meadow blazing star	3	LIALIG	3.02e-03
Linaria vulgaris	Plantaginaceae	Yellow toadflax	2	LINVUL	1.64e-03
Lithospermum canescens	Boraginaceae	Hoary puccoon	5	LITCAN	3.76e-03
Lobelia siphilitica	Campanulaceae	Blue cardinal flower	1	LOBSIP	3.85e-03
Lobelia spicata	Campanulaceae	Pale spike lobelia	2	LOBSPI	-
Lycopus asper	Lamiaceae	Rough bugleweed	2	LYCASP	1.54e-02
Lythrum alatum	Lythraceae	Winged loosestrife	1	LYTALA	5.92e-02
Melilotus albus	Fabaceae	White sweetclover	10	MEAL2	4.36e-01
Medicago lupulina	Fabaceae	Black medick	6	MELU	6.47e-03
Melilotus officinalus	Fabaceae	Yellow sweetclover	9	MEOF	2.38e-01
Medicago sativa	Fabaceae	Alfalfa	5	MESA	3.06e-02
Monarda fistulosa	Lamiaceae	Wild bergamot	6	MONFIS	5.55e-02
Mulgedium pulchellum	Asteraceae	Showy blue lettuce	2	MULPUL	5.92e-04

Oxalis violacea	Oxalidaceae	Violet wood-sorrel	4	OXAVIO	7.29e-04
Packera plattensis	Asteraceae	Prairie ragwort	1	PACPLA	-
Pediomelum argophyllum	Fabaceae	Silverleaf scurfpea	5	PEDARG	1.48e-03
Phlox pilosa	Polemoniaceae	Downy phlox	2	PHLPIL	2.74e-03
Ratibida columnifera	Asteraceae	Upright prairie coneflower	6	RATCOL	7.57e-04
Ratibida pinnata	Asteraceae	Gray-headed coneflower	3	RATPIN	6.40e-03
Rosa acicularis	Rosaceae	Wild rose	4	ROSACI	1.40e-04
Rudbeckia hirta	Asteraceae	Black-eyed susan	2	RUDHIR	1.19e-03
Silene vulgaris	Caryophyllaceae	Bladder campion	1	SILVUL	1.58e-03
Sisyrinchium campestre	Iridaceae	Prairie blue-eyed grass	3	SISCAM	1.70e-03
Sisyrinchium montanum	Iridaceae	Strict blue-eyed grass	1	SISMON	-
Solidago canadensis	Asteraceae	Canada goldenrod	1	SOLCAN	2.31e-02
Solidago gigantea	Asteraceae	Giant goldenrod	6	SOLGIG	2.51e-02
Solidago missouriensis	Asteraceae	Missouri goldenrod	5	SOLMIS	4.58e-02
Solidago ptarmicoides	Asteraceae	Upland white goldenrod	1	SOLPHA	3.63e-02
Solidago rigida	Asteraceae	Stiff goldenrod	2	SOLRIG	6.10e-02
Sonchus arvensis	Asteraceae	Field sow thistle	10	SONARV	1.13e-03
Stachys palustris	Lamiaceae	Marsh woundwort	3	STAPAL	7.57e-03
Symphyotrichum ericoides	Asteraceae	White heath aster	5	SYMERI	8.21e-03
Symphyotrichum lanceolatum	Asteraceae	Lance-leaf aster	3	SYMLAN	4.82e-03
Symphyotrichum novae-angliae	Asteraceae	New England aster	1	SYMNOE	2.79e-02
Symphyotrichum sericeum	Asteraceae	Silky aster	1	SYMSER	2.50e-03
Symphoricarpos occidentalis	Caprifoliaceae	Western snowberry	4	SYOC	5.84e-03
Taraxacum officinalis	Asteraceae	Common dandelion	5	TAROFF	2.80e-04
Tragopogon dubius	Asteraceae	Yellow salsify	2	TRDU	7.00e-05
Trifolium repens	Fabaceae	White clover	2	TRIREP	2.10e-02
Trifolium pratense	Fabaceae	Red clover	5	TRPR2	9.11e-02
Verbena stricta	Verbenaceae	Hoary vervain	2	VERSTR	8.61e-03
Verbascum thapsus	Scrophulariaceae	Common mullien	1	VERTHA	2.10e-03
Verbena urticifolia	Verbenaceae	White vervain	1	VERURT	9.01e-03
Viola nephrophylla	Violaceae	Northern bog violet	1	VIONEP	-
Viola pedatifida	Violaceae	Prairie violet	2	VIOPED	3.37e-04
Zizia aptera	Apiaceae	Heart-leaved golden alexanders	5	ZIZAPT	1.57e-01

1255