Computational anatomy and geometric shape analysis enables analysis of complex

craniofacial phenotypes in zebrafish mutants

Kelly M. Diamond[1], Sara M. Rolfe[1,2], Ronald Y. Kwon[3,4], A. Murat Maga[1,5]

[1] Center for Developmental Biology and Regenerative Medicine, Seattle Children's Research

Institute, Seattle, WA, USA

[2] Friday Harbor Marine Laboratories, University of Washington, San Juan, WA, USA

[3]Department of Orthopedics and Sports Medicine, University of Washington, Seattle, WA, USA

[4]Institute for Stem Cell and Regenerative Medicine, University of Washington, Seattle, WA, USA

[5]Division of Craniofacial Medicine, Department of Pediatrics, University of Washington, Seattle,

WA, USA

Abstract

Computational anatomy (CA) approaches start by estimating a canonical 'template' from a

sample of images. This template is then used as a basis for statistical analysis to quantify

structural differences among groups of interest. Due to the complexity of fish skulls, previous

attempts to classify craniofacial phenotypes have relied on qualitative features or 2D

landmarks. In this work we aim to identify and quantify differences in 3D craniofacial

phenotypes in adult zebrafish mutants. We first estimate a synthetic 'normative' zebrafish

template using microCT scans from a sample pool of wildtype animals using the Advanced

Normalization Tools (ANTs). To validate the accuracy of the template and our CA pipeline, we

compared the otolith volumes derived from the CA approach to manually segmented volumes

of the same set of zebrafish. Our overall CA based segmentation volumes are statistically similar

to our manual segmentations and both results show that mutants have larger otoliths than

their wildtype controls. We apply CA to zebrafish with somatic mutations in *bmp1a*, whose

human ortholog when disrupted is associated with Osteogenesis Imperfecta. Our Generalized

Procrustes Analysis separated the bmp1a and wildtype fish along the first two principal

components which collectively explained 31.8% of the variation in the dataset. Compared to

controls, the phenotypic differences in *bmp1a* fish are concentrated around the operculum and

the orbit. Our CA approach seems to offer a potential pipeline for high throughput screening of

complex fish craniofacial phenotypes, especially those of zebrafish which are an important

model system for testing genome to phenome relationships in the study of development,

evolution, and human diseases.

Introduction

The two main drivers of an organism's phenotype are the underlying genetic makeup and the

environment in which an organism is exposed during development. Current best practices for

quantifying the relationship between genotype and phenotype is to compare the phenotype of

known genetic mutants to their wildtype siblings under identical rearing conditions (Houle et

al., 2010; Watson et al., 2020). These methods minimize the environmental effect, allowing for

differences among groups to be associated with a specific genetic mutation. One area where

these methods are being actively developed is with zebrafish and other laboratory fish models

with mutations in genes whose human orthologs when disrupted cause skeletal diseases (Busse

et al., 2019; Kwon et al., 2019; Watson et al., 2020).

Osteogenesis Imperfecta (OI) is a disease of the collagen matrix and results in brittle

bones and skeletal deformities (Busse et al., 2019). Different forms of OI are caused by

mutations in genes that encode for components of the collagen matrix and associated proteins.

One such gene is Bone Morphogenetic Protein 1 (*BMP1*), which encodes for a secreted protein

involved in procollagen processing. Individuals with mutations in *BMP1* exhibit increased bone

mineral density and recurrent fractures (Asharani et al., 2012). In zebrafish, an ortholog of

BMP1 is *bmp1a*, which shows conserved function in osteogenesis across species and is required

for mature collagen generation downstream of osteoblast maturation in bone (Asharani et al.,

2012). *bmp1a* mutants exhibit ectopic accumulation of minerals in bone, causing malformation

and fractures in adult skeletal structures (Asharani et al., 2012). Severe forms of OI are

frequently associated with craniofacial abnormalities (Dagdeviren et al., 2019). Previous work in

bmp1a zebrafish mutants have identified phenotypic patterns across the axial skeleton that

distinguish *bmp1a* mutants from wildtype siblings (Hur et al., 2017). However, due to the

complicated structure of the fish cranial skeleton, craniofacial abnormalities in zebrafish OI

models have mostly focused on qualitative phenotypes (Gistelinck et al., 2018), and little work

has been done to quantify complex cranial phenotype.

The traditional methods for quantifying cranial morphology use homologous landmark points

on 2D images of the lateral view of the head (Sidlauskas, 2008). However, the requirement for

homologous structures limits landmark placement across the skull, and hence may miss the

phenotypic variation in these areas. Here we propose using an atlas-based computational

anatomy (CA) approach to build a template and then using a pseudo-landmark pipeline to

identify areas of the skull that vary among mutant and wildtype fish. Atlas-based approaches

estimate an unbiased anatomical 'template' from a group of images (Guimond et al., 2006), and

then use this template to the basis to assess shape differences among groups of interest

(Ashburner and Friston, 2000). Atlas-based approaches have been used to characterize

phenotypes in many neuroimaging studies in humans, in fetal mice (KOMP2 project) as well as

in the mouse cranial skeleton (Maga et al., 2017; Toussaint et al., 2020). We define pseudo-

landmarks here as landmarks that are not homologous but evenly cover the surface of our fish

skulls. These methods aim to quantify the cranial phenotype associated with mutations in

zebrafish with minimal user intervention so that large scale studies can examine phenotype-

genotype associations in the skeletal system can be accomplished in a high-throughput fashion.

Methods

Zebrafish microCT scans used in this study were acquired as part of a previous study, for full

animal use information involved in that study see Watson et al., 2020. We used a total of 23

wildtype fish from two clutches ("wildtype fish") to build our atlas and used 12 *bmp1a* somatic

mutants ("*bmp1a* fish"), from a single clutch. Somatic *bmp1a* mutants are mosaic for mutations

in *bmp1a*. Watson et al., 2020 performed a comparison of *bmp1a* somatic and germline

mutants and showed that somatic *bmp1a* mutants recapitulate germline *bmp1a* mutant

phenotypes, but possess additional phenotypic variability due to mosaicism. We focused our

analyses on *bmp1a* somatic mutants as they provide a real-world sample of phenotypic

variability likely to be encountered in CRISPR-based reverse genetic screens (Shah et al., 2015;

Watson et al., 2020).

MicroCT Imaging and Atlas Building

Imaging was performed using a vivaCT40 (Scanco Medical, Switzerland) at 21 micron voxel

resolution (55kVp, 145mA, 1024 samples, 500proj/180˚, 200 ms integration time). Files of

individual fish were reconstructed using Scanco software to create 3D volume files for each fish

in the study. To investigate potential asymmetry patterns, we opted to build a symmetrical

atlas of wildtype fish (N=23) by first reflecting all volumes along the sagittal plane. A symmetric

atlas was generated using the antsMultivariateTemplateConstruction2.sh script as provided by

the Advance Normalization Tools (Avants et al., 2014), using the default settings. Atlas building

script initiates with a linear average of all samples, to which all samples are deformable

registered to (Figure 1). The resultant deformation fields are applied to samples, and a new

average is estimated and then used as a new reference for the next step of registrations. Four

iterations were sufficient to obtain a symmetrical and anatomically detailed template.

Atlas validation

First, we review the resultant atlas qualitatively by investigating the spatial arrangement of

bones in 3D rendering. To quantitatively validate the atlas and our computational anatomy

framework, we first created segmentations of individual otoliths from every sample manually

using the open-source 3D Slicer program (Fedorov et al., 2012). We chose otoliths because they

are dense, spread out along the dorsoventral axis of the crania, and do not touch any bones,

which minimized the potential for error in our manual segmentations that serve as the ground

truth data. The otoliths from the atlas are segmented in the same manner. Next, we

deformably register every sample, including the mutants, to our atlas using the ANTsR package,

and apply the resultant transformation field to our atlas otolith segmentation to map them

directly onto the subject space. From this mapping, we calculated the volumes of CA derived

segmentation and statistically compared them to ground truth. All statistical analysis and image

registrations are done using the R extensions of the ANTs ecosystem (Avants, 2020).

Analysis of ZF cranial shape difference in wild-types and mutants.

Because a standard set of cranial landmarks in fish skulls (Sidlauskas, 2008) are too sparse to

adequately represent the complex shape of fish skulls and to capture potential local shape

differences, we opted to use pseudo-landmarks. Pseudo-landmarks are geometrically

constructed points. SlicerMorph's PseudoLMGenerator provides a convenient interface to

estimate a dense point-cloud from a mesh, using its geometry (Rolfe et al., 2021; Figure 1). For

bilaterally symmetric structures, PseudoLMGenerator allows users to define the axis of

symmetry around which pseudo-landmarks are symmetrically reflected, allowing for

asymmetry analyses. More implementation and technical specifics regarding

PseudoLMGenerator can be found in the cited paper above.

We have segmented the skull as a single structure from our atlas, as well as individual

samples using the Segment Editor module in 3D Slicer by exporting the segment as 3D surface

model. We used the Surface Toolbox module in 3D slicer to decimate the model to create a

smoother surface mesh with more uniform vertex distribution. The 3D skull model from our

symmetric atlas was input into PseudoLMGenerator. Using the original mesh geometry and the

sagittal plane as the axis of symmetry, we generated a dense point cloud. One author (KMD)

then went through the pseudo-landmarks and removed the points that were on both jaws and

gills using the SlicerMorph's MarkupEditor. Both of those structures are highly prone to plastic

post-mortem deformation due to handling and preservation, as such they represent confound

non-biological variation. The ALPACA module in SlicerMorph provides a convenient interface to

transfer points on a reference mesh (in this case the symmetric template) to new subjects using

linear and deformable mesh registration (Porto et al., 2020). We used the default settings and

skipped the scaling option to transfer pseudo-landmarks from the atlas to all meshes in our

sample (Figures 1,2).

Analysis

To explore differences between *bmp1a* and wildtype fish, we first ran a Generalized Procrustes

Analysis on the 372 pseudo-landmark points using the GPA module in SlicerMorph (Rolfe et al.,

2021). Procrustes ANOVA on principal components is used to assess the significance of PCA

results using the geomorph package in R (Adams and Otárola-Castillo, 2013). Visualizations

were performed in the GPA module of SlicerMorph and using ggplot in R (Wickham and Chang,

2007).

Results & Discussion

When analyzing otoliths, we did not find significant differences between manually segmented

volumes and atlas segmented volumes (t=-0.912, p=0.363; Figure 3), though there were some

differences between some of the individual otoliths (Table 2; Figure 3). The most apparent

difference between *bmp1a* and wildtype fish is that *bmp1a* fish have larger otoliths than

wildtype fish, especially for the asteriscus, the largest otoliths in the zebrafish (Table 1; Figure

3). In contrast to bone formation in which the mineral phase is primarily hydroxyapatite,

otoliths are formed via an accumulation of calcium carbonate in the acellular endolymph of the

fish inner ear (Payan et al., 2004). Previous work found higher tissue mineral density in *bmp1a*

fish across the axial skeleton (Hur et al., 2017) and this result suggests potential influence of

bmp1a on other pathways associated with mineralized tissues.

Looking next at the results of our pseudo-landmark pipeline, we again find differences in

cranial phenotype between *bmp1a* and wildtype fish. We found significant differences between

bmp1a and wildtype fish for both PC1, which explained 22% of the variation in the data

(F=3.959; Z=1.316; p=0.036), and PC2, which explained 9.8% of the variation in the data

(F=4.528; Z=1.669; p=0.019; Figure 4). When we visualize the variation observed across our

pseudo-landmark points, we see higher instances of variation in the posterior portion of the

operculum, and around the orbit (longer vectors in Figure 5). As we removed pseudo-landmark

points associated with areas of the skull that varied due to preservation or scanning methods,

these represent areas of interest for exploring how phenotype differs between mutant and

wildtype fishes. PC1 shows greater variation around the dorsal portion of the orbit and the right

posterior operculum (green vectors in Figure 5). Whereas PC2 shows greater variation on the

ventral portion of the orbit and left ventral opercula (dark blue vectors in Figure 5). Together

these results provide evidence for phenotypic effects of the *bmp1a* mutation on the cranial

phenotype of zebrafish. Future work should expand the number of families to ensure this is not

unique to this particular family. We have shown how our pipeline can identify areas of greatest

variation among groups of animals. In combination with additional morphological analyses, we

hope this pipeline will enable researchers to better define the links between genotype and

phenotype.

Acknowledgements

We thank members of the Maga and MSB labs for feedback and development of this project.

This project was partly supported by National Science Foundation grants An Integrated

Platform for Retrieval, Visualization and Analysis of 3D Morphology from Digital Biological

Collections (DBI/1759883) and Biology Guided Neural Networks for discovering phenotypic

traits (OAC/1939505) to AMM. RYK was supported by NIH Grant AR074417.

Competing Interests

The authors declare no competing interests

Author Contributions

Contribution of fish and microCT scans: RYK. Conceptualization and methodology: all authors,

Writing: KMD; Editing and approval: all authors.

Data availability

Atlas and pseudo-landmark points are available at:

https://github.com/SlicerMorph/ZF_Skull_atlas/

Literature Cited

Adams, D.C., Otárola-Castillo, E., 2013. Geomorph: An r package for the collection and analysis

of geometric morphometric shape data. Methods Ecol. Evol. 4, 393–399.

Asharani, P. V., Keupp, K., Semler, O., Wang, W., Li, Y., Thiele, H., Yigit, G., Pohl, E., Becker, J.,

Frommolt, P., Sonntag, C., Altmüller, J., Zimmermann, K., Greenspan, D.S., Akarsu, N.A.,

Netzer, C., Schönau, E., Wirth, R., Hammerschmidt, M., Nürnberg, P., Wollnik, B., Carney,

T.J., 2012. Attenuated BMP1 function compromises osteogenesis, leading to bone fragility

in humans and Zebrafish. Am. J. Hum. Genet. 90, 661–674.

Ashburner, J., Friston, K.J., 2000. Voxel-based morphometry - The methods. Neuroimage 11,

805–821.

Avants, B.B., 2020. ANTsR: ANTs in R: Quantification tools for biomedical images version

0.5.7.5.

Avants, B.B., Tustison, N.J., Johnson, H., 2014. Advanced Normalization Tools (ANTs).

Busse, B., Galloway, J.L., Gray, R.S., Harris, M.P., Kwon, R.Y., 2019. Zebrafish: An Emerging

Model for Orthopedic Research. J. Orthop. Res.

Dagdeviren, D., Tamimi, F., Lee, B., Sutton, R., Rauch, F., Retrouvey, J.-M., 2019. Dental and

craniofacial characteristics caused by the p.Ser40Leu mutation in IFITM5. Am. J. Med.

Genet. 176, 65–70.

Fedorov, A., Beichel, R., Kalpathy-Cramer, J., Finet, J., JC, F.-R., Pujol, S., Bauer, C., Jennings, D.,

Fennessy, F., Sonka, M., Buatti, J., Aylward, S., Miller, J., Pieper, S., Kikinis, R., 2012. 3D

Slicer as an image computing platform for the quantitative Imaging Network. Magn Reson

Imaging 30, 1323–1241.

Gistelinck, C., Kwon, R.Y., Malfait, F., Symoens, S., Harris, M.P., Henke, K., Hawkins, M.B., Fisher,

S., Sips, P., Guillemyn, B., Bek, J.W., Vermassen, P., De Saffel, H., Witten, P.E., Weis, M.A.,

De Paepe, A., Eyre, D.R., Willaert, A., Coucke, P.J., 2018. Zebrafish type I collagen mutants

faithfully recapitulate human type I collagenopathies. Proc. Natl. Acad. Sci. U. S. A. 115,

E8037–E8046.

Guimond, A., Meunier, J., Thirion, J., 2006. Average Brain Models : A Convergence Study To cite

this version : Average Brain Models : A Convergence Study.

Houle, D., Govindaraju, D.R., Omholt, S., 2010. Phenomics: The next challenge. Nat. Rev. Genet.

11, 855–866.

Hur, M., Gistelinck, C.A., Huber, P., Lee, J., Thompson, M.H., Monstad-Rios, A.T., Watson, C.J.,

McMenamin, S.K., Willaert, A., Parichy, D.M., Coucke, P., Kwon, R.Y., 2017. MicroCT-Based

Phenomics in the Zebrafish Skeleton Reveals Virtues of Deep Phenotyping in a Distributed

Organ System. Elife 6, e26014.

Kwon, R.Y., Watson, C.J., Karasik, D., 2019. Using zebrafish to study skeletal genomics. Bone

126, 37–50.

Maga, A.M., Tustison, N.J., Avants, B.B., 2017. A population level atlas of Mus musculus

craniofacial skeleton and automated image-based shape analysis. J. Anat. 231, 433–443.

Payan, P., De Pontual, H., Bœuf, G., Mayer-Gostan, N., 2004. Endolymph chemistry and otolith

growth in fish. Comptes Rendus - Palevol 3, 535–547.

Porto, A., Rolfe, S.M., Maga, A.M., 2020. ALPACA: a fast and accurate approach for automated

landmarking of three-dimensional biological structures. bioRxiv.

Rolfe, S., Pieper, S., Porto, A., Diamond, K., Winchester, J., Shan, S., Kirveslahti, H., Boyer, D.,

Summers, A., Maga, A.M., 2021. SlicerMorph: An open and extensible platform to retrieve,

visualize and analyze 3D morphology. bioRxiv.

Shah, A.N., Davey, C.F., Whitebirch, A.C., Miller, A.C., Moens, C.B., 2015. Rapid reverse genetic

screening using CRISPR in zebrafish. Nat. Methods 12, 535–540.

Sidlauskas, B., 2008. Continuous and arrested morphological diversification in sister clades of

characiform fishes: A phylomorphospace approach. Evolution (N. Y). 62, 3135–3156.

Toussaint, N., Redhead, Y., Liu, W., Fisher, E.M.C., Hallgrimsson, B., Tybulewicz, V.L.J., Schnabel,

J.A., Green, J.B.A., 2020. Application of high-resolution landmark-free morphometrics to a

mouse model of Down Syndrome reveals a tightly localised cranial phenotype. BioRxiv.

Watson, C.J., Monstad-Rios, A.T., Bhimani, R.M., Gistelinck, C., Willaert, A., Coucke, P., Hsu,

Y.H., Kwon, R.Y., 2020. Phenomics-Based Quantification of CRISPR-Induced Mosaicism in

Zebrafish. Cell Syst. 10, 275-286.e5.

Wickham, H., Chang, W., 2007. ggplot2: An Implementation of the Grammar of Graphics. GNU

General Public License. https://cran.r-project.org/web/packages/ggplot2. Cran.

Tables and Figures

Table 1. Welch two sample t-test for difference between mutants and wildtype fish for each pair of manually segmented otolith volumes. We provide the mean volumes(x) for mutants and wildtype groups, degrees of freedom (df), test statistic (t), p value (p), and confidence interval (UCL-LCL).

Otolith	x mutant (mm³)	x wildtype (mm³)	df	t	p	UCL	LCL
Left asteriscus	0.044	0.039	15.975	3.383	0.004	0.002	0.008
Right asteriscus	0.044	0.040	14.62	3.232	0.006	0.002	0.007
Left lapilus	0.026	0.025	18.013	2.000	0.061	-0.0007	0.003
Right lapilus	0.026	0.025	17.693	1.554	0.138	-0.0004	0.003
Left sagitta	0.005	0.005	18.665	2.244	0.037	0.0003	0.009
Right sagitta	0.005	0.005	15.375	1.902	0.076	-0.0005	0.008

280 **Table 2.** Welch two sample t-test for difference between manual and atlas-segmented otolith
281 volumes for mutant and wildtype fish. We provide the mean volumes(x) for each method,
282 degrees of freedom (df), test statistic (t), p value (p), and confidence interval (UCL-LCL).

Otolith	group	x atlas (mm³)	x manual (mm³)	df	t	p	UCL	LCL
Left asteriscus	mutant	0.042	0.044	21.971	-0.902	0.377	-0.006	0.044
Left asteriscus	wildtype	0.037	0.039	25.117	-1.998	0.057	-3.995	6.056
Right asteriscus	mutant	0.043	0.044	21.997	-0.618	0.543	-0.005	0.003
Right asteriscus	wildtype	0.037	0.040	20.891	-2.744	0.012	-0.005	-0.0007
Left lapilus	mutant	0.024	0.026	22.000	-3.074	0.006	-0.005	-0.0009
Left lapilus	wildtype	0.022	0.025	25.581	-5.438	<0.001	-0.004	-0.002
Right lapilus	mutant	0.023	0.026	20.94	-2.490	0.021	-0.005	-0.0005
Right lapilus	wildtype	0.021	0.025	24.294	-5.347	<0.001	-0.005	-0.002
Left sagitta	mutant	0.006	0.005	17.619	1.998	0.061	-0.00004	-0.002
Left sagitta	wildtype	0.005	0.005	19.564	0.799	0.434	-0.0003	0.0007
Right sagitta	mutant	0.006	0.005	16.603	1.934	0.070	-0.0006	0.002
Right sagitta	wildtype	0.005	0.005	15.959	0.236	0.816	-0.0005	0.0005

Figure 1. Pipeline for atlas building, pseudo-landmark generation, and transferring pseudo-

landmarks to individual fish. Starting with microCT scans of wildtype fish ANTs, uses a series of

rigid, affine, and deformable registrations to create an average image, or Atlas. The

PseudoLMGenerator tool in SlicerMorph was used to place 372 pseudo-landmarks on the atlas.

The ALPACA tool in SlicerMorph was used to transfer points from the atlas to wildtype and

bmp1a fish for comparisons between groups.

Figure 2. Left lateral view of fish meshes with pseudo-landmarks generated from the

PseudoLMGenerator and transverse using the ALPACA modules in SlicerMorph. Name color

represents groupings, the atlas is in blue, wildtype fish are in grey and *bmp1a* fish are in orange.

Figure 3. Boxplots of segmented otolith volumes. Wildtype (grey) and *bmp1a* mutant (orange)

boxplots are shown for each of the six otoliths for both the (A) manually segmented volumes

and (B) atlas segmented volumes. For each otolith, mutants have larger median volumes than

wildtype fish (Table 1). Insets show the dorsal view of otolith segments (asterisk in blue, lapilus

in orange, and sagitta in grey) with lighter colors on the left side of the head, and darker colors

on the right.

Figure 4. First two principal components from Generalized Procrustes Analysis. Color shows

bmp1a (orange) or wildtype (grey) fish as well as the location of the atlas (blue) in

morphospace. Larger circles show average position for each group. *bmp1a* and wildtype fish

separate along both axes.

Figure 5. Landmark vector plots for first two principal components from GPA analysis. Vectors

show magnitude and direction of variation along PC1(green) and PC2 (dark blue). Panels show

(A) left lateral, (B) right lateral, (C) dorsal, and (D) ventral orientation of the zebrafish crania.

Click here for animation.

320