Chemical ecology of a tripartite symbiosis

Chemical ecology of a tripartite symbiosis

Nicholas C. Mucci[1], Katarina A. Jones[2], Mengyi Cao[3], Michael R. Wyatt II[4], Shane Foye[5], Sarah
Kauffman[1], Michela Taufer[4], Yoshito Chikaraishi[6], Shawn Steffan[5, 7], Shawn Campagna[2,8], Heidi
Goodrich-Blair[1,3#]

[1]Department of Microbiology, University of Tennessee-Knoxville, Knoxville TN, USA

[2]Department of Chemistry, University of Tennessee-Knoxville, Knoxville TN, USA

[3]Department of Bacteriology, University of Wisconsin-Madison, Madison WI, USA

[4]Department of Electrical Engineering and Computer Science, University of Tennessee-
Knoxville, Knoxville TN, USA

[5]Department of Entomology, University of Wisconsin-Madison, Madison WI, USA

[6]Department of Biogeochemistry, Japan Agency for Marine-Earth Science and Technology,
Yokosuka 237-0061, Japan

[7]US Department of Agriculture, Agricultural Research Service, Madison WI, USA

[8]Biological and Small Molecule Mass Spectrometry Core, University of Tennessee-Knoxville,
Knoxville TN, USA

#Corresponding author: Heidi Goodrich-Blair, hgblair@utk.edu

ABSTRACT

Microbial symbiotic interactions, mediated in part by small molecule signaling, drive

physiological processes of higher order systems, including the acquisition and consumption of

nutrients that support symbiotic partner reproduction. Advances in metabolic analytic

technologies provide new avenues to examine how chemical ecology, or the conversion of

existing biomass to new forms, changes over a symbiotic lifecycle. Here we examine such

processes using the tripartite relationship involving the nematode host *Steinernema*

carpocapsae, its obligate mutualist bacterium, *Xenorhabdus nematophila*, and the insects they

infect together. The nematode infective juveniles infect insects into which they release bacteria

that help suppress insect immunity and kill the insect. The nematode-bacterium pair consume

the insect cadaver and reproduce until nutrients are depleted, causing a new generation of

infective juvenile nematodes, colonized by the bacterial symbiont, to leave the cadaver in

search of insect prey. To begin to understand the processes by which insect biomass is

converted over time to either nematode or bacterium biomass, we took a three-pronged

approach integrating information from trophic, metabolomics, and gene regulation analyses.

Trophic analysis established bacteria as the primary insect consumers, with nematodes at a

trophic position of 4.37, indicating consumption of bacteria and likely also other nematodes.

Metabolic changes associated with bioconversion of *Galleria mellonella* insects were assessed

using multivariate statistical analyses of metabolomics datasets derived from sampling over an

infection time course. Statistically significant, discrete phases were distinguishable from each

other, indicating the insect chemical environment changes reproducibly during bioconversion.

Tricarboxylic acid (TCA) cycle components and amino acids such as proline and leucine were

significantly affected throughout the infection. Hierarchical clustering revealed a similar

molecular abundance fluctuation pattern for nucleic acid, amino acid, and lipid biosynthesis

metabolites. Together, these findings contribute to an ongoing understanding of how symbiont

associations shape chemical environments.

IMPORTANCE

The biological world depends on microbial processes, including symbiotic exchanges

among mutualists and antagonists. Technical advances have enabled investigation of microbial

trophic positions within communities and the identification of metabolites exchanged between

microbes and their hosts. These tools were applied to the tripartite *Xenorhabdus* bacteria-

Steinernema nematode-*Galleria* insect symbiosis. Trophic analyses demonstrate the primary

consumer of the insect are the bacteria, and in turn the nematode consumes the bacteria. This

suggests the *Steinernema-Xenorhabdus* mutualism is a form of agriculture in which the

nematode cultivates their bacterial food source by inoculating them into insect hosts.

Metabolomics analysis revealed a shift in biological material throughout progression of the

lifecycle: active infection, insect death, and conversion of cadaver tissues into bacterial biomass

and nematode tissue. We show that each phase of the lifecycle is metabolically distinct, with

significant differences in tricarboxylic acid cycle and amino acid metabolism. Our findings

demonstrate that symbiotic lifecycles can be defined by reproducible stage-specific chemical

signatures, enhancing our broad understanding of metabolic processes that underpin a three-

way symbiosis.

INTRODUCTION

Symbiotic interactions are ubiquitous in biological systems and have shaped the evolution of life {Margulis, 1981}. These long-term, intimate associations are driven by small molecule signaling between partners. Bacterial populations establish diverse and expansive metabolite-mediated signaling networks that control gene expression and downstream behaviors, such as biofilm formation and the production of host-interacting effectors {Camilli, 2006; Worthington, 2012; Chaston, 2012}. A common mechanism by which bacteria sense and transduce metabolic signals is through transcription factors whose DNA binding affinity or specificity is modulated by binding metabolite ligands. For instance, LysR-type transcription factors, which are conserved across proteobacteria, are characterized by a conserved *N*-terminal DNA-binding domain and a *C*-terminal domain that varies among LysR-type regulator homologs. The latter domain is responsible for ligand metabolite binding and dictates the response specificity of the transcription factor {Budnick, 2020; Henikoff, 1988; Maddocks, 2008}. In a range of bacteria, LysR-type regulators modulate various phenotypes, including virulence, nutrient uptake and metabolic homeostasis, motility, quorum sensing, and antibiotic resistance {Abdelhamed, 2020}. Another diverse family of transcription factors, feast/famine regulatory factors like leucine-responsive regulatory protein (Lrp), will bind and dimerize with amino acids in response to nutrient levels and globally induce transcriptional changes {Yokoyama, 2006}.

Given the key function of metabolites in communicating information about intracellular and extracellular environmental conditions, examining their identities and abundances is critical to understanding biological systems. Metabolomics has enabled such studies and has been used to detect specific small molecules that drive essential cellular processes and inter-kingdom signaling {Tang, 2011; Clarke, 2008; Phelan, 2014}. Further, it is being applied to more complex ecosystems comprising multi-species microbiota colonizing a host {Dong, 2019}. However, to date such studies have been primarily focused on binary conditional comparisons between

treatments, or on single, snapshot sampling of complex interactions. Here, to gain insights

temporal changes in metabolic pathways that occur in complex ecosystems, a longitudinal

analysis of metabolic profiles was conducted in closed ecosystem in which biomass is being

reproducibly converted from one type of living organism to another. The closed ecosystem

comprised an individual insect infected with an entomopathogenic nematode (EPN) and

bacterium (EPNB) pair.

EPNs of the genera *Steinernema* and *Heterorhabditis* associate with mutualistic bacteria

in the genera *Xenorhabdus* and *Photorhabdus,* respectively. An infective juvenile (IJ) stage of

EPN carry their mutualistic bacteria in their intestine as they dwell in the soil seeking insect

hosts to infect. Upon infection, the bacteria are released into the insect blood cavity and

together the nematode and bacterium kill and consume the insect for their own reproduction

before developing into the bacteria-colonized infective stage again to repeat the cycle

{Richards, 2009; Stock, 2019}. EPNBs have been applied as insecticide alternatives to promote

agricultural productivity and to help prevent transmission of insect diseases like dengue and

West Nile virus {Lacey, 2012; Javal, 2019; da Silva, 2020}.

In this study, consumption and bioconversion of the insect *Galleria mellonella* by the

EPN *Steinernema carpocapsae* and its mutualistic bacterial symbiont, *Xenorhabdus*

nematophila was examined from a metabolic perspective. *G. mellonella* is used for laboratory

isolation and propagation of EPNB, is a model host to understand virulence of a variety of

microbial pathogens {Tsai, 2016; Ramarao, 2012} and has a characterized metabolome {Killiny,

2018}. The *S. carpocapsae-X. nematophila* pair was chosen due to the wealth of information

available about them from molecular, cellular, and genetic studies {Martens, 2004; Stock, 2019}.

For example, it is known that *X. nematophila* bacterial effectors and natural products suppress

insect immunity, kill insect blood cells, degrade insect tissues, and defend the insect cadaver

from opportunistic competitors {Herbert, 2007}. Also, *X. nematophila* bacteria are essential for

S. carpocapsae reproduction; in the absence of bacteria fewer nematode IJs emerge from

insect cadavers after reproduction {Mitani, 2004}. Expression of effectors and physiological

adaptation to changing host environments is controlled in *X. nematophila* by transcriptional

regulators that are predicted to sense and respond to prevailing metabolic conditions {Richards,

2008}. For instance, the LysR-type regulator LrhA that is necessary for *X. nematophila* virulence

and controls expression of an extracellular phospholipase that is necessary for insect

degradation {Richards, 2008; Richards, 2009; Richards, 2010}, the sigma factor RpoS that is

necessary for colonizing the IJ stage of the nematode {Vivas, 2001}, the two-component system

CpxRA, and the leucine-responsive regulatory protein Lrp, both of which are necessary for

normal virulence and mutualism behaviors {Herbert, 2007; Herbert-Tran, 2009a; Herbert-Tran,

2009b, Cowles, 2007; Herbert, 2007}, NilR, a lambda like repressor family transcription factor

that negatively regulates genes necessary for nematode colonization {Cowles, 2006} and the

two-component system OmpR/EnvZ that negatively controls *X. nematophila* swarming motility

behavior and exoenzyme production {Park and Forst, 2006}. Further, *X. nematophila* displays

phenotypic heterogeneity with respect to behaviors important for adaptation to host

environments. For instance, "primary form" [1°] *X. nematophila* can be distinguished from

"secondary form" [2°] by its motility, antibiotic and natural products secretion, and hemolytic and

lipolytic activities and additional phenotypic variants arise over the course of insect infection

{Volgyi, 1989; Givaudan, 1995; Cambon, 2019}.

The goal of this study was to begin to understand the overall metabolic transformations,

or bioconversion processes occurring within a closed yet complex biological ecosystem. [15]N

isotopic enrichment analyses were performed to establish the relative trophic positions of the

insect, *G. mellonella*, the nematode, *S. carpocapsae*, and the bacterium, *X. nematophila*, so that

the relative roles in bioconversion of each ecosystem member could be established. Then, a

metabolomics analysis using an ultra-high performance liquid chromatography high-resolution

mass spectrometry (UHPLC-HRMS) metabolomics technique was conducted over a 16-day

time course after *S. carpocapsae-X. nematophila* infection of *G. mellonella*, encompassing a

complete bioconversion of insect tissues to the bacterial-colonized progeny IJs that emerged

from the insect.

RESULTS

Trophic analysis reveals *S. carpocapsae* nematodes directly feed on *X. nematophila*
bacteria *in vitro* and *in vivo*:

***In vitro* trials**

The trophic identities of the entomopathogenic nematode (*Steinernema carpocapsae*), its

bacterial symbiont (*Xenorhabdus nematophila*), and their host insect were measured empirically

based on ^{15}N isotopic enrichment of amino acids (Table 1 and File S1). To this end, it was first

necessary to establish that the degree of ^{15}N-enrichment between the consumers (nematodes,

bacteria) and their respective diets (e.g., agar growth media, bacteria, or the insect) was

consistent with past studies of inter-trophic enrichment (Chikaraishi, 2009; Chikaraishi, 2011;

Steffan 2013; Steffan, 2015). This past work relied upon compound-specific isotopic analyses of

select amino acid pools, particularly the degree of ^{15}N-enrichment between two amino acids—

glutamic acid (glu) and phenylalanine (phe). The differential enrichment between these two

amino acids provides a measure of inter-trophic enrichment, which is largely attributable to an

organism's assimilation of dietary amino acids (Chikaraishi, 2007; Chikaraishi, 2009; Takizawa,

2020). Such inter-trophic enrichment has been referred to as the trophic discrimination factor

(TDF$_{glu-phe}$), and in carefully controlled feeding studies among diverse consumer groups in the

Animalia, Fungi, and Bacteria, the TDF$_{glu-phe}$ has averaged approximately 7.2‰ (Steffan, 2015;

Steffan, 2017). Here, following controlled-feeding *in vitro* trials, the nematodes and bacteria

were shown to have both registered TDF$_{glu-phe}$ values in line with past findings (Fig. 2A).

Specifically, the mean (± SE) TDF$_{glu-phe}$ value exhibited by nematodes cultured on bacterial

lawns was 7.41 ± 0.22‰ (*N* = 14). When parsed by nematode stage, the TDF$_{glu-phe}$ values of

adult and infective juvenile nematodes were, respectively, 6.96 ± 0.16‰ (*N* = 8) and 8.02 ±

0.36‰ ($N = 6$). Nematodes fed exclusively a diet of homogenized insect biomass produced a

TDF of 7.38 ± 0.05‰ ($N = 3$). Collectively, the nematode TDF was 7.40 ± 0.18‰ ($N = 17$). The

bacterial symbiont, *Xenorhabdus*, which had been cultured on agar growth media, registered a

$TDF_{glu-phe}$ of 6.53 ± 0.20‰ ($N = 6$). The mean $TDF_{glu-phe}$ across both the nematodes and bacteria

in this food-chain was 7.18 ± 0.16‰, which did not represent a significant departure from the

generalized 7.2‰ $TDF_{glu-phe}$ benchmark (t_{22} = -0.14, P = 0.893). Given the degree of inter-trophic

enrichment exhibited in the nematodes and bacteria, these consumer groups were consistent

with the enrichment patterns of heterotrophs across terrestrial, marine, and freshwater systems,

allowing for trophic position estimation using established isotopic protocols (Chikaraishi, 2007;

Chikaraishi, 2009; Chikaraishi, 2011; Steffan, 2013; Steffan, 2015; Steffan, 2017; Takizawa,

2020).

Using compound-specific isotopic analysis of amino acids, the trophic identities of

consumers and their respective diets within the *in vitro* food-chain were measured. At the base

of the food-chain, the agar growth media registered a trophic position ($TP_{glu-phe}$) of 1.0 ± 0.04‰

($N = 3$), and the bacteria feeding upon the agar registered at 1.9 ± 0.03‰ ($N = 6$), which

represented approximately one trophic level higher than their diet. Correspondingly, the adult

and infective juvenile nematodes that had fed upon the bacteria registered, respectively, at 2.90

± 0.02‰ ($N = 8$) and 3.0 ± 0.06‰ ($N = 6$), which, as predicted, was exactly one trophic level

higher than their diet. The homogenate of insect biomass was measured at 2.2 ± 0.02‰ ($N = 6$),

and the nematodes feeding exclusively on this homogenate registered at 3.2 ± 0.01‰ ($N = 4$),

which again demonstrated that when the nematodes consumed a given diet, they registered

one trophic level higher. The *in vitro* food-chain effectively compartmentalized each consumer

group and thereby provided a means to confirm that when the nematodes or bacteria consumed

a given diet, their isotopic compositions enriched consistently and produced predictable trophic

position estimates.

Figure 2: Trophic analyses reveal *Steinernema* nematodes feed on *Xenorhabdus* bacteria A) *In vitro* and B) *In vivo*. Trophic isoclines are represented via numeric $TP_{glu\text{-}phe}$ % ratios. Specific bacterial cultures or animals are displayed as the different shapes shown in the figure legends.

In vivo trials

These findings allowed for the subsequent *in vivo* trials involving insect cadavers (Fig. 2B, Table 1 and File S1). As reported above, the mean $TP_{glu\text{-}phe}$ of an uncolonized insect cadaver was 2.2 ± 0.02 ($N = 6$). When the insect was colonized by bacteria alone, the $TP_{glu\text{-}phe}$ of the insect-

195 bacteria complex was 2.5 ± 0.03 (N = 3). This complex represented the blending of consumer

196 and diet (as described in Steffan, 2017), wherein the consumer (i.e., the *Xenorhabdus* bacterial

197 population) was suffused within and throughout its diet (the insect cadaver). Given that both the

198 bacterial and insect biomass were available within the cadaver, this established the basis for the

199 question as to what a developing nematode would consume/assimilate within the cadaver. The

200 diet of the nematodes (i.e., the insect-bacterial complex) was measured at ~2.5, thus if the

201 nematodes within the cadaver fed randomly on all available substrates, the nematode $TP_{glu-phe}$

202 would be expected to be ~3.5 (i.e., ~2.5 + 1.0). However, the infective juveniles emerging from

203 the cadavers registered a $TP_{glu-phe}$ of 4.6 ± 0.08 (N = 5), a full trophic level higher than expected.

204 **Table 1. Summary of *in vitro* and *in vivo* trophic measurements.**

Consumer[a]	Diet type[b]	$TDF_{glu-phe}$[c]	$TP_{expected}$[c]	$TP_{glu-phe}$[c]
in vitro growth conditions				
None	Yeast-soy lipid agar (YE-YS)	NA	1.0	1.0
Xenorhabdus	Yeast-soy broth (YE-YS)	6.53	2.0	1.9
Steinernema (adults)	*Xenorhabdus* on yeast-soy lipid agar (YE-YS)	6.96	3.0	2.9
Steinernema (IJs)	*Xenorhabdus* on yeast-soy lipid agar (YE-YS)	8.02	3.0	3.0
in vivo growth conditions				
None	*Galleria* (base of food-chain)	NA	2.0	2.2
None	*Galleria* + PBS buffer (positive control)	NA	2.0	2.2
None	Yeast-soy broth (YE-YS)	NA	2.0	2.2
Xenorhabdus (measured as *Galleria-Xenorhabdus* complex)	*Galleria*	NA	2.5	2.5
Steinernema IJs	*Galleria-Xenorhabdus* complex	NA	3.5	4.6

205 [a]Each organism in the ecosystem was assessed for its trophic position as a consumer under controlled *in vitro* conditions

206 or *in vivo* within an insect cadaver. None indicates that the condition tested was diet only, no consumer.

208   The data above demonstrated that adult *Steinernema* nematodes consume their

209 mutualistic bacteria. During this stage, *Xenorhabdus* bacteria colonize the anterior intestinal

210 caecum. To determine if this colonization influences the ability of *Steinernema* nematodes to

211 consume *Xenorhabdus*, the $TP_{glu-phe}$ of adult *Steinernema* cultivated on lawns of either wild type

212 *Xenorhabdus*, or a non-colonizing mutant (Δ*SR1*) was assessed {Cowles, 2008}. Nematodes

213 had the same $TP_{glu-phe}$ regardless of the colonization proficiency of the bacterial diet, indicating

214 that colonization is not required for nematode direct feeding on its symbiotic bacteria (File S1).

216 ***X. nematophila* transcriptional control of metabolic pathways**

The trophic analyses described above establish *X. nematophila* bacteria as the linchpin

organism in the closed ecosystem, responsible for direct consumption of the insect tissue and

serving as a primary food source for its mutualistic host *S. carpocapsae.* To gain insights into

the metabolic pathways utilized by *X. nematophila* in performance of these functions, the global

regulons of several transcription factors was determined using an exploratory microarray

analysis, portions of which have been reported elsewhere (Fig. 2 and Table S1, and File S2)

{Lu, 2011; Lu, 2012; Chaston, 2011; Hussa, 2015}. Microarray analyses were conducted on

mutants lacking genes encoding the transcription factors LrhA, RpoS, NilR, and Lrp, each of

which has a defect in one or more aspects of the *X. nematophila* life cycle {Vivas, 2001; Cowles,

2006; Cowles, 2007; Richards, 2008}. In addition, since the primary to secondary form

phenotypic variation globally influences host-interaction phenotypes, the transcriptional profiles

of these variants were examined from a metabolic perspective. The mutant and secondary form

cells were each compared to their wild type parent or primary form, respectively, using a 2<|fold

change| significance cutoff for differences in transcript levels.

The number of differential transcripts are a fraction of the 3733 averaged total expressed

chromosome ORFs among the strains. The number of genes with differential transcript

abundance in mutant/secondary form compared to wild-type/primary was highest in Δ*lrhA* at

396 genes (10.6%), followed by Δ*lrp*, secondary form, Δ*rpoS*, and Δ*nilR* with 273 (7.3%), 159

(4.3%), 157 (4.2%), and 83 (2.2%) differentially abundant transcripts, respectively (Fig. 2A). The

proportion of genes categorized as being involved in metabolic activity varied amongst the

strains. Through KEGG annotation, the highest proportion of differentially-expressed genes

categorized as metabolic was observed in the Δ*rpoS* strain, at 38.9%, with Δ*lrhA*, Δ*lrp*,

secondary form, and Δ*nilR* having metabolic-related activities at 18.9%, 13.6%, 10.7%, and

9.8%, of the differentially expressed genes in the respective strain. Differential transcript overlap

was observed while comparing the 5 strains (Supplemental figure 1). The largest overlap

between 2 strains was for the secondary form and Δ*lrp* mutant, consisting of mostly amino acid

(*xncB*, the aminotransferase XNC1_2154) and lipid biosynthesis (*fabG*, *xncL*) genes. *X.*

nematophila lacking *lrp* are phenotypically secondary form {Cowles, 2007}. Another large

overlap was observed between Δ*lrp* and Δ*nilR*, which synergistically repress nematode

colonization {Cowles, 2006}. The overlap regulation includes the phosphotransferase

system/ascorbate metabolic (XNC1_2826-2828) genes and prokaryotic defense system genes

(XNC1_3717-3719, 3724, and 3931).

Functional analysis of the differentially expressed metabolic transcripts was performed

through KEGG annotation using BlastKOALA (KEGG Orthology and Links Annotation).

Sequences were aligned against a nonredundant set of prokaryotic KEGG genes using BLAST

searches (Kanehisa, 2016). Consistent with the KEGG annotation analysis noted above, of the

strains tested Δ*rpoS*, Δ*lrhA*, Δ*lrp* were the most strongly impacted with respect to metabolic

pathway transcripts. These three strains displayed differences in carbohydrate and amino acid

metabolic regulation (Fig. 2A). Branches of carbohydrate metabolism, like propanoate, pentose

and glucuronate, and glyoxylate metabolism were impacted by the Δ*lrhA* and Δ*rpoS* mutations

(Fig. 2B). Inositol phosphate metabolism was impacted in Δ*rpoS* and Δ*lrp* strains, while

pyruvate metabolism was impacted in the Δ*lrhA* strain. Butanoate metabolism, a branch of

carbohydrate metabolism where the amino acid ornithine is converted into short-chain fatty

acids, was commonly interrupted for all mutant strains except Δ*nilR*. Amino acid biosynthetic

pathway transcripts were differently regulated between these three strains, with tyrosine, and

the alanine, aspartate, and glutamate biosynthetic pathways similarly disrupted. Histidine and

valine metabolism was uniquely altered by the Δ*rpoS* mutation, glycine, serine, and threonine

metabolism was uniquely altered by the Δ*lrhA* mutation, and phenylalanine metabolism was

uniquely altered by the Δ*lrp* mutation. To investigate the impacts these pathways and others

have on the *Xenorhabdus-Steinernema* lifecycle, a time course metabolomics experiment was

designed to measure the relative quantities of metabolites within them.

Figure 2: Differentially expressed transcripts between *X. nematophila* mutants and their broad functional categorization. A) Quantification of the number of differentially regulated transcripts and how many are considered metabolic (light grey), as determined by KEGG annotation. |Signal fold change| > 2 was used as a cutoff for significance. BlastKOALA functional categorization of the differential metabolic transcripts are adjacent. The color legend is organized by having the most common category listed first. B) Breakdown of the specific amino acid and carbohydrate metabolism pathways that were affected by the mutations, compared among each strain, with # of genes listed. Positive genes represent transcripts higher in the mutant relative to WT, negative genes represent transcripts lower in the mutant relative to WT.

The metabolomic profile of EPNB-infected *G. mellonella*: an overview

Having established that *X. nematophila* bacteria consume infected insect tissue, and in

turn the bacteria are consumed by reproducing and developing nematodes, the temporal

dynamics of metabolic profiles associated with these processes were examined. *G. mellonella*

were infected with *S. carpocapsae* infective juveniles colonized by *X. nematophila* bacteria.

Weights of the whole insect samples were relatively similar (Table S2). Insects were sampled

over a 16-day time-course. As expected, insects began to die by Hour 24 after infection, and

both living and dead insects were sampled at that time point. *X. nematophila* will have

reproduced rapidly in order to suppress the insect immune response and release toxins to kill

the insect host. By the second day post-infection all insects had succumbed to infection and

were dead. Consistent with the initial degradation of the insect cadaver by bacteria predicted by

the trophic analysis conducted above, the nematodes began to reproduce by Day 4, 3 days

after the insects had died from infection (Fig. 3). At Day 7 post-infection, insect cadavers were

placed in a collection trap to encourage the emergence of progeny *S. carpocapsae* IJs. Adults

and IJs were observed at this stage. This is when the 2nd generation of nematodes begin to

emerge, and endotokia matricida is occurring when some of these juveniles will consume their

parents for nutrients. By Day 16 the insect cadavers were largely consumed, and most

remaining IJs will have exited. On Day 16 the *S. carpocapsae* IJs, colonized by *X. nematophila*

symbionts, were collected from the water trap and analyzed with the other samples. The

proportions of nematodes observed at these stages are consistent with previous observations,

in which there are more adults and juveniles during the middle phase compared to IJs which

changes to high numbers of IJs into the late phase {Cao, 2020}. The above observations led us

to design a time course metabolomics experiment in which samples were divided into 3 major

time frames: the early infection, characterized by bacterial replication and the killing of the insect

host, middle infection, characterized by nematode reproduction and nutrient conversion of the

cadaver, and late infection, characterized by nutrient depletion and IJ emergence.

Figure 3: Key moments in the EPNB lifecycle mapped onto important molecules are indicative of the bioconversion of the insect cadaver. The top 15 VIPs>1 metabolites averaged relative abundances were grouped together into 3 categories: purine and pyrimidine components, amino acid components, and other important molecules. Relative metabolite abundance in log scale is displayed on the y-axis of the line graphs.

Metabolites were extracted from individual insects sampled over the infection time

course and analyzed using an untargeted UHPLC-HRMS method. The untargeted metabolic

profiling analysis revealed 13,748 spectral features. Through the mass spectrometric

measurements, a total of 170 of these features were identified based on comparison to known

exact mass-to-charge (*m/z*) ratio and retention times from a database of central energy

metabolites (File S3). Another 3,138 unidentified spectral features were included in the analysis

and putatively annotated based on their exact masses compared to a *Xenorhabdus* secondary

metabolite database (File S4). This database serves as a rich repository to explore secondary

metabolite temporal flux in the tripartite ecosystem of insect, bacteria, and nematode.

Multivariate data analysis shows metabolic profile gradient corresponding to infection

progression

Partial least squares-discriminant analysis (PLS-DA) was performed to observe gross

chemical environment changes over time of insect bioconversion to nematode-bacterium

complex, when combining all detected metabolite data. A three-dimensional PLS-DA plot shows

a progression of distinct metabolic profiles from uninfected insects (black circles) to insects in

which bacteria and nematodes are reproducing (red and yellow gradients), and finally to fully

consumed insects (green gradients) from which bacterial-colonized infective juvenile

populations are emerging (Fig. 4A). Component 1 is 41.1% and contributes the most

significantly to the separation observed in the PLS-DA plot.

To examine which metabolites are responsible for most of the variation represented by

the PLS plots, Variable Importance in Projection (VIP) values for component 1 were calculated.

VIP is a weighted sum of squares of the PLS loadings that considers the amount of explained Y-

variation in each dimension. A VIP score >1 indicates that the metabolite significantly

contributed to time point differentiation. Most of the VIP>1 metabolites exhibited a bimodal

pattern, going from very low in the uninfected insect, to rising in the early bacterial replication

phase, to dropping during the middle nematode reproduction phase, and finally rising very high

in the late nutrient deplete phase (Fig. 4B and File S5). Overall, these metabolites were involved

in nucleotide and nucleoside biosynthesis, NAD^+ biosynthesis, and iron acquisition. These

included the purine and pyrimidine metabolites 7-methylguanosine, guanine, deoxyinosine,

uridine, deoxyuridine, and deoxycytidine. Other top metabolites include kynurenic acid and

anthranilate which are precursors to NAD^+ synthesis. Of the top 15 VIPs, ascorbate was the

only molecule to exhibit a decreased abundance over time, dropping from very high abundance

to very low later in the time course. This vitamin is necessary for neuron development and could

be salvaged from the cadaver to build the nematode nervous systems {May, 2012}.

Kynurenic acid is an intermediate in the kynurenine pathway and is a way for organisms

to synthesize NAD^+ if they cannot *de novo* synthesize the compound through encoding a

quinolinate phosphoribosyltransferase (QPRTase) {Katsyuba, 2018}. *S. carpocapsae*, like *C.*

elegans, lacks a standard QPRTase but encodes the uridine monophosphate

phosphoribosyltransferase (*umps-1*) which synthesizes NAD^+ from the kynurenine pathway

{Rougon-Cardoso, 2016; Lee, 2020}. Significant flux of this metabolite throughout the lifecycle

could be a metabolic signature nematode NAD^+ production.

Phenylacetic acid is a uremic toxin that builds up in kidney patients and is the product of

bacterial metabolisms. In *P. aeruginosa* phenylacetic acid (PAA) accumulates at high cell

density and inhibits the Type III Secretion System (T3SS), which is toxic to host cells {Wang,

2013}. *X. nematophila* does not have a T3SS but does have the evolutionarily related flagellar

export apparatus. The transcription factor Lrp positively regulates the flagellar regulon, as well

as the gene encoding the XlpA lipase, an enzyme associated with the ability of *X. nematophila*

to support *S. carpocapsae* reproduction {Richards, 2008; Richards, 2010}. Although not

detected by microarray analysis, quantitative reverse transcriptase analyses indicate that the

transcription factor LrhA also positively regulates *xlpA* expression {Richards, 2008}. The

relatively higher intensity of phenylacetic acid at later stages of insect bioconversion may signal

inhibition of secretion of bioconversion enzymes.

2,3-dihydroxybenzoate is a compound involved in siderophore biosynthesis non-

ribosomal peptide biosynthesis of siderophore, highlighting a potential importance in iron

acquisition from the cadaver. Iron itself does not appear to be limited in the cadaver but it needs

to be harvested by the bacteria to aid in nematode reproduction {Ciche, 2003}. *X. nematophila*

does not encode *entA* or *entB*, genes that participate in the conversion of 2,3-

dihydroxybenzoate to the enterobactin siderophore pathway {Latham, 2014}. Siderophore

production is necessary for antibiosis in the closely related *Photorhabdus-Heterorhabditis* EPNB

symbiosis, which has characterized *phb* genes that are homologous to the *ent* genes {Ciche,

2003}. The *Photorhabdus phb* genes encode proteins that sequester iron from the cadaver to

fend off soil-dwelling bacterial colonizers that exploit the cadaver, and *X. nematophila* does not

have any homologs to these genes. *X. nematophila* does not completely dominate the bacterial

community of the insect host, suggesting that other taxa such as *Alcaligenes* are possibly

responsible for this concentration shift {Cambon, 2020}.

Figure 4: Distinct chemical environments occur during bioconversion of an insect

cadaver by *S. carpocapsae* and *X. nematophila*. A) Three-dimensional partial least squares-

discriminant analysis (PLS-DA) of time course infection metabolic profiles grouped according to

stage of infection: uninfected (black) and early (red gradient), middle (yellow gradient), and late

infected insects (green gradient). Components contributing to the separation of the profiles are

listed (in %) on the axes. B) The top 15 VIPs contributing to component 1 are listed, where the

relative abundance shifts over the time course shown in a heatmap on the right. The numbers

on top of the heatmap show the time phases: 0 (uninfected), 1 (early), 2 (middle), and 3 (late).

To identify patterns and groups of metabolite abundance changes over time, hierarchical

clustering was performed to reveal groups of metabolites that exhibit similar concentration

changes over the time course of bioconversion. A dendrogram of all 170 identified metabolites

was generated using the absolute value of the spearman correlation between molecular

concentrations, where distance between molecules is defined as 1-|rs| with rs as the spearman

rank correlation between time course data points of said molecules (Fig. 5A). Metabolite

concentration averages were taken for the four time-phases defined: uninfected, early, middle,

and late infection. Metabolite clusters were visualized in a heatmap that displays their pairwise

correlation between each molecule (Supplemental Figure 2). A heatmap that shows the

metabolite clusters, separated by black bars, with the molecule trends in concentration change

over time was generated (Fig. 5B). There were 10 total metabolite clusters identified, each with

a clear molecular concentration pattern in which the metabolites in that cluster exhibited similar

rates of change together over the time phases.

Clusters of metabolites that exhibit similar rates of change for each time phase were

examined to gain an understanding of very broad metabolic pathways affected at each time

phase (File S3). In the early infection phase relative to the previous uninfected phase, there are

increased abundances of Clusters 4 and 5. These clusters contain metabolites involved in

glutathione biosynthesis (glutamate, cysteine, pyroglutamic acid, and $NADP^+$). Glutathione

intermediates are increasing while glutathione itself (Cluster 3) is decreasing. Decreased

abundance of glutathione paired with increased abundance of synthesis intermediates could be

indicative of how the insect is fighting off this pathogen; the insect immune response attempts to

strike an equilibrium between non-specific reactive molecules released through the

phenoloxidase melanization cascade and the antioxidant glutathione which protects against the

reactive molecules {Clark, 2010; Stahlschmidt, 2015}. Also increasing during this phase are

several amino acids, namely those involved in tryptophan metabolism (tryptophan itself and

indole). Metabolites that exhibited continuously decreasing abundances in the early phase

relative to the uninfected phase were mapped onto Clusters 1, 3, 8, 9, and 10. These clusters

contain many compounds, and the highest proportions are involved in purine and pyrimidine

biosynthesis and ascorbate metabolism (myo-Inositol, UDP-glucose, UDP-glucuronate, and

glucarate). From the microarray data, generally the regulator mutants caused transcripts in

ascorbate (*nilR, lrp*) and tryptophan (*lrhA, rpoS*) to decrease. These data from the early phase

represent the clusters of metabolites that could be targeted by the bacteria to induce insect

death and the metabolites used by the insect to fight the losing war.

From the early phase into the middle infection phase, few clusters exhibit an increase in

rates of change. These metabolites are in Clusters 8 and 10, which contain several purine

components (deoxyinosine, xanthosine, and inosine) as well as one of the only B vitamins

detected in this screen, riboflavin (vitamin B_2). Other detected B vitamins, like biotin (vitamin B_7),

pantothenate (vitamin B_5), and 4-pyridoxate (catabolic product of vitamin B_6), are decreasing in

abundance during the middle phase. Many other compounds decreased in abundance during

the middle infection phase, relative to the previous early infection phase. These include amino

acids (arginine, phenylalanine, tyrosine, tryptophan, cysteine, and methionine), which could

reflect that these are the amino acids being incorporated into protein creation for bacterial and

nematode biomass accumulation. Microarray analysis indicates widespread differential

regulation in these amino acid categories, particularly in the secondary form, *lrp*, and *lrhA*

mutants. Other decreasing compounds include pyrimidine intermediates (UMP, CMP, CDP, and

UDP) and ascorbate and sugar acid compounds. These data from the middle phase are

reflective of the compounds that are being syphoned from the cadaver and incorporated into the

nematode lifecycle.

As the insect cadavers entered the late infection phase, nucleic acids (guanine, thymine,

and uracil) and amino acids (arginine, cysteine and methionine, and aspartate) steeply

increased. This suggests these accumulating compounds are available for nematode DNA,

RNA, and protein incorporation, but other factors such as overcrowding in the cadaver or lack of

other necessary resources force the nematode to exit. Decreasing rate of change of metabolites

relative to the previous middle phase included compounds involved with leucine metabolism and

the TCA cycle. These could be more rate-limiting compounds, where their decreasing

abundance could signal to the expanding nematode population that it is time to exit.

A

B

Figure 5: Hierarchical clustering analysis of detected metabolites found ten metabolite cluster exhibit similar rates of change over the infection.
A) Dendrogram corresponding to spearman correlation values for each metabolite. B) Identified metabolite clusters with similar log(rate of change) over the lifecycle for the time phase, compared to the previous time phase. Metabolites with a red box exhibit an increased molecular rate of change and metabolites with a blue box exhibit a decreased molecular rate of change.

Interference of insect tricarboxylic acid (TCA) cycle is critical for infection success and subsequent propagation of nematodes

To identify significant metabolites that are important for infection progression, an ANOVA with post-hoc Tukey's HSD test was performed on metabolite abundances throughout the lifecycle. Two comparisons were examined: metabolite abundances from uninfected insects compared to individual time points, and individual time points compared to the next subsequent time point. As summarized in Figure 6, TCA cycle components significantly ($p<0.05$) fluctuate in relation to uninfected insects as well as between time phases, throughout the time course.

In the early phase of infection, while the insect is still alive and combatting bacteria and nematode invaders using innate immunity, several key TCA cycle intermediates are reduced in abundance relative to an uninfected insect (Fig. 6). This is shown through significantly decreased abundances of citrate in the Hour 12 and Hour 24 living insects compared to the uninfected insect and the Hour 24 dead insects. Although not significant, a similar trend is observed for two other TCA-related metabolites, malate and *sn*-glycerol-3-phosphate, which aids in NAD$^+$ regeneration through the glycerol phosphate shuttle, as well as NAD$^+$, NADH, fumarate {Shi, 2018}. This could mean these metabolites were diverted for the immune response, given the differences between the living and dead insects. As the infection progresses into a middle phase, citrate abundances flux, but generally are decreasing. Into the late phase on Day 10, malate, *sn*-glycerol phosphate, succinate, and citrate all drop, which could suggest carbon is being stored (rather than used) in the IJs before they exit the cadaver.

TCA cycle components were mostly in Cluster 6 (*sn*-glycerol-3-phosphate, NAD$^+$, NADH, citrate and isocitrate) and Cluster 2 (fumarate, alpha-Ketoglutarate). Student t-tests were utilized to determine additional significant components by comparing each broad time phase to the uninfected insects (File S3). Acetylphosphate was found to be approaching significantly high ($p<0.1$) at the early phase and was significantly high ($p<0.05$) at the middle and late phases.

NAD$^+$ was found to be approaching significantly low ($p<0.1$) at the early and middle phases and

was significantly low ($p<0.05$) for the late phase. *Xenorhabdus* spp. cannot synthesize NAD$^+$

and requires nicotinate for growth {Orchard, 2004; Martens, 2005}. Generally, the trend for all

TCA components in the lifecycle seem to be decreasing as the infection progresses with the

exceptions of acetylphosphate, FAD, and succinate.

Figure 6: Infection with *S. carpocapsae* IJs affects insect TCA cycle. Normalized molecular

concentration box plots throughout the lifecycle are shown for all detected metabolites involved

in the TCA cycle. Box plot colors represent which time phase the individual plots belong to for:

uninfected (black), early infection (red, going from earliest, 1hr, to latest, 24 hours dead, time

points), middle infection (yellow, days 2-8), and late infection (green, days 10-16). Lines in the

middle of the boxes indicate the mean molecular concentration. Bolded metabolites indicate

significant concentration shifts during the life cycle. NS indicates detection, but not significantly

affected (through the ANOVA) throughout infection. ND indicates not detectable. Highlighted

The aforementioned *X. nematophila* mutants experienced differences in transcripts

involved in either pyruvate metabolism, glyoxylate metabolism, or the TCA cycle (File S2). Most

of these genes were regulated in the Δ*lrhA* and Δ*rpoS* mutant backgrounds. *aceA* and *aceB* are

negatively regulated, while *aceE* and *aceF* were positively regulated between Δ*lrhA* and Δ*rpoS*

mutant strains and WT. These genes are involved in the glyoxylate shunt which is a pathway

utilized by many bacteria and nematodes to convert 2-carbon compounds into energy

resources, in the absence of bountiful sugars {Maloy, 1982}. Δ*rpoS* mutants upregulate several

succinate dehydrogenase genes which are necessary for oxidative phosphorylation. In *E. coli*

these genes are regulated in response to different environmental conditions like iron and heme

availability {Park, 1995}.

Acetyl-coA is an important node in metabolism, connecting glycolysis, the TCA cycle,

fatty acid, amino acid, and secondary metabolite pathways, and acetate dissimilation {Wolfe,

2015}. RpoS and LrhA positively influence the expression of AceF, PflB, Pta, and AckA. Pta-

AckA comprise the acetate dissimilation (excretion) pathway {Wolfe, 2015}. Coordinated

elevation of these enzymes is predicted to result in lower levels of acetyl-coA and higher levels

of acetylphosphate and acetate, which is excreted and available for use by the nematodes.

Acetylphosphate is a phosphoryl donor for some response regulators and can be a donor for

protein acetylation. Protein acetylation, a ubiquitous post-translational modification in

prokaryotes and eukaryotes, is involved in regulation of many different bacterium-host

interactions like chemotaxis, replication, and acid resistance, as well as regulating bacterial

DNA-binding and protein stability {Ren, 2017}. Acetylphosphate was detected in the

metabolome and generally increased over the infection, as well as being a VIP>1 metabolite for

components 1 and 2. Acetate freely diffuses across membranes and can be incorporated into

biomass of both bacteria and nematodes via the glyoxylate shunt {Alaimo, 2012}. *pflB* is

predicted to encode the pyruvate-formate lyase (PFL) enzyme involved in conversion of

pyruvate and CoA into formate and acetyl-coA and is greatly (>7 |fold change|) downregulated

in the Δ*rpoS* and Δ*lrhA* mutants relative to wild type. PflB converts glucose to formate, and up to

one-third of the carbon procured from glucose is converted through this enzyme in *E. coli*

{Doberenz, 2014}. PFL condenses acetyl-CoA and formate, allowing for the microbes to use

acetate and formate (fermentation products) as the sole carbon sources {Zelcbuch, 2016}.

RpoS and LrhA negatively regulate Acs, which is the acetate assimilation pathway {Wolfe,

2015}. In *E. coli*, Acs activity is inhibited by acetylation of a conserved lysine and its abundance

is negatively regulated by the small RNA SdhX {You, 2014; De Mets, 2019}. RpoS and LrhA

also both negatively regulate the TCA cycle enzymes AcnA and AceAB, and this inhibition is

predicted to result in accumulation of citrate. Citrate and isocitrate progressively decrease in

abundance over the infection cycle and these combined data might indicate that citrate

produced and accumulated by *X. nematophila* bacteria may be a provision for nematodes,

consumed during reproduction.

Analysis of amino acid abundances found proline fluctuates throughout the infection

The amino acid pyroglutamic acid, which is a precursor to glutamate, significantly rises

(as determined through the aforementioned ANOVA tests). in abundance during the late phase,

at Day 16 compared to the uninfected insect and between Day 12 and Day 16. Glutamate is a

precursor for proline metabolism {Watford, 2008}. Proline abundances fluctuated significantly

throughout the time course. Compared to uninfected insects, proline levels were significantly

lower at Day 10 and higher at Day 16, but otherwise were not significantly different. When

comparing each time point to the previous, there was a significantly higher level of proline in

dead insects at Hour 24 relative to living insects at the same time point. However, in dead

insects between Hour 24 and Day 2 the levels of proline dropped again. Thereafter there was a

cycle of increase and decrease in proline abundance (Day 4<Day 6, Day 8>Day 10, Day

12<Day 16), with an overall rise throughout the middle and late phases. Additional metabolites

in the arginine/proline/polyamine metabolism pathways that were significantly different among

samples based on student t-tests (File S3), included ornithine, which was significantly high at

the early and middle infection stages compared to uninfected insects, and hydroxyproline, which

was significantly high at early, middle, and late infection stages compared to uninfected insects.

Hierarchical clustering analysis reveals Cluster 1 (proline and hydroxyproline) and Cluster 5

(glutamate and pyroglutamic acid) contain these compounds. These cluster trends differ in the

early phase, where Cluster 1 decreases while Cluster 5 increases, possibly indicating that these

compounds are getting converted into each other.

Microarray analysis of Δ*lrhA*, Δ*lrp*, Δ*nilR*, Δ*rpoS*, and secondary form *X. nematophila*

mutants reveal similarities to each other in disrupted proline gene regulation compared to their

wild type/primary form counterparts. *putA*, predicted to encode 1-pyrroline-5-carboxylate

dehydrogenase, an enzyme involved in the conversion of proline to glutamate, was negatively

regulated (2<|fold change|) in the Δ*lrhA* and Δ*rpoS* strains, and positively regulated in the

secondary form strain. PutA is necessary for virulence in *Pseudomonas aeruginosa* and

protects the bacterium against oxidative stress {Zheng, 2018}. XNC1_2468 is downregulated in

the Δ*lrp*, Δ*rpoS*, and secondary form mutant backgrounds. This gene encodes a spermidine N1-

acetyltransferase as part of polyamine biosynthesis from ornithine and putrescine. XNC1_2274

and XNC1_3619 are FAD-dependent oxidoreductases involved downstream of proline

metabolism in putrescine utilization and are upregulated in the Δ*lrhA*, Δ*lrp*, Δ*nilR*, and Δ*rpoS*

mutant backgrounds. These genes are mostly involved in downstream proline metabolism,

specifically polyamine biosynthesis. Pathogenic gram-negative bacteria exploit polyamine-

related processes of their host for growth and proliferation, using these host molecules for toxin

activity, biofilm production, and limiting host immune responses {Shah, 2008; Di Martino, 2013}.

XNC1_2154 is predicted to encode an enzyme with L-aspartate:2-oxoglutarate

aminotransferase activity and is downregulated in the secondary form and Δ*lrp* mutant

backgrounds.

Amino acids leucine, isoleucine, and phenylalanine amino acids were significantly higher

in abundance at the late stage of infection. These are essential amino acids which *C. elegans*,

and presumably *S. carpocapsae*, must acquire from its diet (i.e. bacteria) {Zečić, 2019}. Several

X. nematophila genes, including *ilvC* and *ilvI*, involved in leucine/isoleucine biosynthesis were

differentially regulated in the Δ*lrhA* (negatively) and Δ*rpoS* (positively regulates *ilvC* and

negatively regulates *ilvI*) mutant backgrounds, relative to wild type. Δ*rpoS* mutants had higher

levels of *leuA* transcript, a gene predicted to encode a 2-isopropylmalate synthase, a leucine

precursor which is increases throughout the lifecycle. Δ*lrhA* increased transcript abundances of

fadA, fadB, *fadI,* and *fadJ* which participate in the conversion of leucine and isoleucine into fatty

acids and acetyl-CoA. In the Δ*lrp* and Δ*rpoS* mutant backgrounds, *mmsA* was upregulated,

another gene that encodes a methylmalonate-semialdehyde dehydrogenase that participates in

the breakdown of leucine and isoleucine.

DISCUSSION

A comprehensive framework to understand how metabolism shifts during infection

lifecycles of entomopathogenic nematodes was established. Physiologically, it was examined

that *X. nematophila* bacteria consume insect tissues, while *S. carpocapsae* nematodes

consume bacteria. The high $TP_{glu-phe}$ of 4.5 observed in the nematodes emerging from an insect

cadaver suggests the IJs were potentially cannibalizing previous generations of nematodes

and/or feeding upon bacteria that were, themselves, already feeding on previous generations of

nematodes and bacteria, since if the colonizing nematodes were feeding on bacterial and insect

biomass only, they would register at around 3.5 Thus, this high trophic level suggests either

endotokia matricida (or bagging) in which nematode eggs hatch within and consume the

mother. likely during the second generation of nematodes when nutrients are becoming

depleted, or that some other form of cannibalism is occurring within the cadaver {Baliadi, 2004}.

A $TP_{glu-phe}$ of 4.6 is similar to many apex predators, such as large marine carnivores or the rare

top predators observed in terrestrial ecosystems {Chikaraishi, 2011; Chikaraishi, 2014; Steffan,

2013; Ohkouchi, 2017}. This underscores the importance of including microbes in studies of

organismal trophic identity. In effect, the cadavers used in this study may represent microcosms

of the broader communities and ecosystems in which they are embedded. The insect cadavers

were, when alive, herbivores. To find multiple levels of carnivory within a single cadaver

suggests that a nematode-colonized arthropod mirrors the trophic richness of the broader food-

web. The interdigitation of microbial carnivores in a trophic hierarchy—here, nematodes and

bacteria—is likely a much more common feature of food-webs than previously thought {Steffan,

2017, Steffan and Dharampal, 2019}. The microbial trophic identities reported in this study may

necessitate a re-calibration of organismal niche concepts, but in so doing, will facilitate the

unification of the macro- and microbiome in food web ecology.

It should be emphasized that it has been exceedingly uncommon to find higher-order

consumers (TP > 4.0) in a community or ecosystem, given that apex predators feed upon other

predators that have, themselves, had to find and subdue 'lower' carnivores {Lindeman, 1942}. In

classical food web ecology, apex predators are generally considered to be large, fierce, and

rare vertebrates Elton, 1927, Colinvaux and Barnett, 1979}. However, perhaps the assumption

that apex predators exist only within the province of large/fierce/rare vertebrates needs to be

revisited. The high trophic positions exhibited by the nematodes in this study suggest that such

obligate higher-order consumers are more common than previously thought, with multitudes of

apex carnivores existing underfoot in many terrestrial ecosystems. Further, the nematodes can

be viewed as farming their symbionts: acting as shepherds that bring their bacterial flock to a

fresh insect pasture for harvesting of nutrients.

The trophic study established the foundation to understand metabolic shifts occurring in

the cadaver. The time course metabolomics study sought to better understand the process of

bioconversion in the cadaver, how is the insect biomass being converted to bacterial and

nematode biomass? Applying multivariate statistical tests to the infection metabolomics data set

revealed distinct time phase clustering. The variance among the time phases seems to increase

as infection progresses, as the healthy insects degrades into bacteria and nematode tissue.

Nematode IJ samples were also compared to these time phases. These samples form a distinct

cluster away from the other time phases (File S5). The IJ samples completely removed from the

lifecycle seem to have metabolic profiles most similar to the late time phase, which are insect

samples bursting with nematodes (Fig. 3). However, it is important to note that these are input

IJs that have spent weeks outside of an insect cadaver. Reassessing this analysis with output

IJs that are recently from the nematode is important to follow up on.

Metabolic analysis revealed TCA cycle components were among the most significant

results, indicating that their use by the entomopathogenic nematodes is paramount to infection

success and subsequent nematode propagation. Citrate metabolism is ubiquitous in many

intracellular pathogens and has been found to be involved in regulating virulence {Martino,

2018}. Citrate is necessary for virulence and growth of the Gram-negative pathogenic bacteria

Pseudomonas aeruginosa, where NADH levels were reduced when the bacterium was treated

with citrate and host-killing activity was abolished as a result {Perinbam, 2020}. This group

hypothesized that this could be due to decreased flux through the glyoxylate bypass, which has

been found to activate the T3SS in this system {Chung, 2013}. As mentioned, *X. nematophila*

does not encode a T3SS, but does have the evolutionarily related flagellar export apparatus.

Several *X. nematophila* glyoxylate bypass genes, were differentially transcribed between

avirulent genetic mutants and WT. Glyoxylate was not detected in our screen, and whether flux

through this pathway affects virulence should be investigated further. The significant flux of TCA

metabolites during the middle phase may be indicative of the role of the TCA cycle in *S.*

carpocapsae development, as the TCA cycle has been shown to be essential in early

embryogenesis in *C. elegans* {Rahman, 2014; Hada, 2019}. Citrate synthase (*cts-1*) and cyclin-

dependent kinase 1 (*cdk-1*) were inhibited in these studies and halted *C. elegans* development,

and both genes have orthologs in *S. carpocapsae*. Neutral lipids are formed from *sn*-glycerol-3-

phosphate and are the major energy reserve in the closely related *S. feltiae* nematodes {Wright,

1997}. Fats are stored as lipid droplets in *C. elegans* dauer larvae intestines and serve as a

starvation survival mechanism (Mak, 2012; O'Riordan, 1989). The previously mentioned

glyoxylate bypass forms carbohydrates from fatty acids and has been implicated in extending

the lifecycle of *C. elegans* {Edwards, 2013}, highlighting another role of this TCA vs. glyoxylate

switching that could be happening later in the life cycle. Any indication that cholesterol is being

synthesized from these intermediates can be attributed entirely to the insect's wheat germ diet,

as the *X. nematophila*, *S. carpocapsae*, or *G. mellonella* cannot synthesize sterols but require

them to grow {Dutky, 1967; Jing, 2020}.

Additionally, metabolites such as 2,3-dihydroxybenzoate that are not synthesized by *X.*

nematophila or *S. carpocapsae* were found to increase over the infection, past insect death. *X.*

nematophila does not encode the genes that convert 2,3-dihydroxybenzoate to the siderophore

enterobactin, which bind iron to create the ferric enterobactin (FeEnt) complex {Raymond,

2003}. However, *X. nematophila* encodes FepB, a periplasmic enterobactin binding protein, as

well as FepC, FepD, and FepG, which transport the FeEnt into the cell. This increased

abundance of this compound suggests other *G. mellonella* microbiome members that survive

infection synthesize a compound that is paramount in iron extraction and could contribute to the

overall fitness of this symbiosis.

Metabolic analysis also revealed the significance in proline throughout the lifecycle.

Insect hemolymph is rich in proline and is used as a main fuel source in some species of flying

insects because of its ability to oxidize carbohydrates {Teulier, 2016; Goldstrohm, 2003}. Proline

can be a signal molecule inducing secondary metabolite biosynthesis in *Xenorhabdus* species.

Several known virulence factors and antibiotics are regulated via exogenously supplied proline

to *Xenorhabdus* cultures {Crawford, 2010}. *Xenorhabdus* species may have evolved to use

proline in insect hemolymph as a preferred amino acid source capable of enhancing the

bacterium's virulence as well as protecting it from various stressors it meets in the insect body.

Proline catabolism has also been implicated in promoting stress responses and modulating

innate immunity in *C. elegans*, highlighting a possible mechanism by which *S. carpocapsae*

survives reactive oxygen species produced by the insect, bacterium, or themselves {Tang,

2016}. Enhanced understanding of proline changes over time in the EPNB lifecycle highlights

the multiple roles this amino acid is playing for both *Xenorhabdus* virulence and *Steinernema*

protection and reproduction. Additional amino acids were found to exhibit similar concentration

shifts over the lifecycle via the hierarchical clustering analysis. This machine learning technique

can be improved with higher granularity of sample time points, which would strengthen the

software developed for this study and allowing it to be used to study more complex and dynamic

chemical environments.

We have shown how the parasitic EPNB infection shapes the insect host metabolism.

The trophic hierarchy improved how we understand the parasites replicate in the host and

highlights the importance of including microbes in these studies. Through rigorous metabolic

pathway reconstruction and multivariate statistics, these results suggest each phase of the

symbiosis can be characterized by stage-specific chemical signatures. Future targeted

metabolomics experiments on EPNB symbioses should be developed to expand the specific

trends elucidated by this study. This work adds to a growing scientific foundation on how

symbioses, both mutualistic and parasitic, shape the chemical environments they inhabit.

MATERIALS AND METHODS

Conventional nematode and aposymbiotic nematode production

S. carpocapsae nematodes (All strain) were propagated through 5[th] instar larvae of insect *G.*

mellonella and using white trp and conventional IJs were collected by trapping in distilled water

at stored at room temperature for <1.5 months {McMullen, 2014}. To generate aposymbiotic IJs,

X. nematophila $\Delta SR1$ mutant were grown in Luria Broth (LB) media overnight at 30°C on cell

culture wheel and 600μL of overnight bacterial culture were spread onto each of 10mL of lipid

agar to grow into confluent lawn at 25°C for 48 hours. Conventional IJs were surface-sterilized,

seeded onto $\Delta SR1$ mutant lawn on lipid agar plates (5000 IJs per 10mL media), and incubated

at 25°C for 7 days in dark for nematode reproduction. Aposymbiotic IJs were collected by water-

trapping using distilled water and stored at room temperature in dark {McMullen, 2014}.

In vitro **controlled feeding experiment**

To collect bacteria sample feeding on terrestrial C3 plants and yeast-based media, *X.*

nematophila wildtype or $\Delta SR1$ mutant bacteria were grown in the dark yeast soy broth (0.5%

yeast extract, 3% tryptic soy broth, and 0.5% NaCl) modified from the bacterial growth media

from {Buecher and Popiel., 1989} at 30°C on cell culture wheel. Wildtype bacterial overnight

cultures (5mL per condition per biological replicate) were collected into microfuge tubes, spun

down at >5000RPM, and washed for three times using 1x PBS buffer by resuspending and

spinning down the bacterial pellets. Exactly 600μL of $\Delta SR1$ mutant were spread onto yeast-soy

lipid agar plates (0.5% yeast extract, 3% tryptic soy broth, 1.5% agar, 0.2% $MgCl_2$, 0.7% corn

syrup, 0.4% soy bean oil, supplemented with 40μg cholesterol at per liter of media) and

incubated for 48h at 25°C to grow into a confluent lawn. Bacterial lawns were washed off the

agar plate using 1x PBS, pelleted and washed as described above. Three individual tubes of

bacterial culture were used per strain as three independent biological replicates.

To grow nematodes using a controlled diet, approximately 5000 conventional IJs were surface-

sterilized and seeded onto bacterial lawn grown on yeast-soy lipid agar plate as described

above. Three individual yeast-soy lipid agar plates were used as three biological replicates for

each bacterial condition. Three days post seeding IJs, first generation reproductive stage of

nematodes (adult males and females) were collected by flooding the bacterial lawn with 1x PBS

buffer to resuspend the nematodes. The nematode suspension were collected in the glass cell

culture tube and washed for three times by resuspending in 1x PBS buffer. Seven days post

seeding the IJs, second generation IJ progenies were collected using distilled water traps and

washed in water for three times (settle by gravity and resuspension).

***In vivo* feeding experiment and sample collection**

To prepare insect controls, *G. mellonella* 5th instar larvae were injected with 10uL of either 1x

PBS buffer, yeast-soy broth media, or nothing. Three insect larvae were prepared per condition

as three biological replicates.

To collect nematodes directly fed on *Galleria* insect tissues, *S. carpocapsae* axenic eggs were

extracted from adult female nematodes grown on yeast-soy lipid agar plates. Approximately

6000 axenic eggs were seeded to each of the *Galleria*-tissue agar plate (20% (w/v) frozen *G.*

mellonella insects cleaned, blended and filtered; 0.5% (w/v) NaCl; and 1.5% (w/v) agar,

supplemented with 50mg/L Kanamycin). Mixed-stages of nematodes were collected by flooding

the *Galleria*-tissue agar with 1x PBS to resuspend the nematodes, then washed in 1x PBS and

water for 3 times to separate nematodes from insect tissue debris.

To establish controlled feeding experiments *in vivo* for bacteria and nematodes, *X. nematophila*

overnight cultures (in yeast-soy broth) were diluted in 1xPBS buffer, approximately 10^4 bacterial

cells were injected with or without aposymbiotic nematodes (100 IJs per insect). Insect cadaver

injected with bacteria only were directly lyophilized and used as insect-bacteria complex

controls (see methods below). Insects with bacteria and nematodes co-injection mixture were

used to collect IJ progenies by water-trapping, washing (3x in distilled water), and pelleting the

IJ samples. Three to five insects were used for each experimental condition as biological

replicates.

Nematode lyophilization and trophic position analysis

Nematodes from *G. mellonella* were collected by placing infected cadavers in modified White

traps in which nematodes migrate into distilled water. Trapped nematodes were transferred to

Falcon test tubes and allowed settle into a pellet at the bottom of the tube. Nematodes from

plate cultivations were harvested by rinsing with sterile distilled water, transferred to Falcon test

tubes, and allowed to settle. Samples of nematodes were stored in water at 10°C no longer than

a few days until they were lyophilized. For lyophilization, water was decanted off of the sample

until only the undisturbed pellet remained at the bottom of the test tube. The top of the test tube

was covered with a Kimwipe held in place with a rubber band before lyophilization for >48 h in a

Labconco Freezone lyophilizer. During this time, pressures fell below 0.2 millibar, and

temperatures reached -50°C. Once the samples had been thoroughly lyophilized, they were

removed from tubes using a laboratory spatula that was sterilized with ethanol and dried with

kimwipe after every use. Each individual sample was relocated into a sterile 1.5 ml microfuge

tube and stored at room temperature for 1-3 months until shipment to Hokkaido, Japan for

analysis.

Trophic position (TP$_{glu-phe}$) estimates were generated using the following equation:

742
$$TP = \frac{\delta^{15}N_{glu} - \delta^{15}N_{phe} + |\beta|}{\Delta_{glu-phe}} + \lambda$$

where $\delta^{15}N_{glu}$ represents the nitrogen isotopic ratio of glutamic acid, $\delta^{15}N_{phe}$ represents the

nitrogen isotopic ratio of phenylalanine, β corrects for the difference in ^{15}N values between

glutamic acid and phenylalanine within the primary producers of the food web (e.g. $\beta \sim 8.4$‰ for

C3 plants), $\Delta_{glu-phe}$ represents the net trophic discrimination between glutamic acid and

phenylalanine, and λ represents the basal trophic level (=1) of the food web {Chikaraishi, 2014}.

The trophic discrimination factor, $\Delta_{glu-phe}$ (referred to here as the $TDF_{glu-phe}$), represents the net

intertrophic ^{15}N-discrimination between glutamic acid and phenylalanine. Significant differences

between known and observed TP values were examined using univariate ANOVA and

nonparametric tests (paired Wilcoxon signed rank tests where data were heteroscedastic).

Distinguishing among TDF values was accomplished using paired t tests {Steffan, 2015}.

Metabolomics sample collection

As per normal infection protocols, 11 *G. mellonella* larvae (Grubco) were placed in the bottom of

each of six 6 x 1.5 cm petri plates lined with 2 pieces of #1 filter paper. The filter paper was then

inoculated with 1 ml of conventional *S. carpocapsae* IJ stage nematodes (carrying *X.*

nematophila bacteria in their intestinal receptacle) to achieve a final average concentration of 10

IJ/μl. At each specified time point (see below), one *G. mellonella* was taken from each of plates

1-5. All insect samples were flash frozen using a dry ice-ethanol bath, and subsequently stored

at -80ºC. The uninfected, Hour 1 post-infection and Hour 12 post-infection data points were

taken of live *G. mellonella*. Since the *G. mellonella* were starting to succumb to the infection at

Hour 24, one living and one dead insect was taken at this time point. At Day 7 post-infection, a

water trap was set up to enable IJ emergence. At Day 12, the last of the *G. mellonella* from

plates 1-5 was used, so insects representing the Day 16 time point were taken entirely from

plate 6. Input *S. carpocapsae* IJ and *X. nematophila* symbionts were also collected from the lab

stocks and sent for analysis, approximately 50 μl of settled IJ per sample, and a total of 4

samples were sent.

Preparation for mass spectrometry

For metabolite extraction, *G. mellonella* insects were equilibrated to -20˚C for ~1 h, 300 μl of

extraction solution (40:40:20 acetic acid, methanol, and water) was added, and insects were

ground using a pestle that fit snugly into the sample tube. All manipulations were performed in a

cold room and samples were processed in groups of 12. After grinding, to each tube an

additional 1000 μl of extraction solution was added and vortexed for 5-10 sec before being

placed at -20˚C for 20 min. Tubes were centrifuged at 16,200 x g for 5 min, and the supernatant

was decanted to a clean tube. To the original insect sample an additional 200 μl of extraction

solution was added, mixed with a pipette tip, vortexed for 5-10 sec, and incubated at -20˚C for

20 min. After pelleting the supernatant was combined with the first supernatant sample.

Samples were dried (Savant), resuspended, and randomized samples were analyzed

consecutively by mass spectrometry using an established 25 minute method (Lu, 2010).

Metabolomics analysis

An established untargeted metabolomics method utilizing ultra-high performance liquid

chromatography coupled to high resolution mass spectrometry (UHPLC-HRMS) (Thermo

Scientific, San Jose, CA, USA) was used to analyze water-soluble metabolites {Lu, 2010}. A

Synergi 2.6 μm Hydro RP column 100 Å, 100 mm x 2.1 mm (Phenomenex, Torrance, CA) and

an UltiMate 3000 pump (Thermo Fisher) were used to carry out the chromatographic

separations prior to full scan mass analysis by an Exactive Plus Orbitrap MS (Thermo Fisher).

HPLC grade solvents (Fisher Scientific, Hampton, NH, USA) were used. Chromatographic peak

areas for each detected metabolite were integrated using an open-source software package,

Metabolomic Analysis and Visualization Engine (MAVEN) {Clasquin, 2012; Melamud, 2010}.

Area under the curve (AUC) was used for further analyses.

Statistical analysis

PLS-DA plots were generated in MetaboAnalyst 4.0 on August 3rd, 2020. VIP scores were

calculated for each component. When more than components are used to calculate the feature

importance, the average of the VIP scores are used. The other importance measure is based on

the weighted sum of PLS-regression. The weights are a function of the reduction of the sums of

squares across the number of PLS components {Chong, 2020}. Samples were normalized

before processing through MetaboAnalyst based on insect weight. Data was log transformed

and pareto scaling was applied. Two-way ANOVA with multiple comparisons and Tukey post-

hoc tests were completed by taking individual time point metabolite abundances and comparing

their means to the uninfected insect model and each other. Student t-tests were performed by

comparing uninfected samples to each time phase (early, middle, and late infection). Relevant

metabolic pathways were identified in MetaboAnalyst's "Pathway Analysis" module using

Drosophila melanogaster, *Caenorhabditis elegans*, and *Escherichia coli* as KEGG pathway

libraries {Kanehisa, 2017}. Technical details for the hierarchical clustering analyses are

available at: http://doi.org/10.5281/zenodo.3962081

Bacterial strains, plasmids, and culture conditions

File S2 lists strains used for this study with references where they were originally published.

Unless specifically mentioned, *E. coli* were grown in LB broth or on LB plates at 37°C; *X.*

nematophila were grown in LB broth or on LB plates supplemented with 0.1% pyruvate at 30°C

and kept in dark. Where appropriate, the following antibiotic concentrations were used:

ampicillin, 150 μg/ml for *E. coli* and 50 μg/ml for *X. nematophila*; chloramphenicol, 30 μg/ml;

erythromycin, 200 μg/ml; kanamycin, 50 μg/ml and streptomycin, 25 μg/ml. *E. coli* donor strain

S17 (λpir) or Δ*asd* strain BW29427 was used to conjugate plasmids into *X. nematophila*.

Microarray experiment and data analysis

Bacteria cultures were grown overnight in 3 ml of LB supplemented with 0.1% pyruvate and

appropriate antibiotics in culture tubes at 30°C on roller, subcultured 1:100 into 30 ml of LB

supplemented with 0.1% pyruvate and 50 g/ml ampicillin in 125 ml glass flasks and grown for 12

hours to early stationary phase (OD 2-2.1) at 30°C at 150 rpm on shaker. 1 ml of each culture

was used to extract total RNA using Qiagen RNeasy Mini Kit, and on-column DNA digestion

was performed using Qiagen RNase-Free DNase Set according to manufacturer's protocol

(Qiagen, Valencia, CA). The RNA purity was tested by measuring 260 nm/280 nm and 260

nm/230 nm ratios in TE buffer and the values should be over 1.8. RNA integrity was verified by

running 2 g of RNA samples on 1% denaturing agarose gel. The samples were then submitted

to Roche NimbleGen for processing and microarray analysis. Gene signals for *lrhA*, *lrp*, and

secondary form *X. nematophila* were compared to HGB800 using a 2-fold change average

signal strength cutoff. The *rpoS* mutant was compared to HGB007 using the same significance

cutoff. Genes were annotated via the Magnifying Genomes (MaGe) microbial genome

annotation system {Vallenet, 2006}, the STRING database {Szklarczyk, 2019}, as well as

through BlastKOALA {Kanehisa, 2016}.

ACKNOWLEDGEMENTS

This work was supported by a grant to H.G.-B. from the National Science Foundation (IOS-

1353674) and by funds from the University of Tennessee-Knoxville. M.C. was supported by a

UW-Madison Louis and Elsa Thomsen Wisconsin Distinguished Graduate Fellowship and a

Department of Bacteriology Michael Foster Predoctoral Fellowship. We thank Terra Mauer for

her help in insect rearing and providing the *G. mellonella* image used for Figure 1. We thank

Xiaojun Lu for preprocessing the microarray data. We thank Jordan Rogerson for processing

the metabolomics samples prior analysis.

REFERENCES

1. **Abdelhamed H., Ramachandran R., Narayanan L., Ozdemir O., Cooper A., Olivier A. K., Karsi A., Lawrence M. L.** (2020). Contributions of a LysR Transcriptional Regulator to *Listeria monocytogenes* Virulence and Identification of Its Regulons. *J Bacteriol*, 202(10). doi:10.1128/JB.00087-20

2. **Alaimo J. T., Davis S. J., Song S. S., Burnette C. R., Grotewiel M., Shelton K. L., Pierce-Shimomura J. T., Davies A. G., Bettinger J. C.** (2012). Ethanol metabolism and osmolarity modify behavioral responses to ethanol in *C. elegans. Alcohol Clin Exp Res*, 36(11), 1840-1850. doi:10.1111/j.1530-0277.2012.01799.x

3. **Baliadi Y, Yoshiga T, Kondo E.** (2004) Infectivity and post-infection development of infective juveniles originating via endotokia matricida in entomopathogenic nematodes. *Applied Entomology and Zoology*;39:61–69.

4. **Budnick J. A., Sheehan L. M., Ginder M. J., Failor K. C., Perkowski J. M., Pinto J. F., Kohl K. A., Kang L., Michalak P., Luo L., Heindl J. E., Caswell C. C.** (2020). A central role for the transcriptional regulator VtlR in small RNA-mediated gene regulation in *Agrobacterium tumefaciens. Sci Rep*, 10(1), 14968. doi:10.1038/s41598-020-72117-0

5. **Buecher E. J., Popiel I.** (1989). Liquid Culture of the Entomogenous Nematode *Steinernema feltiae* with Its Bacterial Symbiont. *J Nematol*, 21(4), 500-504. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/19287644

6. **Blumer C., Kleefeld A., Lehnen D., Heintz M., Dobrindt U., Nagy G., Michaelis K., Emödy L., Polen T., Rachel R., Wendisch V. F., Unden G**. (2005). Regulation of type 1 fimbriae synthesis and biofilm formation by the transcriptional regulator LrhA of *Escherichia coli. Microbiology,* 151(Pt 10), 3287-3298. doi:10.1099/mic.0.28098-0

7. **Cambon M. C., Lafont P., Frayssinet M., Lanois A., Ogier J. C., Pages S., Parthuisot N., Ferdy J. B., Gaudriault S**. (2020). Bacterial community profile after the

lethal infection of *Steinernema-Xenorhabdus* pairs into soil-reared *Tenebrio molitor*

larvae. *FEMS Microbiol Ecol,* 96(2). doi:10.1093/femsec/fiaa009

8. **Camilli A., Bassler B. L.** (2006). Bacterial small-molecule signaling pathways. *Science,*

311(5764), 1113-1116. doi:10.1126/science.1121357

9. **Cao M., Goodrich-Blair H.** (2020). *Xenorhabdus nematophila* bacteria shift from

mutualistic to virulent Lrp-dependent phenotypes within the receptacles of *Steinernema*

carpocapsae insect-infective stage nematodes. *Environ Microbiol*, 22(12), 5433-5449.

doi:10.1111/1462-2920.15286

10. **Chaston J., Douglas A. E.** (2012). Making the most of "omics" for symbiosis research.

Biol Bull, 223(1), 21-29. doi:10.1086/BBLv223n1p21

11. **Chaston J. M., Suen G., Tucker S. L., Andersen A. W., Bhasin A., Bode E., Bode H.**

B., Brachmann A. O., Cowles C. E., Cowles K. N., Darby C., de Léon L., Drace K.,

Du Z., Givaudan A., Herbert Tran E. E., Jewell K. A., Knack J. J., Krasomil-

Osterfeld K. C., Kukor R., Lanois A., Latreille P., Leimgruber N. K., Lipke C. M., Liu

R., Lu X., Martens E. C., Marri P. R., Médigue C., Menard M. L., Miller N. M.,

Morales-Soto N., Norton S., Ogier J. C., Orchard S. S., Park D., Park Y., Qurollo B.

A., Sugar D. R., Richards G. R., Rouy Z., Slominski B., Slominski K., Snyder H.,

Tjaden B. C., van der Hoeven R., Welch R. D., Wheeler C., Xiang B., Barbazuk B.,

Gaudriault S., Goodner B., Slater S. C, Forst S., Goldman B. S., Goodrich-Blair H.

(2011). The entomopathogenic bacterial endosymbionts *Xenorhabdus* and

Photorhabdus: convergent lifestyles from divergent genomes. *PLoS One.* 6(11):e27909.

doi: 10.1371/journal.pone.0027909. PMID: 22125637; PMCID: PMC3220699.

12. **Chikaraishi Y., Kashiyama Y., Ogawa N. O., Kitazato H., Ohkouchi N.** (2007).

Biosynthetic and metabolic controls of nitrogen isotopic composition of amino acids in

marine macroalgae and gastropods: implications for aquatic food web studies. Mar.

Ecol. Prog. Ser. 342: 85– 90.

13. **Chikaraishi Y., Ogawa N. O., Kashiyama Y., Takano Y., Suga H., Tomitani A.,**

Ohkouchi N. (2009). Determination of aquatic food-web structure based on compound-

specific nitrogen isotopic composition of amino acids. *Limnol. Oceanogr. Methods* 7:

740– 750.

14. **Chikaraishi Y., Ogawa N.O., Doi H., Ohkouchi N.** (2011) ^{15}N/^{14}N ratios of amino acids

as a tool for studying terrestrial food webs: a case study of terrestrial insects (bees,

wasps, and hornets). *Ecol. Res.*;26:835–844.

15. **Chikaraishi Y., Steffan S. A., Ogawa N. O., Ishikawa N. F., Sasaki Y., Tsuchiya M.,**

Ohkouchi N. (2014). High-resolution food webs based on nitrogen isotopic composition

of amino acids. *Ecol Evol,* 4(12), 2423-2449. doi:10.1002/ece3.1103

16. **Chong J., Xia J.** (2020). Using MetaboAnalyst 4.0 for Metabolomics Data Analysis,

Interpretation, and Integration with Other Omics Data. *Methods Mol Biol, 2104*, 337-360.

doi:10.1007/978-1-0716-0239-3_17

17. **Chung J. C., Rzhepishevska O., Ramstedt M., Welch M. (**2013). Type III secretion

system expression in oxygen-limited *Pseudomonas aeruginosa* cultures is stimulated by

isocitrate lyase activity. *Open Biol*, 3(1), 120131. doi:10.1098/rsob.120131

18. **Ciche T. A., Blackburn M., Carney J. R., Ensign J. C.** (2003). Photobactin: a catechol

siderophore produced by *Photorhabdus luminescens*, an entomopathogen mutually

associated with *Heterorhabditis bacteriophora* NC1 nematodes. *Appl Environ Microbiol*,

69(8), 4706-4713. doi:10.1128/aem.69.8.4706-4713.2003

19. **Clark K. D., Lu Z., Strand M. R.** (2010). Regulation of melanization by glutathione in the

moth *Pseudoplusia includens*. *Insect Biochem Mol Biol*, 40(6), 460-467.

doi:10.1016/j.ibmb.2010.04.005

20. **Clasquin M. F., Melamud E., Rabinowitz J. D.** (2012). LC-MS data processing with

MAVEN: a metabolomic analysis and visualization engine. *Curr Protoc Bioinformatics*,

Chapter 14, Unit14 11. doi:10.1002/0471250953.bi1411s37

21. **Cobb N. A.** (1914). Nematodes and their relationships. *Yearbook United States*

Department Agriculture, 457-490.

22. **Colinvaux P. A., Barnett B. D.** (1979). Lindeman and the Ecological Efficiency of

Wolves. *Am. Nat.*; 114: 707-718

23. **Cowles C. E., Goodrich-Blair H.** (2006). *nilR* is necessary for co-ordinate repression of

Xenorhabdus nematophila mutualism genes. *Molecular Microbiology*; 62:760-771.

24. **Cowles C. E., Goodrich-Blair H.** (2008). The *Xenorhabdus nematophila nilABC* genes

confer the ability of *Xenorhabdus* spp. to colonize *Steinernema carpocapsae*

nematodes. J Bacteriol, 190(12), 4121-4128. doi:10.1128/JB.00123-08

25. **Cowles K. N., Cowles C. E., Richards G. R., Martens E. C., Goodrich-Blair H. (**2007).

The global regulator Lrp contributes to mutualism, pathogenesis and phenotypic

variation in the bacterium *Xenorhabdus nematophila*. *Cell Microbiol,* 9(5), 1311-1323.

doi:10.1111/j.1462-5822.2006.00873.x

26. **Crawford J. M., Kontnik R., Clardy J. (**2010). Regulating alternative lifestyles in

entomopathogenic bacteria. *Curr Biol,* 20(1), 69-74. doi:10.1016/j.cub.2009.10.059

27. **da Silva W. J., Pilz-Junior H. L., Heermann R., da Silva O. S.** (2020). The great

potential of entomopathogenic bacteria *Xenorhabdus* and *Photorhabdus* for mosquito

control: a review. *Parasit Vectors*, 13(1), 376. doi:10.1186/s13071-020-04236-6

28. **De Mets F., Van Melderen L., Gottesman S.** (2019). Regulation of acetate metabolism

and coordination with the TCA cycle via a processed small RNA. *Proc Natl Acad Sci U S*

A, 116(3), 1043-1052. doi:10.1073/pnas.1815288116

29. **Dearth S. P., Castro H. F., Venice F., Tague E. D., Novero M., Bonfante P.,**

Campagna S. R. (2018). Metabolome changes are induced in the arbuscular

mycorrhizal fungus *Gigaspora margarita* by germination and by its bacterial

endosymbiont. *Mycorrhiza*, 28(5-6), 421-433. doi:10.1007/s00572-018-0838-8

30. **Di Martino M. L., Campilongo R., Casalino M., Micheli G., Colonna B., Prosseda G.**

(2013). Polyamines: emerging players in bacteria-host interactions. *Int J Med Microbiol,*

303(8), 484-491. doi:10.1016/j.ijmm.2013.06.008

31. **Doberenz C., Zorn M., Falke D., Nannemann D., Hunger D., Beyer L., Ihling C. H.,**

Meiler J., Sinz A., Sawers R. G. (2014). Pyruvate formate-lyase interacts directly with

the formate channel FocA to regulate formate translocation. *J Mol Biol,* 426(15), 2827-

2839. doi:10.1016/j.jmb.2014.05.023

32. **Dong Y., Brewer G. J.** (2019). Global Metabolic Shifts in Age and Alzheimer's Disease

Mouse Brains Pivot at NAD+/NADH Redox Sites. *J Alzheimers Dis*, 71(1), 119-140.

doi:10.3233/JAD-190408

33. **Dutky S. R., Robbins W. F., Thompson J. V.** (1967). The demonstration of sterols as

requirements for the growth, development and reproduction of the DD- 136 nematode.

Nematologica;13:140.

34. **Edwards C. B., Copes N., Brito A. G., Canfield J., Bradshaw P. C.** (2013). Malate and

fumarate extend lifespan in *Caenorhabditis elegans. PLoS One,* 8(3), e58345.

doi:10.1371/journal.pone.0058345

35. **Elton C. S.** (1927). Animal Ecology. Sidgwick & Jackson, London.

36. **Gibson K. E., Silhavy T. J.** (1999). The LysR homolog LrhA promotes RpoS

degradation by modulating activity of the response regulator *sprE. J Bacteriol,* 181(2),

563-571. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/9882671

37. **Givaudan A., Baghdiguian S., Lanois A., Boemare N.** (1995). Swarming and

Swimming Changes Concomitant with Phase Variation in *Xenorhabdus nematophilus.*

Appl Environ Microbiol, 61(4), 1408-1413. doi:10.1128/AEM.61.4.1408-1413.1995

38. **Givaudan A., Lanois A.** (2000). *flhDC*, the flagellar master operon of *Xenorhabdus*

nematophilus: requirement for motility, lipolysis, extracellular hemolysis, and full

virulence in insects. *J Bacteriol,* 182(1), 107-115. doi:10.1128/jb.182.1.107-115.2000

39. **Goldstrohm D. A., Pennington J. E., Wells M. A.** (2003). The role of hemolymph

proline as a nitrogen sink during blood meal digestion by the mosquito *Aedes aegypti*. *J*

Insect Physiol, 49(2), 115-121. doi:10.1016/s0022-1910(02)00267-6

40. **Grabau Z. J., Mauldin M. D., Habteweld A., Carter E. T.** (2020). Nematicide efficacy at

managing *Meloidogyne arenaria* and non-target effects on free-living nematodes in

peanut production. *J Nematol, 52*, 1-10. doi:10.21307/jofnem-2020-028

41. **Hada K., Hirota K., Inanobe A., Kako K., Miyata M., Araoi S., Matsumoto M., Ohta**

R., Arisawa M., Daitoku H., Hanada T., Fukamizu A. (2019). Tricarboxylic acid cycle

activity suppresses acetylation of mitochondrial proteins during early embryonic

development in *Caenorhabditis elegans. J Biol Chem,* 294(9), 3091-3099.

doi:10.1074/jbc.RA118.004726

42. **Harris S. J., Shih Y. L., Bentley S. D., Salmond G. P.** (1998). The *hexA* gene of

Erwinia carotovora encodes a LysR homologue and regulates motility and the

expression of multiple virulence determinants. *Mol Microbiol, 28*(4), 705-717.

doi:10.1046/j.1365-2958.1998.00825.x

43. **Henikoff S., Haughn G. W., Calvo J. M., Wallace J. C.** (1988). A large family of

bacterial activator proteins. *Proc Natl Acad Sci U S A*, 85(18), 6602-6606.

doi:10.1073/pnas.85.18.6602

44. **Herbert E. E., Goodrich-Blair H.** (2007). Friend and foe: the two faces of *Xenorhabdus*

nematophila. Nat Rev Microbiol, 5(8), 634-646. doi:10.1038/nrmicro1706

45. **Heroven A. K., Dersch P.** (2006). RovM, a novel LysR-type regulator of the virulence

activator gene *rovA*, controls cell invasion, virulence and motility of *Yersinia*

pseudotuberculosis. Mol Microbiol, 62(5), 1469-1483. doi:10.1111/j.1365-

2958.2006.05458.x

46. **Hussa E. A., Casanova-Torres A. M., Goodrich-Blair H. (**2015). The global

transcription factor Lrp controls virulence modulation in *Xenorhabdus nematophila*. *J*

Bacteriol; 197:3015-3025.

47. **Joyce S. A., Clarke D. J.** (2003). A *hexA* homologue from *Photorhabdus* regulates

pathogenicity, symbiosis and phenotypic variation. *Mol Microbiol,* 47(5), 1445-1457.

doi:10.1046/j.1365-2958.2003.03389.x

48. **Kanehisa M., Sato Y., Morishima K.** (2016) BlastKOALA and GhostKOALA: KEGG

tools for functional characterization of genome and metagenome sequences. *J. Mol.*

Biol. 428, 726-731.

49. **Kanehisa M., Furumichi M., Tanabe M., Sato Y., Morishima K.** (2017). KEGG: new

perspectives on genomes, pathways, diseases and drugs. *Nucleic Acids Res*, 45(D1),

D353-D361. doi:10.1093/nar/gkw1092

50. **Katsyuba E., Mottis A., Zietak M., De Franco F., van der Velpen V., Gariani K., Ryu**

D., Cialabrini L., Matilainen O., Liscio P., Giacchè N., Stokar-Regenscheit N.,

Legouis D., de Seigneux S., Ivanisevic J., Raffaelli N., Schoonjans K., Pellicciari R.,

Auwerx J. (2018). De novo NAD+ synthesis enhances mitochondrial function and

improves health. *Nature*. 563(7731):354-359. doi: 10.1038/s41586-018-0645-6. PMID:

30356218; PMCID: PMC6448761.

51. **Killiny N.** (2018). Generous hosts: Why the larvae of greater wax moth, *Galleria*

mellonella is a perfect infectious host model? *Virulence*, 9(1), 860-865.

doi:10.1080/21505594.2018.1454172

52. **Lacey L. A., Georgis, R.** (2012). Entomopathogenic nematodes for control of insect

pests above and below ground with comments on commercial production. *J Nematol,*

44(2), 218-225. Retrieved from https://www.ncbi.nlm.nih.gov/pubmed/23482993

53. **Latham J. A., Chen D., Allen K. N., Dunaway-Mariano D.** (2014). Divergence of

substrate specificity and function in the *Escherichia coli* hotdog-fold thioesterase

paralogs YdiI and YbdB. *Biochemistry*, 53(29), 4775-4787. doi:10.1021/bi500333m

54. **Lehnen D., Blumer C., Polen T., Wackwitz B., Wendisch V. F., Unden G.** (2002).

LrhA as a new transcriptional key regulator of flagella, motility and chemotaxis genes in

Escherichia coli. Mol Microbiol, 45(2), 521-532. doi:10.1046/j.1365-2958.2002.03032.x

55. **Lee Y., Jeong H., Park K. H., Kim K. W.** (2020). Effects of NAD(+) in *Caenorhabditis*

elegans Models of Neuronal Damage. Biomolecules, 10(7). doi:10.3390/biom10070993

56. **Lindeman R.L.** (1942). The trophic-dynamic aspect of ecology. *Ecology*, 23, pp. 399-

417

57. **Javal M., Terblanche J. S., Conlong D. E., Malan, A. P.** (2019). First Screening of

Entomopathogenic Nematodes and Fungus as Biocontrol Agents against an Emerging

Pest of Sugarcane, *Cacosceles newmannii* (Coleoptera: Cerambycidae). *Insects*, 10(4).

doi:10.3390/insects10040117

58. **Jing X., Behmer S. T.** (2020). Insect Sterol Nutrition: Physiological Mechanisms,

Ecology, and Applications*. Annu Rev Entomol*, 65, 251-271. doi:10.1146/annurev-ento-

011019-025017

59. **Lu W., Clasquin M. F., Melamud E., Amador-Noguez D., Caudy A. A., Rabinowitz J.**

D. (2010). Metabolomic analysis via reversed-phase ion-pairing liquid chromatography

coupled to a stand alone orbitrap mass spectrometer. *Anal Chem*, 82(8), 3212-3221.

doi:10.1021/ac902837x

60. **Lu X., Goodrich-Blair H., Tjaden B.** (2011). Assessing computational tools for the

discovery of small RNA genes in bacteria. *RNA*; 17:1635-1647.

61. **Lu X.** (2012). Examining Pathogenic and Mutualistic Regulatory Networks in the

Bacterium *Xenorhabdus nematophila.* Ph.D. thesis, UW-Madison

62. **Maddocks S. E., Oyston P. C. F.** (2008). Structure and function of the LysR-type

transcriptional regulator (LTTR) family proteins. *Microbiology*, 154(Pt 12), 3609-3623.

doi:10.1099/mic.0.2008/022772-0

63. **Mak H. Y.** (2012) Lipid droplets as fat storage organelles in *Caenorhabditis elegans*:

thematic review series: lipid droplet synthesis and metabolism: from yeast to man. *J*

Lipid Res.;53(1):28–33.

64. **Maloy S. R., Nunn W. D.** (1982). Genetic regulation of the glyoxylate shunt in

Escherichia coli K-12. *J Bacteriol*, 149(1), 173-180. Retrieved from

https://www.ncbi.nlm.nih.gov/pubmed/7033207

65. **Margulis L.** (1981). *Symbiosis in cell evolution: life and its environment on the early*

Earth. San Francisco: W. H. Freeman.

66. **Martens E. C., Vivas E. I., Heungens K., Cowles C. E., Goodrich-Blair H**. (2004)

Investigating mutualism between entomopathogenic bacteria and nematodes. *Nematol.*

Monographs Persp. 2, 447–462

67. **Martens E. C., Russell F. M., Goodrich-Blair H.** (2005). Analysis of *Xenorhabdus*

nematophila metabolic mutants yields insight into stages of *Steinernema carpocapsae*

nematode intestinal colonization. *Mol Microbiol, 58*(1), 28-45. doi:10.1111/j.1365-

2958.2005.04742.x

68. **Martino G. P., Perez C. E., Magni C., Blancato V. S.** (2018). Implications of the

expression of *Enterococcus faecalis* citrate fermentation genes during infection. *PLoS*

One, 13(10), e0205787. doi:10.1371/journal.pone.0205787

69. **May J. M., Qu Z. C., Meredith M. E.** (2012). Mechanisms of ascorbic acid stimulation of

norepinephrine synthesis in neuronal cells. *Biochem Biophys Res Commun*, 426(1),

148-152. doi:10.1016/j.bbrc.2012.08.054

70. **McMullen J. G. 2nd, Stock S. P**. (2014). *In vivo* and *in vitro* rearing of entomopathogenic

nematodes (*Steinernematidae* and *Heterorhabditidae*). *J Vis Exp,* (91), 52096.

doi:10.3791/52096

71. **Melamud E., Vastag L., Rabinowitz J. D.** (2010). Metabolomic analysis and

visualization engine for LC-MS data. *Anal Chem*, 82(23), 9818-9826.

doi:10.1021/ac1021166

72. **O'Riordan V. B., Burnell A. M.** (1989) Intermediary metabolism in the dauer larva of the

nematode *Caenorhabditis elegans*— 1. Glycolysis, gluconeogenesis, oxidative

phosphorylation and the tricarboxylic acid cycle. *Comp Biochem Physiol B.*;92(2):233–8.

73. **Ohkouchi N., Chikaraishi Y., Close H. G., Fry B., Larsen T., Madigan D. J.,**

McCarthy M. D., McMahon K. W., Nagata T., Naito Y. I., Ogawa N. O., Popp B. N.,

Steffan S., Takano Y., Tayasu I., Wyatt A. S. J., Yamaguchi Y. T., Yokoyama Y.

(2017). Advances in the application of amino acid nitrogen isotopic analysis in ecological

and biogeochemical studies. *Organic Geochemistry*, Volume 113, 150-174.

https://doi.org/10.1016/j.orggeochem.2017.07.009.

74. **Orchard S. S., Goodrich-Blair H.** (2005). Pyrimidine nucleoside salvage confers an

advantage to *Xenorhabdus nematophila* in its host interactions. *Appl Environ Microbiol,*

71(10), 6254-6259. doi:10.1128/AEM.71.10.6254-6259.2005

75. **Park S. J., Tseng C. P., Gunsalus R. P.** (1995). Regulation of succinate

dehydrogenase (*sdhCDAB*) operon expression in *Escherichia coli* in response to carbon

supply and anaerobiosis: role of ArcA and Fnr. *Mol Microbiol,* 15(3), 473-482.

doi:10.1111/j.1365-2958.1995.tb02261.x

76. **Perinbam K., Chacko J. V., Kannan A., Digman M. A., Siryaporn A.** (2020). A Shift in

Central Metabolism Accompanies Virulence Activation in *Pseudomonas aeruginosa*.

mBio, 11(2). doi:10.1128/mBio.02730-18

77. **Peterson C. N., Carabetta V. J., Chowdhury T., Silhavy T. J.** (2006). LrhA regulates

rpoS translation in response to the Rcs phosphorelay system in *Escherichia coli*. *J*

Bacteriol, 188(9), 3175-3181. doi:10.1128/JB.188.9.3175-3181.2006

78. **Phelan V. V., Moree W. J., Aguilar J., Cornett D. S., Koumoutsi A., Noble S. M.,**

Pogliano K., Guerrero C. A., Dorrestein P. C. (2014). Impact of a transposon insertion

in phzF2 on the specialized metabolite production and interkingdom interactions of

Pseudomonas aeruginosa. *J Bacteriol*, 196(9), 1683-1693. doi:10.1128/JB.01258-13

79. **Rahman M. M., Rosu S., Joseph-Strauss D., Cohen-Fix O.** (2014). Down-regulation of

tricarboxylic acid (TCA) cycle genes blocks progression through the first mitotic division

in *Caenorhabditis elegans* embryos. *Proc Natl Acad Sci U S A,* 111(7), 2602-2607.

doi:10.1073/pnas.1311635111

80. **Ramarao N., Nielsen-Leroux C., Lereclus D.** (2012). The insect *Galleria mellonella* as

a powerful infection model to investigate bacterial pathogenesis. *J Vis Exp*(70), e4392.

doi:10.3791/4392

81. **Raymond K. N., Dertz E. A., Kim S. S.** (2003). Enterobactin: an archetype for microbial

iron transport. *Proc Natl Acad Sci U S A*, 100(7), 3584-3588.

doi:10.1073/pnas.0630018100

82. **Ren J., Sang Y., Lu J., Yao Y. F.** (2017). Protein Acetylation and Its Role in Bacterial

Virulence. *Trends Microbiol*, 25(9), 768-779. doi:10.1016/j.tim.2017.04.001

83. **Richards G. R., Goodrich-Blair H.** (2009). Masters of conquest and pillage:

Xenorhabdus nematophila global regulators control transitions from virulence to nutrient

acquisition. *Cell Microbiol, 11*(7), 1025-1033. doi:10.1111/j.1462-5822.2009.01322.x

84. **Richards G. R., Herbert E. E., Park Y., Goodrich-Blair, H. (**2008). *Xenorhabdus*

nematophila lrhA is necessary for motility, lipase activity, toxin expression, and virulence

in *Manduca sexta* insects. *J Bacteriol, 190*(14), 4870-4879. doi:10.1128/JB.00358-08

85. **Rougon-Cardoso A., Flores-Ponce M., Ramos-Aboites H. E., Martínez-Guerrero C.**

E., Hao Y. J., Cunha L., Rodríguez-Martínez J. A., Ovando-Vázquez C., Bermúdez-

Barrientos J. R., Abreu-Goodger C., Chavarría-Hernández N., Simões N., Montiel R.

(2016). The genome, transcriptome, and proteome of the nematode *Steinernema*

carpocapsae: evolutionary signatures of a pathogenic lifestyle. *Sci Rep*, 6, 37536.

doi:10.1038/srep37536

86. **Shah P., Swiatlo E.** (2008). A multifaceted role for polyamines in bacterial pathogens.

Mol Microbiol, 68(1), 4-16. doi:10.1111/j.1365-2958.2008.06126.x

87. **Shi Y., Wang H., Yan Y., Cao H., Liu X., Lin F., Lu J**. (2018). Glycerol-3-Phosphate

Shuttle Is Involved in Development and Virulence in the Rice Blast Fungus *Pyricularia*

oryzae. Front Plant Sci, 9, 687. doi:10.3389/fpls.2018.00687

88. **Stahlschmidt Z., Acker M., Kovalko I., Adamo S.** (2015). The double-edged sword of

immune defence and damage control: Do food availability and immune challenge alter

the balance? *Functional Ecology.* 29. 10.1111/1365-2435.12454.

89. **Steffan S. A., Chikaraishi Y., Horton D. R., Ohkouchi N., Singleton M. E., Miliczky**

E., Hogg D. B., Jones V. P. (2013). Trophic hierarchies illuminated via amino acid

isotopic analysis. *PLoS One*, 8(9), e76152. doi:10.1371/journal.pone.0076152

90. **Steffan S. A., Chikaraishi Y., Currie C. R., Horn H., Gaines-Day H. R., Pauli J. N.,**

Zalapa J. E., Ohkouchi N. (2015). Microbes are trophic analogs of animals. *Proc Natl*

Acad Sci U S A, 112(49), 15119-15124. doi:10.1073/pnas.1508782112

91. **Steffan S. A., Chikaraishi Y., Dharampal P. S., Pauli J. N., Guedot C., Ohkouchi N.**

(2017). Unpacking brown food-webs: Animal trophic identity reflects rampant

microbivory. *Ecol Evol*, 7(10), 3532-3541. doi:10.1002/ece3.2951

92. **Steffan S. A., Dharampal P. S.** (2019). Undead food-webs: Integrating microbes into

the food-chain. *Food Webs*, 18, e00111. https://doi.org/10.1016/j.fooweb.2018.e00111

93. **Stock S. P.** (2019). Partners in crime: symbiont-assisted resource acquisition in

Steinernema entomopathogenic nematodes. *Curr Opin Insect Sci*, 32, 22-27.

doi:10.1016/j.cois.2018.10.006

94. **Szklarczyk D., Gable A. L., Lyon D., Junge A., Wyder S., Huerta-Cepas J.,**

Simonovic M., Doncheva N. T., Morris J. H., Bork P., Jensin L. J., Mering C. V.

(2019). STRING v11: protein-protein association networks with increased coverage,

supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids*

Res, 47(D1), D607-D613. doi:10.1093/nar/gky1131

95. **Takizawa Y., Takano Y., Choi B., Dharampal P. S., Steffan S. A., Ogawa N. O.,**

Ohkouchi N., Chikaraishi Y. (2020). A new insight into isotopic fractionation associated

with decarboxylation in organisms: implications for amino acid isotope approaches in

biogeoscience. *Prog Earth Planet Sci* 7, 50. https://doi.org/10.1186/s40645-020-00364-w

96. **Tang H., Pang S.** (2016). Proline Catabolism Modulates Innate Immunity in

Caenorhabditis elegans. Cell Rep, 17(11), 2837-2844. doi:10.1016/j.celrep.2016.11.038

97. **Tang J.** (2011). Microbial metabolomics. *Curr Genomics, 12*(6), 391-403.

doi:10.2174/138920211797248619

98. **Teulier L., Weber J. M., Crevier J., Darveau C. A. (**2016). Proline as a fuel for insect

flight: enhancing carbohydrate oxidation in hymenopterans. *Proc Biol Sci,* 283(1834).

doi:10.1098/rspb.2016.0333

99. **Tsai C. J., Loh J. M., Proft T.** (2016). *Galleria mellonella* infection models for the study

of bacterial diseases and for antimicrobial drug testing. *Virulence,* 7(3), 214-229.

doi:10.1080/21505594.2015.1135289

100.   **Vallenet D., Labarre L., Rouy Z., Barbe V., Bocs S., Cruveiller S., Lajus A.,**

Pascal G., Scarpelli C., Médigue C. (2006). MaGe: a microbial genome annotation

system supported by synteny results. *Nucleic Acids Res*, 34(1), 53-65.

doi:10.1093/nar/gkj406

101.   **Vivas E.I., Goodrich-Blair H.** (2001) *Xenorhabdus nematophilus* as a model for

1166   host-bacterium interactions: rpoS is necessary for mutualism with nematodes. *Journal of*

1167   *Bacteriology*; 183:4687-4693.

1168   102.   **Volgyi A, Fodor A, Szentirmai A, Forst S.** (1998). Phase Variation in

1169   *Xenorhabdus nematophilus. Appl Environ Microbiol.*;64(4):1188-93. doi:

1170   10.1128/AEM.64.4.1188-1193. PMID: 16349534

1171   103.   **Wang J., Dong Y., Zhou T., Liu X., Deng Y., Wang C., Lee J., Zhang L. H.**

1172   (2013) *Pseudomonas aeruginosa* cytotoxicity is attenuated at high cell density and

1173   associated with the accumulation of phenylacetic acid. *PLoS One.*;8(3):e60187.

1174   doi:10.1371/journal.pone.0060187

1175   104.   **Watford M.** (2008). Glutamine metabolism and function in relation to proline

1176   synthesis and the safety of glutamine and proline supplementation. *J Nutr,* 138(10),

1177   2003S-2007S. doi:10.1093/jn/138.10.2003S

1178   105.   **Worthington R. J., Richards J. J., Melander C.** (2012). Small molecule control

1179   of bacterial biofilms. *Org Biomol Chem*, 10(37), 7457-7474. doi:10.1039/c2ob25835h

1180   106.   **Yokoyama K., Ishijima S. A., Clowney L., Koike H., Aramaki H., Tanaka C.,**

1181   **Makino K., Suzuki M.** (2006). Feast/famine regulatory proteins (FFRPs): *Escherichia*

1182   *coli* Lrp, AsnC and related archaeal transcription factors. *FEMS Microbiol Rev*, 30(1), 89-

1183   108. doi:10.1111/j.1574-6976.2005.00005.x

1184   107.   **You D., Yao L. L., Huang D., Escalante-Semerena J. C., Ye B. C.** (2014).

1185   Acetyl coenzyme A synthetase is acetylated on multiple lysine residues by a protein

1186   acetyltransferase with a single Gcn5-type N-acetyltransferase (GNAT) domain in

1187   *Saccharopolyspora erythraea. J Bacteriol.*;196(17):3169-3178. doi:10.1128/JB.01961-14

1188   108.   **Zečić A., Dhondt I., Braeckman B.P.** (2019). The nutritional requirements of

1189   *Caenorhabditis elegans. Genes Nutr* 14, 15 (2019). https://doi.org/10.1186/s12263-019-

1190   0637-7

109.   **Zelcbuch L., Lindner S. N., Zegman Y., Vainberg Slutskin I., Antonovsky N.,**

Gleizer S., Milo R., Bar-Even A. (2016). Pyruvate Formate-Lyase Enables Efficient

Growth of *Escherichia coli* on Acetate and Formate. *Biochemistry*, 55(17), 2423-2426.

doi:10.1021/acs.biochem.6b00184

110.   **Zheng R., Feng X., Wei X., Pan X., Liu C., Song R., Jin Y., Bai F., Jin S., Wu**

Wu., Cheng Z. (2018). PutA Is Required for Virulence and Regulated by PruR in

Pseudomonas aeruginosa. Front Microbiol, 9, 548. doi:10.3389/fmicb.2018.00548

1198

Supplementary Figure 1: Venn diagram comparing the differentially expressed transcripts between the 5 microarrays. Visualization provided from: http://bioinformatics.psb.ugent.be/webtools/Venn/

Supplemental Figure 2: Heatmap of metabolites clusters with pairwise correlation displayed. Red indicates positive correlation, blue indicates negative correlation.