1

Cancer blues? A validated judgment bias task suggests pessimism in nude mice with

tumors

Resasco, A[1,2]*., MacLellan, A.[3]*, Ayala, M. A.[2], Kitchenham, L[4]., Edwards, A. M.[5], Lam,

S.[6], Dejardin, S.[7] & Mason, G.[3]**

[1] Institute of Cell Biology and Neurosciences, National Scientific and Technical Research

Council-University of Buenos Aires, Autonomous City of Buenos Aires, Argentina.

[2] Laboratory of Experimental Animals, Faculty of Veterinary Sciences, National University

of La Plata, La Plata, Argentina.

[3] Department of Integrative Biology, University of Guelph, Guelph, Canada

[4] Department of Animal Biosciences, University of Guelph, Guelph, Canada

[5] Ontario Agricultural College, University of Guelph, Guelph, Canada

[6] Ontario Veterinary College, University of Guelph, Guelph, Canada

[7] Formerly Department of Animal Biosciences, University of Guelph, Guelph, Canada

These authors contributed equally to this work

** Author for correspondence: Georgia Mason

email: gmason@uoguelph.ca

Abstract

In humans, affective states can bias responses to ambiguous information: a phenomenon termed judgment bias (JB). Judgment biases have great potential for assessing affective states in animals, in both animal welfare and biomedical research. New animal JB tasks require construct validation, but for laboratory mice (*Mus musculus*), the most common research vertebrate, a valid JB task has proved elusive. Here (Experiment 1), we demonstrate construct validity for a novel mouse JB test: an olfactory Go/Go task in which subjects dig for high- or low-value food rewards. In C57BL/6 and Balb/c mice faced with ambiguous cues, latencies to dig were sensitive to high/low welfare housing, environmentally-enriched animals responding with relative 'optimism' through shorter latencies. Illustrating the versatility of a validated JB task across fields of research, it further allowed us to test hypotheses about the mood-altering effects of cancer (Experiment 2). Male nude mice bearing subcutaneous lung adenocarcinomas responded more pessimistically than healthy controls to ambiguous cues. Similar effects were not seen in females, however. To our knowledge, this is the first validation of a mouse JB task and the first demonstration of pessimism in tumor-bearing animals. This task, especially if refined to improve its sensitivity, thus has great potential for investigating mouse welfare, the links between affective state and disease, depression-like states in animals, and hypotheses regarding the neurobiological mechanisms that underlie affect-mediated biases in judgment.

Keywords: laboratory mice, judgment bias, validation, affective state, animal welfare, cancer

1. Introduction

Of the 115+ million animals used annually in biomedical research [1], most are rodents. They are often used to model potentially distressing conditions like cancer, arthritis and psychiatric disorders (e.g. anxiety, depression). But even conventional practices like handling (e.g. [2]) and the use of small, non-enriched cages (e.g. [3–5]) can compromise their wellbeing. These welfare costs can modify experimental outcomes in undesired directions [6]. They also have ethical implications, especially given the poor replicability [7] and translatability of biomedical research [8–10]. Our focus here is a potential method for assessing affective states (emotions and long-term moods [11]) in mice: the most widely used vertebrate in both basic and translational research [12]. Such methods are important for assessing mouse welfare, and for understanding the neurobiological mechanisms underlying normal and pathological affective functioning.

In humans, affective states modulate the interpretation of ambiguous information, a phenomenon known as judgment bias (JB). JB refers to the way that individuals experiencing negative affect (e.g. anxiety, depression) can process ambiguous information (e.g. neutral facial expressions) 'pessimistically', as if negative, while individuals in positive states might demonstrate more 'optimistic' interpretations of the same ambiguous cues [see 13–15]. In animal JB studies, optimism can be operationalized as increased expectations of reward when faced with ambiguous cues, and pessimism, by increased expectations of punishment [16]. Harding et al., [17] pioneered this method of animal JB assessment: rats trained that one cue predicts reward while another predicts punishment, were exposed to ambiguous (intermediate) cues. Rats exposed to unpredictable housing showed pessimistic JBs, treating the ambiguous cues as if predicting punishment. Since this seminal work, JB tasks have

gained popularity as potentially powerful tools for assessing animal affect due to their

sensitivity to changes in both valence and intensity of these states [18]. Thus JB tasks have

been developed for a wide range of species (e.g. dogs, sheep, horses, honeybees), using a

variety of cues (e.g. visual, olfactory, tactile), and across diverse fields of research (e.g.

behavioral biology, neuroscience and animal welfare) [19,20].

For mice, however, validated JB tasks had remained elusive. Valid JB tasks must meet two

technical criteria: that animals discriminate between positive and negative cues, and then

interpret intermediate cues as ambiguous [20,21]. But like any putative indicator of affective

state, they must also demonstrate construct validity: sensitivity to deliberate affect

manipulations (c.f. [22,23]). For mice, previous efforts have either not attempted construct

validation (5/15 experiments [24–26]), or attempted it and failed (10/15 experiments [27–

33]; Table S1). Here, we therefore aimed to validate a novel JB task, manipulating affective

state through the use of highly preferred environmentally enriched cages [34], versus

conventional cages known to induce stress [35], anxiety [36,37], and depression-like effects

[35,38,39]. Environmental enrichment (modification of an animal's environment to improve

well-being and meet species-specific needs [40]), has been used in neuroscience for decades

for its positive effects in neuroplasticity and disease recovery [41]. Morphological and

physiological changes in the brain due to enrichment have also been associated with

improved welfare [42], and JB has been shown to be sensitive to the effects of enrichment in

other species (e.g. rats [43,44]).

In a second experiment, we applied this newly validated task to mice with tumors, to assess

its utility in translational biomedical research. It is well established that cancer can be

debilitating when tumors cause pain and discomfort (e.g. [45]), and rodent welfare guidelines

for oncology already focus on such harms (e.g. [46]). However, tumors are known to reduce

human well-being at much earlier stages: tumors can induce depression-like feelings of

sadness and hopelessness [47,48], even before cancer is diagnosed (e.g. [49,50]), thanks to

elevated pro-inflammatory cytokines [51,52]. Mice with tumors likewise show signs of

depression (e.g. increased anhedonia [53]). And again these reflect inflammatory responses

[54–57], and are manifest before clinical signs emerge [46,58]. However, these subtle

changes have received negligible attention in mouse welfare guidelines. Nor have more

nuanced measures of mood yet been developed for researchers interested in the translational

benefits of mouse models of cancer. To bridge these research gaps, we thus aimed to assess

mood in mice with tumors through judgment bias.

2. Materials, methods and results

2.1. Ethical note

Both experiments were approved by institutional ethics committees. Experiment 1 (AUP

#3700) complied with Canadian Council on Animal Care guidelines, and Experiment 2

(protocol number 42-1-14T) complied with Guidelines for the Welfare and Use of Animals

in Cancer Research [46]. One C57 was removed before testing for barbering a cagemate

(Experiment 1), and one male nude mouse was removed due an eye abscess (Experiment 2).

This report also meets ARRIVE (Animal Research: Reporting of *In Vivo* Experiments)

requirements [59].

2.2. Experiment 1: Validating a Novel JB Task with Housing-Manipulated Affective States

2.2.1. Animals and Housing

Eighteen C57BL/6NCrl ('C57') and 18 Balb/cAnNCrl ('Balb') females were purchased from

Charles River (Raleigh, North Carolina) at 3-4 weeks old. Females were chosen to allow for

the combined use of group housing (important since mice are a social species), and

environmental enrichment without the risk of resource guarding aggression that can be

problematic in male mice [60]. Mice were randomly assigned to open-top enriched or

conventional housing treatments (respectively EH or CH). Here they lived in mixed strain

groups (c.f. [61]), each cage containing one C57 and one Balb, plus two DBA/2NCrl

cagemates used in another experiment. CH comprised transparent polyethylene laboratory

cages (27L x 16W x 12H cm, Allentown Inc.; n = 9), with corn cob bedding (Envigo,

Mississauga, Ontario, Canada), a paper cup and two types of nesting material (crinkled paper

strips and cotton pads; Fig. 1 A). EH cages were large (60L x 60W x 30H cm, n=9), opaque

plastic with one transparent red plastic window, containing a variety of enrichments that

facilitate species-typical behaviors (e.g. hiding, climbing, chewing, and nesting [c.f. 39]; Fig.

1 B and C). Attached to each enriched cage was a standard 'annex' cage that mice could

freely access via a tunnel. Mice were trained to enter the annex cage for a food treat when a

cup full of sweet oat cereal (Cheerios) was shaken; the access tunnel could then be blocked

allowing for ease of catching and handling in the annex. Handling for both treatments always

followed cup or tunnel methods to minimize aversive effects [2]. The room was kept at

21±1°C and 35-55% relative humidity, on a reverse 12:12 hour light cycle (lights off at

06:00). Food (Harlan® Teklad, Global Diet 14% protein) and water were *ad libitum*.

Fig 1. Housing treatments and timeline for Experiment 1. A. Conventional 'shoebox' laboratory cage with a paper cup and two types of nest material; B and C. Upper and front view of the enriched cage, respectively; a standard 'annex' cage was attached to one of the sides of the enriched cage to facilitate handling. D. Timeline and summary of positive, negative and ambiguous training and test trials for Experiment 1. DS(+): positive discriminative stimulus, DS(-): negative discriminative stimulus, AMB: ambiguous mixture (50% vanilla-50%mint), B: banana chip, C: Rodent diet ('chow'), X: no food rewards.

2.2.2. Judgment bias (JB) training and testing

Our olfactory, digging-based task utilized a "Go/Go" design that was divided into three phases (Fig. 1 D). All JB training and testing was conducted under red light in an experimental area of the colony room (separated by a plastic curtain), between 08:00 and 18:00. Mice were pseudorandomly assigned to an experimenter blind to treatment (AM or AR), counterbalancing across housing and strains. Randomization was conducted through an online random order generator [62], unless otherwise noted. Mice were fasted for one hour prior to training or testing throughout all phases to increase motivation for food, while also

maintaining the preference for the high- over the low-value rewards [30]. After fasting, mice

were moved from their home cage to a transport cage (27L x 16W x 12H cm, Allentown Inc)

by a familiar research assistant and placed in the experimental area. The order in which mice

were tested was random across days since EH mice were opportunistically caught in the

annex cage (see above). Between trials, all plastic components of the apparatus were wiped

thoroughly with 70% ethanol and disposable materials were replaced.

The apparatus (Fig. 2) comprised a start compartment and two arms, each containing a scent

dispenser at its entrance (a cotton-filled tissue cassette) and a 6.5L x 6.5W x 4H cm corncob-

filled pot at its far end. To prevent the scent of the buried treats from revealing which pot

was rewarded, an inaccessible treat compartment was located at the bottom of each pot with

perforated plastic to allow odor transmission. Treats included in the inaccessible

compartment were dependent on which treat (if any) was accessible, so that each pot always

included a total of one chow piece and one banana chip across compartments. The whole

apparatus was topped with a transparent plexiglass cover.

Fig 2. Judgment bias apparatus used in Experiments 1 and 2. The dotted line represents the sliding

door that was opened at the beginning of each trial.

Pilot tests identified preferred treats (Desjardin unpubl); dried, sweetened banana chips (Stock and Barrel) were selected as the high-value reward and regular rodent chow was used as low-value reward. Vanilla and mint essences (Fleibor S.R.L, Buenos Aires, Argentina), diluted 1:4 in distilled water [63,64], acted as cues (discriminative stimuli: DS). In each trial, one arm of the apparatus was always unscented (marked with distilled water), predicting a buried low-value reward (rodent chow). DS+ or DS- solution (0.1 ml) was applied to the scent dispenser and corncob of the scented arm, respectively predicting buried high-value rewards, banana chips (in positive trials) or no reward (negative trials) (Table 1). Throughout all phases, to facilitate learning and prevent extinction, if a mouse was still eating a reward when the trial ended she was allowed 30 seconds to eat before being handled. Additionally, if she had not yet found the reward by the end of a trial, the appropriate treat was placed on top of the bedding and the mouse was gently guided to it (and given 30 seconds to eat). Each mouse's DS+ (mint or vanilla) and the side of the scented arm (left or right) were pseudorandomly assigned, counterbalancing across strain, housing and experimenter.

2.2.3. Digging Training

One week before training (when 8-9 weeks old), mice were habituated to digging pots with two being placed in their cages daily for 10 minutes, one containing low-value rewards, the other high-value rewards (see Fig. 1 D for full experimental timeline). Digging training in the apparatus began the following week. Here, treats were placed on top of the corn cob bedding on Day 1 and progressively buried in the following four days until they were completely buried at the bottom of the pot by Day 5. Two positive trials were run per day, each lasting 5 minutes. Mice were allowed to freely explore the apparatus and their latency

to start eating each reward was live recorded. Preferences for banana chips over rodent chow

were confirmed on the final day (see Results). All mice were able to find the rewards by the

end of this phase allowing them to move on to the next stage.

2.2.4. Discrimination Training

This phase introduced the negative trial and lasted 10 days, with two trials per mouse each

day. For Days 1-5, the first trial was positive and the second was negative. During Days 6-

10, the order of the positive and negative trial was randomized daily. Latencies to dig and to

eat in both arms were live scored.

2.2.5. Testing for discrimination learning

To confirm successful discrimination of DS+ and DS-, mice underwent 4 reinforced trials

daily for 2-4 days (the length of this phase being variable and determined by how quickly

each mouse reached discrimination criterion; see below). These trials were divided into two

blocks, with one unreinforced (test) trial in the middle to assess their responses to each DS

(Table S2). The order of positive and negative reinforced trials before and after the test was

randomized for each mouse. Positive and negative test trials were presented in alternating

order across days (e.g. Day One DS+ test, Day 2 DS- test, Day 3 DS+ test, etc.). Test trials

lasted 2 minutes and were videoed. Appropriate rewards were placed on the corncob after

each trial (banana for DS+, chow for DS-), and mice were allowed 30 seconds to eat before

being moved back to their transport cage.

Latency to dig, as well as the total duration of digging, in the first and the full 2 minutes of

test trials were recorded by two observers blind to treatment, and their values were averaged

(with videos showing marked discrepancies being re-scored). The discrimination criterion

set required mice to dig for at least twice as long in the DS+ arm than in the DS- arm in the

first minute of testing (with a minimum DS+ digging time of 3 seconds). Mice who met

discrimination criteria moved on to ambiguous cue testing the following day (see below).

Mice who did not yet meet criteria continued to be presented with unrewarded DS+ and DS-

trials until criteria was met. Mice who did not meet criteria within 4 days were excluded from

ambiguous trials (see results).

On the day of ambiguous testing, mice received one positive and one negative trial in random

order, followed by a video-recorded ambiguous unreinforced trial in which an ambiguous

mixture (50% diluted mint, 50% diluted vanilla) marked the scented arm. Again, videos were

scored by two observers, blind to treatment, for latency to dig and digging duration.

Trial Details					
		Experiment 1		Experiment 2	
	Trial type	Scented Arm	Unscented Arm	Scented Arm	Unscented Arm
Digging and Discrimination Training	Pos training	DS+ / Banana	Water / Chow	DS+ / Almond	Water / Corn flake
	Neg training	DS- / No reward	Water / Chow	DS- / No reward	Water / Corn flake
Testing for Discrimination and JB (All conducted with no buried rewards)	Discrimination criterion (DC)	Mice must dig twice as long in the DS+ pot (Pos test) than DS- pot (Neg test), and dig for at least 3 seconds			
	Pos test	DS+	Water	DS+	Water
	Neg test	DS-	Water	DS-	Water
	Ambiguous test	Mixture	Water	Mixture	Water

Table 1. Summary of the trial details in Experiments 1 and 2. DS(+): positive discriminative stimulus,

DS(-): negative discriminative stimulus, Pos: positive, Neg: negative. See Supplemental Table S2 for

expanded table.

2.2.6. In-cage behavioral observations

Behavioral observations were conducted to check for expected differences in welfare

between EH and CH mice (e.g. higher levels of stereotypic behaviors in the latter; [34,39]).

Data were collected via live scan sampling during the dark, active phase. A silent observer

scanned each cage every 15 minutes for four hours, starting two hours after lights off [c.f.

65]. The first observed behavior for each mouse was categorized according to the ethogram

(Table S3). Since EH mice had more opportunities to be out of sight, each behavior was

calculated as a proportion of visible scans.

2.2.7. Statistical analyses for Experiment 1

Generalized Linear Mixed Models in SAS®9.4 were used, on data transformed where needed

to meet assumptions (normality and homogeneity of residuals). Where assumptions could

not be met, non-parametric tests were used instead (and noted in text). Treat preferences were

confirmed during digging training by assessing latency to eat high- and low-value rewards.

The repeated measures model therefore included Reward, Housing, Strain, DS+ Odor and all

two way interactions, plus Cage (a random effect nested in Housing and DS+ odor) and

Mouse ID (a random effect nested in Cage, Housing, DS+ odor and Strain). To test for

judgment bias, repeated measures models were run to assess both latency to dig and digging

duration in the scented arm for positive, negative and ambiguous test trials. Trial Type,

Housing, Strain, Trial Type*Housing, DS+ odor, Trial Type*Strain, Trial Type*DS+ odor,

Trial Type*Housing*DS+ odor, Cage (a random effect nested in Housing and DS+ odor) and

Mouse ID (a random effect nested in Cage, Housing, DS+ odor and Strain) were always

included in Experiment 1 models. To select which additional main and interactive effects to

include (e.g. Tester ID and its interactions), a stepwise forward selection process using

corrected Akaike's Information Criteria (c.f. [66]) identified the most parsimonious final

models. These were then run using maximum-likelihood estimations (the experiment

becoming unbalanced when not all mice met discrimination criteria, c.f. [20]). Since Housing

was the treatment of interest, simple effects were calculated from the Trial Type*Housing,

using the SLICEDIFF command when calculating the Least Squares Means [67]. One-tailed

Ps were used since only one specific effect would validate the task (c.f.[68]): shorter latencies

and longer digging times for EH than CH mice, in ambiguous trials only. Finally, effect sizes

(Cohen's d) were calculated, and ANOVA tables were used to assess whether

Treatment*Trial Type contributed significant variation. To confirm housing effects, levels of

home cage stereotypic behavior (SB) and time spent inactive but awake (IBA) were assessed.

For SB, terms included in the model were Housing, Strain, Housing*Strain and Cage (as a

random effect nested within housing). Home cage IBA data did not meet assumptions of

normality and homogeneity, so a Wilcoxon rank sum test was used instead.

2.3. Experiment 1 results:

All but 4 C57 mice met discrimination criteria (n=31). Treat preference was confirmed by

the lower latencies to eat the high-value reward over the low-value reward the last day of

digging training ($F_{1,32}$=80.46, p<0.0001, *Cohen's d*=2.215). Housing*Reward ($F_{1,32}$=4.12,

p=0.005) and Strain*Reward were significant ($F_{1,32}$=6.89, p=0.007), but banana was still

preferred in all subgroups (p<0.0001). During tests for JB, Trial Type*DS+ odor was

significant for both latency to dig and digging duration (respectively $F_{2,62}$=5.74, p=0.005 and

$F_{2,62}$=18.88, p<0.0001), Mint DS+ mice unexpectedly treating intermediate odor mixtures as

positive (as if 100% mint), but Vanilla DS+ mice treating intermediate odor mixtures as

ambiguous as required for a valid JB task (Fig. 3 A, Table S4). There were also significant

strain differences in latency (latency: $F_{1,31}$=4.87, p=0.018, *Cohen's d*=-0.799; duration:

$F_{1,31}$=3.82, p=0.030, *Cohen's d*=0.484), C57s from both housing conditions showing shorter

latencies and digging durations than Balbs across trials.

Because only Vanilla mice met the requirement of treating the scent mixture as intermediate

between the DS+ and DS-, simple effects of housing were calculated from the Trial

Type*Housing term, using the SLICEDIFF command ([cf. 67]. Housing influenced digging

latencies in the Vanilla DS+ mice, CH animals being slower than EH to dig in ambiguous

trials (ambiguous: t=2.14, d.f.=91.89, p=0.018, *Cohen's d*=1.083; positive: t=0.39,

d.f.=91.89, p=0.348, *Cohen's d*=0.198; negative: t=0.61, d.f.= 91.89, p=0.273, *Cohen's*

d=0.308: Fig. 3 B). Similar effect did not hold for Mint DS+ mice (ambiguous: t=0.68,

d.f.=90.63, p=0.251, *Cohen's d*=0.372; positive: t=-0.77, d.f.=90.63, p=0.221, *Cohen's d*=-

0.425; negative: t=0.75, d.f.= 90.63, p=0.229, *Cohen's d*=0.410), even though Trial

Type*Housing*DS+ odor and Trial Type*Housing did not account for significant variation

(respectively $F_{2,65.37}$=0.49, p=0.344 and $F_{2,62}$=1.41, p=0.252). Digging duration was not

affected by housing, in contrast (e.g. Vanilla DS+ ambiguous trials: t=-0.38, d.f.=91.37,

p=0.353, *Cohen's d*=-0.191).

Latency data were then re-analyzed using only the first minute of testing, to assess the utility

of a shortened protocol. For Vanilla DS+ mice, SLICEDIFF tests for simple effects of

Housing again showed that CH animals had longer latencies to dig in ambiguous trials than

EH mice (ambiguous: t=2.27, d.f.=92.94, p=0.014, Cohen's d=1.148; positive: t=0.22, d.f.=

92.94, p=0.414, Cohen's d=0.110; negative: t=0.80, d.f.= 92.94, p=0.214, Cohen's d=0.404:

Fig. 3 C), while again the same did not hold for Mint DS+ mice (ambiguous: t=0.88,

d.f.=91.94, p=0.193, Cohen's d=0.482; positive: t=-0.65, d.f.= 91.94, p=0.260, Cohen's d=-

0.357; negative t=0.78, d.f.= 91.94, p=0.220, Cohen's d=0.427). This was again despite Trial

Type*Housing*DS+ odor and Trial Type*Housing not accounting for significant variation

(respectively $F_{2,65.37}$=0.36, p=0.392 and $F_{2,62}$=1.66, p=0.198). Consistent with 2-minute

results, C57s showed shorter latencies in the first minute ($F_{1,31}$=8.49, p=0.003, Cohen's d=-

1.056) and there was a significant effect of Scented arm side ($F_{1,31}$=6.81 p=0.001, Cohen's

d=-0.903).

Analyses of homecage observations confirmed expected housing effects on welfare (c.f.

[38,69]): more stereotypic behavior ($F_{1,17.8}$=25.19, p<0.0001, Cohen's d=1.839) and time

spent 'inactive but awake' (Wilcoxon rank sum test; Z=-2.839 p=0.008 Cohen's d=0.484) in

CH than EH cages. Taken together, these results validated digging latency as a JB indicator

when vanilla is the DS+, and justified using a shortened, '1min' protocol in Experiment 2.

Fig 3. Digging latency least square means (± standard error) during positive, negative and ambiguous

test trials. A. 2min digging latency in mice receiving mint (M, n=15) or vanilla (V, n=16) as the

positive discriminative stimulus (DS+) (data logarithmically transformed), B. 2min digging latency

in Vanilla DS+ mice from conventional (CH, n=7) or enriched (EH, n=9) housing (data

logarithmically transformed), C. 1min digging latency in the same subjects (data Box Cox

transformed).

2.4. Experiment 2: *Applying the Task to Mice with Tumors*

2.4.1. Animals and Housing

Twenty male and 19 female 'nude' mice (stock NLAE:NIH(S)-*Fox1*[nu/nu]), free from common/zoonotic mouse pathogens, were obtained at 4-5 weeks old from La Plata National University's Faculty of Veterinary Sciences and randomly allocated to cancer or cancer-free treatments (Fig. 4 B). We employed human A549 cell line lung adenocarcinoma tissue (from tumors grown in other mice: [70]): widely used in oncology [71] and associated with inflammatory cytokines [72]. The donor mouse was killed by cervical dislocation, and the tumor was aseptically removed and placed into a petri dish with Minimum Essential Medium, where it was divided into 2mm^2 pieces. These pieces were immediately transplanted subcutaneously into the lateral abdominal area of mice anesthetized with ketamine/xylazine (100/10 mg/kg i.p., respectively). During the procedure, mice also received 0.03 ml of 1% lidocaine at the site of the incision and a single dose of 10 mg/kg of tramadol (administered subcutaneously). Post-operative care and ulterior veterinary inspection were performed as previously described [70].

2.4.2. Simplified JB protocol

Preference tests again identified high- and low-value rewards [73] (see Results). Training occurred on a clean bench in the colony room, under white light. A simplified protocol was used (Fig. 4 A, Table S2) in which vanilla was now used as the DS+ for all mice, since only vanilla DS+ mice in Experiment 1 interpreted intermediate mint-vanilla odor mixtures as ambiguous and responded to them with JB. Here, digging training was identical to Experiment 1 but lasted only 4 days, discrimination training lasted 9 days and involved 2, 3-

minute trials per day, and testing lasted 3 days with mice meeting discrimination criteria

being tested for responses to ambiguous cues as in Experiment 1. Daily veterinary checks

assessed clinical signs of disease (c.f. [46])

Fig 4: Timeline of positive, negative and ambiguous training and test trials for Experiment 2 (A),

including a photograph of a mouse bearing a subcutaneous tumor (B).

DS (+): positive discriminative stimulus (vanilla), DS (-): negative discriminative stimulus (mint),

AMB: ambiguous mixture (50% vanilla-50%mint), A: almond, C: cornflake, X: no food rewards.

2.4.3. Statistical analyses for Experiment 2

Data were analyzed with repeated measures Generalized Linear Mixed Models as in

Experiment 1, but model selection was not required due to the simpler design. Trial Type,

Treatment (cancer status), Sex and their two- and 3-way interactions were therefore all

included as fixed effects. Cage and Mouse ID, both nested within sex and treatment, were

included as random effects. The simple effects of cancer status in each Trial Type were again

tested using SLICEDIFF commands (now with two-tailed Ps). Sex differences in tumor

volume and treat preference were also checked. For these, data did not meet assumptions of

normality and homogeneity so a Wilcoxon rank sum test was used.

2.5. Experiment 2 Results:

All but 3 male and 7 female mice met discrimination criteria (n=29). As for Experiment 1, high-value rewards were preferred over low-value rewards (males: Wilcoxon rank sum test; Z =-4.823, $p<0.0001$, *Cohen's d*=0.234; females: Wilcoxon rank sum test; Z =-3.683, $p=0.001$, *Cohen's d*=1.292). Trial Type*Treatment*Sex was significant ($F_{2,58.68}=8.77$, $p<0.001$) because cancer status had different effects on males and females (Fig. 5 A, Table S4). In females, who treated the ambiguous cue as negative, only a trend for increased digging latency in negative trials for tumor-bearing animals was detected (t=-1.71, d.f.=80.65, $p=0.093$, *Cohen's d*=1.001). In males, who in terms of latency treated the ambiguous cue as intermediate, the simple effect of cancer status was significant in ambiguous trials: tumor-bearing mice had longer latencies than controls (t=-2.93, d.f.=81.45, $p=0.005$, *Cohen's d*=1.425). Unexpectedly, these mice also showed shorter latencies in negative trials (t=2.27, d.f.=81.83, $p=0.027$, *Cohen's d*=-1.023). (Fig. 5 B).

No mice presented clinical signs: all remained active with good body condition scores [74], normal gaits, normal skin condition over the tumors, and no nasal or ocular discharge. Following testing, tumor volumes were within accepted ranges [46], and similar between sexes (Wilcoxon rank sum test; Z =-0.7415, $p=0.4698$, *Cohen's d*=1.292).

397　Fig 5. Digging latencies (± standard error) during positive, negative and ambiguous test trials (data

398　logarithmically transformed). *A*. Digging latencies in female (F, n= 12) and male (M, n= 17) nude

399　mice, *B*. Digging latencies in control (C, n=8) and tumor-bearing (T, n=9) males.

3. Discussion

To our knowledge, Experiment 1 represents the first successful construct validation of a

mouse judgment bias (JB) task. Although sometimes omitted (5 studies in Table S1), or

downplayed [26,75,76], the construct validation of any new indicator of affective state is a

crucial step. This is especially important for animal JB tasks, which differ greatly from the

unconditioned human tasks that inspired them [21] and which, perhaps as a consequence, can

sometimes produce counterintuitive results [e.g. 77]. In mice, for instance, altered affective

states have often failed to influence JB (10 studies in Table S1). We suspect that a

combination of factors contributed to the success of this, the 11th published validation

attempt. First, affective state was manipulated through a highly preferred housing system that

consistently produces robust differences in affect [34,36,38,39], resulting in large treatment

effects. In these complex environments, with numerous places to hide or escape, stressful

catching that otherwise could have masked treatment effects [78] was also avoided by

training the mice to enter the 'annex cages'. Second, the task was naturalistic and tailored to

mouse olfactory and digging skills [79,80]. Third, we utilized a sensitive Go/Go design

[21,22]; and rewards were pre-checked for their relative value. In addition, our initial

counter-balanced design allowed us to detect an important asymmetry: when faced with a

50:50 mixture of scents, only mice trained to use vanilla as a DS+ treated this cue as

ambiguous (a technical requirement for a valid JB task). Mice trained to use mint as the DS+

instead of treating these mixtures as 100% mint. This result identified Vanilla DS+ mice as

the only potential candidates for validation of the current JB task (an issue we revisit below,

in the context of future research directions).

For these Vanilla DS+ mice, latency to dig in ambiguous trials proved sensitive to well-being,

thus showing construct validation: CH mice showed the predicted "pessimism", taking longer

to commence digging. Digging duration, however, proved insensitive to affect-modulated

JB: although the expected graded response across trials was detected for this variable, no

significant effect of housing was observed. This is perhaps because during ambiguous trials,

subjects were influenced by the absence of reward, having learned in the positive and

negative unreinforced test trials that if a treat was not detected early, further digging was not

beneficial. However, it might also be that digging duration is only impacted by more severe

affect manipulations (as we explore further below). Importantly, we were also able to

demonstrate the versatility of this novel JB task across strains. Though strain differences

were observed in Experiment 1, C57s being faster to begin digging while showing shorter

digging durations, such effects were consistent across trial type, and housing treatments. This

suggests that when technical criteria are met -- mice being able to discriminate between

positive and negative cues, and interpreting intermediate cues as ambiguous -- the task has

potential generalizability across strains. Moreover, the nude mice of Experiment 2 (with a

Swiss genetic background), were also able to meet these technical criteria even after a

shortened protocol.

In Experiment 2, an abbreviated form of this task (the labor-intensiveness of JB testing being

a challenge [81,82]) thus tested hypotheses about the affective impacts of cancer. In males

faced with ambiguous cues, tumor implantation reduced digging latencies indicating negative

JBs. This, the first evidence of pessimism in tumor-bearing animals, is consistent with the

low mood commonly reported in human patients and the depression-like behavioral changes

commonly found in rodent models (see Introduction). The degree of pessimism in tumor-

bearing males was surprising however, ambiguous cues were treated even more negatively

than negative trials. This effect may parallel the influence that human affective states have

on response latency during JB tasks: individuals in some negative states are slower to make

decisions when presented with ambiguous cues [83,84]. It is also unclear why females

showed no tumor-induced JB. It is possible that this reflects genuine sex differences (since

male mice seem more adversely affected than females later in disease progression [85]). But

this might instead indicate floor effects, since even control females seemed to treat

ambiguous cues as negative, perhaps because all female mice were experiencing negative

affect from being kept in barren CH (c.f. Experiment 1). The results from tumor-bearing mice

thus confirmed our new task's potential for assessing affective state in biomedical research.

But they also highlighted needs for further validation and refinement.

Together, our results thus identify a valid JB task with great promise, but also show that its

sensitivity needs improving. Future avenues could include reducing potential floor effects in

replicate cancer research by housing nude mice with enrichment [c.f. 73]. Reinvestigating

the construct validity of digging duration would also be warranted. Incidental findings

revealed that despite Experiment 1's null results, Experiment 2's tumor-bearing males

(pessimistic according to JB latencies) also showed shorter digging durations in ambiguous

trials, suggesting that this measure too was sensitive to low mood (see Figure S1 and Table

S5). Future studies could thus evaluate more negative modifications of affective state in

replicate validation work, to assess whether digging duration is sensitive to these more

challenging manipulations (Experiment 1's null results then being floor effects).

Experimenting with different training odors, and also different mixtures as ambiguous cues,

is also now warranted. To illustrate, Experiment 1's Mint DS+ mice treated 50/50 mixtures

as 100% mint (i.e. positive) and even to human noses these mixtures smelled strongly

'minty'. This suggests that when using a mint DS+, more vanilla-skewed mixtures are needed

for them to be treated as intermediate. Further, when faced with the same mixtures, female

nude mice trained with Vanilla as DS+ treated them as *negative*, suggesting that for these

mice, 'near positive' ambiguous cues skewed toward vanilla would be needed for a mixture

to be treated as intermediate. Pilot tests to identify appropriate odor mixtures, and/or use of

multiple ambiguous probes could help mitigate such effects. Exposing subjects to a full

spectrum of ambiguous cues (e.g. near positive, intermediate, near negative) could also allow

for distinction between different types of negative states (e.g. anxiety- and depression-like

responses, see [18,21]). And as a final avenue for future research, since housing and cancer

status are both long term manipulations, future work assessing the sensitivity of this JB task

to acute changes in affect would be of great benefit.

Such methodological refinements are important given the great research value of a validated

murine JB task. As a humane technique potentially sensitive to both positive and negative

affect [11,86], it could be hugely useful for assessing and improving mouse husbandry.

Further, cognitive biases play important roles in human affective disorders, yet relatively

little is known about the neurophysiological correlates and underlying mechanisms [84].

Since mice are the dominant model species in cognitive and behavioral neuroscience, a

validated murine JB task opens the doors to further investigation and understanding of these

mechanisms. It could be valuable in biomedical research too. In oncology, for instance, a

validated JB task could help investigate how cancer influences affective states (as in our

preliminary work). But it could also help investigate the contradictory effects of opiates on

cancer pathogenicity [c.f. 87]; the adverse effects of chemotherapy on wellbeing [88]; and

the role of mood in the cancer-protective effects of housing rodents with warmth [89],

companions [90] and enrichments [91]: fascinating topics with both animal welfare and

clinical implications.

4. Conclusion

In summary, this novel JB task has proven to be a valid indicator of affective state. For C57

and Balb females, CH animals showed predicted pessimistic responses to ambiguous cues

when their latency to dig was assessed. For tumor-bearing male nude mice, this task also indicated negative states through pessimistic responses to ambiguity, even before the occurrence of clinical signs of disease. Together, these results highlight the potential value of this novel murine JB task across diverse fields of research. This indicator of affective state is easy to implement, economic, and has already proven effective in three common mouse strains. But in addition to its utility, this JB task is also humane: an uncommon feature in available indicators of mouse affect, which often involve aversive experiences (e.g. exposure to open fields, elevated platforms, electrical shocks). This JB task thus offers a welfare-friendly alternative to standard indicators of affect. Although replication and further refinements to improve sensitivity are still needed, a valid JB task has potential in animal welfare assessment and addressing fundamental questions about affective state.

5. Acknowledgments

The authors would like to acknowledge that Experiment 1 was completed on the ancestral lands of the Attawandaron people and the treaty lands and territory of the Mississaugas of the Credit. The authors would also like to thank the mice, Dr. Jim Petrick for providing comments on the manuscript and our wonderful animal care technicians Michaela Randal and Michelle Cieplak.

6. Funding Sources

An NSERC Natural Sciences and Engineering Research Council (Canada) Discovery grant to GJM (no. 05828 / 145607139), and a grant from the National University of La Plata to MAA (Argentina) (no. 11/V253).

References

[1]   J. Keen, Wasted money in United States biomedical and agricultural animal research, in: K. Herrmann, K. Jayne (Eds.), Animal Experimentation: Working Towards a Paradigm Change, Brill, Leiden, The Netherlands, 2019: pp. 244–272.

[2]   K. Gouveia, J.L. Hurst, Optimising reliability of mouse performance in behavioural testing: The major role of non-aversive handling, Sci. Rep. 7 (2017) 44999. https://doi.org/10.1038/srep44999.

[3]   J.P. Balcombe, Laboratory environments and rodents' behavioural needs: A review, Lab. Anim. 40 (2006) 217–235. https://doi.org/10.1258/002367706777611488.

[4]   V. Baumans, The impact of the environment on laboratory animals, in: M.L. Andersen, S. Tufik (Eds.), Rodent Model as Tools in Ethical Biomedical Research, Springer, 2016: pp. 13–22.

[5]   I.A.S. Olsson, K. Dahlborn, Improving housing conditions for laboratory mice: A review of "environmental enrichment," Lab. Anim. 36 (2002) 243–270. https://doi.org/10.1258/002367702320162379.

[6]   T. Poole, Happy animals make good science, Lab. Anim. 31 (1997) 116–124. https://doi.org/10.1258/002367797780600198.

[7]   L.P. Freedman, I.M. Cockburn, T.S. Simcoe, The economics of reproducibility in preclinical research, PLoS Biol. 13 (2015) e1002165. https://doi.org/10.1371/journal.pbio.1002165.

[8]   I.W.Y. Mak, N. Evaniew, M. Ghert, Lost in translation: Animal models and clinical

547  trials in cancer treatment, Am. J. Transl. Res. 6 (2014) 114–118.

548 [9] C.H. Wong, K.W. Siah, A.W. Lo, Estimation of clinical trial success rates and

549  related parameters, Biostatistics. 20 (2019) 273–286.

550  https://doi.org/10.1093/biostatistics/kxx069.

551 [10] J.P. Garner, The significance of meaning: Why do over 90% of behavioral

552  neuroscience results fail to translate to humans, and what can we do to fix it?, ILAR

553  J. 55 (2014) 438–456. https://doi.org/10.1093/ilar/ilu047.

554 [11] M. Mendl, O.H.P. Burman, R.M.A. Parker, E.S. Paul, Cognitive bias as an indicator

555  of animal emotion and welfare: Emerging evidence and underlying mechanisms,

556  Appl. Anim. Behav. Sci. 118 (2009) 161–181.

557  https://doi.org/10.1016/j.applanim.2009.02.023.

558 [12] European Commission, Report From the Commission to the European Parlaiment

559  and the Council, 2020. https://eur-lex.europa.eu/legal-

560  content/EN/TXT/HTML/?uri=CELEX:52020DC0016&from=EN.

561 [13] I. Blanchette, A. Richards, The influence of affect on higher level cognition: A

562  review of research on interpretation, judgement, decision making and reasoning,

563  Cogn. Emot. 24 (2010) 561–595. https://doi.org/10.1080/02699930903132496.

564 [14] C. MacLeod, I.L. Cohen, Anxiety and the Interpretation of Ambiguity: A Text

565  Comprehension Study, J. Abnorm. Psychol. 102 (1993) 238–247.

566  https://doi.org/10.1037/0021-843X.102.2.238.

567 [15] A. Mathews, C. MacLeod, Cognitive approaches to emotions, Anu. Rev. Psychol. 45

568  (1994) 25–50.

569   [16]   C. Douglas, M. Bateson, C. Walsh, A. Bédué, S.A. Edwards, Environmental

570   enrichment induces optimistic cognitive biases in pigs, Appl. Anim. Behav. Sci. 139

571   (2012) 65–73. https://doi.org/10.1016/j.applanim.2012.02.018.

572   [17]   E.J. Harding, E.S. Paul, M. Mendl, Cognitive bias and affective state, Nature. 427

573   (2004) 312. https://doi.org/10.1038/427312a.

574   [18]   M. Mendl, O.H.P. Burman, R.M.A. Parker, E.S. Paul, Cognitive bias as an indicator

575   of animal emotion and welfare: Emerging evidence and underlying mechanisms,

576   Appl. Anim. Behav. Sci. 118 (2009) 161–181.

577   https://doi.org/10.1016/j.applanim.2009.02.023.

578   [19]   M. Mendl, E. Paul, Getting to the heart of animal welfare. The study of animal

579   emotion, Stichting Animales, Netherlands, 2017.

580   [20]   L. Gygax, The A to Z of statistics for testing cognitive judgement bias, Anim.

581   Behav. 95 (2014) 59–69. https://doi.org/10.1016/j.anbehav.2014.06.013.

582   [21]   S. Roelofs, H. Boleij, R.E. Nordquist, F.J. van der Staay, Making decisions under

583   ambiguity: Judgment bias tasks for assessing emotional state in animals, Front.

584   Behav. Neurosci. 10 (2016) 1–16. https://doi.org/10.3389/fnbeh.2016.00119.

585   [22]   M. Lagisz, J. Zidar, S. Nakagawa, V. Neville, E. Sorato, E.S. Paul, M. Bateson, M.

586   Mendl, H. Løvlie, Optimism, pessimism and judgement bias in animals: A

587   systematic review and meta-analysis, Neurosci. Biobehav. Rev. 118 (2020) 3–17.

588   https://doi.org/10.1016/j.neubiorev.2020.07.012.

589   [23]   J. Ahloy-Dallaire, J. Espinosa, G. Mason, Play and optimal welfare: Does play

590   indicate the presence of positive affective states?, Behav. Processes. 156 (2018) 3–

15. https://doi.org/10.1016/j.beproc.2017.11.011.

[24]   S. Jones, E.S. Paul, P. Dayan, E.S.J. Robinson, M. Mendl, Pavlovian influences on

learning differ between rats and mice in a counter-balanced Go/NoGo judgement

bias task, Behav. Brain Res. 331 (2017) 214–224.

https://doi.org/10.1016/j.bbr.2017.05.044.

[25]   S. Hintze, L. Melotti, S. Colosio, J.D. Bailoo, M. Boada-Saña, H. Würbel, E.

Murphy, A cross-species judgement bias task: Integrating active trial initiation into a

spatial Go/No-go task, Sci. Rep. 8 (2018) 1–13. https://doi.org/10.1038/s41598-018-

23459-3.

[26]   V. Krakenberg, I. Woigk, L. Garcia Rodriguez, N. Kästner, S. Kaiser, N. Sachser,

S.H. Richter, Technology or ecology? New tools to assess cognitive judgement bias

in mice, Behav. Brain Res. 362 (2019) 279–287.

https://doi.org/10.1016/j.bbr.2019.01.021.

[27]   H. Boleij, J. van't Klooster, M. Lavrijsen, S. Kirchhoff, S.S. Arndt, F. Ohl, A test to

identify judgement bias in mice, Behav. Brain Res. 233 (2012) 45–54.

https://doi.org/10.1016/j.bbr.2012.04.039.

[28]   V. Kloke, R.S. Schreiber, C. Bodden, J. Möllers, H. Ruhmann, S. Kaiser, K.P.

Lesch, N. Sachser, L. Lewejohann, Hope for the best or prepare for the worst?

Towards a spatial cognitive bias test for mice, PLoS One. 9 (2014) e105431.

https://doi.org/10.1371/journal.pone.0105431.

[29]   J. Novak, J.D. Bailoo, L. Melotti, J. Rommen, H. Würbel, An exploration based

cognitive bias test for mice: Effects of handling method and stereotypic behaviour,

PLoS One. 10 (2015) e0130718. https://doi.org/10.1371/journal.pone.0130718.

[30]   J. Novak, K. Stojanovski, L. Melotti, T. Reichlin, R. Palme, H. Würbel, Effects of

stereotypic behaviour and chronic mild stress on judgement bias in laboratory mice,

Appl. Anim. Behav. Sci. 174 (2016) 162–172.

https://doi.org/10.1016/j.applanim.2015.10.004.

[31]   J. Novak, J.D. Bailoo, L. Melotti, H. Würbel, Effect of cage-induced stereotypies on

measures of affective state and recurrent perseveration in CD-1 and C57BL/6 mice,

PLoS One. 11 (2016) e0153203. https://doi.org/10.1371/journal.pone.0153203.

[32]   J.D. Bailoo, E. Murphy, M. Boada-Saña, J.A. Varholick, S. Hintze, C. Baussière,

K.C. Hahn, C. Göpfert, R. Palme, B. Voelkl, H. Würbel, Effects of cage enrichment

on behavior, welfare and outcome variability in female mice, Front. Behav.

Neurosci. 12 (2018). https://doi.org/10.3389/fnbeh.2018.00232.

[33]   V. Krakenberg, V.T. von Kortzfleisch, S. Kaiser, N. Sachser, S.H. Richter,

Differential Effects of Serotonin Transporter Genotype on Anxiety-Like Behavior

and Cognitive Judgment Bias in Mice, Front. Behav. Neurosci. 13 (2019) 263.

https://doi.org/10.3389/fnbeh.2019.00263.

[34]   S.C. Tilly, J. Dallaire, G.J. Mason, Middle-aged mice with enrichment-resistant

stereotypic behaviour show reduced motivation for enrichment, Anim. Behav. 80

(2010) 363–373. https://doi.org/10.1016/j.anbehav.2010.06.008.

[35]   S. Chourbaji, C. Zacher, C. Sanchis-Segura, R. Spanagel, P. Gass, Social and

structural housing conditions influence the development of a depressive-like

phenotype in the learned helplessness paradigm in male mice, Behav. Brain Res. 164

(2005) 100–106. https://doi.org/10.1016/j.bbr.2005.06.003.

[36]   A. Adcock, E. Choleris, M. Denommé, H. Khan, L. Levison, A. MacLellan, B.

Nazal, L. Niel, E. Nip, G. Mason, Where are you from? Female mice raised in

enriched or conventional cages differ socially, and can be discriminated by other

mice, Behav. Brain Res. 400 (2021) 113025.

https://doi.org/10.1016/j.bbr.2020.113025.

[37]   N. Benaroya-Milshtein, N. Hollander, A. Apter, T. Kukulansky, N. Raz, A. Wilf, I.

Yaniv, C.G. Pick, Environmental enrichment in mice decreases anxiety, attenuates

stress responses and enhances natural killer cell activity, Eur. J. Neurosci. 20 (2004)

1341–1347. https://doi.org/10.1111/j.1460-9568.2004.03587.x.

[38]   C. Fureix, M. Walker, L. Harper, K. Reynolds, A. Saldivia-woo, G. Mason,

Stereotypic behaviour in standard non-enriched cages is an alternative to depression-

like responses in C57BL/6 mice, Behav. Brain Res. 305 (2016) 186–190.

https://doi.org/10.1016/j.bbr.2016.02.005.

[39]   E. Nip, A. Adcock, B. Nazal, A. MacLellan, L. Niel, E. Choleris, L. Levison, G.

Mason, Why are enriched mice nice ? Investigating how environmental enrichment

reduces agonism in female C57BL / 6 , DBA / 2 , and BALB / c mice, Appl. Anim.

Behav. Sci. 217 (2019) 73–82. https://doi.org/10.1016/j.applanim.2019.05.002.

[40]   V. Baumans, The laboratory mouse, in: R.C. Hubrecht, J. Kirkwood (Eds.), UFAW

Handbook on the Care and Management of Laboratory and Other Research Animals,

8th ed., Wiley-Blackwell, Oxford, UK, 2010: pp. 276–310.

[41]   G. Kempermann, Environmental enrichment, new neurons and the neurobiology of

individuality, Nat. Rev. Neurosci. 20 (2019) 235–245.

https://doi.org/10.1038/s41583-019-0120-x.

[42] C. Poirier, M. Bateson, F. Gualtieri, E.A. Armstrong, G.C. Laws, T. Boswell, T. V.

Smulders, Validation of hippocampal biomarkers of cumulative affective experience,

Neurosci. Biobehav. Rev. 101 (2019) 113–121.

https://doi.org/10.1016/j.neubiorev.2019.03.024.

[43] S.H. Richter, A. Schick, C. Hoyer, K. Lankisch, P. Gass, B. Vollmayr, A glass full

of optimism: Enrichment effects on cognitive bias in a rat model of depression,

Cogn. Affect. Behav. Neurosci. 12 (2012) 527–542. https://doi.org/10.3758/s13415-

012-0101-2.

[44] N.M. Brydges, M. Leach, K. Nicol, R. Wright, M. Bateson, Environmental

enrichment induces optimistic cognitive bias in rats, Anim. Behav. 81 (2011) 169–

175. https://doi.org/10.1016/j.anbehav.2010.09.030.

[45] J. V. Roughan, C.A. Coulter, P.A. Flecknell, H.D. Thomas, K.J. Sufka, The

conditioned place preference test for assessing welfare consequences and potential

refinements in a mouse bladder cancer model, PLoS One. 9 (2014) e103362.

https://doi.org/10.1371/journal.pone.0103362.

[46] P. Workman, E.O. Aboagye, F. Balkwill, A. Balmain, G. Bruder, D.J. Chaplin, J.A.

Double, J. Everitt, D.A.H. Farningham, M.J. Glennie, L.R. Kelland, V. Robinson,

I.J. Stratford, G.M. Tozer, S. Watson, S.R. Wedge, S.A. Eccles, V. Navaratnam, S.

Ryder, Guidelines for the welfare and use of animals in cancer research, Br. J.

Cancer. 102 (2010) 1555–1577. https://doi.org/10.1038/sj.bjc.6605642.

[47] A. Schrepf, S.K. Lutgendorf, L.M. Pyter, Pre-treatment effects of peripheral tumors

on brain and behavior: Neuroinflammatory mechanisms in humans and rodents,

Brain. Behav. Immun. 49 (2015) 1–17. https://doi.org/10.1016/j.bbi.2015.04.010.

[48] R.J. Dunlop, C.W. Campbell, Cytokines and advanced cancer, J. Pain Symptom

Manage. 20 (2000) 214–232. https://doi.org/10.1016/S0885-3924(00)00199-8.

[49] L. Van Esch, J.A. Roukema, M.F. Ernst, G.A.P. Nieuwenhuijzen, J. De Vries,

Combined anxiety and depressive symptoms before diagnosis of breast cancer, J.

Affect. Disord. 136 (2012) 895–901. https://doi.org/10.1016/j.jad.2011.09.012.

[50] G.D. Friedman, J.S. Skilling, N. V. Udaltsova, L.H. Smith, Early symptoms of

ovarian cancer: A case-control study without recall bias, Fam. Pract. 22 (2005) 548–

553. https://doi.org/10.1093/fampra/cmi044.

[51] B. Bortolato, T.N. Hyphantis, S. Valpione, G. Perini, M. Maes, G. Morris, M.

Kubera, C.A. Köhler, B.S. Fernandes, B. Stubbs, N. Pavlidis, A.F. Carvalho,

Depression in cancer: The many biobehavioral pathways driving tumor progression,

Cancer Treat. Rev. 52 (2017) 58–70. https://doi.org/10.1016/j.ctrv.2016.11.004.

[52] F. Colotta, P. Allavena, A. Sica, C. Garlanda, A. Mantovani, Cancer-related

inflammation, the seventh hallmark of cancer: Links to genetic instability,

Carcinogenesis. 30 (2009) 1073–1081. https://doi.org/10.1093/carcin/bgp127.

[53] M.G. Nashed, E.P. Seidlitz, B.N. Frey, G. Singh, Depressive-like behaviours and

decreased dendritic branching in the medial prefrontal cortex of mice with tumors: A

novel validated model of cancer-induced depression, Behav. Brain Res. 294 (2015)

25–35. https://doi.org/10.1016/j.bbr.2015.07.040.

[54] D.M. Norden, R. Devine, S. Bicer, R. Jing, P.J. Reiser, L.E. Wold, J.P. Godbout,

D.O. McCarthy, Fluoxetine prevents the development of depressive-like behavior in

a mouse model of cancer related fatigue, Physiol. Behav. 140 (2015) 230–235.

https://doi.org/10.1016/j.physbeh.2014.12.045.

[55] D.M. Lamkin, S.K. Lutgendorf, D. Lubaroff, A.K. Sood, T.G. Beltz, A.K. Johnson,

Cancer induces inflammation and depressive-like behavior in the mouse: Modulation

by social housing, Brain. Behav. Immun. 25 (2011) 555–564.

https://doi.org/10.1016/j.bbi.2010.12.010.

[56] M. Yang, J. Kim, J.S. Kim, S.H. Kim, J.C. Kim, M.J. Kang, U. Jung, T. Shin, H.

Wang, C. Moon, Hippocampal dysfunctions in tumor-bearing mice, Brain. Behav.

Immun. 36 (2014) 147–155. https://doi.org/10.1016/j.bbi.2013.10.022.

[57] A.M. Casaril, M. Domingues, S.R. Bampi, D. de Andrade Lourenço, T.Â.

Smaniotto, N. Segatto, B. Vieira, F.K. Seixas, T. Collares, E.J. Lenardão, L.

Savegnago, The antioxidant and immunomodulatory compound 3-[(4-

chlorophenyl)selanyl]-1-methyl-1H-indole attenuates depression-like behavior and

cognitive impairment developed in a mouse model of breast tumor, Brain. Behav.

Immun. 84 (2020) 229–241. https://doi.org/10.1016/j.bbi.2019.12.005.

[58] D.M. Norden, S. Bicer, Y. Clark, R. Jing, C.J. Henry, L.E. Wold, P.J. Reiser, J.P.

Godbout, D.O. McCarthy, Tumor growth increases neuroinflammation, fatigue and

depressive-like behavior prior to alterations in muscle function, Brain. Behav.

Immun. 43 (2015) 76–85. https://doi.org/10.1016/j.bbi.2014.07.013.

[59] N. Percie du Sert, V. Hurst, A. Ahluwalia, S. Alam, M.T. Avey, M. Baker, W.J.

Browne, A. Clark, I.C. Cuthill, U. Dirnagl, M. Emerson, P. Garner, S.T. Holgate,

D.W. Howells, N.A. Karp, S.E. Lazic, K. Lidster, C.J. MacCallum, M. Macleod, E.J.

Pearl, O.H. Petersen, F. Rawle, P. Reynolds, K. Rooney, E.S. Sena, S.D. Silberberg,

T. Steckler, H. Würbel, The ARRIVE guidelines 2.0: Updated guidelines for

reporting animal research*, J. Cereb. Blood Flow Metab. 40 (2020) 1769–1777.

https://doi.org/10.1177/0271678X20943823.

[60] E.M. Weber, J.A. Dallaire, B.N. Gaskill, K.R. Pritchett-Corning, J.P. Garner,

Aggression in group-housed laboratory mice: Why can't we solve the problem?, Lab

Anim. (NY). 46 (2017) 157–161. https://doi.org/10.1038/laban.1219.

[61] M. Walker, C. Fureix, R. Palme, J.A. Newman, J. Ahloy Dallaire, G. Mason, Mixed-

strain housing for female C57BL/6, DBA/2, and BALB/c mice: Validating a split-

plot design that promotes refinement and reduction, BMC Med. Res. Methodol. 16

(2016). https://doi.org/10.1186/s12874-016-0113-7.

[62] M. Haahr, RANDOM.ORG: True Random Number Service, (2021).

https://www.random.org.

[63] J. Chapuis, D.A. Wilson, Cholinergic modulation of olfactory pattern separation,

Neurosci. Lett. 545 (2013) 50–53. https://doi.org/10.1016/j.neulet.2013.04.015.

[64] J. Chapuis, Y. Cohen, X. He, Z. Zhang, S. Jin, F. Xu, D.A. Wilson, Lateral

entorhinal modulation of piriform cortical activity and fine odor discrimination, J.

Neurosci. 33 (2013) 13449–13459. https://doi.org/10.1523/JNEUROSCI.1387-

13.2013.

[65] L. Harper, E. Choleris, K. Ervin, C. Fureix, K. Reynolds, M. Walker, G. Mason,

Stereotypic mice are aggressed by their cagemates, and tend to be poor

demonstrators in social learning tasks, Anim. Welf. 24 (2015) 463–473.

[66] K.P. Burnham, D.R. Anderson, Model selection and multimodel inference: a

practical information-theoretic approach, 2nd ed., Springer, New York, 2002.

[67] J. Wei, R.J. Carroll, K.K. Harden, G. Wu, Comparisons of treatment means when

factors do not interact in two-factorial studies, Amino Acids. 42 (2012) 2031–2035.

https://doi.org/10.1007/s00726-011-0924-0.

[68] G.D. Ruxton, M. Neuhäuser, When should we use one-tailed hypothesis testing?,

Methods Ecol. Evol. 1 (2010) 114–117. https://doi.org/10.1111/j.2041-

210x.2010.00014.x.

[69] G. Mason, J. Rushen, eds., Stereotypic Animal Behaviour—Fundamentals and

Applications to Welfare, 2nd ed., CABI, Wallingford, Oxford, 2006.

[70] A. Resasco, A.C. Carranza Martin, M.A. Ayala, S.L. Diaz, C. Carbone, Non-

aversive photographic measurement method for subcutaneous tumours in nude mice,

Lab. Anim. 53 (2019) 352–361. https://doi.org/10.1177/0023677218793450.

[71] L. Korrodi-Gregório, V. Soto-Cerrato, R. Vitorino, M. Fardilha, R. Pérez-Tomás,

From proteomic analysis to potential therapeutic targets: Functional profile of two

lung cancer cell lines, A549 and SW900, widely studied in pre-clinical research,

PLoS One. 11 (2016) e0165973. https://doi.org/10.1371/journal.pone.0165973.

[72] Q. Huang, L. Duan, X. Qian, J. Fan, Z. Lv, X. Zhang, J. Han, F. Wu, M. Guo, G. Hu,

J. Du, C. Chen, Y. Jin, IL-17 Promotes Angiogenic Factors IL-6, IL-8, and Vegf

Production via Stat1 in Lung Adenocarcinoma, Sci. Rep. 6 (2016) 36551.

https://doi.org/10.1038/srep36551.

[73] A. Resasco, Impacto del desarrollo de la línea tumoral A549 en el bienestar de

ratones de la cepa NLAE:NIH(S)-*Fox1*$^{nu/nu}$, Universidad Nacional De La Plata,

2018.

http://sedici.unlp.edu.ar/bitstream/handle/10915/87590/Documento_completo.pdf?se

quence=1.

[74] M.H. Ullman-Culleré, C.J. Foltz, Body condition scoring: A rapid and accurate

method for assessing health status in mice, Lab. Anim. Sci. 49 (1999) 319–323.

[75] A. Verjat, P. Devienne, H.G. Rödel, C. Féron, More exploratory house mice judge

an ambiguous situation more negatively, Anim. Cogn. 24 (2021) 53–64.

https://doi.org/10.1007/s10071-020-01414-y.

[76] H.A.T. Nguyen, C. Guo, J.R. Homberg, Cognitive Bias Under Adverse and

Rewarding Conditions: A Systematic Review of Rodent Studies, Front. Behav.

Neurosci. 14 (2020) 1–12. https://doi.org/10.3389/fnbeh.2020.00014.

[77] E. Verbeek, D. Ferguson, C. Lee, Are hungry sheep more pessimistic? The effects of

food restriction on cognitive bias and the involvement of ghrelin in its regulation,

Physiol. Behav. 123 (2014) 67–75. https://doi.org/10.1016/j.physbeh.2013.09.017.

[78] E. Nip, The Long-Term Effects of Environmental Enrichment on Agonism in

Female C57BL/6, BALB/c, and DBA/2 Mice, University of Guelph, 2018.

[79] N. Latham, G. Mason, From house mouse to mouse house: The behavioural biology

of free-living Mus musculus and its implications in the laboratory, Appl. Anim.

Behav. Sci. 86 (2004) 261–289. https://doi.org/10.1016/j.applanim.2004.02.006.

[80]   J.W. Young, L.E. Kerr, J.S. Kelly, H.M. Marston, C. Spratt, K. Finlayson, J.

Sharkey, The odour span task: A novel paradigm for assessing working memory in

mice, Neuropharmacology. 52 (2007) 634–645.

https://doi.org/10.1016/j.neuropharm.2006.09.006.

[81]   S. Jones, V. Neville, L. Higgs, E.S. Paul, P. Dayan, E.S.J. Robinson, M. Mendl,

Assessing animal affect: an automated and self-initiated judgement bias task based

on natural investigative behaviour, Sci. Rep. 8 (2018) 12400.

https://doi.org/10.1038/s41598-018-30571-x.

[82]   N.M. Brydges, L. Hall, A shortened protocol for assessing cognitive bias in rats, J.

Neurosci. Methods. 286 (2017) 1–5. https://doi.org/10.1016/j.jneumeth.2017.05.015.

[83]   E.S. Paul, I. Cuthill, G. Kuroso, V. Norton, J. Woodgate, M. Mendl, Mood and the

speed of decisions about anticipated resources and hazards, Evol. Hum. Behav. 32

(2011) 21–28. https://doi.org/10.1016/j.evolhumbehav.2010.07.005.

[84]   A. Schick, M. Wessa, B. Vollmayr, C. Kuehner, P. Kanske, Indirect assessment of

an interpretation bias in humans: Neurophysiological and behavioral correlates,

Front. Hum. Neurosci. 7 (2013) 1–11. https://doi.org/10.3389/fnhum.2013.00272.

[85]   P.F. Cosper, L.A. Leinwand, Cancer causes cardiac atrophy and autophagy in a

sexually dimorphic manner, Cancer Res. 71 (2011) 1710–1720.

https://doi.org/10.1158/0008-5472.CAN-10-3145.

[86]   E.S. Paul, E.J. Harding, M. Mendl, Measuring emotional processes in animals: The

utility of a cognitive approach, Neurosci. Biobehav. Rev. 29 (2005) 469–491.

https://doi.org/10.1016/j.neubiorev.2005.01.002.

[87] B. Afsharimani, K. Kindl, P. Good, J. Hardy, Pharmacological options for the
management of refractory cancer pain—what is the evidence?, Support. Care
Cancer. 23 (2015) 1473–1481. https://doi.org/10.1007/s00520-015-2678-9.

[88] L.J. Wood, L.M. Nail, A. Gilster, K.A. Winters, C.R. Elsea, Cancer chemotherapy-
related symptoms: Evidence to suggest a role for proinflammatory cytokines, Oncol.
Nurs. Forum. 33 (2006) 536–542. https://doi.org/10.1188/06.ONF.535-542.

[89] K.M. Kokolus, M.L. Capitano, C.T. Lee, J.W.L. Eng, J.D. Waight, B.L. Hylander, S.
Sexton, C.C. Hong, C.J. Gordon, S.I. Abrams, E.A. Repasky, Baseline tumor growth
and immune control in laboratory mice are significantly influenced by
subthermoneutral housing temperature, Proc. Natl. Acad. Sci. U. S. A. 110 (2013)
20176–20181. https://doi.org/10.1073/pnas.1304291110.

[90] G.L. Hermes, B. Delgado, M. Tretiakova, S.A. Cavigelli, T. Krausz, S.D. Conzen,
M.K. McClintock, Social isolation dysregulates endocrine and behavioral stress
while increasing malignant burden of spontaneous mammary tumors, Proc. Natl.
Acad. Sci. U. S. A. 106 (2009) 22393–22398.
https://doi.org/10.1073/pnas.0910753106.

[91] L. Cao, X. Liu, E.J.D. Lin, C. Wang, E.Y. Choi, V. Riban, B. Lin, M.J. During,
Environmental and Genetic Activation of a Brain-Adipocyte BDNF/Leptin Axis
Causes Cancer Remission and Inhibition, Cell. 142 (2010) 52–64.
https://doi.org/10.1016/j.cell.2010.05.029.

831