Influence of warming temperatures on coregonine embryogenesis within and among species

Taylor R. Stewart[1,5], Mikko Mäkinen[2], Chloé Goulon[3], Jean Guillard[3], Timo J. Marjomäki[2],

Emilien Lasne[4], Juha Karjalainen[2], and Jason D. Stockwell[5]

[1]Department of Biology, University of Vermont, Burlington, VT, USA

[2]University of Jyväskylä, Jyväskylä, Finland

[3]University Savoie Mont Blanc, INRAE, CARRTEL, Thonon-les-Bains, France

[4]UMR ESE Agrocampus Ouest-INRAE, Rennes, France

[5]Rubenstein Ecosystem Science Laboratory, University of Vermont, Burlington, VT, USA

Correspondence: Taylor R. Stewart, Department of Biology, Rubenstein Ecosystem Science

Laboratory, University of Vermont, 3 College St, Burlington, VT 05401, USA. Email:

taylor.stewart@uvm.edu

ABSTRACT:

The greatest response of lakes to climate change has been the increase in water temperatures on a

global scale. The responses of many lake fishes to warming water temperatures are projected to

be inadequate to counter the speed and magnitude of climate change, leaving some species

vulnerable to decline and extinction. We experimentally evaluated the responses of embryos

from a group of cold, stenothermic fishes (Salmonidae Coregoninae) – within conspecifics

across lake systems, between congeners within the same lake system, and among congeners

across lake systems – to a thermal gradient using an incubation method that enabled global

comparisons. Study groups included cisco (*Coregonus artedi*) from lakes Superior and Ontario

(USA), and vendace (*C. albula*) and European whitefish (*C. lavaretus*) from Lake Southern

Konnevesi (Finland). All species spawn in the fall and their embryos incubate over winter before

hatching in spring. Embryos were incubated at water temperatures of 2.0, 4.5, 7.0, and 9.0°C,

and the responses to the incubation temperatures were quantified for life-history (*i.e.,* embryo

survival and incubation period) and morphological traits (*i.e.,* length-at-hatch and yolk-sac

volume). We found contrasting reaction norms to temperature in embryo survival and similar

reaction norms to temperature for incubation period, length-at-hatch, and yolk-sac volume in

conspecific and congeneric coregonines. For example, congeneric responses differed in

embryonic survival in the same system, suggesting species differences in adaptability to

warming winter temperatures. Differential levels of parental effects were found within and

among study groups and traits suggesting population biodiversity may provide more flexibility

for populations to cope with changing inter-annual environmental conditions. Our results suggest

coregonines may have a wide range of embryo responses to changing winter conditions as a

result of climate change.

Keywords: climate change, embryo incubation, water temperature, thermal habitat, plasticity,

reaction norm, parental effect, Coregonus

INTRODUCTION:

Freshwater lakes are one of the most sensitive ecosystems to climate change (Jenny et al., 2020; Woolway et al., 2020). One of the greatest threats of climate change to lakes is the unprecedented rise of water temperatures on a global scale (Austin & Colman, 2007; Maberly et al., 2020; O'Reilly et al., 2015; Woolway et al., 2017), although the rise is not projected to be consistent across regions, seasons, or lake types (McCullough et al., 2019; O'Reilly et al., 2015). The greatest seasonal increase in water temperature of seasonally ice-covered lakes is projected to take place during the spring (Schindler et al., 1990; Winslow et al., 2017), and the greatest seasonal increase in air temperature is expected during winter in northern Europe and North America (Christensen et al., 2007).

Warming spring water temperatures and increases in the length of the frost-free season can prolong annual growing seasons with warmer summers, longer and warmer autumns, and shorter ice-cover duration (Sharma et al., 2019, 2020). Temperature is an abiotic master factor for aquatic ecosystems because water temperature directly affects the physical and chemical properties of water, and phenology, reproductive events, metabolic rates, growth, and survival of aquatic organisms (Brett, 1979; Brown et al., 2004; Busch et al., 2012; Gillooly et al., 2002; Little et al., 2020; Ohlberger et al., 2007). Although the broader impacts of climate-derived changes in lake dynamics remain unclear (Shatwell et al., 2019), the responses of many lake organisms are projected to be inadequate to counter the speed and magnitude of climate change, leaving some species vulnerable to decline and extinction (Hoffmann & Sgrò, 2011). These pressures present challenges for biodiversity conservation and sustainability of ecosystem services. To navigate challenges, a foundational understanding of the primary threats to aquatic

ecosystems and organisms across a range of spatial scales from local to global is needed

(Halpern et al., 2015; Langhans et al., 2019; Vörösmarty et al., 2010).

The effects of increasing temperature on lake fishes are predicted to lead to declines in cold-

water species and increases in warm-water species (Comte et al., 2013; Hansen et al., 2017).

Species that possess narrow optimal thermal ranges, live near their thermal limits, or have long

development times at cold temperatures are at-risk under warming climate scenarios as

temperature can have strong direct and indirect effects at early-life stages (Blaxter, 1991; Dahlke

et al., 2020; Ficke et al., 2007; Lim et al., 2017; Mari et al., 2016; Pepin, 1991). Unlike their

marine counterparts, most freshwater fishes are restricted to their lake system, where their ability

to evade the effects of climate change is impeded due to the isolated nature of lakes (Ficke et al.,

2007) and limited swimming capacity during early-life stages (Downie et al., 2020; Herbing,

2002). Fundamental questions for eco-evolutionary and conservation biologists in a global

change context include how lake fishes will respond to rising water temperatures and what

adaptive mechanisms may be involved (Hairston et al., 2005; Kinnison & Hairston, 2007;

Pelletier et al., 2009). Shifts in physiology of lake fish populations living close to their upper

thermal limits will be required if species are to persist under increasingly stressful thermal

conditions (Howells et al., 2016; Woolsey et al., 2015).

Freshwater whitefishes, Salmonidae Coregoninae (hereafter coregonines), are of great socio-

economic value (Ebener et al., 2008; Lynch et al., 2015, 2016; Nyberg et al., 2001; Vonlanthen

et al., 2009, 2012), and are also considered to be critically sensitive to the effects of climate

change because they are cold, stenothermic fishes (Elliott & Bell, 2011; Isaak, 2014; Jeppesen et

al., 2012; Jonsson & Jonsson, 2014; Karjalainen et al., 2015; Karjalainen, Jokinen, et al., 2016;

Stockwell et al., 2009). Coregonine fisheries worldwide have experienced population declines

due to highly variable and weak year-class strengths (Anneville et al., 2015; Myers et al., 2015;

Nyberg et al., 2001; Vonlanthen et al., 2012). In the 20th century, causes of decline included

fishing and stocking practices (Anneville et al., 2015) and eutrophication causing poor

incubation conditions (Müller, 1992; Vonlanthen et al., 2012). Today, the trophic state of lakes

and fisheries management practices are improving, but coregonines continue to be the focus of

reintroduction, restoration, and conservation efforts in many lakes (Bronte et al., 2017; Favé &

Turgeon, 2008; Zimmerman & Krueger, 2009). Reasons for declining recruitment are unknown,

but climate change, increasing water temperatures, and habitat degradation are hypothesized as

causal factors (Anneville et al., 2015; Jeppesen et al., 2012; Karjalainen et al., 2015; Karjalainen,

Jokinen, et al., 2016; Marjomäki et al., 2004; Nyberg et al., 2001).

Coregonines generally spawn during late-autumn, embryos incubate over winter, and hatch in

late-spring (Karjalainen et al., 2015; Stockwell et al., 2009). The time between fertilization and

hatching is inversely related to water temperature (Colby & Brooke, 1970, 1973; Karjalainen,

Jokinen, et al., 2016; Luczynski & Kirklewska, 1984; Pauly & Pullin, 1988). Rising spring water

temperatures trigger hatching in autumn-spawned coregonine embryos (Häkkinen et al., 2002;

Karjalainen et al., 2015; Urpanen et al., 2005). The length of the newly-hatched larvae is

negatively correlated with the temperature of incubation (Colby & Brooke, 1970, 1973;

Karjalainen et al., 2015; Luczynski & Kirklewska, 1984). The long period between spawning

and hatching exposes coregonines to a variety of thermal conditions, potentially resulting in a

wide range of environmentally-induced phenotypes or plastic responses (Karjalainen et al., 2015;

Karjalainen, Jokinen, et al., 2016; Karjalainen, Urpanen, et al., 2016). Coregonines are known to

be highly plastic (Muir et al., 2013) and some species (*e.g.,* vendace *Coregonus albula* and

European whitefish *C. lavaretus*) have demonstrated the ability to respond to winter temperature

changes within the limits of phenotypic plasticity and through genetic adaptive changes, such as

different embryo developmental rates under short and long winter conditions (Karjalainen et al.,

2015; Karjalainen, Jokinen, et al., 2016).

Geographic variation is also important to consider with phenotypic plasticity. Many fishes in

high-latitudes are adapted to relatively colder waters, extensive periods of ice cover, and strong

seasonal daylight variations (Reist et al., 2006). Thus, in high-latitude environments, populations

can show differential long-term adaptation to climates across a latitudinal gradient (Chavarie et

al., 2010; Conover & Present, 1990; Wilder et al., 2020; Yamahira & Conover, 2002). For

example, a number of fishes have demonstrated an inverse relationship between length of the

growing season and reproductive characteristics (*i.e.,* countergradient variation; Conover and

Present 1990, Conover and Schultz 1995, Schultz et al. 1996, 1998, Billerbeck et al. 2000,

Jonassen 2000, Yamahira and Conover 2002, Chavarie et al. 2010). Fishes at high latitudes

experience lower temperatures overall and shorter growing seasons and should exhibit lower

standard metabolic rates, growth rates, and smaller size-at-age than individuals at low latitudes

(Reist et al., 2006; White et al., 2012). However, for cold-water stenothermic fishes, water

temperatures at low latitudes may exceed their optimal range for significant portions of the

growing season, or the amount of optimal thermal habitat decrease, while water temperatures at

high latitudes may remain near the optimum for maximal growth efficiency throughout the

growing season (Conover & Schultz, 1995). Because water temperature has a great influence on

fish physiology and varies across latitudes, a wide range of responses by populations to

increasing temperatures across latitudes is possible (Reist et al., 2006). Coregonines occur

broadly across northern latitudes and are an ideal group to test how cold-water fishes may

respond to climate-driven shifts in environmental variables, such as water temperature. Large-

scale experimental studies can aid in understanding the adaptive thermal capacity of fishes from

different latitudes and what level of adaptive response is needed to mitigate the effects of

changing local environments (Hoffmann & Sgrò, 2011).

Our objective was to experimentally analyze the response of coregonine embryos – within

conspecifics across lake systems, between congeners within the same lake system, and among

congeners across all lake systems – to a thermal gradient using an incubation method that

enabled global comparisons. We hypothesized that coregonines have differential levels of

phenotypic plasticity in life-history and morphological traits of embryos in response to warming

winter incubation conditions based on putative adaptation to their local environments. We

predicted coregonines that share the same thermal environment respond similarly and

geographically distinct groups with different thermal environments respond dissimilarly to

increasing incubation temperatures.

METHODS:

Study Species and Locations

We used a cross-lake, cross-continent, cross-species approach to evaluate the responses and

thermal tolerances of coregonine embryos to changing thermal regimes. Wild-caught populations

of cisco (*C. artedi*) in Lake Superior (LS-Cisco; USA/Canada) and Lake Ontario (LO-Cisco;

USA/Canada), and vendace (LK-Vendace) and European whitefish (LK-Whitefish) in Lake

Southern Konnevesi (Finland; Figure 1) were sampled.

Cisco is one of the most widespread of the North American species of coregonines (Eshenroder

et al., 2016) and were one of the most abundant fish in the Great Lakes (Yule et al., 2013). Cisco

is found in north-central to eastern United States and throughout most of Canada, with the lower

Great Lakes close to its southernmost extent (Scott & Crossman, 1973). Cisco spawning is

initiated when water temperatures decrease to 4-5°C in late Autumn (Eshenroder et al., 2016;

Pritchard, 1931) and occurs at different spawning depths. Spawning can occur at depths ranging

from 1-5 m in Lake Ontario (Paufve, 2019; Pritchard, 1931) and 10-64 m in Lake Superior

(Dryer & Beil, 1964; Paufve, 2019). Experimental thermal optima for normal cisco embryo

development is between 2 and 8°C (Brooke & Colby, 1980; Colby & Brooke, 1970). However,

temperature data at historical spawning grounds indicate that *in-situ* incubations typically occur

between 1 and 4°C, with Lake Ontario warmer than Lake Superior (Figure 2; unpublished data).

Vendace and European whitefish are widely distributed in Northern Europe (Sipponen et al.,

2006). Vendace spawning is initiated when water temperatures decrease to 6°C at the end of

October in Lake Southern Konnevesi and lasts 2-3 weeks (Karjalainen & Marjomäki, 2018).

European whitefish spawn 2-3 weeks later in November when water temperatures decrease to 4-

5°C (Karjalainen et al., 2015). Spawning of vendace occurs in the littoral and sublittoral zone of

lakes and eggs dispersed widely at depths less than 20 m (Heikkilä et al., 2006; Karjalainen et

al., In press). Whitefish are known to spawn at shallower depths than vendace. Embryos are

incubated at 1-2°C until the beginning of April, when water temperatures have gradually

180 increased, and hatching peaks after ice-out at 4-6°C (Figure 2, Karjalainen et al. 2015). Although

181 whitefish spawn later than vendace, their hatching occurs at the same time or slightly earlier than

182 vendace (Karjalainen et al., 2015).

184 <u>Adult Collections</u>

185 Adults were sampled using multi-mesh gillnets in Lake Superior, trap nets in Lake Ontario, and

186 seines (vendace) and multi-mesh gillnets (European whitefish) in Lake Southern Konnevesi.

187 Adult field collections occurred during coregonine spawning periods for Lake Ontario and Lake

188 Superior. On Lake Southern Konnevesi, adults were collected prior to spawning and held in

189 aquaculture pools with water fed directly from the lake until spawning was initiated. All

190 sampling, fertilization, and experimental work for study groups on each continent were

191 conducted at a single laboratory in North America (University of Vermont (UVM), USA) and

192 Europe (University of Jyväskylä (JYU), Finland). Experiments were performed during the 2018-

193 19 season in Finland and the 2019-20 season in the USA.

195 For clarity, our operational use of a study group is to represent a single species within a single

196 lake (*e.g.,* cisco in Lake Superior).

198 <u>Crossing Design and Fertilization</u>

199 Eggs and milt were stripped from 12 females and 16 males from each study group and artificially

200 fertilized under a blocked, nested full-sib, half-sib fertilization design to create a maximum of 48

201 full-sibling families nested within half-siblings per group. The crossing design maximized the

202 amount of genetic variation and minimized the potential loss of multiple families if a female or

male produced poor quality gametes, for a given total number of families, compared to a full-

factorial design. Adults used in the experiment were divided into three or four fertilization

blocks. A single block consisted of four males each paired to three unrelated females, where all

offspring of a given female were full siblings. Fertilizations were performed block by block to

ensure germ cell survival.

Approximately 200 eggs per female were fertilized with an equal amount of milt (5-15 μl) from

each male in the block. After the addition of milt, water was added to activate the germ cells and

gently mixed for one minute. The embryos were rinsed with water 2-3 times until the water was

clear. Temperature of the water used during all fertilizations was ca. 4°C. Reconstructed fresh

water was used during fertilizations (OECD ISO 6341:2012) to standardize the chemical

properties of the water used among study groups and between labs. Embryos were transported in

coolers either by shipping overnight for Lake Superior or driven same-day for Lake Ontario. A

temperature logger recorded air temperature inside the cooler during transport (Lake Superior:

mean (SD) = 2.80°C (0.21); Lake Ontario: mean (SD) = 3.28°C (0.37)). No embryo transport

was required for Lake Southern Konnevesi. Demographic data (*e.g.,* total length, mass, and egg

diameter) were collected on adults. Fertilization success was determined by haphazardly taking

10 embryos from each family and assessing under microscopy within 72-hours post-fertilization

(Oberlercher & Wanzenböck, 2016). If fertilization was low (<30%), the family was removed

from the experimental setup.

 <u>Rearing Conditions</u>

Embryos from successfully fertilized families were individually distributed into 24-well cell

culture microplates and incubated in 2 ml of reconstructed fresh water (OECD ISO 6341:2012).

Reconstructed fresh water was used during incubation to maintain sterility, prevent bacterial

growth in the wells, and eliminate the need for fungicide treatments on the embryos. A total of

36 embryos per family were used for Lake Southern Konnevesi species and 48 embryos per

family for each of Lake Ontario and Lake Superior cisco. Families were randomly distributed

across three or four microplates (*i.e.,* 12 eggs per family per microplate and two families per 24-

well microplate). Microplates from each study group were incubated at target constant

temperatures of 2.0 (coldest), 4.5 (cold), 7.0 (warm), and 9.0°C (warmest) and randomly placed

in climate-controlled chambers at UVM (Memmert[®] IPP260Plus) and climate-controlled rooms

at JYU (Huurre[®]). Experimental incubation temperature treatments were chosen to mimic *in-situ*

mean temperatures and to exceed optimum embryonic development temperatures. Forced airflow

was used in both the climate-controlled chambers and rooms to ensure equal air circulation

around the microplates. All microplates were covered to minimize evaporation. Microplate

orientation and position were rotated weekly to eliminate any temperature heterogeneity within

the chambers and rooms. Water temperatures were recorded hourly with loggers (HOBO[®] Water

Temperature Pro v2 at UVM and Escort iMini at JYU) and daily mean water temperatures

calculated. Incubations took place in the dark, with the exception of short (< 60 mins)

maintenance periods. Microplates were checked weekly for dead eggs and the eye-up stage.

During the hatch period, microplates were checked on a two-day cycle for newly hatched larvae.

For cisco, all newly hatched larvae were photographed for life-history and morphological traits

(Nikon[®] D5600 and Nikon[®] AF-S DX 18-55mm lens). Egg size, total length, and yolk-sac axes

were measured from images using Olympus® LCmicro. For Lake Southern Konnevesi, the larvae

were preserved in ethanol at hatch and flushed and soak in distilled water for 15 min before

measuring the total length and fresh mass under the microscope (Karjalainen, 1992).

Mean water temperature during incubations was maintained near the target incubation

temperature for the cold and warm treatments at each lab. Mean incubation water temperatures

for the cold and warmest treatments were lower than the target incubation temperature at JYU,

but not at UVM (Table 1).

All work described here was approved for ethical animal care under University of Vermont's

Institutional Animal Care and Use Committee (Protocol # PROTO202000021).

<u>Life-History and Morphological Traits</u>

Embryo survival was estimated as the percent of embryos surviving between the eye-up and

hatch stages to rule out unfertilized eggs which can bias mortality estimates. Incubation period

was assessed by two variables: the number of days from fertilization to hatching (days post-

fertilization; DPF) and the sum of the degree-days (accumulated degree-days; ADD). Total

length-at-hatch (mm) and yolk-sac volume (YSV; mm^3) were measured from five individuals per

family at, or as close as possible to, 50% hatching for each family. Yolk-sac volume was

calculated assuming the shape of an ellipse (Blaxter, 1963):

$$YSV = \frac{\pi}{6} ab^2 ,$$

268   where a = length of the yolk sac (mm) and b = height of the yolk sac (mm).

 <u>Statistical Analyses and Estimation of Variance Components</u>

Embryo survival was analyzed as a binomial response variable, and incubation period, length-at-

hatch, and yolk-sac volume at hatching as continuous response variables. Early embryo mortality

and variable fertilization success induced from fertilization failure produced inequalities in the

number of offspring among families and an unbalanced design. The sample size for incubation

period is a function of embryo survival and subsequently resulted in an unbalanced design. All

non-proportional data were checked visually for approximate normality using histograms and Q-

Q plots before the analysis with parametric models. A cubic transformation was applied to LAH

for cisco and a cubic root transformation was applied to DPF, ADD, and YSV to normalize the

distributions. Therefore, binary data *(i.e.,* embryo survival) were analyzed with binomial

generalized linear mixed-effects models (LMM) and variables with distributions not strongly

deviating from normal *(i.e.,* incubation period, length-at-hatch, and yolk-sac volume) were

analyzed with restricted maximum likelihood LMMs with the lme4 package (Bates et al., 2015).

To eliminate any confounding effects between continents, conspecific ciscos were analyzed

independently from congeners in Lake Southern Konnevesi, resulting in two models: Great

Lakes (Lake Superior and Lake Ontario) cisco and Lake Southern Konnevesi vendace and

European whitefish. Population (for cisco only), species (for Lake Southern Konnevesi only),

and incubation temperature were included as fixed effects and female, male, family (female and

male combination), and fertilization block as random effects. Because embryos were raised

independently, the replication unit in the statistical models is the individual embryo. Although

incubation temperature was treated as fixed variable based on our experimental design *(i.e.,* few

treatment levels with high replication), we acknowledge that temperature is a continuous,

independent variable in nature. All traits were examined for population or species, depending on

the continent, and incubation temperature effects in addition to individual parental effects

(female, male, and family effects), fertilization block, and all possible interactions with

backward, stepwise effect-selection using the *buildmer* package (Voeten, 2020). The maximal

model for each trait was selected by comparing a model including or lacking the term of interest

to the reference model based on changes in log-likelihood, Akaike information criterion,

Bayesian information criterion, and change in explained deviance. Significance values for the

mixed-effects model parameters (*i.e.,* population, species, incubation temperature, interaction

effects, and any random-effects selected) were determined using a likelihood ratio test between

the maximal model and reduced models with the model parameter of interest removed.

The mixed-effects model output does not produce significance values for model effects;

therefore, significance for population, species, incubation temperature, interaction effects, and

any random-effects selected were determined using a likelihood ratio test between the maximal

model and reduced models with the model effect of interest removed.

To allow for interspecific comparisons, the response to temperature for each trait was

standardized to the optimal temperature for each study group. Based on literature data (cf.

above), the coldest incubation temperature treatment (2.0°C and 2.2°C; Table 1) was assumed to

be the optimal incubation temperature. For each trait, the within-family mean was calculated for

each temperature treatment and the percent change from the optimal temperature estimated.

Standard error was calculated as the among-family variation in percent change.

The phenotypic variance components were partitioned into random effects for female, male,

female x male, and random residual variance components using mixed-effects models with the

fullfact package (Houde & Pitcher, 2019) for each study group and incubation temperature

treatment. Negative variance components were treated as zero (Neff & Pitcher, 2005). The

percent of total phenotypic variation was used to calculate the Pearson correlation coefficient

between each variance component and the increase in incubation temperature for each study

group. A threshold of ±0.7 was used to categorize correlations either positive or negative, with

all values in between as no correlation. European whitefish from Lake Southern Konnevesi were

removed from this analysis due to a low embryo survival and a low number of families.

All analyses were performed in R version 4.0.3 (R Core Team, 2020).

RESULTS:

<u>Spawning Adults</u>

Total lengths and fresh mass of spawning adults used for gamete collection varied widely among

study groups (Table 2). LK-Vendace were notably smaller than all other study groups. The

remaining study groups varied less in size, but LK-Whitefish were smaller than LS-Cisco and

LO-Cisco.

The LK-Vendace females had the smallest egg diameters and LO-Cisco females had the largest

egg diameters among the study groups (Table 3). LK-Whitefish and LS-Cisco egg diameters

were similar (Table 3).

<u>Life-History and Morphological Traits and Variance Components</u>

All cisco traits had significant interaction effects between population and incubation temperature

(maximum $P < 0.001$; Tables 4 and 5). All vendace and European whitefish traits had significant

interaction effects between species and incubation temperature (maximum $P = 0.002$; Tables 4

and 5). The interaction effects precluded any interpretation of main effects, but did suggest

contrasting norms of reaction for the model groups. Below we describe the interaction effects.

All random effects (*i.e.,* female, male, and female x male) were significant (maximum $P = 0.038$)

except male for LAH and female x male for LAH and YSV in cisco (Tables 4 and 5). All

statistical model results can be found in Tables 4 and 5.

Embryo Survival

Embryo survival was highest among all study groups at the coldest temperature and lowest at the

warmest temperature (Figure 3). The effect of temperature for cisco depended on population

because embryo survival was higher for LO-Cisco (99.3%) than LS-Cisco (80.0%) at the coldest

temperature but not different between populations (<0.1%) at the warmest temperature (Figure

3). For Lake Southern Konnevesi, the effect of temperature depended on species because the

difference in embryo survival between LK-Vendace and LK-Whitefish was less pronounced at

the coldest temperature (29.0%) than at the warmest temperature (50.5%; Figure 3). LK-Vendace

and LK-Whitefish embryo survival had a differential temperature response as LK-Whitefish had

a greater decrease (74.4%) than LK-Vendace (17.7%) from the coldest to warmest incubation

temperatures. LK-Whitefish had the strongest, decreasing response to increasing incubation

temperatures compared to all other study groups (Figure 3).

In the phenotypic variance component analysis, the residual error was the largest component of

phenotypic variation in embryo survival (means >55.2%) for all study groups (Figure 4, SI Table

1). The mean female variance had the highest percentage, excluding error, of the phenotypic

variation in embryo survival for LK-Vendace (17.4%), LS-Cisco (24.1%), and LO-Cisco

(19.9%; Figure 4, SI Table 1). The female variance component correlations for embryo survival

had either negative or no correlations to increasing temperature; however, male and error

variances had positive and no correlations suggesting that as the female component decreases at

higher temperatures the importance of the male component and error increases (Table 6).

Incubation Period (days post-fertilization)

The number of days post-fertilization to hatching was highest for all study groups at the coldest

temperature and decreased as temperature increased (Figure 3). For cisco, DPF was higher for

LO-Cisco (179.2 days) than LS-Cisco (154.3 days) at the coldest temperature and the difference

between populations was less pronounced at the warmest temperature (5.0 days; Figure 3).

For Lake Southern Konnevesi, the effect of temperature depended on species because the

difference in DPF between LK-Vendace and LK-Whitefish was less pronounced at the coldest

temperature (8.9 days) than at warmest temperature (27.3 days; Figure 3). All study groups had

similar responses to temperature, with between 54.2 to 68.3% decreases in DPF from the coldest

to warmest treatments. However, LS-Cisco, LO-Cisco, and LK-Whitefish had a greater decrease

in DPF (66.1, 68.3, 65.3%, respectively), than LK-Vendace (54.2%; Figure 3).

In the phenotypic variance component analysis, the residual error was the largest component of

phenotypic variation in DPF (means >60.8%) for LK-Vendace and LS-Cisco (Figure 4, SI Table

1). The mean female variance was the largest phenotypic variation component in DPF for LO-

Cisco (47.1%). LK-Vendace and LS-Cisco had similar mean female variances for DPF across all

temperatures, with 28.1 and 21.0%, respectively (Figure 4, SI Table 1). The DPF correlations for

female effect had a negative response to temperature for LS-Cisco and LO-Cisco (Table 6).

Incubation Period (accumulated degree-days)

Accumulated degree-days were highest for all study groups at 6.9°C (Figure 3). The effect of

temperature for cisco depended on population because ADD was higher for LO-Cisco (531.9 and

547.7 ADD) than LS-Cisco (461.0 and 492.5 ADD) at the cold and warm temperatures,

respectively, and the differences between populations were less pronounced at the coldest and

warmest temperatures (49.2 and 41.3 ADD, respectively; Figure 3). LS-Cisco and LO-Cisco

ADD responded similarly to increasing incubation temperature. For Lake Southern Konnevesi,

the effect of temperature depended on species because the difference in ADD between LK-

Vendace and LK-Whitefish was less pronounced at the coldest temperature (7.7 ADD) than at

the warm temperature (198.1 ADD; Figure 3). LK-Vendace and LK-Whitefish ADD had a

differential temperature response as LK-Vendace had a greater increase (198.4%) than LK-

Whitefish (159.4%) from the coldest to warm treatment. LK-Vendace had the strongest,

increasing response to increasing incubation temperatures compared to all other study groups

(Figure 3).

In the phenotypic variance component analysis and correlations, ADD had a similar response as

DPF among all study groups as the data only had a different temperature scaling factor (Figure 4,

SI Table 1, Table 6).

Length-at-Hatch

All study groups had a common, decreasing response in LAH as temperature increased (Figure

5). For cisco, LAH was higher for LO-Cisco (11.32 and 9.75 mm) than LS-Cisco (10.21 and 8.68

mm) at the coldest and warmest temperatures, respectively, and the difference between

populations was less pronounced at the cold and warm temperatures (0.99 and 0.90 mm,

respectively; Figure 5). LS-Cisco and LO-Cisco responded to increasing incubation temperature

with a 15.9 and 13.8% respective decrease in LAH from the coldest to warmest treatments. For

Lake Southern Konnevesi, the effect of temperature depended on species because the difference

in LAH between LK-Vendace and LK-Whitefish was more pronounced at the cold and warm

temperatures (2.73 and 2.72 mm, respectively) than at the coldest and warmest temperatures

(2.68 and 2.61 mm, respectively; Figure 5). LK-Vendace and LK-Whitefish each responded

similarly to temperature with a 9.0 and 9.2% respective decrease in LAH from the coldest to

warmest treatments. LS-Cisco and LO-Cisco LAH had a stronger, decreasing response to

increasing incubation temperatures than LK-Vendace and LK-Whitefish (Figure 5).

In the phenotypic variance component analysis, the residual error was the largest component of

phenotypic variation in LAH (means >49.2%) for all study groups (Figure 4, SI Table 2). The

mean female variance had the highest percentage, excluding error, of the phenotypic variation in

LAH for LK-Vendace (40.6%), LS-Cisco (38.2%), and LO-Cisco (17.1%; Figure 4, SI Table 2).

The LAH correlations for each study group had a similar response to temperature with negative

or no female and male correlations and positive or no female x male correlations (Table 6).

Yolk-sac Volume

Yolk-sac volume was highest for all study groups at 9.0°C and decreased as temperature decreased (Figure 5). For cisco, the difference in YSV was similar between populations at the warmest incubation temperature (0.04 mm^3) but diverged as incubation temperature decreased; YSV in LO-Cisco (0.40 mm^3) was smaller than LS-Cisco (0.64 mm^3) at the coldest temperature (Figure 5). Yolk-sac volume in LS-Cisco and LO-Cisco responded differently to incubation temperature, with a 203.6 and 311.0% respective increase from the coldest to warmest treatment. For Lake Southern Konnevesi, the effect of temperature depended on species because the difference in YSV between LK-Vendace and LK-Whitefish was less pronounced at the coldest temperature (0.20 mm^3) than at the warmest temperature (1.07 mm^3; Figure 5). LK-Vendace and LK-Whitefish had the strongest response to temperature with an increase in YSV of 445.0 and 536.6% from the coldest to warmest treatment, respectively. LK-Vendace and LK-Whitefish had a stronger, increasing response to increasing incubation temperatures in YSV than LS-Cisco and LO-Cisco (Figure 5).

In the phenotypic variance component analysis, the residual error was the largest component of phenotypic variation in YSV (means >53.9%) for all study groups (Figure 4, SI Table 2). The mean YSV female variance was the highest percentage, excluding error, of the phenotypic variation for LK-Vendace (23.9%), LS-Cisco (20.5%), and LO-Cisco (23.9%; Figure 4, SI Table 2). The YSV correlations for female, female x male, and error variance components had differential responses to temperature with positive female, positive female x male, and negative error correlations for LK-Vendace, while LS-Ciso had inverse correlations to LK-Vendace (Table 6). All LO-Cisco variance components had no correlation to temperature (Table 6).

DISCUSSION:

Our incubation experiments demonstrated both similar and contrasting reaction norms to temperature for life-history and morphological traits in conspecific and congeneric coregonines. First, we found contrasting responses to temperature in embryo survival within and among study groups (*i.e.,* conspecifics and congeners). Second, incubation periods (both DPF and ADD) responded similarly to increasing temperature (negative response for DPF and positive response for ADD) among study groups, however, LK-Vendace had the strongest response and longest incubations across all temperatures. Third, all study groups had similar negative responses to temperature for LAH and positive responses in YSV, with the strongest responses for LAH in LS-Cisco and LO-Cisco and for YSV in LK-Vendace and LK-Whitefish. Lastly, differential levels of parental effects were found within and among study groups and traits.

Embryo survival had an overall negative correlation with increasing temperature among all study groups. However, populations of ciscos and congeneric species from the same lake (*i.e.,* vendace and European whitefish) had contrasting levels of response to temperature. LK-Whitefish embryo survival had the strongest, negative response to temperature (74.4% survival loss) and all other study groups were impacted less (< 26% survival loss) by increasing temperatures. Temperature is a strong driver of coregonine embryo development (Karjalainen et al., 2015) and survival (Brooke & Colby, 1980; Colby & Brooke, 1970; Luczynski & Kirklewska, 1984) but study groups showed contrasted levels of sensitivity to increased incubation temperatures reflecting specific physiological adaptations. Additionally, our experiment, temperature aside, provided near-optimal incubation conditions to individually reared embryos and these conditions

are idealized compared to what occurs in the wild. For instance, embryos in the wild are

deposited on the substratum and are exposed to deposited sediment that can impact survival

(Müller, 1992). Interaction between temperature and sediments are likely, and temperature

increases may act as a catalyzer of embryo sensitivity to sediment stress (Mari et al., 2016, In

press). Even though temperature did negatively impact embryo survival in our experiment, the

effect of temperature in the wild, in combination with other factors, could be even stronger.

Our result that incubation periods from LK-Vendace and LK-Whitefish were longer and had a

strong response to increasing temperatures, even at the warmest incubation temperatures,

supports previous findings that vendace and European whitefish from Lake Southern Konnevesi

have a high degree of developmental flexibility (Karjalainen et al., 2015). The contrasting

response between vendace and European whitefish was likely due to species differences

(Karjalainen et al., 2015) and ecotypes (*i.e.,* benthic versus pelagic; Mcphee et al., 2012).

Additionally, the different magnitude of temperature responses among all congeners suggests a

differential level of developmental plasticity to increasing incubation temperatures among

species and locations. Long, relatively cold incubations may require a shorter period of spring

warming for individuals to initiate hatching, while short, relatively warm incubations may

require a longer period of warmer spring conditions to hatch (Karjalainen et al., 2015). If winter

water temperatures rise as embryos incubate, the ability to match optimal spring nursery

conditions may be weakened (Cushing, 1990; Karjalainen et al., 2015; Myers et al., 2015).

Populations that are more resilient to increasing or variable winter incubation temperatures may

have a better opportunity to regulate ontogeny and control the time of hatching.

Fish spawning strategies are variable, ranging in frequency from daily to once in a lifetime and

in timing from the same time each year to across all seasons (McBride et al., 2015). For many

species, spawning strategies and breeding patterns are constrained by the adult body condition,

gonadogenesis, and the environment (Jørgensen et al., 2006; McBride et al., 2015; Muir et al.,

2014; van Damme et al., 2009). In this context, the short duration of cisco embryo incubation

periods when exposed to high temperatures was notable. High-latitude populations typically

spawn earlier in autumn and may have the opportunity to delay spawning later into the season,

while still providing an adequate incubation period for embryo development, if water

temperatures continue to rise and do not inhibit ovulation as a result of climate change. However,

low-latitude coregonine populations already spawn in late-autumn and early-winter (Eshenroder

et al., 2016; Stockwell et al., 2009), which begs the question: do low-latitude populations have

capacity to spawn later in the winter if temperatures continue to rise? Winter spawning may lead

to less vulnerability to contemporary climate change. For instance, Atlantic herring (*Clupea*

harengus) have both autumn- and winter-spawning stocks in the North Sea that share the same

summer feeding grounds and start oocyte development at the same time (van Damme et al.,

2009). Similarly, coregonines can exhibit contrasting spawning strategies.

Sympatric coregonine species with autumn, winter, and spring-spawning stocks co-occur in

several northern- and central-Eurasian lakes (Eronen & Lahti, 1988; Ohlberger et al., 2008;

Schulz et al., 2006; Schulz & Freyhof, 2003) and allopatric spring-spawning stocks of cisco are

found in Lac des Écorces (southwestern Quebec; Hénault & Fortin, 1989, 1991; Pariseau et al.,

1983). Winter and spring spawners continue oocyte development through autumn which results

in a lower number of larger eggs compared to the autumn-spawning stocks (Eronen & Lahti,

1988; Hénault & Fortin, 1991). Oocyte development is driven by body energy content, and

winter- and spring-spawning stocks may give iteroparous females the chance to mitigate the

disproportionate energy demand toward somatic growth during the summer when metabolic

demands are higher. Consequently, changes in the environment and the condition of an

individual spawning adult could affect future coregonine spawning strategies. Our results suggest

that the cisco embryos examined may not have the developmental plasticity to mitigate the

effects of increased water temperatures during incubation. In this context, adjusting the time of

spawning may be a more efficient long-term life-history strategy than the embryos adapting to

increased temperatures. However, further work is needed to evaluate the reproductive plasticity

of coregonine adults in the face of climate change.

Lake morphology is also important to consider for the question of a winter- or spring-spawning

adaptation; deeper lakes could sufficiently provide cold thermal refuges at greater depths if

suitable spawning habitat is available. Spring-spawning ciscos in Lac des Écorces, where a 4°C

summer stratum does not exist, initiate spawning when spring water temperatures reach 6°C at

depths ranging from 20-30 m (Hénault & Fortin, 1989, 1991). This strategy of spawning in

deeper, colder water allows for normal embryogenesis throughout the summer to mitigate high

water temperatures during the summer period at shallow depths. Model projections of suitable

thermal and oxythermal habitat for cisco indicate deeper and less eutrophic lakes will likely

provide the best cold-water habitat as water temperatures and land uses change (Herb et al.,

2014; Jacobson et al., 2010; Schmitt et al., 2020). While deep lakes may possess acceptable

thermal refugia for coregonines, access to and requirements for suitable spawning and incubation

habitat is unknown for most populations.

In addition to lower survival and shorter incubations as temperature increases, we also found

both similar and contrasting responses to temperature in morphological traits (*i.e.,* length-at-

hatch and yolk-sac volume) among study groups. The contrast in morphological traits among

study groups and locations is likely related to different egg sizes at fertilization. Smaller eggs

will produce smaller larvae, requiring a lower growth and development rate and less demand on

maternal yolk than larger eggs. The demand for yolk and egg size are positively related and

temperature during embryogenesis is positively related to metabolic rate (Hodson & Blunt, 1986;

Kamler, 2008). The steady decline in YSV for LK-Vendace and LK-Whitefish as incubation

temperatures decreased, but less variability in LAH across temperatures, suggests a high

efficiency to convert yolk into somatic tissues across all temperatures where maximal LAH is

constrained by egg size. In contrast, both cisco populations had a steady decline in YSV and

increase in LAH as incubation temperatures decreased, suggesting a low yolk conversion

efficiency at warmer temperatures. All study groups had a decrease in YSV with time as basal

metabolism consumed yolk as a function of the length of incubation. Regardless of the mode, our

results suggest a synergistic relationship among species, location, egg size, incubation period,

and incubation temperature in determining the phenotype of LAH and YSV.

The trade-off between LAH and YSV is well documented in larval fish physiology (Blaxter,

1991). Climate change impacts may only exacerbate the importance of each morphological trait

in determining either a match or mismatch between larval coregonines and their zooplankton

prey. Using winter water temperatures collected from Lake Superior, we can show the theoretical

interactions among water temperature, spawning period, incubation period, LAH, and YSV

supported by our results (Figure 6). If spawning does shift to a later period from rising water

temperatures, we would expect the size of eggs to increase and the fecundity of females to

decrease from environmentally-induced stress on gametogenesis (Figure 6). While our

experiments used constant incubation temperatures due to logistical constraints, the impact

different spring warming rates can have on the time of hatching and the size of larvae should not

be ignored. Lake Southern Konnevesi vendace and European whitefish previously exhibited

flexibility in embryo development rates and feeding windows under different warming scenarios

(Karjalainen et al., 2015). Such complex responses challenge our ability to predict the

downstream impacts changing autumn, winter, and spring water temperatures may have on

embryo and larval phenotypic plasticity.

Traits of embryos depend not only on species, population, and incubation temperature but also

on parental and transgenerational effects (Blaxter, 1963, 1991; Kekäläinen et al., 2018). Our

results suggest that both female and male effects controlled a portion of early-stage offspring

trait phenotypes in coregonines. The variability in phenotypes induced by parental effects can

provide more flexibility for a population to cope with changing inter-annual environmental

conditions, prevent full year-class failure, and ensure population persistence (Karjalainen,

Jokinen, et al., 2016; Oomen & Hutchings, 2015; Wright & Trippel, 2009). In fishes, the female

effect is usually more pronounced than male and female x male interaction effects, and is

stronger in traits directly related to egg size (Huuskonen et al., 2011; Kennedy et al., 2007;

Nagler et al., 2000), and our results support this trend. Residual error estimates, however,

remained high. Intersexual selection and mate pairing has been proposed as an important

component affecting coregonine offspring fitness (Huuskonen et al., 2011; Karjalainen &

Marjomäki, 2018; Wedekind et al., 2008), and may play a role in conserving natural biodiversity

within populations (Anneville et al., 2015). The long-term stability of commercially exploited

stocks, which can experience fisheries-induced evolution, has been linked to population diversity

(Freshwater et al., 2019; Schindler et al., 2010). Spawning stocks that comprise individuals of

variable sizes and ages (*e.g.,* portfolio effect; Schindler et al., 2010) may contribute differently to

spawning, offspring performance, and recruitment (Figge, 2004; Luck et al., 2003), and are

likely an important consideration as the rapid rate of climate change adds additional stressors on

populations.

The methods we used provide reproducible experimental conditions (*e.g.,* uniform water source

between laboratories, no moving water, minimal embryo disturbance, etc.) and standardized

results that can be compared to future experiments that examine temperature responses across a

wider range of populations. Additionally, efforts to include more northerly populations of cisco

from Canada were thwarted due to restrictions on transport of live embryos across an

international border. This further highlights the importance of the standardized experimental

methods we used to allow for future large-scale, cross-laboratory experiments. However, our

results do suggest that some form of latitudinal variation is likely present and promote fruitful

opportunities for future large-scale experimental research on coregonines and other cold,

stenothermic fishes.

Additionally, interpreting the impacts of parental responses within an environmental context

continues to be important to determine how parental effects may mitigate species' responses to

rapid climate change. The existence of varying parental responses raises questions concerning

possible causal mechanisms. Genomic studies will be needed to better understand what is

genetically impacted by increasing temperatures, how it is impacted, and when during

development (*i.e.,* when is temperature most critical; Chen et al., 2018; Narum et al., 2013). A

mechanistic understanding of thermal response from populations across latitudes will be

essential to predict the vulnerability of species and populations to climate change. Additionally,

stocking practices are important to consider, including supportive breeding, and may affect the

adaptive potential of populations through an artificial selection process (Anneville et al., 2015;

Christie et al., 2012; Ford, 2002) and their ability to respond to environmental changes.

Water temperature is fundamental in regulating fish physiology, and environmental variation

during development can play a large role in generating variability in offspring through

phenotypic plasticity (Little et al., 2020). How coregonines respond during the critical embryonic

stage is important to determine whether the capacity to respond to climate change and the

projected increases in their incubation temperatures exists. Knowing how populations have

adapted historically to environmental variability will help us understand the range of possible

responses to climate change and assist managers to keep coregonines out of hot water.

ACKNOWLEDGMENTS:

We thank the staff at the Wisconsin Department of Natural Resources Bayfield Fisheries Field

Station, United States Geological Survey (USGS) Tunison Laboratory of Aquatic Science, New

York State Department of Environmental Conservation Cape Vincent Fisheries Station, and

Konnevesi Research Station and local fishers for conducting field collections of spawning adults.

We also thank Rachel Taylor, Mark Vinson, Dan Yule, Caroline Rosinski, Jonna Kuha, and

Rosanna Sjövik for help with fertilizations and experiment maintenance. This work was funded

by the USGS under Grant/Cooperative Agreement No. (G16AP00087) to the Vermont Water

Resources and Lakes Studies Center. We acknowledge INRAE, French National Research

Institute for Agriculture, Food, and Environment, the UMR CARRTEL (INRAE - USMB), and

the National Science Foundation (award number 1829451) for supporting a workshop to develop

this experiment.

LITERATURE CITED:

Anneville, O., Lasne, E., Guillard, J., Eckmann, R., Stockwell, J. D., Gillet, C., & Yule, D. L.

(2015). Impact of Fishing and Stocking Practices on Coregonid Diversity. *Food and*

Nutrition Sciences, *06*(11), 1045–1055. https://doi.org/10.4236/fns.2015.611108

Austin, J. A., & Colman, S. M. (2007). Lake Superior summer water temperatures are increasing

more rapidly than regional temperatures: A positive ice-albedo feedback. *Geophysical*

Research Letters, *34*(6), 1–5. https://doi.org/10.1029/2006GL029021

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models

Using lme4. *Journal of Statistical Software; Vol 1, Issue 1 (2015)*.

https://www.jstatsoft.org/v067/i01

Billerbeck, J. M., Schultz, E. T., & Conover, D. O. (2000). Adaptive variation in energy

acquisition and allocation among latitudinal populations of the Atlantic silverside.

Oecologia, *122*(2), 210–219. https://doi.org/10.1007/PL00008848

Blaxter, J. H. S. (1963). The influence of egg size on herring larvae (Clupea harengus L). *J.*

Cons. Int. Explor. Mer, *28*, 211–240. https://doi.org/10.1093/icesjms/28.2.211

Blaxter, J. H. S. (1991). The effect of temperature on larval fishes. *Netherlands Journal of*

660 *Zoology*, *42*(2–3), 336–357. https://doi.org/10.1163/156854291X00379

661 Brett, J. R. (1979). Environmental factors and growth. In *Fish physiology, vol. VIII.*

662 *Bioenergetics and growth*. Academic press. https://doi.org/10.1016/S1546-5098(08)60033-

663 3

664 Bronte, C. R., Bunnell, D. B., David, S. R., Gordon, R., Gorsky, D., Millard, M. J., Read, J.,

665 Stein, R. A., & Vaccaro, L. (2017). *Report from the workshop on coregonine restoration*

666 *science*. US Geological Survey. https://doi.org/10.3133/ofr20171081

667 Brooke, L. T., & Colby, P. J. (1980). Development and survival of embryos of lake herring at

668 different constant oxygen concentrations and temperatures. *The Progressive Fish-Culturist*,

669 *42*(1), 3–9. https://doi.org/10.1577/1548-8659(1980)42[3:DASOEO]2.0.CO;2

670 Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). Toward a

671 metabolic theory of ecology. *Ecology*, *85*(7), 1771–1789. https://doi.org/10.1890/03-9000

672 Busch, S., Kirillin, G., & Mehner, T. (2012). Plasticity in habitat use determines metabolic

673 response of fish to global warming in stratified lakes. *Oecologia*, *170*(1), 275–287.

674 https://doi.org/10.1007/s00442-012-2286-z

675 Chavarie, L., Dempson, J. B., Schwarz, C. J., Reist, J. D., Power, G., & Power, M. (2010).

676 Latitudinal variation in growth among Arctic charr in eastern North America: Evidence for

677 countergradient variation? *Hydrobiologia*, *650*(1), 161–177. https://doi.org/10.1007/s10750-

678 009-0043-z

679 Chen, Z., Farrell, A. P., Matala, A., & Narum, S. R. (2018). Mechanisms of thermal adaptation

680 and evolutionary potential of conspecific populations to changing environments. *Molecular*

681 *Ecology*, *27*(3), 659–674. https://doi.org/10.1111/mec.14475

682 Christensen, J. H., Hewitson, B., Busuioc, A., Chen, A., Gao, X., Held, R., Jones, R., Kolli, R.

K., Kwon, W. K., & Laprise, R. (2007). Regional climate projections. In *Climate Change,*

2007: The Physical Science Basis. Contribution of Working group I to the Fourth

Assessment Report of the Intergovernmental Panel on Climate Change, University Press,

Cambridge, Chapter 11 (pp. 847–940).

Christie, M. R., Marine, M. L., French, R. A., & Blouin, M. S. (2012). Genetic adaptation to

captivity can occur in a single generation. *Proceedings of the National Academy of*

Sciences, *109*(1), 238–242. https://doi.org/10.1073/pnas.1111073109

Colby, P. J., & Brooke, L. T. (1970). Survival and development of lake herring (Coregonus

artedii) eggs at various incubation temperatures. *Biology of Coregonid Fishes*, 417–428.

Colby, P. J., & Brooke, L. T. (1973). Effects of Temperature on Embryonic-Development of

Lake Herring (Coregonus-Artedii). *Journal of the Fisheries Research Board of Canada*,

30(6), 799–810. https://doi.org/10.1139/f73-135

Comte, L., Buisson, L., Daufresne, M., & Grenouillet, G. (2013). Climate-induced changes in the

distribution of freshwater fish: Observed and predicted trends. *Freshwater Biology*, *58*(4),

625–639. https://doi.org/10.1111/fwb.12081

Conover, D. O., & Present, M. C. (1990). Countergradient variation in growth rate:

compensation for lenth of the growing season among Atlantic silversides from different

latitudes. *Oecologica*, *83*(3), 316–324. https://doi.org/Doi 10.1007/Bf00317554

Conover, D. O., & Schultz, E. T. (1995). Phenotypic similarity and the evolutionary significance

of countergradient variation. *Trends in Ecology & Evolution*, *10*(6), 248–252.

https://doi.org/10.1016/S0169-5347(00)89081-3

Cushing, D. H. (1990). Plankton production and year-class strength in fish populations: An

update of the match/mismatch hypothesis. *Advances in Marine Biology*, *26*(C), 249–293.

https://doi.org/10.1016/S0065-2881(08)60202-3

Dahlke, F. T., Wohlrab, S., Butzin, M., & Pörtner, H.-O. (2020). Thermal bottlenecks in the life

cycle define climate vulnerability of fish. *Science*, *369*(6499), 65–70.

https://doi.org/10.1126/science.aaz3658

Downie, A. T., Illing, B., Faria, A. M., & Rummer, J. L. (2020). Swimming performance of

marine fish larvae: review of a universal trait under ecological and environmental pressure.

Reviews in Fish Biology and Fisheries, 1–16. https://doi.org/10.1007/s11160-019-09592-w

Dryer, W. R., & Beil, J. (1964). Life history of lake herring in Lake Superior. *Fish. Bull*, *63*(3),

493–530.

Ebener, M. P., Stockwell, J. D., Yule, D. L., Gorman, O. T., Hrabik, T. R., Kinnunen, R. E.,

Mattes, W. P., Oyadomari, J. K., Schreiner, D. R., Geving, S., Scribner, K., Schram, S. T.,

Seider, M. J., & Sitar, S. P. (2008). Status of cisco (Coregonus artedi) in Lake Superior

during 1970-2006 and management and research considerations. *Ann Arbor, Michigan:*

Great Lakes Fishery Commission, Lake Superior Technical Report, *1*, 126.

Elliott, J. A., & Bell, V. A. (2011). Predicting the potential long-term influence of climate

change on vendace (Coregonus albula) habitat in Bassenthwaite Lake, U.K. *Freshwater*

Biology, *56*(2), 395–405. https://doi.org/10.1111/j.1365-2427.2010.02506.x

Eronen, T., & Lahti, E. (1988). Life cycle of winter spawning vendace (Coregonus albula L.) in

Lake Kajoonjärvi, eastern Finland. *Finnish Fisheries Research*, *9*, 197–203.

Eshenroder, R. L., Vecsei, P., Gorman, O. T., Yule, D. L., Pratt, T. C., Mandrak, N. E., Bunnell,

D. B., & Muir, A. M. (2016). Ciscoes (Coregonus, subgenus Leucichthys) of the Laurentian

Great Lakes and Lake Nipigon. In *Canadian Journal of Fisheries and Aquatic Sciences*.

Great Lakes Fishery Commission.

http://www.glfc.org/pubs/misc/Ciscoes_of_the_Laurentian_Great_Lakes_and_Lake_Nipigo

n.pdf

Favé, M.-J., & Turgeon, J. (2008). Patterns of genetic diversity in Great Lakes bloaters

(Coregonus hoyi) with a view to future reintroduction in Lake Ontario. *Conservation*

Genetics, *9*(2), 281–293. https://doi.org/10.1007/s10592-007-9339-6

Ficke, A. D., Myrick, C. A., & Hansen, L. J. (2007). Potential impacts of global climate change

on freshwater fisheries. *Reviews in Fish Biology and Fisheries*, *17*(4), 581–613.

https://doi.org/10.1007/s11160-007-9059-5

Figge, F. (2004). Bio-folio: applying portfolio theory to biodiversity. *Biodiversity &*

Conservation, *13*(4), 827–849. https://doi.org/10.1023/B:BIOC.0000011729.93889.34

Ford, M. J. (2002). Selection in captivity during supportive breeding may reduce fitness in the

wild. *Conservation Biology*, *16*(3), 815–825. https://doi.org/10.1046/j.1523-

1739.2002.00257.x

Freshwater, C., Anderson, S. C., Holt, K. R., Huang, A., & Holt, C. A. (2019). Weakened

portfolio effects constrain management effectiveness for population aggregates. *Ecological*

Applications, *29*(7), e01966. https://doi.org/10.1002/eap.1966

Gillooly, J. F., Charnov, E. L., West, G. B., Savage, V. M., & Brown, J. H. (2002). Effects of

size and temperature on developmental time. *Nature*, *417*(6884), 70–73.

https://doi.org/doi.org/10.1038/417070a

Hairston, N. G., Ellner, S. P., Geber, M. A., Yoshida, T., & Fox, J. A. (2005). Rapid evolution

and the convergence of ecological and evolutionary time. *Ecology Letters*, *8*(10), 1114–

1127. https://doi.org/10.1111/j.1461-0248.2005.00812.x

Häkkinen, J., Vehniäinen, E., Ylönen, O., Heikkilä, J. J., Soimasuo, M., Kaurola, J., Oikari, A.,

& Karjalainen, J. (2002). The effects of increasing UV-B radiation on pigmentation, growth

and survival of coregonid embryos and larvae. *Environmental Biology of Fishes*, *64*(4),

451–459. https://doi.org/10.1023/A:1016165802417

Halpern, B. S., Frazier, M., Potapenko, J., Casey, K. S., Koenig, K., Longo, C., Lowndes, J. S.,

Rockwood, R. C., Selig, E. R., Selkoe, K. A., & Walbridge, S. (2015). Spatial and temporal

changes in cumulative human impacts on the world's ocean. *Nature Communications*, *6*, 1–

7. https://doi.org/10.1038/ncomms8615

Hansen, G. J. A., Read, J. S., Hansen, J. F., & Winslow, L. A. (2017). Projected shifts in fish

species dominance in Wisconsin lakes under climate change. *Global Change Biology*,

23(4), 1463–1476. https://doi.org/10.1111/gcb.13462

Heikkilä, J. J., Huuskonen, H., & Karjalainen, J. (2006). Location of spawning grounds of

vendace (Coregonus albula (L.)): implication for dispersion of newly hatched larvae.

Internationale Vereinigung Für Theoretische Und Angewandte Limnologie:

Verhandlungen, *29*(4), 1725–1728. https://doi.org/10.1080/03680770.2006.11902983

Hénault, M., & Fortin, R. (1989). Comparison of meristic and morphometric characters among

spring-and fall-spawning ecotypes of cisco (Coregonus artedii) in southern Quebec, Canada.

Canadian Journal of Fisheries and Aquatic Sciences, *46*(1), 166–173.

https://doi.org/10.1139/f89-022

Hénault, M., & Fortin, R. (1991). Early life stages, growth, and reproduction of spring-spawning

ciscoes (Coregonus artedii) in Lac des Écorces, Quebec. *Canadian Journal of Zoology*,

69(6), 1644–1652. https://doi.org/10.1139/z91-229

Herb, W. R., Johnson, L. B., Jacobson, P. C., & Stefan, H. G. (2014). Projecting cold-water fish

habitat in lakes of the glacial lakes region under changing land use and climate regimes.

Canadian Journal of Fisheries and Aquatic Sciences, *71*(9), 1334–1348.

https://doi.org/10.1139/cjfas-2013-0535

Herbing, I. H. von. (2002). Effects of temperature on larval fish swimming performance: the

importance of physics to physiology. *Journal of Fish Biology*, *61*(4), 865–876.

https://doi.org/10.1111/j.1095-8649.2002.tb01848.x

Hodson, P. V, & Blunt, B. R. (1986). The effect of time from hatch on the yolk conversion

efficiency of rainbow trout, Salmo gairdneri. *Journal of Fish Biology*, *29*(1), 37–46.

https://doi.org/10.1111/j.1095-8649.1986.tb04924.x

Hoffmann, A. A., & Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*

Australia, *470*(7335), 479–485. https://doi.org/10.1038/nature09670

Houde, A. L., & Pitcher, T. E. (2019). *fullfact: Full Factorial Breeding Analysis*. https://cran.r-

project.org/package=fullfact

Howells, E. J., Abrego, D., Meyer, E., Kirk, N. L., & Burt, J. A. (2016). Host adaptation and

unexpected symbiont partners enable reef□building corals to tolerate extreme temperatures.

Global Change Biology, *22*(8), 2702–2714. https://doi.org/10.1111/gcb.13250

Huuskonen, H., Kekäläinen, J., Panda, B., Shikano, T., & Kortet, R. (2011). Embryonic survival

and larval predator-avoidance ability in mutually ornamented whitefish. *Biological Journal*

of the Linnean Society, *103*(3), 593–601. https://doi.org/10.1111/j.1095-8312.2011.01664.x

Isaak, D. J. (2014). Climate Change and the Future of Freshwater Fisheries. In *Future of*

Fisheries: Perspectives for Emerging Professionals.

Jacobson, P. C., Stefan, H. G., & Pereira, D. L. (2010). Coldwater fish oxythermal habitat in

Minnesota lakes: influence of total phosphorus, July air temperature, and relative depth.

Canadian Journal of Fisheries and Aquatic Sciences, *67*(12), 2002–2013.

798 https://doi.org/10.1139/F10-115

799 Jenny, J.-P., Anneville, O., Arnaud, F., Baulaz, Y., Bouffard, D., Domaizon, I., Bocaniov, S. A.,

800  Chèvre, N., Dittrich, M., & Dorioz, J.-M. (2020). Scientists' Warning to Humanity: Rapid

801  degradation of the world's large lakes. *Journal of Great Lakes Research*.

802  https://doi.org/10.1016/j.jglr.2020.05.006

803 Jeppesen, E., Mehner, T., Winfield, I. J., Kangur, K., Sarvala, J., Gerdeaux, D., Rask, M.,

804  Malmquist, H. J., Holmgren, K., Volta, P., Romo, S., Eckmann, R., Sandström, A., Blanco,

805  S., Kangur, A., Ragnarsson Stabo, H., Tarvainen, M., Ventelä, A. M., Søndergaard, M., …

806  Meerhoff, M. (2012). Impacts of climate warming on the long-term dynamics of key fish

807  species in 24 European lakes. In *Hydrobiologia* (Vol. 694, Issue 1, pp. 1–39).

808  https://doi.org/10.1007/s10750-012-1182-1

809 Jonassen, T. (2000). Geographic variation in growth and food conversion efficiency of juvenile

810  Atlantic halibut related to latitude. *Journal of Fish Biology*, *56*(2), 279–294.

811  https://doi.org/10.1006/jfbi.1999.1159

812 Jonsson, B., & Jonsson, N. (2014). Early environment influences later performance in fishes.

813  *Journal of Fish Biology*, *85*(2), 151–188. https://doi.org/10.1111/jfb.12432

814 Jørgensen, C., Ernande, B., Fiksen, Ø., & Dieckmann, U. (2006). The logic of skipped spawning

815  in fish. *Canadian Journal of Fisheries and Aquatic Sciences*, *63*(1), 200–211.

816  https://doi.org/10.1139/f05-210

817 Kamler, E. (2008). Resource allocation in yolk-feeding fish. *Reviews in Fish Biology and*

818  *Fisheries*, *18*(2), 143. https://doi.org/10.1007/s11160-007-9070-x

819 Karjalainen, J. (1992). Effects of different preservation methods on total length and weight of

820  larval vendace (Coregonus albula (L.)). *Nordic Journal of Freshwater Research (Sweden)*.

Karjalainen, J., Jokinen, L., Keskinen, T., & Marjomäki, T. J. (2016). Environmental and genetic

effects on larval hatching time in two coregonids. *Hydrobiologia*, *780*(1), 135–143.

https://doi.org/10.1007/s10750-016-2807-6

Karjalainen, J., Keskinen, T., Pulkkanen, M., & Marjomäki, T. J. (2015). Climate change alters

the egg development dynamics in cold-water adapted coregonids. *Environmental Biology of*

Fishes, *98*(4), 979–991. https://doi.org/10.1007/s10641-014-0331-y

Karjalainen, J., & Marjomäki, T. J. (2018). Communal pair spawning behaviour of vendace

(Coregonus albula) in the dark. *Ecology of Freshwater Fish*, *27*(2), 542–548.

https://doi.org/10.1111/eff.12368

Karjalainen, J., Tuloisela, M., Nyholm, K., & Marjomäki, T. J. Vendace (Coregonus albula)

disperse their eggs widely during spawning. *Annales Zoologici Fennici*, *In press*.

Karjalainen, J., Urpanen, O., Keskinen, T., Huuskonen, H., Sarvala, J., Valkeajärvi, P., &

Marjomäki, T. J. (2016). Phenotypic plasticity in growth and fecundity induced by strong

population fluctuations affects reproductive traits of female fish. *Ecology and Evolution*,

6(3), 779–790. https://doi.org/10.1002/ece3.1936

Kekäläinen, J., Oskoei, P., Janhunen, M., Koskinen, H., Kortet, R., & Huuskonen, H. (2018).

Sperm pre-fertilization thermal environment shapes offspring phenotype and performance.

Journal of Experimental Biology, *221*(20). https://doi.org/10.1242/jeb.181412

Kennedy, J., Geffen, A. J., & Nash, R. D. M. (2007). Maternal influences on egg and larval

characteristics of plaice (Pleuronectes platessa L.). *Journal of Sea Research*, *58*(1), 65–77.

https://doi.org/10.1016/j.seares.2007.01.003

Kinnison, M. T., & Hairston, N. G. (2007). Eco-evolutionary conservation biology:

Contemporary evolution and the dynamics of persistence. *Functional Ecology*, *21*(3), 444–

454. https://doi.org/10.1111/j.1365-2435.2007.01278.x

Langhans, S. D., Jähnig, S. C., Lago, M., Schmidt-Kloiber, A., & Hein, T. (2019). The potential

of ecosystem-based management to integrate biodiversity conservation and ecosystem

service provision in aquatic ecosystems. *Science of The Total Environment*, *672*, 1017–

1020. https://doi.org/10.1016/j.scitotenv.2019.04.025

Lim, M. Y.-T., Manzon, R. G., Somers, C. M., Boreham, D. R., & Wilson, J. Y. (2017). The

effects of fluctuating temperature regimes on the embryonic development of lake whitefish

(Coregonus clupeaformis). *Comparative Biochemistry and Physiology Part A: Molecular &*

Integrative Physiology, *214*, 19–29. https://doi.org/10.1016/j.cbpa.2017.08.010

Little, A. G., Loughland, I., & Seebacher, F. (2020). What do warming waters mean for fish

physiology and fisheries? *Journal of Fish Biology*, *97*, 328–340.

https://doi.org/10.1111/jfb.14402

Luck, G. W., Daily, G. C., & Ehrlich, P. R. (2003). Population diversity and ecosystem services.

Trends in Ecology & Evolution, *18*(7), 331–336. https://doi.org/10.1016/S0169-

5347(03)00100-9

Luczynski, M., & Kirklewska, A. (1984). Dependence of Coregonus albula embryogenesis rate

on the incubation temperature. *Aquaculture*, *42*(1), 43–55. https://doi.org/10.1016/0044-

8486(84)90312-0

Lynch, A. J., Cooke, S. J., Deines, A. M., Bower, S. D., Bunnell, D. B., Cowx, I. G., Nguyen, V.

M., Nohner, J., Phouthavong, K., Riley, B., Rogers, M. W., Taylor, W. W., Woelmer, W.,

Youn, S.-J., & Beard, T. D. (2016). The social, economic, and environmental importance of

inland fish and fisheries. *Environmental Reviews*, *24*(2), 115–121.

https://doi.org/10.1139/er-2015-0064

Lynch, A. J., Taylor, W. W., Beard, T. D., & Lofgren, B. M. (2015). Climate change projections

for lake whitefish (Coregonus clupeaformis) recruitment in the 1836 Treaty Waters of the

Upper Great Lakes. *Journal of Great Lakes Research*, *41*(2), 415–422.

https://doi.org/10.1016/j.jglr.2015.03.015

Maberly, S. C., O'Donnell, R. A., Woolway, R. I., Cutler, M. E. J., Gong, M., Jones, I. D.,

Merchant, C. J., Miller, C. A., Politi, E., & Scott, E. M. (2020). Global lake thermal regions

shift under climate change. *Nature Communications*, *11*(1), 1–9.

Mari, L., Daufresne, M., Guillard, J., Evanno, G., & Lasne, E. Elevated temperature and

deposited sediment jointly affect early life history traits in southernmost arctic charr

populations. *Canadian Journal of Fisheries and Aquatic Sciences*, *In press*.

https://doi.org/10.1139/cjfas-2020-0256

Mari, L., Garaud, L., Evanno, G., & Lasne, E. (2016). Higher temperature exacerbates the impact

of sediments on embryo performances in a salmonid. *Biology Letters*, *12*(12), 20160745.

https://doi.org/10.1098/rsbl.2016.0745

Marjomäki, T. J., Auvinen, H., Helminen, H., Huusko, A., Sarvala, J., Valkeajärvi, P., Viljanen,

M., & Karjalainen, J. (2004). Spatial synchrony in the inter-annual population variation of

vendace (Coregonus albula (L.)) in Finnish lakes. *Annales Zoologici Fennici*, *41*(1), 225–

240. https://www.jstor.org/stable/23736206

McBride, R. S., Somarakis, S., Fitzhugh, G. R., Albert, A., Yaragina, N. A., Wuenschel, M. J.,

Alonso☐Fernández, A., & Basilone, G. (2015). Energy acquisition and allocation to egg

production in relation to fish reproductive strategies. *Fish and Fisheries*, *16*(1), 23–57.

https://doi.org/10.1111/faf.12043

McCullough, I. M., Cheruvelil, K. S., Collins, S. M., & Soranno, P. A. (2019). Geographic

patterns of the climate sensitivity of lakes. *Ecological Applications*, *29*(2), e01836.

https://doi.org/10.1002/eap.1836

Mcphee, M. V, Noakes, D. L. G., & Allendorf, F. W. (2012). Developmental rate: A unifying

mechanism for sympatric divergence in postglacial fishes? *Current Zoology*, *58*(1), 21–34.

https://doi.org/10.1093/czoolo/58.1.21

Muir, A. M., Arts, M. T., Koops, M. A., Johnson, T. B., Krueger, C. C., Sutton, T. M., &

Rosenfeld, J. (2014). Reproductive life-history strategies in lake whitefish (Coregonus

clupeaformis) from the Laurentian Great Lakes. *Canadian Journal of Fisheries and*

Aquatic Sciences, *71*(8), 1256–1269. https://doi.org/10.1139/cjfas-2013-0254

Muir, A. M., Vecsei, P., Pratt, T. C., Krueger, C. C., Power, M., & Reist, J. D. (2013).

Ontogenetic shifts in morphology and resource use of cisco Coregonus artedi. *Journal of*

Fish Biology, *82*(2), 600–617. https://doi.org/10.1111/jfb.12016

Müller, R. (1992). Trophic state and its implications for natural reproduction of salmonid fish. In

The Dynamics and Use of Lacustrine Ecosystems (pp. 261–268). Springer.

Myers, J. T., Yule, D. L., Jones, M. L., Ahrenstorff, T. D., Hrabik, T. R., Claramunt, R. M.,

Ebener, M. P., & Berglund, E. K. (2015). Spatial synchrony in cisco recruitment. *Fisheries*

Research, *165*, 11–21. https://doi.org/10.1016/j.fishres.2014.12.014

Nagler, J. J., Parsons, J. E., & Cloud, J. G. (2000). Single pair mating indicates maternal effects

on embryo survival in rainbow trout, Oncorhynchus mykiss. *Aquaculture*, *184*(1–2), 177–

183. https://doi.org/10.1016/S0044-8486(99)00309-9

Narum, S. R., Campbell, N. R., Meyer, K. A., Miller, M. R., & Hardy, R. W. (2013). Thermal

adaptation and acclimation of ectotherms from differing aquatic climates. *Molecular*

Ecology, *22*(11), 3090–3097. https://doi.org/10.1111/mec.12240

Neff, B. D., & Pitcher, T. E. (2005). Genetic quality and sexual selection: an integrated

framework for good genes and compatible genes. *Molecular Ecology*, *14*(1), 19–38.

https://doi.org/10.1111/j.1365-294X.2004.02395.x

Nyberg, P., Bergstrand, E., Degerman, E., & Enderlein, O. (2001). Recruitment of pelagic fish in

an unstable climate: studies in Sweden's four largest lakes. *Ambio*, *30*(8), 559–564.

https://doi.org/10.1579/0044-7447-30.8.559

O'Reilly, C. M., Rowley, R. J., Schneider, P., Lenters, J. D., Mcintyre, P. B., & Kraemer, B. M.

(2015). Rapid and highly variable warming of lake surface waters around the globe.

Geophysical Research Letters, *42*(24), 1–9. https://doi.org/10.1002/2015GL066235

Oberlercher, T. M., & Wanzenböck, J. (2016). Impact of electric fishing on egg survival of

whitefish, Coregonus lavaretus. *Fisheries Management and Ecology*, *23*(6), 540–547.

https://doi.org/10.1111/fme.12197

Ohlberger, J., Mehner, T., Staaks, G., & Hölker, F. (2008). Is ecological segregation in a pair of

sympatric coregonines supported by divergent feeding efficiencies? *Canadian Journal of*

Fisheries and Aquatic Sciences, *65*(10), 2105–2113. https://doi.org/10.1139/F08-120

Ohlberger, J., Staaks, G., & Hölker, F. (2007). Effects of temperature, swimming speed and body

mass on standard and active metabolic rate in vendace (Coregonus albula). *Journal of*

Comparative Physiology B, *177*(8), 905–916. https://doi.org/10.1139/F08-120

Oomen, R. A., & Hutchings, J. A. (2015). Variation in spawning time promotes genetic

variability in population responses to environmental change in a marine fish. *Conservation*

Physiology, *3*(1), 1–12. https://doi.org/10.1093/conphys/cov027

Pariseau, R., Dumont, P., & Migneault, J.-G. (1983). Découverte, dans le sud-ouest du Québec,

d'une population de cisco de lac, Coregonus artedii, frayant au printemps. *Canadian*

Journal of Zoology, *61*(10), 2365–2368. https://doi.org/10.1139/z83-314

Paufve, M. R. (2019). *Diversity in spawning habitat across Great Lakes Cisco populations*

[Master's thesis]. Cornell University.

Pauly, D., & Pullin, R. S. V. (1988). Hatching time in spherical, pelagic, marine fish eggs in

response to temperature and egg size. *Environmental Biology of Fishes*, *22*(4), 261–271.

https://doi.org/10.1007/BF00004892

Pelletier, F., Garant, D., & Hendry, A. P. (2009). Eco-evolutionary dynamics. In *Philosophical*

Transactions of the Royal Society B: Biological Sciences (Vol. 364, Issue 1523, pp. 1483–

1489). The Royal Society. https://doi.org/10.1098/rstb.2009.0027

Pepin, P. (1991). Effect of temperature and size on development, mortality, and survival rates of

the pelagic early life history stages of marine fish. *Canadian Journal of Fisheries and*

Aquatic Sciences, *48*(3), 503–518. https://doi.org/10.1139/f91-065

Pritchard, A. L. (1931). Spawning habits and fry of the cisco (Leucichthys artedi) in Lake

Ontario. *Contributions to Canadian Biology and Fisheries*, *6*(1), 225–240.

https://doi.org/10.1139/f31-009

R Core Team. (2020). *R: A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.r-project.org/

Reist, J. D., Wrona, F. J., Prowse, T. D., Power, M., Dempson, J. B., Beamish, R. J., King, J. R.,

Carmichael, T. J., & Sawatzky, C. D. (2006). General effects of climate change on Arctic

fishes and fish populations. *AMBIO: A Journal of the Human Environment*, *35*(7), 370–380.

https://doi.org/10.1579/0044-7447(2006)35[370:GEOCCO]2.0.CO;2

Schindler, D. E., Hilborn, R., Chasco, B., Boatright, C. P., Quinn, T. P., Rogers, L. A., &

Webster, M. S. (2010). Population diversity and the portfolio effect in an exploited species.

Nature, *465*(7298), 609–612. https://doi.org/10.1038/nature09060

Schindler, D. W., Beaty, K. G., Fee, E. J., Cruikshank, D. R., DeBruyn, E. R., Findlay, D. L.,

Linsey, G. A., Shearer, J. A., Stainton, M. P., & Turner, M. A. (1990). Effects of Climatic

Warming on Lakes of the Central Boreal Forest. *Science*, *250*(4983), 967–970.

https://doi.org/10.1126/science.250.4983.967

Schmitt, J. D., Vandergoot, C. S., O'Malley, B. P., & Kraus, R. T. (2020). Does Lake Erie still

have sufficient oxythermal habitat for cisco Coregonus artedi? *Journal of Great Lakes*

Research, *46*(2), 330–338. https://doi.org/https://doi.org/10.1016/j.jglr.2020.01.019

Schultz, E. T., Conover, D. O., & Ehtisham, A. (1998). The dead of winter: size-dependent

variation and genetic differences in seasonal mortality among Atlantic silverside

(Atherinidae: \textit{Menidia menidia}) from different latitudes. *Canadian Journal of*

Fisheries and Aquatic Sciences, *55*(5), 1149–1157. https://doi.org/10.1139/cjfas-55-5-1149

Schultz, E. T., Reynolds, K. E., & Conover, D. O. (1996). Countergradient Variation in Growth

Among Newly Hatched Fundulus Heteroclitus: Geographic Differences Revealed by

Common-Environment Experiments. *Functional Ecology*, *10*(3), 366.

https://doi.org/10.2307/2390285

Schulz, M., & Freyhof, J. (2003). Coregonus fontanae, a new spring-spawning cisco from Lake

Stechlin, northern Germany (Salmoniformes: Coregonidae). *Ichthyological Exploration of*

Freshwaters, *14*(3), 209–216.

Schulz, M., Freyhof, J., Saint-Laurent, R., Østbye, K., Mehner, T., Bernatchez, L.,

Saint☐Laurent, R., Østbye, K., Mehner, T., & Bernatchez, L. (2006). Evidence for

independent origin of two spring☐spawning ciscoes (Salmoniformes: Coregonidae) in

Germany. *Journal of Fish Biology*, *68*(A), 119–135. https://doi.org/10.1111/j.0022-

1112.2006.01039.x

Scott, W. B., & Crossman, E. J. (1973). *Freshwater Fishes of Canada; Bulletin 184*.

Sharma, S., Blagrave, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R.,

Straile, D., Weyhenmeyer, G. A., & Winslow, L. A. (2019). Widespread loss of lake ice

around the Northern Hemisphere in a warming world. *Nature Climate Change*, *9*(3), 227.

https://doi.org/10.1038/s41558-018-0393-5

Sharma, S., Meyer, M. F., Culpepper, J., Yang, X., Hampton, S. E., Berger, S. A., Brousil, M. R.,

Fradkin, S. C., Higgins, S. N., & Jankowski, K. J. (2020). Integrating perspectives to

understand lake ice dynamics in a changing world. *Journal of Geophysical Research:*

Biogeosciences, *125*(8), 1–18. https://doi.org/10.1029/2020JG005799

Shatwell, T., Thiery, W., & Kirillin, G. (2019). Future projections of temperature and mixing

regime of European temperate lakes. *Hydrology and Earth System Sciences*, *23*(3), 1533–

1551. https://doi.org/10.5194/hess-23-1533-2019

Sipponen, M., Muje, K., Marjomäki, T. J., Valkeajärvi, P., & Karjalainen, J. (2006). Interlocked

use of inland fish resources: a new management strategy under private property rights.

Fisheries Management and Ecology, *13*(5), 299–307. https://doi.org/10.1111/j.1365-

2400.2006.00506.x

Stockwell, J. D., Ebener, M. P., Black, J. A., Gorman, O. T., Hrabik, T. R., Kinnunen, R. E.,

Mattes, W. P., Oyadomari, J. K., Schram, S. T., Schreiner, D. R., Seider, M. J., Sitar, S. P.,

& Yule, D. L. (2009). A Synthesis of Cisco Recovery in Lake Superior: Implications for

Native Fish Rehabilitation in the Laurentian Great Lakes. *North American Journal of*

Fisheries Management, *29*(3), 626–652. https://doi.org/10.1577/M08-002.1

Urpanen, O., Huuskonen, H., Marjomäki, T. J., & Karjalainen, J. (2005). Growth and size-

selective mortality of vendace (Coregonus albula (L.)) and whitefish (C. lavaretus L.)

larvae. *Boreal Environment Research*, *10*(3), 225–238.

van Damme, C. J. G., Dickey-Collas, M., Rijnsdorp, A. D., & Kjesbu, O. S. (2009). Fecundity,

atresia, and spawning strategies of Atlantic herring (Clupea harengus). *Canadian Journal of*

Fisheries & Aquatic Sciences, *66*(12). https://doi.org/10.1139/F09-153

Voeten, C. C. (2020). *buildmer: Stepwise Elimination and Term Reordering for Mixed-Effects*

Regression. https://cran.r-project.org/package=buildmer

Vonlanthen, P., Bittner, D., Hudson, A. G., Young, K. A., Müller, R., Lundsgaard-Hansen, B.,

Roy, D., Di Piazza, S., Largiader, C. R., Seehausen, O., Largiadèr, C. R., & Seehausen, O.

(2012). Eutrophication causes speciation reversal in whitefish adaptive radiations. *Nature*,

482(7385), 357–362. https://doi.org/10.1038/nature10824

Vonlanthen, P., Roy, D., Hudson, A. G., Largiadèr, C. R., Bittner, D., & Seehausen, O. (2009).

Divergence along a steep ecological gradient in lake whitefish (Coregonus sp.). *Journal of*

Evolutionary Biology, *22*(3), 498–514. https://doi.org/10.1111/j.1420-9101.2008.01670.x

Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P.,

Glidden, S., Bunn, S. E., Sullivan, C. A., & Liermann, C. R. (2010). Global threats to

human water security and river biodiversity. *Nature*, *467*(7315), 555–561.

https://doi.org/10.1038/nature09440

Wedekind, C., Evanno, G., Urbach, D., Jacob, A., & Müller, R. (2008). 'Good-genes' and

'compatible-genes' effects in an Alpine whitefish and the information content of breeding

tubercles over the course of the spawning season. *Genetica*, *132*(2), 199–208.

https://doi.org/10.1007/s10709-007-9164-3

White, C. R., Alton, L. A., & Frappell, P. B. (2012). Metabolic cold adaptation in fishes occurs

at the level of whole animal, mitochondria and enzyme. *Proceedings of the Royal Society B:*

Biological Sciences, *279*(1734), 1740–1747. https://doi.org/10.1098/rspb.2011.2060

Wilder, A. P., Palumbi, S. R., Conover, D. O., & Therkildsen, N. O. (2020). Footprints of local

adaptation span hundreds of linked genes in the Atlantic silverside genome. *Evolution*

Letters, *4*(5), 430–443. https://doi.org/doi.org/10.1002/evl3.189

Winslow, L. A., Read, J. S., Hansen, G. J. A., Rose, K. C., & Robertson, D. M. (2017).

Seasonality of change: Summer warming rates do not fully represent effects of climate

change on lake temperatures. *Limnology and Oceanography*, *62*(5), 2168–2178.

https://doi.org/10.1002/lno.10557

Woolsey, E. S., Keith, S. A., Byrne, M., Schmidt-Roach, S., & Baird, A. H. (2015). Latitudinal

variation in thermal tolerance thresholds of early life stages of corals. *Coral Reefs*, *34*(2),

471–478. https://doi.org/10.1007/s00338-014-1253-z

Woolway, R. I., Dokulil, M. T., Marszelewski, W., Schmid, M., Bouffard, D., & Merchant, C. J.

(2017). Warming of Central European lakes and their response to the 1980s climate regime

shift. *Climatic Change*, *142*(3–4), 505–520. https://doi.org/10.1007/s10584-017-1966-4

Woolway, R. I., Kraemer, B. M., Lenters, J. D., Merchant, C. J., O'Reilly, C. M., & Sharma, S.

(2020). Global lake responses to climate change. *Nature Reviews Earth & Environment*, 1–

16. https://doi.org/10.1038/s43017-020-0067-5

Wright, P. J., & Trippel, E. A. (2009). Fishery☐induced demographic changes in the timing of

spawning: consequences for reproductive success. *Fish and Fisheries*, *10*(3), 283–304.

https://doi.org/10.1111/j.1467-2979.2008.00322.x

Yamahira, K., & Conover, D. O. (2002). Intra- vs . Interspecific Latitudinal Variation in

Growth☐: Adaptation to Temperature or Seasonality? *Ecology*, *83*(5), 1252–1262.

https://doi.org/10.1890/0012-9658(2002)083[1252:IVILVI]2.0.CO;2

Yule, D. L., Moore, S. A., Ebener, M. P., Claramunt, R. M., Pratt, T. C., Salawater, L. L., &

Connerton, M. J. (2013). Morphometric variation among spawning cisco aggregations in the

Laurentian Great Lakes: are historic forms still present. *Advances in Limnology*, *64*, 119–

132. https://doi.org/10.1127/1612-166X/2013/0064-0022

Zimmerman, M. S., & Krueger, C. C. (2009). An Ecosystem Perspective on Re-establishing

Native Deepwater Fishes in the Laurentian Great Lakes. *North American Journal of*

Fisheries Management, *29*(5), 1352–1371. https://doi.org/10.1577/M08-194.1

TABLES:

Table 1. Mean (SD) water temperatures during embryo incubations at the University of Vermont (UVM) and University of Jyväskylä (JYU).

Laboratory	Incubation Temperature Treatment (°C)			
	2.0 Coldest	4.5 Cold	7.0 Warm	9.0 Warmest
UVM	2.0 (0.5)	4.4 (0.2)	6.9 (0.2)	8.9 (0.3)
JYU	2.2 (1.5)	4.0 (0.7)	6.9 (0.5)	8.0 (0.6)

Table 2. Mean (SD) total length (TL) and fresh mass (FM) of the female and males from Lake Southern Konnevesi (LK-Vendace (*Coregonus albula*) and LK-Whitefish (*C. lavaretus*)), Lake Superior (LS-Cisco (*C. artedi*)), and Lake Ontario (LO-Cisco).

Sex	LK-Vendace		LK-Whitefish		LS-Cisco		LO-Cisco	
	TL (mm)	FM (g)	TL (mm)	FM (g)	TL (mm)	FM (g)	TL (mm)	FM (g)
Female	144.67 (16.51)	18.36 (5.95)	256.57 (11.63)	117.00 (19.16)	428.92 (44.40)	676.02 (181.51)	380.33 (24.18)	567.59 (122.89)
Male	140.83 (9.22)	13.85 (2.27)	285.75 (40.86)	171.34 (87.22)	400.25 (34.35)	523.82 (134.65)	366.56 (25.30)	443.29 (103.16)

Table 3. Mean (SD) egg diameter of females with the number of eggs measured (N) from Lake Southern Konnevesi (LK-Vendace (*Coregonus albula*) and LK-Whitefish (*C. lavaretus*)), Lake Superior (LS-Cisco (*C. artedi*)), and Lake Ontario (LO-Cisco).

Population	Egg diameter (mm)	N
LK-Vendace	1.58 (0.11)	273
LK-Whitefish	2.13 (0.12)	70
LS-Cisco	2.14 (0.12)	140
LO-Cisco	2.30 (0.08)	240

Table 4. Likelihood ratio test output for each model selected for embryo survival and incubation period (number of days post-fertilization (DPF) and accumulated degree days (°C; ADD)) from Lakes Superior and Ontario cisco (*Coregonus artedi*) and Lake Southern Konnevesi vendace (*C. albula*) and European whitefish (*C. lavaretus*). t indicates temperature, pop indicates population, and sp indicates species. The full model that was selected is bolded for each trait and species.

Trait	Species	Model	Effect Tested	df	χ^2	p-value
Embryo Survival	Cisco	**t + pop + t:pop + female:male + female**				
		pop + female:male + female	t	3	443.54	< 0.00
		t + female:male + female	pop	1	600.61	< 0.00
		t + pop + female:male + female	t:pop	3	198.56	< 0.00
		t + pop + t:pop + female	female:male	1	181.47	< 0.00
		t + pop + t:pop + female:male	female	1	23.36	< 0.00
	Vendace & European Whitefish	**t + sp + t:sp + female:male + female**				
		sp + female:male + female	t	3	223.54	< 0.00
		t + female:male + female	sp	1	993.43	< 0.00
		t + sp + female:male + female	t:sp	3	52.94	< 0.00
		t + sp + t:sp + female	female:male	1	1042.9	< 0.00
		t + sp + t:sp + female:male	female	1	1015.8	< 0.00
Incubation Period (DPF)	Cisco	**t + pop + t:pop + female:male + female + male**				
		pop + female:male + female + male	t	3	31,183.13	< 0.00
		t + female:male + female + male	pop	1	3,651.86	< 0.00
		t + pop + female:male + female + male	t:pop	3	315.04	< 0.00
		t + pop + t:pop + female + male	female:male	1	58.62	< 0.00
		t + pop + t:pop + female:male + male	female	1	65.04	< 0.00
		t + pop + t:pop + female:male + female	male	1	14.22	<0.00
	Vendace & European Whitefish	**t + sp + t:sp + female:male + female + male**				
		sp + female:male + female + male	t	3	7,178.37	< 0.00
		t + female:male + female + male	sp	1	904.95	< 0.00
		t + sp + female:male + female + male	t:sp	3	353.67	< 0.00
		t + sp + t:sp + female + male	female:male	1	13.55	< 0.00
		t + sp + t:sp + female:male + male	female	1	36.07	< 0.00
		t + sp + t:sp + female:male + female	male	1	4.77	0.029
Incubation Period (ADD)	Cisco	**t + pop + t:pop + female:male + female + male**				
		pop + female:male + female + male	t	3	15,519.48	< 0.00
		t + female:male + female + male	pop	1	3,586.18	< 0.00
		t + pop + female:male + female + male	t:pop	3	137.36	< 0.00
		t + pop + t:pop + female + male	female:male	1	54.58	< 0.00
		t + pop + t:pop + female:male + male	female	1	63.20	< 0.00
		t + pop + t:pop + female:male + female	male	1	15.34	< 0.00
	Vendace & European	**t + sp + t:sp + female:male + female + male**				
		sp + female:male + female + male	t	3	3,571.72	< 0.00

	Whitefish	t + female:male + female + male	sp	1	763.19	< 0.00
		t + sp + female:male + female + male	t:sp	3	466.23	< 0.00
		t + sp + t:sp + female + male	female:male	1	12.70	< 0.00
		t + sp + t:sp + female:male + male	female	1	36.79	< 0.00
		t + sp + t:sp + female:male + female	male	1	4.33	0.038

Table 5. Likelihood ratio test output for each model selected for length-at-hatch (mm) and yolk-sac volume (mm^3) from Lakes Superior and Ontario cisco (*Coregonus artedi*), Lake Southern Konnevesi vendace (*C. albula*), and Lake Southern Konnevesi European whitefish (*C. lavaretus*). t indicates temperature and pop indicates population. The full model that was selected is bolded for each trait and species.

Trait	Species	Model	Effect Tested	df	χ^2	p-value
Length-at-Hatch	Cisco	**t + pop + t:pop + female**				
		pop + female	t	3	886.79	< 0.001
		t + female	pop	1	628.15	< 0.001
		t + pop + female	t:pop	3	19.03	< 0.001
		t + pop + t:pop	female	1	161.40	< 0.001
	Vendace & European Whitefish	**t + sp + t:sp + female:male + female**				
		sp + female:male + female	t	3	467.20	< 0.001
		t + female:male + female	sp	1	2091.33	< 0.001
		t + sp + female:male + female	t:sp	3	14.49	0.002
		t + sp + t:sp + female	female:male	1	13.83	< 0.001
		t + sp + t:sp + female:male	female	1	48.81	< 0.001
Yolk-sac Volume	Cisco	**t + pop + t:pop + female**				
		pop + female	t	3	1,163.87	< 0.001
		t + female	pop	1	246.56	< 0.001
		t + pop + female	t:pop	3	99.25	< 0.001
		t + pop + t:pop	female	1	410.42	< 0.001
	Vendace & European Whitefish	**t + sp + t:sp + female + male**				
		sp + family + female	t	3	980.02	< 0.001
		t + family + female	sp	1	805.34	< 0.001
		t + sp + family + female	t:sp	3	107.58	< 0.001
		t + sp + t:sp + female	family	1	33.33	< 0.001
		t + sp + t:sp + family	female	1	6.96	0.008

Table 6. Phenotypic variation component correlation directions from increasing incubation temperature for embryo survival (%), incubation period (number of days post-fertilization; DPF), incubation period (accumulated degree-days; ADD), length-at-hatch (mm), and yolk-sac volume (mm^3) from Lake Southern Konnevesi vendace (LK-Vendace (*Coregonus albula*)), Lake Superior cisco (LS-Cisco (*C. artedi*)), and Lake Ontario cisco (LO-Cisco). - indicates a negative correlation, + indicates a positive correlation, and 0 indicates no correlation.

| Trait | Study Group | Correlation Direction | | | |
		Dam	Sire	Dam:Sire	Error
Embryo Survival	LK-Vendace	-	+	0	+
	LS-Cisco	-	0	0	0
	LO-Cisco	0	+	0	+
Incubation Period (DPF)	LK-Vendace	0	0	+	0
	LS-Cisco	-	0	0	0
	LO-Cisco	-	0	0	+
Incubation Period (ADD)	LK-Vendace	0	-	+	0
	LS-Cisco	-	0	0	0
	LO-Cisco	-	-	0	+
Length-at-Hatch	LK-Vendace	0	-	+	0
	LS-Cisco	-	0	0	0
	LO-Cisco	-	0	+	0
Yolk-sac Volume	LK-Vendace	+	0	+	-
	LS-Cisco	-	+	-	+
	LO-Cisco	0	0	0	0

FIGURES:

Figure 1. Map showing the location of each lake (LS = Lake Superior; LO = Lake Ontario; LK = Lake Southern Konnevesi) sampled in North America (A) and Europe (B).

Figure 2. Winter water temperatures from Lake Southern Konnevesi, Lake Superior, and Lake Ontario. Lake Southern Konnevesi and Lake Superior data were recorded using *in-situ* sensors on the lakebed (10-m deep). Lake Ontario data was recorded using remote sensing sea surface temperatures. X indicates *ca.* spawning and O indicates *ca.* hatching.

Figure 3. Mean embryo survival (%) and incubation period (number of days post-fertilization (DPF) and accumulated degree days (°C; ADD)) at each incubation temperature (°C; left) and standardized temperature responses within each study group (%; right) from Lake Southern Konnevesi (LK-Vendace (*Coregonus albula*) and LK-Whitefish (*C. lavaretus*)), Lake Superior (LS-Cisco (*C. artedi*)), and Lake Ontario (LO-Cisco). Error bars indicate standard error.

Figure 4. Mean percent of total phenotypic variation across incubation temperatures for embryo survival, incubation period (number of days post-fertilization (DPF) and accumulated degree days (°C; ADD)), length-at-hatch (mm), and yolk-sac volume (mm^3) from Lake Southern Konnevesi vendace (LK-Vendace (*Coregonus albula*), Lake Superior cisco (LS-Cisco (*C. artedi*)), and Lake Ontario cisco (LO-Cisco). Error bars indicate standard error.

Figure 5. Mean length-at-hatch (mm; LAH) and yolk-sac volume (mm^3; YSV) at each incubation temperature (°C; left) and standardized temperature responses within each study group (%; right) from Lake Southern Konnevesi (LK-Vendace (*Coregonus albula*) and LK-Whitefish (*C. lavaretus*)), Lake Superior (LS-Cisco (*C. artedi*)), and Lake Ontario (LO-Cisco). Error bars indicate standard error.

Figure 6. Theoretical winter incubation periods and responses of embryo demographics under normal (2.0°C; blue) and hypothetical warm (5.0°C; orange) winter thermal regimes. The shaded regions indicate spawning periods (left) and hatching periods (right) that may occur between 4-5°C. The 2.0°C temperature regime is water temperature data collected from Lake Superior at 10-m depth in 2018.