The cohesin regulator Stag1 promotes cell plasticity through heterochromatin regulation.

Dubravka Pezic[1], Sam Weeks[1], Wazeer Varsally[1], Pooran S. Dewari[2], Steven Pollard[2], Miguel R. Branco[3], Suzana Hadjur[1]*

1 Research Department of Cancer Biology, Cancer Institute, University College London, 72 Huntley Street, London, United Kingdom

2 MRC Centre for Regenerative Medicine, University of Edinburgh, Edinburgh, United Kingdom

3 Blizard Institute, Barts and The London School of Medicine and Dentistry, QMUL, London, United Kingdom

* Correspondence: s.hadjur@ucl.ac.uk

SUMMARY

Fundamental biological processes such as embryo development and stem cell control rely on cellular plasticity. We present a role for the cohesin regulator, Stag1 in cellular plasticity control via heterochromatin regulation. Stag1 localises to heterochromatin domains and repetitive sequences in embryonic stem (ES) cells and contains intrinsically disordered regions in its divergent terminal ends which promote heterochromatin compaction. ES cells express Stag1 protein isoforms lacking the disordered ends and fluctuations in isoform abundance skews the cell state continuum towards increased differentiation or reprogramming. The role for Stag1 in heterochromatin condensates and nucleolar function is dependent on its unique N-terminus. $Stag1^{N\Delta}$ ESCs have decompacted chromatin and reprogram towards totipotency, exhibiting MERVL de-repression, reduced nucleolar transcription and decreased translation. Our results move beyond protein-coding gene regulation via chromatin loops into a new role for Stag1 in heterochromatin and nucleolar function and offer fresh perspectives on its contribution to cell identity and disease.

KEY WORDS

Cellular plasticity, heterochromatin, repetitive elements, nucleolus, Stag1, translation

INTRODUCTION

Cellular populations consist of mixtures of cells across a continuum of states and such heterogeneity underlies responsiveness to changing conditions. Fundamental biological processes including embryo development, stem cell control and cancer rely on conversion between states. While transcriptional states are ever-increasingly well described (Wagner et al., 2016), the mechanisms that control the stability of a given state are less well understood. The chromatin landscape with its inherent dynamics (Cho et al., 2018; Finn et al., 2019; Nozaki et al., 2017; Ricci et al., 2015) and complex 3-dimensional (3D) organization (Bonev et al., 2017; Cardozo Gizzi et al., 2019; Dekker and Mirny, 2016; Nagano et al., 2017; Schlesinger and Meshorer, 2019) can act as a mechanism to support transcriptional heterogeneity (Mateo et al., 2019; Rodriguez et al., 2019) and the chromatin proteins that regulate this landscape may play key roles in cell plasticity (Meshorer et al., 2006).

Genomes are partitioned into distinct functional domains within the nucleus (Bickmore and van Steensel, 2013; Bonev and Cavalli, 2016). A prominent feature of nuclear organization is the compact heterochromatin that accumulates at the periphery of the nucleus, around the nucleolus and at distinct foci within the nucleoplasm (Guelen et al., 2008; Németh et al., 2010; Padeken and Heun, 2014; Quinodoz et al., 2018). Heterochromatin is formed at repetitive sequences, is tightly associated with repressive histone modifications like methylation of histone H3 at lysine 9 (H3K9me2,3), and a specific set of proteins, including Heterochromatin Protein 1 (HP1) that together condense chromatin to maintain repression (Allshire and Madhani, 2018). Further, sequestration of condensed chromatin into dynamic phase-separated condensates (Larson et al., 2017; Strom et al., 2017) contributes to heterochromatin-mediated silencing. Heterochromatin organization plays a key role in cell identity and is rapidly remodelled during ES cell differentiation and embryo development. Decompaction of heterochromatin and subsequent de-repression of repetitive elements drives reprogramming towards totipotent embryos, while progressive compaction is associated with terminal differentiation (Ahmed et al., 2010; Borsos and Torres-Padilla, 2016; Martin et al., 2006; Meshorer et al., 2006; Novo et al., 2016).

Cohesin is a ubiquitously expressed, multi-subunit protein complex that has
fundamental roles in cell biology including sister chromosome cohesion, 3D chromatin
topology and regulation of cell identity (Cuartero et al., 2018; Horsfield et al., 2007; Kline
et al., 2018; Leiserson et al., 2015; Romero-Pérez et al., 2019; Viny et al., 2019). Much
of our understanding of how cohesin contributes to cell identity has been studied in the
context of its roles in protein-coding gene expression and 3D organization of interphase
chromatin structure (Hadjur et al., 2009; Kagey et al., 2010; Mishiro and Tsutsumi, 2009;
Misulovin et al., 2007; Parelho et al., 2008; Phillips-Cremins et al., 2013; Rao et al., 2014;
Vietri Rudan et al., 2015; Wendt et al., 2008). Indeed, loss of cohesin and its regulators
results in a dramatic loss of chromatin topology at the level of Topologically Associated
Domains (TAD) and chromatin loops, with only modest changes to gene expression
(Haarhuis et al., 2017; Rao et al., 2017; Schwarzer et al., 2017; Seitan et al., 2013;
Sofueva et al., 2013; Wutz et al., 2017; Zuin et al., 2014). This suggests that cohesin's
roles in development and disease extend beyond gene expression regulation and
highlight the need to re-evaluate how cohesin regulators shape the structure and function
of the genome.

The association of cohesin with chromosomes is tightly controlled by several
regulators, including the Stromalin Antigen protein (known as Stag or SA), which has
been widely implicated in cell identity regulation and disease development (Cuadrado et
al., 2019; Lehalle et al., 2017; Leiserson et al., 2015; Soardi et al., 2017; Viny et al., 2019;
Yuan et al., 2019). Stag proteins interact with the Rad21 subunit of cohesin and mediate
its association with DNA and CTCF (Hara et al., 2014; Li et al., 2020; Orgil et al., 2015;
Xiao et al., 2011). Mammalian cells have three Stag paralogs, Stag1, 2 and 3. These
show >90% conservation of sequence in their central domain yet perform distinct
functions (Canudas and Smith, 2009; Kojic et al., 2018; Remeseiro et al., 2012a; Winters
et al., 2014). It is likely that the divergent N- and C-terminal regions provide functional
specificity. For example, the N-terminus of Stag1 contains a unique AT-hook (Bisht et al.,
2013) which is required for its preferential participation in telomere cohesion (Canudas
and Smith, 2009). The underlying mechanisms by which Stag proteins and their divergent
ends influence cell identity are still largely unknown.

Here we report that Stag1 is the dominant paralogue in ES cells and supports

pluripotency by regulating heterochromatin organization. We discover that ES cells

regulate the level of Stag1 protein and the proportion of its divergent N and C-terminal

ends, which contain disordered regions. This naturally occurring Stag1 protein

heterogeneity supports a continuum of functionally distinct cellular states within the

population. Changing the balance in the levels of Stag1 isoforms leads to conversion

between cell states, with the loss of the N-terminus favoring a reprogrammed, totipotent

state and decompaction of heterochromatin condensates and the loss of the C-terminus

priming cells towards exit from pluripotency through gene expression deregulation. These

results define specialised, non-redundant roles for the divergent ends. Mechanistically,

the N-terminus of Stag1 represses reprogramming to totipotent 2-cell-like (2C-L) cells by

maintaining nucleolar structure and function. We uncover Nucleolin and Trim28 as direct

interactors of Stag1 and show that cells selectively expressing Stag1 isoforms lacking the

N-terminal AT-hook domain exhibit reduced nascent nucleolar transcription and a

decrease in global translation. Our results take us beyond protein coding gene regulation

via chromatin loops into a new role for Stag1 in the regulation of heterochromatin and

nucleolar structure and function. Importantly, by identifying changes to translation control

upon Stag1 loss in stem cells, we open a new perspective by which Stag proteins and

cohesin regulation can impact cell identity and disease.

RESULTS

A functional change in cohesin regulation in cells of different potential.

We analyzed the expression levels of cohesin regulators in embryonic stem cells (ES) at different stages of pluripotency. During the transition between naïve (2i) and primed (EpiLC) pluripotency *in vitro*, levels of the core cohesin subunits Smc1 and Smc3 do not change, while Stag1 becomes downregulated and Stag2 becomes upregulated (Figures 1A, S1A, B). This was confirmed at the protein level, where we observe a 2-3-fold higher level of chromatin-associated Stag1 compared to Stag2 in naïve ES cells, while Stag2 levels are 5-10-fold higher in EpiLCs (Figures 1B, S1C). These results, together with similar observations (Cuadrado et al., 2019), identify Stag1 as the dominant paralog in naïve ES cells and suggest that a switch between Stag1 and Stag2 may represent a functionally relevant change in cohesin regulation at different stages of pluripotency.

Stag1 supports the naïve pluripotent state.

To investigate the functional importance of Stag1 in the regulation of pluripotency, we established a Stag1 RNA knockdown (KD, 'siSA1-SP', Methods) strategy using siRNAs. This resulted in a significant reduction of Stag1 at the mRNA and protein levels (4-5-fold, 8-10-fold, respectively), in both serum-grown (FCS) and naïve (2i) ES cells without affecting the cell cycle (Figures 1C, S1D, E). Using Nanog as a marker of naïve pluripotency, we observed a significant downregulation of Nanog mRNA and protein levels within 24hrs of Stag1 KD in all ES populations (Figures 1D, E, S1F), suggesting that Stag1 may be functionally required for pluripotency. Indeed, a global analysis of the ES transcriptome upon siRNA-mediated Stag1 KD revealed that 375 genes were up- and 205 genes were down-regulated by at least 2-fold (Figure 1F). Among the downregulated group were several genes known to have key roles in the maintenance of pluripotency, including Nanog, Tbx3, Esrrb, Klf2, Klf4, Prdm14, Tfcp2l1, Lefty2. Notably, we did not detect a change in expression of Oct4 or Sox2, and Zfp42 was minimally affected. Moreover, we also observed both an upregulation of genes associated with exit from the pluripotent state (Dppa3, Fgf5) as well as differentiation-specific genes such as Pou3f1

(Oct6) and Sox11 (Figure 1F). Single-gene analysis revealed consistent, albeit low fold-
change trends across our biological replicates, thus we used Gene Set Enrichment
Analysis (GSEA) (Mootha et al., 2003; Subramanian et al., 2005) to detect modest but
coordinate changes in the expression of groups of functionally related genes. This
revealed a robust gene signature of exit from pluripotency and enrichment for genes
associated with primed pluripotency upon Stag1 KD across all biological replicates
(Figures 1G, S1G).

The loss of the naïve transcriptional programme upon Stag1 KD suggests that ES
cells may require Stag1 for the maintenance of self-renewal. To test this, we plated cells
in self-renewal conditions at clonal density and determined the proportion of
undifferentiated cells upon Stag1 KD by measuring the area occupied by the colonies
with high alkaline phosphatase activity (AP+). In scrambled siRNA-treated controls, 52%
of plated cells retain their naïve state, identified by AP+ colonies (Figure S1H). However,
upon Stag1 KD, both the proportion of AP+ colonies and the area they occupy decreased
by 20%, indicating that ES cells have a reduced self-renewal ability in the absence of
Stag1 (Figures 1H, S1H). As independent validation of these results, we used
CRISPR/Cas9 to knock-in an mNeonGreen-FKBP12^{F36V} tag (Nabet et al., 2018) at the C-
terminus of both alleles of the endogenous Stag1 locus (SA1$^{NG\_FKBP}$) in ES cells (Figures
1I, S1I-K). Upon dTAG addition, Stag1 protein is robustly degraded in SA1$^{NG\_FKBP}$ ES
clones (Figure 1I, S1K). As was observed with siRNA treatment, dTAG-mediated
degradation of Stag1 led to a 50% reduction of self-renewal potential (Figure 1J).
Together, our results are consistent with a requirement for Stag1 in the control of naïve
pluripotency and provide an opportunity to discover the mechanisms of Stag1 actions.

STAG1 localizes to AT-rich heterochromatin.

To begin to understand how STAG1 contributes to pluripotency, we investigated the
subcellular localization of endogenous STAG1. Live cell imaging of Hoechst-labelled
Stag1$^{NG\_FKBP}$ ES cells revealed the expected and predominant localisation of STAG1 in

the nucleus (Figures 2A, B). Interestingly, STAG1 was not uniformly distributed within the

nucleoplasm. In addition to a dispersed nucleoplasmic localisation pattern, we observed

STAG1 colocalization with Hoechst-dense regions. These included colocalization at large

Hoechst-dense foci (Figure 2A, top cell), within the interior of the nucleolus (Figure 2A,

top cell) and at the periphery of the nucleolus (Figure 2A, all cells). The mean intensity of

STAG1 (as measured by mNeonGreen signal) was significantly enriched within Hoechst-

dense foci compared to the whole nucleus, and the signal was sensitive to treatment with

dTAG (Figure 2B). We made similar observations of STAG1 localization at DAPI-dense

foci in cells expressing Dox-inducible GFP-tagged full-length Stag1 (SA1^{FL}-GFP) (Figure

S2A, B). We note that repressive heterochromatin domains are readily observed by

staining with AT-rich DNA dyes (ie. DAPI and Hoechst) and are organized around the

nucleolus, in discreet foci within the nucleoplasm, or tethered to the nuclear periphery

(Padeken and Heun, 2014; Quinodoz et al., 2018). Thus, the profile of STAG1 within ES

cells is consistent with its localization to AT-rich heterochromatin. Given the presence of

an AT-hook within STAG1 and the importance of heterochromatin regulation in

development, we investigated whether STAG1 may have a role in heterochromatin

structure.

STAG1 interacts with heterochromatin proteins and repetitive DNA.

Since STAG1 was localised to nuclear heterochromatin domains, we investigated

whether it was also interacting with heterochromatin proteins and bound to genomic

sequences known to form heterochromatin, such as repeats. Constitutive

heterochromatin is characterized by the binding of HP1α to H3K9me2/) and plays a

critical role in silencing of repetitive DNA elements (Allshire and Madhani, 2018) and

nuclear organization (Larson et al., 2017). The periphery of the nucleolus accumulates

marks of constitutive heterochromatin coincident with transcriptionally inactive rDNA

repeats.   Nucleolin is a major nucleolar protein which controls the organization of

nucleolar chromatin, rDNA transcription and ribosome assembly and plays important

roles during development and in ES cells (Kresoja-Rakic and Santoro, 2019; Percharde

et al., 2018). We observed nuclear colocalization between STAG1 and HP1α in dox-

induced SA1^{FL}-GFP cells around DAPI-dense foci (Figures S2A, B). Further, using
chromatin coImmunoprecipitation (coIP), we show that STAG1 interacts with both HP1α
and Nucleolin in ES cells (Figure 2C, S2C).

Previous studies have analysed STAG1 binding profiles in mouse ES cells and
have primarily focused on its association with protein-coding genes (Cuadrado et al.,
2019). A thorough investigation of STAG1 binding to repetitive sequences has not been
conducted. Thus, we re-analysed STAG1 chromatin immunoprecipitation followed by
sequencing (ChIP-seq) experiments to calculate the proportion of STAG1 peaks that
overlapped genes (based on promoter and exon features), repeats (within the Repeat
Masker annotation) introns and intergenic regions not already represented by repeats or
genes. Of the 18,600 STAG1 peaks identified, the majority (76%) are bound to genomic
elements that are distinct from protein-coding genes with a significant proportion of
binding sites at repetitive elements and intergenic regions (Figure 2D). Together with the
localizations observed by microscopy, this suggests that the role of STAG1 in ES cells
may extend beyond protein-coding gene regulation. While STAG1 binding is not enriched
compared to all genomic repeats, we asked whether specific repeat families might be
enriched for STAG1 binding above random expectation (Deniz et al., 2020). We found
several repeat families to be significantly enriched for STAG1 peaks, including those
within the DNA transposon and Retrotransposon classes, both known to form constitutive
heterochromatin. Specifically, STAG1 was enriched at SINE B2-Mm2, (previously shown
to be enriched at TAD borders (Dixon et al., 2012)) and B3 elements, LINE1 elements
(L1Tf, L1A), and several LTR families, two of which have been previously shown to be
associated with CTCF (LTR41, LTR55) (Schwalie et al., 2013) (Figures 2E and S2D). The
enrichment of Stag1 at non-coding and repetitive sequences has not been previously
described and points to a novel role for STAG1 in the regulation of repeats in ES cells.

Stag1 regulates nuclear organization of heterochromatin.

Since H3K9me3 is the defining histone modification of silent heterochromatin, we
assessed the impact of STAG1 loss and overexpression on H3K9me3 in ES cells. While
global levels of H3K9me3 were unchanged upon Stag1 KD (Figure S2E),

immunofluorescence (IF) of H3K9me3 revealed changes to heterochromatin

organization. H3K9me3 foci displayed a greater variation in volume compared to

scrambled control treated cells, suggesting that Stag1 is required for proper H3K9me3

compaction (Figures 2F, G). We detected similarly variable changes to global chromatin

accessibility, as measured by DNase I digestion. In four out of six experiments, ES cells

treated with Stag1 siRNAs revealed a tendency towards increased accessibility, whereas

two experiments showed increased compaction (Figure 2H, S2F). On the other hand,

Dox-inducible STAG1 expression led to a dramatic condensation of H3K9me3 into large

nuclear puncta compared to non-induced cells (Figures 2I, J), where STAG1 could also

be seen to be colocalized with H3K9me3 in the condensate. This phenotype led us to

investigate whether STAG1 may have characteristics found in other proteins known to

play a role in heterochromatin phase separation(Larson et al., 2017). We used the

PONDR tool (Obradovic et al., 2003) to assess potential intrinsically disordered regions

(IDR) (Banani et al., 2017) within STAG1. STAG1 has an overall PONDR score of 0.4397,

and both the N-and C-terminal divergent regions contain sequences with a high

propensity for intrinsic disorder (Figure 2K). Interestingly, these coincide with known

STAG1 domains, most notably the N-terminal AT-hook. Together our results uncover a

novel role for STAG1 in forming or maintaining heterochromatin structures in ES cells and

suggest that the terminal ends may play important roles therein.

Stag1 expression is highly regulated in ES cells.

STAG1 levels are highest in naïve 2i-grown and lower in FCS-grown ES cells, a culture

condition that supports a mix of naïve and primed cells (Figure S1B). This prompted us

to investigate whether STAG1 is regulated at the transcriptional level. To address this,

we employed a series of approaches to comprehensively characterize Stag1 mRNAs and

discovered widespread regulation of Stag1 transcription in ES cells. First, we used RACE

(Rapid Amplification of cDNA Ends) to characterize the starts and ends of Stag1 mRNAs.

Using 5' RACE, we uncovered four novel alternative transcription start sites (TSS) in ES

cells; one located 50kb upstream of the canonical Stag1 TSS (referred to as 'SATS', and

previously identified in (Feng et al., 2016)) (Figures 3A, D, S3A), one between canonical

exon 1 and exon 2 (referred to as alternative exon 1 or altex1) (Figure 3D), one at exon

6, and one at exon 7 (Figures 3B, D, S3A, B (for increased exposure). Interestingly, the

novel TSS located at exon 7 (e7) was preceded by a sequence located *in trans* to the

Stag1 gene, carrying simple repeats and transcription factor binding sites (Figure S3C).

While the frequency of this alternative TSS was significantly lower than the other TSSs,

it was identified in multiple RACE replicates, indicating it may have important functions in

a subset of the ES population. We also discovered widespread alternative splicing in the

5' region of Stag1, with particularly frequent skipping of exons 2 and 3 (e2/3Δ) and exon

5 (e5Δ) (Figures 3D, S3A). Using 3' RACE, we detected an early termination site in intron

25 and inclusion of an alternative exon 22 introducing an early STOP codon, as well as

several 3'UTRs (Figures 3C, D, S3D).

Next, PCR- and Sanger sequencing-based clonal screening confirmed that the

newly discovered 5' and 3' ends represent true Stag1 transcript ends, validated the

existence of the e2/3Δ and e5Δ isoforms, and uncovered an isoform lacking exon 31

(e31Δ) (Figures 3D, S3E). To determine the complete sequences of the Stag1 transcript

isoforms and to use a non-PCR-based approach, we performed long-read PacBio Iso-

seq from 2i ES RNA. This confirmed the diversity of the Stag1 5' and 3'UTRs, the e31Δ

isoform, multiple TSSs, including SATS, and early termination events, including in i22 and

i25 (Figure S3F). Importantly, these transcripts all had polyA tails, in support of their

protein-coding potential. Finally, we validated and quantified the newly discovered

splicing events by calculating the frequency (percentage spliced in (PSI)) of exon splicing

in our and published RNA-seq data using VAST-tools, a previously developed

computational method (Tapial et al., 2017) (Figure 3E, Table S2). Together, these results

point to a previously unappreciated diversity of endogenous Stag1 transcripts in ES cells,

highlighting the importance of Stag1 regulation in stem cell populations.

The chromatin landscape reflects Stag1 transcriptional regulation.

Visual inspection of the genome topology around the *Stag1* locus in our existing 2i ES

and neural stem (NS) cell Hi-C datasets (Barrington et al.) revealed that the *STAG1* gene

undergoes significant 3-D reorganization as cells differentiate (Figure 3F). First, the entire

STAG1 TAD switches from the active to the repressive compartment during

differentiation, in line with the decrease in Stag1 levels described above. Furthermore,

we observed several changes to the sub-TAD architecture which corresponded to the

newly discovered Stag1 TSSs and TTSs described above (Figure 3F, compare 'SA1

transcripts' track with Hi-C maps). We quantified these changes by designing site-specific

UMI-4C baits guided by the Hi-C topology. In naïve ES cells, the genomic region

containing the alternative SATS TSS has several Nanog binding sites and makes

numerous contacts with the *Stag1* gene body (Figure 3F, 'Nanog bait 1'). Meanwhile, in

NS cells, the SATS TSS is no longer active and becomes isolated away from the *STAG1*

gene body due to the reinforcement of the TAD border at the canonical promoter of *Stag1*

(Figure 3F, 'CTCF bait'). In addition, a cluster of low-occupancy CTCF binding sites within

the *Stag1* gene mark an ES-specific contact insulation point giving rise to a sub-TAD. We

note that the observed Stag1 early termination events align with these CTCF sites. These

results suggest that 3D chromatin topology may play a direct role in facilitating the

transcriptional output of *Stag1*.

Multiple Stag1 protein isoforms are expressed in ES cells.

Stag1 transcript diversity was intriguing because many of the events were either specific

to ES cells or enriched compared to MEFs and NSCs (Figure 3A, E, S3E) and further,

because the transcript variants are predicted to produce STAG1 protein isoforms with

distinct structural features and molecular weights (Figure 3D, G). For example, the

truncation of the N terminus (e2/3Δ, e5Δ, e6 TSS and e7 TSS), and thus loss of the AT

hook (amino acid 3-58), could impact STAG1 association with DNA. Meanwhile, C-

terminal truncated Stag1 isoforms (altex22, i25 end, e31Δ) could affect STAG1-cohesin

interactions. Interestingly, the evolutionarily conserved Stag-domain ('SCD', AA 296-381)

(Orgil et al., 2015), shown to play a role in CTCF interaction (Li et al., 2020), would be

retained in all the transcripts identified here. Importantly, these events would yield STAG1

isoforms lacking either the N- or C-terminal disordered regions and could thus impact the

ability of STAG1 to form condensate-like structures (Figure 3H).

Immunoprecipitation (IP) of endogenous STAG1 revealed multiple bands
corresponding to the predicted molecular weights for several protein isoforms and
identified by mass spectrometry to contain Stag1 peptides (Figure 3I and Table S3),
whose levels were reduced between naïve and primed cells (Figure S3G) and sensitive
to Stag1 KD, alongside the canonical, full-length isoform (Figure 3J). Treatment of
Stag1$^{NG\_FKBP}$ ES cells with dTAG followed by Western blotting of chromatin-associated
proteins with an antibody to the v5 tag further confirmed the sensitivity of the isoforms to
dTAG-mediated degradation, validating the presence of N-terminally truncated STAG1
protein variants (Figure 3K). Overall, our results indicate that complex transcriptional
regulation gives rise to multiple Stag1 transcripts and protein isoforms with distinct
regulatory regions and coding potential, the majority of which are expressed specifically
in naïve ESCs. Our discovery of such naturally occurring STAG1 isoforms highlights the
importance of STAG1 in ES cells and offers a unique opportunity to define the ES-specific
functions of the divergent N- and C-terminal ends of STAG1 in the context of pluripotency
gene expression and heterochromatin regulation.
320

Skewing the abundance of Stag1 isoforms promotes transitions between cell
states.
To study the functional consequences of STAG1 isoform expression changes on ES cells,
we took advantage of our detailed understanding of Stag1 transcript diversity to design
custom siRNA pools to selectively target, or retain, specific isoforms (Figure 4A).
Alongside the siRNAs from Figure 1 (SmartPool, SP), we designed siRNAs to specifically
target the SATS 5'UTR (esiSATS), the 5' end (siSA1-5p) or the 3' end (siSA1-3p) of Stag1
mRNA (see Methods). We anticipated that the KD panels would not completely abolish
all Stag1 transcript variants, but rather change the relative proportions, in effect skewing
the levels of the N- and C-terminal ends of Stag1. 3p siRNAs were expected to
downregulate full-length and N-term truncated isoforms and retain C-term truncated
isoforms. Meanwhile 5p siRNAs would specifically retain N-term truncated isoforms.
siRNAs to the 5p and 3p ends of Stag1 reduce full-length Stag1 mRNA and protein
with similar efficiency to SP KDs (Figures 4B, C, S4A), while esiSATS reduces Stag1 by

50-60%, indicating that the SATS TSS functions to enhance expression of Stag1

specifically in naïve ES cells. We confirmed that Stag1 isoform proportions were altered

upon siRNA treatment using RNA-seq, RACE and immunoprecipitation. RNA-seq reads

aligning to Stag1 in the different siRNA treatments were quantified to represent the

residual N-terminal, middle and C-terminal read proportions (Figure 4D). Residual reads

in the SP and 3p KDs aligned primarily to the N-terminus and were depleted from the C-

term. While the 5p KD had the least read retention in the N-terminus, supporting the

expectation that residual transcripts in the 5p KD have full C-terminal ends (Figure 4D).

These results were further supported by quantifying specific splicing events from the KD

RNA-seq datasets using VAST-tools (Figure S4B). In parallel, RACE was used to

validate changes to the proportions of Stag1 isoforms. 5' RACE in ES cells treated with

5p siRNA revealed downregulation of full-length Stag1 transcript while several N-terminal

truncated isoforms were upregulated compared to untreated cells (Figures 4E, S3A).

Similarly, transcripts terminating at the canonical 3' end of Stag1 are strongly reduced in

the SP and 3p siRNA KD samples and to a lesser extent in the 5p KD, while the transcript

terminating in i25 is substantially enriched upon 3p KD (Figure 4E). Immunoprecipitation

of STAG1 using an antibody which recognizes the N-terminus in Stag1$^{NG\_FKBP}$ ES cells

treated with dTAG reveals the enrichment of STAG1-CtermΔ isoforms (Figure 4F,

arrows). Thus, the siRNA panel used here provide us with a powerful tool to modulate the

proportion of the divergent ends of STAG1 in ES cells and study their potential roles in

cell fate regulation.

The C-terminus of STAG1 has a specific role in maintenance of the naïve

pluripotent state.

We analyzed pluripotency gene expression in the different KDs using RT-qPCR and RNA-

seq. RT-qPCR results suggested that 5p and 3p KDs may have differential effects on

Nanog expression in serum-grown ES cells (Figure S4C). Namely, there was a consistent

tendency towards Nanog downregulation in 3p KD while 5p KD had little effect on Nanog.

GSEA supported the differential effect of the SATS, 3p and 5p KDs on naïve and primed pluripotency signatures. In support of STAG1 playing a role in pluripotency, reducing Stag1 levels by targeting the ES-specific SATS promoter leads to downregulation of the naïve pluripotency gene signature and upregulation of the primed signature (Figure 4G), reminiscent of the phenotype from SP KD (Figure 1G). A similar, more prominent loss of the naïve signature was observed in 3p KD, while the opposite was true for 5p KD cells where the naïve signature was maintained (Figure 4G).

The distinct gene expression signatures of the 3p and 5p KDs are reflected in differences in cellular phenotypes. Cells treated with 3p siRNAs exhibited a further loss of self-renewal potential, consistent with the loss of the naïve pluripotency signature, with 20% of colonies exhibiting AP-staining compared to 30% of colonies in the SP KDs, and reduction of the area occupied by AP+ colonies by 40-60% (Figures 4H, S4D). This was not evident in the 5p KD, where the effect on self-renewal was similar to scrambled control treated cells (Figures 4H, S4D). Interestingly, unlike siRNA to Stag1, esiSATS results in a variable effect on self-renewal (ranging from between 5-35% reduction in AP+ area) (Figure 4H), likely because the SATS TSS is expressed in the most naïve cells of the population, the frequency of which varies significantly between FCS populations. Our results further confirm the importance of STAG1 in self-renewal and point to a specific role for the C-terminal of Stag1 in the maintenance of the naïve pluripotent state.

The N-terminus of Stag1 regulates heterochromatin and conversion to totipotent state.

Both the N-termΔ and C-termΔ STAG1 isoforms lost prominent IDRs (Figure 3H), thus we investigated the effect of these on heterochromatin structure. ES cells treated with 5p siRNAs exhibited variable H3K9me3 foci volumes, similar to the effect of SP KD (Figure 2G), implicating the N-terminus in H3K9me3 condensation (Figures S5A, B). On the other hand, H3K9me3 foci volumes in 3p KD cells were not significantly different from scr control cells. DNase I digestion of chromatin further confirmed global changes to chromatin accessibility upon 5p, but not 3p, KD whereby four of six experiments revealed chromatin decompaction (Figure S5C). We built up on these results by generating ES

cell lines expressing dox-inducible SA1$^{e5\Delta}$-GFP or SA1$^{i25-\Delta C}$-GFP, as representative of the

N-terminal, AT-hookΔ and the C-termΔ groups respectively. H3K9me3 foci in SA1$^{i25-\Delta C}$-

GFP-expressing cells revealed increased condensation compared to control cells, with a

similar effect as SA1^{FL}-GFP. Meanwhile, the condensation of H3K9me3 foci in SA1$^{e5\Delta}$-

GFP-expressing cells was significantly attenuated from SA1^{FL}-GFP cells (Figures 5A, B).

Thus, while both the N- and C-termini of STAG1 play a role in heterochromatin structure,

loss of the basic N-terminus (which interacts with DNA via the AT-hook region(Bisht et

al., 2013)), significantly affects condensate structure, indicating its importance in

heterochromatin compaction.

Since STAG1 was bound to repeats, which play significant roles in ES cell fate

determination, and are involved in heterochromatin regulation, we also profiled the

expression of LINE-1, IAPEz and MERVL elements in our KD panel. Surprisingly, MERVL

elements were significantly derepressed in 5p KD, but not in SP or 3p KD conditions

(Figures 5C and S5D). The effect was specific to MERVL, we saw no significant change

to the expression of LINE1-T or IAPEz (Figure 5C), despite the enrichment of STAG1 at

L1T elements (Figure 2D). Re-activation of MERVL is a hallmark of a rare subpopulation

of totipotent cells (termed two-cell-like, 2C-L) which spontaneously arise in ES cell

cultures and exhibit unique molecular and transcriptional features including chromatin

decompaction, reminiscent of our 5p KD (Eckersley-Maslin et al., 2016; Ishiuchi et al.,

2015; Macfarlan et al., 2012). Indeed, changes to MERVL expression in the 5p KD was

accompanied by increased expression of additional hallmark 2C-L genes and chimeric

transcripts (Dux, Gm6763, AW822073 and Gm4981) in serum-grown and 2i cells (Figures

5D, S5E). We note that the effect of 5p KD on 2C-L genes is modest in serum-grown ES

cells and significantly stronger in naïve ES cells. Notably, all 2C-L genes analyzed

remained unchanged in 3p KD conditions with a modest but insignificant upregulation in

SP KD. Despite the modest but coordinated changes we observed by qRT-PCR of 2C-L

genes (Figure 5D), GSEA using a published 2C gene set (Percharde et al., 2018)

revealed a specific enrichment among the upregulated genes in 5p KDs that was not

observed in 3p KDs (Figure 5E, S5F). Together our results point to a specific role for the
N-terminus of STAG1 in totipotency regulation.

To investigate this further, we asked whether the loss of the N-terminus of STAG1
drives conversion of ES cells towards a totipotent 2C-L state. We obtained ES cells
expressing a dox-inducible *Dux-HA*-expression construct together with a MERVL-linked
GFP reporter (Hendrickson et al., 2017) and used flow cytometry to measure the number
of GFP-positive cells in our different Stag1 KD conditions (Figures 5F, G). Chaf1 is a
chromatin accessibility factor previously shown to support conversion of ES cells towards
totipotency (Ishiuchi et al., 2015). In support of the upregulation of the 2C gene set in 5p
KD cells, we observed an 8-9% increase in the proportion of GFP-positive cells in 5p KD
conditions compared to scramble treated controls, similar to the effect of Chaf1 KD
(Figures 5F, G). There was a modest increase in GFP+ cells upon SP KD and no effect
upon 3p KD. Chaf1 and 5p double KD had an additive effect on the proportion of GFP-
positive cells, suggesting that the two proteins function in complementary pathways for
conversion towards totipotency.

A 2C-specific Stag1 promoter.

Our results point to the N-terminus of Stag1 having a protective function in the conversion
of ES cells towards totipotency. Given the known role for increased chromatin
accessibility during conversion to 2C (Ishiuchi et al., 2015) and the importance of the N-
terminus in heterochromatin organization reported here, we propose that the Stag1 N-
terminus, and specifically the AT-hook within it, is involved in heterochromatin silencing
by forming or maintaining a condensed heterochromatin structure. This predicts that the
2C-L cells which spontaneously arise within the ES population could express Stag1NΔ
protein isoforms to support their preferred chromatin status and reinforce their cell state.
To formally address this, we induced *DuxHA*-expression in the MERVL-GFP ES cells and
performed 5' RACE on sorted GFP+ and GFP- cells (Figure 5H). We enriched several of
the previously identified N-term truncated Stag1 transcripts in the GFP+ population

including e2/3Δ and e5Δ isoforms. Importantly, we also identified a transcript starting at

e7, similar to the one previously found in 5p KD ES cells (Figure 3B, S3C). Remarkably

however, the sequence preceding the TSS in e7 in *Dux*-induced cells was an MT2-

MERVL element, creating a chimeric, LTR-driven transcript akin to others specifically

expressed in the 2C-L state (Figures 5H, I). Thus, the 2C-L state selectively expresses

an N-term truncated Stag1 isoform which in turn supports the maintenance or emergence

of that state.

Stag1 regulates 2C fate via changes to nucleolar structure and function.

As ES cells preferentially expressing STAG1NΔ isoforms led to conversion towards

totipotent cell states and STAG1 is localised to the nucleolus and interacting with

Nucleolin, we asked whether the STAG1-induced 2C-L state could be explained by the

association of STAG1 with the Nucleolin/Trim28 complex known to derepress Dux targets

(Percharde et al., 2018). While STAG1 proteins directly interact with Nucleolin as well as

Trim28 (Figures 5J), Dux itself is weakly or variably deregulated in Stag1 KDs (Figure

5D). Similarly, while MERVL is robustly derepressed upon 5p KD (Figures 5C, F), Stag1

protein is not enriched at MERVL elements (Figure 2D), suggesting that the observed

effects are unlikely to be via *Dux*-mediated derepression of targets and may be indirect.

Thus, we investigated whether the Stag1-induced 2C-L state could be explained by

changes to nucleolar structure or function. In this context, we examined the consensus

sequence of the rDNA locus and note that there are several SINE elements located within

the intergenic spacer (IGS). Analysis of Stag1 ChIP-seq alignments to this region is

complicated by the repetitive nature of the region, however there was evidence of

possible Stag1 binding to the B3 elements within the IGS, suggesting that Stag1 may

directly support nucleolar structure and function (Figure S6B).

We explored this hypothesis by investigating the effect of Stag1 loss on rRNA

transcription. ES cells were pulsed with 5-ethynyl uridine (EU) which becomes actively

incorporated into nascent RNA and enables detection of newly synthesized RNA either

spatially by immunofluorescence or globally by flow cytometry. Samples for immunofluorescence were co-stained with an antibody to Nucleolin to quantify changes in nascent rRNA transcription. Cells treated with scramble siRNA showed a distinct nucleolar structure and the EU signal could be seen throughout the nucleus, with a strong enrichment within the nucleolus as expected from rRNA expression (Figures 6A, B). A significant reduction in nascent rRNA signal was observed in all KD conditions compared to the si scr control. Interestingly, while the medians between the three siSA1 KDs were not dramatically different, the effect of the 5p KD on rRNA signal distribution was significantly different from the 3p KD, revealing different roles for the N and C-terminal ends of Stag1 in nucleolar function. In parallel, we used flow cytometry to validate the above and quantify the effect on global levels of transcription upon Stag1 loss. In support of the immunofluorescence results, we observed a reduction in nascent transcription in all siSA1 KDs compared to scrambled control (Figures 6C, D). SP and 5p KD reduced nascent transcription by 39% and 47% respectively, the bulk of which is likely from the observed change to rRNA expression. Meanwhile, cells treated with 3p siRNA exhibited a 16% reduction in global transcription compared to scrambled controls, further implying that the N- and C-terminal ends of Stag1 likely target different pools of newly synthesized transcripts.

Finally, given the changes to nascent rRNA transcription and the known role for translation in 2C-L state conversion (Eckersley-Maslin et al., 2016; Hung et al., 2013), we measured effects on translation. Indeed, previous results showed a stronger effect of Stag1 KD on Nanog protein compared to Nanog mRNA levels (Figures 1D, E). ES cells were pulsed with L-homopropargylglycine (HPG), an amino acid analog of methionine and flow cytometry was used to quantify the impact of Stag1 KD on global protein synthesis. The proportion of cells that had reduced incorporation of HPG increased significantly in the SP and 5p siRNA treated cells compared to scramble controls (32% and 35% of si Scr) (Figures 6E, F). We did observe a mild affect on global nascent translation in 3p KD treated cells (16% of si scr), although this was not significantly different from scramble controls (Figures 6E, F). Overall, our work has uncovered a novel

506   role for Stag1 in translation control and proposes that Stag1 may act directly at the rDNA

507   locus or via Nucleolin to influence the regulation of rRNA expression. To our knowledge,

508   a role for Stag1 in translation regulation has not been previously described and offers a

509   completely new perspective on how Stag1 impacts cell fate, moving beyond its known

510   roles in gene expression by regulating chromatin loops.

DISCUSSION

Cell fate transitions during early development are accompanied by extensive transcriptional and epigenetic reprogramming. In addition to changes in protein-coding gene expression, repetitive sequences play an integral role in developmental programmes (Jachowicz et al., 2017; Macfarlan et al., 2012; Percharde et al., 2018). Repetitive sequences are spatially clustered into heterochromatin domains to facilitate their regulation (Allshire and Madhani, 2018; Padeken and Heun, 2014). How cohesin function, or Stag proteins contribute to heterochromatin or repeat regulation has not been extensively studied in mammalian cells, despite the importance of both for cell identity.

While previous studies have implicated Stag proteins in cell fate decisions (Cuadrado et al., 2019; Viny et al., 2019), they have primarily focused on the impact to protein-coding gene regulation. By discovering a novel role for Stag1 in heterochromatin structure, repetitive elements and translation control, we significantly expand our understanding of Stag1 functions and thus deliver new insight into the mechanisms by which Stag proteins and cohesin regulation impact cell identity and lead to disease. Further, by comprehensively characterising naturally occurring Stag1 protein isoforms, we identify distinct roles for the N- and C-termini of Stag1 which shed mechanistic light on the reported distinct functions of the Stag paralogs (Cuadrado and Losada, 2020). For example, the N-terminal containing AT-hook of Stag1 represses heterochromatin condensation and its loss leads to global changes in chromatin compaction and reprogramming of ES cells towards totipotent 2C-L cells. Our observation that the AT-hook domain is required for heterochromatin compaction may explain the dominance of Stag1 in stem cell populations since Stag2 does not have an AT-hook and heterochromatin must be tightly regulated during early development.

We have characterized diverse transcription–regulatory events at the Stag1 locus in stem cell populations giving rise to protein isoforms with distinct regulatory domains. Importantly, this extensive Stag1 protein heterogeneity supports a continuum of cell states within the population since experimentally induced imbalances in isoform proportions skews cell fate probabilities. Using RACE in cells enriched for the 2C-L fate,

we find evidence for skewed expression of particular Stag1 isoforms, arguing that

individual ES cells may naturally predominantly express particular Stag1 protein isoforms.

Interestingly, the proportions of Stag1 isoforms likely originate in part from the stochastic

process of splicing (Fiszbein and Kornblihtt, 2017; Gabut et al., 2011; Salomonis et al.,

2010), which is itself further randomized by the fluctuating chromatin landscape (Mateo

et al., 2019; Rodriguez et al., 2019), acting as a feedback mechanism for further Stag1

diversity. Thus, the naturally occurring Stag1 isoforms described here may act as intrinsic

sources of noise that both directly and indirectly support cellular plasticity.

We propose that a balance of Stag1 protein isoforms is required for variation in

gene and repeat expression and thus, a continuum of cellular states. The N-terminal

containing AT-hook of Stag1 supports global chromatin compaction and heterochromatin

condensation, suggesting that it plays an important role in clustering repetitive sequences

into heterochromatin condensates for their regulation. Indeed, we observe dramatic

changes to rRNA expression in 5p KD cells where isoforms lacking the N-terminus are

specifically retained. Moreover, we note that our live cell microscopy also revealed

condensate-like structures within the nucleoplasm. It is possible that the C-terminus of

Stag1 regulates euchromatin condensate formation to support pluripotency gene

networks. Several lines of evidence support this. First, many of the genes that are

downregulated upon 3p KD are super-enhancer-associated genes which are known to

form condensate structures (Sabari et al., 2018; Whyte et al., 2013). Further, global

nascent transcription is reduced in 3p KD cells, although rRNA is not as affected as in the

5p KD, suggesting that the C-term of Stag1 may have a role in regulating a different

subset of nascent transcripts. Thus, the Stag1 protein heterogeneity discovered here

provides the necessary fluctuations in the levels and composition of the Stag1 disordered

regions to impact the stability of both euchromatin and heterochromatin condensates,

thereby supporting plasticity within the population.

Stag1 knockout (Stag1$^{\Delta/\Delta}$) ES cells give rise to mice which survive to E13.5

(Remeseiro et al., 2012b). At first this observation seems at odds with our report that

Stag1 is required for pluripotency. However, our observations may in fact explain why the

Stag1$^{\Delta/\Delta}$ mouse model does not exhibit early embryonic lethality. In this model, only the

5' region of Stag1 was targeted, meaning that the Stag1 isoforms lacking the N-terminus

may still be retained in the targeted ES cells. This is consistent with our results showing

that 5p KD cells have not lost their ability to self-renew nor is their pluripotency gene

signature affected. It further suggests that changes to heterochromatin may exist in these

cells.

The role for Stag1 in heterochromatin structure described here may be quite

general, and the fact that Stag1 binds to HP1a and repetitive sequences known to form

constitutive heterochromatin suggests that this is true. Indeed, the AT-hook domain of

Stag1 has been shown to be important for telomere cohesion (Bisht et al., 2013). Here

we have specifically uncovered a role for Stag1 in nucleolar structure and function in stem

cells. Transcriptionally inactive rDNA arrays are associated with the periphery of the

nucleolus and accumulate marks of constitutive heterochromatin while actively

transcribed rDNA loci are looped inside the nucleolus (Padeken and Heun, 2014). We

observe Stag1 at both the interior and periphery, suggesting that it may be involved in

multiple aspects of rDNA and nucleolar regulation. Stag1 is enriched at SINE elements

and we note that the intergenic spacer region of the rDNA consensus contains multiple

B3 and B2_Mm2 sequences, strongly suggesting that the changes in nascent rRNA

transcription are direct effects of Stag1 loss and implicates Stag1 in the formation or the

maintenance of the nucleolus. Indeed, CTCF has previously been implicated in nucleolar

structure by tethering insulators to the periphery (Yusufzai et al., 2004). As well as

impacting global translation, the change to nucleolar structure may also indirectly support

the emergence of the 2C state by destabilizing complexes such as Nucleolin/Trim28

which are required for the derepression of repeat elements (Percharde et al., 2018).

Finally, a role for Stag proteins or cohesin in translation regulation has, to our

knowledge, not been described before despite the fact that translation has important roles

in development (Buszczak et al., 2014). How Stag1 contributes to translation in ES cells

and whether it plays similar roles in other stem or progenitor populations are important

future questions. Our results offer new perspectives on how Stag proteins and cohesin

regulation contribute to cell identity during development and in disease.

Acknowledgments

This work would not be possible without the support of a Senior Research Fellowship from the Wellcome Trust awarded to S.H. (106985/Z/15/Z). We would like to thank Sally Lowell and Mattias Malaguti for advice throughout the project. We are grateful to the members of the Hadjur lab for critical discussions and reading of the manuscript. We thank M. Irima for help with VAST-tools pipelines; B. Cairns for the inducible Dux-HA, MERVL-GFP cell line; W.Reik for a second MERVL-reporter ES cell line; H. Rowe for advice on 2C-L cells and J. Vaquerizas for advice on repeat expression analysis. Thank you to Y. Guo and J. Manji in the Cancer Institute CRUK Centre FACS and Imaging core facilities for their invaluable assistance.

Author Contributions

D.P. and S.H. conceived the project. D.P. designed and performed all the experiments on ES cells with assistance from S.W. S.W. performed all protein analysis, generated the SA1-NG-FKBP ES cell line, performed the Spinning Disk microscopy and helped with the siRNA knockdown experiments. W.V. performed all bioinformatic analyses with the exception of the Stag1 enrichments at repeat elements, which was done by M.B. P.D. and S.P. provided advice on CRISPR targeting. D.P. and S.H. formatted all figures and wrote the manuscript with input from all authors.

Declaration of Interests

The authors declare no competing interests.

Figure legends

Figure 1. STAG1 is required for pluripotency.

(A) Log2 fold change of pluripotency factors and cohesin subunit gene expression assessed by qRT-PCR during embryonic stem cell (ES) differentiation towards Epiblast-like (EpiLC) cells. Multiple primer pairs were used for SA1 and SA2 mRNA. Data are derived from two biological replicates.

(B) Whole cell protein extracts (WCL) from serum-grown (FCS) or naïve (2i) ES cells, EpiLC and mouse embryonic fibroblasts (MEF) and analyzed by western blot (WB) for levels of SA1, SA2 and Smc3.

(C) WB analysis of SA1 levels in WCL, cytoplasmic and chromatin fractions upon treatment with scrambled siRNAs (siScr) or SmartPool SA1 siRNAs (siSA1) for 24hrs in 2i ES cells. Tubulin (Tub) and H3 serve as fractionation and loading controls.

(D) Relative expression of Nanog mRNA by qRT-PCR in 2i- (left panel, n=19) or FCS-grown (right panel, n=20) ES cells after no (UT), si scr or si SA1 treatment. Whiskers and boxes indicate all and 50% of values respectively. Central line represents the median. Asterisks indicate a statistically significant difference as assessed using 2-tailed T-test. * p<0.05, ** p<0.005, n.s., not significant.

(E) Mean Fluorescence Intensity (MFI) of Nanog protein assessed by Immunofluorescence in 2i ES cells treated with si scr, a second siRNA control to Luciferase (esi Luc) and si SA1. Cells were stained for Nanog, SA1 and counterstained with DAPI. Quantifications and statistical analysis were done as above. Each treatment was repeated a minimum of two times (n>100/condition). **** p<0.0005.

(F) Volcano plot displaying the statistical significance (-log2 p-value) versus magnitude of change (log2 fold change) from RNA-sequencing data produced in FCS ES cells treated with si scr or siSA1 for 24hrs (n=3). Vertical blue dashed lines represent changes of 2-fold. Selected genes associated with pluripotency and differentiation have been highlighted.

(G) Enrichment score (ES) plots from Gene Set Enrichment analysis (GSEA) using curated Naïve or Primed gene sets (see Methods). Negative and positive normalized (NES) values point to the gene set being over-represented in the top most down- or up-regulated genes in SA1 KD ES cells, respectively. Vertical bars refer to individual genes in the gene set and their position reflects the contribution of each gene to the NES.

(H) Area occupied by AP+ colonies in ES cells treated with si scr and si SA1 from three
independent biological replicates where n>50 colonies/condition were counted.

(I) CRISPR/Cas9 was used to knock-in a NeonGreen-v5-FKBP tag on both alleles of
endogenous Stag1 at the C-terminus (SA1$^{NG-FKBP}$). The resultant Stag1 protein is 42kDa
larger. Shown are known features of SA1 including the N-terminal AT-hook (AT) and the
stromalin conserved domain (SCD). WB analysis of SA1 levels in a targeted ES clone
after treatment with DMSO-only or dTAG for 16 hrs. Tubulin (Tub) serves as a loading
control.

(J) Analysis of the area of AP+ colonies as above in wildtype ES and in a SA1$^{NG-FKBP}$ ES
clone treated with DMSO-only or dTAG for 16 hrs. Data is represented as mean ± SEM
of three independent experiments and relative to the DMSO control.

See also Figure S1.

Figure 2. STAG1 associates with AT-rich heterochromatin.

(A) Live-cell Spinning Disk confocal images of three SA1$^{NG-FKBP}$ cells counterstained with
Hoechst. Arrows indicate notable regions of overlap of SA1 and Hoechst, including inside
the nucleolus and at Hoechst-dense foci (top cell), and at the nucleolar periphery (bottom
cells).

(B) Mean Intensity of SA1-NeonGreen signal from (A) within the whole nucleus or
Hoechst-dense foci as identified using Imaris in the SA1$^{NG-FKBP}$ clone treated with DMSO
or dTAG. n=270 cells/condition.

(C) co-Immunoprecipitation of endogenous SA1 with Smc3, HP1a and Nucleolin in ES
cells.

(D) Percentage of SA1 peaks (unique reads) at promoters, exons, transposable elements
introns and intergenic sequences.

(E) STAG1 ChIP-seq data (unique and multimapping reads) aligned to full-length Repeat
elements of the indicated SINE, LINE and LTR families.

(F) Confocal images of Immunofluorescence (IF) to SA1 and H3K9me3 in siscr or siSA1
treated ES cells, counterstained with DAPI.

(G) Imaris quantification of the volume of H3K9me3 foci from (F). Box plots and statistical
analysis were done as above. Each treatment was repeated twice (n>100/condition).
p<0.05.

(H) Global chromatin accessibility as detected by DNase I digestion of genomic DNA at
different concentrations from 200,000 cells/condition in siScr or siSA1 ES cells.

(I) Confocal images of IF to GFP and H3K9me3 in ES cells expressing full-length SA1
tagged with GFP (SA1-FL), counterstained with DAPI.

(J) Imaris quantification of the volume of H3K9me3 foci from (I). Box plots and statistical
analysis were done as above. n>50/condition. p<0.0005.

(K) PONDR (Predictor of Natural Disordered Regions) analysis of SA1 protein using VSL2
predictor at http://www.pondr.com showing consecutive stretches of disordered regions
corresponding to the N- and C-terminus of Stag1.

See also Figure S2.

Figure 3. Diverse Transcription-Regulatory Control of Stag1 in ES cells.

(A, B) 5' Rapid Amplification of cDNA ends (RACE) for SA1 in ES and EpiLC cells. Arrows
indicate bands which were cloned and sequenced. In A, red star indicates SATS TSS and
red arrow indicates canonical TSS. In B: red indicates full length Stag1 with both SATS
and can TSSs; dark blue indicates alternatively spliced variants, skipping of various exons
in 5' region; light blue indicates the TSSs at e6, e7.

(C) 3' RACE for SA1 in ES cells. Arrows indicate bands which were cloned and
sequenced. Red indicates canonical full-length end; green indicates end in i25.

(D) Top, schematic of the Stag1 gene annotation in mm10 and the identified TSS and
TTSs from RACE indicated. Bottom, aligned sequence clones from the PCR mini-screen
and their predicted impact on the SA1 protein, right. Green arrows and red bars within the
transcripts indicate start of the coding sequence and the TTS respectively. Shown also
are the regions which code for the AT hook and SCD domains.

(E) Percent Spliced In (PSI) calculations based on VAST-Tools analysis of RNA-seq from
multiple 2i (blue) and FCS (red) datasets (see Methods for details of libraries). Data are
shown relative to Neural Stem cell (NSC) frequencies to highlight the events that are ES-
specific.

(F) Genome topology at the Stag1 locus. Hi-C contact maps in ES (2i) and NS cells of the 900kb region on chromosome 9 containing the Stag1 topologically associated domain (TAD). TADs are denoted with a vertical line and as repressed (orange) or active (blue). Shown also are tracks for Genes, Nanog and CTCF ChIP-seq as well as a track indicating the directionality of CTCF binding sites (red, forward; blue, reverse). Aligned to the Gene track are also the SA1 transcripts discovered above where red represents the untranslated regions and blue the coding body. UMI-4C-seq viewpoints (asterisks on the ChIP tracks) are positioned to the leftmost CTCF site ('CTCF bait') and to the Nanog site 40 kb upstream of the Stag1 canonical TSS ('Nanog bait'). For each bait, UMI information for each cell type is shown as well as the comparative plots where red represents an enrichment of contacts in ES compared to NS.

(G) Top, cartoon depicting functional domains within Stag1 protein, including the AT-hook (aa 3-58); Stromalin conserved domain (SCD, aa 296-381) and the C-terminus. Bottom, the predicted Stag1 protein isoforms based on transcript analysis with estimated sizes for each isoform. Purple boxes in the 105kDa and 90kDa isoforms represent retained introns.

(H) PONDR tracks as before shown for the N-terminal truncated (top) and C-terminal truncated (bottom) transcripts.

(I) Chromatin immunoprecipitation of SA1 from ES cells.

(J) WB analysis of SA1 isoforms in chromatin fractions from ES cells treated with siscr and siSA1. H3 serves as a fraction and loading control.

(K) Chromatin immunoprecipitation for the v5 tag in SA1[NG-FKBP] ES cells treated with DMSO-only or dTAG. Note, SA1 bands now run 42kDa higher due to the addition of the tag.

See also Figure S3, Table S1 and Table S2.

Figure 4. Fluctuations in the levels of the Stag1 isoforms skews cell fates.

(A) Schematic of the siRNA pools used in this study. esiRNA SATS represents 'enzymatically-prepared' siRNAs (see Methods).

(B) Relative expression of SA1 mRNA by qRT-PCR in FCS ES cells after no treatment (UT), or upon si scr, si SA1 SP, si 3p, si 5p or esi SATS treatment. Data is represented as mean ± SEM, n=3. * $p<0.05$

(C) WB analysis of SA1 levels in FCS ES WCL upon treatment with the same siRNA panel from (B). Tubulin serves as a loading control.

(D) RNA-seq reads (as transcripts per million, TPM) aligning to sectioned Stag1 in datasets from the various siRNA pools and presented as relative to untreated ES RNA-seq. N-terminal reads include SATS and exons 1-8, Mid reads are from exons 12-19 and C-terminal reads include exons 20-25 and exons 26-34.

(E) Left, 5' and Right, 3' RACE for SA1 in ES cells treated with the siRNA panels. Arrows indicate bands which were cloned and sequenced and colour-coded as before.

(F) Chromatin immunoprecipitation using an N-terminal Stag1 antibody in SA1[NG-FKBP] ES cells treated with DMSO-only or dTAG for 16 hrs. Green arrow indicates residual C-terminal truncated Stag1 isoforms.

(G) Enrichment score (ES) plots from GSEA using Naïve or Primed gene sets as in Figure 1G and RNA-seq data from the different siRNA treated ES cell samples.

(H) Area occupied by AP+ colonies in ES cells treated with the siRNA panel from three independent biological replicates. n>50 colonies/condition were counted.

See also Figure S4.

Figure 5. Stag1 N-terminus represses heterochromatin and promotes transition towards totipotency.

(A) Confocal images of IF to GFP and H3K9me3 in ES cells expressing SA1-FL-GFP, SA1-exon5Δ-GFP or SA1-CtermΔ-GFP. Nuclei were counterstained with DAPI.

(B) Imaris quantification of the volume of H3K9me3 foci per cell from (A). Box plots and statistical analysis were done as before. n>50/condition. ** $p<0.005$, ****$p<0.0005$.

(C, D) Relative expression of (C) repeat elements; MERVL, LINE1-T and IAPEz RNA (n=8 biological replicates each) and (D) 2C-L genes; Gm6763 (n=6) and Dux (n=5) by qRT-PCR in FCS ES cells after treatment with the siRNA panel. Boxes and whiskers as before. * $p<0.05$, n.s, not significant as per 2-tailed T-test. NB. The p-value for Gm6763 expression in 5p KD vs UT = 0.07.

(E) Enrichment score (ES) plots from GSEA using a published 2C-L gene set (Methods) and RNA-seq data from the 3p and 5p siRNA treated ES cell samples.

(F) Representative FACS analysis of the proportion of ES cells expressing a MERVL-
GFP reporter in the different siRNA treated cells and including siRNA to Chaf1 as a
positive control. Percentage of MERVL-GFP+ cells based on Flo-Jo analysis is shown in
red.

(G) Proportion of MERVL-GFP+ cells in the different siRNA conditions relative to the
siChaf1 positive control. Data is represented as mean ± SEM of four independent
experiments. * $p<0.05$, ** $p<0.005$, n.s, not significant as per 2-tailed T-test.

(H) 5' RACE for SA1 in Dux-HA, MERVL-GFP ES cells sorted for GFP+ cells. Arrows
indicate bands which were cloned and sequenced and colour-coded as before.

(I) Sequence of the 5'RACE product identifying a novel Stag1 TSS with direct splicing of
exon7 to MT2_MERVL element.

(J) Co-Immunoprecipitation of SA1 with Nucleolin and Trim28 in ES cells.

See also Figure S5.

Figure 6. rRNA expression and nascent translation upon Stag1 loss.

(A) Confocal images of IF to Nucleolin (Ncl) and Nascent RNA in ES cells treated with the
siRNA panel. Nuclei were counterstained with DAPI. Note the accumulation of nascent
RNA within the nucleolus.

(B) Imaris quantification from (A) of the mean intensity of nascent RNA (EU-488) within
the nucleoli, as defined by Nucleolin signal. Box plots and statistical analysis were done
as before. Data are from two biological replicates, n>50/condition, except for siSA1 5p
where n>35. ** $p<0.005$, ****$p<0.0005$ as per 2-tailed T-test.

(C, D) Global analysis of Nascent transcription by measuring EU-488 incorporation using
Flow cytometry. (C) representative Flo-Jo analysis of EU incorporation in ES cells treated
with the siRNA KD panel. (D) Quantification of the change in EU incorporation relative to
si Scr treated cells. Data are from two biological replicates *$p<0.05$, **$p<0.005$ as per 2-
tailed T-test.

(E, F) Global analysis of Nascent translation by measuring HPG-597 incorporation using
Flow cytometry. (E) Representative Flo-Jo analysis of HPG incorporation in ES cells
treated with the siRNA KD panel. (F) Quantification of the change in EU incorporation

relative to si Scr treated cells. Data are from four biological replicates *p<0.05, **p<0.005
as per 2-tailed T-test.

See also Figure S6.

METHODS

Embryonic stem cell culture and siRNA-mediated knockdown.

Male mouse E14 embryonic stem (ES) cells were cultured in serum (FCS) or naïve (2i) conditions. Serum-cultured cells were grown on 0.1% gelatin-coated plates in GMEM, 10% FCS (Sigma), NEAA, Na Pyruvate, 0.1 mM ßMercaptoethanol (BMe), Glutamax, and freshly added LIF (1:10,000). 2i-cultured cells were grown on plates coated with Fibronectin, in DMEM:F12/Neurobasal 1:1, KnockOut Serum Replacement, N2, B27, Glutamax, 1μM PD0325901, 3μM CHIR9902, 0.1 mM BMe, and freshly added LIF as above. DuxHA/MERVL-GFP cells were cultured in 2i conditions. siRNAs were purchased from Horizon Discovery (previously Dharmacon) or Sigma (for 'enzymatically-derived' esiRNAs). siRNA knockdowns (KDs) were performed for 24hr with the exception of those in Figure 5 which were performed for 72hr. Knockdowns were performed in 6-well plates where 200,000 cells were seeded for 72 hr KDs, and 400,000 for 24 hr KD. 50pmol siRNAs were transfected using RNAiMax Lipofectamine at the time of seeding, and after 48 hrs for 72hr timepoints. Two siRNA controls were used, scrambled (scr) was D-001810-10 and Luciferase (esiLuc) control purchased from Sigma. siSA1 'SmartPool' (SP) was derived from equimolar ratios of commercial siRNAs (D-041989-02, -04, -05, -06, -07, -08). siSA1 5p was a custom Duplex siRNA sequence (AGGAGCAGGUCGUGGAAGAUU). siSA1 3p was derived from equimolar ratios of commercial siRNAs J-041989-05, -07, -08. esiRNA to SATS was purchased from Sigma as a custom-made product to the entire SATS 5'UTR (mm10 chr9:100,597,794-100,598,109).

qRT-PCR analysis

Total RNA was isolated using Monarch RNA prep kit (NEB). Reverse transcription was performed on 0.5 μg DNase-treated total RNA using Lunascript RT (NEB) in 20μl reactions. qPCR was performed using 2x SensiFAST SYBR No-ROX kit (Bioline) in 20 μl reactions using 1μl of RT reaction as input and 0.4μM each primer.

Alkaline Phosphatase (AP) assay and quantification

Cells were seeded in 6 well plates and transfected with siRNAs at the time of plating as above. After 24 hrs, cells were collected for RNA isolation and KD efficiency analyzed by qRT-PCR. Cells from each condition were counted and 1,000 cells per well seeded into a new 6-well plate. Cells were re-transfected after 48 hrs using 5 pmol of siRNAs. Cells were fed every day. Four days

after seeding cells at clonal density, the cells were assayed for alkaline phosphatase (AP)

expression using StemTAG Alkaline Phosphatase staining kit (Cell Biolabs CBA-300). AP stained

cells were imaged in 6-well plates using a M7000 Imaging System (Zeiss) with a 4X objective and

a Trans-illumination brightfield light source. For quantification, AP-high and AP-low colonies from

each condition were counted. Area occupied by AP-high colonies was also measured using

ImageJ, and plotted as fraction of total area of all colonies.

RACE (Rapid Amplification of cDNA Ends) and PCR mini screen

RACE was performed using GeneRacer kit (RLM RACE, Invitrogen L1500). 2μg of total RNA was

used as input. Final products were amplified by nested PCR, using Kapa 2x MasterMix. First PCR

was done in a 50μl reaction using 1μl RT as input, 25 cycles. DNA was purified using Qiagen

PCR Purification kit, and nested PCR was performed on a tenth of the first PCR for 30 cycles.

Viewpoint for 5'RACE was in exon 2 (Fig 3A) or exon 8 (Fig 3B) of Stag1. Viewpoint for 3'RACE

was in exon 23 (Fig 3C). RACE primer details can be found in Table S3. PCR products were

excised from the gel, A-tailed using Klenow exo- (NEB) and cloned into pCR4-TOPO vector

(Invitrogen). At least three clones were sequenced per PCR product. For the PCR Mini-Screen,

forward primers at either SATS or canonical 5' UTR were used with reverse primers either at the

end of Stag1 canonical coding sequence, or at the end of coding sequence in intron 25 (see Table

S3). PCR was performed using Kapa 2x MasterMix. DNA was excised from the gel, A tailed, and

cloned into pCR4-TOPO. At least six clones per PCR product were Sanger-sequenced.

Sequences from the PCR Mini-screen were aligned using Minimap2 (2.14-r884) in 'splice' mode

to ensure long read splice alignment (Fig 3D and S3A).

PONDR Predictions

Internally disordered regions were predicted using VSL2 predictor at http://www.pondr.com.

CRISPR-Mediated Stag1 Knock-in Cell Line Generation

The guide RNA targeting Stag1 3' terminal coding region was designed using Tagin Software

(http://tagin.stembio.org) and purchased from IDT. Lyophilised gRNA was rehydrated in RNA

duplex buffer (100μM). The single stranded oligodeoxynucleotides (ssODN) encoding

mNeonGreen (mNG)-V5-FKBP12^{F36V} and the left and right homology arms was designed using

the software tool ChopChop (https://chopchop.cbu.uib.no) and purchased as a High-Copy Amp-

resistant plasmid from Twist Bioscience. 2.2μl gRNA (100μM) was mixed with 2.2μl tracrRNA

ATTO 550nm (IDT) and annealed together. The RNA duplex was then incubated with $20\mu g$ S.p

Cas9 Nuclease V3 (IDT) for 10min at room temperature and stored on ice prior to transfection.

Linearised KI sequence was mixed with 100% DMSO and denatured at 95°C for 5min. The

ssODN was plunged immediately into ice. The RNP complex was mixed with confluent 2i-grown

ES cells re-suspended in P3 transfection buffer (Lonza) before being transferred to an

electroporation microcuvette well (Lonza). Transfection was performed using a 4D Amaxa

electroporator. Post-nucleofection, the cells were seeded into a fibronectin-coated 6 well plate

with fresh ES media. The media was changed daily for four days before being expanded into a

T75 flask. Confluent ES cells were FACS sorted for GFP+ population (BD FACS Aria Fusion Cell

Sorter) and sparsely seeded into 10 cm plates. Clones were manually picked into 96 well plates

and expanded for selection by v5 IF, genotyping and Sanger sequencing.

Dox-inducible Stag1-GFP isoform cell lines

Stag1 isoforms were cloned into pCW57.1 vector (Addgene 41393), modified using Gibson

assembly to include an EGFP tag at the 3'end of the Gateway cassette, using Gateway

recombination by LR clonase. For primers used to clone the isoforms see Supplementary Table

S3. Plasmids were transfected into 2i-grown ES cells using Lipofectamine 3000 and cells grown

in Puromycin-supplemented media ($1\mu g/ml$) for ten days to make stable lines. Isoform expression

was induced using $2\mu g/ml$ Doxycycline for 24 hrs, and the population enriched for GFP-positive

cells using FACS. For IF experiments, isoforms were induced by adding Dox for 48 hours.

Protein Lysates, Fractionations and Western blotting.

Whole cell lysates (WCL) were collected by lysis in RIPA buffer (150mM NaCl, 1% NP-40

detergent, 0.5% Sodium Deoxycholate, 0.1% SDS, 25mM Tris-HCl pH 7.4, 1mM DTT) and

sonicated at 4°C for x5 30 second cycles using Diagenode Bioruptor. Insoluble material was

pelleted and the supernatant lysate was quantified using BSA Assay (Thermo Scientific). For

cellular fractionations, a cellular ratio of 5×10^{6} cells/$80\mu l$ buffer was maintained throughout the

protocol. Cells were re-suspended in Cell Membrane Lysis Buffer (0.1% Triton X, 10mM HEPES

pH 7.9, 10mM KCl, 1.5mM MgCl2, 0.34M sucrose, 10% glycerol, 1mM DTT), incubated on ice for

5min and centrifuged for 5min at 3700rpm to collect the cytoplasmic sample. The pellet was

washed and then re-suspended in Nuclear Lysis Buffer (3mM EDTA, 0.2mM EGTA, 1mM DTT)

and incubated on ice for 1 hr. Nuclear lysis was aided by sonication with a handheld homogeniser

(VWR) for 10sec at 10min intervals. The nucleoplasmic supernatant and chromatin pellet were

separated by centrifugation at 9000rpm for 10min at 4°C. The chromatin pellet was re-suspended

in 160µl 2X Laemmli Buffer (Bio-Rad). Equal volumes of each fraction were used for Western

Blotting (WB). Cytoplasmic and nucleoplasmic protein samples were diluted in 2X Laemmli Buffer

and boiled for 5min at 95°C, then loaded on a 4-20% SDS-PAGE gel (Bio-rad) or a 3-8% Tris

Acetate gel (Invitrogen). Proteins were wet transferred onto a PDVF membrane (Millipore) and

assessed for successful transfer with Ponceau Red (Sigma). The membrane was blocked with

10% milk and incubated with primary antibodies in 1% milk, 0.1% Tween-PBS overnight at 4°C.

Membranes were imaged with SuperSignal West Femto Maximum Sensitivity (Thermo) on an

ImageQuant.

Chromatin Co-Immunoprecipitation (co-IP)

Cells were re-suspended in 0.1% NP-40-PBS (1ml/$1x10^7$ cells) with 1X Protease Inhibitors

(Roche) and 1mM DTT, and centrifuged at 1500rpm for 2min at 4°C. The pellet was re-suspended

in Nuclear Lysis Buffer (3mM EDTA, 0.2mM EGTA, 1X Protease Inhibitors, 1mM DTT), vortexed

for 30sec before being incubated on a rotator for 30min at 4°C and centrifuged at 6500g for 5min

at 4°C to isolate the glassy chromatin pellet. This was re-suspended in High Salt Chromatin

Solubilisation Buffer (50mM Tris-HCl pH 7.5, 1.5mM MgCl2, 300mM KCl, 20% glycerol, 1mM

EDTA, 0.1% NP-40, 1mM Pefabloc, 1X Protease Inhibitors, 1mM DTT) with Benzonase (Sigma)

($6U/1x10^7$) and incubated on rotator for 30min at 4°C. Chromatin was digested with 3x 10sec

sonication at 30% intensity with a Vibra-Cell probe. The supernatant was collected by

centrifugation at 1300rpm for 30min at 4°C, and then diluted to 200mM KCl concentration with no

KCL buffer. 30µl of Dynabeads (Invitrogen) were used per co-IP. Beads were washed 2x in

200mM KCl IP Buffer, re-suspended in IP Buffer with 10µg of the IP antibody, or an IgG-containing

serum to match the species of the IP antibody and placed on rotator for 5h at 4°C. Beads were

washed 3x in IP buffer and then incubated in 1mg chromatin lysate on a rotator overnight at 4°C.

The beads were washed, re-suspended in 2X Laemmli Buffer (Bio-Rad), boiled for 10min at 95°C

and used for WB as above.

Immunofluorescence and Microscopy

ES cells were cultured on fibronectin or gelatin-coated cover glass in 6-well plates. Cells were

fixed in 4% Paraformaldehyde for 5min and incubated in 0.1% Triton X-PBS for 10min before

being washed and blocked in 10% FCS-PBS for 20min. Primary antibodies were diluted in 10%

FCS, 0.1% Saponin (Sigma) and incubated overnight at 4°C. The next day, the cells were

incubated with an Alexa fluorophore-conjugated secondary antibody diluted in 10% FCS, 0.1% Saponin for 1 hr at room temperature, washed and mounted on cover slides with ProLong Diamond Antifade Mountant with DAPI (Invitrogen). Z-stacks imaging of fixed cells was done using a LSM 880 confocal microscope (Zeiss) with a 63X oil objective. Analysis was performed using Imaris 9.6 (Oxford instruments). Live cell imaging was performed using a 3i Spinning Disc confocal microscope (Zeiss). Stag1-mNG-V5-FKBP12^{F36V} cells were seeded in an 8-chambered coverglass (Lab-Tek II) and DMSO or dTAG (500nM) were added for 24hr before imaging. Directly prior to imaging, cells were incubated with Hoechst 33342 (BD Pharmingen) for 45min, and then replaced with fresh 2i ES media. Cells were imaged as confocal Z-stacks using DAPI and GFP lasers with a 63X objective and 1.4 Numerical Aperture.

Antibodies used in this study

Protein	Catalogue No.	Company	Figure references
Stag1/SA1, N-term epitope	ab4455	Abcam	1B, C, I, S1C, K, 2C, S2C, E, 3J, S3G, 4C, F, 5J
Stag1/SA1, C-term epitope	ab4457	Abcam	2F, S5A, 3I
Stag2/SA2	A300-158A	Bethyl	1B, S1C
Smc3	ab9263	Abcam	1B, 2C
Nanog	ab70482	Abcam	1E, S1F
Tubulin (Tub)	T5168	Sigma	1C, 1I, S2E, 4C, S6A
Actin	Mab8929	Novus	S1C
H3	ab1791	Abcam	1C
v5	14-6796-82	Invitrogen	3K
HP1a	2616	Cell Signalling	2C, S2B, C
Nucleolin (Ncl)	ab22758	Abcam	2C, 5J, 6A, S6A
H3K9me3	ab8898	Abcam	2F, I, S2E, 5A, S5A
H3K4me3	ab8580	Abcam	S2E
Alexa488-anti-GFP (GFP)	A-21311	ThermoFisher	2I, S2A, B, 5A
Trim28	MA1-2023	ThermoFisher	5J

Chromatin accessibility analysis by DNase I treatment.

DNase I digestion was performed as in Huo et al (Mol Cell, 2020), with modifications. 200,000 cells per condition were resuspended in DNaseI digestion buffer (50mM Tris-HCl pH 7.5, 5mM

MgCl$_2$, 0.1 mM CaCl$_2$, 0.2% Triton X, 5mM Na butyrate, protease inhibitor) and incubated for 10

min at room temp. DNase I (50U/μl, ThermoFisher ES0523) was diluted in 1X DNase buffer and

added to cells to have the following Units/μl: 1.25, 2.5, 5, 7.5 and 10. Cells were incubated at 37C

for 10 min in the thermoblock, shaking at 1,000 rpm. To stop the digestion, 10μl 0.5M EDTA and

10ul 10% SDS was added and incubated 10-15min at room temperature. 400μl TE buffer was

added, followed by 10μl PureLink RNaseA (ThermoFisher, 20mg/ml, 12091021), and incubated

at 37C for 1hr. Proteinase K digestion was then performed by adding 100μg of Proteinase K and

incubating at 55C for 3hr to overnight. To isolate DNA, 30μl of 5M NaCl and 525 μl Isopropanol

was added, DNA precipitated at room temperature for 15 min, pelleted by high speed

centrifugation at 4C for 20min, dried, resuspended in 20μl of TE buffer and loaded on 1% agarose

gel. The gel was stained using SYBR Green.

Nascent transcription and translation analysis

For nascent transcription analysis, we used the Click-iT® RNA Alexa Fluor® 488 HCS Assay

(Invitrogen C10327). ES cells were labelled with 1mM EU for 45min at 37C in fresh ES

media. Cells were fixed in solution or onto coverslips with 3.7% paraformaldehyde and

permeabilised with 0.5% Triton-X solution. Cells were incubated with the Click-iT reaction

cocktail for 30min. Cells were then either processed further for Immunofluorescence as

per methods described above (directly to the blocking step) or analysed by flow cytometry

on a BD Fortessa X20. For the Nascent translation analysis, Click-iT™ HPG Alexa Fluor™ 594

Protein Synthesis Assay Kit (Invitrogen C10429) was used. Cells were pre-incubated in

Methionine-free media for 30 min in the 37C incubator before addition of L-homopropargylglycine

(HPG) at 50μM. Cells were incubated with HPG for 30 min, then collected, fixed, permeabilized,

and stained using Click-It reaction in low retention tubes. HPG incorporation was measured by

Flow Cytometry. FACS analysis (in Figures 5,6) was done with FloJo software (version 10.7.1).

Next generation Sequencing and Analysis

Genomic data generated in this study (RNA-seq, PacBio-seq and UMI4C-seq) was submitted to

GEO with the Accession GSE160390.

RNA sequencing (RNA-seq) library preparation and sequencing

ES cells were treated for 24hrs with siRNA pools to Stag1 (SA1) and two sets of control siRNAs,

scrambled (SCR) and Luciferase (Luc). There are three replicate sets for SP KD and two for the

siRNA pools (SATS, 3p, 5p). Total RNA was isolated using NEB Monarch RNA prep kit. 1μg of

total RNA was rRNA-depleted using NEBNext rRNA depletion kit (Human/Mouse/Rat). Libraries

were prepared from 10-50ng rRNA-depleted total RNA, depending on availability of material,

using NEBNext Ultra II directional RNAseq kit according to manufacturer's instructions using 8

cycles of PCR. All ESC FCS libraries were rRNA depleted and only the ESC 2i libraries were

PolyA-enriched before library prep. Two rounds of PolyA+ enrichment were performed. RNA-seq

libraries were sequenced on the Illumina HiSeq3000 platform, 75bp paired-end or single-end

reads. Reads were quality controlled using FASTQC. RNA-seq data was processed using the

RNA-seq Nextflow pipeline (v19.01.0), with the following parameters –aligner hisat2 –genome

mm10, with –reverse_stranded specified for paired-end samples. FeatureCounts output was

parsed through edgeR (v3.16.5) and DESeq2 (v1.14.1) to generate normalised expression

counts. The normalised counts for RNAseq (Figure 1) were calculated in edgeR. Low expressed

genes were removed (rowSum cpm <2 across SCR and SA1SP replicates), normalisation factors

were calculated using calcNormFactors and dispersions estimated using estimateDisp. The

edgeR volcano plot statistics were calculated using the exactTest and topTags functions. To

generate the normalised counts for RNAseq experiments required to calculate the log2FC GSEA

ranked lists, the FeatureCounts output for all experiments was combined into a single table and

read into DESeq2. A DESeq2 object was built using the function DESeqDataSetFromMatrix and

estimation of size factors and dispersions were calculated using the DEseq function. Normalised

counts were calculated using the 'counts' function. Low expressed genes (rowSum normalised

count <10 across all samples) were removed.

1009

GSEA

Broad Institute GSEAPreranked (v4.0.3) was used to determine the enrichment of curated

genesets within our RNA-seq data. For each sample a ranked list was generated with genes

ranked in descending order by their log2FC value using normalised expression scores from

DEseq2. Log2FC per gene was calculated between the KD and its respective SCR using the

following calculation: Log2(normalised_counts KD +1) - log2(normalised_counts SCR +1). In the

case of experiments with multiple KD replicates, the average log2 normalised count was used.

Three gene sets were assayed in this study, 'naïve pluripotency', 'primed pluripotency' and '2C

signatures'. The naïve and primed pluripotency gene sets were curated in-house from Fidalgo M

et al. (CSC, 2016) where genes were selected if they had ≥2 fold change. The naïve and primed

gene sets contained 661 and 580 genes respectively. The 2C signatures gene set (147 genes)

was obtained from Percharde M et al. (Cell, 2018). Gene sets were classed as having significant

enrichment if the p-value was $\leq$0.05 and the normalised enrichment score (NES) exceeded +/- 1.

1023

VAST-TOOLS

VAST-TOOLS was used to generate Percent Spliced In (PSI) scores, a statistic which represents

how often a particular exon is spliced into a transcript using the ratio between reads which include

and exclude said exon. Paired-end RNA-seq datasets were submitted to VAST-TOOLS (v2.1.3)

using the Mmu genome (Tapial J et al, Gen Res 2017). Briefly, reads are split into 50nt words

with a 25nt sliding window. The 50nt words are aligned to a reference genome using Bowtie to

obtain unmapped reads. These unmapped reads are then aligned to a set of predefined exon-

exon junction (EJJ) libraries allowing for the quantification of alternative exon events. The output

was further interrogated using a script which searches all hypothetical EEJ combinations between

potential donors and acceptors within Stag1. PSI scores could be obtained providing there was

at least a single read within our RNAseq data that supported one of these potential events. Some

datasets were combined to have enough reads for the analysis. See Table S1 for PSI values and

names of RNA-seq libraries used for analysis in Figure 3E and S4B.

1037

1038   **Quantifying sectioned Stag1**

1039   Stag1 was split into 5 sections; SATS, e1-e8, e12-e19, e20-e25, e26-e34. Using Kallisto

1040   (v0.46.1), raw RNAseq reads were used to quantify each section of Stag1. Kallisto was run in

1041   quant mode, using the –rf-stranded parameter, outputting a TPM per Stag1 section. A line plot

1042   was generated showing TPM in relative to UT.

1043

PacBio library, sequencing and analysis

ES cells were cultured in naïve 2i conditions and PolyA-enriched mRNAs were hybridized to a

custom Biotinylated oligonucleoltide probe set. Post-capture, mRNAs were amplified using the

Clontech SMARTer PCR cDNA Synthesis Kit with 9 cycles and used in the SMRTbell library prep

according to manufacturers instructions. The library was sequenced on the SMRTseq 2000

platform. PacBio reads were processed through the SMRTLINK v8.0.0 IsoSeq3 pipeline. 403,995

Circular consensus sequences (CCS) were generated using default parameters (--minPasses =

1, --min-rq = 0.8, CCS Polish = No). Further refining through lima (removal of adapters and correct

orientation of sequences), poly-A trimming and concatemer removal resulted in 265,106 full length

non-chimeric (FLNC) reads. FLNC reads were aligned to the mm10 genome using Minimap2 with

the following parameters (-ax splice, -uf, -k14).

1055

ChIP-seq Analysis

Previously published Stag1 Chromatin Immunoprecipitation-sequencing (ChIP-seq) datasets

from ES 2i cells (GSE126659, only Replicate 1 and 2 libraries) were trimmed using trim_galore

and aligned to mm10 using bowtie2. Peak detection was performed with MACS2 using uniquely

reads (MAPQ≥2). Peaks were overlapped with genomic features in a hierarchical manner

(promoters > exons > repeats > introns > intergenic), and overlap frequency was compared with

a randomly shuffled version of the peaks. To identify repeat families enriched for STAG1 peaks,

a previously described pipeline was used (Deniz O et al. Nat Comm, 2020) that compares family-

levels overlap frequency with that observed in 1,000 permutations of random peak shuffling.

Coverage profiles across specific TE families were generated using HOMER and including multi-

mapping reads (MAPQ<2).

1067

UMI-4C library preparation.

$1x10^7$ cells were fixed at RT for 10min in 1% formaldehyde and fixation was quenched with 0.125M

Glycine for 5min. Cells were then lysed on ice in 10ml Lysis Buffer (10mM NaCl, 10mM Tris-HCl

pH 8.0, 0.25% NP40, protease inhibitor) for 30min, followed by 10 strokes of douncing using a

tight pestle. Nuclei were pelleted, 8min 700 rcf, washed in 1ml 1.2X DpnII buffer in Protein LoBind

tubes (Eppendorf) and resuspended in 500 μl 1.2X DpnII buffer. **15ul of 10% SDS** was added

and incubated for 1hr at 37°C shaking at 650 rcf. **50ul of 20% TritonX** was added to quench the

SDS and incubated for 15 min at 37°C with shaking. **750U of DpnII was added and incubated**

overnight at 37°C **with interval shaking. The next morning, nuclei were** pelleted at 4°C by

650 rcf for 5 min and resuspended in 500μl 1X DpnII buffer. 500U DpnII was added and incubated

for an additional four hours. The nuclei were washed twice in 100 μl of 1X T4 Ligase Buffer and

resuspended in 200 μl Ligase Buffer. **6ul of T4 DNA Ligase was added and incubated for 3hr**

at 16°C. Nuclei were then pelleted, resuspended in 200 μl 1x fresh Ligase Buffer, 6μl of T4 DNA

Ligase added, and incubated overnight at 16°C. Samples were treated with 20μl of ProtK (NEB

Molecular Biology Grade), incubated for 3 hrs at 55°C and 5 hrs at 65°C to reverse crosslinks.

Samples were treated with **RNase A** (PureLink, Invitrogen) for 1 hr at 37°C and DNA was

extracted and precipitated overnight. For library preparation, 3x5μg of ligated DNA was sonicated

using Covaris (10% duty cycle, intensity 5, cycle burst 200, 70sec). Samples were end-repaired

using DNA PolII Klenow Large Fragment (NEB), A-tailed using Klenow (exo-) (NEB), and Illumina

indexed adapters ligated using Quick DNA Ligase (NEB). Reactions were denatured at 95°C for

3 min, placed on ice, and purified using 1.2X SizeSelect AmpPure beads to recover ssDNA.

Libraries were amplified using GoTaq (Promega), with 20 cycles for PCR1 and 15 cycles for

nested PCR2 on 50% material from 1st PCR. For custom UMI bait sequences, see Table S3.

1091

Hi-C and UMI-4C-seq analysis

Hi-C libraries were analysed as previously described (Barrington 2019). UMI-4C tracks were

processed using the 'umi4cPackage' pipeline (v0.0.0.9000) (Schwartzman, O et al. Nat Meth

2017). Briefly, raw reads are parsed through the UMI-4C pipeline, those reads containing the bait

and padding sequence are retained and de-multiplexed. Reads lacking the padding sequence are

considered non-specific and are removed from further analysis. Retained reads are split based

on a match to the restriction enzyme sequence to create a segmented fastq file. The first 10 bases

of read 2 are extracted and attached to the segments derived from each read pair. Mapping to

mm10 is done with Bowtie2. Read pairs that have reverse complement segments are mapped to

a restriction fragment ID, with the fragment ID, strand and distance from each end represented

within a fragment-chain table. UMI filtering is used to determine the number of molecules

supporting each ligation event. The resulting UMI-4C tracks are then imported into R, and data

from multiple bait replicates can be merged by summing the molecule counts per ligated fragment,

at which point contact intensity profiles and domainograms around the viewpoint can be

generated (see Figure 3).   The contact intensity profile represents the mean number of ligations

within a genomic window, with the resolution of the contact intensity profile being determined by

the window size (set to 15 here). The domainogram reports the mean contact per fend at a series

of window sizes, a stacked representation of contact intensity values in increasing window sizes

from 10 to 300 fragment ends, their colour can be used to identify peak locations. ES and NSC

contact profiles were compared after normalisation to correct for bias (see Schwartzman et al for

further details). For the compared profiles, the total molecule count for restriction fragment ends

for each are calculated at three ranges around the viewpoint. One profile is selected as a

reference and the second is scaled to the first using the ratio in total molecule counts between

the two profiles as the scaling factor. Below the contact profile is the profile resolution indicator,

which shows the number of fends required to include at least 15 UMI molecules. The darker the

colour, the larger the window size required. The domainogram at the bottom represents the log2

ratio between the domainogram values of the compared profiles and highlights locations where

ES has more contacts than NSC or vice versa.

1120

Supplementary titles and legends.

Figure S1. Stag1 is required for pluripotency, Related to Figure 1.

(A) Cartoon of the cohesin complex including the core trimer subunits of Smc1a, Smc3 and Rad21 complexed with either Stag1 or Stag2.

(B) Relative expression of Stag1 and Stag2 mRNA by qRT-PCR in 2i- or FCS-grown ES cells, EpiLCs and MEFs. Data is represented as mean ± SEM of two independent experiments and relative to Actin control expression.

(C) WCL from naïve (2i) ES and EpiLCs, sorted for cells in the G1 phase and analyzed by WB for levels of SA1 and SA2. Actin serves as a loading control.

(D) Relative expression of Stag1 mRNA by qRT-PCR in FCS- (left panel, n=20) or 2i-grown (right panel, n=19) ES cells upon treatment with si scr or si SA1. Whiskers and boxes indicate all and 50% of values respectively. Central line represents the median.

(E) Cell cycle analysis of Hoechst-stained 2i ES cells after treatment with siScr or siSA1 siRNAs for 24hrs. Shown are the percentages of cells in G1 or G2 phases. These are the same cells that were used for the RNA-sequencing experiments shown in Figure 1F.

(F) MFI of Stag1 protein assessed by IF in 2i ES cells treated with si scr and si SA1. Cells were stained for SA1 and counterstained with DAPI. Data is from the second biological replicate (replicate 1 is in Fig 1E), n>100 cells/condition **** p<0.0005.

(G) Enrichment score (ES) plots from GSEA using Naïve or Primed gene sets as in Figure 1G and RNA-seq data from the other two siSA1 treated ES cell samples.

(H) AP+ colonies in ES cells (purple), as a percentage of all colonies (pink and purple) treated with siscr and si SA1. Data are from three independent biological replicates and merged. See Fig 1H for the individual experiments.

(I-K) CRISPR/Cas9 targeting of the C-terminus of endogenous Stag1. (I) Schematic of the targeted locus and location of primers used for genotyping. (J) Shown are four ES cell clones representing integration into both alleles, one allele and a wildtype clone. (K) IF using the v5 tag in a homozygote clonal line treated with dTAG for 16 hrs shows complete loss of the NeonGreen signal.

Figure S2. Stag1 associates with heterochromatin, Related to Figure 2.

(A, B) Confocal IF images in dox-inducible full-length SA1-GFP ES cells (SA1-FL-GFP)
stained with (A) GFP and (B) HP1a. Nuclei were counterstained with DAPI. NB,
colocalization of SA1-GFP with DAPI-dense foci and HP1a.

(C) Replicate experiment for co-Immunoprecipitation of endogenous SA1 with Smc3,
HP1a and Nucleolin in ES cells.

(D) Enrichment of Stag1 ChIP at additional LTR and DNA transposon elements.

(E) WCL from ES treated with the siRNA panel and analyzed by WB for levels of SA1,
H3K9me3 and K3K4me3. Tubulin serves as a loading control.

(F) Global chromatin accessibility as detected by DNase I digestion of genomic DNA in
siScr or siSA1 ES cells. NB, In two of six biological replicates, we observed increased
compaction upon SA1 loss, as is shown here.

1163

Figure S3. Transcription-Regulatory Control of Stag1 in ESC, Related to Figure 3.

(A) Aligned Stag1 transcript variants identified from 5'RACE in Figure 3A, B. Arrows refer
to the bands on the RACE gels which were cloned and sequenced. NB, the diversity of
skipping events that all result in a functional loss of the 5' end of Stag1.

(B) Over-exposure of the 5'RACE gel shown in Figure 3B to better show the small RACE
products.

(C) Close-up of the 5' RACE sequence that identified a new TSS at exon 7 spliced directly
to a sequence in trans carrying regulatory elements.

(D) Close-up of the 3' RACE sequence that identified a new alternative TTS in intron 25
(sequence shown in dark blue).

(E) Initial PCR screen in ES 2i and MEFs using various combinations of forward (5')
primers (SATS, canonical TSS, Alt exon 1 TSS) and reverse (3') primers (canonical TTS,
Alt intron 25 TTS). NB. SATS is only expressed in ES cells; canonical, full-length Stag1
is more expressed in ES compared to MEFs; and the alternative intron 25 TTS is most
often expressed with a canonical TSS.

(F) Top, Strategy for mRNA capture of cohesin genes for Pac-Bio long read RNA-
sequencing. Bottom, full-length transcripts sequenced on the PacBio platform includes
many isoforms already discovered using RACE and PCR cloning methods.

(G) Chromatin Immunoprecipitation of SA1 in ES and EpiLCs.

1183

Figure S4. Fluctuations in Stag1 isoforms skews cell fates, Related to Figure 4.

(A) Relative expression of Stag1 mRNA by qRT-PCR in FCS- (leftmost and rightmost
panels, n=7) or 2i-grown (middle panel, n=6) ES cells upon si scr or the si SA1 panel.
Whiskers and boxes as before. NB, all siRNAs knockdown Stag1 levels to a similar extent
with the exception of esiRNA SATS which reduced Stag1 by ~40-50%.

(B) Percent Spliced In (PSI) calculations based on VAST-Tools analysis of RNA-seq from
ES cells treated with the siRNA panel. Data are shown relative to untreated ES cells.

(C) Relative expression of Nanog mRNA by qRT-PCR in FCS-grown ES cells upon si scr
or the si SA1 panel (n=13). Whiskers and boxes as before. NB, the modest, but different
influence of the 5p and the 3p KDs on Nanog levels.

(D) AP+ colonies in ES cells (purple), as a percentage of all colonies (pink and purple)
treated with the siRNA panel. Data are from two independent biological replicates and
merged. See Fig 4H for the individual experiments.

1197

Figure S5. Stag1 N-terminus represses heterochromatin and reprograms cells,
Related to Figure 5.

(A) Confocal IF images of GFP and H3K9me3 in ES cells treated with siRNA panel. Nuclei
were counterstained with DAPI.

(B) Imaris quantification of the volume of H3K9me3 foci per cell from (A). Box plots and
statistical analysis were done as before. Data are from two biological replicates,
n>50/condition. ****$p<0.0005$.

(C) Global chromatin accessibility as detected by DNase I digestion of genomic DNA in
siScr or siSA1 3p or 5p ES cells. NB. There was no effect on accessibility in the 3p KD
but a consistent decompaction observed in the 5p KD.

(D, E) Relative expression of (C) MERVL and (D) genes associated with the 2C-L state
by qRT-PCR in 2i-grown ES cells after 72hr of treatment with si scr or the si SA1 panel.
Data are represented as mean +/- SEM from n=3 biological replicates. NB, the effect of
the 5p KD on the expression of these hallmark 2C-L genes/repeats is much more
significant than in the FCS-grown ES cells shown in Figure 5C, D.

(F) Enrichment score (ES) plots from GSEA using 2C gene sets as in Figure 5E and
RNA-seq data from the different siRNA treated ES cell samples.

1215

Figure S6. Stag1 deregulates nascent rRNA expression, Related to Figure 6.

(A) WCL from ES cells treated with the siRNA panel and analyzed by WB for global levels
of Nucleolin. Tubulin serves as a loading control.

(B) Top, cartoon of the consensus Mus musculus ribosomal DNA (rDNA) (GenBank:
BK000964.3), showing the ribosomal genes and the intergenic spacer (IGS) region which
contains several SINE elements (Red, B2_Mm2; Green, B3). Bottom, Stag1 ChIP
replicates aligned to this region. NB, possible Stag1 binding to the B3 elements in the
IGS.

1224

REFERENCES

Ahmed, K., Dehghani, H., Rugg-Gunn, P., Fussner, E., Rossant, J., and Bazett-Jones, D.P.
(2010). Global chromatin architecture reflects pluripotency and lineage commitment in the early
mouse embryo. PLoS ONE *5*, e10531.

Allshire, R.C., and Madhani, H.D. (2018). Ten principles of heterochromatin formation and
function. Nat. Rev. Mol. Cell Biol. *19*, 229–244.

Banani, S.F., Lee, H.O., Hyman, A.A., and Rosen, M.K. (2017). Biomolecular condensates:
organizers of cellular biochemistry. Nat. Rev. Mol. Cell Biol. *18*, 285–298.

Barrington, C., Georgopoulou, D., Pezic, D., et al. (2019) Enhancer accessibility and CTCF
occupancy underlie asymmetric TAD architecture and cell type specific genome topology.
Nature Communications.

Bickmore, W.A., and van Steensel, B. (2013). Genome architecture: domain organization of
interphase chromosomes. Cell *152*, 1270–1284.

Bisht, K.K., Daniloski, Z., and Smith, S. (2013). SA1 binds directly to DNA through its unique
AT-hook to promote sister chromatid cohesion at telomeres. J. Cell. Sci. *126*, 3493–3503.

Bonev, B., and Cavalli, G. (2016). Organization and function of the 3D genome. Nat. Rev.
Genet. *17*, 661–678.

Bonev, B., Cohen, N.M., Szabo, Q., Fritsch, L., Papadopoulos, G.L., Lubling, Y., Xu, X., Lv, X.,
Hugnot, J.-P., Tanay, A., et al. (2017). Multiscale 3D Genome Rewiring during Mouse Neural
Development. Cell *171*, 557.e1–557.e24.

Borsos, M., and Torres-Padilla, M.-E. (2016). Building up the nucleus: nuclear organization in
the establishment of totipotency and pluripotency during mammalian development. Genes Dev.
30, 611–621.

Buszczak, M., Signer, R.A.J., and Morrison, S.J. (2014). Cellular Differences in Protein
Synthesis Regulate Tissue Homeostasis. Cell *159*, 242–251.

Canudas, S., and Smith, S. (2009). Differential regulation of telomere and centromere cohesion
by the Scc3 homologues SA1 and SA2, respectively, in human cells. J Cell Biol *187*, 165–173.

Cardozo Gizzi, A.M., Cattoni, D.I., Fiche, J.-B., Espinola, S.M., Gurgo, J., Messina, O.,
Houbron, C., Ogiyama, Y., Papadopoulos, G.L., Cavalli, G., et al. (2019). Microscopy-Based
Chromosome Conformation Capture Enables Simultaneous Visualization of Genome
Organization and Transcription in Intact Organisms. Molecular Cell *74*, 212–222.e215.

Cho, W.-K., Spille, J.-H., Hecht, M., Lee, C., Li, C., Grube, V., and Cisse, I.I. (2018). Mediator
and RNA polymerase II clusters associate in transcription-dependent condensates. Science
361, 412–415.

Cuadrado, A., and Losada, A. (2020). Specialized functions of cohesins STAG1 and STAG2 in

3D genome architecture. Curr. Opin. Genet. Dev. *61*, 9–16.

Cuadrado, A., Giménez-Llorente, D., Kojic, A., Rodriguez-Corsino, M., Cuartero, Y., Martín-
Serrano, G., Gomez-Lopez, G., Marti-Renom, M.A., and Losada, A. (2019). Specific
Contributions of Cohesin-SA1 and Cohesin-SA2 to TADs and Polycomb Domains in Embryonic
Stem Cells. Cell Rep *27*, 3500–3510.e3504.

Cuartero, S., Weiss, F.D., Dharmalingam, G., Guo, Y., Ing-Simmons, E., Masella, S., Robles-
Rebollo, I., Xiao, X., Wang, Y.-F., Barozzi, I., et al. (2018). Control of inducible gene expression
links cohesin to hematopoietic progenitor self-renewal and differentiation. Nat Immunol *19*, 932–
941.

Dekker, J., and Mirny, L. (2016). The 3D Genome as Moderator of Chromosomal
Communication. Cell *164*, 1110–1121.

Deniz, Ö., Ahmed, M., Todd, C.D., Rio-Machin, A., Dawson, M.A., and Branco, M.R. (2020).
Endogenous retroviruses are a source of enhancers with oncogenic potential in acute myeloid
leukaemia. Nature Communications *11*, 3506–3514.

Dixon, J.R., Selvaraj, S., Yue, F., Kim, A., Li, Y., Shen, Y., Hu, M., Liu, J.S., and Ren, B. (2012).
Topological domains in mammalian genomes identified by analysis of chromatin interactions.
Nature *485*, 376–380.

Eckersley-Maslin, M.A., Svensson, V., Krueger, C., Stubbs, T.M., Giehr, P., Krueger, F.,
Miragaia, R.J., Kyriakopoulos, C., Berrens, R.V., Milagre, I., et al. (2016). MERVL/Zscan4
Network Activation Results in Transient Genome-wide DNA Demethylation of mESCs. Cell Rep
17, 179–192.

Feng, G., Tong, M., Xia, B., Luo, G.Z., Wang, M., Xie, D., Wan, H., Zhang, Y., Zhou, Q., and
Wang, X.J. (2016). Ubiquitously expressed genes participate in cell-specific functions via
alternative promoter usage. EMBO Rep. *17*, 1304–1313.

Finn, E.H., Pegoraro, G., Brandão, H.B., Valton, A.-L., Oomen, M.E., Dekker, J., Mirny, L., and
Misteli, T. (2019). Extensive Heterogeneity and Intrinsic Variation in Spatial Genome
Organization. Cell *176*, 1502–1515.e1510.

Fiszbein, A., and Kornblihtt, A.R. (2017). Alternative splicing switches: Important players in cell
differentiation. Bioessays *39*, 1600157.

Gabut, M., Samavarchi-Tehrani, P., Wang, X., Slobodeniuc, V., O'Hanlon, D., Sung, H.-K.,
Alvarez, M., Talukder, S., Pan, Q., Mazzoni, E.O., et al. (2011). An alternative splicing switch
regulates embryonic stem cell pluripotency and reprogramming. Cell *147*, 132–146.

Guelen, L., Pagie, L., Brasset, E., Meuleman, W., Faza, M.B., Talhout, W., Eussen, B.H., de
Klein, A., Wessels, L., de Laat, W., et al. (2008). Domain organization of human chromosomes
revealed by mapping of nuclear lamina interactions. Nature *453*, 948–951.

Haarhuis, J.H.I., van der Weide, R.H., Blomen, V.A., Yáñez-Cuna, J.O., Amendola, M., van
Ruiten, M.S., Krijger, P.H.L., Teunissen, H., Medema, R.H., van Steensel, B., et al. (2017). The
Cohesin Release Factor WAPL Restricts Chromatin Loop Extension. Cell *169,* 693–707.e14.

Hadjur, S., Williams, L.M., Ryan, N.K., Cobb, B.S., Sexton, T., Fraser, P., Fisher, A.G., and
Merkenschlager, M. (2009). Cohesins form chromosomal cis-interactions at the developmentally
regulated IFNG locus. Nature *460*, 410–413.

Hara, K., Zheng, G., Qu, Q., Liu, H., Ouyang, Z., Chen, Z., Tomchick, D.R., and Yu, H. (2014).
Structure of cohesin subcomplex pinpoints direct shugoshin-Wapl antagonism in centromeric
cohesion. Nature Publishing Group *21*, 864–870.

Hendrickson, P.G., Doráis, J.A., Grow, E.J., Whiddon, J.L., Lim, J.-W., Wike, C.L., Weaver,
B.D., Pflueger, C., Emery, B.R., Wilcox, A.L., et al. (2017). Conserved roles of mouse DUX and
human DUX4 in activating cleavage-stage genes and MERVL/HERVL retrotransposons. Nat.
Genet. *49*, 925–934.

Horsfield, J.A., Anagnostou, S.H., Hu, J.K.-H., Cho, K.H.Y., Geisler, R., Lieschke, G., Crosier,
K.E., and Crosier, P.S. (2007). Cohesin-dependent regulation of Runx genes. Development
134, 2639–2649.

Hung, S.S.C., Wong, R.C.B., Sharov, A.A., Nakatake, Y., Yu, H., and Ko, M.S.H. (2013).
Repression of global protein synthesis by Eif1a-like genes that are expressed specifically in the
two-cell embryos and the transient Zscan4-positive state of embryonic stem cells. DNA Res *20*,
391–402.

Ishiuchi, T., Enriquez-Gasca, R., Mizutani, E., Bošković, A., Ziegler-Birling, C., Rodriguez-
Terrones, D., Wakayama, T., Vaquerizas, J.M., and Torres-Padilla, M.-E. (2015). Early
embryonic-like cells are induced by downregulating replication-dependent chromatin assembly.
Nature Publishing Group *22*, 662–671.

Jachowicz, J.W., Bing, X., Pontabry, J., Bošković, A., Rando, O.J., and Torres-Padilla, M.-E.
(2017). LINE-1 activation after fertilization regulates global chromatin accessibility in the early
mouse embryo. Nat. Genet. *49*, 1502–1510.

Kagey, M.H., Newman, J.J., Bilodeau, S., Zhan, Y., Orlando, D.A., van Berkum, N.L., Ebmeier,
C.C., Goossens, J., Rahl, P.B., Levine, S.S., et al. (2010). Mediator and cohesin connect gene
expression and chromatin architecture. Nature *467*, 430–435.

Kline, A.D., Moss, J.F., Selicorni, A., Bisgaard, A.-M., Deardorff, M.A., Gillett, P.M., Ishman,
S.L., Kerr, L.M., Levin, A.V., Mulder, P.A., et al. (2018). Diagnosis and management of Cornelia
de Lange syndrome: first international consensus statement. Nat. Rev. Genet. *19*, 649–666.

Kojic, A., Cuadrado, A., De Koninck, M., Giménez-Llorente, D., Rodriguez-Corsino, M., Gomez-
Lopez, G., Le Dily, F., Marti-Renom, M.A., and Losada, A. (2018). Distinct roles of cohesin-SA1
and cohesin-SA2 in 3D chromosome organization. Nature Publishing Group *25*, 496–504.

Kresoja-Rakic, J., and Santoro, R. (2019). Nucleolus and rRNA Gene Chromatin in Early
Embryo Development. Trends Genet. *35*, 868–879.

Larson, A.G., Elnatan, D., Keenen, M.M., Trnka, M.J., Johnston, J.B., Burlingame, A.L., Agard,
D.A., Redding, S., and Narlikar, G.J. (2017). Liquid droplet formation by HP1α suggests a role
for phase separation in heterochromatin. Nature *547*, 236–240.

Lehalle, D., Mosca-Boidron, A.-L., Begtrup, A., Boute-Benejean, O., Charles, P., Cho, M.T.,
Clarkson, A., Devinsky, O., Duffourd, Y., Duplomb-Jego, L., et al. (2017). STAG1 mutations
cause a novel cohesinopathy characterised by unspecific syndromic intellectual disability. J Med
Genet *54*, 479–488.

Leiserson, M.D.M., Vandin, F., Wu, H.-T., Dobson, J.R., Eldridge, J.V., Thomas, J.L.,
Papoutsaki, A., Kim, Y., Niu, B., McLellan, M., et al. (2015). Pan-cancer network analysis
identifies combinations of rare somatic mutations across pathways and protein complexes. Nat.
Genet. *47*, 106–114.

Li, Y., Haarhuis, J.H.I., Sedeño Cacciatore, Á., Oldenkamp, R., van Ruiten, M.S., Willems, L.,
Teunissen, H., Muir, K.W., de Wit, E., Rowland, B.D., et al. (2020). The structural basis for
cohesin-CTCF-anchored loops. Nature *578*, 472–476.

Macfarlan, T.S., Gifford, W.D., Driscoll, S., Lettieri, K., Rowe, H.M., Bonanomi, D., Firth, A.,
Singer, O., Trono, D., and Pfaff, S.L. (2012). Embryonic stem cell potency fluctuates with
endogenous retrovirus activity. Nature *487*, 57–63.

Martin, C., Beaujean, N., Brochard, V., Audouard, C., Zink, D., and Debey, P. (2006). Genome
restructuring in mouse embryos during reprogramming and early development. Dev. Biol. *292*,
317–332.

Mateo, L.J., Murphy, S.E., Hafner, A., Cinquini, I.S., Walker, C.A., and Boettiger, A.N. (2019).
Visualizing DNA folding and RNA in embryos at single-cell resolution. Nature *568*, 1–24.

Meshorer, E., Yellajoshula, D., George, E., Scambler, P.J., Brown, D.T., and Misteli, T. (2006).
Hyperdynamic plasticity of chromatin proteins in pluripotent embryonic stem cells. Dev. Cell *10*,
105–116.

Mishiro, T., and Tsutsumi, S. (2009). Architectural roles of multiple chromatin insulators at the
human apolipoprotein gene cluster. Embo J. *28*, 1234–1245.

Misulovin, Z., Schwartz, Y.B., Li, X.-Y., Kahn, T.G., Gause, M., MacArthur, S., Fay, J.C., Eisen,
M.B., Pirrotta, V., Biggin, M.D., et al. (2007). Association of cohesin and Nipped-B with
transcriptionally active regions of the Drosophila melanogaster genome. Chromosoma *117*, 89–
102.

Mootha, V.K., Lindgren, C.M., Eriksson, K.-F., Subramanian, A., Sihag, S., Lehar, J.,
Puigserver, P., Carlsson, E., Ridderstråle, M., Laurila, E., et al. (2003). PGC-1alpha-responsive
genes involved in oxidative phosphorylation are coordinately downregulated in human diabetes.
Nat. Genet. *34*, 267–273.

Nabet, B., Roberts, J.M., Buckley, D.L., Paulk, J., Dastjerdi, S., Yang, A., Leggett, A.L., Erb,
M.A., Lawlor, M.A., Souza, A., et al. (2018). The dTAG system for immediate and target-
specific protein degradation. Nature Chemical Biology *14*, 1–16.

Nagano, T., Lubling, Y., Várnai, C., Dudley, C., Leung, W., Baran, Y., Mendelson Cohen, N.,
Wingett, S., Fraser, P., and Tanay, A. (2017). Cell-cycle dynamics of chromosomal organization
at single-cell resolution. Nature *547*, 61–67.

Németh, A., Conesa, A., Santoyo-Lopez, J., Medina, I., Montaner, D., Péterfia, B., Solovei, I., Cremer, T., Dopazo, J., and Längst, G. (2010). Initial genomics of the human nucleolus. PLoS Genet. *6*, e1000889.

Novo, C.L., Tang, C., Ahmed, K., Djuric, U., Fussner, E., Mullin, N.P., Morgan, N.P., Hayre, J., Sienerth, A.R., Elderkin, S., et al. (2016). The pluripotency factor Nanog regulates pericentromeric heterochromatin organization in mouse embryonic stem cells. Genes Dev. *30*, 1101–1115.

Nozaki, T., Imai, R., Tanbo, M., Nagashima, R., Tamura, S., Tani, T., Joti, Y., Tomita, M., Hibino, K., Kanemaki, M.T., et al. (2017). Dynamic Organization of Chromatin Domains Revealed by Super-Resolution Live-Cell Imaging. Molecular Cell *67*, 282–293.e287.

Obradovic, Z., Peng, K., Vucetic, S., Radivojac, P., Brown, C.J., and Dunker, A.K. (2003). Predicting intrinsic disorder from amino acid sequence. Proteins *53 Suppl 6*, 566–572.

Orgil, O., Matityahu, A., Eng, T., Guacci, V., Koshland, D., and Onn, I. (2015). A conserved domain in the scc3 subunit of cohesin mediates the interaction with both mcd1 and the cohesin loader complex. PLoS Genet. *11*, e1005036.

Padeken, J., and Heun, P. (2014). Nucleolus and nuclear periphery: velcro for heterochromatin. Curr. Opin. Cell Biol. *28*, 54–60.

Parelho, V., Hadjur, S., Spivakov, M., Leleu, M., Sauer, S., Gregson, H.C., Jarmuz, A., Canzonetta, C., Webster, Z., Nesterova, T., et al. (2008). Cohesins functionally associate with CTCF on mammalian chromosome arms. Cell *132*, 422–433.

Percharde, M., Lin, C.-J., Yin, Y., Guan, J., Peixoto, G.A., Bulut-Karslioglu, A., Biechele, S., Huang, B., Shen, X., and Ramalho-Santos, M. (2018). A LINE1-Nucleolin Partnership Regulates Early Development and ESC Identity. Cell *174*, 391–405.e19.

Phillips-Cremins, J.E., Sauria, M.E.G., Sanyal, A., Gerasimova, T.I., Lajoie, B.R., Bell, J.S.K., Ong, C.-T., Hookway, T.A., Guo, C., Sun, Y., et al. (2013). Architectural protein subclasses shape 3D organization of genomes during lineage commitment. Cell *153*, 1281–1295.

Quinodoz, S.A., Ollikainen, N., Tabak, B., Palla, A., Schmidt, J.M., Detmar, E., Lai, M.M., Shishkin, A.A., Bhat, P., Takei, Y., et al. (2018). Higher-Order Inter-chromosomal Hubs Shape 3D Genome Organization in the Nucleus. Cell *174*, 744–757.e24.

Rao, S.S.P., Huang, S.-C., Hilaire, B.G.S., Engreitz, J.M., Perez, E.M., Kieffer-Kwon, K.-R., Sanborn, A.L., Johnstone, S.E., Bascom, G.D., Bochkov, I.D., et al. (2017). Cohesin Loss Eliminates All Loop Domains. Cell *171*, 305–309.e324.

Rao, S.S.P., Huntley, M.H., Durand, N.C., Stamenova, E.K., Bochkov, I.D., Robinson, J.T., Sanborn, A.L., Machol, I., Omer, A.D., Lander, E.S., et al. (2014). A 3D map of the human genome at kilobase resolution reveals principles of chromatin looping. Cell *159*, 1665–1680.

Remeseiro, S., Cuadrado, A., Carretero, M., Martínez, P., Drosopoulos, W.C., Cañamero, M., Schildkraut, C.L., Blasco, M.A., and Losada, A. (2012a). Cohesin-SA1 deficiency drives aneuploidy and tumourigenesis in mice due to impaired replication of telomeres. Embo J. *31*,

2076–2089.

Remeseiro, S., Cuadrado, A., López, G.G., Pisano, D.G., and Losada, A. (2012b). A unique role
of cohesin-SA1 in gene regulation and development. Embo J. *31*, 2090–2102.

Ricci, M.A., Manzo, C., García-Parajo, M.F., Lakadamyali, M., and Cosma, M.P. (2015).
Chromatin fibers are formed by heterogeneous groups of nucleosomes in vivo. Cell *160*, 1145–
1158.

Rodriguez, J., Ren, G., Day, C.R., Zhao, K., Chow, C.C., and Larson, D.R. (2019). Intrinsic
Dynamics of a Human Gene Reveal the Basis of Expression Heterogeneity. Cell *176*, 213–
226.e218.

Romero-Pérez, L., Surdez, D., Brunet, E., Delattre, O., and Grünewald, T.G.P. (2019). STAG
Mutations in Cancer. Trends Cancer *5*, 506–520.

Sabari, B.R., Dall'Agnese, A., Boija, A., Klein, I.A., Coffey, E.L., Shrinivas, K., Abraham, B.J.,
Hannett, N.M., Zamudio, A.V., Manteiga, J.C., et al. (2018). Coactivator condensation at super-
enhancers links phase separation and gene control. Science *361*, eaar3958–13.

Salomonis, N., Schlieve, C.R., Pereira, L., Wahlquist, C., Colas, A., Zambon, A.C., Vranizan, K.,
Spindler, M.J., Pico, A.R., Cline, M.S., et al. (2010). Alternative splicing regulates mouse
embryonic stem cell pluripotency and differentiation. Proc. Natl. Acad. Sci. U.S.a. *107*, 10514–
10519.

Schlesinger, S., and Meshorer, E. (2019). Open Chromatin, Epigenetic Plasticity, and Nuclear
Organization in Pluripotency. Dev. Cell *48*, 135–150.

Schwalie, P.C., Ward, M.C., Cain, C.E., Faure, A.J., Gilad, Y., Odom, D.T., and Flicek, P.
(2013). Co-binding by YY1 identifies the transcriptionally active, highly conserved set of CTCF-
bound regions in primate genomes. Genome Biol. *14*, R148–15.

Schwarzer, W., Abdennur, N., Goloborodko, A., Pekowska, A., Fudenberg, G., Loe-Mie, Y.,
Fonseca, N.A., H Haering, C., Mirny, L., and Spitz, F. (2017). Two independent modes of
chromatin organization revealed by cohesin removal. Nature *551*, 51–56.

Seitan, V.C., Faure, A.J., Zhan, Y., McCord, R.P., Lajoie, B.R., Ing-Simmons, E., Lenhard, B.,
Giorgetti, L., Heard, E., Fisher, A.G., et al. (2013). Cohesin-based chromatin interactions enable
regulated gene expression within preexisting architectural compartments. Genome Res. *23*,
2066–2077.

Soardi, F.C., Machado-Silva, A., Linhares, N.D., Zheng, G., Qu, Q., Pena, H.B., Martins,
T.M.M., Vieira, H.G.S., Pereira, N.B., Melo-Minardi, R.C., et al. (2017). Familial STAG2 germline
mutation defines a new human cohesinopathy. NPJ Genom Med *2*, 7–11.

Sofueva, S., Yaffe, E., Chan, W.-C., Georgopoulou, D., Rudan, M.V., Mira-Bontenbal, H.,
Pollard, S.M., Schroth, G.P., Tanay, A., and Hadjur, S. (2013). Cohesin-mediated interactions
organize chromosomal domain architecture. Embo J. *32*, 3119–3129.

Strom, A.R., Emelyanov, A.V., Mir, M., Fyodorov, D.V., Darzacq, X., and Karpen, G.H. (2017).

Phase separation drives heterochromatin domain formation. Nature *547*, 241–245.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A.,
Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene set enrichment
analysis: a knowledge-based approach for interpreting genome-wide expression profiles. Proc
Natl Acad Sci USA *102*, 15545–15550.

Tapial, J., Ha, K.C.H., Sterne-Weiler, T., Gohr, A., Braunschweig, U., Hermoso-Pulido, A.,
Quesnel-Vallières, M., Permanyer, J., Sodaei, R., Marquez, Y., et al. (2017). An atlas of
alternative splicing profiles and functional associations reveals new regulatory programs and
genes that simultaneously express multiple major isoforms. Genome Res. *27*, 1759–1768.

Vietri Rudan, M., Barrington, C., Henderson, S., Ernst, C., Odom, D.T., Tanay, A., and Hadjur,
S. (2015). Comparative Hi-C reveals that CTCF underlies evolution of chromosomal domain
architecture. CellReports *10*, 1297–1309.

Viny, A.D., Bowman, R.L., Liu, Y., Lavallée, V.-P., Eisman, S.E., Xiao, W., Durham, B.H.,
Navitski, A., Park, J., Braunstein, S., et al. (2019). Cohesin Members Stag1 and Stag2 Display
Distinct Roles in Chromatin Accessibility and Topological Control of HSC Self-Renewal and
Differentiation. Cell Stem Cell *25*, 682–696.e688.

Wagner, A., Regev, A., and Yosef, N. (2016). Revealing the vectors of cellular identity with
single-cell genomics. Nature Publishing Group *34*, 1145–1160.

Wendt, K.S., Yoshida, K., Itoh, T., Bando, M., Koch, B., Schirghuber, E., Tsutsumi, S., Nagae,
G., Ishihara, K., Mishiro, T., et al. (2008). Cohesin mediates transcriptional insulation by
CCCTC-binding factor. Nature *451*, 796–801.

Whyte, W.A., Orlando, D.A., Hnisz, D., Abraham, B.J., Lin, C.Y., Kagey, M.H., Rahl, P.B., Lee,
T.I., and Young, R.A. (2013). Master transcription factors and mediator establish super-
enhancers at key cell identity genes. Cell *153*, 307–319.

Winters, T., McNicoll, F., and Jessberger, R. (2014). Meiotic cohesin STAG3 is required for
chromosome axis formation and sister chromatid cohesion. Embo J. *33*, 1256–1270.

Wutz, G., Várnai, C., Nagasaka, K., Cisneros, D.A., Stocsits, R.R., Tang, W., Schoenfelder, S.,
Jessberger, G., Muhar, M., Hossain, M.J., et al. (2017). Topologically associating domains and
chromatin loops depend on cohesin and are regulated by CTCF, WAPL, and PDS5 proteins.
Embo J. *36*, 3573–3599.

Xiao, T., Wallace, J., and Felsenfeld, G. (2011). Specific sites in the C terminus of CTCF
interact with the SA2 subunit of the cohesin complex and are required for cohesin-dependent
insulation activity. Mol. Cell. Biol. *31*, 2174–2183.

Yuan, B., Neira, J., Pehlivan, D., Santiago-Sim, T., Song, X., Rosenfeld, J., Posey, J.E., Patel,
V., Jin, W., Adam, M.P., et al. (2019). Clinical exome sequencing reveals locus heterogeneity
and phenotypic variability of cohesinopathies. Genet Med *21*, 663–675.

Yusufzai, T.M., Tagami, H., Nakatani, Y., and Felsenfeld, G. (2004). CTCF tethers an insulator
to subnuclear sites, suggesting shared insulator mechanisms across species. Molecular Cell *13*,

291–298.

Zuin, J., Dixon, J.R., van der Reijden, M.I.J.A., Ye, Z., Kolovos, P., Brouwer, R.W.W., van de
Corput, M.P.C., van de Werken, H.J.G., Knoch, T.A., van IJcken, W.F.J., et al. (2014). Cohesin
and CTCF differentially affect chromatin architecture and gene expression in human cells. Proc.
Natl. Acad. Sci. U.S.a. *111*, 996–1001.

1492

A esiRNA SATS | SmartPool (SP) | ex4 (5pKD) | Stag1 (SA1) gene | ex 21 | ex22/23 | ex 29 | ex 30 | 3pKD | chr9 mm10

B SA1 rel expression (ES-FCS) — UT, si scr, siSA1 SP, siSA1 3p, siSA1 5p, esi SATS

C SA1^NG-FKBP — UT, si scr, si SA1 SP, esi SATS, si SA1 5p, si SA1 3p; SA1 (150 kDa), Tubulin

D TPM rel to UT — si scr, siSA1 SP, siSA1 3p, siSA1 5p; SATS, ex1-8, ex12-19, ex20-25, ex26-34; Sectioned SA1: 5' / middle / 3'

E 5' RACE — wt ES, siSA1 5p; 3' RACE — si scr, si SA1 SP, si SA1 3p, si SA1 5p

F SA1^NG-FKBP — +DMSO, +dTAG; inp, SA1 IP, IgG

G NAIVE PLURIPOTENCY / PRIMED PLURIPOTENCY
- esi SATS: NES=-1.15 / NES=1.17
- siSA1 3p: NES=-1.69 / NES=1.54
- siSA1 5p: NES=1.56 / NES=-1.28
- UP in Stag1 KD / DOWN in Stag1 KD

H si scr, siSA1 SP, esi SATS, siSA1 3p, siSA1 5p; Exp1, Exp2, Exp3

A DAPI | GFP | H3K9me3 | merge

wt | SA1^FL | SA1^Δe5 | SA1^ΔCterm

B H3K9me3 foci volume/cell (μm³)
wt (GFP-) | SA1^FL | SA1^ΔN | SA1^ΔC

C MERVL | LINE1-T | IAPEz — relative expression

D Gm6763 | Dux — relative expression
UT | si scr | siSA1 SP | siSA1 3p | siSA1 5p

E SA1 3'KD | SA1 5'KD — 2C GENE SET Enrichment Score (ES)
NES=-1.33 | NES=1.53

F MERVL-GFP
si scr 0.085% | si Luc 0.12% | si Chaf1 1.03% | siSA1 SP 0.27% | siSA1 3p 0.13% | siSA1 5p 0.82% | siSA1 5p + si Chaf1 1.87%

G MERVL-GFP+ fold change rel to Chaf1
UT | si scr | si Chaf1 | siSA1 SP | siSA1 3p | siSA1 5p | siSA1 5p +Chaf1

H DuxHA / MERVL-GFP — GFP- | GFP+

I
GGACACTGACATGGACTGAAGGAGTAGAAAAGCTCTTGCTCTAGCTCCATCGCCTGCTTGCTGTGTGAGACTGAGTAACTGCTAGATCCTTGGACTTCCATTCACAGCTGCGACTGAACAATTGTTGGGAATTGGGCTGCCGACTGACAGTGGTGACTACCCCCTTACATGCCTGGTCCTCAGTGGAAAAAATTTCGTTCCAATTTTTGTGAATTTATTGGAGTCCTGATTCGACAGTGTCAATATAGCATAATTTATGATGAATATATGATGGACACCGTAATTTCCCTTTTGACTGGTTTGTCAGACTCCCAAGTCAGAGCTTTTAGGCATACAAGTACCCTTGCTGCAATGAAGCTGATGCTGCTCTGGTGAATGTTGCTTTAAACCTCAGTATTCATCAAGATAATACACAGAGGCAATATGAAGCTGAGAGAAATAAAATGATTGGGAAGAGAGCCAATGAAAGACTGGAGTTACTACTTCAGAAACGTAAAGAG

- 5'GeneRacer oligo (green)
- MT2_MERVL (red)
- Stag1 exon 7 + exon 8 (teal)

J SA1 — input | IP | IgG
Stag1 | Nucleolin | Trim28