Neurological mechanism of sensory deficits after developmental exposure to non-dioxin-like polychlorinated biphenyls (PCBs)

Nadja R. Brun[1], Jennifer M. Panlilio[1], Kun Zhang[2,3], Yanbin Zhao[2,3], Evgeny Ivashkin[4,5], John J. Stegeman[1], Jared V. Goldstone[1]

[1] Department of Biology, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

[2] School of Environmental Science and Engineering, Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai 200240, China

[3] Shanghai Institute of Pollution Control and Ecological Security, Shanghai 200092, China

[4] Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA, USA

[5] A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

Address correspondence to Nadja Brun: nbrun@whoi.edu

Abstract

The most abundant polychlorinated biphenyl (PCB) congeners found in the environment and in humans are the *ortho*-substituted, non-dioxin-like (NDL) congeners, especially PCB153. While evidence indicates that exposure to PCB153 and other NDL-PCBs in early vertebrate development can contribute to neurobehavioral disorders, the basis of neurobiological effects of NDL-PCBs is poorly understood. We assessed a range of effects of several NDL-PCB congeners (PCB153, PCB52, PCB118, and PCB138) on synaptic transmission and cellular morphologies linked to the proper execution of a sensorimotor response in zebrafish, taking advantage of the well-established neural circuit in this model. Startle responses were dramatically delayed in 6-day old larvae exposed to NDL-PCB. In contrast, exposure to the dioxin-like PCB126 did not delay startle response times. Morphological and biochemical data showed that exposure to NDL-PCBs during development induces swelling of afferent sensory neurons and disrupts dopaminergic and GABAergic signaling associated with motor movement. This study demonstrates that connections between neurotoxic mechanisms and processing of sensory stimuli occur at low concentrations of NDL-PCBs, similar to those present in human and animal samples. The effects on important and broadly conserved signaling mechanisms in vertebrates suggest that NDL-PCBs can contribute to neurodevelopmental abnormalities, relevant for both exposed human populations and wildlife.

Keywords

1 Introduction

Humans and wildlife are exposed to a wide range of contaminants in the environment, and the role of these chemical pollutants in developmental neurotoxicity, cognitive impairment, and neurodegenerative disorders has become a global environmental health concern (Landrigan et al., 2018). Among the top industrial chemicals thought to be involved in developmental neurotoxicity are polychlorinated biphenyls (PCBs) (Grandjean and Landrigan, 2014), a mix of 209 different structures, including planar dioxin-like (DL) congeners and non-planar non-dioxin-like (NDL) congeners. PCBs are persistent legacy contaminants that have been produced and used for decades to serve as insulators, coatings, and caulks, and there is worldwide PCB contamination of wildlife tissue and human cord blood, amniotic fluid, and mother's milk (Ennaceur et al., 2008; Lancz et al., 2015a). The developing nervous system is generally sensitive to chemical exposure at doses much lower than those affecting adult nervous system function (Rice and Barone, 2000). For instance, prenatal exposure to PCBs is associated with a decrease in motor skills in infants, while postnatal exposure seems to have no obvious effects (Ribas-Fitó et al., 2001). Thus, identifying the mechanisms underpinning the effects of neurotoxic PCBs during gestation and early life stages is of critical importance in attaining successful treatments and remediating health effects in humans and wildlife due to environmental exposure.

The NDL-PCBs are the most abundant congeners in human and environmental samples. The NDL-PCB138 and PCB153, are particularly abundant. Median PCB153 concentrations measured in human cord blood serum range from 6.4 to 110 ng g^{-1} lipid (Bergonzi et al., 2009; Herbstman et al., 2007; Lancz et al., 2015b), far greater than the levels of DL congeners. Evidence also suggests the NDL-PCBs contribute substantially to neurotoxic and behavioral effects (Boix et al., 2010; Boucher et al., 2009). However, understanding of the mechanisms

and effects of NDL-PCBs, especially during the more sensitive early stages of development,

significantly lags that of DL-PCBs.

Increased serum PCB concentrations in children are associated with deficits in cochlear

functions (Jusko et al., 2014; Trnovec et al., 2010), fine motor skills (Walkowiak et al., 2001),

and cognitive impairment (Boucher et al., 2009; Chen et al., 1992). Developmental exposure of

animals to NDL-PCBs has been linked to behavioral changes, including hyperactivity,

impulsiveness, and impairment of cognitive function (e.g., impairment of the ability to learn the Y

maze task) (Berger et al., 2001; Johansen et al., 2014; Schantz et al., 2003). NDL-PCB

developmental exposure also adversely affects auditory function in adult rats, including elevated

thresholds of brainstem auditory evoked potentials, auditory deficits at low frequencies,

audiogenic seizures, and reduced auditory startle responses (Crofton et al., 2000; Goldey et al.,

1995; Lilienthal et al., 2015, 2011; Poon et al., 2015). The mechanism(s) involved in these

effects on cognitive and auditory systems are not known.

Zebrafish are increasingly used as model organism to uncover the mechanisms of neurotoxicity

that are relevant for humans and other vertebrates. Zebrafish remain transparent throughout

early development, facilitating the use of transgenic and other molecular approaches to label

and observe specific classes of neurons and to follow their functional development. Fish have

relatively simple neural circuits, particularly for the auditory system. In fish, the auditory-evoked

startle response plays a critical role in animal survival, and involves cellular functions, sensory-

motor pathways, and monoamine neurotransmitters that are conserved among vertebrates

(Stewart et al., 2014). Many of the pathways and genes that regulate zebrafish startle behavior

also operate in human neurological disorders (Howe et al., 2013; Maximino and Herculano,

2010; Meserve et al., 2020; Tropepe and Sive, 2003; Wolman et al., 2015). In this study, we use

the zebrafish startle response to decipher the neuro-disruptive basis of NDL-PCBs on the

auditory system during the early stages of development. To this end, we examined the effects of

different NDL-PCBs (PCB153, PCB52, PCB118, PCB138), in comparison to the DL-PCB126

and at different environmentally relevant concentrations, dissecting various steps in the neural

circuitry underlying the acoustic startle response.

87   2   Materials and Methods

88   2.1   Animal model

In addition to other attributes, zebrafish have relatively simple neural circuits, particularly for the

auditory system. Many of the components of the auditory circuit are homologous in mammals

and fish, including hair cells, neural transmission, and lower motorneurons, while the

mechanosensory lateral line system and Mauthner (M) cells are exclusively found in fish and a

cochlea is only found in mammals. In fish, vibro-acoustic stimuli are first sensed by the hair cell

bundles in the auditory, vestibular, and lateral-line systems and rapidly encoded into trains of

action potentials in the afferent neurons (**Figure 1**) (Hudspeth, 1985). The signal detected

through the sensory hair cells of the lateral line is transmitted through the anterior and posterior

lateral line ganglia whereas the auditory input is transmitted through the VIIIth statoacoustic

nerve, all of which synapse to the lateral dendrite of the M-cell and two segmental homologs

(Nakayama, 2004; Pujol-Martí and López-Schier, 2013). The signal is propagated down the M-

cell axon in the spinal cord to activate the motoneurons leading to a unilateral contraction of the

trunk musculature.

Figure 1. Startle response circuit in zebrafish. The proper execution of the startle response relies on hair cells to sense the auditory stimulus, Mauthner-cells to integrate the signals, and motor neurons and musculature to execute the startle behavior.

2.2 Zebrafish husbandry and early-life stage exposure

Zebrafish (*Danio rerio*) were maintained at 28.5 °C and on a 12:12 h light-dark cycle according to the guidelines of the Zebrafish Model Organism Database (ZFIN, http://zfin.org) and institutional guidelines at WHOI and the experimental procedures approved by the Institutional Animal Care and Use Committee (ID Number BI21981.01). Adult zebrafish were fed brine shrimp and pellets every day. In this study, we used wild-type zebrafish of the AB strain obtained from Zebrafish International Resource Center (ZIRC), as well as several transgenic lines: the photoconvertible calcium indicator line *Tg[elavl3:CaMPARI(W391F+V398L)]*[if9] kindly provided by Jessica Plavicki (Brown University, USA), and the transgenic lines *Tg(cntn1b:EGFP-CAAX)* (Panlilio et al., 2021) labeling axons, *Tg(olig2:EGFP)*[vu12] (Park et al., 2007; Shin et al., 2003) marking oligodendrocyte precursor cell bodies, *Tg(sox10:mRFP)*

(Takada et al., 2010) marking oligodendrocyte lineage cells, as well as *Tg(mbp:EGFP-CAAX)* (Almeida et al., 2011) *and Tg(mbp:EGFP)* (gift from Dr. Kelly Monk, generated by Dr. Charles Kaufman in the laboratory of Dr. Leonard Zon, Harvard Medical School, Boston, MA) marking myelin sheaths. Zebrafish eggs were obtained by pairwise breeding and kept in 0.3x Danieau's (17 mM NaCl, 0.2 mM KCl, 0.12 mM $MgSO_4$, 0.18 mM $Ca(NO_3)_2$ and 1.5 mM HEPES, pH 7.6) in glass vials at a density of 10 individuals per 10 mL, at a constant water temperature of 28 ± 1°C and standard 14-h light/10-h dark cycle. The medium was supplemented with 1-Phenyl-2-thiourea (PTU, 0.003% w/v) from 24 hours post-fertilization (hpf) onwards to prevent pigment formation when necessary for imaging. Fertilized eggs were used for exposures starting at 3 hpf to solvent control (0.1% DMSO, v/v), 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153), 2,2′,5,5′-PCB, 2,2′,5,5′-tetrachlorobiphenyl (PCB52), 2,3',4,4',5-pentachlorobiphenyl (PCB118), 2,2',3,4,4',5'-hexachlorobiphenyl (PCB138), or 3,3',4,4',5-hexachlorobiphenyl (PCB126). Additionally, one exposure scenario with PCB153 was started at 24 hpf, to assess whether effects are induced during or after initial neuronal cell differentiation. Sensory neurons send their projection along the ectoderm as early as 16 hpf (Kimmel, 1993). The lowest exposure concentration for PCB153 was 10 nM (3.6 ng mL^{-1}) which is estimated to result in ~5 ng g^{-1} (5 ppm) per zebrafish larvae (comparing to PCB95 uptake levels; (Ranasinghe et al., 2019)). PCB153 mean concentrations found in maternal cord blood serum are 0.27 ng mL^{-1} or 110 ng g^{-1} lipid (Lancz et al., 2015b)). Other concentrations used were 30, 100, and 1000 nM which are found in fish of contaminated ecosystems (Gräns et al., 2015). For PCB52, PCB118, and PCB138, a concentration of 1000 nM was chosen to compare to the highest effect concentration of PCB153. The DL-PCB126 is cardioteratogenic in the low nanomolar range. To avoid malformations, a concentration of 1 nM was chosen. Exposures were refreshed daily until four days post-fertilization (dpf) and all larvae were kept in Danieau's from 5 to 6 dpf. All imaging and behavioral assays were performed at 6 dpf, except for the touch assay, which was performed at 26 hpf.

'

The influence of neurotransmitters on the startle behavior was assessed by exposures to

compounds known to interfere with monoamines. At 6 dpf, larvae were evaluated for their startle

behavior and then immersed in the dopamine precursors L-DOPA (1 mM) or L-tyrosine (1 mM),

dopamine receptor agonist quinpirole (10 µM), dopamine D2-receptor antagonist and GABA

receptor modulator haloperidol (10 µM), endogenous serotonin 5-HT (10 µM), and serotonin

precursor 5-HTP (1 mM) for 30 min in the well plate before assessing the startle behavior again.

Additionally, the effect of neurotransmitter modulation during development on startle response

was assessed by exposing larvae from 3 to 120 hpf to L-DOPA (1 mM), L-tyrosine (1 mM), and

quinpirole (10 µM).

2.3 Behavioral Analysis

Larval locomotion was monitored using the Noldus Daniovison system (Noldus Information

Technology, Leesburg, VA, USA) using 6-dpf larvae distributed in 48-well plates (1 larva per

well). After an acclimation period of 40 minutes in the illuminated chamber, spontaneous

swimming activity was tracked for 10 minutes. Following the spontaneous swimming, three

alternating cycles of dark and light conditions (10 minutes each) were run. The experiment was

repeated three times with cohorts from separate breeding events (n = 24 per group and

breeding event). Video data were recorded with 30 frames per second via an infrared camera.

Obtained data were analyzed with the supplied software EthoVision XT® 12 (Noldus. Inc.).

The activity of Rohon-Beard sensory neurons was tested at 26 hpf by lightly touching DMSO-

and 1000 nM PCB153-exposed embryos (n = 10 per group and breeding event) using a piece of

fishing line affixed to a glass pipette. Each embryo was touched ten times and the response was

scored as follows: 0, no response; 0.5, light tail flicking response, 1, coiling in the opposite

direction of the touch stimulus, yielding a score of maximum 10 per embryo. The experiment

was repeated twice with cohorts from separate breeding events.

Vibro-acoustic startle response was assessed as described previously (Panlilio et al., 2021).

Briefly, the acoustic stimuli were delivered by a vibrational exciter producing acoustic vibrations

at a frequency of a 1000 Hz for a duration of 2 ms at four different amplitudes (32, 38, 41, 43

dB). For each amplitude, the stimulus was delivered four times, spaced 20 seconds apart, to a 4

x 4 well-plate mounted on top of the speaker, each well containing an individual larva. The

response was tracked with a high-speed camera (Edgertronic SC1) at 1000 frames s^{-1} for 250

ms (13 ms before and 237 ms after the stimulus). PulsePal was used to synchronize the camera

recording to the acoustic stimuli using a TTL pulse. Automated analysis of larval movement

kinematics (response frequency, latency, turning angle) was performed using the FLOTE

software package (Burgess and Granato, 2007). Each treatment group was assessed at least

three times with eggs from different clutches. The total replicate numbers are stated in the

Table S1 (Supplemental Material).

A startle response in fish can be executed with a short latency C-bend (SLC) or long latency C-

bend (LLC), with the SLC being the more frequent response at higher stimulus intensities. The

cut-off time between SLC and LLC can vary depending on environmental conditions such as

temperature and has been determined empirically for this experiment using a Gaussian mixture

model (Panlilio et al., 2020)(**Figure S1**). Responses within 15 ms were categorized as SLC,

latencies greater than 15 ms as LLC. Following this classification, the median response type

(Relative Startle Bias) for each responsive fish was determined for each stimulus intensity (32,

38, 41, and 43 dB). Bias per individual fish was calculated as: (frequency of SLC – frequency of

LLC) / total responses. Pure SLC responses result in a value of +1, while pure LLC responses

are a value of -1 (Jain et al., 2018).

Electric field pulses (EFPs) directly activate the M-cells (Tabor et al., 2014). Responsiveness of

M-cells was assessed by applying EFPs (4.4 V cm^{-1} for 2 ms, square pulse) to head-restrained

larvae (n = 46 per treatment group). Seven consecutive EFPs were applied with a recovery time

of 20 s in between pulses. For each run, one control and one exposed larva were embedded in

1.5% agarose side-by-side and their tails freed. Response frequency, latency, and bend angle

were tracked using FLOTE. For all behavioral assays, only larvae with an inflated swim bladder

were used.

2.4   Larval growth

Larval length (from snout to the tail-fin end) was measured in DMSO (n = 57) and 1000 nM

PCB153 (n = 51) exposed larvae using ImageJ. The experiment was performed three times with

embryos from different clutches.

2.5   FM1-43 labeling

The vital fluorescent dye FM1-43 (n-(3-triethylammoniumpropyl)-4-(4-(dibutylamino)-styryl)

pyridinium dibromide; Invitrogen) specifically labels hair cells in the lateral line and inner ear of

vertebrates of by entering through open mechanotransduction channels and filling the

cytoplasm of hair cells (Meyers et al., 2003). At 6 dpf, zebrafish larvae of 1000 nM PCB153-

exposed (n = 32) and unexposed (n = 28) groups were immersed in Danieau's supplemented

with 3 µM FM1-43 for 30 s, rinsed three times in Danieau's, anesthetized in 1 mM 3-amino-

benzoic acid (Tricaine), and mounted in 1.5% low melting agarose for confocal imaging of the

L3 neuromast. The experiment was performed three times independently.

2.6   Immunohistochemistry

Whole-mount immunohistochemistry of zebrafish larvae and brains was performed following

standard procedures. Briefly, 10 AB larvae per exposure group were treated as described above

and subsequently fixed in 4% paraformaldehyde (PFA) in PBS overnight at 4 ˚C. For labelling

brain tissue, larval brains were dissected before continuing with the protocol. Multiple rinsing

steps in PBS containing either 0.5% Triton-X100 or 0.1% Tween-20 (PBST) for larvae or brains,

respectively, was followed by an antigen retrieval step in 150 mM Tris HCl (pH 9.0) at 70 °C

(Inoue and Wittbrodt, 2011), and a permeabilization step using proteinase K (10 µg/mL for 3

minutes). Subsequently, the samples were blocked in blocking solution (10% Normal Goat

Serum, 1% DMSO, 1% BSA, 88% PBST) for 1 h and then stained with blocking solution

supplemented with the primary antibody for three days at 4 ˚C. The primary anti-neurofilament

mouse polyclonal antibody (3A10; Developmental Hybridoma Bank, 1:100 dilution) was used to

label a subset of hindbrain spinal cord projecting neurons such as the M-cell neurons (DMSO *n*

= 23, PCB153 *n* = 21). The anti-5HT rabbit polyclonal antibody (Sigma-Aldrich, #S5545, 1:500)

was used to label serotonergic cells (DMSO *n* = 16, PCB153 *n* = 13). The anti-acetylated alpha-

tubulin mouse polyclonal antibody (Sigma-Aldrich, #T6793, 1:1000 dilution) was used to label

kinocilia and hair cell innervating neurons (DMSO *n* = 25, PCB153 *n* = 24). After several

washing steps, in PBST, the samples were incubated with Alexa Fluor 488- or 594-conjugated

goat-anti-mouse or goat-anti-rabbit secondary antibodies (Abcam, 1:500 dilution) overnight at 4

˚C and rinsed several times in PBST before mounting in ProLong Diamond Antifade mountant

(Invitrogen) for imaging. For visualization of skeletal muscle fibers (DMSO *n* = 11, PCB153 *n* =

11), larvae were fixed as above, followed by permeabilization in 5% Triton-X for overnight and

incubation in phalloidin Alexa-546 conjugate (Invitrogen, 1:500). All staining experiments were

repeated three times with larvae from independent clutches except the phalloidin staining, which

were performed once.

2.7 Formation of hair cell innervating neurons, primary motoneuron, and
oligodendrocyte lineage cells

The *Tg(cntn1b:EGFP-CAAX)* transgenic line was used to image the cell bodies of neurons that

innervate the hair cells at the L3 position in the lateral line (DMSO *n* = 22, PCB153 *n* = 20, two

independent experiments). The axons of the main caudal primary motoneurons were visualized

using the *Tg(olig2:EGFP)* transgenic line (*n* = 4 per treatment group, one experiment). The

double transgenic *Tg(olig2:EGFP) x Tg(sox10:mRFP)* was used to quantify oligodendrocyte

lineage cells in the spinal cord of in DMSO (n = 22) and 1000 nM PCB153 (n = 16) treated

larvae (Shin et al., 2003). All larvae were anesthetized in tricaine (0.16% MS222) before

mounting laterally in 1.5% low melting agarose on a glass-bottom microscopy dish (MatTek).

Fluorescently labelled cell bodies were imaged at the distal end of the intestine, around the L3

neuromast. The experiments were performed three times independently.

2.8 Calcium imaging of freely swimming and startling larvae

Neuronal activity in DMSO- and PCB153-treated larvae was visualized using the CaMPARI

(*elavl3*:CaMPARI) transgenic zebrafish line with an engineered fluorescent protein that

permanently photoconverts from green to red in the presence of elevated calcium and UV light

(Fosque et al., 2015). Photoconversion was achieved by mounting an LED light (405 nm;

LEDSupply) between the speaker and a custom-made 3-well plate containing the freely

swimming larvae (DMSO- and PCB153-treated). Larvae of the same treatment were divided into

two batches: one was photoconverted in medium at room temperature (RT) and the other one at

4˚C to elicit high neuronal activity. Photoconversion was achieved by shining the LED light for

10 sec either with one vibrational stimulus of 43 dB at the start or without stimulus.

Subsequently, larvae were anesthetized, embedded in agarose, and whole brains imaged using

the 10x objective. The experiment was performed twice (n = 11-16 per treatment group).

2.9 Image processing

Whole-brain, hair cells, M-cell, and motor neurons were imaged on a Zeiss LSM 800 confocal

system with a 10x/0.45 Plan-Apochromat, 20x/0.8 Plan-Apochromat, or 63x/1.2 C-Apochromat

water immersion objective. Airy scan was applied for hair cell imaging at 63x. Z-stacks were

merged to a single plane by maximal intensity projection using Fiji (Schindelin et al., 2012). The

number of *olig2* cells dorsal of the spinal cord (DMSO n = 22, 1000 nM PCB153 n = 16) was

counted in confocal image stacks using the 3D projection of the Zeiss Zen blue software.

Myelination was evaluated after imaging the spinal cord (DMSO n = 26, 1000 nM PCB153 n =

27) using the 20x objective and scored for deficits according to Panlilio et al. 2020.

Representative myelination was imaged at 40x. The intensity of serotonergic neurons in the

brain (DMSO n = 16, 1000 nM PCB153 n = 13) was measured performing a Sum Slices Z-

projection followed by drawing a ROI around the 5-HT immuno-responsive cells, and per image,

three random squares (area size = 10.65 x 10.65) within the non-fluorescent area of the ROI for

background subtraction (for orientation see **Figure S12**). Similarly, the FM1-43 intensity was

measured drawing a ROI around the fluorescent hair cells and measuring the background in

three random squares (DMSO n = 20, 1000 nM PCB153 n = 20). Corrected total cell

fluorescence (CTCF) was calculated according to the formula: Integrated density - (ROI x Mean

fluorescence of background readings). For quantification of active neurons in the brain of freely

swimming and startling CaMPARI zebrafish larvae, Sum Slices Z-projection were analyzed for

red fluorescence intensity in the rostral hindbrain (for orientation see **Figure S13**).

The volume of hair cell innervating neurons was calculated by measuring the volume of

thresholded pixels (30-255, Otsu method) in image stacks using an imageJ macro looping

through the stack and summing the area measurements, then multiplying this sum by the depth

of each slice (DMSO n = 30, 1000 nM PCB153 n = 29). The presence of primary motor neurons

was confirmed using the Simple Neurite Tracer plugin in Fiji (DMSO n = 21, 1000 nM PCB153 n

= 18). All image evaluations and quantifications were done without the knowledge of the

experimental condition of the specimens.

2.10 Neurotransmitter analysis

The analysis of neurotransmitters, including histamine, L-histidine, L-glutamic acid, gamma-

aminobutyric acid (GABA), acetylcholine, L-DOPA, 3-methoxytyramine, serotonin, 5-

hydroxyindole-3-acetic acid, glycine, L-glutamine, L-valine, 3-hydroxytyramine and choline, was

performed with a method developed in a previous study (Zhao et al., 2020). Briefly, the

zebrafish embryo samples were spiked with 100 ppb isotope-labeled internal standards and

then homogenized. The homogenized tissue samples were extracted with 200 µL of 80:20 v/v

ACN/Milli-Q water containing 0.1% formic acid and 1% ascorbic acid and 200 µL ACN. After

protein precipitation, the supernatants were dried and redissolved in 100 µL of Milli-Q water

containing 0.1% formic acid for UHPLC-MS/MS analysis. Neurotransmitters were quantified by

a high-performance liquid chromatography system coupled to a Q-Exactive plus quadrupole-

Orbitrap mass spectrometer (UHPLC-Q-Orbitrap, Thermo fisher, US). The separation of

neurotransmitters was carried out by an Acquity UPLC HSS T3 column with Milli-Q water

containing 0.1% formic acid and ACN containing 0.1% formic acid as mobile phase. The MS

acquisition was performed in the full scan mode and parallel reaction monitoring mode. The

experiment was performed once (n = 4 per treatment group).

2.11 Statistical Analyses

All statistical analyses were conducted using Graph Pad Prism 8.4 except some of the startle

behavior analysis, which was completed in R. All quantitative data were tested for normality

using the Shapiro-Wilk test, and if there were unequal number of subjects between groups, the

data were tested for homogeneity in variances using the Brown-Forsythe test. All treatment

groups were compared to respective control groups.

Responsiveness to auditory stimuli (response or not) and startle type (SLC vs. LLC) was

modelled in R (version 4.0.2) using a binomial generalized linear mixed effect model (glmer()

from the lme4 R package) (Bates et al., 2015). The response to electrical stimulus was modeled

using glm() from the lme4 R package. For the auditory response, we included treatment as a

fixed effect and the replicate experimental trials as a random effect. The model specification for

"fraction responding" was as follows: Responders ~ TreatmentGroup + (TreatmentGroup|trial).

The model specification for "startle bias" was as follows: SLCorLLCresponse ~ TreatmentGroup

+ stimdB + (1|trial). The data were analyzed fitting the model for each stimulus intensity

independently. The model specification for the electrical stimulation are as follows: Responders

~ TreatmentGroup.

To assess treatment differences of normal data with three or more groups (fluorescence

intensity CaMPARI), ANOVA comparisons were performed followed by a multiple comparison

using the Tukey's correction to adjust the critical values. A Tukey-corrected p-value < .05 was

considered statistically significant. For normal data with heterogeneity within-group variances

(LLC bend angle), a Welch ANOVA followed by a Dunnett's T3 multiple comparison test was

performed. For non-normal data with three or more groups (SLC bend angle), a Kruskal-Wallis

followed by the Dunn's multiple comparison test was performed. For normal data with two

groups (length, mean latency, bend angle in e-stim, oligodendrocyte lineage cell count, 5-HT

intensity), an unpaired two-tailed t-test with Welch's correction (Welch's t- test) was performed.

The nonparametric scoring data of the touch assay and latency in e-stim was analyzed using a

two-tailed Mann-Whitney test. The data are presented as mean ± SD with single data points

(replicates) superimposed on the graph unless specified in the figure legend.

3   Results

The startle response circuit is of wide relevance and well-established for assessing the influence

of specific factors on neuronal cell development, neuromuscular connectivity, and transmission

(Burgess and Granato, 2007; Medan and Preuss, 2014; Meserve et al., 2020; Panlilio et al.,

2020). We used the startle response circuit to assess the effects of NDL-PCBs on the formation

of neuronal populations and neurotransmission and its consequences for mechanosensory

disruptions, thus encompassing hearing and touch.

3.1 Behavioral effects

Rohon-Beard cells are the earliest mechanosensory neurons that develop. They are located in the dorsal region of the embryonic spinal cord and appear around 24 hpf (Roberts, 2000). A few days into development, Rohon-Beard cells also relay sensory information to the M-cells (Liu and Hale, 2017). In our experiments, a tactile stimulus on the trunk at 26 hpf evoked a C-shaped body flexion in both the DMSO- and the 1000 nM PCB153-treated embryos (U = 183, p = .599; **Figure S2**), indicating that the formation and function of Rohon-Beard cells were not impaired by PCB153.

Exposure to contaminants during development can result in reduced growth, which can affect kinematics in swimming behavior (Voesenek et al., 2018). PCB153-exposed larvae developed normally with no difference in body length on day six in comparison to the control group, in all three trials (Trial 1: t(25.98) = .661, p = .514; Trial 2: t(37.98) = .443, p = .660; Trial 3: t(37.43) = .645, p = .523; **Figure S3**). At 6 dpf, spontaneous and light-dark-stimulated swimming activity was assessed. Larvae exposed to PCB 153 (1000 nM) had lower cumulative locomotion activity over ten minutes (M = 1182 mm, SD = 382) in comparison to the control group (M = 1510 mm, SD = 402), t(132.4) = 4.854, p < .0001 (**Figure S4A**). Zebrafish larvae typically increase locomotion in dark conditions following light exposure. Once stimulated with a rapid loss of illumination during the dark phase, both treatment groups exhibited similar swimming activity (**Figure S4B**).

The vibro-acoustic startle response can be executed either rapidly, within 15 ms (short latency C-bend; SLC), or beginning at longer times and extending for up to 100 ms (long latency C-bend; LLC). The mean latency value, representing the population response including SLC and LLC responses, was 21 ± 20 ms for control larvae at the highest stimulus intensity (**Figure S1**, **Figure 2A, B**). By contrast, the NDL-PCB-treated larvae that responded to the vibro-acoustic stimulus showed significantly longer average latencies in response to the same auditory

stimulus given the controls, $F(5, 907 = 14.61, p < .0001)$. PCB153-treated (1000 nM) larvae

responded at a mean latency of 68 ± 32 ms, PCB138 at 52 ± 37 ms, PCB118 at 45 ± 27 ms,

and PCB52 at 39 ± 32 ms. For PCB153, a dose-dependent response was observed, with mean

latencies increasing with dose (10 nM: 26 ± 27 ms, 30 nM: 50 ± 36 ms, 100 nM: 50 ± 30 ms,

1000 nM: 68 ± 32 ms). The lowest PCB153 concentration tested (10 nM) was not significantly

different from controls. Unlike NDL-PCB treated larvae, the DL-PCB126-treated larvae had a

mean latency (22 ± 22 ms), which was not significantly different from controls.

Examining the SLC and LLC responses separately can give insight into the underlying

mechanism of effect. A greater stimulus intensity increases the probability of evoking SLC,

shifting the response bias from LLC to SLC (Troconis et al., 2017). Here, we also observed a

shift to SLC with increasing stimulus intensity in solvent control (DMSO-treated) larvae (**Figure**

2C, Figure S5A). Larvae were tested four times at each stimulus intensity; thus, a larva is

categorized as having SLC bias if it responded 3 or 4 times out of 4 stimuli with an SLC. At the

highest stimulus intensity that we tested (43 dB), the majority of control larvae (300/448)

responded with a SLC within 7 ± 3 ms. 43 larvae showed no bias towards SLC or LLC (2 times

SLC and 2 times LLC, bias = 0). Those that did not perform an SLC (n = 105) exhibited an LLC

with a mean latency of 54 ± 29 ms (**Figure 2C**). In contrast, all NDL-PCB-treated larvae (except

for 10 nM PCB153) retained a response bias towards LLC at all stimulus intensities tested,

which was significantly different from the control ($p < .001$ for all stimulus intensities). Lowering

the PCB153 dose gradually increased the shift towards SLC, yet 10 nM PCB153 was still

significantly different from the control (p <.01 for all stimulus intensities). Exposures starting

after many structures of the brain and the neural tube have developed, at 24 hpf (Smith and

Kimelman, 2019), had the same effect on startle latency (**Figure S5B)**.

The proportion of control larvae responding to the stimulus (either SLC or LLC) increased from

75.2 ± 33.0% (n = 450) to 95.8 ± 17.0% (n = 457) from the lowest to highest stimulus intensity,

whereas the proportion of NDL-PCB153-, PCB138-, and PCB52-treated larvae that responded

remained below 70% (except for 10 nM PCB153) at all intensities and were significantly

different (p < .01) from the control (**Figure 2D**, **Figure S6A**). In contrast to those exposed to

NDL-PCBs, the larvae exposed to the DL-PCB126 and NDL-PCB118 had a response latency

and rate similar to the control larvae.

The different startle response bends exhibit different kinematics, with SLCs exhibiting larger

bend angles and faster angular velocities than LLCs (Burgess and Granato, 2007). Thus, LLC

and SLC kinematics were analyzed separately. PCB52-treated larvae had increased SLC bend

angles (p > .0001) at two intermediate stimulus intensities (38 and 41 dB) while PCB126 larvae

had increased SLC bend angles at the three highest stimulus intensities (**Figure S7**).

Furthermore, PCB153 (1000 nM) exposure resulted in an increased SLC bend angle at the

highest stimulus intensity for those few larvae that exhibited an SLC (n = 26 of 138, 43 dB, p <

.0456). LLC bend angle was significantly decreased across all stimulus intensities for PCB118.

It is not clear how bend angle is associated with altered signaling and cellular function.

Figure 2. Effect of PCBs on startle response in 6 dpf zebrafish. (**A**) Representative short-latency C-bend (SLC) and long-latency C-bend (LLC) startle response of a control zebrafish larva at 6 dpf in comparison to a typical response of a PCB153 treated larva. (**B**) Vibro-acoustic startle latency at highest stimulus intensity (43 dB) in vehigle control (*n* = 479), NDL-PCB153 1000 nM (*n* = 179), NDL-PCB153 100 nM (*n* = 253), NDL-PCB153 30 nM (*n* = 152), NDL-PCB153 10 nM (*n* = 119), NDL-PCB138 (*n* = 139), and DL-PCB126 (*n* = 97) exposed larvae. (**C**) Bias of SLC and LLC and (**D**) response rate of different exposure groups at different vibro-acoustic intensities. Values are presented as mean ± SEM. All data points are biologically independent replicates from at least three independent experiments. Significant differences to DMSO controls (p < 0.01) is indicated by open symbols.

3.2   Cellular structures

The proper execution of the startle response relies on hair cells that sense the auditory stimulus, M-cells to integrate the signals, and motor neurons and musculature to execute the startle behavior. Morphological analysis showed that M-cells, motor neurons, and musculature all appeared unaffected by PCB153, but the hair cell innervating neurons appeared to be swollen (**Figure 3**). The L3 neuromast volume of control larvae was on average 569 µm^3 (*SD* = 125),

whereas PCB153-exposed larvae (1000 nM) had significantly larger L3 neuromast volumes 848

µm^3 (*SD* = 185), *t*(48.93) = 6.765, *p* < .0001 (**Figure 3B**). However, the number of hair cells in

L3 neuromasts (**Figure 3C**) and the intensity of FM1-43 dye internalized by hair cells (**Figure**

3D) remained unchanged in PCB153-treated larvae in comparison to the control.

Figure 3. Effect of PCB153 on morphology and function of lateral line hair cells and

innervating neurons. (**A**) Diagram indicating L3 neuromast. (**B**) Hair cell innervating neurons at

neuromast L3 shown using *Tg(cntn1b:EGFP-CAAX)*. Volume of hair cell innervating neurons of

DMSO (*n* = 30) and PCB153 (*n* = 29) treated larvae. Scale bar = 20 µm. (**C**) Whole-mount immune

staining of kinocilia and hair cell innervating neurons (anti-acetylated tubulin antibody; green) and

counterstain of stereocilia bundles and muscle tissue (actin; magenta) of DMSO (*n* = 26) and

PCB153 (*n* = 24) treated larvae. The hair cell number was determined based on the number of

kinocilia. Scale bar = 10 µm (**D**) Uptake of FM-143 dye as indication of functional hair cells of

DMSO (*n* = 32) and PCB153 (*n* = 28) treated larvae. Scale bar = 10 µm. All data points are

biologically independent replicates from at three independent experiments. Asterisks indicate

significant differences to controls (***p < 0.0001).

The presence of M-cells is necessary for the SLC startle response. Staining with anti-3A10

confirmed the presence of M-cells in all PCB153-exposed larvae (21/21) (**Figure 4A**). We tested

the functional responsiveness of the M-cells by applying electric field pulses to head-restrained

larvae. This electrical stimulation bypasses the sensory system and directly activates the M-cell

(Tabor et al., 2014). There was no difference detected in response time between control and

1000 nM PCB153-exposed larvae (U = 907.5, p = .489), demonstrating that the signal

transmission from the M-cell to the motor neurons is not disrupted in PCR153-exposed fish

(**Figure S8A**). The fraction responding also was similar in both the DMSO-treated (290/299) and

the PCB153-treated (209/219) groups (GLM, p = .427; **Figure 4C, Figure S8B**). However, the

C-bend evoked by the electrical stimulation was larger in PCB153-exposed larvae (M = 101.4,

SD = 38.60) than in the control larvae (M = 83.12, SD = 24.23). An unpaired t-test with Welch-s

correction indicated that this difference was statistically significant, t(74.27) = 2.679, p =.0091

(**Figure S8C**). This difference could indicate a PCB153-evoked functional alteration of the M-cell

axon or motor neuron.

Figure 4. Effect of PCB153 on morphology and function of the Mauthner-cells. (**A**) Diagram
indicating localization of Mauthner-cells in the hindbrain. Brain tissue immunostaining with anti-
neurofilament 3A10 antibody labeling reticulospinal neurons in control and PCB153 treated larvae

at 6 dpf (ventral view). Mauthner (arrow) cells were present in all brain tissues (control: n = 23,

PCB153: n = 21). Scale bar = 50 μm. (**B**) Diagram of electrical stimulation performed to assess

the functionality of Mauthner-cells. (**C**) Percentage of responders to electrical stimulation. All data

points are biologically independent replicates from at least three independent experiments.

For rapid and effective axonal signal transmission, proper myelination of axons is key. We did

not find PCB153 to reduce cells of the oligodendrocyte lineage in the spinal cord, (t(36) = .553,

p < .584) nor any obvious differences in myelination itself (**Figure S9**).

Proper function and activation of the motor neurons is crucial for completion of the startle circuit,

leading to muscle contractions and the characteristic swimming behaviors. The main caudal

primary (CaP), middle primary (MiP), and rostral primary (RoP) motor neurons did not show any

distinct structural differences between exposed and control larvae (**Figure S10**). The fast

muscle fibers in the myotome were not found to be differentially striated in either treatment

groups (**Figure S11**). No structural difference in the motor neurons or the myotome was

apparent in PCB153-exposed larvae, suggesting that the increased bend angle does not

originate from any overt structural differences in the motor neurons or muscle fibers.

3.3   Altered neural activity

The release of neurotransmitters into the synaptic cleft is a characteristic process in signal

transmission. Both serotonergic and dopaminergic neuromodulators can affect startle latency

(Jain et al., 2018), and evidence has been found that NDL-PCBs alter these neuromodulators in

mammals (e.g. Castoldi et al. 2006; Enayah et al. 2018; Wigestrand et al. 2013). We observed

that PCB153 (1000 nM) altered the levels of several neurotransmitters in 6 dpf zebrafish larvae

(**Figure 5A**). Significantly increased level of GABA (t(6) = 9.719, p < .001), choline-chloride (t(6)

= 8.459, p < .001), 3-methoxytyramine hydrochloride (t(6) = 3.803, p = .009), and L-valine (t(6) =

3.134, p = .020) were measured in PCB-exposed larvae in comparison to control larvae; while

L-Glutamine was decreased in comparison to control larvae (t(6) = 5.966, p < .001).

To assess the interaction between PCB-exposure and changes in neuromodulatory activity, we

pharmacologically manipulated the neurotransmitter levels using dopamine, serotonin, and

GABA receptor agonists and antagonists. In control larvae, a 30 min exposure to dopamine

precursor L-DOPA and D2-receptor agonist Quinpirole, as well as serotonin precursor 5-HTP,

shifted the response to LLC, at multiple intensities tested ($p < .01$; **Figure 5C**). In contrast, the

dopamine antagonist haloperidol exposure induced a shift to SLC in controls across all stimulus

intensities ($p < .01$; **Figure 5C**). Haloperidol induced the same shift to SLC in PCB153-exposed

larvae ($p < .001$), suggesting that the startle circuit is functional but repressed by PCB (**Figure**

5C). The response rate remained unchanged after exposure to neurotransmitter modulators

except 5-HT, which caused an increase in response probability across all stimulus intensities in

control larvae ($p < .01$); Quinpirole, which caused a decrease in response probability at the two

lowest stimulus intensities in control larvae ($p < .01$); and Haloperidol, which caused an increase

in response probability at 41 dB in PCB153-treated larvae ($p < .01$; **Figure S6**). It is noteworthy,

that co-exposure to L-DOPA and PCB153 throughout development increased the response rate

of PCB153-treated larvae significantly ($p < .01$), yet only induced a significant shift towards SLC

at 38 dB ($p < .01$; **Figure S5C and S6C**).

Figure 5. Involvement of neurotransmitters in startle latency. (**A**) Neurotransmitter levels in

6 dpf larvae exposed to 1000 nM PCB153 in comparison to control ($n = 4$). (**B**) Representative

maximum intensity z-projection from confocal stack after photoconversions of freely swimming 6

dpf CaMPARI larvae. Scale bar = 100 µm. Ratio of red to green fluorescence intensity in the

rostral hindbrain (indicated by dashed line in image) in DMSO ($n = 14$) and 1000 nM PCB153 (n

= 14) exposed startling larvae in 4 ˚C (cold) medium. Dashed lines indicate median per trial. Two

independent experiments. Asterisks indicate significant differences to controls (*p < 0.05, **p <

0.01, and ***p < 0.001). (**C**) Startle bias shift induced by 30-minute exposure to serotonin,

dopamine, and GABA neurotransmitter modulators in 6 dpf control and 100 nM PCB153 exposed

larvae. Control ($n = 138-172$), 5-HTP ($n = 67-80$), L-DOPA ($n = 76-94$), Haloperidol (Control $n =$

123-162, PCB153 $n = 37-61$), and Bicuculline (Control $n = 90-162$, PCB153 $n = 83-137$). All data

points are biologically independent replicates from at least three independent experiments.

Significant differences to respective controls ($p < 0.01$) are indicated by open symbols.

In the fish brain, a distinct population of serotonergic neurons is found in the posterior

tuberculum/hypothalamus, forming a prominent horseshoe-like pattern that is one of several

brain regions where 5-HT is synthesized (Oikonomou et al., 2019). To assess whether the

development of serotonergic cells in the brain of PCB153 larvae was affected, we performed

immunohistochemistry using an antibody for 5-HT (**Figure S12**). The intensity of the fluorescent

signal in the hypothalamus was not different, comparing PCB153 (M = 4.70 x 10^6, SD = 2.10 x

10^6) with the control (M = 4.38 x 10^6, SD = 8.1 x 10^5) group. This suggests that the serotonergic

neurons in this region were not affected.

During specific behaviors or sensory experiences, only a small percentage of neurons are

active. We analyzed the active neuronal cell populations in PCB153-treated and -untreated

larvae, focusing on the rostral hindbrain, where the M-cells are located, to assess whether there

were changes in neural activity. We examined neuronal activity using the CaMPARI zebrafish

line, which contains a protein that upon UV illumination permanently photoconverts from green

to red fluorescence in cells with high calcium levels (Fosque et al., 2015). This genomic tool

enabled us to look for active neuronal cell populations in larvae during normal swimming

movement, and in larvae experiencing a vibroacoustic stimulated startle response, both in

embryo medium at room temperature (24 °C), and at a colder temperature (4 °C). A higher red

to green ratio (F_{red}/F_{green}) was measured comparing cold medium with room temperature

medium: swimming control larvae in cold medium had a higher F_{red}/F_{green} ratio (M = 2.30, SD =

0.51) in comparison to swimming control larvae at RT (M = 0.96, SD = 0.17), confirming the

potential of cold medium to elicit higher neuronal activity (**Figure S13**). However, this difference

was diminished in PCB153-treated, yet still significant in trial 1: $F(1, 18)$ = 121, *adj. p* < .001 and

trial 2: $F(1, 32)$ = 65.2, *adj. p* = .041. Importantly, the difference between swimming control and

PCB153-treated larvae in cold medium was significant in both trial 1: $F(1, 18)$ = 121, *adj. p* =

.028 and trial 2 $F(1, 32)$ = 65.2, *adj. p* < .001 (**Figure 5**). Similar differences in neuronal activity

between control and PCB-153 treated larvae were measured for startling larvae during

photoconversion (**Figure S13**).

4   Discussion

Despite most use ending decades ago, PCBs remain an important and persistent pollutant

impacting both wildlife and humans. To date, mechanistic studies of neurodevelopmental effects

of NDL-PCBs suggest two major targets: (1) alterations in neurotransmission, particularly

involving dopamine and serotonin, and consequently disruptions to Ca^{2+} signal transduction

(Campagna et al., 2011; Dervola et al., 2015; Enayah et al., 2018; Fonnum and Mariussen,

2009; Langeveld et al., 2012), and (2) loss of outer hair cells (Crofton et al., 2000; Trnovec et

al., 2010) or cochlear development (Goldey et al., 1995). However, such studies have not

addressed how PCBs exert developmental neurobehavioral deficits, and thus we do not know

whether neurobehavioral disorders resulting from prenatal exposure originate in a structural

effect in the auditory system, a neurotransmitter imbalance, or both. Furthermore, PCBs are

prevalent in aquatic ecosystems, yet we do not know the behavioral effects in fish and the

specificity of NDL-PCB congeners for these effects. In this study we used the zebrafish startle

response to comprehensively investigate the effects of different PCBs on neuron development

and signal transmission and to assess effects at environmentally relevant concentrations for

both humans and aquatic life.

The morphology of the neuronal cell body (perikaryal) and neurite provides essential information

on the developmental and physiological abnormalities in the neural circuits. We found that in

PCB153-exposed larvae, the neurons that innervate the hair cells had an increased volume

(**Figure 3**), indicating that PCB153 alters neuronal structure and hair cell function. Swelling of

afferent terminals is associated with hearing deficits in both mammalian models (Puel et al.,

1991) and zebrafish (Sebe et al., 2017). The increased volume may originate from increased

numbers of bulged (swollen) afferent neurite termini or increased arborization (branching). For

example, *tmie* mutant zebrafish, which are deaf with defective hair-cell function (Gleason et al.,

2009), have more complex peripheral arbors in afferent axons and an increased number of

neurites. In wildtype zebrafish, neurites with swollen axonal terminals are associated with hair

cells of both polarities rather than just a single polarity, suggesting a dysregulation of selection

in synaptic targets (Faucherre et al., 2010). At the sub-cellular level, neuronal swelling is due to

increased cationic permeability in neurons, causing an osmotic imbalance that leads to water

entry and cytotoxic edema (Rungta et al., 2015). While we examined a number of morphological

features of the startle circuit anatomy (hair cell, afferent neurons, M-cell, oligodendrocyte

lineage cells, myelination, motor neuron, muscle), only the volume of the neurons that innervate

the hair cells showed a measurable deviation in PCB153-exposed larvae. We thus infer that the

enlargement of axon termini is contributing to the neurotoxic potential of NDL-PCBs – an

understudied target of PCB toxicity. A few studies associate NDL-PCB exposure with altered

dendritic arborization (Lein et al., 2007; Yang et al., 2009). Axonal swelling is a pathology

appearing in many neurodegenerative disease (Coleman, 2005), and exposure to NDL-PCBs

may be a contributing factor in such diseases.

The startle response in fish and amphibians is a fundamental mechanism involved in predator

escape. Acoustic stimuli can trigger two distinct motor behaviors: a short-latency C-bend (SLC**;**

Figure 2A) initiated within 5-15 ms of the stimulus, or a long-latency C-bend (LLC; **Figure 2A**),

initiated within 15-100 ms (Burgess and Granato, 2007). At the structural level, laser-ablation of

the M-cell abolishes SLC responses without affecting LLC responses (Lacoste et al., 2015),

suggesting that the LLC is a 'backup' response (Burgess and Granato, 2007). The finding that

NDL-PCB-exposed larvae predominantly exhibit LLC without loss of M-cells (**Figure 4A**),

suggests that the M-cells are receiving a delayed input, are not firing properly, or that the signal

propagation downstream of the M-cell is disrupted. To distinguish these, we directly activated

the M-cells by applying electrical stimulation to control larvae with their heads restrained, and

found that unlike auditory stimulation, PCB153-exposed fish had latencies indistinguishable

from the controls. Furthermore, direct stimulation of PCB153-exposed larvae resulted in a

normal response rate and response time, although the bend angle was significantly increased.

These results indicate a proper function of M-cells in NDL-PCB-treated larvae. However, the

changes in bend angles we observed during these responses could indicate alterations

downstream of the M-cells.

There is less known about the mechanisms leading to an aberrant turning angle. Since we

found no obvious structural impairment downstream of the M-cell (oligodendrocyte lineage cells,

myelination, motor neuron, musculature), we speculate that the increased bend angle in

electrically stimulated larvae exposed to PCB153 may be linked to a sub-cellular disruption such

as gap junctions. It is known that PCBs can alter gap junction function in cells of many organs

(Bager et al., 1997; Kang et al., 1996; Machala et al., 2003) including NDL-PCB specific

inhibition of gap junctional intercellular communication in neural crest cells (Nyffeler et al., 2018)

and neuronal stem cells (PCB153; (Kang et al., 2001)). Gap junctions are intercellular channels

formed by connexins (Cx). RNA sequencing of zebrafish larvae exposed to PCB153 in a similar

exposure regime (4-120 hpf) and concentration (10 µM) significantly altered the expression of

connexin genes cx28.9, cx42, and cx35b (Aluru et al., 2020). Cx35 is expressed on the

Mauthner neuron lateral dendrites in zebrafish (Jain et al., 2018) and plays a crucial role in

regulating spinal motor activity during fast motor behavior of adult mosquitofish (Serrano-Velez

et al., 2014). We thus propose that gap junction function could be disrupted, which may be a

future study direction using single-cell RNA sequencing or genetically encoded optogenetics

that has yet to be developed in a model organism (Dong et al., 2018).

Neurotransmitters also play a key role in the startle response. A previous screening of more

than 1000 pharmacologically active compounds identified serotonergic and dopaminergic

modulators among the largest class of drugs that shift bias between SLC and LLC startle

behavior (Jain et al., 2018). Specifically, acute exposure to the D3R agonist (7OHD) shifted the

startle bias from LLC to SLC, while exposure to a D3R antagonist (or a 5-HT1AR agonist)

shifted bias toward LLC behavior (Jain et al., 2018). Our results extend these findings, showing

that 30 minutes exposure to L-DOPA or 5-HTP induces a shift from SLC to LLC at higher

stimulus intensities, as opposed to the D2R antagonist haloperidol, which evoked a shift to a

stronger SLC bias at all stimulus intensities (**Figure 5C**). NDL-PCBs are thought to affect

dopaminergic pathways (Mariussen and Fonnum, 2001; Tanaka et al., 2018; Wigestrand et al.,

2013), which is partly supported by our measurements of catecholamines and traces amines.

Specifically, we found that the extracellular dopamine metabolite 3-methoxytyramine

hydrochloride and the inhibitory transmitter GABA were significantly increased (**Figure 5A**). The

interplay between dopamine and GABA receptors activity is complex. In the mammalian system,

dopamine is a direct-acting modulator of $GABA_ARs$ (Hoerbelt et al., 2015). The simultaneous

presence of a dopamine metabolite and GABA in PCB153-exposed zebrafish larvae may

indicate that increased levels of GABA are a downstream effect of dysregulated dopamine

activity. With both dopamine and GABA playing critical roles in motor activity, we hypothesize

that PCB153 impairs dopaminergic neurotransmission and subsequently increases GABA

levels, leading to the predominant LLC response. One of the most crucial regulators of synapse

development and function mediating the GABAergic, dopaminergic, and serotonergic synaptic

transmission is the brain derived neurotrophic factor (BDNF). PCB153 induced the expression

of *bdnf* in zebrafish larvae (Aluru et al. 2020) suggesting BDNF as potential target for

transcriptional regulation. Furthermore, we propose that GABA is involved in the swelling of the

afferent terminals (**Figure 3**) as GABA activates Cl⁻ channels leading to an influx of Cl⁻

accompanied by H_2O (Cesetti et al., 2012; Rungta et al., 2015). The osmotic neuronal swelling

alters neuronal activity. This study thus provides new evidence that GABA is involved in the

neurobehavioral mode of action of higher chlorinated NDL-PCBs, as has been for lower

chlorinated congeners (Fernandes et al., 2010).

An important finding of this study is that PCB153-exposed larvae can execute SLC when treated

with haloperidol, a D2-antagonist that can increase dopamine turnover in acute exposure

(Magnusson et al., 1987). There is a large body of evidence showing that NDL-PCBs, including

PCB153, modulate dopaminergic neurotransmission by inhibiting the dopamine precursor

enzyme tyrosine hydroxylase and dopamine synthesis itself, in both neural crest cells and adult

laboratory animal models including non-human primates (Dervola et al., 2015; Enayah et al.,

2018; Fonnum and Mariussen, 2009; Seegal et al., 1990). In zebrafish larvae, PCB153

exposure during development elicited differential expression of nine genes related to the

dopamine pathway, including tyrosine hydroxylase 2 (th2) (Aluru et al., 2020). Yet linking such

effects to behavior is rare. Tanaka et al. (2018) suggested that the NDL-PCB dominant mixture

Aroclor 1254 and BDE-47 functionally inhibited dopaminergic neurons in early zebrafish

embryos, as increased activity in 26 hpf embryos was inhibited by supplementation with L-

tyrosine and L-DOPA (Tanaka et al., 2018).

The restoration of the delayed startle phenotype by haloperidol, together with the increased

levels of a dopamine metabolite and GABA in PCB153-exposed larvae (**Figure 5A**), indicates

that disruption of dopaminergic and GABAergic signaling between neurons of the auditory circuit

is the dominant effect of PCB153-mediated aberrant motor behavior. These signaling processes

are widely conserved across vertebrate species, and are important in human neurological

diseases. For example, a recent study found that multiple gene mutations causing defective

startle response in zebrafish were also associated with disorders of the locomotor system in

humans (Meserve et al., 2020).

The localization of the PCB-induced disruption of dopaminergic signal transmission is likely

downstream of the hair cells, potentially in the hindbrain. Although dopaminergic neurons

innervate lateral line neuromasts, where D1R antagonist reduces the hair cell activity (Toro et

al., 2015), we found no difference in the hair cell activity of PCB153-exposed larvae (**Figure**

3D). This is in contrast to the hindbrain, where neuronal activity was reduced in PCB153

exposed CaMPARI zebrafish larvae (**Figure 5**), indicating a reduced capacity for release or

turnover of calcium. Intracellular Ca^{2+} signaling in neurons controls neurotransmitter secretion

among many other cellular processes and thus may be indirectly involved in dopamine or GABA

regulation. Some NDL-PCBs (PCB 95, 138) are known to inhibit voltage-gated calcium channels

(Langeveld et al., 2012) contributing to interference with calcium homeostasis and consequently

neurotransmission and synaptic plasticity (Brunelli et al., 2012; Campagna et al., 2011; Gafni et

al., 2003; Ta et al., 2006).

The startle response to threatening stimuli is vital for survival. PCB153-exposed zebrafish larvae

predominantly responded with a LLC, which not only delays the response, increasing the

chance to be caught by a predator, but also usually does not displace the animal from its

original location (**Figure 2A** and Jain et al., 2018). While survival of predator attack is likely

context-dependent and species-specific, escaping fast and away from a predator increases

survival in the field (McCormick et al., 2018). This suggests that exposure to NDL-PCBs may

alter selection pressure and thereby shift population structure. We have observed a significantly

delayed startle response at concentrations as low as 10 nM (3.6 µg L^{-1}). PCBs are associated

with sediments, where concentrations of PCB153 frequently reach multiple µg kg^{-1} (Lai et al.,

2015). PCB concentrations in organisms may be even higher due to extensive bioconcentration

(Killifish from New Bedford Harbor have 6.5 to 7 µg of PCB153 per g wet weight; Gräns et al.

2015). All NDL-PCBs tested in this study increased the startle latency. Thus, in the environment,

where PCBs occur in mixtures of numerous congeners that likely have additive effects, the

effect concentration is suggested to be lower than that of single congeners. Taken together,

these results highlight the importance of the neurotoxic potential of NDL-PCBs in an ecological

context.

In conclusion, the comprehensive approach presented here suggests that NDL-PCBs alter

dopamine signaling, leading to an increase in GABA that ultimately results in swelling of axonal

terminals in the auditory circuit. While the involvement of connexins remains speculative, our

findings provide a foothold into the dopamine dysregulating potential of NDL-PCBs in sensory-

motor processing in developing vertebrates, and link that to motor behavior. In a direct

comparison, the present study demonstrates that the adverse behavioral effects are induced by

all NDL-PCBs tested, but not by the classical DL-PCB126. Whether this is true for compounds

that are structurally similar to NDL-PCBs, such as some PBDEs, is not known and will require

further research. A key finding is that the effect of NDL-PCBs on the startle latency occurs at

concentrations relevant for both environmental and human exposures, and therefore may

contribute to the selection pressure in the environment and neurotoxicological disease in

human. This study provides valuable clues for tackling health effects in humans and wildlife

caused by environmental exposure.

CRediT authorship contribution statement

Nadja R. Brun: Conceptualization, Methodology, Formal analysis, Investigation, Writing - original

draft, Writing - review & editing, Visualization, Funding acquisition. **Jennifer M. Panlilio:**

Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing - review &

editing, Funding acquisition. **Kun Zhang:** Methodology, Investigation, Writing - review & editing,

Funding acquisition. **Yanbin Zhao:** Methodology, Investigation, Writing - review & editing,

Funding acquisition. **Evgeny Ivashkin:** Methodology, Investigation. **John J. Stegeman:**

Conceptualization, Writing - review & editing, Funding acquisition. **Jared V. Goldstone:**

Conceptualization, Writing - review & editing, Project administration.

Competing Financial Interests

The authors declare they have no actual or potential competing financial interests.

Funding

This work was supported by the Swiss National Science Foundation P2EZP2_165200 (NRB), the

Boston University Superfund Research Program NIH 5P42ES007381 (JJS and JVG), the Woods

Hole Center for Oceans and Human Health (NIH: P01ES021923 and P01ES028938; NSF: OCE-

1314642 and OCE-1840381) (NRB, JMP and JJS), and the National Natural Science Foundation

of China 22006099 (KZ and YZ) and the Shanghai Pujiang Program 19PJ1404900 (KZ and YZ).

Acknowledgments

The authors would like to thank Neel Aluru for helpful discussion and providing the confocal

microscope and Ed Levin and Mark Hahn for critically reading the manuscript and making

valuable suggestions.

References

Almeida, R.G., Czopka, T., Ffrench-Constant, C., Lyons, D.A., 2011. Individual axons regulate
the myelinating potential of single oligodendrocytes in vivo. Development 138, 4443–50.
https://doi.org/10.1242/dev.071001

Aluru, N., Krick, K.S., Mcdonald, A.M., Karchner, S.I., 2020. Developmental exposure to
PCB153 (2,2′,4,4′,5,5′-hexachlorobiphenyl) alters circadian rhythms and the expression of
clock and metabolic genes. Toxicol. Sci. 173, 41–52. https://doi.org/10.1093/toxsci/kfz217

Bager, Y., Lindebro, M.C., Martel, P., Chaumontet, C., Wärngård, L., 1997. Altered function,
localization and phosphorylation of gap junctions in rat liver epithelial, IAR 20, cells after
treatment with PCBs or TCDD. Environ. Toxicol. Pharmacol. 3, 257–266.
https://doi.org/10.1016/S1382-6689(97)00021-5

Bates, D., Mächler, M., Bolker, B., Walker, S., 2015. Fitting Linear Mixed-Effects Models using lme4. J. Stat. Softw. 67, 1–48. https://doi.org/10.18637/jss.v067.i01

Berger, D.F., Lombardo, J.P., Jeffers, P.M., Hunt, A.E., Bush, B., Casey, A., Quimby, F., 2001. Hyperactivity and impulsiveness in rats fed diets supplemented with either Aroclor 1248 or PCB-contaminated St. Lawrence river fish. Behav. Brain Res. 126, 1–11. https://doi.org/10.1016/S0166-4328(01)00244-3

Bergonzi, R., Specchia, C., Dinolfo, M., Tomasi, C., De Palma, G., Frusca, T., Apostoli, P., 2009. Distribution of persistent organochlorine pollutants in maternal and foetal tissues: Data from an Italian polluted urban area. Chemosphere 76, 747–754. https://doi.org/10.1016/j.chemosphere.2009.05.026

Boix, J., Cauli, O., Felipo, V., 2010. Developmental exposure to polychlorinated biphenyls 52, 138 or 180 affects differentially learning or motor coordination in adult rats. Mechanisms involved. Neuroscience 167, 994–1003. https://doi.org/10.1016/j.neuroscience.2010.02.068

Boucher, O., Muckle, G., Bastien, C.H., 2009. Prenatal exposure to polychlorinated biphenyls: A neuropsychologic analysis. Environ. Health Perspect. 117, 7–16. https://doi.org/10.1289/ehp.11294

Brunelli, L., LLansola, M., Felipo, V., Campagna, R., Airoldi, L., De Paola, M., Fanelli, R., Mariani, A., Mazzoletti, M., Pastorelli, R., 2012. Insight into the neuroproteomics effects of the food-contaminant non-dioxin like polychlorinated biphenyls. J. Proteomics 75, 2417–2430. https://doi.org/10.1016/J.JPROT.2012.02.023

Burgess, H.A., Granato, M., 2007. Sensorimotor gating in larval zebrafish. J. Neurosci. 27, 4984–4994. https://doi.org/10.1523/JNEUROSCI.0615-07.2007

Campagna, R., Brunelli, L., Airoldi, L., Fanelli, R., Hakansson, H., Heimeier, R.A., De Boever, P., Boix, J., Llansola, M., Felipo, V., Pastorelli, R., 2011. Cerebellum Proteomics addressing the cognitive deficit of rats perinatally exposed to the food-relevant polychlorinated biphenyl 138. Toxicol. Sci. 123, 170–179. https://doi.org/10.1093/toxsci/kfr156

Castoldi, A.F., Blandini, F., Randine, G., Samuele, A., Manzo, L., Coccini, T., 2006. Brain monoaminergic neurotransmission parameters in weanling rats after perinatal exposure to methylmercury and 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153). Brain Res. 1112, 91–98. https://doi.org/10.1016/j.brainres.2006.07.022

Cesetti, T., Ciccolini, F., Li, Y., 2012. GABA not only a neurotransmitter: Osmotic regulation by GABA AR signaling. Front. Cell. Neurosci. https://doi.org/10.3389/fncel.2012.00003

Chen, Y.-C.J., Guo, Y.-L., Hsu, C.-C., Rogan, W.J., 1992. Cognitive development of Yu-Cheng ('Oil Disease') children prenatally exposed to heat-degraded PCBs. JAMA J. Am. Med. Assoc. 268, 3213. https://doi.org/10.1001/jama.1992.03490220057028

Coccini, T., Roda, E., Castoldi, A.F., Poli, D., Goldoni, M., Vettori, M.V., Mutti, A., Manzo, L., 2011. Developmental exposure to methylmercury and 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153) affects cerebral dopamine D1-like and D2-like receptors of weanling and

pubertal rats. Arch. Toxicol. 85, 1281–1294. https://doi.org/10.1007/s00204-011-0660-y

Coleman, M., 2005. Axon degeneration mechanisms: Commonality amid diversity. Nat. Rev.
Neurosci. https://doi.org/10.1038/nrn1788

Crofton, K.M., Ding, D.-L., Padich, R., Taylor, M., Henderson, D., 2000. Hearing loss following
exposure during development to polychlorinated biphenyls: A cochlear site of action. Hear.
Res. 144, 196–204. https://doi.org/10.1016/S0378-5955(00)00062-9

Dervola, K.S.N., Johansen, E.B., Walaas, S.I., Fonnum, F., 2015. Gender-dependent and
genotype-sensitive monoaminergic changes induced by polychlorinated biphenyl 153 in the
rat brain. Neurotoxicology 50, 38–45. https://doi.org/10.1016/J.NEURO.2015.07.004

Dong, A., Liu, S., Li, Y., 2018. Gap junctions in the nervous system: Probing functional
connections using new imaging approaches. Front. Cell. Neurosci. 12, 1–9.
https://doi.org/10.3389/fncel.2018.00320

Enayah, S.H., Vanle, B.C., Fuortes, L.J., Doorn, J.A., Ludewig, G., 2018. PCB95 and PCB153
change dopamine levels and turn-over in PC12 cells. Toxicology 394, 93–101.
https://doi.org/10.1016/j.tox.2017.12.003

Ennaceur, S., Gandoura, N., Driss, M.R., 2008. Distribution of polychlorinated biphenyls and
organochlorine pesticides in human breast milk from various locations in Tunisia: Levels of
contamination, influencing factors, and infant risk assessment. Environ. Res. 108, 86–93.
https://doi.org/10.1016/J.ENVRES.2008.05.005

Faucherre, A., Baudoin, J.P., Pujol-Martí, J., López-Schier, H., 2010. Multispectral four-
dimensional imaging reveals that evoked activity modulates peripheral arborization and the
selection of plane-polarized targets by sensory neurons. Development 137, 1635–1643.
https://doi.org/10.1242/dev.047316

Fernandes, E.C.A., Hendriks, H.S., Van Kleef, R.G.D.M., Van Den Berg, M., Westerink, R.H.S.,
2010. Potentiation of the human GABAA receptor as a novel mode of action of lower-
chlorinated non-dioxin-like PCBs. Environ. Sci. Technol. 44, 2864–2869.
https://doi.org/10.1021/es902321a

Fonnum, F., Mariussen, E., 2009. Mechanisms involved in the neurotoxic effects of
environmental toxicants such as polychlorinated biphenyls and brominated flame
retardants. J. Neurochem. 111, 1327–1347. https://doi.org/10.1111/j.1471-
4159.2009.06427.x

Fosque, B.F., Sun, Y., Dana, H., Yang, C., Ohyama, T., Tadross, M.R., Patel, R., Zlatic, M.,
Kim, D.S., Ahrens, M.B., Jayaraman, V., Looger, L.L., Schreiter, E.R., 2015. Labeling of
active neural circuits in vivo with designed calcium integrators. Science (80-.). 347, 755–
760.

Gafni, J., Wong, P.W., Pessah, I.N., 2003. Non-coplanar 2,2',3,5',6-pentachlorobiphenyl (PCB
95) amplifies ionotropic glutamate receptor signaling in embryonic cerebellar granule
neurons by a mechanism involving ryanodine receptors. Toxicol. Sci. 77, 72–82.
https://doi.org/10.1093/toxsci/kfh004

Gleason, M.R., Nagiel, A., Jamet, S., Vologodskaia, M., López-Schier, H., Hudspeth, A.J., 2009.
The transmembrane inner ear (Tmie) protein is essential for normal hearing and balance in
the zebrafish. Proc. Natl. Acad. Sci. U. S. A. 106, 21347–21352.
https://doi.org/10.1073/pnas.0911632106

Goldey, E.S., Kehn, L.S., Lau, C., Rehnberg, G.L., Crofton, K.M., 1995. Developmental
exposure to polychlorinated biphenyls (Aroclor 1254) reduces circulating thyroid hormone
concentrations and causes hearing deficits in rats. Toxicol. Appl. Pharmacol. 135, 77–88.
https://doi.org/10.1006/TAAP.1995.1210

Grandjean, P., Landrigan, P.J., 2014. Neurobehavioural effects of developmental toxicity.
Lancet Neurol. 13, 330–338. https://doi.org/10.1016/S1474-4422(13)70278-3

Gräns, J., Wassmur, B., Fernández-Santoscoy, M., Zanette, J., Woodin, B.R., Karchner, S.I.,
Nacci, D.E., Champlin, D., Jayaraman, S., Hahn, M.E., Stegeman, J.J., Celander, M.C.,
2015. Regulation of pregnane-X-receptor, CYP3A and P-glycoprotein genes in the PCB-
resistant killifish (Fundulus heteroclitus) population from New Bedford Harbor. Aquat.
Toxicol. 159, 198–207. https://doi.org/10.1016/j.aquatox.2014.12.010

Herbstman, J.B., Sjödin, A., Apelberg, B.J., Witter, F.R., Patterson, D.G., Halden, R.U., Jones,
R.S., Park, A., Zhang, Y., Heidler, J., Needham, L.L., Goldman, L.R., 2007. Determinants
of prenatal exposure to polychlorinated biphenyls (PCBs) and polybrominated diphenyl
ethers (PBDEs) in an urban population. Environ. Health Perspect. 115, 1794–1800.
https://doi.org/10.1289/ehp.10333

Hoerbelt, P., Lindsley, T.A., Fleck, M.W., 2015. Dopamine directly modulates GABAA receptors.
J. Neurosci. 35, 3525–3536. https://doi.org/10.1523/JNEUROSCI.4390-14.2015

Howe, K., Clark, M.D., Torroja, C.F., Torrance, J., Berthelot, C., Muffato, M., Collins, J.E.,
Humphray, S., McLaren, K., Matthews, L., McLaren, S., Sealy, I., Caccamo, M., Churcher,
C., Scott, C., Barrett, J.C., Koch, R., Rauch, G.-J., White, S., Chow, W., Kilian, B., Quintais,
L.T., Guerra-Assunção, J. a, Zhou, Y., Gu, Y., Yen, J., Vogel, J.-H., Eyre, T., Redmond, S.,
Banerjee, R., Chi, J., Fu, B., Langley, E., Maguire, S.F., Laird, G.K., Lloyd, D., Kenyon, E.,
Donaldson, S., Sehra, H., Almeida-King, J., Loveland, J., Trevanion, S., Jones, M., Quail,
M., Willey, D., Hunt, A., Burton, J., Sims, S., McLay, K., Plumb, B., Davis, J., Clee, C.,
Oliver, K., Clark, R., Riddle, C., Elliot, D., Eliott, D., Threadgold, G., Harden, G., Ware, D.,
Begum, S., Mortimore, B., Mortimer, B., Kerry, G., Heath, P., Phillimore, B., Tracey, A.,
Corby, N., Dunn, M., Johnson, C., Wood, J., Clark, S., Pelan, S., Griffiths, G., Smith, M.,
Glithero, R., Howden, P., Barker, N., Lloyd, C., Stevens, C., Harley, J., Holt, K.,
Panagiotidis, G., Lovell, J., Beasley, H., Henderson, C., Gordon, D., Auger, K., Wright, D.,
Collins, J., Raisen, C., Dyer, L., Leung, K., Robertson, L., Ambridge, K., Leongamornlert,
D., McGuire, S., Gilderthorp, R., Griffiths, C., Manthravadi, D., Nichol, S., Barker, G.,
Whitehead, S., Kay, M., Brown, J., Murnane, C., Gray, E., Humphries, M., Sycamore, N.,
Barker, D., Saunders, D., Wallis, J., Babbage, A., Hammond, S., Mashreghi-Mohammadi,
M., Barr, L., Martin, S., Wray, P., Ellington, A., Matthews, N., Ellwood, M., Woodmansey,
R., Clark, G., Cooper, J.D., Cooper, J., Tromans, A., Grafham, D., Skuce, C., Pandian, R.,
Andrews, R., Harrison, E., Kimberley, A., Garnett, J., Fosker, N., Hall, R., Garner, P., Kelly,
D., Bird, C., Palmer, S., Gehring, I., Berger, A., Dooley, C.M., Ersan-Ürün, Z., Eser, C.,
Geiger, H., Geisler, M., Karotki, L., Kirn, A., Konantz, J., Konantz, M., Oberländer, M.,
Rudolph-Geiger, S., Teucke, M., Lanz, C., Raddatz, G., Osoegawa, K., Zhu, B., Rapp, A.,

Widaa, S., Langford, C., Yang, F., Schuster, S.C., Carter, N.P., Harrow, J., Ning, Z.,
Herrero, J., Searle, S.M.J., Enright, A., Geisler, R., Plasterk, R.H. a, Lee, C., Westerfield,
M., de Jong, P.J., Zon, L.I., Postlethwait, J.H., Nüsslein-Volhard, C., Hubbard, T.J.P.,
Roest Crollius, H., Rogers, J., Stemple, D.L., 2013. The zebrafish reference genome
sequence and its relationship to the human genome. Nature 496, 498–503.
https://doi.org/10.1038/nature12111

Hudspeth, A., 1985. The cellular basis of hearing: the biophysics of hair cells. Science (80-.).
230, 745–752. https://doi.org/10.1126/science.7414321

Inoue, D., Wittbrodt, J., 2011. One for all-a highly efficient and versatile method for fluorescent
immunostaining in fish embryos. PLoS One 6, 1–7.
https://doi.org/10.1371/journal.pone.0019713

Jain, R.A., Wolman, M.A., Marsden, K.C., Nelson, J.C., Shoenhard, H., Echeverry, F.A., Szi, C.,
Bell, H., Skinner, J., Cobbs, E.N., Sawada, K., Zamora, A.D., Pereda, A.E., Granato, M.,
2018. A forward genetic screen in zebrafish identifies the G-protein-coupled receptor CaSR
as a modulator of sensorimotor decision making. Curr. Biol. 28, 1357-1369.e5.
https://doi.org/10.1016/j.cub.2018.03.025

Johansen, E.B., Fonnum, F., Lausund, P.L., Walaas, S.I., Bærland, N.E., Wøien, G.,
Sagvolden, T., 2014. Behavioral changes following PCB 153 exposure in the
Spontaneously Hypertensive rat - an animal model of Attention-Deficit/Hyperactivity
disorder. Behav. Brain Funct. 10, 1–19. https://doi.org/10.1186/1744-9081-10-1

Jusko, T.A., Sisto, R., Iosif, A.-M., Moleti, A., Wimmerová, S., Lancz, K., Tihányi, J., Šovčiková,
E., Drobná, B., Palkovičová, L., Jurečková, D., Thevenet-Morrison, K., Verner, M.-A.,
Sonneborn, D., Hertz-Picciotto, I., Trnovec, T., 2014. Prenatal and postnatal serum PCB
concentrations and cochlear function in children at 45 months of age. Environ. Health
Perspect. 122, 1246–1252. https://doi.org/10.1289/ehp.1307473

Kang, K.-S., Park, J.-E., Ryu, D.-Y., Lee, Y.-S., 2001. Effects and neuro-toxic mechanisms of
2,2',4,4',5,5'-hexachlorobiphenyl and endosulfan in neuronal stem cells. J. Vet. Med. Sci.
63, 1183–1190. https://doi.org/10.1292/jvms.63.1183

Kang, K.S., Wilson, M.R., Hayashi, T., Chang, C.C., Trosko, J.E., 1996. Inhibition of gap
junctional intercellular communication in normal human breast epithelial cells after
treatment with pesticides, PCBs, and PBBs, alone or in mixtures. Environ. Health Perspect.
104, 192–200. https://doi.org/10.1289/ehp.96104192

Kimmel, C.B., 1993. Patterning the brain of the zebrafish embryo. Annu. Rev. Neurosci. 16,
707–732. https://doi.org/10.1146/annurev.ne.16.030193.003423

Lacoste, A.M.B., Schoppik, D., Robson, D.N., Haesemeyer, M., Portugues, R., Li, J.M.,
Randlett, O., Wee, C.L., Engert, F., Schier, A.F., 2015. A convergent and essential
interneuron pathway for mauthner-cell-mediated escapes. Curr. Biol. 25, 1526–1534.
https://doi.org/10.1016/j.cub.2015.04.025

Lai, Z., Li, X., Li, H., Zhao, L., Zeng, Y., Wang, C., Gao, Y., Liu, Q., 2015. Residual distribution
and risk assessment of polychlorinated biphenyls in surface sediments of the Pearl River

Delta, South China. Bull. Environ. Contam. Toxicol. 95, 37–44.
https://doi.org/10.1007/s00128-015-1563-z

Lancz, K., Hertz-Picciotto, I., Jusko, T.A., Murínová, Ľ., Wimmerová, S., Šovčíková, E., Dedík, L., Strémy, M., Drobná, B., Farkašová, D., Trnovec, T., 2015a. Duration of breastfeeding and serum PCB 153 concentrations in children. Environ. Res. 136, 35–39. https://doi.org/10.1016/j.envres.2014.09.036

Lancz, K., Palkovičováa, L., Patayová, H., Drobná, B., Wimmerová, S., Šovčíková, E., Kováč, J., Farkašová, D., Hertz-Picciotto, I., Jusko, T.A., Trnovec, T., 2015b. Ratio of cord to maternal serum PCB concentrations in relation to their congener-specific physicochemical properties. Int J Hyg Env. Heal. 218, 91–98. https://doi.org/doi:10.1016/j.ijheh.2014.08.003

Landrigan, P.J., Fuller, R., Acosta, N.J.R., Adeyi, O., Arnold, R., Basu, N., Baldé, A.B., Bertollini, R., Bose-O'Reilly, S., Boufford, J.I., Breysse, P.N., Chiles, T., Mahidol, C., Coll-Seck, A.M., Cropper, M.L., Fobil, J., Fuster, V., Greenstone, M., Haines, A., Hanrahan, D., Hunter, D., Khare, M., Krupnick, A., Lanphear, B., Lohani, B., Martin, K., Mathiasen, K. V., McTeer, M.A., Murray, C.J.L., Ndahimananjara, J.D., Perera, F., Potočnik, J., Preker, A.S., Ramesh, J., Rockström, J., Salinas, C., Samson, L.D., Sandilya, K., Sly, P.D., Smith, K.R., Steiner, A., Stewart, R.B., Suk, W.A., van Schayck, O.C.P., Yadama, G.N., Yumkella, K., Zhong, M., 2018. The Lancet Commission on pollution and health. Lancet 391, 462–512. https://doi.org/10.1016/S0140-6736(17)32345-0

Langeveld, W.T., Meijer, M., Westerink, R.H.S., 2012. Differential effects of 20 non-dioxin-like PCBs on basal and depolarization-evoked intracellular calcium levels in PC12 cells. Toxicol. Sci. 126, 487–496. https://doi.org/10.1093/toxsci/kfr346

Lein, P.J., Yang, D., Bachstetter, A.D., Tilson, H.A., Harry, G.J., Mervis, R.F., Kodavanti, P.R.S., 2007. Ontogenetic alterations in molecular and structural correlates of dendritic growth after developmental exposure to polychlorinated biphenyls. Environ. Health Perspect. 115, 556–563. https://doi.org/10.1289/ehp.9773

Lilienthal, H., Heikkinen, P., Andersson, P.L., van der Ven, L.T.M., Viluksela, M., 2011. Auditory effects of developmental exposure to purity-controlled polychlorinated biphenyls (PCB52 and PCB180) in rats. Toxicol. Sci. 122, 100–111. https://doi.org/10.1093/toxsci/kfr077

Lilienthal, H., Korkalainen, M., Andersson, P.L., Viluksela, M., 2015. Developmental exposure to purity-controlled polychlorinated biphenyl congeners (PCB74 and PCB95) in rats: Effects on brainstem auditory evoked potentials and catalepsy. Toxicology 327, 22–31. https://doi.org/10.1016/J.TOX.2014.11.004

Liu, Y.C., Hale, M.E., 2017. Local spinal cord circuits and bilateral mauthner cell activity function together to drive alternative startle behaviors. Curr. Biol. 27, 697–704. https://doi.org/10.1016/j.cub.2017.01.019

Machala, M., Bláha, L., Vondráček, J., Trosko, J.E., Scott, J., Upham, B.L., 2003. Inhibition of gap junctional intercellular communication by noncoplanar polychlorinated biphenyls: Inhibitory potencies and screening for potential Mode(s) of Action. Toxicol. Sci. 76, 187–195. https://doi.org/10.1093/TOXSCI

Magnusson, O., Mohringe, B., Thorell, G., Lake-Bakaar, D.M., 1987. Effects of the dopamine D2
selective receptor antagonist remoxipride on dopamine turnover in the rat brain after acute
and repeated administration. Pharmacol. Toxicol. 60, 368–373.
https://doi.org/10.1111/j.1600-0773.1987.tb01529.x

Mariussen, E., Fonnum, F., 2001. The effect of polychlorinated biphenyls on the high affinity
uptake of the neurotransmitters, dopamine, serotonin, glutamate and GABA, into rat brain
synaptosomes. Toxicology 159, 11–21. https://doi.org/10.1016/S0300-483X(00)00374-7

Maximino, C., Herculano, A.M., 2010. A review of monoaminergic neuropsychopharmacology in
zebrafish. Zebrafish 7, 359–378. https://doi.org/10.1089/zeb.2010.0669

McCormick, M.I., Fakan, E., Allan, B.J.M., 2018. Behavioural measures determine survivorship
within the hierarchy of whole-organism phenotypic traits. Funct. Ecol. 32, 958–969.
https://doi.org/10.1111/1365-2435.13033

Medan, V., Preuss, T., 2014. The Mauthner-cell circuit of fish as a model system for startle
plasticity. J. Physiol. Paris 108, 129–140. https://doi.org/10.1016/j.jphysparis.2014.07.006

Meserve, J.H., Nelson, J.C., Marsden, K.C., Hsu, J., Echeverry, F.A., Jain, A., Wolman, M.A.,
Pereda, A.E., Granato, M., 2020. A forward genetic screen identifies Dolk as a regulator of
startle magnitude through the potassium channel subunit Kv1.1. bioRxiv.

Meyers, J.R., MacDonald, R.B., Duggan, A., Lenzi, D., Standaert, D.G., Corwin, J.T., Corey,
D.P., 2003. Lighting up the senses: FM1-43 loading of sensory cells through nonselective
ion channels. J. Neurosci. 23, 4054–65. https://doi.org/10.1523/JNEUROSCI.23-10-
04054.2003

Nakayama, H., 2004. Common sensory inputs and differential excitability of segmentally
homologous reticulospinal neurons in the hindbrain. J. Neurosci. 24, 3199–3209.
https://doi.org/10.1523/JNEUROSCI.4419-03.2004

Nyffeler, J., Chovancova, P., Dolde, X., Holzer, A.K., Purvanov, V., Kindinger, I., Kerins, A.,
Higton, D., Silvester, S., van Vugt-Lussenburg, B.M.A., Glaab, E., van der Burg, B.,
Maclennan, R., Legler, D.F., Leist, M., 2018. A structure–activity relationship linking non-
planar PCBs to functional deficits of neural crest cells: new roles for connexins. Arch.
Toxicol. 92, 1225–1247. https://doi.org/10.1007/s00204-017-2125-4

Oikonomou, G., Altermatt, M., Zhang, R. wei, Coughlin, G.M., Montz, C., Gradinaru, V., Prober,
D.A., 2019. The Serotonergic Raphe Promote Sleep in Zebrafish and Mice. Neuron 103,
686-701.e8. https://doi.org/10.1016/j.neuron.2019.05.038

Panlilio, J.M., Aluru, N., Hahn, M.E., 2020. Developmental Neurotoxicity of the Harmful Algal
Bloom Toxin Domoic Acid: Cellular and Molecular Mechanisms Underlying Altered
Behavior in the Zebrafish Model. Environ. Health Perspect. 128, 117002.
https://doi.org/10.1289/EHP6652

Panlilio, J.M., Jones, I.T., Salanga, M.C., Aluru, N., Hahn, M.E., 2021. Developmental exposure
to domoic acid disrupts startle response behavior and circuitry. bioRxiv.
https://doi.org/10.1101/2021.01.08.425996

Park, H.-C., Shin, J., Roberts, R.K., Appel, B., 2007. An olig2 reporter gene marks
oligodendrocyte precursors in the postembryonic spinal cord of zebrafish. Dev. Dyn. 236,
3402–3407. https://doi.org/10.1002/dvdy.21365

Poon, E., Bandara, S.B., Allen, J.B., Sadowski, R.N., Schantz, S.L., 2015. Developmental PCB
exposure increases susceptibility to audiogenic seizures in adulthood. Neurotoxicology 46,
117–124. https://doi.org/10.1016/j.neuro.2014.12.007

Puel, J.L., Pujol, R., Ladrech, S., Eybalin, M., 1991. α-Amino-3-hydroxy-5-methyl-4-isoxazole
propionic acid electrophysiological and neurotoxic effects in the guinea-pig cochlea.
Neuroscience 45, 63–72. https://doi.org/10.1016/0306-4522(91)90103-U

Pujol-Martí, J., López-Schier, H., 2013. Developmental and architectural principles of the lateral-
line neural map. Front. Neural Circuits 7, 47. https://doi.org/10.3389/fncir.2013.00047

Ranasinghe, P., Thorn, R.J., Seto, R., Creton, R., Bridges, W.C., Chapman, S.C., Lee, C.M.,
2019. Embryonic exposure to 2,2′,3,5′,6-pentachlorobiphenyl (PCB-95) causes
developmental malformations in zebrafish. Environ. Toxicol. Chem. 39, 162–170.
https://doi.org/10.1002/etc.4587

Ribas-Fitó, N., Sala, M., Kogevinas, M., Sunyer, J., 2001. Polychlorinated biphenyls (PCBs) and
neurological development in children: A systematic review. J. Epidemiol. Community
Health 55, 537–546. https://doi.org/10.1136/jech.55.8.537

Rice, D., Barone, S., 2000. Critical periods of vulnerability for the developing nervous system:
Evidence from humans and animal models. Environ. Health Perspect. 108, 511–533.
https://doi.org/10.2307/3454543

Roberts, A., 2000. Early functional organization of spinal neurons in developing lower
vertebrates. Brain Res. Bull. https://doi.org/10.1016/S0361-9230(00)00392-0

Rungta, R.L., Choi, H.B., Tyson, J.R., Malik, A., Dissing-Olesen, L., Lin, P.J.C., Cain, S.M.,
Cullis, P.R., Snutch, T.P., Macvicar, B.A., 2015. The cellular mechanisms of neuronal
swelling underlying cytotoxic edema. Cell 161, 610–621.
https://doi.org/10.1016/j.cell.2015.03.029

Schantz, S.L., Widholm, J.J., Rice, D.C., 2003. Effects of PCB exposure on neuropsychological
function in children. Environ. Health Perspect. 111, 357–576.
https://doi.org/10.1289/ehp.5461

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch,
S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.Y., White, D.J., Hartenstein, V., Eliceiri,
K., Tomancak, P., Cardona, A., 2012. Fiji: An open-source platform for biological-image
analysis. Nat. Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

Sebe, J.Y., Cho, S., Sheets, L., Rutherford, M.A., von Gersdorff, H., Raible, D.W., 2017. Ca^{2+}-
permeable AMPARs mediate glutamatergic transmission and excitotoxic damage at the
hair cell ribbon synapse. J. Neurosci. 37, 6162–6175.
https://doi.org/10.1523/jneurosci.3644-16.2017

Seegal, R.F., Bush, B., Shain, W., 1990. Lightly chlorinated ortho-substituted PCB congeners
decrease dopamine in nonhuman primate brain and in tissue culture. Toxicol. Appl.
Pharmacol. 106, 136–44.

Serrano-Velez, J.L., Rodriguez-Alvarado, M., Torres-Vazquez, I.I., Fraser, S.E., Yasumura, T.,
Vanderpool, K.G., Rash, J.E., Rosa-Molinar, E., 2014. Abundance of gap junctions at
glutamatergic mixed synapses in adult Mosquitofish spinal cord neurons. Front. Neural
Circuits 8, 66. https://doi.org/10.3389/fncir.2014.00066

Shin, J., Park, H.-C., Topczewska, J.M., Mawdsley, D.J., Appel, B., 2003. Neural cell fate
analysis in zebrafish using <i>olig2<i> BAC transgenics. Methods Cell Sci. 25, 7–14.
https://doi.org/10.1023/B:MICS.0000006847.09037.3a

Smith, N.L., Kimelman, D., 2019. Establishing the body plan: The first 24 hours of zebrafish
development, The Zebrafish in Biomedical Research: Biology, Husbandry, Diseases, and
Research Applications. Elsevier. https://doi.org/10.1016/B978-0-12-812431-4.00007-5

Stenberg, M., Hamers, T., Machala, M., Fonnum, F., Stenius, U., Lauy, A.A., van Duursen,
M.B.M., Westerink, R.H.S., Fernandes, E.C.A., Andersson, P.L., 2011. Multivariate toxicity
profiles and QSAR modeling of non-dioxin-like PCBs - An investigation of in vitro screening
data from ultra-pure congeners. Chemosphere 85, 1423–1429.
https://doi.org/10.1016/j.chemosphere.2011.08.019

Stewart, A.M., Braubach, O., Spitsbergen, J., Gerlai, R., Kalueff, A. V., 2014. Zebrafish models
for translational neuroscience research: From tank to bedside. Trends Neurosci.
https://doi.org/10.1016/j.tins.2014.02.011

Ta, T.A., Feng, W., Molinski, T.F., Pessah, I.N., 2006. Hydroxylated xestospongins block
inositol-1,4,5-trisphosphate-induced Ca^{2+} release and sensitize Ca^{2+}-induced Ca^{2+} release
mediated by ryanodine receptors. Mol. Pharmacol. 69, 532–538.
https://doi.org/10.1124/mol.105.019125.used

Tabor, K.M., Bergeron, S.A., Horstick, E.J., Jordan, D.C., Aho, V., Porkka-Heiskanen, T.,
Haspel, G., Burgess, H.A., 2014. Direct activation of the Mauthner cell by electric field
pulses drives ultrarapid escape responses. J. Neurophysiol. 112, 834–844.
https://doi.org/10.1152/jn.00228.2014

Takada, N., Kucenas, S., Appel, B., 2010. Sox10 is necessary for oligodendrocyte survival
following axon wrapping. Glia 58, 996–1006. https://doi.org/10.1002/glia.20981

Tanaka, Y., Fujiwara, M., Shindo, A., Yin, G., Kitazawa, T., Teraoka, H., 2018. Aroclor 1254 and
BDE-47 inhibit dopaminergic function manifesting as changes in locomotion behaviors in
zebrafish embryos. Chemosphere 193, 1207–1215.
https://doi.org/10.1016/j.chemosphere.2017.11.138

Toro, C., Trapani, J.G., Pacentine, I., Maeda, R., Sheets, L., Mo, W., Nicolson, T., 2015.
Dopamine modulates the activity of sensory hair cells. J. Neurosci. 35, 16494–16503.
https://doi.org/10.1523/JNEUROSCI.1691-15.2015

Trnovec, T., Šovčíková, E., Pavlovčinová, G., Jakubíková, J., Jusko, T.A., Hus'ták, M.,

Jurečková, D., Palkovičová, L., Kočan, A., Drobná, B., Lancz, K., Wimmerová, S., 2010.
Serum PCB concentrations and cochlear function in 12-year-old children. Environ. Sci.
Technol. 44, 2884–2889. https://doi.org/10.1021/es901918h

Troconis, E.L., Ordoobadi, A.J., Sommers, T.F., Aziz-Bose, R., Carter, A.R., Trapani, J.G.,
2017. Intensity-dependent timing and precision of startle response latency in larval
zebrafish. J. Physiol. 595, 265–282. https://doi.org/10.1113/JP272466

Tropepe, V., Sive, H.L., 2003. Can zebrafish be used as a model to study the
neurodevelopmental causes of autism? Genes, Brain Behav.
https://doi.org/10.1034/j.1601-183X.2003.00038.x

Vettori, M.V., Goldoni, M., Caglieri, A., Poli, D., Folesani, G., Ceccatelli, S., Mutti, A., 2006.
Antagonistic effects of methyl-mercury and PCB153 on PC12 cells after a combined and
simultaneous exposure. Food Chem. Toxicol. 44, 1505–1512.
https://doi.org/10.1016/j.fct.2006.04.009

Voesenek, C.J., Muijres, F.T., Van Leeuwen, J.L., 2018. Biomechanics of swimming in
developing larval fish. J. Exp. Biol. 221. https://doi.org/10.1242/jeb.149583

Walkowiak, J., Wiener, J.-A., Fastabend, A., Heinzow, B., Krämer, U., Schmidt, E., Steingrüber,
H.-J., Wundram, S., Winneke, G., 2001. Environmental exposure to polychlorinated
biphenyls and quality of the home environment: effects on psychodevelopment in early
childhood. Lancet 358, 1602–1607. https://doi.org/10.1016/S0140-6736(01)06654-5

Wigestrand, M.B., Stenberg, M., Walaas, S.I., Fonnum, F., Andersson, P.L., 2013. Non-dioxin-
like PCBs inhibit [3H]WIN-35,428 binding to the dopamine transporter: A structure–activity
relationship study. Neurotoxicology 39, 18–24.
https://doi.org/10.1016/J.NEURO.2013.07.005

Wolman, M.A., Bell, H., Hogenesch, J.B., Granato, M., Jain, R.A., Skinner, J., Hayer, K.E.,
Marsden, K.C., 2015. A Genome-wide Screen Identifies PAPP-AA-Mediated IGFR
Signaling as a Novel Regulator of Habituation Learning. Neuron 85, 1200–1211.
https://doi.org/10.1016/j.neuron.2015.02.025

Yang, D., Kim, K.H., Phimister, A., Bachstetter, A.D., Ward, T.R., Stackman, R.W., Mervis, R.F.,
Wisniewski, A.B., Klein, S.L., Kodavanti, P.R.S., Anderson, K.A., Wayman, G., Pessah,
I.N., Lein, P.J., 2009. Developmental exposure to polychlorinated biphenyls interferes with
experience-dependent dendritic plasticity and ryanodine receptor expression in weanling
rats. Environ. Health Perspect. 117, 426–435. https://doi.org/10.1289/ehp.11771

Zhao, Y., Liang, J., Meng, H., Yin, Y., Zhen, H., Zheng, X., Shi, H., Wu, X., Zu, Y., Wang, B.,
Fan, L., Zhang, K., 2020. Rare earth elements lanthanum and praseodymium adversely
affect neural and cardiovascular eevelopment in zebrafish (Danio rerio). Environ. Sci.
Technol. https://doi.org/10.1021/acs.est.0c06632

1076