Neurological mechanism of sensory deficits after exposure to non-dioxin-like polychlorinated biphenyls (PCBs) in zebrafish larvae

Nadja R. Brun[1], Jennifer M. Panlilio[1], Kun Zhang[2,3], Yanbin Zhao[2,3], Evgeny Ivashkin[4,5], John J. Stegeman[1], Jared V. Goldstone[1]

[1] Department of Biology, Woods Hole Oceanographic Institution, Woods Hole, MA, USA

[2] School of Environmental Science and Engineering, Shanghai Jiao Tong University, 800 Dongchuan Road, Shanghai 200240, China

[3] Shanghai Institute of Pollution Control and Ecological Security, Shanghai 200092, China

[4] Josephine Bay Paul Center for Comparative Molecular Biology and Evolution, Marine Biological Laboratory, Woods Hole, MA, USA

[5] A.N. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Moscow, Russia

Address correspondence to

Nadja Brun, nbrun@whoi.edu

Woods Hole Oceanographic Institution, Mail Stop 32
266 Woods Hole Road
Woods Hole, MA 02543

Competing Financial Interests

The authors declare they have no actual or potential competing financial interests.

Abstract

Background: The most abundant polychlorinated biphenyl (PCB) congeners found in the environment and in humans are the *ortho*-substituted, non-dioxin-like (NDL) congeners, especially PCB153. While evidence indicates that exposure to PCB153 and other NDL-PCBs in early vertebrate development can contribute to neurobehavioral disorders, the basis of neurobiological effects of NDL-PCBs is poorly understood.

Methods: We assessed a range of effects of several NDL-PCB congeners (PCB153, PCB52, PCB118, and PCB138) on cellular morphologies and synaptic transmission linked to the proper execution of a sensorimotor response in zebrafish, taking advantage of the well-established neural circuit in this model. Specifically, we imaged hair cells, neurons, and muscle tissue to evaluate their presence and morphology. We measured neurotransmitter levels in larvae and assessed the mechanosensory startle response in co-exposures to PCB153 and neurotransmitter modulators to evaluate the involvement of neurotransmitters in the behavioral response. The startle response was also assessed in dioxin-like (DL)-PCB126 exposed larvae.

Results: Startle responses were dramatically delayed in 6-day old larvae exposed to NDL-PCB. In contrast, exposure to the DL PCB126 did not delay startle response times. Morphological and biochemical data showed that exposure to NDL-PCBs during development induces swelling of afferent sensory neurons and disrupts dopaminergic and GABAergic signaling associated with motor movement.

Discussion: This study demonstrates that connections between neurotoxic mechanisms and processing of sensory stimuli occur at low concentrations of NDL-PCBs, similar to those present in human and animal samples. The effects on important and broadly conserved signaling mechanisms in vertebrates suggest that NDL-PCBs can contribute to neurodevelopmental abnormalities, relevant for both exposed human populations and wildlife.

Introduction

Humans and wildlife are exposed to a wide range of contaminants in the environment, and the role of these chemical pollutants in developmental neurotoxicity, cognitive impairment, and neurodegenerative disorders has become a global environmental health concern (Landrigan et al. 2018). Among the top industrial chemicals thought to be involved in developmental neurotoxicity are polychlorinated biphenyls (PCBs) (Grandjean and Landrigan 2014), a mix of 209 different structures, including planar dioxin-like (DL) congeners and non-planar non-dioxin-like (NDL) congeners. PCBs are persistent legacy contaminants that have been produced and used for decades to serve as insulators, coatings, and caulks, and there is worldwide PCB contamination of wildlife tissue and human cord blood, amniotic fluid, and mother's milk (Ennaceur et al. 2008; Lancz et al. 2015a). The developing nervous system is generally sensitive to chemical exposure at doses much lower than those affecting adult nervous system function (Rice and Barone 2000). For instance, prenatal exposure to PCBs is associated with a decrease in motor skills in infants, while postnatal exposure seems to have no obvious effects (Ribas-Fitó et al. 2001). Thus, identifying the mechanisms underpinning the effects of neurotoxic PCBs during gestation and early life stages is of critical importance in attaining successful treatments and remediating health effects in humans and wildlife due to environmental exposure.

The NDL-PCBs are the most abundant congeners in human and environmental samples. The NDL-PCB138 and PCB153, are particularly abundant. Median PCB153 concentrations measured in human cord blood serum range from 6.4 to 110 ng g^{-1} lipid (Bergonzi et al. 2009; Herbstman et al. 2007; Lancz et al. 2015b), far greater than the levels of DL congeners. Evidence also suggests the NDL-PCBs contribute substantially to neurotoxic and behavioral effects (Boix et al. 2010; Boucher et al. 2009). However, understanding of the mechanisms and effects of NDL-PCBs, especially during the more sensitive early stages of development, significantly lags that of DL-PCBs.

Increased serum PCB concentrations in children are associated with deficits in cochlear

functions (Jusko et al. 2014; Trnovec et al. 2010), fine motor skills (Walkowiak et al. 2001),

and cognitive impairment (Boucher et al. 2009; Chen et al. 1992). Developmental exposure

of rats to NDL-PCBs has been linked to behavioral changes, including hyperactivity,

impulsiveness, and impairment of cognitive function (e.g., impairment of the ability to learn

the Y maze task) (Berger et al. 2001; Boix et al. 2010; Johansen et al. 2014). NDL-PCB

developmental exposure also adversely affects auditory function in adult rats, including

elevated thresholds of brainstem auditory evoked potentials, auditory deficits at low

frequencies, audiogenic seizures, and reduced auditory startle responses (Crofton et al.

2000; Goldey et al. 1995; Lilienthal et al. 2011, 2015; Poon et al. 2015). The mechanism(s)

involved in these effects on cognitive and auditory systems are not known.

Zebrafish are increasingly used as model organism to uncover the mechanisms of

neurotoxicity that are relevant for humans and other vertebrates. Zebrafish remain

transparent throughout early development, facilitating the use of transgenic and other

molecular approaches to label and observe specific classes of neurons and to follow their

functional development. Fish have relatively simple neural circuits, particularly for the

auditory system. In fish, the auditory-evoked startle response plays a critical role in animal

survival, and involves cellular functions, sensory-motor pathways, and monoamine

neurotransmitters that are conserved among vertebrates (Stewart et al. 2014). Many of the

pathways and genes that regulate zebrafish startle behavior also operate in human

neurological disorders (Howe et al. 2013; Maximino and Herculano 2010; Meserve et al.

2020; Tropepe and Sive 2003; Wolman et al. 2015). In this study, we use the zebrafish

startle response to decipher the neuro-disruptive basis of NDL-PCBs on the auditory system

during the early stages of development. To this end, we examined the effects of different

NDL-PCBs (PCB153, PCB52, PCB118, PCB138), in comparison to the DL-PCB126 and at

different environmentally relevant concentrations, dissecting various steps in the neural

circuitry underlying the acoustic startle response.

Materials and Methods

Animal model

In addition to other attributes, zebrafish have relatively simple neural circuits, particularly for the auditory system. Many of the components of the auditory circuit are homologous in mammals and fish, including hair cells, neural transmission, and lower motorneurons, while the mechanosensory lateral line system and Mauthner (M) cells are exclusively found in fish and a cochlea is only found in mammals. In fish, vibro-acoustic stimuli are first sensed by the hair cell bundles in the auditory, vestibular, and lateral-line systems and rapidly encoded into trains of action potentials in the afferent neurons (**Figure 1**) (Hudspeth 1985). The signal detected through the sensory hair cells of the lateral line is transmitted through the anterior and posterior lateral line ganglia whereas the auditory input is transmitted through the VIIIth statoacoustic nerve, all of which synapse to the lateral dendrite of the M-cell and two segmental homologs (Nakayama 2004; Pujol-Martí and López-Schier 2013). The signal is propagated down the M-cell axon in the spinal cord to activate the motoneurons leading to a unilateral contraction of the trunk musculature.

Zebrafish husbandry and early-life stage exposure

Zebrafish (*Danio rerio*) were maintained at 28.5 °C and on a 12:12 h light-dark cycle according to the guidelines of the Zebrafish Model Organism Database (ZFIN, http://zfin.org) and institutional guidelines at WHOI and the experimental procedures approved by the Institutional Animal Care and Use Committee (ID Number BI21981.01). Adult zebrafish were fed brine shrimp and pellets every day. In this study, we used wild-type zebrafish of the AB strain obtained from Zebrafish International Resource Center (ZIRC), as well as several transgenic lines: the photoconvertible calcium indicator line *Tg[elavl3:CaMPARI(W391F+V398L)]*[if9] kindly provided by Jessica Plavicki (Brown University, USA), and the transgenic lines *Tg(cntn1b:EGFP-CAAX)* (Panlilio et al. 2021) labeling axons, *Tg(olig2:EGFP)*[vu12] (Park et al. 2007; Shin et al. 2003) marking oligodendrocyte precursor cell bodies, *Tg(sox10:mRFP)* (Takada et al. 2010) marking oligodendrocyte lineage cells, as

well as *Tg(mbp:EGFP-CAAX)* (Almeida et al. 2011) *and Tg(mbp:EGFP)* (gift from Dr. Kelly

Monk, generated by Dr. Charles Kaufman in the laboratory of Dr. Leonard Zon, Harvard

Medical School, Boston, MA) marking myelin sheaths. Zebrafish eggs were obtained by

pairwise breeding and kept in 0.3x Danieau's (17 mM NaCl, 0.2 mM KCl, 0.12 mM $MgSO_4$,

0.18 mM $Ca(NO_3)_2$ and 1.5 mM HEPES, pH 7.6) in glass vials at a density of 10 individuals

per 10 mL, at a constant water temperature of 28 ± 1°C and standard 14-h light/10-h dark

cycle. The medium was supplemented with 1-Phenyl-2-thiourea (PTU, 0.003% w/v) from 24

hours post-fertilization (hpf) onwards to prevent pigment formation when necessary for

imaging. Fertilized eggs were used for exposures starting at 3 hpf to solvent control (0.1%

DMSO, v/v), 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153), 2,2′,5,5′-PCB, 2,2′,5,5′-

tetrachlorobiphenyl (PCB52), 2,3',4,4',5-pentachlorobiphenyl (PCB118), 2,2',3,4,4',5'-

hexachlorobiphenyl (PCB138), or 3,3',4,4',5-hexachlorobiphenyl (PCB126). Additionally, one

exposure scenario with PCB153 was started at 24 hpf, to assess whether effects are

induced during or after initial neuronal cell differentiation. Sensory neurons send their

projection along the ectoderm as early as 16 hpf (Kimmel 1993). The lowest exposure

concentration for PCB153 was 10 nM (3.6 ng mL^{-1}) which is estimated to result in ~5 ng g^{-1}

(5 ppm) per zebrafish larvae (comparing to PCB95 uptake levels; (Ranasinghe et al. 2019)).

PCB153 mean concentrations found in maternal cord blood serum are 0.27 ng mL^{-1} or 110

ng g^{-1} lipid (Lancz et al. 2015b)). Other concentrations used were 30, 100, and 1000 nM

which are found in fish of contaminated ecosystems (Gräns et al. 2015). For PCB52,

PCB118, and PCB138, a concentration of 1000 nM was chosen to compare to the highest

effect concentration of PCB153. The DL-PCB126 is cardioteratogenic in the low nanomolar

range. To avoid malformations, a concentration of 1 nM was chosen. Exposures were

refreshed daily until four days post-fertilization (dpf) and all larvae were kept in Danieau's

from 5 to 6 dpf. All imaging and behavioral assays were performed at 6 dpf, except for the

touch assay, which was performed at 26 hpf.

The influence of neurotransmitters on the startle behavior was assessed by exposures to

compounds known to interfere with monoamines. At 6 dpf, larvae were evaluated for their

startle behavior and then immersed in the dopamine precursors L-DOPA (1 mM) or L-

tyrosine (1 mM), dopamine receptor agonist quinpirole (10 µM), dopamine D2-receptor

antagonist and GABA receptor modulator haloperidol (10 µM), endogenous serotonin 5-HT

(10 µM), and serotonin precursor 5-HTP (1 mM) for 30 min in the well plate before assessing

the startle behavior again. Additionally, the effect of neurotransmitter modulation during

development on startle response was assessed by exposing larvae from 3 to 120 hpf to L-

DOPA (1 mM), L-tyrosine (1 mM), and quinpirole (10 µM).

Behavioral Analysis

Larval locomotion was monitored using the Noldus Daniovison system (Noldus Information

Technology, Leesburg, VA, USA) using 6-dpf larvae distributed in 48-well plates (1 larva per

well). After an acclimation period of 40 minutes in the illuminated chamber, spontaneous

swimming activity was tracked for 10 minutes. Following the spontaneous swimming, three

alternating cycles of dark and light conditions (10 minutes each) were run. The experiment

was repeated three times with cohorts from separate breeding events (n = 24 per group and

breeding event). Video data were recorded with 30 frames per second via an infrared

camera. Obtained data were analyzed with the supplied software EthoVision XT® 12

(Noldus. Inc.).

The activity of Rohon-Beard sensory neurons was tested at 26 hpf by lightly touching

DMSO- and 1000 nM PCB153-exposed embryos (n = 10 per group and breeding event)

using a piece of fishing line affixed to a glass pipette. Each embryo was touched ten times

and the response was scored as follows: 0, no response; 0.5, light tail flicking response, 1,

coiling in the opposite direction of the touch stimulus, yielding a score of maximum 10 per

embryo. The experiment was repeated twice with cohorts from separate breeding events.

Vibro-acoustic startle response was assessed as described previously (Panlilio et al. 2021).

Briefly, the acoustic stimuli were delivered by a vibrational exciter producing acoustic

vibrations at a frequency of a 1000 Hz for a duration of 2 ms at four different amplitudes (32,

38, 41, 43 dB). For each amplitude, the stimulus was delivered four times, spaced 20

seconds apart, to a 4 x 4 well-plate mounted on top of the speaker, each well containing an

individual larva. The response was tracked with a high-speed camera (Edgertronic SC1) at

1000 frames s^{-1} for 250 ms (13 ms before and 237 ms after the stimulus). PulsePal was used

to synchronize the camera recording to the acoustic stimuli using a TTL pulse. Automated

analysis of larval movement kinematics (response frequency, latency, turning angle) was

performed using the FLOTE software package (Burgess and Granato 2007). Each treatment

group was assessed at least three times with eggs from different clutches. The total replicate

numbers are stated in the **Table S1 (Supplemental Material)**.

A startle response in fish can be executed with a short latency C-bend (SLC) or long latency

C-bend (LLC), with the SLC being the more frequent response at higher stimulus intensities.

The cut-off time between SLC and LLC can vary depending on environmental conditions

such as temperature and has been determined empirically for this experiment using a

Gaussian mixture model (Panlilio et al. 2020)(**Figure S1**). Responses within 15 ms were

categorized as SLC, latencies greater than 15 ms as LLC. Following this classification, the

median response type (Relative Startle Bias) for each responsive fish was determined for

each stimulus intensity (32, 38, 41, and 43 dB). Bias per individual fish was calculated as:

(frequency of SLC – frequency of LLC) / total responses. Pure SLC responses result in a

value of +1, while pure LLC responses are a value of -1 (Jain et al. 2018).

Electric field pulses (EFPs) directly activate the M-cells (Tabor et al. 2014). Responsiveness

of M-cells was assessed by applying EFPs (4.4 V cm^{-1} for 2 ms, square pulse) to head-

restrained larvae (*n* = 46 per treatment group). Seven consecutive EFPs were applied with a

recovery time of 20 s in between pulses. For each run, one control and one exposed larva

were embedded in 1.5% agarose side-by-side and their tails freed. Response frequency,

latency, and bend angle were tracked using FLOTE. For all behavioral assays, only larvae

with an inflated swim bladder were used.

Larval growth

Larval length (from snout to the tail-fin end) was measured in DMSO (n = 57) and 1000 nM

PCB153 (n = 51) exposed larvae using ImageJ. The experiment was performed three times

with embryos from different clutches.

FM1-43 labeling in hair cells

The vital fluorescent dye FM1-43 (n-(3-triethylammoniumpropyl)-4-(4-(dibutylamino)-styryl)

pyridinium dibromide; Invitrogen) specifically labels hair cells in the lateral line and inner ear

of vertebrates of by entering through open mechanotransduction channels and filling the

cytoplasm of hair cells (Meyers et al. 2003). At 6 dpf, zebrafish larvae of 1000 nM PCB153-

exposed (n = 32) and unexposed (n = 28) groups were immersed in Danieau's

supplemented with 3 µM FM1-43 for 30 s, rinsed three times in Danieau's, anesthetized in 1

mM 3-amino-benzoic acid (Tricaine), and mounted in 1.5% low melting agarose for confocal

imaging of the L3 neuromast. The experiment was performed three times independently.

Immunohistochemistry to stain M-cells, kinocilia, and muscle fibers

Whole-mount immunohistochemistry of zebrafish larvae and brains was performed following

standard procedures. Briefly, 10 AB larvae per exposure group were treated as described

above and subsequently fixed in 4% paraformaldehyde (PFA) in PBS overnight at 4 ˚C. For

labelling brain tissue, larval brains were dissected before continuing with the protocol.

Multiple rinsing steps in PBS containing either 0.5% Triton-X100 or 0.1% Tween-20 (PBST)

for larvae or brains, respectively, was followed by an antigen retrieval step in 150 mM Tris

HCl (pH 9.0) at 70 °C (Inoue and Wittbrodt 2011), and a permeabilization step using

proteinase K (10 µg/mL for 3 minutes). Subsequently, the samples were blocked in blocking

solution (10% Normal Goat Serum, 1% DMSO, 1% BSA, 88% PBST) for 1 h and then

stained with blocking solution supplemented with the primary antibody for three days at 4 ˚C.

The primary anti-neurofilament mouse polyclonal antibody (3A10; Developmental Hybridoma

Bank, 1:100 dilution) was used to label a subset of hindbrain spinal cord projecting neurons

such as the M-cell neurons (DMSO n = 23, PCB153 n = 21). The anti-5HT rabbit polyclonal

antibody (Sigma-Aldrich, #S5545, 1:500) was used to label serotonergic cells (DMSO n = 16,

PCB153 n = 13). The anti-acetylated alpha-tubulin mouse polyclonal antibody (Sigma-

Aldrich, #T6793, 1:1000 dilution) was used to label kinocilia and hair cell innervating neurons

(DMSO n = 25, PCB153 n = 24). After several washing steps, in PBST, the samples were

incubated with Alexa Fluor 488- or 594-conjugated goat-anti-mouse or goat-anti-rabbit

secondary antibodies (Abcam, 1:500 dilution) overnight at 4 ˚C and rinsed several times in

PBST before mounting in ProLong Diamond Antifade mountant (Invitrogen) for imaging. For

visualization of skeletal muscle fibers (DMSO n = 11, PCB153 n = 11), larvae were fixed as

above, followed by permeabilization in 5% Triton-X for overnight and incubation in phalloidin

Alexa-546 conjugate (Invitrogen, 1:500). All staining experiments were repeated three times

with larvae from independent clutches except the phalloidin staining, which were performed

once.

Imaging of hair cell innervating neurons, primary motoneuron, and oligodendrocyte
lineage cells

The *Tg(cntn1b:EGFP-CAAX)* transgenic line was used to image the cell bodies of neurons

that innervate the hair cells at the L3 position in the lateral line (DMSO n = 22, PCB153 n =

20, two independent experiments). The axons of the main caudal primary motoneurons were

visualized using the *Tg(olig2:EGFP)* transgenic line (n = 4 per treatment group, one

experiment). The double transgenic *Tg(olig2:EGFP) x Tg(sox10:mRFP)* was used to quantify

oligodendrocyte lineage cells in the spinal cord of in DMSO (n = 22) and 1000 nM PCB153

(n = 16) treated larvae (Shin et al. 2003). All larvae were anesthetized in tricaine (0.16%

MS222) before mounting laterally in 1.5% low melting agarose on a glass-bottom microscopy

dish (MatTek). Fluorescently-labeled cell bodies were imaged at the distal end of the

intestine, around the L3 neuromast. The experiments were performed three times

independently.

Calcium imaging of freely swimming and startling larvae

Neuronal activity in DMSO- and PCB153-treated larvae was visualized using the CaMPARI

(*elavl3*:CaMPARI) transgenic zebrafish line with an engineered fluorescent protein that

permanently photoconverts from green to red in the presence of elevated calcium and UV

light (Fosque et al. 2015). Photoconversion was achieved by mounting an LED light (405 nm;

LEDSupply) between the speaker and a custom-made 3-well plate containing the freely

swimming larvae (DMSO- and PCB153-treated). Larvae of the same treatment were divided

into two batches: one was photoconverted in medium at room temperature (RT) and the

other one at 4˚C to elicit high neuronal activity. Photoconversion was achieved by shining the

LED light for 10 sec either with one vibrational stimulus of 43 dB at the start or without

stimulus. Subsequently, larvae were anesthetized, embedded in agarose, and whole brains

imaged using the 10x objective. The experiment was performed twice (n = 11-16 per

treatment group).

Image processing

Whole-brain, hair cells, M-cell, and motor neurons were imaged on a Zeiss LSM 800

confocal system with a 10x/0.45 Plan-Apochromat, 20x/0.8 Plan-Apochromat, or 63x/1.2 C-

Apochromat water immersion objective. Airy scan was applied for hair cell imaging at 63x. Z-

stacks were merged to a single plane by maximal intensity projection using Fiji (Schindelin et

al. 2012). The number of *olig2* cells dorsal of the spinal cord (DMSO n = 22, 1000 nM

PCB153 n = 16) was counted in confocal image stacks using the 3D projection of the Zeiss

Zen blue software. Myelination was evaluated after imaging the spinal cord (DMSO n = 26,

1000 nM PCB153 n = 27) using the 20x objective and scored for deficits according to Panlilio

et al. 2020. Representative myelination was imaged at 40x. The intensity of serotonergic

neurons in the brain (DMSO n = 16, 1000 nM PCB153 n = 13) was measured performing a

Sum Slices Z-projection followed by drawing a ROI around the 5-HT immuno-responsive

cells, and per image, three random squares (area size = 10.65 x 10.65) within the non-

fluorescent area of the ROI for background subtraction (for orientation see **Figure S12**).

Similarly, the FM1-43 intensity was measured drawing a ROI around the fluorescent hair

cells and measuring the background in three random squares (DMSO n = 20, 1000 nM

PCB153 n = 20). Corrected total cell fluorescence (CTCF) was calculated according to the

formula: Integrated density - (ROI x Mean fluorescence of background readings). For

quantification of active neurons in the brain of freely swimming and startling CaMPARI

zebrafish larvae, Sum Slices Z-projection were analyzed for red fluorescence intensity in the

rostral hindbrain (for orientation see **Figure S13**).

The volume of hair cell innervating neurons was calculated by measuring the volume of

threshold pixels (30-255, Otsu method) in image stacks using an imageJ macro looping

through the stack and summing the area measurements, then multiplying this sum by the

depth of each slice (DMSO n = 30, 1000 nM PCB153 n = 29). The presence of primary

motor neurons was confirmed using the Simple Neurite Tracer plugin in Fiji (DMSO n = 21,

1000 nM PCB153 n = 18). All image evaluations and quantifications were done without the

knowledge of the experimental condition of the specimens.

Neurotransmitter analysis

The analysis of neurotransmitters, including histamine, L-histidine, L-glutamic acid, gamma-

aminobutyric acid (GABA), acetylcholine, L-DOPA, 3-methoxytyramine, serotonin, 5-

hydroxyindole-3-acetic acid, glycine, L-glutamine, L-valine, 3-hydroxytyramine and choline,

was performed with a method developed in a previous study (Zhao et al. 2020). Briefly, the

zebrafish embryo samples were spiked with 100 ppb isotope-labeled internal standards and

then homogenized. The homogenized tissue samples were extracted with 200 µL of 80:20

v/v ACN/Milli-Q water containing 0.1% formic acid and 1% ascorbic acid and 200 µL ACN.

After protein precipitation, the supernatants were dried and redissolved in 100 µL of Milli-Q

water containing 0.1% formic acid for UHPLC-MS/MS analysis. Neurotransmitters were

quantified by a high-performance liquid chromatography system coupled to a Q-Exactive

plus quadrupole-Orbitrap mass spectrometer (UHPLC-Q-Orbitrap, Thermo fisher, US). The

separation of neurotransmitters was carried out with an Acquity UPLC HSS T3 column with

Milli-Q water containing 0.1% formic acid and ACN containing 0.1% formic acid as mobile

phase. The MS acquisition was performed in the full scan mode and parallel reaction

monitoring mode. The experiment was performed once (n = 4 per treatment group).

Statistical Analyses

All statistical analyses were conducted using Graph Pad Prism 8.4 except some of the startle

behavior analysis, which was completed in R. All quantitative data were tested for normality

using the Shapiro-Wilk test, and if there were unequal number of subjects between groups,

the data were tested for homogeneity in variances using the Brown-Forsythe test. All

treatment groups were compared to respective control groups.

Responsiveness to auditory stimuli (response or not) and startle type (SLC vs. LLC) was

modeled in R (version 4.0.2) using a binomial generalized linear mixed effect model (glmer()

from the lme4 R package) (Bates et al. 2015). The response to electrical stimulus was

modeled using glm() from the lme4 R package. For the auditory response, we included

treatment as a fixed effect and the replicate experimental trials as a random effect. The

model specification for "fraction responding" was as follows: Responders ~ TreatmentGroup

+ (TreatmentGroup|trial). The model specification for "startle bias" was as follows:

SLCorLLCresponse ~ TreatmentGroup + stimdB + (1|trial). The data were analyzed fitting

the model for each stimulus intensity independently. The model specification for the electrical

stimulation are as follows: Responders ~ TreatmentGroup.

To assess treatment differences of normal data with three or more groups (fluorescence

intensity CaMPARI), ANOVA comparisons were performed followed by a multiple

comparison using the Tukey's correction to adjust the critical values. A Tukey-corrected p-

value < .05 was considered statistically significant. For normal data with heterogeneity

within-group variances (LLC bend angle), a Welch ANOVA followed by a Dunnett's T3

multiple comparison test was performed. For non-normal data with three or more groups

(SLC bend angle), a Kruskal-Wallis followed by the Dunn's multiple comparison test was

performed. For normal data with two groups (length, mean latency, bend angle in e-stim,

oligodendrocyte lineage cell count, 5-HT intensity), an unpaired two-tailed t-test with Welch's

correction (Welch's t- test) was performed. The nonparametric scoring data of the touch

assay and latency in e-stim was analyzed using a two-tailed Mann-Whitney test. The data

are presented as mean ± SD with single data points (replicates) superimposed on the graph

unless specified in the figure legend.

Results

The startle response circuit is of wide relevance and well-established for assessing the

influence of specific factors on neuronal cell development, neuromuscular connectivity, and

transmission (Burgess and Granato 2007; Medan and Preuss 2014; Meserve et al. 2020;

Panlilio et al. 2020). We used the startle response circuit to assess the effects of NDL-PCBs

on the formation of neuronal populations and neurotransmission and its consequences for

mechanosensory disruptions, thus encompassing hearing and touch.

Behavioral effects

Rohon-Beard cells are the earliest mechanosensory neurons that develop. They are located

in the dorsal region of the embryonic spinal cord and appear around 24 hpf (Roberts 2000).

A few days into development, Rohon-Beard cells also relay sensory information to the M-

cells (Liu and Hale 2017). In our experiments, a tactile stimulus on the trunk at 26 hpf evoked

a C-shaped body flexion in both the DMSO- and the 1000 nM PCB153-treated embryos (U =

183, p = .599; **Figure S2**), indicating that the formation and function of Rohon-Beard cells

were not impaired by PCB153.

Exposure to contaminants during development can result in reduced growth, which can

affect kinematics in swimming behavior (Voesenek et al. 2018). PCB153-exposed larvae

developed normally with no difference in body length on day six in comparison to the control

group, in all three trials (Trial 1: $t(25.98)$ = .661, p = .514; Trial 2: $t(37.98)$ = .443, p = .660;

Trial 3: $t(37.43)$ = .645, p = .523; **Figure S3**). At 6 dpf, spontaneous and light-dark-stimulated

swimming activity was assessed. Larvae exposed to PCB 153 (1000 nM) had lower

cumulative locomotion activity over ten minutes (M = 1182 mm, SD = 382) in comparison to

the control group (M = 1510 mm, SD = 402), $t(132.4)$ = 4.854, p < .0001 (**Figure S4A**).

Zebrafish larvae typically increase locomotion in dark conditions following light exposure.

Once stimulated with a rapid loss of illumination during the dark phase, both treatment

groups exhibited similar swimming activity (**Figure S4B**).

The vibro-acoustic startle response can be executed either rapidly, within 15 ms (short

latency C-bend; SLC), or beginning at longer times and extending for up to 100 ms (long

latency C-bend; LLC). The mean latency value, representing the population response

including SLC and LLC responses, was 21 ± 20 ms for control larvae at the highest stimulus

intensity (**Figure S1**, **Figure 2A, B**). By contrast, the NDL-PCB-treated larvae that

responded to the vibro-acoustic stimulus showed significantly longer average latencies in

response to the same auditory stimulus given the controls, $F(5, 907$ = 14.61, p < .0001).

PCB153-treated (1000 nM) larvae responded at a mean latency of 68 ± 32 ms, PCB138 at

52 ± 37 ms, PCB118 at 45 ± 27 ms, and PCB52 at 39 ± 32 ms. For PCB153, a dose-

dependent response was observed, with mean latencies increasing with dose (10 nM: 26 ±

27 ms, 30 nM: 50 ± 36 ms, 100 nM: 50 ± 30 ms, 1000 nM: 68 ± 32 ms). The lowest PCB153

concentration tested (10 nM) was not significantly different from controls. Unlike NDL-PCB

treated larvae, the DL-PCB126-treated larvae had a mean latency (22 ± 22 ms), which was

not significantly different from controls.

Examining the SLC and LLC responses separately can give insight into the underlying

mechanism of effect. A greater stimulus intensity increases the probability of evoking SLC,

shifting the response bias from LLC to SLC (Troconis et al. 2017). Here, we also observed a

shift to SLC with increasing stimulus intensity in solvent control (DMSO-treated) larvae

(**Figure 2C, Figure S5A**). Larvae were tested four times at each stimulus intensity; thus, a

larva is categorized as having SLC bias if it responded 3 or 4 times out of 4 stimuli with an

SLC. At the highest stimulus intensity that we tested (43 dB), the majority of control larvae

(300/448) responded with a SLC within 7 ± 3 ms. 43 larvae showed no bias towards SLC or

LLC (2 times SLC and 2 times LLC, bias = 0). Those that did not perform an SLC (n = 105)

exhibited an LLC with a mean latency of 54 ± 29 ms (**Figure 2C**). In contrast, all NDL-PCB-

treated larvae (except for 10 nM PCB153) retained a response bias towards LLC at all

stimulus intensities tested, which was significantly different from the control (p < .001 for all

stimulus intensities). Lowering the PCB153 dose gradually increased the shift towards SLC,

yet 10 nM PCB153 was still significantly different from the control (p <.01 for all stimulus

intensities). Exposures starting after many structures of the brain and the neural tube have

developed, at 24 hpf (Smith and Kimelman 2019), had the same effect on startle latency

(**Figure S5B**).

The proportion of control larvae responding to the stimulus (either SLC or LLC) increased

from 75.2 ± 33.0% (n = 450) to 95.8 ± 17.0% (n = 457) from the lowest to highest stimulus

intensity, whereas the proportion of NDL-PCB153-, PCB138-, and PCB52-treated larvae that

responded remained below 70% (except for 10 nM PCB153) at all intensities and were

significantly different (p < .01) from the control (**Figure 2D**, **Figure S6A**). In contrast to those

exposed to NDL-PCBs, the larvae exposed to the DL-PCB126 and NDL-PCB118 had a

response latency and rate similar to the control larvae.

The different startle response bends exhibit different kinematics, with SLCs exhibiting larger

bend angles and faster angular velocities than LLCs (Burgess and Granato 2007). Thus,

LLC and SLC kinematics were analyzed separately. PCB52-treated larvae had increased

SLC bend angles (p > .0001) at two intermediate stimulus intensities (38 and 41 dB) while

PCB126 larvae had increased SLC bend angles at the three highest stimulus intensities

(**Figure S7**). Furthermore, PCB153 (1000 nM) exposure resulted in an increased SLC bend

angle at the highest stimulus intensity for those few larvae that exhibited an SLC (n = 26 of

138, 43 dB, p < .0456). LLC bend angle was significantly decreased across all stimulus

intensities for PCB118. It is not clear how bend angle is associated with altered signaling and

cellular function.

Cellular structures

The proper execution of the startle response relies on hair cells that sense the auditory

stimulus, M-cells to integrate the signals, and motor neurons and musculature to execute the

startle behavior. Morphological analysis showed that M-cells, motor neurons, and

musculature all appeared unaffected by PCB153, but the hair cell innervating neurons

appeared to be swollen (**Figure 3**). The L3 neuromast volume of control larvae was on

average 569 µm^3 (SD = 125), whereas PCB153-exposed larvae (1000 nM) had significantly

larger L3 neuromast volumes 848 µm^3 (SD = 185), t(48.93) = 6.765, p < .0001 (**Figure 3B**).

However, the number of hair cells in L3 neuromasts (**Figure 3C**) and the intensity of FM1-43

dye internalized by hair cells (**Figure 3D**) remained unchanged in PCB153-treated larvae in

comparison to the control.

The presence of M-cells is necessary for the SLC startle response. Staining with anti-3A10

confirmed the presence of M-cells in all PCB153-exposed larvae (21/21) (**Figure 4A**). We

tested the functional responsiveness of the M-cells by applying electric field pulses to head-

restrained larvae. This electrical stimulation bypasses the sensory system and directly

activates the M-cell (Tabor et al. 2014). There was no difference detected in response time

between control and 1000 nM PCB153-exposed larvae (U = 907.5, p = .489), demonstrating

that the signal transmission from the M-cell to the motor neurons is not disrupted in PCR153-

exposed fish (**Figure S8A**). The fraction responding also was similar in both the DMSO-

treated (290/299) and the PCB153-treated (209/219) groups (GLM, p = .427; **Figure 4C,**

Figure S8B). However, the C-bend evoked by the electrical stimulation was larger in

PCB153-exposed larvae (M = 101.4, SD = 38.60) than in the control larvae (M = 83.12, SD =

24.23). An unpaired t-test with Welch-s correction indicated that this difference was

statistically significant, $t(74.27) = 2.679$, $p = .0091$ (**Figure S8C**). This difference could

indicate a PCB153-evoked functional alteration of the M-cell axon or motor neuron.

For rapid and effective axonal signal transmission, proper myelination of axons is key. We

did not find PCB153 to reduce cells of the oligodendrocyte lineage in the spinal cord, $(t(36) =$

$.553$, $p < .584$) nor any obvious differences in myelination itself (**Figure S9**).

Proper function and activation of the motor neurons is crucial for completion of the startle

circuit, leading to muscle contractions and the characteristic swimming behaviors. The main

caudal primary (CaP), middle primary (MiP), and rostral primary (RoP) motor neurons did not

show any distinct structural differences between exposed and control larvae (**Figure S10**).

The fast muscle fibers in the myotome were not found to be differentially striated in either

treatment groups (**Figure S11**). No structural difference in the motor neurons or the

myotome was apparent in PCB153-exposed larvae, suggesting that the increased bend

angle does not originate from any overt structural differences in the motor neurons or muscle

fibers.

Neuronal activity

The release of neurotransmitters into the synaptic cleft is a characteristic process in signal

transmission. Both serotonergic and dopaminergic neuromodulators can affect startle latency

(Jain et al. 2018), and evidence has been found that NDL-PCBs alter these neuromodulators

in mammals (e.g. Castoldi et al. 2006; Enayah et al. 2018; Wigestrand et al. 2013). We

observed that PCB153 (1000 nM) altered the levels of several neurotransmitters in 6 dpf

zebrafish larvae (**Figure 5A**). Significantly increased level of GABA ($t(6) = 9.719$, $p < .001$),

choline-chloride ($t(6) = 8.459$, $p < .001$), 3-methoxytyramine hydrochloride ($t(6) = 3.803$, $p =$

$.009$), and L-valine ($t(6) = 3.134$, $p = .020$) were measured in PCB-exposed larvae in

comparison to control larvae; while L-Glutamine was decreased in comparison to control

larvae ($t(6) = 5.966$, $p < .001$).

To assess the interaction between PCB-exposure and changes in neuromodulatory activity,

we pharmacologically manipulated the neurotransmitter levels using dopamine, serotonin,

and GABA receptor agonists and antagonists. In control larvae, a 30 min exposure to

dopamine precursor L-DOPA and D2-receptor agonist Quinpirole, as well as serotonin

precursor 5-HTP, shifted the response to LLC, at multiple intensities tested ($p < .01$; **Figure**

5C). In contrast, the dopamine antagonist haloperidol exposure induced a shift to SLC in

controls across all stimulus intensities ($p < .01$; **Figure 5C**). Haloperidol induced the same

shift to SLC in PCB153-exposed larvae ($p < .001$), suggesting that the startle circuit is

functional but repressed by PCB (**Figure 5C**). The response rate remained unchanged after

exposure to neurotransmitter modulators except 5-HT, which caused an increase in

response probability across all stimulus intensities in control larvae ($p < .01$); Quinpirole,

which caused a decrease in response probability at the two lowest stimulus intensities in

control larvae ($p < .01$); and Haloperidol, which caused an increase in response probability at

41 dB in PCB153-treated larvae ($p < .01$; **Figure S6**). It is noteworthy, that co-exposure to L-

DOPA and PCB153 throughout development increased the response rate of PCB153-treated

larvae significantly ($p < .01$), yet only induced a significant shift towards SLC at 38 dB ($p <$

$.01$; **Figure S5C and S6C**).

In the fish brain, a distinct population of serotonergic neurons is found in the posterior

tuberculum/hypothalamus, forming a prominent horseshoe-like pattern that is one of several

brain regions where 5-HT is synthesized (Oikonomou et al. 2019). To assess whether the

development of serotonergic cells in the brain of PCB153 larvae was affected, we performed

immunohistochemistry using an antibody for 5-HT (**Figure S12**). The intensity of the

fluorescent signal in the hypothalamus was not different, comparing PCB153 ($M = 4.70$ x

10^6, $SD = 2.10$ x 10^6) with the control ($M = 4.38$ x 10^6, $SD = 8.1$ x 10^5) group. This suggests

that the serotonergic neurons in this region were not affected.

During specific behaviors or sensory experiences, only a small percentage of neurons are

active. We analyzed the active neuronal cell populations in PCB153-treated and -untreated

larvae, focusing on the rostral hindbrain, where the M-cells are located, to assess whether

there were changes in neural activity. We examined neuronal activity using the CaMPARI

zebrafish line, which contains a protein that upon UV illumination permanently photoconverts

from green to red fluorescence in cells with high calcium levels (Fosque et al. 2015). This

genomic tool enabled us to look for active neuronal cell populations in larvae during normal

swimming movement, and in larvae experiencing a vibroacoustic stimulated startle response,

both in embryo medium at room temperature (24 °C), and at a colder temperature (4 °C). A

higher red to green ratio (F_{red}/F_{green}) was measured comparing cold medium with room

temperature medium: swimming control larvae in cold medium had a higher F_{red}/F_{green} ratio

(M = 2.30, SD = 0.51) in comparison to swimming control larvae at RT (M = 0.96, SD =

0.17), confirming the potential of cold medium to elicit higher neuronal activity (**Figure S13**).

However, this difference was diminished in PCB153-treated, yet still significant in trial 1: $F(1,$

$18)$ = 121, *adj. p* < .001 and trial 2: $F(1, 32)$ = 65.2, *adj. p* = .041. Importantly, the difference

between swimming control and PCB153-treated larvae in cold medium was significant in

both trial 1: $F(1, 18)$ = 121, *adj. p* = .028 and trial 2 $F(1, 32)$ = 65.2, *adj. p* < .001 (**Figure 5**).

Similar differences in neuronal activity between control and PCB-153 treated larvae were

measured for startling larvae during photoconversion (**Figure S13**).

Discussion

Despite most use ending decades ago, PCBs remain an important and persistent pollutant

impacting both wildlife and humans. To date, mechanistic studies of neurodevelopmental

effects of NDL-PCBs suggest two major targets: (1) alterations in neurotransmission,

particularly involving dopamine and serotonin, and consequently disruptions to Ca^{2+} signal

transduction (Campagna et al. 2011; Dervola et al. 2015; Enayah et al. 2018; Fonnum and

Mariussen 2009; Langeveld et al. 2012), and (2) loss of outer hair cells (Crofton et al. 2000;

Trnovec et al. 2010) or cochlear development (Goldey et al. 1995). However, such studies

have not addressed how PCBs exert developmental neurobehavioral deficits, and thus we

do not know whether neurobehavioral disorders resulting from prenatal exposure originate in

a structural effect in the auditory system, a neurotransmitter imbalance, or both. In this study

we used the zebrafish startle response to comprehensively investigate the effects of different

PCBs on neuron development and signal transmission and to assess effects at

environmentally relevant concentrations for both humans and aquatic life. All NDL-PCBs

delayed the acoustic startle response in zebrafish larvae, which coincided with axonal

swelling and dysregulated GABA and dopamine metabolite levels. Co-exposures to the

dopamine D2-receptor antagonist and GABA receptor modulator drug haloperidol, alleviated

the delay in startle response.

The morphology of the neuronal cell body (perikarya) and neurite provides essential

information on the developmental and physiological abnormalities in the neural circuits. We

found that in PCB153-exposed larvae, the neurons that innervate the hair cells had an

increased volume (**Figure 3**), indicating that PCB153 alters neuronal structure and hair cell

function. Swelling of afferent terminals is associated with hearing deficits in both mammalian

models (Puel et al. 1991) and zebrafish (Sebe et al. 2017). The increased volume may

originate from increased numbers of bulged (swollen) afferent neurite termini or increased

arborization (branching). For example, *tmie* mutant zebrafish, which are deaf with defective

hair-cell function (Gleason et al. 2009), have more complex peripheral arbors in afferent

axons and an increased number of neurites. In wildtype zebrafish, neurites with swollen

axonal terminals are associated with hair cells of both polarities rather than just a single

polarity, suggesting a dysregulation of selection in synaptic targets (Faucherre et al. 2010).

At the sub-cellular level, neuronal swelling is due to increased cationic permeability in

neurons, causing an osmotic imbalance that leads to water entry and cytotoxic edema

(Rungta et al. 2015). While we examined a number of morphological features of the startle

circuit anatomy (hair cell, afferent neurons, M-cell, oligodendrocyte lineage cells,

myelination, motor neuron, muscle), only the volume of the neurons that innervate the hair

cells showed a measurable deviation in PCB153-exposed larvae. We thus infer that the

enlargement of axon termini is contributing to the neurotoxic potential of NDL-PCBs – an

understudied target of PCB toxicity. A few studies associate NDL-PCB exposure with altered

dendritic arborization (Lein et al. 2007; Yang et al. 2009). Axonal swelling is a pathology

appearing in many neurodegenerative disease (Coleman 2005), and exposure to NDL-PCBs

may be a contributing factor in such diseases.

The startle response in fish and amphibians is a fundamental mechanism involved in

predator escape. Acoustic stimuli can trigger two distinct motor behaviors: a short-latency C-

bend (SLC**; Figure 2A**) initiated within 5-15 ms of the stimulus, or a long-latency C-bend

(LLC; **Figure 2A**), initiated within 15-100 ms (Burgess and Granato 2007). At the structural

level, laser-ablation of the M-cell abolishes SLC responses without affecting LLC responses

(Lacoste et al. 2015), suggesting that the LLC is a 'backup' response (Burgess and Granato

2007). The finding that NDL-PCB-exposed larvae predominantly exhibit LLC without loss of

M-cells (**Figure 4A**), suggests that the M-cells are receiving a delayed input, are not firing

properly, or that the signal propagation downstream of the M-cell is disrupted. To distinguish

these, we directly activated the M-cells by applying electrical stimulation to control larvae

with their heads restrained, and found that unlike auditory stimulation, PCB153-exposed fish

had latencies indistinguishable from the controls. Furthermore, direct stimulation of PCB153-

exposed larvae resulted in a normal response rate and response time, although the bend

angle was significantly increased. These results indicate a proper function of M-cells in NDL-

PCB-treated larvae. However, the changes in bend angles we observed during these

responses could indicate alterations downstream of the M-cells.

There is less known about the mechanisms leading to an aberrant turning angle. Since we

found no obvious structural impairment downstream of the M-cell (oligodendrocyte lineage

cells, myelination, motor neuron, musculature), we speculate that the increased bend angle

in electrically stimulated larvae exposed to PCB153 may be linked to a sub-cellular

disruption such as gap junctions. It is known that PCBs can alter gap junction function in

cells of many organs (Bager et al. 1997; Kang et al. 1996; Machala et al. 2003) including

NDL-PCB specific inhibition of gap junctional intercellular communication in neural crest cells

(Nyffeler et al. 2018) and neuronal stem cells (PCB153; (Kang et al. 2001)). Gap junctions

are intercellular channels formed by connexins (Cx). RNA sequencing of zebrafish larvae

exposed to PCB153 in a similar exposure regime (4-120 hpf) and concentration (10 µM)

significantly altered the expression of connexin genes cx28.9, cx42, and cx35b (Aluru et al.

2020). Cx35 is expressed on the Mauthner neuron lateral dendrites in zebrafish (Jain et al.

2018) and plays a crucial role in regulating spinal motor activity during fast motor behavior of

adult mosquitofish (Serrano-Velez et al. 2014). We thus propose that gap junction function

could be disrupted, which may be a future study direction using single-cell RNA sequencing

or genetically encoded optogenetics that has yet to be developed in a model organism (Dong

et al. 2018).

Neurotransmitters also play a key role in the startle response. A previous screening of more

than 1000 pharmacologically active compounds identified serotonergic and dopaminergic

modulators among the largest class of drugs that shift bias between SLC and LLC startle

behavior (Jain et al. 2018). Specifically, acute exposure to the D3R agonist (7OHD) shifted

the startle bias from LLC to SLC, while exposure to a D3R antagonist (or a 5-HT1AR

agonist) shifted bias toward LLC behavior (Jain et al. 2018). Our results extend these

findings, showing that 30 minutes exposure to L-DOPA or 5-HTP induces a shift from SLC to

LLC at higher stimulus intensities, as opposed to the D2R antagonist haloperidol, which

evoked a shift to a stronger SLC bias at all stimulus intensities (**Figure 5C**). NDL-PCBs are

thought to affect dopaminergic pathways (Mariussen and Fonnum 2001; Tanaka et al. 2018;

Wigestrand et al. 2013), which is partly supported by our measurements of catecholamines

and traces amines. Specifically, we found that the extracellular dopamine metabolite 3-

methoxytyramine hydrochloride and the inhibitory transmitter GABA were significantly

increased (**Figure 5A**). The interplay between dopamine and GABA receptors activity is

complex. In the mammalian system, dopamine is a direct-acting modulator of $GABA_ARs$

(Hoerbelt et al. 2015). The simultaneous presence of a dopamine metabolite and GABA in

PCB153-exposed zebrafish larvae may indicate that increased levels of GABA are a

downstream effect of dysregulated dopamine activity. With both dopamine and GABA

playing critical roles in motor activity, we hypothesize that PCB153 impairs dopaminergic

neurotransmission and subsequently increases GABA levels, leading to the predominant

LLC response. One of the most crucial regulators of synapse development and function

mediating the GABAergic, dopaminergic, and serotonergic synaptic transmission is the brain

derived neurotrophic factor (BDNF). PCB153 induced the expression of *bdnf* in zebrafish

larvae (Aluru et al. 2020) suggesting BDNF as potential target for transcriptional regulation.

Furthermore, we propose that GABA is involved in the swelling of the afferent terminals

(**Figure 3**) as GABA activates Cl^- channels leading to an influx of Cl^- accompanied by H_2O

(Cesetti et al. 2012; Rungta et al. 2015). The osmotic neuronal swelling alters neuronal

activity. This study thus provides new evidence that GABA is involved in the neurobehavioral

mode of action of higher chlorinated NDL-PCBs, as has been for lower chlorinated

congeners (Fernandes et al. 2010).

An important finding of this study is that PCB153-exposed larvae can execute SLC when

treated with haloperidol, a D2-antagonist that can increase dopamine turnover in acute

exposure (Magnusson et al. 1987). There is a large body of evidence showing that NDL-

PCBs, including PCB153, modulate dopaminergic neurotransmission by inhibiting the

dopamine precursor enzyme tyrosine hydroxylase and dopamine synthesis itself, in both

neural crest cells and adult laboratory animal models including non-human primates (Dervola

et al. 2015; Enayah et al. 2018; Fonnum and Mariussen 2009; Seegal et al. 1990). In

zebrafish larvae, PCB153 exposure during development elicited differential expression of

nine genes related to the dopamine pathway, including tyrosine hydroxylase 2 (th2) (Aluru et

al. 2020). Yet linking such effects to behavior is rare. Tanaka et al. (2018) suggested that the

NDL-PCB dominant mixture Aroclor 1254 and BDE-47 functionally inhibited dopaminergic

neurons in early zebrafish embryos, as increased activity in 26 hpf embryos was inhibited by

supplementation with L-tyrosine and L-DOPA (Tanaka et al. 2018).

The restoration of the delayed startle phenotype by haloperidol, together with the increased

levels of a dopamine metabolite and GABA in PCB153-exposed larvae (**Figure 5A**),

indicates that disruption of dopaminergic and GABAergic signaling between neurons of the

auditory circuit is the dominant effect of PCB153-mediated aberrant motor behavior. These

signaling processes are widely conserved across vertebrate species, and are important in

human neurological diseases. For example, a recent study found that multiple gene

mutations causing defective startle response in zebrafish were also associated with

disorders of the locomotor system in humans (Meserve et al. 2020).

The localization of the PCB-induced disruption of dopaminergic signal transmission is likely

downstream of the hair cells, potentially in the hindbrain. Although dopaminergic neurons

innervate lateral line neuromasts, where D1R antagonist reduces the hair cell activity (Toro

et al. 2015), we found no difference in the hair cell activity of PCB153-exposed larvae

(**Figure 3D**). This is in contrast to the hindbrain, where neuronal activity was reduced in

PCB153 exposed CaMPARI zebrafish larvae (**Figure 5**), indicating a reduced capacity for

release or turnover of calcium. Intracellular Ca^{2+} signaling in neurons controls

neurotransmitter secretion among many other cellular processes and thus may be indirectly

involved in dopamine or GABA regulation. Some NDL-PCBs (PCB 95, 138) are known to

inhibit voltage-gated calcium channels (Langeveld et al. 2012) contributing to interference

with calcium homeostasis and consequently neurotransmission and synaptic plasticity

(Brunelli et al. 2012; Campagna et al. 2011; Gafni et al. 2003; Ta et al. 2006).

The startle response to threatening stimuli is vital for survival. PCB153-exposed zebrafish

larvae predominantly responded with a LLC, which not only delays the response, increasing

the chance to be caught by a predator, but also usually does not displace the animal from its

original location (**Figure 2A** and Jain et al., 2018). While survival of predator attack is likely

context-dependent and species-specific, escaping fast and away from a predator increases

survival in the field (McCormick et al. 2018). This suggests that exposure to NDL-PCBs may

alter selection pressure and thereby shift population structure. We have observed a

significantly delayed startle response at concentrations as low as 10 nM (3.6 µg L^{-1}). PCBs

are associated with sediments, where concentrations of PCB153 frequently reach multiple

µg kg^{-1} (Lai et al. 2015). PCB concentrations in organisms may be even higher due to

extensive bioconcentration (Killifish from New Bedford Harbor have 6.5 to 7 µg of PCB153

per g wet weight; Gräns et al. 2015). All NDL-PCBs tested in this study increased the startle

latency. Thus, in the environment, where PCBs occur in mixtures of numerous congeners

that likely have additive effects, the effect concentration is suggested to be lower than that of

single congeners. Taken together, these results highlight the importance of the neurotoxic

potential of NDL-PCBs in an ecological context.

In conclusion, the comprehensive approach presented here suggests that NDL-PCBs alter

dopamine signaling, leading to an increase in GABA that ultimately results in swelling of

axonal terminals in the auditory circuit. While the involvement of connexins remains

speculative, our findings provide a foothold into the dopamine dysregulating potential of

NDL-PCBs in sensory-motor processing in developing vertebrates, and link that to motor

behavior. In a direct comparison, the present study demonstrates that the adverse

behavioral effects are induced by all NDL-PCBs tested, but not by the classical DL-PCB126.

Whether this is true for compounds that are structurally similar to NDL-PCBs, such as some

PBDEs, is not known and will require further research. A key finding is that the effect of NDL-

PCBs on the startle latency occurs at concentrations relevant for both environmental and

human exposures, and therefore may contribute to the selection pressure in the environment

and neurotoxicological disease in human. This study provides valuable clues for tackling

health effects in humans and wildlife caused by environmental exposure.

CRediT authorship contribution statement

Nadja R. Brun: Conceptualization, Methodology, Formal analysis, Investigation, Writing - original draft, Writing - review & editing, Visualization, Funding acquisition. **Jennifer M. Panlilio:** Conceptualization, Methodology, Software, Formal analysis, Investigation, Writing - review & editing, Funding acquisition. **Kun Zhang:** Methodology, Investigation, Writing - review & editing, Funding acquisition. **Yanbin Zhao:** Methodology, Investigation, Writing - review & editing, Funding acquisition. **Evgeny Ivashkin:** Methodology, Investigation. **John J. Stegeman:** Conceptualization, Writing - review & editing, Funding acquisition. **Jared V. Goldstone:** Conceptualization, Writing - review & editing, Project administration.

Funding

This work was supported by the Swiss National Science Foundation P2EZP2_165200 (NRB), the Boston University Superfund Research Program NIH 5P42ES007381 (JJS and JVG), the Woods Hole Center for Oceans and Human Health (NIH: P01ES021923 and P01ES028938; NSF: OCE-1314642 and OCE-1840381) (NRB, JMP and JJS), and the National Natural Science Foundation of China 22006099 (KZ and YZ) and the Shanghai Pujiang Program 19PJ1404900 (KZ and YZ).

Acknowledgments

The authors would like to thank Neel Aluru for helpful discussion and providing the confocal microscope and Ed Levin and Mark Hahn for critically reading the manuscript and making valuable suggestions.

References

Almeida RG, Czopka T, Ffrench-Constant C, Lyons DA. 2011. Individual axons regulate the myelinating potential of single oligodendrocytes in vivo. Development 138:4443–50; doi:10.1242/dev.071001.

Aluru N, Krick KS, Mcdonald AM, Karchner SI. 2020. Developmental exposure to PCB153 (2,2′,4,4′,5,5′-hexachlorobiphenyl) alters circadian rhythms and the expression of clock

and metabolic genes. Toxicol Sci 173:41–52; doi:10.1093/toxsci/kfz217.

Bager Y, Lindebro MC, Martel P, Chaumontet C, Wärngård L. 1997. Altered function,
localization and phosphorylation of gap junctions in rat liver epithelial, IAR 20, cells after
treatment with PCBs or TCDD. Environ Toxicol Pharmacol 3:257–266;
doi:10.1016/S1382-6689(97)00021-5.

Bates D, Mächler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models using
lme4. J Stat Softw 67:1–48; doi:10.18637/jss.v067.i01.

Berger DF, Lombardo JP, Jeffers PM, Hunt AE, Bush B, Casey A, et al. 2001. Hyperactivity
and impulsiveness in rats fed diets supplemented with either Aroclor 1248 or PCB-
contaminated St. Lawrence river fish. Behav Brain Res 126:1–11; doi:10.1016/S0166-
4328(01)00244-3.

Bergonzi R, Specchia C, Dinolfo M, Tomasi C, De Palma G, Frusca T, et al. 2009.
Distribution of persistent organochlorine pollutants in maternal and foetal tissues: Data
from an Italian polluted urban area. Chemosphere 76:747–754;
doi:10.1016/j.chemosphere.2009.05.026.

Boix J, Cauli O, Felipo V. 2010. Developmental exposure to polychlorinated biphenyls 52,
138 or 180 affects differentially learning or motor coordination in adult rats. Mechanisms
involved. Neuroscience 167:994–1003; doi:10.1016/j.neuroscience.2010.02.068.

Boucher O, Muckle G, Bastien CH. 2009. Prenatal exposure to polychlorinated biphenyls: A
neuropsychologic analysis. Environ Health Perspect 117:7–16; doi:10.1289/ehp.11294.

Brunelli L, LLansola M, Felipo V, Campagna R, Airoldi L, De Paola M, et al. 2012. Insight
into the neuroproteomics effects of the food-contaminant non-dioxin like polychlorinated
biphenyls. J Proteomics 75:2417–2430; doi:10.1016/J.JPROT.2012.02.023.

Burgess HA, Granato M. 2007. Sensorimotor gating in larval zebrafish. J Neurosci 27:4984–
4994; doi:10.1523/JNEUROSCI.0615-07.2007.

Campagna R, Brunelli L, Airoldi L, Fanelli R, Hakansson H, Heimeier RA, et al. 2011.
Cerebellum Proteomics addressing the cognitive deficit of rats perinatally exposed to
the food-relevant polychlorinated biphenyl 138. Toxicol Sci 123:170–179;
doi:10.1093/toxsci/kfr156.

Castoldi AF, Blandini F, Randine G, Samuele A, Manzo L, Coccini T. 2006. Brain
monoaminergic neurotransmission parameters in weanling rats after perinatal exposure
to methylmercury and 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153). Brain Res 1112:91–
98; doi:10.1016/j.brainres.2006.07.022.

Cesetti T, Ciccolini F, Li Y. 2012. GABA not only a neurotransmitter: Osmotic regulation by
GABA AR signaling. Front Cell Neurosci 6; doi:10.3389/fncel.2012.00003.

Chen Y-CJ, Guo Y-L, Hsu C-C, Rogan WJ. 1992. Cognitive development of Yu-Cheng ('Oil
Disease') children prenatally exposed to heat-degraded PCBs. JAMA J Am Med Assoc
268:3213; doi:10.1001/jama.1992.03490220057028.

Coccini T, Roda E, Castoldi AF, Poli D, Goldoni M, Vettori MV, et al. 2011. Developmental
exposure to methylmercury and 2,2′,4,4′,5,5′-hexachlorobiphenyl (PCB153) affects

cerebral dopamine D1-like and D2-like receptors of weanling and pubertal rats. Arch
Toxicol 85:1281–1294; doi:10.1007/s00204-011-0660-y.

Coleman M. 2005. Axon degeneration mechanisms: Commonality amid diversity. Nat Rev
Neurosci 6:889–898; doi:10.1038/nrn1788.

Crofton KM, Ding D-L, Padich R, Taylor M, Henderson D. 2000. Hearing loss following
exposure during development to polychlorinated biphenyls: A cochlear site of action.
Hear Res 144:196–204; doi:10.1016/S0378-5955(00)00062-9.

Dervola KSN, Johansen EB, Walaas SI, Fonnum F. 2015. Gender-dependent and genotype-
sensitive monoaminergic changes induced by polychlorinated biphenyl 153 in the rat
brain. Neurotoxicology 50:38–45; doi:10.1016/J.NEURO.2015.07.004.

Dong A, Liu S, Li Y. 2018. Gap junctions in the nervous system: Probing functional
connections using new imaging approaches. Front Cell Neurosci 12:1–9;
doi:10.3389/fncel.2018.00320.

Enayah SH, Vanle BC, Fuortes LJ, Doorn JA, Ludewig G. 2018. PCB95 and PCB153
change dopamine levels and turn-over in PC12 cells. Toxicology 394:93–101;
doi:10.1016/j.tox.2017.12.003.

Ennaceur S, Gandoura N, Driss MR. 2008. Distribution of polychlorinated biphenyls and
organochlorine pesticides in human breast milk from various locations in Tunisia: Levels
of contamination, influencing factors, and infant risk assessment. Environ Res 108:86–
93; doi:10.1016/J.ENVRES.2008.05.005.

Faucherre A, Baudoin JP, Pujol-Martí J, López-Schier H. 2010. Multispectral four-
dimensional imaging reveals that evoked activity modulates peripheral arborization and
the selection of plane-polarized targets by sensory neurons. Development 137:1635–
1643; doi:10.1242/dev.047316.

Fernandes ECA, Hendriks HS, Van Kleef RGDM, Van Den Berg M, Westerink RHS. 2010.
Potentiation of the human GABAA receptor as a novel mode of action of lower-
chlorinated non-dioxin-like PCBs. Environ Sci Technol 44:2864–2869;
doi:10.1021/es902321a.

Fonnum F, Mariussen E. 2009. Mechanisms involved in the neurotoxic effects of
environmental toxicants such as polychlorinated biphenyls and brominated flame
retardants. J Neurochem 111:1327–1347; doi:10.1111/j.1471-4159.2009.06427.x.

Fosque BF, Sun Y, Dana H, Yang C, Ohyama T, Tadross MR, et al. 2015. Labeling of active
neural circuits in vivo with designed calcium integrators. Science 347: 755–760.

Gafni J, Wong PW, Pessah IN. 2003. Non-coplanar 2,2',3,5',6-pentachlorobiphenyl (PCB 95)
amplifies ionotropic glutamate receptor signaling in embryonic cerebellar granule
neurons by a mechanism involving ryanodine receptors. Toxicol Sci 77:72–82;
doi:10.1093/toxsci/kfh004.

Gleason MR, Nagiel A, Jamet S, Vologodskaia M, López-Schier H, Hudspeth AJ. 2009. The
transmembrane inner ear (Tmie) protein is essential for normal hearing and balance in
the zebrafish. Proc Natl Acad Sci U S A 106:21347–21352;
doi:10.1073/pnas.0911632106.

Goldey ES, Kehn LS, Lau C, Rehnberg GL, Crofton KM. 1995. Developmental exposure to
polychlorinated biphenyls (Aroclor 1254) reduces circulating thyroid hormone
concentrations and causes hearing deficits in rats. Toxicol Appl Pharmacol 135:77–88;
doi:10.1006/TAAP.1995.1210.

Grandjean P, Landrigan PJ. 2014. Neurobehavioural effects of developmental toxicity.
Lancet Neurol 13:330–338; doi:10.1016/S1474-4422(13)70278-3.

Gräns J, Wassmur B, Fernández-Santoscoy M, Zanette J, Woodin BR, Karchner SI, et al.
2015. Regulation of pregnane-X-receptor, CYP3A and P-glycoprotein genes in the
PCB-resistant killifish (Fundulus heteroclitus) population from New Bedford Harbor.
Aquat Toxicol 159:198–207; doi:10.1016/j.aquatox.2014.12.010.

Herbstman JB, Sjödin A, Apelberg BJ, Witter FR, Patterson DG, Halden RU, et al. 2007.
Determinants of prenatal exposure to polychlorinated biphenyls (PCBs) and
polybrominated diphenyl ethers (PBDEs) in an urban population. Environ Health
Perspect 115:1794–1800; doi:10.1289/ehp.10333.

Hoerbelt P, Lindsley TA, Fleck MW. 2015. Dopamine directly modulates GABAA receptors. J
Neurosci 35:3525–3536; doi:10.1523/JNEUROSCI.4390-14.2015.

Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. 2013. The
zebrafish reference genome sequence and its relationship to the human genome.
Nature 496:498–503; doi:10.1038/nature12111.

Hudspeth A. 1985. The cellular basis of hearing: the biophysics of hair cells. Science
230:745–752; doi:10.1126/science.7414321.

Inoue D, Wittbrodt J. 2011. One for all-a highly efficient and versatile method for fluorescent
immunostaining in fish embryos. PLoS One 6:1–7; doi:10.1371/journal.pone.0019713.

Jain RA, Wolman MA, Marsden KC, Nelson JC, Shoenhard H, Echeverry FA, et al. 2018. A
forward genetic screen in zebrafish identifies the G-protein-coupled receptor CaSR as a
modulator of sensorimotor decision making. Curr Biol 28:1357-1369.e5;
doi:10.1016/j.cub.2018.03.025.

Johansen EB, Fonnum F, Lausund PL, Walaas SI, Bærland NE, Wøien G, et al. 2014.
Behavioral changes following PCB 153 exposure in the Spontaneously Hypertensive rat
- an animal model of Attention-Deficit/Hyperactivity disorder. Behav Brain Funct 10:1–
19; doi:10.1186/1744-9081-10-1.

Jusko TA, Sisto R, Iosif A-M, Moleti A, Wimmerová S, Lancz K, et al. 2014. Prenatal and
postnatal serum PCB concentrations and cochlear function in children at 45 months of
age. Environ Health Perspect 122:1246–1252; doi:10.1289/ehp.1307473.

Kang K-S, Park J-E, Ryu D-Y, Lee Y-S. 2001. Effects and neuro-toxic mechanisms of
2,2',4,4',5,5'-hexachlorobiphenyl and endosulfan in neuronal stem cells. J Vet Med Sci
63:1183–1190; doi:10.1292/jvms.63.1183.

Kang KS, Wilson MR, Hayashi T, Chang CC, Trosko JE. 1996. Inhibition of gap junctional
intercellular communication in normal human breast epithelial cells after treatment with
pesticides, PCBs, and PBBs, alone or in mixtures. Environ Health Perspect 104:192–
200; doi:10.1289/ehp.96104192.

Kimmel CB. 1993. Patterning the brain of the zebrafish embryo. Annu Rev Neurosci 16:707–
732; doi:10.1146/annurev.ne.16.030193.003423.

Lacoste AMB, Schoppik D, Robson DN, Haesemeyer M, Portugues R, Li JM, et al. 2015. A
convergent and essential interneuron pathway for mauthner-cell-mediated escapes.
Curr Biol 25:1526–1534; doi:10.1016/j.cub.2015.04.025.

Lai Z, Li X, Li H, Zhao L, Zeng Y, Wang C, et al. 2015. Residual distribution and risk
assessment of polychlorinated biphenyls in surface sediments of the Pearl River Delta,
South China. Bull Environ Contam Toxicol 95:37–44; doi:10.1007/s00128-015-1563-z.

Lancz K, Hertz-Picciotto I, Jusko TA, Murínová Ľ, Wimmerová S, Šovčíková E, et al. 2015a.
Duration of breastfeeding and serum PCB 153 concentrations in children. Environ Res
136:35–39; doi:10.1016/j.envres.2014.09.036.

Lancz K, Palkovičováa L, Patayová H, Drobná B, Wimmerová S, Šovčíková E, et al. 2015b.
Ratio of cord to maternal serum PCB concentrations in relation to their congener-
specific physicochemical properties. Int J Hyg Env Heal 218:91–98;
doi:doi:10.1016/j.ijheh.2014.08.003.

Landrigan PJ, Fuller R, Acosta NJR, Adeyi O, Arnold R, Basu N, et al. 2018. The Lancet
Commission on pollution and health. Lancet 391:462–512; doi:10.1016/S0140-
6736(17)32345-0.

Langeveld WT, Meijer M, Westerink RHS. 2012. Differential effects of 20 non-dioxin-like
PCBs on basal and depolarization-evoked intracellular calcium levels in PC12 cells.
Toxicol Sci 126:487–496; doi:10.1093/toxsci/kfr346.

Lein PJ, Yang D, Bachstetter AD, Tilson HA, Harry GJ, Mervis RF, et al. 2007. Ontogenetic
alterations in molecular and structural correlates of dendritic growth after developmental
exposure to polychlorinated biphenyls. Environ Health Perspect 115:556–563;
doi:10.1289/ehp.9773.

Lilienthal H, Heikkinen P, Andersson PL, van der Ven LTM, Viluksela M. 2011. Auditory
effects of developmental exposure to purity-controlled polychlorinated biphenyls
(PCB52 and PCB180) in rats. Toxicol Sci 122:100–111; doi:10.1093/toxsci/kfr077.

Lilienthal H, Korkalainen M, Andersson PL, Viluksela M. 2015. Developmental exposure to
purity-controlled polychlorinated biphenyl congeners (PCB74 and PCB95) in rats:
Effects on brainstem auditory evoked potentials and catalepsy. Toxicology 327:22–31;
doi:10.1016/J.TOX.2014.11.004.

Liu YC, Hale ME. 2017. Local spinal cord circuits and bilateral mauthner cell activity function
together to drive alternative startle behaviors. Curr Biol 27:697–704;
doi:10.1016/j.cub.2017.01.019.

Machala M, Bláha L, Vondráček J, Trosko JE, Scott J, Upham BL. 2003. Inhibition of gap
junctional intercellular communication by noncoplanar polychlorinated biphenyls:
Inhibitory potencies and screening for potential Mode(s) of Action. Toxicol Sci 76:187–
195; doi:10.1093/TOXSCI.

Magnusson O, Mohringe B, Thorell G, Lake-Bakaar DM. 1987. Effects of the dopamine D2
selective receptor antagonist remoxipride on dopamine turnover in the rat brain after

acute and repeated administration. Pharmacol Toxicol 60:368–373; doi:10.1111/j.1600-
0773.1987.tb01529.x.

Mariussen E, Fonnum F. 2001. The effect of polychlorinated biphenyls on the high affinity
uptake of the neurotransmitters, dopamine, serotonin, glutamate and GABA, into rat
brain synaptosomes. Toxicology 159:11–21; doi:10.1016/S0300-483X(00)00374-7.

Maximino C, Herculano AM. 2010. A review of monoaminergic neuropsychopharmacology in
zebrafish. Zebrafish 7:359–378; doi:10.1089/zeb.2010.0669.

McCormick MI, Fakan E, Allan BJM. 2018. Behavioural measures determine survivorship
within the hierarchy of whole-organism phenotypic traits. Funct Ecol 32:958–969;
doi:10.1111/1365-2435.13033.

Medan V, Preuss T. 2014. The Mauthner-cell circuit of fish as a model system for startle
plasticity. J Physiol Paris 108:129–140; doi:10.1016/j.jphysparis.2014.07.006.

Meserve JH, Nelson JC, Marsden KC, Hsu J, Echeverry FA, Jain A, et al. 2020. A forward
genetic screen identifies Dolk as a regulator of startle magnitude through the potassium
channel subunit Kv1.1. bioRxiv.

Meyers JR, MacDonald RB, Duggan A, Lenzi D, Standaert DG, Corwin JT, et al. 2003.
Lighting up the senses: FM1-43 loading of sensory cells through nonselective ion
channels. J Neurosci 23:4054–65; doi:10.1523/JNEUROSCI.23-10-04054.2003.

Nakayama H. 2004. Common sensory inputs and differential excitability of segmentally
homologous reticulospinal neurons in the hindbrain. J Neurosci 24:3199–3209;
doi:10.1523/JNEUROSCI.4419-03.2004.

Nyffeler J, Chovancova P, Dolde X, Holzer AK, Purvanov V, Kindinger I, et al. 2018. A
structure–activity relationship linking non-planar PCBs to functional deficits of neural
crest cells: new roles for connexins. Arch Toxicol 92:1225–1247; doi:10.1007/s00204-
017-2125-4.

Oikonomou G, Altermatt M, Zhang R wei, Coughlin GM, Montz C, Gradinaru V, et al. 2019.
The serotonergic raphe promote sleep in zebrafish and mice. Neuron 103:686-701.e8;
doi:10.1016/j.neuron.2019.05.038.

Panlilio JM, Aluru N, Hahn ME. 2020. Developmental neurotoxicity of the harmful algal
bloom toxin domoic acid: Cellular and molecular mechanisms underlying altered
behavior in the zebrafish Model. Environ Health Perspect 128:117002;
doi:10.1289/EHP6652.

Panlilio JM, Jones IT, Salanga MC, Aluru N, Hahn ME. 2021. Developmental exposure to
domoic acid disrupts startle response behavior and circuitry. bioRxiv;
doi:10.1101/2021.01.08.425996.

Park H-C, Shin J, Roberts RK, Appel B. 2007. An olig2 reporter gene marks oligodendrocyte
precursors in the postembryonic spinal cord of zebrafish. Dev Dyn 236:3402–3407;
doi:10.1002/dvdy.21365.

Poon E, Bandara SB, Allen JB, Sadowski RN, Schantz SL. 2015. Developmental PCB
exposure increases susceptibility to audiogenic seizures in adulthood. Neurotoxicology

46:117–124; doi:10.1016/j.neuro.2014.12.007.

Puel JL, Pujol R, Ladrech S, Eybalin M. 1991. α-Amino-3-hydroxy-5-methyl-4-isoxazole
propionic acid electrophysiological and neurotoxic effects in the guinea-pig cochlea.
Neuroscience 45:63–72; doi:10.1016/0306-4522(91)90103-U.

Pujol-Martí J, López-Schier H. 2013. Developmental and architectural principles of the
lateral-line neural map. Front Neural Circuits 7:47; doi:10.3389/fncir.2013.00047.

Ranasinghe P, Thorn RJ, Seto R, Creton R, Bridges WC, Chapman SC, et al. 2019.
Embryonic exposure to 2,2′,3,5′,6-pentachlorobiphenyl (PCB-95) causes developmental
malformations in zebrafish. Environ Toxicol Chem 39:162–170; doi:10.1002/etc.4587.

Ribas-Fitó N, Sala M, Kogevinas M, Sunyer J. 2001. Polychlorinated biphenyls (PCBs) and
neurological development in children: A systematic review. J Epidemiol Community
Health 55:537–546; doi:10.1136/jech.55.8.537.

Rice D, Barone S. 2000. Critical periods of vulnerability for the developing nervous system:
Evidence from humans and animal models. Environ Health Perspect 108:511–533;
doi:10.2307/3454543.

Roberts A. 2000. Early functional organization of spinal neurons in developing lower
vertebrates. Brain Res Bull 53:585–593; doi:10.1016/S0361-9230(00)00392-0.

Rungta RL, Choi HB, Tyson JR, Malik A, Dissing-Olesen L, Lin PJC, et al. 2015. The cellular
mechanisms of neuronal swelling underlying cytotoxic edema. Cell 161:610–621;
doi:10.1016/j.cell.2015.03.029.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. 2012. Fiji:
An open-source platform for biological-image analysis. Nat Methods 9:676–682;
doi:10.1038/nmeth.2019.

Sebe JY, Cho S, Sheets L, Rutherford MA, von Gersdorff H, Raible DW. 2017. Ca^{2+}-
permeable AMPARs mediate glutamatergic transmission and excitotoxic damage at the
hair cell ribbon synapse. J Neurosci 37:6162–6175; doi:10.1523/jneurosci.3644-
16.2017.

Seegal RF, Bush B, Shain W. 1990. Lightly chlorinated ortho-substituted PCB congeners
decrease dopamine in nonhuman primate brain and in tissue culture. Toxicol Appl
Pharmacol 106: 136–44.

Serrano-Velez JL, Rodriguez-Alvarado M, Torres-Vazquez II, Fraser SE, Yasumura T,
Vanderpool KG, et al. 2014. Abundance of gap junctions at glutamatergic mixed
synapses in adult Mosquitofish spinal cord neurons. Front Neural Circuits 8:66;
doi:10.3389/fncir.2014.00066.

Shin J, Park H-C, Topczewska JM, Mawdsley DJ, Appel B. 2003. Neural cell fate analysis in
zebrafish using <i>olig2<i> BAC transgenics. Methods Cell Sci 25:7–14;
doi:10.1023/B:MICS.0000006847.09037.3a.

Smith NL, Kimelman D. 2019. *Establishing the body plan: The first 24 hours of zebrafish
development*. Elsevier.

Stenberg M, Hamers T, Machala M, Fonnum F, Stenius U, Lauy AA, et al. 2011. Multivariate
toxicity profiles and QSAR modeling of non-dioxin-like PCBs - An investigation of in vitro
screening data from ultra-pure congeners. Chemosphere 85:1423–1429;
doi:10.1016/j.chemosphere.2011.08.019.

Stewart AM, Braubach O, Spitsbergen J, Gerlai R, Kalueff A V. 2014. Zebrafish models for
translational neuroscience research: From tank to bedside. Trends Neurosci 37:264–
278; doi:10.1016/j.tins.2014.02.011.

Ta TA, Feng W, Molinski TF, Pessah IN. 2006. Hydroxylated xestospongins block inositol-
1,4,5-trisphosphate-induced Ca^{2+} release and sensitize Ca^{2+}-induced Ca^{2+} release
mediated by ryanodine receptors. Mol Pharmacol 69:532–538;
doi:10.1124/mol.105.019125.used.

Tabor KM, Bergeron SA, Horstick EJ, Jordan DC, Aho V, Porkka-Heiskanen T, et al. 2014.
Direct activation of the Mauthner cell by electric field pulses drives ultrarapid escape
responses. J Neurophysiol 112:834–844; doi:10.1152/jn.00228.2014.

Takada N, Kucenas S, Appel B. 2010. Sox10 is necessary for oligodendrocyte survival
following axon wrapping. Glia 58:996–1006; doi:10.1002/glia.20981.

Tanaka Y, Fujiwara M, Shindo A, Yin G, Kitazawa T, Teraoka H. 2018. Aroclor 1254 and
BDE-47 inhibit dopaminergic function manifesting as changes in locomotion behaviors
in zebrafish embryos. Chemosphere 193:1207–1215;
doi:10.1016/j.chemosphere.2017.11.138.

Toro C, Trapani JG, Pacentine I, Maeda R, Sheets L, Mo W, et al. 2015. Dopamine
modulates the activity of sensory hair cells. J Neurosci 35:16494–16503;
doi:10.1523/JNEUROSCI.1691-15.2015.

Trnovec T, Šovčíková E, Pavlovčinová G, Jakubíková J, Jusko TA, Husťák M, et al. 2010.
Serum PCB concentrations and cochlear function in 12-year-old children. Environ Sci
Technol 44:2884–2889; doi:10.1021/es901918h.

Troconis EL, Ordoobadi AJ, Sommers TF, Aziz-Bose R, Carter AR, Trapani JG. 2017.
Intensity-dependent timing and precision of startle response latency in larval zebrafish.
J Physiol 595:265–282; doi:10.1113/JP272466.

Tropepe V, Sive HL. 2003. Can zebrafish be used as a model to study the
neurodevelopmental causes of autism? Genes, Brain Behav 2:268–281;
doi:10.1034/j.1601-183X.2003.00038.x.

Vettori MV, Goldoni M, Caglieri A, Poli D, Folesani G, Ceccatelli S, et al. 2006. Antagonistic
effects of methyl-mercury and PCB153 on PC12 cells after a combined and
simultaneous exposure. Food Chem Toxicol 44:1505–1512;
doi:10.1016/j.fct.2006.04.009.

Voesenek CJ, Muijres FT, Van Leeuwen JL. 2018. Biomechanics of swimming in developing
larval fish. J Exp Biol 221; doi:10.1242/jeb.149583.

Walkowiak J, Wiener J-A, Fastabend A, Heinzow B, Krämer U, Schmidt E, et al. 2001.
Environmental exposure to polychlorinated biphenyls and quality of the home
environment: effects on psychodevelopment in early childhood. Lancet 358:1602–1607;

doi:10.1016/S0140-6736(01)06654-5.

Wigestrand MB, Stenberg M, Walaas SI, Fonnum F, Andersson PL. 2013. Non-dioxin-like
PCBs inhibit [3H]WIN-35,428 binding to the dopamine transporter: A structure–activity
relationship study. Neurotoxicology 39:18–24; doi:10.1016/J.NEURO.2013.07.005.

Wolman MAA, Jain RAA, Marsden KCC, Bell H, Skinner J, Hayer KEE, et al. 2015. A
genome-wide screen identifies PAPP-AA-mediated IGFR signaling as a novel regulator
of habituation learning. Neuron 85:1200–1211; doi:10.1016/j.neuron.2015.02.025.

Yang D, Kim KH, Phimister A, Bachstetter AD, Ward TR, Stackman RW, et al. 2009.
Developmental exposure to polychlorinated biphenyls interferes with experience-
dependent dendritic plasticity and ryanodine receptor expression in weanling rats.
Environ Health Perspect 117:426–435; doi:10.1289/ehp.11771.

Zhao Y, Liang J, Meng H, Yin Y, Zhen H, Zheng X, et al. 2020. Rare earth elements
lanthanum and praseodymium adversely affect neural and cardiovascular development
in zebrafish (Danio rerio). Environ Sci Technol; doi:10.1021/acs.est.0c06632.

1001

1002

 Figures

Statoacoustic ganglion
Mauthner-cells
Hair cells
Posterior lateral line
Spinal cord
Motor neurons

1004   **Figure 1. Startle response circuit in zebrafish.** The proper execution of the startle response
1005   relies on hair cells to sense the auditory stimulus, Mauthner-cells to integrate the signals, and
1006   motor neurons and musculature to execute the startle behavior.

Figure 2. Effect of PCBs on startle response in 6 dpf zebrafish. (A) Representative short-latency C-bend (SLC) and long-latency C-bend (LLC) startle response of a control zebrafish larva at 6 dpf in comparison to a typical response of a PCB153 treated larva. (B) Vibro-acoustic startle latency at highest stimulus intensity (43 dB) in vehicle control (n = 479), NDL-PCB153 1000 nM (n = 179), NDL-PCB153 100 nM (n = 253), NDL-PCB153 30 nM (n = 152), NDL-PCB153 10 nM (n = 119), NDL-PCB138 (n = 139), and DL-PCB126 (n = 97) exposed larvae. (C) Bias of SLC and LLC and (D) response rate of different exposure groups at different vibro-acoustic intensities. Values are presented as mean ± SEM. All data points are biologically independent replicates from at least three independent experiments. Significant differences to DMSO controls (p < 0.01) are indicated by open symbols.

Figure 3. Effect of PCB153 on morphology and function of lateral line hair cells and innervating neurons. (**A**) Diagram indicating L3 neuromast. (**B**) Hair cell innervating neurons at neuromast L3 shown using *Tg(cntn1b:EGFP-CAAX)*. Volume of hair cell innervating neurons of DMSO (*n* = 30) and PCB153 (*n* = 29) treated larvae. Scale bar = 20 μm. (**C**) Whole-mount immune staining of kinocilia and hair cell innervating neurons (anti-acetylated tubulin antibody; green) and counterstain of stereocilia bundles and muscle tissue (actin; magenta) of DMSO (*n* = 26) and PCB153 (*n* = 24) treated larvae. The hair cell number was determined based on the number of kinocilia. Scale bar = 10 μm (**D**) Uptake of FM-143 dye as indication of functional hair cells of DMSO (*n* = 32) and PCB153 (*n* = 28) treated larvae. Scale bar = 10 μm. All data points are biologically independent replicates from at three independent experiments. Asterisks indicate significant differences to controls (***p < 0.0001).

A

B

C

Figure 4. Effect of PCB153 on morphology and function of the Mauthner-cells. (**A**)
Diagram indicating localization of Mauthner-cells in the hindbrain. Brain tissue immunostaining
with anti-neurofilament 3A10 antibody labeling reticulospinal neurons in control and PCB153
treated larvae at 6 dpf (ventral view). Mauthner (arrow) cells were present in all brain tissues
(control: n = 23, PCB153: n = 21). Scale bar = 50 µm. (**B**) Diagram of electrical stimulation
performed to assess the functionality of Mauthner-cells. (**C**) Percentage of responders to
electrical stimulation. All data points are biologically independent replicates from at least three
independent experiments.

Figure 5. Involvement of neurotransmitters in startle latency. (**A**) Neurotransmitter levels in 6 dpf larvae exposed to 1000 nM PCB153 in comparison to control (*n* = 4). (**B**) Representative maximum intensity z-projection from confocal stack after photoconversions of freely swimming 6 dpf CaMPARI larvae. Scale bar = 100 μm. Ratio of red to green fluorescence intensity in the rostral hindbrain (indicated by dashed line in image) in DMSO (*n* = 14) and 1000 nM PCB153 (*n* = 14) exposed startling larvae in 4 °C (cold) medium. Dashed lines indicate median per trial. Two independent experiments. Asterisks indicate significant differences to controls (*p < 0.05, **p < 0.01, and ***p < 0.001). (**C**) Startle bias shift induced by 30-minute exposure to serotonin, dopamine, and GABA neurotransmitter modulators in 6 dpf control and 100 nM PCB153 exposed larvae. Control (*n* = 138-172), 5-HTP (*n* = 67-80), L-DOPA (*n* = 76-94), Haloperidol (Control *n* = 123-162, PCB153 *n* = 37-61), and Bicuculline (Control *n* = 90-162, PCB153 *n* = 83-137). All data points are biologically independent replicates from at least three independent experiments. Significant differences to respective controls (*p* < 0.01) are indicated by open symbols.