Comparison of library preparation and sequencing depths for direct sequencing of

***Bordetella pertussis* positive samples**

Winkie Fong[1,3,4], Keenan Pey[1,3,4], Rebecca Rockett[1,3], Rosemarie Sadsad[1,2,3,4], Vitali

Sintchenko[1,3,4], and Verlaine Timms[1,3]

[1]Centre for Infectious Diseases and Microbiology – Public Health, Westmead Hospital,

Westmead, NSW, Australia

[2]Centre for Infectious Diseases and Microbiology Laboratory Services, Westmead Hospital,

Westmead, NSW, Australia

[3]Westmead Clinical School, The University of Sydney, Westmead, NSW, Australia

[4]Marie Bashir Institute for Infectious Diseases and Biosecurity, The University of Sydney,

Camperdown, NSW, Australia

Corresponding author: Winkie Fong
Email: wfon4473@sydney.edu.au
17

<u>Abstract</u>

Whooping cough, or pertussis, is a highly transmissible respiratory infection caused by

Bordetella pertussis. Due to the high burden of pertussis, vaccine programmes were

introduced internationally and in Australia since the 1950s. This has resulted in a significant

decrease of pertussis infections. However, since the 1990s the number of pertussis

notifications has increased considerably. Currently circulating *B. pertussis* strains differ in

vaccine antigen composition compared to strains that circulated in the pre-vaccination era.

These genetic differences are thought to contribute, in part, to the re-emergence of pertussis

in Australia and around the world. Whole genome sequencing (WGS) can resolve minute

differences in circulating strains and provides unparalleled resolution of vaccine antigens.

This high-resolution snapshot can provide clues that enable more targeted public health

interventions. However, pertussis is primarily diagnosed with culture-independent diagnostic

assays which offer fast turnaround result times and reduced laboratory costs, eliminating the

need to culture isolates. Current WGS methods require a cultured isolate, resulting in an

absence of *B. pertussis* genome sequences in the post vaccination era. This scarcity has, in

turn, limited understanding of currently circulating strains and respective vaccine antigen

compositions.

Recent advancements of WGS technologies have allowed direct sequencing of clinical

specimens without the need for a cultured isolate. However, recovering reliable sequence

data from clinical samples of low bacterial load infections such as *B. pertussis* is a pressing

challenge. We sought to increase the yield of *B. pertussis* sequences direct from a clinical

sample by evaluating widely available WGS library preparation methods.

We report that the Illumina DNA prep library preparation kit combined with deep sequencing

allowed the detection of important surveillance information such as allelic variations in the *B.*

pertussis vaccine antigens. Further, our method generates high coverage over the 23S

ribosomal RNA of *B. pertussis* enabling macrolide resistance to be easily determined.

Overall, this method can improve surveillance of *B. pertussis,* by monitoring changes in

vaccine antigens, detecting antimicrobial resistance and guiding Public Health control

interventions.

<u>Introduction</u>

Bordetella pertussis, the primary causative agent of whooping cough, is a highly contagious

respiratory pathogen that causes mild symptoms in adults but severe infection with a high

mortality rate in infants. Global vaccination efforts have significantly reduced pertussis

mortality, however, pertussis remains endemic globally, even in countries with high vaccine

coverage[1]. The cause of the recent resurgence is hypothesised to include waning host

immunity[2] and vaccine escape[3]. Investigation of the persistence of pertussis has been

challenged by the lack of *B. pertussis* isolates. The introduction of highly sensitive and fast

PCR-based diagnostics has rendered culture of *B. pertussis* largely redundant, and limited the

ability to perform phenotypic strain typing and antibiotic susceptibility testing[4]. The re-

emergence of pertussis, along with the emergence of macrolide resistant *B. pertussis*,

demonstrates how critical it is that we continue to monitor *B. pertussis* strains, despite a

decline in the number of culture isolates.

Whole genome sequencing (WGS) of bacteria with epidemic potential provides highly

nuanced data[5, 6] used in outbreak control where it can identify clusters and potential

transmission chains[6]. Read depth, defined as the number of reads over a genome position, and

genome coverage, the percentage of read coverage across the genome, are important

parameters in direct sequencing protocols as they provide assurance of the accuracy and

quality of the consensus sequence. When this high-resolution technology is applied directly

to a clinical sample the overwhelming majority of sequences obtained are of human origin. In

a respiratory sample, the number of human cells poses the greatest obstacle to resolving

sufficient sequencing reads for a low-load pathogen, such as *B. pertussis*[6]. Studies have

previously shown that pre-treatment of nasopharyngeal aspirates (NPA) can reduce human

DNA levels, and increase target pathogen DNA and therefore pathogen specific sequencing

reads[4, 7]. However, despite the increased yield of *B. pertussis* from a saponin treated sample,

microbial genome coverage and read depth was still insufficient in providing informative

molecular typing data[4]. One way to improve sequence coverage is increase the number of

total reads per samples, known as deep sequencing. Deep sequencing aims to increase the

read depth and the accuracy of detecting mutations over positions in genes of interest by

generating more unique reads across the genome[8]. However, read depth and coverage can

also be enhanced by selecting the appropriate library preparation method[9].

Library preparation kits play a critical role in coverage yield, as the method of fragmentation

used can introduce significant biases. For *B. pertussis*, G+C bias is a major limiting factor in

transposase and PCR-based library preparation methods as it results in a loss of reads and low

coverage over high G+C content areas of the genome[10-13]. Considering *B. pertussis* has an

average G+C content of 67%, library preparation methods (e.g., Nextera XT) can bias library

construction to human DNA present and not produce libraries as efficiently from *B. pertussis*

DNA present in a clinical sample[10-13]. Moreover, in contrast to mechanical fragmentation-

based library preparation methods, enzymatic fragmentation contains slightly greater

insertion biases due to sequence preferences of the transposase[14-16]. While there is little

impact of these insertion biases on rapid, parallel sequencing of isolates, the effect of this bias

on sequencing of a high G+C content pathogen directly from clinical samples cannot be

ignored and requires further investigation[12, 17].

This study examines the influence of library preparation on culture-independent sequencing

of *B. pertussis*. We compare library preparation kits from Illumina, the most widely used

platform in Public Health Microbiology laboratories and assess the read depth and genome

coverage of a low-load, high GC pathogen. Further, by comparing libraries at increased

sequencing depths we investigate whether accurate genomic information relevant for public

health control of *B. pertussis* can be recovered directly from a clinical sample.

<u>Methods</u>

Nasopharygeal aspirates (NPA) were collected from the Centre of Infectious Diseases and

Microbiology – Laboratory Services (CIDMLS), NSW Health Pathology, Westmead

Hospital, Sydney. These NPA specimens were submitted for clinical testing of other

infectious diseases and were negative for *B. pertussis*. The samples were pooled and 200 μL

was taken for baseline extractions with the Qiagen DNeasy Blood and Tissue (QIAGEN,

Germany) DNA Extraction Kit (further described below) to confirm the NPA was *B.*

pertussis negative by PCR. The pooled NPA were then spiked with 0.5 McFarland

suspension of *B. pertussis* (ATCC®9797™ 18323) corresponding to approximately 1.4×10^6

CFU/mL (BP10-2), 1.4×10^5 CFU/mL (BP10-3), and 1.4×10^4 CFU/mL (BP10-4), then

stored at -20°C. The specific dilutions were chosen as C_T cycles by *IS481* real-time PCR

(rtPCR) demonstrate these are within the normal range of clinical samples[18]. Further clinical

NPA (n=2) that were positive for *B. pertussis* and collected by CIDMLS with the highest C_T

cycle were included for standard and deep sequencing. Details of *IS481* PCR are presented in

the Supplementary Material.

Extraction and DNA Quality Control

The spiked and clinical samples were treated with 0.025% Saponin and TurboDNase as

previously described[4, 7]. The spiked samples were then extracted with the Qiagen DNeasy

Blood and Tissue (QIAGEN, Germany) DNA Extraction Kit for "Purification of Total DNA

from Animal Tissues (Spin-Column Protocol)" with modifications. The modifications

included lysis of the NPA with Proteinase K at 56°C for 1.5 hours. The DNA extracts were

split into four aliquots for use in rtPCR targeting *IS481* and *ERV3* as previously described[4],

and for use in library preparation method comparison (Supplementary Material).

Library Preparation and Sequencing

The three Illumina (Illumina, USA) library preparation kits selected were; NexteraXT DNA

Library Preparation Kit v2.5, TruSeq DNA Nano Library Preparation Kit and the Illumina

DNA prep Library Preparation Kit. Libraries were constructed following the manufacturer's

protocol. The two clinical NPA samples were prepared with Illumina DNA prep only.

Fragment size and quality was assessed by the Agilent TapeStation 4200 using the High-

Sensitivity DNA ScreenTape Assay (Agilent, USA) on TruSeq input DNA pre- and post-

sonication, and Nextera XT, Illumina DNA prep and TruSeq post-library construction.

Library quantification was assessed using the KAPA (Roche Diagnostics, Switzerland) assay

on all libraries for normalisation and final pool concentration. Illumina DNA prep libraries

were sequenced on the Illumina NextSeq 500 instrument in conjunction with other routine

samples that generate between 3-5 million reads per sample. TruSeq and Illumina DNA prep

libraries were sequenced together on the Illumina MiniSeq High-Throughput platform. These

were considered as sequenced on a standard level. In addition, Illumina DNA prep libraries

were sequenced again on the Illumina NextSeq 500 with a High-Throughput flow cell for

deep sequencing to generate around 80 million reads per sample.

Analysis

Quality Control

Analysis began with FastQC[19](v 0.11.3) to assess the sequencing quality of raw reads,

followed by trimming the ends of poor-quality reads with Trimmomatic (v0.36)[20] using

optimised parameters (Leading:3 Trailing:3 SlidingWindow:4:20 Minlen:36).

Mapping

Trimmed reads were then mapped to the Human genome GRCh38.p12

(GCA_000001405.27) by Burrows-Wheelers Aligner (BWA v0.7.12) using default settings[21].

Three strains of *B. pertussis* were utilised as mapping reference genomes – Tohama I

reference genome (NC_002929.2), ATCC®9797™ 18323 whole genome (NC_018518.1) as

this was the strain used to spike the NPA, or B1917 (NZ_CP009751.1) as this is a

representative strain of currently circulating *B. pertussis*[22].

Gene and Genome Counts

160 Percentage of each representative organism (Human and *B. pertussis*) was determined by the

161 number of reads mapped to the respective genomes divided by the number of reads of the

162 total sample.

$$\% \text{ of } B.\,pertussis \text{ reads} = \frac{\text{No. of } B.\,pertussis \text{ reads}}{\text{No. of Total reads}}$$

Manual visualisations of coverage across genes of interest such as the 23S ribosomal RNA,

vaccine antigen genes: *ptx, prn,* and *fhaB* encoding regions were performed using Qiagen

CLC Genomics Workbench 12 (Qiagen, Germany). The list of all genes of interest has been

provided in Supplementary Material.

HTSeq (v11.2)[23] with default 'union' settings was used to count the number of unique reads

mapped over these genes of interest, mapped reads over multiple genes were discarded.

Samtools depth (v1.9)[24] and BEDtools intersectBed and genomecov (v2.25.0)[25] were used to

calculate the average read depth, gene and genome coverage and plotted with ggplot2

(v3.3.0)[26].

<u>Results</u>

The first question this study aimed to address was whether the method of library preparation

had any impact on the sequencing efficiency of a high G+C, low load microbial pathogen

such as *B. pertussis*.

Table 1: Standard sequencing comparison of library preparation methods by average total number of reads across duplicates, percentage of human and *B. pertussis* reads, average coverage depth of the *B. pertussis* Tohama I reference genome and total coverage of both *B. pertussis* Tohama I and ATCC®9797™ 18323 genomes.

Library Preparation Method	CFU/mL	Post-depletion *IS481* PCR (C_T)	Post-depletion *ERV3* PCR (C_T)	Total Reads	% of Human reads	% of *B. pertussis* reads	Average Coverage Depth	Coverage of the *B. pertussis* Tohama I Genome (%)	Coverage of the *B. pertussis* ATCC Genome (%)
Nextera XT									
BP10-2	1.4×10^6	23.53	36.48	1,465,653	40.9%	0.44%	1.36	57.02%	56.11%
BP10-3	1.4×10^5	28.72	36.35	3,446,052	36.7%	1.29%	0.48	3.77%	3.77%
BP10-4	1.4×10^4	32.75	36.09	1,690,120	55.8%	0.30%	0.18	0.77%	0.78%
Illumina DNA prep									
BP10-2	1.4×10^6	Same as above		5,207,437	51.0%	4.25%	4.25	89.16%	93.28%
BP10-3	1.4×10^5	Same as above		7,635,405	36.4%	0.89%	0.89	13.31%	13.78%
BP10-4	1.4×10^4	Same as above		3,095,938	57.8%	0.37%	0.01	1.26%	1.27%
TruSeq									
BP10-2	1.4×10^6	Same as above		Not Sequenced					
BP10-3	1.4×10^5	Same as above		Not Sequenced					
BP10-4	1.4×10^4	Same as above		514,032	67.5%	0.33%	0.34	0.33%	0.34%

NexteraXT DNA Library Preparation using standard sequencing

As expected, the percentage of *B. pertussis* reads identified decreased as CFU decreased (Table 1). Genome mapping to both *B. pertussis* (Tohama I and ATCC®9797™ 18323) genomes show only $56.5 \pm 0.5\%$ of the genomes are covered by these reads. Deeper analysis into coverage over the vaccine antigen and 23S ribosomal RNA showed poor coverage across the vaccine antigen genes, with BP10-2-1 failing (zero reads) to present reads to 10 out of 14 genes and BP10-2-2 failing 3 out of 14. BP10-3-1 contained reads to all genes, 170 reads mapped to *fhaB* which accounted for 85.6% of the gene. BP10-3-2, BP10-4-1 and BP10-4-2 failed to produce reads for 10, 10 and 11 genes out of 14, respectively.

Illumina DNA Library Preparation using standard sequencing

Similar to Nextera XT, the percentage of *B. pertussis* reads recovered decreased with decreasing CFU. However, unlike Nextera XT, genome mapping of BP10-2 showed Illumina DNA prep reads covered $91.2 \pm 2.1\%$ of both Tohama I and ATCC®9797™ 18323 genomes, thus, increasing whole genome coverage by 34.7%. Only one sample of BP10-4 was sequenced due to BP10-4-2 failing to pass library QC checkpoints with low library concentrations at the correct size of 300-600 bp. Compared to Nextera XT, Illumina DNA prep resulted in a more even distribution of reads across the vaccine antigens and 23S ribosomal RNA (Figure 1 and Figure 3). BP10-2 Illumina DNA prep at a standard sequencing level was able to provide 4.1 X coverage across all 14 vaccine antigen genes. BP10-3 and BP10-4 produced zero coverage for 6 and 10 genes respectively out of the 14. To test whether the library preparation result could be enhanced by deep sequencing, we sequenced these libraries (BP10-2, BP10-3 and BP10-4) at a concentration estimated to generate 80 million reads.

TruSeq DNA Nano Library Preparation using standard sequencing

Most samples of the TruSeq library preparation showed promising library concentration (236

– 3,350 pg/µL), however the majority of the library fragments were only 92-106 bp in size,

which were too small to be sequenced. Only BP10-4-2 had a library peak at 490 bp with a

concentration of 37.6 pg/µL, hence only this sample was sequenced. BP10-4-2 produced

127,330 reads with 426 reads to *B. pertussis,* these reads provided a depth of 0.34 X across

0.33% of the genome. TruSeq at standard sequencing level of BP10-4-2 failed to yield any

reads to genes of interest (Supplementary Material)

Figure 1: Comparison of Standard Sequencing (SS) gene coverage map over the sequences of the vaccine antigens – *cyaA, fhaB, fim2, fim3, prn, ptxA and ptxP*. The figure also compares the Illumina DNA prep (FX) (Red) and Nextera XT (XT) (Teal) kits at a SS level. The TruSeq (TN) kit was not presented in this figure as no reads were obtained across these regions. Coverage was capped at a maximum of 40x coverage.

Comparisons of Illumina DNA prep at Standard and Deep Sequencing Levels

Based on the results above, all Illumina DNA prep libraries were re-sequenced on a High-Throughput Flow cell on the NextSeq 500 platform, delivering a total of 493,937,514 reads to 5 samples – BP10-2 (n=2), BP10-3 (n=2) and BP10-4 (n=1). Average read depth and coverage across the genome are presented in

Table **2**, and library quantification C_T cycles are presented in the Supplementary Material.

The number of reads in deep sequencing was amplified 16-18 times and accordingly

improved average read depth across the genome by 5-16-fold. In the Illumina DNA prep deep

sequencing level, BP10-2 reads provided full 100% coverage of all 14 genes with 50-90X

depth, however BP10-3 and BP10-4 had partial coverage only.

Comparisons of standard and deep sequencing across the vaccine antigen genes of all

Illumina DNA prep libraries are presented in Figure 2. Given that a mutation (A → G)

present in position 2037 of *B. pertussis* Tohama I has been accepted as indicative of

macrolide resistance in *B. pertussis*, coverage of this region was investigated (Figure 3). All

kits provided coverage over position 2037 ranging from 66 X coverage in BP10-2-1 standard

level Nextera XT to 2795 X coverage in BP10-2-1 deep sequencing Illumina DNA prep.

Table 2: Summary of Illumina DNA prep deep sequencing average reads between duplicate samples.

	CFU/mL	Post-depletion *IS481* PCR (C_T)	Post-depletion *ERV3* PCR (C_T)	KAPA PCR (C_T)	Total Reads	% of human DNA reads	% of *B. pertussis* DNA reads	Average read depth	Coverage across *B. pertussis* Tohama I (%)
BP10-2	1.4×10^6	23.53	36.48	15.44	82,375,087	53.44%	2.71%	67.93	95.22
BP10-3	1.4×10^5	28.72	36.35	14.32	138,165,091	49.76%	2.05%	16.07	4.68
BP10-4	1.4×10^4	32.73	36.09	16.32	52,857,158	60.18%	0.45%	3.86	1.81
BP12	N/A	25.71	-	19.04	48,289,160	27.84%	2.20%	20.00	98.82
BP16	N/A	22.30	38.56	15.96	6,773,770	31.18%	4.14%	9.70	95.68

Figure 2. Gene coverage map for genes coding for vaccine antigens. Sequencing at Standard

Sequencing (Blue) and Deep Sequencing (Red) levels of Illumina DNA prep (FX) library are

shown. Coverage in this figure was capped at 125, only small regions of the genes contained

reads exceeding 400X coverage.

Figure 3. Coverage of the 23S rRNA gene of *B. pertussis*, comparing library preparation kits

- Illumina DNA prep (FX) and Nextera XT (XT), and Deep Sequencing (DS) and Standard

Sequencing (SS) depths, with varying *B. pertussis* spiked CFU concentrations (BP10-2 1.4 x

10^6 CFU/mL and BP10-3 1.4 x 10^5 CFU/mL duplicates).

Trials on Clinical NPA specimens

Standard sequencing using Illumina DNA prep of the two clinical samples BP12 and BP16,
yielded 3,443,046 and 514,032 total reads, with 70,960 (2.06%) and 24,141 (4.69%) *B.*
pertussis reads, respectively. These reads produced an average coverage depth of 2.54 X for
BP12 and 1.57 X for BP16. Furthermore, BP12 reads covered 50.2% of the B1917
(NZ_CP009751.1) genome, while BP16 covered 42.5%. When these libraries were
sequenced at a deeper level the genome coverage increased by +48.7% and +53.2% for BP12
and BP16 respectively

Table **2** and Figure 4. Analysis of the genes of interest showed variability in recovery with

most genes having more than 92% coverage, with an average depth of 32.91 X and 10.98 X

respectively (Supplementary Material).

Figure 4: Whole genome coverage map of the positive strand in BP12 and BP16 at Standard

Sequencing (SS) (Blue and Pink) and Deep Sequencing (DS) Levels (Purple and Green),

mapped to B1917 (NZ_CP009751.1). Panel A represents BP12 and Panel B represents BP16.

Scale of both samples have been capped at 50 X depth.

197
<u>Discussion</u>

In the era of culture-independent diagnostic testing, the primary consequence is the loss of

routine laboratory culture. Without cultured isolates, many gold-standard phenotypic typing

systems have been lost. Therefore, further laboratory classification of currently circulating

strains has become unavailable and this in turn affects public health outbreak control

strategies. The aim of this study was to compare commercially available library preparation

methods and sequencing depths to enhance our ability to directly sequence *Bordetella*

pertussis-positive nasopharyngeal aspirates. Comparisons were made based on sequencing

read depths, average genome coverage and the ability to accurately extract sequencing data

from regions of interest such as the vaccine antigens.

This study demonstrated that different library preparation methods can impact the recovery of

important genomic information from culture-independent sequencing. Nextera XT libraries at

a standard sequencing depth produced uneven coverage across the genome, only accounting

for ~50% of the *B. pertussis* Tohama I genome, compared to more than 80% of the genome

when libraries are prepared with Illumina DNA prep. Overall, the Illumina DNA prep library

preparation kit generated better average coverage across the *B. pertussis* genome at both the

standard and deep sequencing depths. Sequencing at a deeper level only amplified depth and

increased confidence to detect SNPs. In line with what others have shown[4], we found that

sequencing at a standard level with any library preparation method, was insufficient in

obtaining accurate molecular typing information.

We report variations in total reads between all Illumina DNA prep samples most likely due to

library normalisation and final library pooling. Interestingly, we noted that in this study the

library concentrations ranged between 3 C_T cycles from each other and these C_T cycles

reflected how many sequencing reads were dedicated to each sample. This variation is

generally acceptable at a standard sequencing scale, hence further investigations into the

effect of library concentration and normalisation would need to be performed to optimise

deep sequencing protocols and ensure an even distribution of sequencing resources for all

samples.

The TruSeq DNA Nano library kit was chosen based on its fragmentation method –

mechanical instead of enzymatic to reduce the inherent G+C bias of the transposase.

However, post-library preparation QC determined very poor concentrations of library

fragments in the appropriate range (400-500bp). TruSeq libraries potentially failed due to low

starting concentration of DNA given the sample had been host DNA depleted. Fragment size

analysis was performed before and after sonication on BP10-2 samples, however low input

DNA concentration made it difficult to observe fragment size peaks efficiently. Sequencing

of one of the TruSeq libraries (BP10-4-2) proceeded due to the presence of a peak in the 400-

500bp region. TruSeq libraries were constructed with other samples with higher starting

DNA concentration, which sequenced successfully, hence library preparation methods and

reagents were not the problem. Therefore, TruSeq was not an appropriate kit for low DNA

concentrations as selection of sonication parameters is difficult to optimise for host depleted

clinical samples with low concentration of DNA.

As expected, decreasing *B. pertussis* CFU in spiked samples were able to demonstrate that

changes in bacterial load can influence the amount of sequencing reads recovered from a

sample. Based on routine rtPCR results, clinical *IS481*-positive specimens have an average

C_T cycle of 30.92, suggesting a CFU load of 1.4 x 10^4 (CFU/mL) equivalent to sample BP10-

4, is a typical *B. pertussis* load for clinical NPA samples. As such, the higher number of *B.*

pertussis cells in BP10-2 of 1.4 x10^6 CFU is an uncommon occurrence in a clinical sample.

The two clinical cases enrolled in this study had a very high initial C_T cycle of 12.15 (BP12)

and 17.97 (BP16), and their total read counts of 48,289,160 and 6,773,770, respectively,

demonstrate the importance of quality clinical sample collection ensuring high load of target

bacterial DNA prior to sequencing to allow sufficient genome coverage.

To achieve adequate depth and coverage of the *B. pertussis* genome in a low bacterial load

NPA sample would require further enrichment of the clinical sample and/or library.

However, our results indicate that a *B. pertussis* positive sample with a CFU equivalent to or

more than 1.4 x10^6 CFU would, without enrichment, generate coverage and depth sufficient

for genomic surveillance analyses. By combining host DNA depletion protocols (Saponin),

an appropriate library preparation method (Illumina DNA prep) and applying a deep

sequencing approach (>50 million reads) with a library quantification result of more than 15

C_T cycles, should result in 20X coverage at least, across the regions of interest if not the

whole genome of *B. pertussis*. While sequencing at a deeper level costs more compared to

sequencing a pure isolate, it is now an option when there are few isolates to sequence in this

era of PCR diagnostics.

This study was limited by the number of replicates that could be performed as a result of

cost-limitations; hence results are subjected to sample bias. Numbers of NPA samples are

also slowly declining as preferences to nasopharyngeal swabs are more popular. The volume

of liquid in NPA are also very limited due it its use in previous diagnostic protocols and

therefore replicates of clinical samples could not be performed. Trialling the protocol on

more common respiratory sample types such as nasopharyngeal and throat swabs would also

expand its use.

Future research should focus on improving sequencing depth and coverage over regions of

interest, particularly genes coding for vaccine antigens and other molecular typing targets.

Despite the expense, further deep sequencing trials on larger number of clinical NPA are

required to validate the protocol and determine information yield across sample types.

In conclusion, our study demonstrated the feasibility of direct sequencing of *B. pertussis* from

clinical NPA specimens with a high bacterial load and the recognition of potentially

actionable targets in *B. pertussis* genome. This can be achieved through the combination of

optimised sample and library preparation followed by deep sequencing.

Author Statements

The study was conceptualised by WF, VT, and VS. Laboratory work was performed by WF,

RR and VT. Bioinformatics and genome analysis was executed by WF, KP and RS. The

manuscript was written by WF and reviewed and edited by KP, RR, RS, VT and VS.

The authors declare no conflict of interest

Reads mapping to *Bordetella pertussis* ATCC9797 18323 for spiked nasopharyngeal

aspirates and BP1917 for clinical nasopharyngeal aspirates have been uploaded to SRA under

BioProject: PRJNA694997

Consent for the images in this article has been provided by the authors.

This work was supported by the Prevention Research Support Program, funded by the New

South Wales Ministry of Health. Special thanks to Illumina for providing a complimentary

Illumina DNA prep Library Preparation Kit. The authors are grateful to the staff of Centre for

Infectious Diseases and Microbiology Laboratory Services, NSW Health Pathology for their

technical assistance and expertise. Computational analysis was performed on the University

of Sydney High Performance Cluster, with the assistance of the Sydney Informatics Hub. The

authors would like to acknowledge the Microbial Genomics Reference Laboratory, Centre for

Infectious Disease and Microbiology –Public Health, Westmead Hospital, for their assistance

with genome sequencing and bioinformatics analysis.

Nasopharyngeal aspirates were collected by the Centre for Infectious Diseases and

Microbiology Laboratory services under the Western Sydney Local Health District Research

Ethics and Governance committee. Project identifier: 2019/PID02294

Consent was not obtained from patients, as these NPA were left-over from previous

diagnostic testing, and otherwise discarded. These NPA samples were pooled to prevent any

identification during sequencing, and no record of identifiable data was collected.

319
<u>References</u>

1.   Crowcroft NS, Stein C, Duclos P, Birmingham M. How best to estimate the global
burden of pertussis? The Lancet Infectious Diseases. 2003;3(7):413-8. doi:10.1016/s1473-
3099(03)00669-8
2.   Schwartz KL, Kwong JC, Deeks SL, Campitelli MA, Jamieson FB, Marchand-Austin
A, et al. Effectiveness of pertussis vaccination and duration of immunity. CMAJ.
2016;188(16):E399-E406. doi:10.1503/cmaj.160193
3.   Bart MJ, van Gent M, van der Heide HG, Boekhorst J, Hermans P, Parkhill J, et al.
Comparative genomics of prevaccination and modern Bordetella pertussis strains. BMC
Genomics. 2010;11:627. doi:10.1186/1471-2164-11-627
4.   Fong W, Rockett R, Timms V, Sintchenko V. Optimization of sample preparation for
culture-independent sequencing of Bordetella pertussis. Microb Genom. 2020.
doi:10.1099/mgen.0.000332
5.   Yang J, Yang F, Ren L, Xiong Z, Wu Z, Dong J, et al. Unbiased parallel detection of
viral pathogens in clinical samples by use of a metagenomic approach. J Clin Microbiol.
2011;49(10):3463-9. doi:10.1128/JCM.00273-11
6.   Bachmann NL, Rockett RJ, Timms VJ, Sintchenko V. Advances in Clinical Sample
Preparation for Identification and Characterization of Bacterial Pathogens Using
Metagenomics. Front Public Health. 2018;6:363. doi:10.3389/fpubh.2018.00363
7.   Hasan MR, Rawat A, Tang P, Jithesh PV, Thomas E, Tan R, et al. Depletion of
Human DNA in Spiked Clinical Specimens for Improvement of Sensitivity of Pathogen
Detection by Next-Generation Sequencing. J Clin Microbiol. 2016;54(4):919-27.
doi:10.1128/JCM.03050-15
8.   Depledge DP, Kundu S, Jensen NJ, Gray ER, Jones M, Steinberg S, et al. Deep
sequencing of viral genomes provides insight into the evolution and pathogenesis of varicella

zoster virus and its vaccine in humans. Mol Biol Evol. 2014;31(2):397-409.
doi:10.1093/molbev/mst210
9.   Rhodes J, Beale MA, Fisher MC. Illuminating choices for library prep: a comparison
of library preparation methods for whole genome sequencing of Cryptococcus neoformans
using Illumina HiSeq. PLoS One. 2014;9(11):e113501. doi:10.1371/journal.pone.0113501
10.   Kozarewa I, Ning Z, Quail MA, Sanders MJ, Berriman M, Turner DJ. Amplification-
free Illumina sequencing-library preparation facilitates improved mapping and assembly of
(G+C)-biased genomes. Nat Methods. 2009;6(4):291-5. doi:10.1038/nmeth.1311
11.   Quail MA, Kozarewa I, Smith F, Scally A, Stephens PJ, Durbin R, et al. A large
genome center's improvements to the Illumina sequencing system. Nat Methods.
2008;5(12):1005-10. doi:10.1038/nmeth.1270
12.   Adey A, Morrison HG, Asan, Xun X, Kitzman JO, Turner EH, et al. Rapid, low-
input, low-bias construction of shotgun fragment libraries by high-density in vitro
transposition. Genome Biol. 2010;11(12):R119. doi:10.1186/gb-2010-11-12-r119
13.   Hillier LW, Marth GT, Quinlan AR, Dooling D, Fewell G, Barnett D, et al. Whole-
genome sequencing and variant discovery in C. elegans. Nat Methods. 2008;5(2):183-8.
doi:10.1038/nmeth.1179
14.   Green B, Bouchier C, Fairhead C, Craig NL, Cormack BP. Insertion site preference of
Mu, Tn5, and Tn7 transposons. Mob DNA. 2012;3(1):3. doi:10.1186/1759-8753-3-3
15.   Lan JH, Yin Y, Reed EF, Moua K, Thomas K, Zhang Q. Impact of three Illumina
library construction methods on GC bias and HLA genotype calling. Hum Immunol.
2015;76(2-3):166-75. doi:10.1016/j.humimm.2014.12.016
16.   Kia A, Gloeckner C, Osothprarop T, Gormley N, Bomati E, Stephenson M, et al.
Improved genome sequencing using an engineered transposase. BMC Biotechnol.
2017;17(1):6. doi:10.1186/s12896-016-0326-1
17.   Sato MP, Ogura Y, Nakamura K, Nishida R, Gotoh Y, Hayashi M, et al. Comparison
of the sequencing bias of currently available library preparation kits for Illumina sequencing
of bacterial genomes and metagenomes. DNA Res. 2019;26(5):391-8.
doi:10.1093/dnares/dsz017
18.   Timms VJ, Fong W, Jeoffreys NJ, Sintchenko V. Evaluation of the BioGX BD-Max
PCR assay for detection of pathogenic Bordetella. Pathology. 2019;51(3):323-4.
doi:10.1016/j.pathol.2018.10.018
19.   Andrews S. FastQC: A Quality Control Tool for High Throughput Sequence Data
2010 [Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/
20.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina
sequence data. Bioinformatics. 2014;30(15):2114-20. doi:10.1093/bioinformatics/btu170
21.   Li H, Durbin R. Fast and accurate short read alignment with Burrows-Wheeler
transform. Bioinformatics. 2009;25(14):1754-60. doi:10.1093/bioinformatics/btp324
22.   Bart MJ, Zeddeman A, van der Heide HG, Heuvelman K, van Gent M, Mooi FR.
Complete Genome Sequences of Bordetella pertussis Isolates B1917 and B1920,
Representing Two Predominant Global Lineages. Genome Announc. 2014;2(6).
doi:10.1128/genomeA.01301-14
23.   Anders S, Pyl PT, Huber W. HTSeq--a Python framework to work with high-
throughput sequencing data. Bioinformatics. 2015;31(2):166-9.
doi:10.1093/bioinformatics/btu638
24.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25(16):2078-9.
doi:10.1093/bioinformatics/btp352
25.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 2010;26(6):841-2. doi:10.1093/bioinformatics/btq033

26.   Wickham H. ggplot2: Elegant Graphics for Data Analysis. New York: Springer-
Verlag; 2016.
27.   Bruinsma S, Burgess J, Schlingman D, Czyz A, Morrell N, Ballenger C, et al. Bead-
linked transposomes enable a normalization-free workflow for NGS library preparation.
BMC Genomics. 2018;19(1):722. doi:10.1186/s12864-018-5096-9
400