Long-reads are revolutionizing 20 years of insect genome sequencing

Scott Hotaling[1]*, John S. Sproul[2], Jacqueline Heckenhauer[3,4], Ashlyn Powell[5], Amanda M.

Larracuente[2], Steffen U. Pauls[3,4,6], Joanna L. Kelley[1], and Paul B. Frandsen[3,5,7]*

Affiliations:

[1] School of Biological Sciences, Washington State University, Pullman, WA, USA

[2] Department of Biology, University of Rochester, Rochester, NY, USA

[3] LOEWE Centre for Translational Biodiversity Genomics (LOEWE-TBG), Frankfurt, Germany

[4] Department of Terrestrial Zoology, Entomology III, Senckenberg Research Institute and Natural

History Museum Frankfurt, Frankfurt, Germany

[5] Department of Plant and Wildlife Sciences, Brigham Young University, Provo, UT, USA

[6] Institute for Insect Biotechnology, Justus-Liebig-University, Giessen, Germany

[7] Data Science Lab, Smithsonian Institution, Washington, DC, USA

*Authors for Correspondence:

Scott Hotaling, School of Biological Sciences, Washington State University, Pullman, WA, 99164,

USA; Email: scott.hotaling@wsu.edu: Phone: (828) 507-9950

Paul B. Frandsen, Department of Plant and Wildlife Sciences, Brigham Young University, Provo,

UT, 84602, USA; Email: paul_frandsen@byu.edu; Phone: (804) 422-2283

Abstract:

The first insect genome (*Drosophila melanogaster*) was published two decades ago. Today,

nuclear genome assemblies are available for a staggering 601 insect species representing 20

orders. In this study, we analyzed the most-contiguous assembly for each species and provide a

"state of the field" perspective, emphasizing taxonomic representation, assembly quality, gene

completeness, and sequencing technologies. Relative to species richness, genomic efforts have

been biased towards four orders (Diptera, Hymenoptera, Collembola, and Phasmatodea),

Coleoptera are underrepresented, and 11 orders still lack a publicly available genome assembly.

The average insect genome assembly is 439.2 megabases in length with 87.5% of single-copy

benchmarking genes intact. Most notable has been the impact of long-read sequencing;

assemblies that incorporate long-reads are ~48x more contiguous than those that do not. We

offer four recommendations as we collectively continue building insect genome resources: (1)

seek better integration between independent research groups and consortia, (2) balance future

sampling between filling taxonomic gaps and generating data for targeted questions, (3) take

advantage of long read sequencing technologies, and (4) expand and improve gene annotations.

Keywords: Insecta, Arthropoda, arthropod genomics, long-read sequencing, Pacific

Biosciences, Oxford Nanopore

Significance statement:

Since the first insect genome was sequenced ~20 years ago, sequencing technologies and the

availability of insect genome assemblies have both advanced dramatically. In this study, we

curated, analyzed, and summarized the field of insect genomics in terms of taxonomic

representation, assembly quality, gene completeness, and sequencing technology. We show that

601 insect species have genome assemblies available, with some groups heavily

overrepresented (e.g., Diptera) relative to others (e.g., Coleoptera). The major takeaway of our

study is that genome assemblies produced with long reads are ~48x more contiguous than short

read assemblies.

Body:

Since the publication of the *Drosophila melanogaster* genome (Adams, et al. 2000) sequencing and analytical technologies have developed rapidly, bringing the power of genome sequencing to an ever-expanding pool of researchers. More than 600 insects have now had their nuclear genome sequenced and made publicly available in the GenBank repository (Sayers, et al. 2021). While representing just 0.06% of the ~1 million described insects (Stork 2018), this breadth of insect genome sequencing still spans ~480 million years of evolution (Misof, et al. 2014) and roughly two orders of genome size from the tiny 99 megabase (Mb) genome of *Belgica antarctica* (Kelley, et al. 2014) to the massive genome of *Locusta migratoria* at 6.5 gigabases (Gb; Wang, et al. 2014).

Accumulating genomic resources have transformed biological research and precipitated major advances in our understanding of the origins of biodiversity (Hug, et al. 2016; McGee, et al. 2020; McKenna, et al. 2019; Seehausen, et al. 2014). Considerable progress has been driven by large-scale consortia [e.g., Human Genome Project (Collins, et al. 2003); Vertebrate Genome Project (Rhie, et al. 2021)] and for insects, the most prominent consortium has been the i5K initiative to sequence genomes for 5,000 different arthropods (i5K Consortium 2013; Robinson, et al. 2011). The rise of long-read sequencing technologies—primarily Oxford Nanopore and Pacific Biosciences (PacBio)—have also changed the landscape of genome sequencing by providing an economical means for high-throughput generation of reads that are commonly 25 kilobases (Kb) or longer (Amarasinghe, et al. 2020), thereby greatly increasing the average size of sequences used in assemblies. Genome sequencing efforts in insects, however, have not been spread evenly. Aquatic insects, as a group, are underrepresented relative to their terrestrial counterparts (Hotaling, et al. 2020). And, some orders (e.g., Diptera) are represented by far more genome assemblies than their species diversity alone would warrant—likely reflecting the model organisms within them—while many orders still have no genomic representation.

Figure 1. Taxonomic representation, contiguity, and the timeline of availability for the most-contiguous

nuclear genome assembly for 601 insect species in GenBank as of November 2020. Only one assembly

per named species or subspecies is included. (a) *The taxonomic diversity of available insect genome*

assemblies. Observed versus expected numbers of genome assemblies represent the total number of available

assemblies versus those that would be expected given the proportion that each order comprises of all described

insect diversity. Significance was assessed with Fisher's exact tests. One order is underrepresented

(Coleoptera) while four orders are overrepresented (Diptera, Hymenoptera, Collembola, Phasmatodea). Eleven

orders (light red silhouettes) have no publicly available genome assembly. A breakdown of sequencing

technology by order is shown in Fig. S1. (b) *Genome contiguity versus total assembly length.* Contiguity was

assessed with contig N50, the mid-point of the contig distribution where 50% of the genome is assembled into

contigs of a given length or longer. The inset plot shows a comparison of contig N50 distributions for short-read

(n = 365) versus long-read (n = 126) assemblies. Significance was assessed with a Welch's T-test. A finer-scale

breakdown by sequencing technology is shown in Fig. S2. (c) *The timeline of genome assembly availability for*

insects according to the GenBank publication date. A steady increase in contiguity is largely precipitated by the

rise of long-read sequencing. Labeled in (b) and (c): well-known or outlier genome assemblies in terms of either

model status, assembly size, or contiguity. Groups of species in the same genus are labeled with black circles.

(d) *Contig N50 by taxonomic group.* Generally, taxa were grouped into orders except when 10 or more

assemblies were available for a lower taxonomic level (family or genus). As in (b) and (c), each point represents

a single insect genome assembly.

Here, we curated and analyzed the best available genome assembly for 601 insects (species or

subspecies). We provide a "state of the field" perspective emphasizing taxonomic representation,

assembly quality, gene completeness, and sequencing technology. We focused on taxonomic

breadth rather than within-group efforts [e.g., The *Anopheles gambiae* 1000 Genomes

Consortium (2017)] to gain a more holistic overview of the field. Following similar studies (e.g.,

Hotaling, et al. 2020; Misof, et al. 2014; Petersen, et al. 2019)*,* we defined insects to include all

groups within the subphylum Hexapoda. We downloaded metadata from GenBank for all nuclear

hexapod genome assemblies on an order-by-order basis (Sayers, et al. 2021; accessed 2

November 2020). We culled this data set to only include the assembly with the highest contig

N50 for each taxon and downloaded these assemblies for analysis. We acknowledge that this

filtering approach may introduce biases towards the present-day for assemblies that have been

improved over the years. Assemblies were classified as "short-read", "long-read", or "not

provided" based on whether only short-reads (e.g., Illumina) were used, any amount of long-read

sequences (e.g., PacBio) were used, or no information was provided. If an assembly used both

short- and long-reads (a "hybrid" assembly), it was classified as a long-read assembly in our

analysis.

To test if insect orders were under- or overrepresented in terms of genome assembly availability,

we compared the observed number of taxa with assemblies to the expected number given the

described diversity for a given order. We obtained totals for the number of insects described

overall and for each order from previous studies (Bellinger, et al. 2020; Zhang 2011). We

assessed significance between observed and expected representation with Fisher's exact tests.

To assess gene completeness, we ran "Benchmarking Universal Single Copy Orthologs"

(BUSCO) v.4.1.4 (Seppey, et al. 2019) on each assembly using the 1,367 reference genes in the

OrthoDB v.10 Insecta gene set (Kriventseva, et al. 2019). It should be noted that Collembola

genome assemblies may have received slightly lower BUSCO scores in this analysis because

non-insect hexapod genomes were not used to generate the Insecta gene set. We tested for

differences in distributions of contig N50 or assembly size between short- and long-read

assemblies with Welch's T-tests. Next, using the BUSCO gene set, we tested whether longer

genes were more likely to be missing or fragmented depending on sequencing technology (short-

or long-read) with Spearman's correlations. We defined BUSCO gene length as the full

nucleotide sequence for the protein-coding portions of the consensus "ancestral" genes included

in the OrthoDB v.10 Insecta gene set. An extended version of the methods and the scripts used

for analysis are provided in the Supplementary Materials and GitHub repository

(https://github.com/pbfrandsen/insect_genome_assemblies).

As of November 2020, 601 different insect species representing 20 orders had nuclear genome

assemblies available in GenBank. These data were dominated by Diptera (n = 169 assemblies),

Hymenoptera (n = 164), and Lepidoptera (n = 118; Fig. 1a). Four orders were overrepresented

relative to their species diversity: Collembola, Diptera, Hymenoptera, and Phasmatodea (P,

Fisher's < 0.03; Fig. 1a). Coleoptera, with 387,100 described species (Zhang 2011), was

significantly underrepresented (41 assemblies versus ~228 expected; P, Fisher's < 0.01). Six

orders were represented by only one genome assembly and 11 orders had no publicly available

assembly. This lack of representation was particularly striking for Neuroptera (5,868 described

species, Zhang 2011).

On average, insect genome assemblies were 439.2 Mb in length (SD = 448.4 Mb; Fig. 2a) with a

mean contig N50 of 1.09 Mb (SD = 4.01 Mb) and 87.5% (SD = 21%) BUSCO completeness

(single and duplicated genes, combined). Substantial variation existed in all three metrics,

however, with assemblies ranging from the highly incomplete assembly of *Piezodorus guildini* at

just 3.2 Mb (contig N50 = 1.5 Kb, BUSCO completeness = 0.2%) to the exceptionally high-quality

140.7 Mb assembly of *D. melanogaster* (contig N50 = 22.4 Mb, BUSCO completeness = 99.9%;

Fig. 2, Table S1). For orders represented by >10 taxa, Hymenoptera assemblies were the most

complete (BUSCO completeness = 94%, SD = 14.3%) and Lepidoptera the least (74.6%, SD =

28.2%; Fig. 2b). At 15.3%, Lepidoptera had the lowest percentage of long-read assemblies (Fig.

S1) and *Heliconius* assemblies were particularly fragmented (Fig. 1d). For families represented

by >10 taxa, Drosophilidae assemblies were the most complete (BUSCO completeness = 98.4%,

SD = 2%) followed closely by Apidae assemblies (97.9%, SD = 3.7%; Fig. 1d, 2b). As expected,

assemblies with higher contig N50 lengths were also more complete (Fig. 2f) but assembly size had little to no effect on gene completeness (Fig. S3).

Figure 2. Variation in assembly size and Benchmarking Universal Single-Copy Ortholog (BUSCO) gene completeness across Insecta. (a) Assembly size for all insects, grouped by order then family. To improve visualization, the upper display limit was set to 2.8 gigabases (Gb). Four genome assemblies exceeded this value and are labeled with gray text (in Gb). Taxa silhouettes were either handmade or taken from PhyloPic (http://phylopic.org). (b) BUSCO results for each insect genome assembly. Each horizontal bar represents one assembly (*n* = 601 species) and corresponds to the same taxon in the assembly size plot to the left in (a). (c-e) Long-read versus short-read genome assembly comparisons of (c) complete BUSCOs (single and duplicated combined), (d) fragmented BUSCOs, and (e) duplicated BUSCOs only. Significance was assessed with Welch's T-tests. (f) A comparison of BUSCO completeness versus contig N50. Each point represents the best available assembly for one taxon and groups of taxa in the same genus are labeled with black circles. Unsurprisingly, more contiguous genome assemblies also exhibit greater gene completeness. (g) Longer genes are more likely to be fragmented in insect genome assemblies, regardless of the technology used. However, a much stronger correlation exists between short-read assemblies and fragmentation of longer genes (Spearman's ρ: 0.24, *P* < 2.2e-16) than for long-read assemblies (Spearman's ρ: 0.08, *P* = 0.002). Unlike in (c-e), each circle in (g) represents the percent of fragmentation for that BUSCO gene across all long- or short-read assemblies. Thus, each gene is included twice (once for each technology). All 1,367 BUSCO genes in the OrthoDB v.10 Insecta gene set (Kriventseva, et al. 2019) were used except one 2.02 Kb gene that was missing in >70% of assemblies and subsequently removed from analysis and visualization. BUSCO gene lengths varied from 198 base pairs to 9.01 Kb.

The type(s) of sequence data used for genome assembly were obtained for ~82% of assemblies (long-read = 126, short-read = 365; Table S1). Long-read assemblies were more contiguous than short-read assemblies (Fig. 1b; *P,* Welch's T-test < 0.0001), averaging contig N50 values that were ~4.4 Mb higher despite no difference in assembly size (*P,* Welch's T-test = 0.12; Fig. S4). Gene regions were also far more complete in long-read assemblies (mean BUSCO completeness = 96%, SD = 7%) versus those generated from short-reads (89.1%, SD = 19%; *P,* Welch's T-test < 1e-8; Fig. 2c) with 70% fewer fragmented genes (*P,* Welch's T-test < 1e-11; Fig. 2d). Long-read assemblies, however, had ~2.6x more duplicated genes (4.4% vs. 1.7%; *P,*

Welch's T-test = 0.003; Fig. 2e). Longer BUSCO genes were also more likely to be fragmented

in both short-read (Spearman's ρ: 0.24, $P < 2.2e-16$) and long-read assemblies (Spearman's ρ:

0.08, $P = 0.002$; Fig. 2g) but they were less likely to be missing in both when compared to

shorter genes (short-read: Spearman's ρ: -0.08, $P = 0.002$; long-read: Spearman's ρ: -0.18, $P =$

9.7e-12; Fig. S5).

The rate at which new insect genome assemblies are becoming available is clearly accelerating

(Fig. 1c). Nearly 50% ($n = 292$) of the best-available insect assemblies were accessioned in

2019–2020 (Tables S1-S2). The same period also represented a high-water mark of contiguity

(mean contig N50, 2019–2020 = 1.77 Mb; Table S2). Much of the increase in contiguity was

driven by long-read assemblies which rose in frequency from 0% of all assemblies in 2011–2012

to 36.1% in 2019–2020. The contiguity of long-read assemblies also sharply increased in 2017

(Fig. S6, Table S2).

We have entered a new era of insect genome biology. Since 2019, a new species has had its

genome assembly deposited in GenBank every 2.3 days. These new assemblies are, on

average, markedly more contiguous than those of just a few years ago. As we continue

developing these resources, we offer four recommendations: first, we should recognize the

community-driven nature of these data and seek better integration between research groups and

consortia in terms of data sharing, best practices, and taxonomic focus. Progress towards these

goals is occurring (e.g., a proposed metric system for describing genome assembly quality with

associated benchmark standards from the Earth BioGenome Project, Lewin, et al. 2018) and will

accelerate as more researchers integrate these standards into their own workflows. Second, new

sequencing efforts should strive to balance sampling that fills taxonomic gaps and improves

existing resources with targeted sampling motivated by specific questions. Both approaches are

valuable and not mutually exclusive. The former—filling taxonomic gaps—is critical to broadly

understanding the evolution of insects, the most diverse animal group on Earth. While the

latter—targeted, question-driven sequencing—is critical to understanding specific aspects of

genome biology which are often best answered using dense sampling of specific groups.

Importantly, success for this recommendation will depend, in part, on our first recommendation.

Better integration and communication will limit redundancy of efforts where the same species'

genome is sequenced by multiple groups simultaneously. Third, we echo the findings of the

Vertebrate Genome Project (Rhie, et al. 2021)—long-read assemblies are vastly more

contiguous than short-read approaches—and recommend that these technologies be embraced

by insect genome scientists. And, fourth, as of 2019, only 40% of insect genome assemblies had

corresponding gene annotations in GenBank (Li, et al. 2019). Expanding and refining the

availability of gene annotations for insects will drive corresponding increases in the scale of

taxonomic comparisons that are possible for many analyses. Overcoming this challenge of

annotation quality and availability can be subdivided into two more specific calls: (1) whenever

possible, annotations should be made available alongside genome assemblies in GenBank or

similar public repositories and (2) researchers should consider using the NCBI Eukaryotic

Genome Annotation Pipeline (Thibaud-Nissen, et al. 2016) to limit variation introduced by

differing annotation approaches and maximize compatibility.

Beyond resource development, we must continue to leverage this data set to conduct new

studies of insect genome biology and evolution. These efforts are beginning to emerge and are

paying dividends. For instance, 76 arthropod genome assemblies were used to better

understand 500 million years of evolution by characterizing changes in gene and protein content

in a temporal and phylogenetic context, including the identification of novel gene families that

arose during diversification with links to key adaptations including flight (Thomas, et al. 2020).

Similarly, a study of 195 insect genomes revealed the high diversity of transposable elements

across insects with varying levels of conservation depending upon phylogenetic position (Gilbert,

et al. 2020). With genome assemblies representing 600+ taxa and ~480 million years of

evolution available in a public repository, the power and promise of insect genome research has

never been greater. While our focus was on insects, long-reads are likely revolutionizing genome

science in virtually all taxonomic groups with untapped genomic potential existing in public

repositories across the Tree of Life. The rise of long-read assemblies will, in particular, spur new

understanding of previously difficult to characterize aspects of the genome (e.g., genome

structure, highly repetitive regions). By continuing to build, curate, and make genomic resources

publicly available, we will gain tremendous insight into genome biology and evolution at broad

phylogenetic scales. We will also create a more inclusive and equitable discipline by expanding

access to resources for scientists whose participation has historically been limited by financial or

technological barriers.

Acknowledgements:

S.H. and J.L.K. were supported by NSF award OPP-1906015. J.H and S.U.P. were supported by

the LOEWE-Centre for Translational Biodiversity Genomics which is funded by the Hessen State

Ministry of Higher Education, Research and the Arts. J.S.S. was supported by an NSF

Postdoctoral Research Fellowship in Biology (DBI-1811930) and an NIH General Medical

Sciences award (R35GM119515) to A.M.L.

Data availability:

The data underlying this article are available in the Supplementary Materials, primarily in Table

S1, with associated scripts for analysis on GitHub:

https://github.com/pbfrandsen/insect_genome_assemblies.

References:

Adams MD, et al. 2000. The genome sequence of Drosophila melanogaster. Science 287: 2185-

2195.

Amarasinghe SL, et al. 2020. Opportunities and challenges in long-read sequencing data

analysis. Genome Biology 21: 1-16.

Bellinger PF, Christiansen KA, Janssens F 2020. Checklist of the Collembola of the World.

http://www.collembola.org.

Collins FS, Morgan M, Patrinos A 2003. The Human Genome Project: lessons from large-scale

biology. Science 300: 286-290.

Consortium AgG 2017. Genetic diversity of the African malaria vector Anopheles gambiae.

Nature 552: 96.

Gilbert C, Peccoud J, Cordaux R 2020. Transposable Elements and the Evolution of Insects.

Annual Review of Entomology 66.

Hotaling S, Kelley JL, Frandsen PB 2020. Aquatic insects are dramatically underrepresented in

genomic research. Insects 11: 601.

Hug LA, et al. 2016. A new view of the tree of life. Nature Microbiology 1: 1-6.

i5K Consortium 2013. The i5K Initiative: advancing arthropod genomics for knowledge, human

health, agriculture, and the environment. Journal of Heredity 104: 595-600.

Kelley JL, et al. 2014. Compact genome of the Antarctic midge is likely an adaptation to an

extreme environment. Nat Commun 5: 4611. doi: 10.1038/ncomms5611

Kriventseva EV, et al. 2019. OrthoDB v10: sampling the diversity of animal, plant, fungal, protist,

bacterial and viral genomes for evolutionary and functional annotations of orthologs.

Nucleic Acids Res 47: D807-D811.

Lewin HA, et al. 2018. Earth BioGenome Project: Sequencing life for the future of life.

Proceedings of the National Academy of Sciences 115: 4325-4333.

Li F, et al. 2019. Insect genomes: progress and challenges. Insect molecular biology 28: 739-

758.

McGee MD, et al. 2020. The ecological and genomic basis of explosive adaptive radiation.

Nature 586: 75-79.

McKenna DD, et al. 2019. The evolution and genomic basis of beetle diversity. Proceedings of

the National Academy of Sciences 116: 24729-24737.

Misof B, et al. 2014. Phylogenomics resolves the timing and pattern of insect evolution. Science

346: 763-767. doi: 10.1126/science.1257570

Petersen M, et al. 2019. Diversity and evolution of the transposable element repertoire in

arthropods with particular reference to insects. BMC Evol Biol 19: 1-15.

Rhie A, et al. 2021. Towards complete and error-free genome assemblies of all vertebrate

species. Nature 592: 737-746.

Robinson GE, et al. 2011. Creating a buzz about insect genomes. Science 331: 1386.

Sayers EW, et al. 2021. GenBank. Nucleic Acids Res 48: D84-D86.

Seehausen O, et al. 2014. Genomics and the origin of species. Nature Reviews Genetics 15:

176-192.

Seppey M, Manni M, Zdobnov EM. 2019. BUSCO: assessing genome assembly and annotation

completeness. In. Gene prediction: Springer. p. 227-245.

Stork NE 2018. How many species of insects and other terrestrial arthropods are there on Earth?

Annual Review of Entomology 63: 31-45.

Thibaud-Nissen F, et al. 2016. The NCBI Eukaryotic Genome Annotation Pipeline. Journal of

Animal Science 94: 184-184.

Thomas GW, et al. 2020. Gene content evolution in the arthropods. Genome Biology 21: 1-14.

Wang X, et al. 2014. The locust genome provides insight into swarm formation and long-distance

flight. Nat Commun 5: 1-9.

Zhang Z-Q. 2011. Animal biodiversity: An outline of higher-level classification and survey of

taxonomic richness: Magnolia press.