Endothelial cells elicit a pro-inflammatory response to SARS-CoV-2 without productive viral

infection

<u>Running Title: Blood vessels are not infected by SARS-CoV-2.</u>

Lilian Schimmel[1*] Keng Yih Chew[2*] Claudia Stocks[1,3] Teodor Yordanov[1] Patricia Essebier[1]

Arutha Kulasinghe[4,5] James Monkman[4,5] Anna Flavia Ribeiro dos Santos Miggiolaro[6] Caroline

Cooper[7] Lucia de Noronha[8] Anne K. Lagendijk[1] Kate Schroder[1,3] Larisa I. Labzin[1,3#] Emma J.

Gordon[1#] & Kirsty R. Short[2#]

1. Division of Cell and Developmental Biology, Institute for Molecular Bioscience, The

University of Queensland, Brisbane, QLD, Australia

2. School of Chemistry and Molecular Biosciences, The University of Queensland, Brisbane,

QLD, Australia

3. IMB Centre for Inflammation and Disease Research, Institute for Molecular Bioscience, The

University of Queensland, Brisbane, QLD, Australia

4. Queensland University of Technology, School of Biomedical Sciences, Faculty of Health,

Brisbane, Queensland, Australia.

5. Translational Research Institute, Brisbane, Queensland, Australia.

6. Postgraduate Program of Health Sciences, School of Medicine, Hospital Marcelino

Champagnat - Pontifícia Universidade Católica do Paraná (PUCPR), Brazil

7. Pathology Queensland, Princess Alexandra Hospital, Brisbane, Qld, Australia

8. School of Medicine & Center of Education, Research and Innovation – Hospital

Marcelino Champagnat - Pontifícia Universidade Católica do Paraná (PUCPR). Brazil

*/# authors contributed equally

Correspondence:

k.short@uq.edu.au

e.gordon@imb.uq.edu.au

ABSTRACT

<u>Objectives:</u> Thrombotic and microvascular complications are frequently seen in deceased COVID-19

patients. However, whether this is caused by direct viral infection of the endothelium or inflammation-

induced endothelial activation remains highly contentious. <u>Methods:</u> Here, we use patient autopsy

samples, primary human endothelial cells and an *in vitro* model of the pulmonary epithelial-endothelial

cell barrier to show that primary human endothelial cells express very low levels the SARS-CoV-2

receptor ACE2 and the protease TMPRSS2. <u>Results:</u> Accordingly, endothelial cells can only be infected

when SARS-CoV-2 is present at very high concentrations. However, this is not a productive infection

(i.e. no infectious virus is produced) and viral entry induces an inflammatory response. We also show

that SARS-CoV-2 does not infect endothelial cells in 3D vessels under flow conditions. We further

demonstrate that in a co-culture model endothelial cells are not infected with SARS-CoV-2. They do

however sense and respond to infection in the adjacent epithelial cells, increasing ICAM-1 expression

and releasing pro-inflammatory cytokines. <u>Conclusions</u>: Taken together, these data suggest that *in vivo,*

endothelial cells are unlikely to be infected with SARS-CoV-2 and that infection is only likely to occur

if the adjacent pulmonary epithelium is denuded (basolateral infection) or a high viral load is present in

the blood (apical infection). In such a scenario, whilst SARS-CoV-2 infection of the endothelium can

occur, it does not contribute to viral amplification. However, endothelial cells are still likely to play a

key role in SARS-CoV-2 pathogenesis by sensing adjacent infection and mounting a pro-inflammatory

response to SARS-CoV-2.

Keywords: SARS-CoV-2, endothelial cells, COVID-19, inflammation, blood vessels.

INTRODUCTION

SARS-CoV-2 causes diverse clinical syndromes, ranging from asymptomatic infection to fatal disease. Of patients hospitalised with COVID-19 (the clinical manifestation of SARS-CoV-2 infection), approximately 30% of individuals go on to develop severe disease associated with progressive lung damage [1]. This is associated with a breakdown of the vascular barrier, oedema, endotheliitis, thrombosis and inflammatory cell infiltration [1, 2]. Many studies have identified thrombotic and microvascular complications in deceased patients, suggesting that vascular pathology is a major driver of severe disease [3, 4]. Other major pathological events include arterial and venous thromboembolism, kidney disease and neurological disorders [5, 6], suggesting that SARS-CoV-2 activates the vasculature throughout the body, potentially resulting in multi-organ failure. However, whether this is caused by direct viral infection of the endothelium or inflammation-induced endothelial activation remains highly contentious.

SARS-CoV-2 cellular uptake is mediated by binding of the spike glycoprotein to ACE2 (angiotensin-converting enzyme 2) and NRP1 (Neuropilin-1) at the cell surface [7-10]. Host surface proteases such as TMPRSS2 cleave full-length spike protein (S0) at its S2' site. Cleavage at the S2' site can facilitate the fusion of viral and cellular membranes to deliver the viral RNA into the cytosol [1, 11]. Alternative routes of virus uptake have also been reported in cells with low expression of TMPRSS2: this involves endo-lysosomal pathway and cathepsins [12]. Numerous studies suggest that endothelial cells express ACE2 [3, 13-17], Neuropilin receptors [18-22] and TMPRSS2 [23], suggesting that viral infection of endothelial cells is theoretically possible. However, others have suggested that ACE2 is highly expressed in microvascular pericytes [14, 24, 25] and that pericyte injury in response to viral infection may lead to endothelial dysfunction.

Evidence regarding direct infection of the endothelium with SARS-CoV-2 remains equivocal [26, 27]. Autopsy studies on COVID-19 patients have suggested the presence of viral particles in the vascular beds of different organs [3, 28, 29] and others have shown SARS-CoV-2 infection of cultured endothelial cells [30, 31] and human blood vessel organoids [32]. Autopsy studies of deceased COVID-19 patients have

also shown SARS-CoV-2 RNA to be enriched in pulmonary endothelial cells [33]. In contrast, several

studies have failed to find evidence of endothelial infection in deceased COVID-19 patients [27] and called

into question the validity of identifying virally-infected cells based on the presence of the SARS-CoV-

2 spike protein [25, 34, 35]. Animal models of SARS-CoV-2 infection have also failed to identify any

obvious signs of endothelial infection [36, 37], despite clear endothelial dysfunction and thrombosis [38].

When and how endothelial cells may be exposed to SARS-CoV-2 *in vivo* also remains unclear. As a

respiratory virus, SARS-CoV-2 encounters pulmonary epithelial cells prior to any interaction with the

endothelium. SARS-CoV-2 infects pulmonary epithelial cells apically and new viral particles are also

released apically into the lumen of the lung [39]. It is possible that the infected epithelial layer loses barrier

function, allowing endothelial infection from the basolateral side of epithelia that is adjacent to alveolar

endothelium. Alternatively, SARS-CoV-2 may infect endothelial cells apically via the blood, although

viremia appears to be rare in COVID-19 patients [40]. The question therefore remains whether endothelial

dysfunction in COVID-19 is the result of direct viral infection of the endothelium.

Here, using patient autopsy samples and *in vitro* models, we show that whilst SARS-CoV-2 can enter

endothelial cells if high viral titres are present, this infection is not productive. Rather, in response to

direct or indirect viral exposure, endothelial cells mount a pro-inflammatory response characterised by

increased expression of ICAM-1 and secretion of CXCL-10 and IL-6. Taken together, our results

provide clarification on the intensely debated topic of endothelial infection by SARS-CoV-2, to reveal

that the endotheliopathy and thrombocytopathy in patients with severe COVID-19 is likely to be due to

the inflammatory response, rather than direct viral infection.

MATERIALS AND METHODS

In vivo samples: Lung tissue from 10 SARS-CoV-2 infected patients was sectioned and Hematoxylin and Eosin (H&E) staining performed. RNAscope probes (ACDbio, US) targeting SARS-CoV-2 spike mRNA (#848561-C3), were used as per manufacturer instructions for automation on Leica Bond RX. DNA was visualised with Syto13, and SARS-CoV-2 spike probe with opal 690 (1:1500). Fluorescent images were acquired with Nanostring Mars prototype DSP at 20x. Autopsy and biopsy materials were obtained from the Pontificia Universidade Catolica do Parana PUCPR the National Commission for Research Ethics (CONEP) under ethics approval numbers 2020001792/30188020.7.1001.0020 and 2020001934/30822820.8.000.0020. The study was also approved under University of Queensland HREC ratification.

Cell culture: Human umbilical vein endothelial cells (HUVECs), and human microvascular endothelial cells from lungs (HMVEC-L) purchased from Lonza (Cat# CC-2935, and CC-2527 respectively) were cultured until passage 8 in EGM-Plus or EGM-2MV medium, supplemented with singlequots (Lonza Cat# CC-5035, CC-3102). Calu-3 cells purchased from ATCC (Cat# HTB-55) were maintained in MEM (Invitrogen), containing 10% (v/v) heat-inactivated foetal bovine serum (Cytiva), 100 U/ml penicillin and streptomycin (Life Technologies Australia), and grown in EGM-2MV for endothelial co-culture experiments. Baby hamster kidney cells (BHK-21) were a kind gift from Prof. Dr. R.G. Parton. Cell lysates of human nasal epithelial cells (HNEp) were a kind gift from Prof. Peter Sly.

Monocultures of HUVEC, HMVEC or Calu-3 cells were performed on 24 well cell culture inserts (Corning 6.5 mm Transwell, 0.4 µm polycarbonate membrane Cat#3413) coated with 5 µg/ml fibronectin (FN) (Sigma). Approximately 25,000 HUVEC, 25,000 HMVEC and 50,000 Calu-3 cells were seeded per insert on either the apical or basolateral side as indicated and grown in EGM-Plus (Lonza), EGM-2MV (Lonza) or MEM (Invitrogen) respectively. Endothelial cells were treated with 100ng/ml TNFα (Life Technologies Australia) or 10ng/ml IFNß (Lonza) for as long as indicated.

Calu-3 and HMVEC were co-cultured on 24 well cell culture inserts (Corning 6.5 mm Transwell, 0.4

μm polycarbonate membrane Cat#3413) coated with 5 μg/ml FN (Sigma) in EGM-2MV (Lonza).

Approximately 25,000 HMVECs were seeded on an inverted transwell membrane for 1 hr to attach.

Transwell membranes were turned over to normal position and approximately 50,000 Calu-3 cells were

seeded in the top compartment of the membrane. Cells were grown for 48 hr prior to infections.

Viral infection: SARS-CoV-2 isolate hCoV-19/Australia/QLD02/2020 was provided by Queensland

Health Forensic & Scientific Services, Queensland Department of Health. Virus was grown on Vero

cells and titred [41]. Sanger sequencing was used to confirm that no mutations occurred in the spike gene

relative to the original clinical isolate. Cells were infected with 6×10^4 plaque forming units (PFU) for

1 hour at 37 degrees 5% Co2. The viral inoculum was then removed, and the medium was replaced with

DMEM (Invitrogen) or MEM (Invitrogen) containing 2% FBS. Alternatively, 2×10^6 PFUs were added

to the cell culture and cells were incubated at 37 degrees 5% CO_2. All studies with SARS-CoV-2 were

performed under physical containment 3 (PC3) conditions and were approved by the University of

Queensland Biosafety Committee (IBC/374B/SCMB/2020).

Antibodies: The following antibodies were used: mouse anti-ACE2 (Santa Cruz, sc-390851, IF1:200)

goat anti-ACE2 (R&D systems, AF933, WB 1:1000), mouse anti-dsRNA (Millipore, MABE1134,

IF1:100), rabbit anti-SARS-CoV nucleocapsid protein/NP (Sino Biological, 40143-R040, WB1:1000

IF1:200), rabbit anti-TMPRSS2 (Abcam, ab92323, WB1:1000), rabbit anti-cleaved caspase 3 (Cell

Signaling, 9661 IF1:30 after pre-labeling), rabbit anti-GAPDH (Cell Signaling, 2118, WB1:2000),

mouse anti-ICAM-1 (R&D systems, BBA3, IF1:200), mouse anti-ICAM-1 (Santa Cruz Biotechnology,

sc-8439, WB1:1000), Phalloidin-Alexa555 (Cytoskeleton, PHDH1-A, IF1:500), Phalloidin-Alexa670

(Cytoskeleton, PHDN1-A, IF1:200). Pre-labeling of rabbit anti-cleaved caspase 3 with Alexa Fluor 555

was performed using Zenon Rabbit IgG labelling kit (Z-25305) according to manufacturer protocol.

Immunofluorescence Staining: Immunofluorescence staining was in general performed on cells

cultured on 12 mm glass coverslips or on 24 well cell culture inserts (Corning 6.5 mm Transwell, 0.4

μm polycarbonate membrane Cat#3413) coated with 5 μg/ml FN (Sigma), washed with $PBS^{+/+}$ (PBS

supplemented with 1 mM $CaCl_2$ and 0.5 mM $MgCl_2$), fixed for 10 min in 4% PFA (Sigma), blocked

and permeabilized for 30 min with 3% BSA, 0.3% Triton X-100 (Sigma). Primary antibodies were

incubated in 1.5% BSA for 1 hr at RT, secondary antibodies linked to Alexa fluorophores (all

Invitrogen) were incubated in 1.5% BSA for 1 hr at RT and incubated with 4% PFA (Sigma) for another

24 hr in case of SARS-CoV-2 infected cells, after each step samples were washed 3x with $PBS^{+/+}$ and

mounted in ProlongGold + DAPI solution (Cell Signaling Technologies, CST 8961S). Z-stack image

acquisition was performed on a confocal laser scanning microscope (Zeiss LSM880) using a 40x NA

1.3 or 63x NA 1.4 oil immersion objective.

3D Microfluidic cell culture: Microfluidic devices were generated as previously described [42] using

2.5mg/ml Collagen (R&D Systems) for the ECM. Cells were seeded 48 hr prior to virus infection and

were kept on a rocker at 37C° and 5%CO_2. The tubes were fixed sequentially in 1% formaldehyde,

containing 0.05% Triton for 90 sec, followed by 4% formaldehyde for 15 min at 37C°. Tubes were

permeabilised with 0.5% Triton for 10 min at 37C° and blocked for 4 hr in 10% goat serum in PBS at

4C° prior to staining. Immunofluorescence staining was performed using NP, dsRNA and Phalloidin-

Alexa670 antibodies. Z-stack image acquisition was performed on a confocal laser scanning microscope

(Zeiss Axiovert 200 inverted microscope with LSM 710 Meta Confocal Scanner) using 40X NA 1.1

water immersion objective.

Western blotting: For total cell lysates, cells were washed once with PBS and lysed with RIPA buffer

(50 mM Tris, 150 mM NaCl, 1 mM EDTA, 1% Triton X-100, 0.1% SDS, 1% sodium deoxycholate,

protease inhibitor, pH 8.0). Pierce BCA protein assay kit (Thermo Scientific) was used to equalize

protein amounts and SDS-sample buffer containing 100 mM DTT (Astral Scientific) was added and

samples were boiled at 95°C for 10 minutes to denature proteins. Proteins were separated on 4-15%

mini protean TGX precast gels (Biorad) in running buffer (200 mM Glycine, 25 mM Tris, 0.1% SDS

(pH8.6)), transferred to nitrocellulose membrane (BioRad Cat#1620112) in blot buffer (48 nM Tris, 39

nM Glycine, 0.04% SDS, 20% MeOH) and subsequently blocked with 5% (w/v) BSA in Tris-buffered

saline with Tween 20 (TBST) for 30 minutes. The immunoblots were analysed using primary antibodies

incubated overnight at 4°C and secondary antibodies linked to horseradish peroxidase (HRP)

(Invitrogen), and after each step immunoblots were washed 4x with TBST. HRP signals were visualized

by enhanced chemiluminescence (ECL) (BioRad) and imaged with a Chemidoc (BioRad).

RNA extraction and cDNA synthesis: RNA was extracted using the Qiagen RNeasy Minikit (Qiagen)

or NucleoZOL (BIOKÉ, The Netherlands) according to manufacturer's guidelines. cDNA was

synthesized using random hexamers and the high-capacity cDNA reverse transcription kit (Life

Technologies) according to the manufacturer's guidelines.

Quantitative PCR (qPCR): qPCR was performed using SYBR Green reagent (Applied Biosystems).

All primer sequences are listed in Table 1. qPCR conditions were applied according to the

manufacturer's instructions using QuantStudio 6 Flex Real- Time PCR System (ThermoFisher

Scientific, Waltham, Massachusetts, MA). HPRT or Glyceraldehyde 3-phosphate dehydrogenase

(GAPDH) was used as a housekeeping gene and relative gene expression was determined using the

comparative cycle threshold ($\Delta\Delta$Ct) method (Livak, 2001).

Table 1: Primers used in the present study

NAME	SEQUENCE (5'- 3')
SARS-COV2 (MPRO) VIRUS FW	GAGACAGGTGGTTTCTCAATCG
SARS-COV2 (MPRO) VIRUS RV	ACGGCAATTCCAGTTTGAGC
GAPDH FW	CGAGATCCCTCCAAAATCAA
GAPDH RV	TTCACACCCATGACGAACAT
HPRT FW	TCAGGCAGTATAATCCAAAGATGGT
HPRT RV	AGTCTGGCTTATATCCAACACTTCG
ACE2 FW	TCACGATTGTTGGGACTCTGC
ACE2 RV	TCGCTTCATCTCCCACCACT

NEUROPILLIN FW	TGGACCGACCACTCCCAACG
NEUROPILLIN RV	AGAGCCCCAGCCAAATTCACAG
TMPRSS2 FW	GGAAGTTCATGGGCAGCAAG
TMPRSS2 RV	AGGCGAACACACCGATTCTC

Viral titres: Viral titres in cell culture supernatant determined by a plaque assay on Vero cells, according

to previously described protocols [41].

Cytokine titres: Cytokine titres in were determined using an Alphalisa kit (Perkin Elmer) according to

the manufacturer's instructions. The sum of cytokines detected in the upper and lower compartments of

the transwell is shown for each of the monocultured samples, while cytokine levels were kept separated

for the upper and lower compartment for the co-cultured samples.

Image analysis: Analysis of immunofluorescent images was performed using ImageJ version 2.0.0-rc-

69/1.52n. A mask of the total cell area was created using default intensity threshold on Phalloidin

staining. Within the mask, Raw Integrated Density of either ACE2, NP, ICAM-1 was measured, and

value was corrected for total mask area resulting in RawIntDen/Area. For dsRNA, MaxEntropy

threshold was performed on dsRNA staining to remove background speckles before measuring Raw

Integrated Density within the mask. Analysis of Western blot images was performed using Gel Analyzer

Tool of ImageJ version 2.0.0-rc-69/1.52n. The background signal was subtracted and values were

normalized to corresponding GAPDH loading control. Imaris (Bitplane) version 8 was used to create

the XZ projection in Figure 4B''. Extrusion of cells was quantified in Supplemental Figure 1C using

orthogonal views in ImageJ version 2.0.0-rc-69/1.52n.

Statistical analysis: All statistical analysis was performed using Graphpad Prism version 9.0. Data are

presented as mean±s.e.m. with individual data points indicated and colour coded per independent

experimental replicate. Statistical significance was determined using Kruskal Wallis test for multiple

comparisons or Mann-Whitney test between mock and SARS-CoV-2 treated conditions. *P<0.05,

**P<0.01, ***P<0.001 and ****P<0.0001.

RESULTS

Endothelial cell infection is not readily detected in deceased COVID-19 patients

To determine the prevalence of *in vivo* SARS-CoV-2 endothelial cell infection, autopsy lung sections

were obtained from 10 deceased COVID-19 patients and probed for SARS-CoV-2 spike mRNA using

RNAscope. Two of the 10 patients were positive for SARS-CoV-2 RNA in the lungs. In these

individuals, spike mRNA could not be detected in pulmonary endothelial cells (endothelial cells defined

by H & E staining (Figure 1). These data suggest that the endothelium is not generally the primary site

of viral replication *in vivo* in COVID-19 patients (patient data available in Supplemental Table 1).

Primary endothelial cells express low levels of ACE2 and TMPRSS2

To further investigate the contribution of endothelial cells to the pathogenesis of SARS-CoV-2, we

established *in vitro* cultures of primary human umbilical vein endothelial cells (HUVECs) and human

microvascular endothelial cells from the lung (HMVEC-L). Multiple studies have reported that

endothelial cells across different vascular beds express ACE2 and TMPRSS2, the host cell receptor and

protease that are required for efficient cell infection by SARS-CoV-2 [10, 11]. We found that both HUVECs

and HMVEC-Ls expressed comparable ACE2 and TMPRSS-2 mRNA to that observed in immortalised

epithelial cells known to be susceptible to SARS-CoV-2 infection (Calu-3) (Figure 2A). Neuropilin-1,

which enhances viral entry through its interaction with the spike multibasic cleavage site [7, 8], was readily

detected in endothelial cells by qPCR (Figure 2A) in line with established data [19]. The expression of

ACE2 protein detected by immunofluorescence (Figure 2B) was significantly reduced compared to

epithelial cells (Calu-3) (Figure 2C). We also detected ACE2 and TMPRSS2 protein in HUVECs and

HMVEC-Ls by western blot (Figure 2D-E). The specificity of our ACE2 antibody was confirmed by

western blot of BHK-21 cells (which are known to be negative for ACE2) [43].These data suggest that

while vascular endothelial cells can theoretically be infected by SARS-CoV-2, infection is likely to be

inefficient compared to epithelial cells, because of weak receptor and protease expression in endothelial

cells.

Endothelial cells are not productively infected by SARS-CoV-2

SARS-CoV-2 may theoretically enter endothelial cells via the apical surface (should the virus enter the

blood stream) or the basolateral surface (should the virus be present at the basolateral surface of the

adjacent epithelial cells). We therefore assessed whether endothelial cells were susceptible to SARS-

CoV-2 infection when exposed either apically (Supplemental Figure 1A) or basolaterally (Supplemental

Figure 1B) to 6 x 10^4 PFUs. Infectious virus was not detected in the supernatant of either HUVECs or

HMVEC-Ls at 24, 48 nor 72 hours post-infection (Supplemental Figure 1C). In contrast, Calu-3 cells

demonstrated robust viral replication and release (Supplemental Figure 1C). Similarly, no increase in

SARS-CoV-2 nucleocapsid protein within infected endothelial cells relative to mock was detected by

immunofluorescence (Supplemental Figure 1D-G) or western blot (Supplemental Figure 1H-I), and no

detectable alterations in endothelial cell morphology were evident after viral exposure, as assessed by

phalloidin immunostaining (Supplemental Figure 1D, F). In contrast, Calu-3 cells displayed a

significant increase in nucleocapsid protein expression after viral exposure. Together, these data suggest

that endothelial cells *in vitro* are not efficiently infected with SARS-CoV-2.

We next sought to establish if the low levels of ACE2 and TMPRSS2 on endothelial cells could be

overcome by a higher viral titre, to mediate infection. Accordingly, endothelial cells were infected either

apically or basolaterally with 2 x 10^6 PFUs of SARS-CoV-2 and viral replication was again assessed

over time. Infectious virus did not increase over time in endothelial cells, rather viral titres decreased

over time, consistent with a degradation of input virus (Figure 3A). In contrast, productive viral

replication was observed in Calu-3 cells over time (Figure 3A, Supplemental Figure 2A), while, as

expected, non-susceptible, ACE2-negative BHK-21 cells showed no productive viral replication

(Supplemental Figure 2A). To determine if endothelial cells were susceptible to viral entry, if not

replication, we assessed viral nucleocapsid protein by immunofluorescence in HMVEC-L (Figure 3B-

C) positive control Calu-3 and negative control BHK-21 (Supplemental Figure 2D-E) and western blot

(Figure 3E-F, Supplemental Figure 2B-C). Viral nucleocapsid protein expression was significantly

increased after both apical and basolateral SARS-CoV-2 infection, revealing that SARS-CoV2 can

indeed enter endothelial cells *in vitro*. We observed that the majority of infected, nucleocapsid protein-

positive endothelial cells appeared to be extruding from the cell monolayer in an apical manner (Figure

3B', 3B'', 3C). These results were not due to dead cells present in the viral inoculum as BHK-21 cells

did not show any nucleocapsid-positive extruded cells (Supplemental Figure 2D). Moreover, extruding

endothelial cells were positive for cleaved caspase 3, indicating that SARS-CoV-2 entry into endothelial

cells induces apoptosis (Figure 3G-H). These results demonstrate that while SARS-CoV-2 can enter

endothelial cells this infection is not productive and leads to apoptosis and extrusion of these apoptotic

cells from the endothelial monolayer.

Endothelial cells are not productively infected by SARS-CoV-2 in 3D vessels

To determine whether endothelial cells can be infected in 3D vessels under flow, we cultured both

HMVEC-L and Vero cells in microfabricated tubes [42] and added SARS-CoV2 to the luminal surface

(Figure 4A). After 24h, Vero cells were readily infected and viral replication was observed, in contrast

to HMVEC-L which shows viral input levels (Figure 4B). In agreement, NP was not detected in

HMVEC-L tubes, whereas Vero cells displayed clear NP staining at the luminal surface (Figure 4C-F).

This data reveals that fluid flow over endothelial cells does not alter the ability of SARS-CoV-2 to

infect cells.

Endothelial cells mount a modest immune response to SARS-CoV-2

As severe COVID-19 disease is associated with an elevated cytokine response [44], we sought to assess

whether the abortive SARS-CoV-2 infection observed in endothelial cells induces a pro-inflammatory

response. We assessed expression of the leukocyte adhesion molecule, ICAM-1, which is well

established to be induced under inflammatory conditions in the endothelium [45]. Immunofluorescence

showed that HMVEC-L exposed both basally and apically to SARS-CoV-2 significantly increased

ICAM-1 expression, except for in nucleocapsid protein-positive HMVEC-L (likely as they are apoptotic

and extruded from the monolayer) (Figure 5A-C). The moderate basal ICAM-1 expression observed in

mock infected cells is likely due to the fact that the cells are kept for 72h without a media change (as

seen in PBS controls (Figure 5B). However, this response was not as robust as when HMVEC-L were

exposed to the established inflammatory cytokine TNFα (Figure 5B-C). Western blot analysis

confirmed that HMVEC-L exposed to SARS-CoV-2 displayed a trending increase in ICAM-1

expression (Figure 5D, E). In addition to upregulating inflammatory adhesion molecules, we analysed

whether endothelial cells can be the source of pro-inflammatory cytokines during SARS-CoV-2

infection. As IL-6 and CXCL10 are elevated in patients with severe COVID-19 [46], we analysed whether

endothelial cells released these pro-inflammatory cytokines when infected either apically or

basolaterally with SARS-CoV-2 for 72h. HMVEC-Ls released modest levels of CXCL10 (Figure 5F)

and IL-6 (Figure 5G). While these modest levels of cytokine release from endothelial cells upon SARS-

CoV-2 infection are lower than CXCL10 or IL-6 induced in response to high concentrations of

recombinant TNF or IFNß, (Supplemental Figure 2F-G), this shows that endothelial cells express high

basal levels of IL-6 and that both cytokines are increased during infection. Taken together, these results

show that endothelial cells do not support a productive SARS-CoV-2 infection but still mount a modest

pro-inflammatory response to the virus.

Infection of epithelial-endothelial co-cultures elicits an immune response in endothelial cells

To create a more physiologically relevant *in vitro* model of the pulmonary endothelium, we adapted our

previously described co-culture model of the lung epithelial-endothelial barrier [47], where epithelial

(Calu-3) cells are seeded on top of a transwell membrane while endothelial cells (HMVEC-L) are

seeded on the basolateral side (Figure 6C). To mirror a respiratory infection, SARS-CoV-2 was added

to the upper compartment and both epithelial and endothelial cells were analysed at 72h post-infection.

In contrast to our observations in endothelial cell monocultures, no viral nucleocapsid protein or viral

dsRNA was detected in HMVEC-L, whereas SARS-CoV-2 nucleocapsid protein and viral dsRNA

(indicating actively replicating virus) was clearly detected in infected Calu-3 cells (Figure 6A,B; 6E,F).

These results were confirmed by the presence of higher levels of viral RNA as measured by qPCR in

the Calu-3 compared to the HMVEC-L (Fig 6G) and higher titres of infectious virions in the

supernatants of the upper (Calu-3) compartment of the transwell (Figure 6H). Despite lack of

endothelial infection, we observed that HMVEC-L cells expressed ICAM-1 expression when virus was

added to the upper (Calu-3) compartment, suggesting that the HMVEC-L respond to adjacent epithelial

infection (Figure 6A, D). These HMVEC-L also responded to the infection in the adjacent Calu-3 cells

by increasing CXCL10 secretion (detected in lower compartment of the co-culture) (Figure 6I). Both

Calu-3 and HMVEC-L appeared to secrete IL-6 in response to SARS-CoV-2 infection, as IL-6 levels

were increased in both compartments (significantly in the upper, trending in the lower) (Figure 6J). In

summary, in a co-culture model of the pulmonary epithelium, endothelial cells are not directly infected

with the virus, but respond to adjacent epithelial infection by upregulating ICAM-1 expression and

releasing pro-inflammatory cytokines. This suggests that endothelial inflammation and dysfunction

during *in vivo* SARS-CoV-2 infection is most likely to occur in response to infection of the adjacent

epithelial layer.

DISCUSSION

SARS-CoV-2 can induce vascular dysfunction and thrombotic events in patients with severe COVID-

19. However, the cellular and molecular mechanisms by which SARS-CoV-2 infection triggers such

pathologies have so far remained elusive. Here, we have shown that endothelial cells do not support

productive replication of SARS-CoV-2, but that they induce pro-inflammatory cytokines and adhesion

molecules upon either direct or indirect exposure to SARS-CoV-2 virions. This supports the hypothesis

that the endothelial dysfunction observed in COVID-19 patients is likely to be largely mediated by

inflammatory signalling pathways.

As COVID-19 case numbers rose, reports of COVID-19 coagulopathy and vascular dysfunction also

began to accumulate. Whether SARS-CoV-2 directly infects the vasculature to drive this vascular

pathology has thus remained a topic of intense debate. Multiple studies show viral particles surrounding

the vasculature [3, 25, 43], yet it is unclear whether this infection is truly specific to the endothelium or

occurs within the perivascular compartment. Our *in vivo* analyses of lung tissue samples from deceased

COVID-19 patients indicates the endothelium remains uninfected. Nevertheless, we showed that *in*

vitro, endothelial cells are theoretically susceptible to SARS-CoV-2 infection, as they express ACE2,

TMPRSS2 and NRP1, albeit at significantly lower levels than epithelial cells. This is in line with other

studies that showed expression of SARS-CoV-2 receptors by the endothelium in human tissue or in

cultured cells [14, 17, 23, 31]. However, we did not detect endothelial cell infection following exposure to 6

x 10^4 PFUs of SARS-CoV-2, whereas epithelial cells were readily infected at the same viral dose. These

results support those of Wang et al. [31], who showed that epithelial cells are more susceptible to infection

than endothelial cells. When exposed to higher viral titres, endothelial cells were positive for viral NP

protein, suggesting effective viral entry. However, no infectious virions were detected in the endothelial

cell supernatant, indicating that this infection was abortive. It is interesting to speculate as to why

SARS-CoV-2 infection is productive in a monoculture of epithelial, but not endothelial cells. This may

relate to cell-dependent differences in viral entry. For example, in epithelial cells, there may be

sufficient surface TMPRSS2 levels available to cleave the spike protein at the S2' site and mediate

viral-host cell membrane fusion. This may in turn allow for efficient release of viral RNA into the

cytoplasm and thus viral replication. In contrast, in cells with lower TMPRSS2 expression (such as

endothelial cells), SARS-CoV-2 may enter the cell via endocytosis, with S2' cleavage being mediated

by endosomal cathepsins [12]. Endosomal entry may not only be inefficient, but it may also activate the

cellular anti-viral response and thereby limit productive viral replication [48, 49]. Alternatively, a host of

other cellular co-factors expressed preferentially in epithelial cells may support selective SARS-CoV-

2 replication, or endothelial-specific restriction factors may limit effective viral replication.

In contrast to endothelial cells in a monoculture, viral proteins could not be detected in endothelial cells

grown in a co-culture with SARS-CoV-2-positive epithelial cells, despite being exposed to the same

dose of virus. These data may reflect the fact that SARS-CoV-2 typically infects and buds from

epithelial cells in an apical manner, resulting in limited exposure of the endothelium to infectious

virions. Extrapolating these data to the *in vivo* situation would suggest that a basolateral (abortive)

infection of the endothelium by SARS-CoV-2 is only likely to occur when the epithelial barrier of the

lung is severely damaged, exposing the endothelium to incoming virions.

While endothelial cells (either in a monoculture or in a co-culture) did not produce infectious virions,

they did respond to SARS-CoV-2 infection. Analysis of infected endothelial monolayers revealed that

cells positive for viral proteins were extruded in an apical fashion, without any apparent disruption to

the monolayer, as assessed by phalloidin expression. This is in contrast to work from Buzhdygan and

colleagues, who demonstrated that purified SARS-CoV-2 spike protein can induce the breakdown of

the vascular barrier in brain microvascular endothelial cells [50]. Discrepancies between these results may

be due to different sources of primary endothelial cells (lung versus brain), or the differences in response

of the endothelium to a purified viral protein vs infectious virus [1, 3, 24, 29, 51-53]. However, if infected cells

are promptly removed from exposed vessels as our data suggests, then this may explain why detection

of viral particles in the endothelium of patients has varied between studies [4, 54].

In the absence of direct endothelial infection, hyperinflammation is likely to contribute to endothelial

dysfunction observed in COVID-19 patients. SARS-CoV-2 triggered an upregulation of endothelial

ICAM-1 expression when HMVEC-Ls were directly exposed to virus, or co-cultured with infected

epithelial cells. ICAM-1 enables immune cells to effectively extravasate into tissues, so its upregulation

is in keeping with the observed influx of inflammatory monocytes, neutrophils and other immune cells

into the lungs in severe COVID-19 patients [55]. This suggests that locally produced pro-inflammatory

cytokines stimulate the endothelium to induce adhesion marker expression. Our previous studies with

influenza virus showed that endothelial cells can themselves be a source of these pro-inflammatory

cytokines [56], and indeed we also observed SARS-CoV-2-induced release of IL-6 and CXCL10 in co-

cultures. Precisely how SARS-CoV-2 is sensed by the endothelium in either the monoculture or co-

culture system remains to be determined. Given that SARS-CoV-2 does not replicate in endothelial

cells, we speculate that endothelial cells are unlikely to sense this virus through cytosolic RNA receptors

such as RIG-I or MDA5, as these immune sensors would detect viral replication intermediates. A more

likely scenario is that the endothelial cells respond to danger signals from neighbouring infected

epithelial or endothelial cells. These data add to a growing body of literature that indicate that

endothelial cell-directed inflammation plays a key role in the pathogenesis of SARS-CoV-2 [57]. It is

tempting to speculate that, similar to influenza virus, the SARS-CoV-2-induced pro-inflammatory state

may induce endothelial cells to express tissue factor, which in turn induces a pro-coagulant state,

microvascular leakage and pulmonary haemorrhage [58], all of which have been described in patients with

severe COVID-19. Furthermore, regardless of direct or indirect infection, elevated IL-6 correlates with

increased fibrinogen [59] and although still controversial, elevated fibrinogen levels have been detected

in critically ill patients [2].

The present study was subject to several limitations. Firstly, coagulation, thrombosis and induction of

angiogenesis in the lungs of deceased COVID-19 patients has been described in addition to endothelial

dysfunction [3]. This angiogenic response is thought to be predominantly mediated through

intussusceptive angiogenesis, where a new blood vessel is formed by splitting of an existing vessel.

This effect appears to be specific to COVID-19 patients, as lungs from decreased influenza patients do

not display increased angiogenic features. Whether this is due to relative hypoxia in the lungs remains

unclear, although elevated angiogenic growth factors such as VEGF-A and VEGF-C have been

associated with COVID-19 [3, 44]. In this study we have not addressed the effect of SARS-CoV-2 exposure

on the angiogenic capacity of endothelial cells in culture. Given the intimate association between

inflammation, endothelial dysfunction and angiogenesis [45, 60], these are critical aspects of COVID-19

pathology that remain to be addressed in subsequent studies. Similarly, due to the very low levels of

ACE2 present on endothelial cells it was not possible to determine if this was more pronounced on the

apical or basolateral surface of the endothelial cell.

An additional limitation of the present study is that, due to the reductionist nature of the *in vitro* systems

used herein, we were unable to address the contribution of immune cells to endothelial cell dysfunction

during SARS-CoV-2 infection. Multiple studies attribute the sustained inflammatory response directly

to immune cells, wherein macrophage activation, monocyte NLRP3 inflammasome signalling,

complement activation and extrusion of neutrophil extracellular traps are implicated [46, 61-63].

Accordingly, our data may best reflect the initial stages of infection when only a limited number of

tissue-resident leukocytes may be present. Future studies will necessitate the addition of leukocytes to

our co-culture system, to determine whether their presence results in a further induction of inflammation

and markers of severe disease.

Here, we have conclusively shown that *in vitro,* endothelial cells are not productively infected by SARS-

CoV-2 but that they mount an inflammatory response after direct or indirect exposure to the virus,

characterised by increased cytokine secretion and expression of adhesion molecules. Our results thus

suggest a key role of the endothelium in the pathogenesis of COVID-19, and that targeting the

inflammatory response may present the best opportunity to prevent endothelial dysfunction.

ACKNOWLEDGEMENTS AND FUNDING STATEMENT

We thank Dr. Fernando Guimaraes for assisting with acquiring samples necessary for this study. This

work is supported by the National Health and Medical Research Council of Australia (Fellowship

1141131 to KS, Fellowship 1124162 to LL, Project Grant APP1158002 to EG) and the Australian

Research Council (Fellowship DE180100512 to KRS, Fellowship DE170100167 to EG, Discovery

Project DP200100737 to EG and AL), The National Heart Foundation of Australia (Future Leader

Fellowship 104692 to EG) and UQ Early Career Researcher Grants (UQECR2058733 to LS,

UQECR2058045 to LL). Imaging was performed in IMB Microscopy incorporating the Cancer

Ultrastructure and Function Facility funded by the Australian Cancer Research Foundation.

CONFLICT STATEMENT

KS is a co-inventor on patent applications for NLRP3 inhibitors which have been licensed to Inflazome

Ltd, a company headquartered in Dublin, Ireland. Inflazome is developing drugs that target the NLRP3

inflammasome to address unmet clinical needs in inflammatory disease. KS served on the Scientific

Advisory Board of Inflazome in 2016–2017, and serves as a consultant to Quench Bio, USA and

Novartis, Switzerland.

DATA AVAILABILITY STATEMENT

All raw data is available upon request.

REFERENCES

1.   Teuwen LA, Geldhof V, Pasut A, Carmeliet P. COVID-19: the vasculature unleashed. *Nat Rev*

Immunol 2020; **20**: 389-391.

2.   Gu SX, Tyagi T, Jain K*, et al.* Thrombocytopathy and endotheliopathy: crucial contributors to

COVID-19 thromboinflammation. *Nature Reviews Cardiology* 2020.

3.   Ackermann M, Verleden SE, Kuehnel M*, et al.* Pulmonary Vascular Endothelialitis,

Thrombosis, and Angiogenesis in Covid-19. *N Engl J Med* 2020; **383**: 120-128.

4.   Goshua G, Pine AB, Meizlish ML*, et al.* Endotheliopathy in COVID-19-associated

coagulopathy: evidence from a single-centre, cross-sectional study. *Lancet Haematol* 2020; **7**:

e575-e582.

5.   Sardu C, Gambardella J, Morelli MB, Wang X, Marfella R, Santulli G. Hypertension,

Thrombosis, Kidney Failure, and Diabetes: Is COVID-19 an Endothelial Disease? A

Comprehensive Evaluation of Clinical and Basic Evidence. *J Clin Med* 2020; **9**.

6.   Gupta A, Madhavan MV, Sehgal K*, et al.* Extrapulmonary manifestations of COVID-19.

Nature medicine 2020; **26**: 1017-1032.

7.   Cantuti-Castelvetri L, Ojha R, Pedro LD*, et al.* Neuropilin-1 facilitates SARS-CoV-2 cell entry

and infectivity. *Science* 2020.

8.   Daly JL, Simonetti B, Klein K*, et al.* Neuropilin-1 is a host factor for SARS-CoV-2 infection.

Science 2020.

9.   Zhou P, Yang X-L, Wang X-G*, et al.* A pneumonia outbreak associated with a new coronavirus

of probable bat origin. *Nature* 2020; **579**: 270-273.

10.   Matheson NJ, Lehner PJ. How does SARS-CoV-2 cause COVID-19? *Science* 2020; **369**: 510-

511.

11.   Hoffmann M, Kleine-Weber H, Schroeder S*, et al.* SARS-CoV-2 Cell Entry Depends on ACE2

and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 2020; **181**: 271-

280.e278.

12.   Laporte M, Stevaert A, Raeymaekers V*, et al.* The SARS-CoV-2 and other human coronavirus

509   spike proteins are fine-tuned towards temperature and proteases of the human airways. *bioRxiv*

510   2020: 2020.2011.2009.374603.

511   13.   Ferrario CM, Jessup J, Chappell MC*, et al.* Effect of angiotensin-converting enzyme inhibition

512   and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation*

513   2005; **111**: 2605-2610.

514   14.   Nicin L, Abplanalp WT, Mellentin H*, et al.* Cell type-specific expression of the putative SARS-

515   CoV-2 receptor ACE2 in human hearts. *Eur Heart J* 2020; **41**: 1804-1806.

516   15.   Sluimer JC, Gasc JM, Hamming I*, et al.* Angiotensin-converting enzyme 2 (ACE2) expression

517   and activity in human carotid atherosclerotic lesions. *J Pathol* 2008; **215**: 273-279.

518   16.   Lu R, Zhao X, Li J*, et al.* Genomic characterisation and epidemiology of 2019 novel

519   coronavirus: implications for virus origins and receptor binding. *Lancet* 2020; **395**: 565-574.

520   17.   Hikmet F, Méar L, Edvinsson Å, Micke P, Uhlén M, Lindskog C. The protein expression profile

521   of ACE2 in human tissues. *Molecular Systems Biology* 2020; **16**: e9610.

522   18.   Soker S, Takashima S, Miao HQ, Neufeld G, Klagsbrun M. Neuropilin-1 is expressed by

523   endothelial and tumor cells as an isoform-specific receptor for vascular endothelial growth

524   factor. *Cell* 1998; **92**: 735-745.

525   19.   Kitsukawa T, Shimono A, Kawakami A, Kondoh H, Fujisawa H. Overexpression of a

526   membrane protein, neuropilin, in chimeric mice causes anomalies in the cardiovascular system,

527   nervous system and limbs. *Development* 1995; **121**: 4309-4318.

528   20.   Gerhardt H, Ruhrberg C, Abramsson A, Fujisawa H, Shima D, Betsholtz C. Neuropilin-1 is

529   required for endothelial tip cell guidance in the developing central nervous system.

530   *Developmental dynamics : an official publication of the American Association of Anatomists*

531   2004; **231**: 503-509.

532   21.   Jones EA, Yuan L, Breant C, Watts RJ, Eichmann A. Separating genetic and hemodynamic

533   defects in neuropilin 1 knockout embryos. *Development* 2008; **135**: 2479-2488.

534   22.   Aspalter IM, Gordon E, Dubrac A*, et al.* Alk1 and Alk5 inhibition by Nrp1 controls vascular

535   sprouting downstream of Notch. *Nature communications* 2015; **6**.

536   23.   Aimes RT, Zijlstra A, Hooper JD, *et al.* Endothelial cell serine proteases expressed during

537   vascular morphogenesis and angiogenesis. *Thromb Haemost* 2003; **89**: 561-572.

538   24.   Chen L, Li X, Chen M, Feng Y, Xiong C. The ACE2 expression in human heart indicates new

539   potential mechanism of heart injury among patients infected with SARS-CoV-2.

540   *Cardiovascular research* 2020; **116**: 1097-1100.

541   25.   McCracken IR, Saginc G, He L, *et al.* Lack of Evidence of Angiotensin-Converting Enzyme 2

542   Expression and Replicative Infection by SARS-CoV-2 in Human Endothelial Cells. *Circulation*

543   2021; **143**: 865-868.

544   26.   Fosse JH, Haraldsen G, Falk K, Edelmann R. Endothelial Cells in Emerging Viral Infections.

545   *Frontiers in Cardiovascular Medicine* (Review).2021; **8**.

546   27.   Bryce C, Grimes Z, Pujadas E, *et al.* Pathophysiology of SARS-CoV-2: targeting of endothelial

547   cells renders a complex disease with thrombotic microangiopathy and aberrant immune

548   response. The Mount Sinai COVID-19 autopsy experience. *medRxiv* 2020:

549   2020.2005.2018.20099960.

550   28.   Schaefer I-M, Padera RF, Solomon IH, *et al.* In situ detection of SARS-CoV-2 in lungs and

551   airways of patients with COVID-19. *Modern Pathology* 2020; **33**: 2104-2114.

552   29.   Varga Z, Flammer AJ, Steiger P, *et al.* Endothelial cell infection and endotheliitis in COVID-

553   19. *Lancet* 2020; **395**: 1417-1418.

554   30.   Amraie R, Napoleon MA, Yin W, *et al.* CD209L/L-SIGN and CD209/DC-SIGN act as

555   receptors for SARS-CoV-2 and are differentially expressed in lung and kidney epithelial and

556   endothelial cells. *bioRxiv* 2020: 2020.2006.2022.165803.

557   31.   Wang P, Luo R, Zhang M, *et al.* A cross-talk between epithelium and endothelium mediates

558   human alveolar–capillary injury during SARS-CoV-2 infection. *Cell Death & Disease* 2020;

559   **11**: 1042.

560   32.   Monteil V, Kwon H, Prado P, *et al.* Inhibition of SARS-CoV-2 Infections in Engineered Human

561   Tissues Using Clinical-Grade Soluble Human ACE2. *Cell* 2020; **181**: 905-913.e907.

562   33.   Delorey TM, Ziegler CGK, Heimberg G, *et al.* A single-cell and spatial atlas of autopsy tissues

563   reveals pathology and cellular targets of SARS-CoV-2. *bioRxiv* 2021: 2021.2002.2025.430130.

34.   Ko CJ, Harigopal M, Gehlhausen JR, Bosenberg M, McNiff JM, Damsky W. Discordant anti-SARS-CoV-2 spike protein and RNA staining in cutaneous perniotic lesions suggests endothelial deposition of cleaved spike protein. *Journal of Cutaneous Pathology* **n/a**.

35.   Baeck M, Hoton D, Marot L, Herman A. Chilblains and COVID-19: why SARS-CoV-2 endothelial infection is questioned. *British Journal of Dermatology* **n/a**.

36.   Sia SF, Yan L-M, Chin AWH*, et al.* Pathogenesis and transmission of SARS-CoV-2 in golden hamsters. *Nature* 2020; **583**: 834-838.

37.   Rockx B, Kuiken T, Herfst S*, et al.* Comparative pathogenesis of COVID-19, MERS, and SARS in a nonhuman primate model. *Science* 2020; **368**: 1012-1015.

38.   Aid M, Busman-Sahay K, Vidal SJ*, et al.* Vascular Disease and Thrombosis in SARS-CoV-2-Infected Rhesus Macaques. *Cell* 2020; **183**: 1354-1366.e1313.

39.   Milewska A, Kula-Pacurar A, Wadas J*, et al.* Replication of Severe Acute Respiratory Syndrome Coronavirus 2 in Human Respiratory Epithelium. *Journal of Virology* 2020; **94**: e00957-00920.

40.   Andersson MI, Arancibia-Carcamo CV, Auckland K*, et al.* SARS-CoV-2 RNA detected in blood products from patients with COVID-19 is not associated with infectious virus. *Wellcome Open Res* 2020; **5**: 181.

41.   Gordon DE, Jang GM, Bouhaddou M*, et al.* A SARS-CoV-2 protein interaction map reveals targets for drug repurposing. *Nature* 2020; **583**: 459-468.

42.   Polacheck WJ, Kutys ML, Tefft JB, Chen CS. Microfabricated blood vessels for modeling the vascular transport barrier. *Nat Protoc* 2019; **14**: 1425-1454.

43.   Conceicao C, Thakur N, Human S*, et al.* The SARS-CoV-2 Spike protein has a broad tropism for mammalian ACE2 proteins. *PLOS Biology* 2020; **18**: e3001016.

44.   Lucas C, Wong P, Klein J*, et al.* Longitudinal analyses reveal immunological misfiring in severe COVID-19. *Nature* 2020; **584**: 463-469.

45.   Pober JS, Sessa WC. Evolving functions of endothelial cells in inflammation. *Nat Rev Immunol* 2007; **7**: 803-815.

46.   Rodrigues TS, de Sá KSG, Ishimoto AY*, et al.* Inflammasomes are activated in response to

592   SARS-CoV-2 infection and are associated with COVID-19 severity in patients. *Journal of*

593   *Experimental Medicine* 2020; **218**.

594   47.   Kasper J, Hermanns MI, Bantz C*, et al.* Inflammatory and cytotoxic responses of an alveolar-

595   capillary coculture model to silica nanoparticles: Comparison with conventional monocultures.

596   *Particle and Fibre Toxicology* 2011; **8**: 6.

597   48.   Kummer S, Avinoam O, Kräusslich H-G. IFITM3 Clusters on Virus Containing Endosomes

598   and Lysosomes Early in the Influenza A Infection of Human Airway Epithelial Cells. *Viruses*

599   2019; **11**: 548.

600   49.   Winstone H, Lista MJ, Reid A*, et al.* The polybasic cleavage site in the SARS-CoV-2 spike

601   modulates viral sensitivity to Type I IFN and IFITM2. *bioRxiv* 2020: 2020.2012.2019.423592.

602   50.   Buzhdygan TP, DeOre BJ, Baldwin-Leclair A*, et al.* The SARS-CoV-2 spike protein alters

603   barrier function in 2D static and 3D microfluidic in vitro models of the human blood-brain

604   barrier. *bioRxiv* 2020: 2020.2006.2015.150912.

605   51.   Bradley BT, Maioli H, Johnston R*, et al.* Histopathology and ultrastructural findings of fatal

606   COVID-19 infections in Washington State: a case series. *Lancet* 2020; **396**: 320-332.

607   52.   Manne BK, Denorme F, Middleton EA*, et al.* Platelet gene expression and function in patients

608   with COVID-19. *Blood* 2020; **136**: 1317-1329.

609   53.   Schurink B, Roos E, Radonic T*, et al.* Viral presence and immunopathology in patients with

610   lethal COVID-19: a prospective autopsy cohort study. *Lancet Microbe* 2020; **1**: e290-e299.

611   54.   Meizlish M, Pine A, Goshua G*, et al.* Circulating Markers of Angiogenesis and

612   Endotheliopathy in COVID-19. *medRxiv* 2020: 2020.2006.2029.20140376.

613   55.   Bost P, Giladi A, Liu Y*, et al.* Host-Viral Infection Maps Reveal Signatures of Severe COVID-

614   19 Patients. *Cell* 2020; **181**: 1475-1488.e1412.

615   56.   Short KR, Kasper J, van der Aa S*, et al.* Influenza virus damages the alveolar barrier by

616   disrupting epithelial cell tight junctions. *Eur Respir J* 2016; **47**: 954-966.

617   57.   Bernard I, Limonta D, Mahal LK, Hobman TC. Endothelium Infection and Dysregulation by

618   SARS-CoV-2: Evidence and Caveats in COVID-19. *Viruses* 2020; **13**.

58. Goeijenbier M, van Gorp ECM, Van den Brand JMA*, et al.* Activation of coagulation and tissue

fibrin deposition in experimental influenza in ferrets. *BMC Microbiology* 2014; **14**: 134.

59. Ranucci M, Ballotta A, Di Dedda U*, et al.* The procoagulant pattern of patients with COVID-

19 acute respiratory distress syndrome. *J Thromb Haemost* 2020; **18**: 1747-1751.

60. McDonald DM, Yao LC, Baluk P. Dynamics of airway blood vessels and lymphatics: lessons

from development and inflammation. *Proc Am Thorac Soc* 2011; **8**: 504-507.

61. Merad M, Martin JC. Pathological inflammation in patients with COVID-19: a key role for

monocytes and macrophages. *Nat Rev Immunol* 2020; **20**: 355-362.

62. Middleton EA, He XY, Denorme F*, et al.* Neutrophil extracellular traps contribute to

immunothrombosis in COVID-19 acute respiratory distress syndrome. *Blood* 2020; **136**: 1169-

1179.

63. Zuo Y, Yalavarthi S, Shi H*, et al.* Neutrophil extracellular traps in COVID-19. *JCI Insight*

2020; **5**.

 FIGURE LEGENDS

Figure 1. In vivo sections of lung show endothelial cells are not infected. Lung tissue from 4

deceased COVID-19 patients was sectioned and stained for Hematoxylin and Eosin

(H&E)(0). Consecutive sections were stained using RNAscope probes targeting either SARS-CoV-

2 spike mRNA or negative control probe and Syto13 for DNA. Patients 1 and 2 are positive for SARS-

CoV-2 spike mRNA (enlargement 1 and 2). Further enlargements of numbered insets (3-6) show vessels

outlined with red dashed line are negative for SARS-CoV-2 spike mRNA in the endothelial layer of the

vessels. Negative control probe shows non-specific background staining.

Figure 2. Endothelial cells express very low levels of ACE2 and TMPRSS2 receptors. A) qPCR

shows presence of mRNA for ACE2, TMPRSS2 and NRP1 in HUVEC, HMVEC-L and Calu-3 cells.

n=3 independent experiments. **B)** Representative immunofluorescent images of HUVEC, HMVEC-L

and Calu-3 cells stained for ACE2 (shown as single channel in top panel) (green), Phalloidin (magenta)

and DAPI (blue). Scalebar 30 μm. **C)** Quantification of ACE2 staining intensity. n=15 images; 5 images

per independent experiments. **D)** Western blot analysis showing ACE2 protein levels are absent in

negative control BHK-21 cells, lower in HUVEC and HMVEC-L compared to Calu-3 cells and

HUVEC + ACE2 overexpression and Human Nasal Epithelium cells (HNEp) as positive controls in top

panel. TMPRSS2 protein levels are lower in HUVEC and HMVEC-L compared to Calu-3 cells in

bottom panel **E)** Quantification of protein levels for ACE2 and TMPRSS2 in HUVEC, HMVEC-L and

Calu-3. n=5 (HUVEC and HMVEC-L), n=3 (Calu-3) independent experiments. **D)** Data are presented

as mean±s.e.m. with individual data points indicated and colour coded per independent experimental

replicate. Statistical significance was determined using a Kruskal Wallis test between Calu-3 and all

others (A, C, E). *P<0.05, ****P<0.0001.

Figure 3. Endothelial cells can be infected with 2 x 10^6 PFU of SARS-CoV-2, but infection is

abortive. A) Viral replication shown as number of PFU per ml of supernatant from SARS-CoV-2

infected HUVEC, HMVEC-L and Calu-3 cells at 24h, 48h and 72h after infection. n=2 (HUVEC and

HMVEC-L), n=3 (Calu-3) independent experiments. **B)** Representative immunofluorescent images of

HMVEC-L stained for NP (shown as single channel in top panel) (green), phalloidin (magenta) and DAPI (blue) with mock or SARS-CoV-2 infection from either apical or basolateral side of the cells at 72h after infection. Scalebar 50 µm. **B')** Enlargement of white box in B shows Phalloidin and DAPI single channels and merge. Scalebar 10 µm. **B'')** XZ projection of merged image in B', with apical side on top of image and basolateral side on the bottom. Scalebar 10 µm. **C)** Quantification of NP staining intensity in HMVEC-L. n=15 images from 3 independent experiments. **D)** Quantification of % NP positive cells that are extruded upon SARS-CoV-2 infection of HMVEC-Ls. n=15 images from 3 independent experiments. **E)** Western blot analysis showing NP protein levels in HUVEC, HMVEC-L and Calu-3 cells after 72h of infection. **F)** Quantification of NP protein levels in HUVEC, HMVEC-L and Calu-3. n=3 independent experiments. **G)** Representative immunofluorescent images of HMVEC-L and Calu-3 stained for NP (shown as single channel in top left panel) (green), cleaved caspase 3 (CC 3) (shown as single channel in top right panel)(magenta) and DAPI (blue) with mock or SARS-CoV-2 infection from apical side of the cells at 72h after infection. Scalebar 50 µm. **H)** Quantification of % NP positive cells that are positive for apoptosis marker cleaved caspase 3. n=15 images from 3 independent experiments. Data are presented as mean±s.e.m. with individual data points indicated and colour coded per independent experimental replicate. Statistical significance was determined using Kruskal Wallis test between 24h and other time points (A) or Mann-Whitney test between mock and + SARS-CoV-2 (C, F, H). *P<0.05, ***P<0.001, ****P<0.0001.

Figure 4. Endothelial cells are not infected with 2 x 10⁶ PFU of SARS-CoV-2 in 3D. A) Schematic of microfabricated device used to culture 3D vessel tubes under oscillatory flow. Cross section shows luminized tube covered with endothelial cells surrounded by extracellular matrix and virus can be flowed through the lumen of those tubes. **B)** Viral replication shown as number of PFU per ml of supernatant from SARS-CoV-2 infected HMVEC-L and Vero cells at 24h after infection. n=3 (HMVEC-L), n=2 (Ver0) independent experiments. Representative immunofluorescent images of **C)** HMVEC-L and **D)** Vero stained for NP (shown as single channel in top panel) (green), phalloidin (magenta) and DAPI (blue) with mock or SARS-CoV-2 infection in the lumen of the tube at 24h after infection. Scalebar 50 µm. YZ projection of **E)** HMVEC-L (merged image in C) and **F)** Vero (merged

image in D), with luminal side towards the right of image and basolateral side towards the right of

images. Scalebar 20 μm.

Figure 5. Endothelial cells mount an inflammatory response upon infection. Representative

immunofluorescent images of HMVEC-L stained for ICAM-1 (shown as single channel in top panel)

(green), NP (magenta), phalloidin (grey) and DAPI (blue) with **A)** mock or SARS-CoV-2 infection

from either apical or basolateral side of the cells at 72h after infection or with **B)** control and TNFα

treatment for 72h. Scalebar 50 μm. **C)** Quantification of ICAM-1 staining intensity in HMVEC-L. n=15

images from 3 independent experiments. **D)** Western blot analysis showing ICAM-1 and NP protein

levels in HMVEC-L and Calu-3 cells after 72h of infection. **E)** Quantification of ICAM-1 protein levels

in HMVEC-L and Calu-3. n=3 independent experiments. Measurement of cytokines with AlphaLISA

Immunoassay kit for **F)** CXCL10 and **G)** IL6 in the supernatant of HMVEC-L with mock or SARS-

CoV-2 infection from either apical or basolateral side of the cells at 72h after infection. n=4 independent

experiments. Data are presented as mean±s.e.m. with individual data points indicated and colour coded

per independent experimental replicate. Statistical significance was determined using Mann-Whitney

test between mock and + SARS-CoV-2 (C, E, F, G). *P<0.05, ****P<0.0001.

Figure 6. Co-culture shows infection of Calu-3 and inflammation of HMVEC-L. A) Representative

immunofluorescent images of co-cultured Calu-3 and HMVEC-L stained for NP (shown as single

channel in top panel) (magenta), ICAM-1 (shown as single channel in middle panel) (green), phalloidin

(grey) and DAPI (blue) with mock or SARS-CoV-2 infection at 72h after infection (2 x 10^6 PFU).

Scalebar 50 μm. **B)** Quantification of NP staining intensity in Calu-3 and HMVEC-L. n=9 images from

3 independent experiments. **C)** Schematic of Calu-3 and HMVEC-L co-cultures on transwell

membranes. **D)** Quantification of ICAM-1 staining intensity in Calu-3 and HMVEC-L. n=9 images

from 3 independent experiments. **E)** Representative immunofluorescent images of co-cultured Calu-3

and HMVEC-L stained for dsRNA (shown as single channel in top panel) (magenta), phalloidin (grey)

and DAPI (blue) with mock or SARS-CoV-2 infection at 72h after infection. Scalebar 50 μm. **F)**

Quantification of dsRNA staining intensity in Calu-3 and HMVEC-L. n=9 images from 3 independent

experiments. **G)** qPCR shows presence of viral RNA (MPRO) in SARS-CoV-2 infected Calu-3 and

HMVEC-L co-cultured cells represented as fold change relative to mock infection at 72h after infection.

n=3 independent experiments. **H)** Viral replication shown as number of PFU per ml of supernatant from

SARS-CoV-2 infected Calu-3 and HMVEC-L co-cultured cells at 72h after infection. n=3 independent

experiments. Measurement of cytokines with AlphaLISA Immunoassay kit for **I)** CXCL10 and **J)** IL6

in the supernatant of Calu-3 and HMVEC-L co-cultured cells with mock or SARS-CoV-2 infection at

72h after infection. n=4 independent experiments. Data are presented as mean±s.e.m. with individual

data points indicated and colour coded per independent experimental replicate. Statistical significance

was determined using Mann-Whitney test between mock and + SARS-CoV-2 (B, D, F, G, H, I, J).

*P<0.05, **P<0.01, ****P<0.0001.

Supplemental Table 1. Patient information. Additional information on cohort of COVID-19 patients.

Supplemental Figure 1. Endothelial cells are not productively infected with 6 x 10^4 PFU of SARS-

CoV-2. Schematic of **A)** apical and **B)** basolateral infection of cells cultured on transwell membranes.

C) Viral replication shown as number of PFU per ml of supernatant from SARS-CoV-2 infected

HUVEC, HMVEC-L and Calu-3 cells at 24h, 48h and 72h after infection. n=2 (HUVEC and HMVEC-

L), n=3 (Calu-3) independent experiments. Representative immunofluorescent images of **D)** HUVEC,

E) Calu-3 or **F)** HMVEC-L stained for NP (shown as single channel in top panel) (green), phalloidin

(magenta) and DAPI (blue) with mock or SARS-CoV-2 infection from either apical or basolateral side

of the cells at 48h after infection. Scalebar 50 μm. **G)** Quantification of NP staining intensity in

HUVEC, HMVEC-L and Calu-3. n=9 images from 3 independent experiments. **H)** Western blot

analysis showing NP protein levels in HUVEC, HMVEC-L and Calu-3 cells after 48h of infection. **I)**

Quantification of protein levels for NP in HUVEC, HMVEC-L and Calu-3. n=2 independent

experiments. Data are presented as mean±s.e.m. with individual data points indicated and colour coded

per independent experimental replicate. Statistical significance was determined using Kruskal Wallis

test between 24h and other time points (C) or Mann-Whitney test between mock and + SARS-CoV-2

(G, I). *P<0.05, **P<0.01

Supplemental Figure 2. Positive and negative controls for infection with 6 x 10⁴ PFU of SARS-CoV-2. A) Viral replication shown as number of PFU per ml of supernatant from SARS-CoV-2 infected BHK-2 and Calu-3 cells at 72h after infection. n=3 independent experiments. **B)** Western blot analysis showing NP protein levels in BHK-21 and Calu-3 cells after 72h of apical infection with mock or SARS-CoV-2. **C)** Quantification of protein levels for NP in BHK-21 and Calu-3. n=3 independent experiment. **D)** Representative immunofluorescent images of BHK-21 and Calu-3 cells stained for NP (shown as single channel in top panel) (green), phalloidin (magenta) and DAPI (blue) with mock or SARS-CoV-2 infection from apical side of the cells at 72h after infection. Scalebar 50 μm. **E)** Quantification of NP staining intensity in BHK-21 and Calu-3. n=9 images from 3 independent experiments. Maximum secretion levels of **F)** CXCL10 and **G)** IL6 in the supernatant of HMVEC-L with mock, TNFα or IFNß treatment from the apical side of the cells at 4h and 16h after treatment. n=2 independent experiments. Data are presented as mean±s.e.m. with individual data points indicated and colour coded per independent experimental replicate. Statistical significance was determined using Mann-Whitney test between mock and + SARS-CoV-2 (C, E). ****P<0.0001

Supplemental Figure 3. Full scans of Western blots. Full scans of Western blots shown in the paper figures. Used lanes are labelled as in the corresponding figure, with other lanes labelled as X. Dashed line indicates where membrane was cut. Underlined proteins indicate which blot was used in the corresponding figure and time indicates exposure time.

Figure 1

Figure 2

A Receptor mRNA expression

B HUVEC, HMVEC-L, Calu-3 — ACE2; ACE2/Phalloidin/DAPI

C ACE2 intensity

D Western blots: BHK-21, HUVEC, HMVEC-L, Calu-3, HUVEC + ACE2, HNEp — ACE2, non-specific band, Actin, Actin long exposure; HUVEC, HMVEC-L, Calu-3 — TMPRSS2, GAPDH

E WB quantification

Figure 3

Figure 4

Figure 5

A HMVEC-L

Apical | Basolateral
Mock | + SARS-CoV-2 | Mock | + SARS-CoV-2

ICAM-1

ICAM-1/NP Phalloidin/DAPI

B HMVEC-L

Apical
PBS | + TNFα

ICAM-1

ICAM-1 Phalloidin/DAPI

C ICAM-1 intensity

mock · PBS ■
+SARS-CoV-2 ▲ + TNFα ▼

RawIntDen/Area x100

**** * ****

Apical Basolateral Control
HMVEC-L

D

HMVEC-L | Calu-3
Apical | Basolateral | Apical

Mock +SARS-CoV-2 Mock +SARS-CoV-2 Mock +SARS-CoV-2

110kDa — ICAM-1
50kDa — NP
37kDa — GAPDH

E ICAM-1 expression

mock · +SARS-CoV-2 ▲

Intensity corrected for GAPDH (a.u.)

*

Apical Basolateral | Apical
HMVEC-L | Calu-3

F CXCL10 secretion

mock · +SARS-CoV-2 ▲

CXCL10 (ng/mL)

Apical Basolateral
Site of infection

G IL6 secretion

mock · +SARS-CoV-2 ▲

IL6 (ng/mL)

* *

Apical Basolateral
Site of infection

Figure 6

Supplemental Table 1: Patient information

*representative images derived from this patient

Patient ID	Diagnosis	Sex	Age	Co-morbidities	Histology	D-Dimer at admission
LN1*	COVID-19	F	87	Systemic Arterial Hypertension, Dyslipidemia, Hypothyroidism, Senile dementia	Acute phase DAD	2014
LN2	COVID-19	M	53	Class II obesity	Acute phase DAD	425
LN3	COVID-19	F	85	Systemic Arterial Hypertension, Brain Stroke, Vascular Dementia	Acute phase DAD	N/A
LN4	COVID-19	M	73	Type 2 Diabetes Mellitus, Chronic Kidney Disease Dialysis, Atrial Fibrillation, Coronary Disease, Heart Failure, Peripheral Obstructive Artery Disease	Reactive Type 2 pneumocyte hyplerplasia	3436
LN6	COVID-19	M	80	Systemic Arterial Hypertension, Coronary Disease, Heart Failure, Class III obesity	Reactive Type 2 pneumocyte hyplerplasia	816
LN7	COVID-19	M	81	Systemic Arterial Hypertension, Chronic Kidney Disease Dialysis, Brain Stroke	Reactive Type 2 pneumocyte hyplerplasia	13662
LN8	COVID-19	N/A	N/A	N/A	Organising pneumonia	N/A
LN9	COVID-19	M	86	Prostate cancer, Abdominal Aortic Aneurysm, Giant Cells Arteritis	Organising pneumonia	11184
LN10	COVID-19	M	46	Dyslipidemia	Organising pneumonia	594
LN11	COVID-19	F	93	Type 2 Diabetes Mellitus, Systemic Arterial Hypertension, Dyslipidemia, Senile dementia	Acute phase DAD	6544

Supplemental Figure 1

Supplemental Figure 2

A Viral replication

B BHK-21 / Calu-3 — Apical — Mock / +SARS-CoV-2 — NP 50kDa, Actin 42kDa

C NP expression — mock / +SARS-CoV-2

D BHK-21 / Calu-3 — Apical — Mock / + SARS-CoV-2 — NP; NP/Phalloidin/DAPI; 50 µm

E NP intensity — mock / +SARS-CoV-2 — ****

F CXCL10 secretion — 4 h / 16 h — PBS, TNFα (100 ng/ml), IFNβ (10 ng/ml)

G IL6 secretion — 4 h / 16 h — PBS, TNFα (100 ng/ml), IFNβ (10 ng/ml)

Supplemental Figure 3
Fig 2D top panel

Fig 3D

Fig 2D bottom panel

Fig 5D

Supplemental Figure 3

Fig S1H

Fig S2B